

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Methicillin-Resistant *Staphylococcus aureus* (MRSA) in Food-Producing and Companion Animals and Food Products

Jungwhan Chon, Kidon Sung and Saeed Khan

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/66645>

Abstract

Methicillin-resistant *Staphylococcus aureus* (MRSA) has become a growing concern in companion and food-producing animals. The presence of multidrug-resistance with a wide range of extracellular enterotoxin genes, virulence factors, and Panton-Valentine leukocidin (*pvl*) cytotoxin genes confer life-threatening traits on MRSA and makes them highly pathogenic and difficult to treat. Clonal complex 398 (CC398), a predominant clonal lineage of livestock-associated-MRSA in domestic animals and retail meat, is capable of infecting humans. In order to monitor and prevent MRSA contamination, it is critical to understand its source and transmission dynamics. In this review, we describe MRSA in food-producing animals (pig, cattle, chicken), horses, pet animals (dogs, cats), and food products (pork, beef, chicken, milk, and fish).

Keywords: MRSA, companion animals, food-producing animals, food products, CC398

1. Introduction

Staphylococcus aureus is a pathogen that causes both human and animal infections and food intoxication [1–3]. It causes simple infections, such as furuncle, boil, stye, impetigo, carbuncle, and keratitis, and serious infections, including septicemia, necrotizing pneumonia, endocarditis, osteomyelitis, and pericarditis [4–7]. Shortly after the introduction of methicillin in clinical practice to control penicillin-resistant staphylococci, the first methicillin-resistant *S. aureus* (MRSA) was isolated [8]. MRSA is one of the most important hospital-acquired pathogens that are resistant to various antimicrobials, thereby making their treatment complicated [9]. MRSA

in humans is usually divided into two groups: hospital-associated MRSA (HA-MRSA) and community-associated MRSA (CA-MRSA) [10]. A third group of MRSA, known as livestock-associated MRSA (LA-MRSA), now has emerged and infects livestock, pets, and wild animals.

LA-MRSA was first detected in milk with bovine mastitis from Belgium in 1972 [11–13]. Thereafter, MRSA reports in various food and companion animals, such as pigs, cattle, chickens, dogs, cats, and horses, have increased [11, 14]. A novel strain of MRSA belonging to multi-locus sequencing type (MLST) 398 (ST398) and related strains collectively grouped into clonal complex 398 (CC 398) have been frequently found in pigs, chickens, veal calves, dairy cattle, horses, dogs, and milk in various countries [11]. Both methicillin-susceptible *S. aureus* (MSSA) and MRSA have been associated with companion and food production animals [15–19]. The most significant of these is intramammary infection of dairy cattle leading to mastitis, which causes a substantial economic loss to the dairy industry worldwide [6, 20, 21]. The CC398 *S. aureus* isolate was more prevalent in nasal swabs of pig and cattle farmers than of non-farming human controls [22, 23]. An examination of livestock-associated MRSA (LA-MRSA) in human case isolates in the Netherlands indicated an increase from 0% in 2002 to greater than 21% in mid-2006 [23] and 35% in 2009 [24]. In most European countries, CC398 remains the most commonly identified type of LA-MRSA [15, 25–27]. However, the epidemiology of LA-MRSA differs in other geographic areas. A different strain of LA-MRSA, CC9, appears to be the prominent type in several Asian countries [28–32]. Poultry may harbor CC398 strains [16, 33, 34] but CC5 [33, 35] and other types unrelated to CC398 have also been reported [36]. The diversity of LA-MRSA in the USA appears to be higher than that identified in Europe or Asia, with reports of both CC398 as well as a variety of “human” types of *S. aureus* in livestock.

LA-MRSA infections among livestock animals and associated farmers are of great concern as these sources could potentially serve as reservoirs for zoonotic infections [14]. Contamination of food with enterotoxin producing *S. aureus* leads to over 240,000 cases of food-borne illness in the United States annually. Although most *S. aureus*-related food-poisoning incidents are self-limiting and go away within 2 days, some serious infections have been reported as well [4, 5]. A large number of the reported staphylococcal food-poisoning outbreaks can be traced back to a human source harboring *S. aureus* producing certain staphylococcal enterotoxins (SEs) [1, 37]. Most of the LA-MRSA strains, particularly the ST398 group, do not appear to code for any of the known SEs [11, 38–42]. However, genes for SEs B, K, and Q have been detected in MRSA CC398 strains isolated from geographically diverse pig farms in Germany [43]. The acquisition of enterotoxin genes along with the virulence factors, such as Panton-Valentine leukocidin (*pvl*) genes by LA-MRSA may eventually pose a threat to humans, suggesting that animals have the potential to be a source of primary contamination as well [16, 44].

2. MRSA in food-producing animals

2.1. Porcine MRSA

In 2005, pigs were reported as an animal reservoir of *S. aureus* in France, including MRSA [17]. Pig farmers were more frequently colonized than nonfarmers and one of the most

prevalent strains of *S. aureus* in pig farmers was CC398 [45–49]. Forty-five percent of veterinarians attending pig farms in the Netherlands were positive for MRSA [48]. In Belgium and Denmark, the prevalence of MRSA in veterinarians was 9.5% and 1.4%, respectively [47]. German researchers reported that MRSA ST398 that carried SCC_{mec} IV or V, accessory gene regulator type I and capsule type 5 [25] in pig primary production herds was higher in their country (45–70%) than in the rest of the European Union [45, 46]. In an interesting report, 12.5% of attendees at an international meeting concerning pig health carried MRSA and 91.2% (31/34) of them CC398 [50]. While early studies on farms and of meat identified CC398 strains in animals, farm workers, and meat products [51, 52], other studies also documented CC398 in populations with no obvious livestock contact [53–55]. The emergence of this strain was also reported in pigs and pig farmers in the Netherlands [22, 56, 57], Denmark [58], Germany [59], and Canada [60]. Detection rates of MRSA in breeding and production herds were 46%, 43.5%, and 40% in Spain, Germany, and Belgium, respectively, but no single strain of MRSA was found in Finland and Denmark in 2008 [61]. The majority of LA-MRSA lineage belonged to ST398, accounting for 92.5% of the MRSA isolates. Other ST types, human-associated MRSA ST1 and cattle-associated ST97 in finishing holdings, and ST9 in the same animal species in Europe [62], CC9, and CC49 in Switzerland [63] have also been reported.

Various farm types in the Netherlands were reported to have MRSA in 23–71% of their pigs, and it was especially high in farms with finishing pigs (pigs are almost ready to be sent to market) [49, 57, 64]. The presence of MRSA is dependent on pig production type and herd size and increases from 31 to 86% depending upon small-, medium-, and large-sized farms carrying <250, >500, and >1000 animals, respectively [45, 62, 64, 65]. MRSA prevalence also varied with farm type, e.g., fattening and closed (farrow-to-finish) farms exhibited 94 and 56% MRSA, respectively [66]. Transportation from farm to slaughterhouse [38, 67], lairage [38], national and international trade [57], and slaughter house employees all have been reported to enhance MRSA contamination and may play important roles in transmission of the bacteria [68]. It was proposed that MRSA contamination in piglets is dependent on the status of sows [69]. When a sow was colonized with MRSA, 100% piglets were MRSA-positive. However, 84% piglets were MRSA-positive when there was no MRSA contamination in a sow. Higher numbers of MRSA were isolated in suckling (52.9%) and weanling piglets (53.4%) than sows (38.3%) [64]. Prevalence of MRSA in pigs has been linked to their age but the data are not conclusive. MRSA were identified in 100% of 9–12-week-old pigs, whereas in adult animals it decreased to 36% [51]. On the other hand, Weese et al. [69] reported that MRSA was more prevalent in post-weaning (85%) than preweaning pigs (34.5%) in pig farms without antimicrobial treatment. Percent of MRSA colonization in Canadian piglets on days 1, 28, 56, and 70 were 1, 34, 50, and 42%, respectively [69]. Khanna et al. [60] reported no variation in MRSA prevalence based on age groups. MRSA does not seem to cause serious infection in pigs, but there have been a few reports of MRSA from exudative epidermitis lesions of piglets on a breeding farm [70] and in pigs suffering from infection of the urinary-genital tract, skin infection, and metritis-mastitis-agalactia syndrome [71]. While MRSA ST398 isolated from diseased pigs did not carry the major virulence genes, such as toxic shock syndrome toxin 1, *pvl*, and exfoliative toxins, they carried some virulence genes, such as α - and δ -hemolysins, proteases, capsule type-specific genes, microbial surface components recognizing adhesive matrix

molecules, biofilm-associated, and enterotoxin genes [44]. Their MLST, *spa*, and SCC*mec* types were identified as ST398, t011, and IV, respectively.

The epidemiology of livestock-associated *S. aureus* in the USA appears to be notably different than that in European countries. Most of the porcine MRSA isolates in Canada, Europe, Peru, and USA were CC398 [65]. A human epidemic clone, Canadian MRSA-2 (CMRSA-2, USA100, CC5), was found in nasal and rectal swabs of pigs in Canada [60]. This isolate was the most common cause of health care-associated infections in Canada. CMRSA-5 (USA500, ST8) also isolated from retail pork in Canada, is a human epidemic strain that also has been documented in horses and horse personnel [72]. Three *spa* types (t011, t034, t108) within CC398 were the most frequent in breeding and production herds in Europe, and t108 was most popular only in the Netherlands among European countries. On the contrary, prevalence of t899 in Italian breeding and production herds was between 24% and 27% [62, 65]. In Italy, CC1 and CC97 lineages among MLST types that do not belong to CC398 were significantly high in the primary production of pigs [62]. In one study carried out in a jail setting in Texas, CC398 isolates made up of 13.2% of all MSSA identified within this population. Apart from CC398, other human strains of *S. aureus* have also been found in USA livestock. Studies carried out on swine farms in the USA have identified human strains within the noses of live animals [55, 73, 74] and farm dust [75]. Several papers have reported CC5 strains rather than CC398-associated types to be the dominant strain isolated from pig farms in both Iowa and Ohio [75, 76], whereas others have found CC398 to be the most common molecular type [51, 76]. Studies on workers on pig farms and in processing plants found substantial diversity within *S. aureus* isolates, including CC398, CC5, and CC8 strains, among others [77–79]. MRSA attributed to ST5 was recently reported in pigs in the USA [38]. A different swine-associated MRSA strain, CC9, is circulating among pigs and pig farmers in China and Malaysia [29, 30, 80, 81]. MRSA ST22, known as human epidemic clone EMRSA-15 in the UK, was also discovered in pigs and in hospitalized patients in Singapore with an elevated frequency [82, 83]. High frequency of MRSA CC9 with *spa* types t899 and t4358 was reported in porcine samples in Asian countries, whereas it was not found as much in Europe or the USA [28–30, 84]. In Thailand, porcine MRSA ST9 isolates had a unique *spa* type (t337) and SCC*mec* type (SCC*mec* IX) that were different from other LA-MRSA ST9 strains found in Asian countries [28]. In 2012, Lim et al. [85] reported two ST types, livestock-associated ST398 and human-associated ST72, from pigs, which was the first finding of ST398 in Korean pigs.

2.2. Bovine MRSA

The first report of MRSA in farm animals was published in the early 1970s, when the bacteria were isolated from the milk of dairy cows with mastitis in Belgium [86] and clustered in the CC398 group [87]. Devriese and Hommez [88] suspected that these samples were most likely contaminated by humans. In the past few years, MRSA has been isolated from cows or their milk in Korea [89–91], Hungary, Mexico, and the Netherlands [89, 92, 93]. There have also been numerous reports of MRSA from cows or their milk in Brazil, Italy, Pakistan, Nigeria, Turkey, and the USA [94–96]. Subsequently, several reports have described bovine udder infections caused by LA-MRSA CC398 [97]. In Dutch farms, MRSA was detected in 18–31% of veal calves [98]. In 2010, the European Union reported that 20% of veal calves in Germany

carried MRSA [99]. A survey of 51 veal calf farms in the Netherlands indicated that an average of 38% of farmers and 16% of family members were colonized with LA-MRSA [19, 100]. Recently, another group of LA-MRSA strains (CC130, CC425, and CC1943) that were initially thought to be bovine-specific lineages emerged in humans [101]. A number of multidrug-resistant MRSA were isolated from bovine mastitis in Germany [97, 102], the majority of which were MRSA ST398 related to animal strains, but an isolate of the clonal complex group CC8 was identified as a human epidemic MRSA strain Irish-01 [103]. In 2010, Hata et al. first discovered MRSA in cow's milk in Japan and the genotypes (ST5-SCC*mec* II) were the same as or similar to human strains [104]. There has been a dramatic surge in human CC398 infection and colonization in the Netherlands, increasing from 0% in 2002 to more than 21% in 2006 [23]. A recent Dutch study indicated that the annual incidence of MRSA in humans more than tripled from 2001 to 2006 where 23%, 26%, 16%, and 10% of the patients acquired MRSA from a foreign hospital, animals, nosocomial transmission, and the community, respectively [105]. The presence of MRSA CC398 in pig farms with a concomitant increase in CC398 infections in humans clearly suggests that pigs or cattle are specifically a risk factor and CC398 MRSA colonization and prevalence in humans is associated with animal contact [23, 46, 106, 107]. The above argument is further strengthened by the findings that the CC398 carrier status of farm workers decreased dramatically when they took a break from direct animal care duties [19, 100]. People who visited farms to collect samples for a shorter duration carried MRSA transiently as compared to those who had prolonged visits, suggesting that a prolonged contact with animals is probably an important factor for higher rates of colonization [49].

While a majority of the MRSA collected from dairy cattle belonged to ST398 [89], other ST types, such as ST1-t286-SCC*mec* IV, ST72-t324 [108], ST59-t437-V [91], ST10-t127-SCC*mec* IVa genotype [92], SCC*mec* types IVg [109], CC97, t4795, and t1730 [110], and a *mecA* variant (*mecA*_{LGA251}) known as *mecC*, are also reported from MRSA CC130 and ST425 isolates [111]. The *mecC* type was also detected in Danish MRSA CC130 isolated from a cow and the genotypic characteristics, such as *spa* type (t843), MLVA (MT429) and PFGE profiles of bovine isolates were the same as the human isolates, implying transmission between humans and ruminants [112]. The geographic variation in the prevalence and origin of CC398 colonization and incidence of infection in humans is quite interesting. MRSA has been detected in retail beef, but nasal and fecal sampling of nearly 500 Canadian feedlot cattle, shortly before slaughter, detected no MRSA [113]. Moreover, while CC398 MRSA infection in humans is a leading cause of CA-MRSA infection in some European countries, it is rare in North America despite the presence of CC398 in livestock [114]. The reasons for the low incidence of CC398 infections in the USA may include differences in direct and indirect contact with food animals, much lower population density in North American pig-rearing regions, and the common presence of other competing MRSA strains in people in the general population. Although some studies suggested that the MRSA present in cattle is bovine-specific, most of the reports indicated that MRSA found in cattle were derived from humans [89, 92, 115, 116]. Bovine, porcine, canine, feline, and equine MRSA isolates containing the *pvl* gene and other virulence factors, such as *chp*, *scn*, *seb*, *sek*, and *seq*, toxic shock syndrome toxin 1 (*tsst1* or *tst*) gene [91, 117], hemolysin, protease, superantigen-like protein, capsule, and biofilm-associated genes [118–121], may pose a potential threat to public health.

2.3. Poultry MRSA

CC398 is not limited to large livestock animals alone; it has also been reported in poultry [18, 122], manure from chicken farms and soil fertilized with this manure [123]. However, the numbers of MRSA ST398-t011-SCC_{mec} V isolated from chickens are lower (0–28%) than in pigs (82–92%) on the same farm [124]. In another study from Belgium [109], MRSA present in 0.8–1.8% layers and broilers (chickens raised for meat) was clustered into two ST types, ST398 (t011, t899) and ST239 (t037). Two other studies from Belgian broiler farms [18, 125] reported 12.8–14.3% of randomly selected Belgian broiler farms to be positive for CC398. In 2010, the Federal Institute for Risk Assessment reported a contamination rate of 32% in turkey meat in Berlin, Germany [126]. Similar contamination frequencies were reported from Canada and the USA [72, 127], as well as from Taiwan [128]. Moon et al. [91] reported that the *S. aureus* isolated from chicken carcasses contained 1.3% MRSA, which was more than the MRSA isolated from other animal carcasses (0.3%) in Korea. Poultry-associated *S. aureus* isolates belonging to genotypes other than CC398 have also been reported from different geographic regions [36, 129–131]. Mulders et al. [132] reported that 6.9% of MRSA present in broiler chickens in the Netherlands represented the ST9-t1430 genotype. A single *spa* type, t1456, in poultry was seen and distinguishable from the *spa* types of ST398 observed in other animals in Belgium [125]. Genotypic and antimicrobial patterns between 14 MRSA isolates from broilers and pigs were identical [133]. The MRSA isolates from Korea had the genotype of ST692-t2247-III [91] and the MRSA isolates from Hong Kong exhibited the genotypes ST9-t899-IV [81] and CC9 (t899, t1234) [134]. A study from Denmark analyzed the isolates from infected poultry and detected a predominant common human epidemic clone CC5 [129]. Using a population genomics approach, Lowder et al. [36] examined the origin of *S. aureus* isolates from diseased and healthy poultry from four continents and found that the majority of isolates belonged to a single clonal complex CC5 belonging to a known human-associated lineage. The poultry isolates were more closely related to each other than to human CC5 isolates, but were most similar to a subclade of CC5 that was circulating in Polish hospitals in the 1980s. In a study conducted in Korea, 930 food samples were collected, and four strains of the CA-MRSA CC5 human clone were identified [135]. In a human case study, a 63-year-old Dutch woman who owned a chicken farm developed a life-threatening endocarditis; the infecting MRSA isolate was identified as CC398 [136], similar to an isolate found in a pig farm nearby and to MRSA isolates previously found in other pig farms in the Netherlands.

2.4. Other meat products

S. aureus is found frequently in a variety of retail meat products. A Dutch Food Safety Agency analyzed 2217 samples of various kinds of meats from the retail stores and found that 11.9% of 2217 samples had MRSA [137]. The distribution of MRSA within various meat types was listed as follows: beef, 10.6%; veal, 15.2%; lamb and mutton, 6.2%; pork, 10.7%; chicken, 16.0%; turkey, 35.3%; fowl, 3.4%; and game, 2.2%. Of all the MRSA isolates, 85% of the isolates belonged to ST398; the other STs were possibly of human origin [137]. Another Dutch survey found that 46% of retail meat samples contained *S. aureus* strains, of which two (2%) were MRSA: one was CC398 and the other was USA300 [138]. Studies in Switzerland and Japan showed the prevalence of *S. aureus* in meat products to be 23 and 65%, respectively [122, 139].

A USA survey of 120 retail meat samples indicated that 39.2% contained *S. aureus* strains, 5% of which were MRSA of the types USA100 (ST5) and USA300 (ST8) [127]. A Canadian survey [72] found that 7.7% of retail meat samples harbored MRSA; 30% belonged to the clonal complex CC398, 40% were CC8, and 30% were CC5, a strain commonly found in humans in both the USA and Canada.

2.5. Milk

In general, the occurrence of MRSA in bovine mastitis isolates is well studied and its prevalence seems to be very low [140]. Following the initial reports of isolation of MRSA from mastitic cows [86], sporadic cases of MRSA in dairy cattle were detected among *S. aureus* isolates from clinical or subclinical mastitis. In one of the studies from Korea [109], MRSA were isolated from the milk of cows with an isolation ratio of 0.18%. In one report on dairy farms in Belgium, a high percentage (15%) of MRSA was found in lactating cows [140]; these cows had a previous history of MRSA. The long-term low prevalence of MRSA mastitis is quite surprising, given the number of years since the first identification of MRSA in cattle and the close contact of humans with the udders of dairy cattle. In Germany, the highest proportion of positive samples (45%) was found in nasal swabs from veal calves at slaughter and the lowest rate was 4.1% in bulk tank milk. Most isolates, irrespective of origin, were from *spa* types t011 and t034 belonging to the clonal complex CC398 [141]. The finding of LA-MRSA CC398 in tank milk suggests udder colonization and possibly cases of subclinical mastitis in dairy cattle in Germany [141]. Close contact of dairy cattle with humans could lead to a transfer of strains between them. In one of the reports from Hungary, MRSA isolates from mastitic cows and a worker were found identical by phenotypic and genotypic analysis indicating a transfer between cows and human [92].

2.6. Fish

Fish is not a normal host for staphylococci and its presence on fish is either due to disease in the fish, contamination, or poor personnel hygiene. The first report of the isolation of MRSA from Tilapia was published in 2010, where 559 *S. aureus* isolates from the brain, eyes, and kidneys of tilapia from 11 farms collected for a period of 2 years were analyzed and 50% were identified as MRSA [142]. In another study [143] from Korea that analyzed 165 *S. aureus* strains isolated from different food samples between 2003 and 2006, four were identified as MRSA. Two of these were from beef and two from fish. The two fish isolates, one from sea bass and other from rockfish, were identified as ST1 and ST72, respectively. An analysis of 200 ready to eat (RTE) fish samples collected from 10 shops belonging to four supermarket chains in Japan, 5 were MRSA and 5 others were identified as coagulase-negative MRSA [144]. Molecular typing of two MRSA isolates by *spa* sequencing and MLST typing identified t1767 and ST8, respectively. Interestingly, MRSA ST8 strains have been predominantly isolated from humans in the USA and Europe but are of rare occurrence in foods in Japan. It is not certain if the MRSA in fish was from human or fish origin. In another report from Greece, one hundred samples from RTE fish products were examined and two were reported to have MRSA belonging to the *spa* types t316 (ST359) and t548 (ST5) [145]. In a recent report, a patient developed foot infection with MRSA after a fish pedicure [146], but the origin of the MRSA

could not be determined in this case. Reports of fish from Egypt, India, and Yemen have also been reported to harbor as much as 3.5% MRSA [147]. In this report, two of the MRSA isolates were found to harbor the enterotoxin genes *seg* and *sei*. Since the global spread of multi-drug-resistant bacteria has increased in the past decade, the finding of enterotoxigenic MRSA in fish should be of concern. The global trade of fish increases the possibility of intercontinental transmission of multidrug-resistant and enterotoxigenic *S. aureus* and its potential influence on consumer health worldwide should be monitored.

3. MRSA in food processing environment

After carcasses leave the slaughter-house chillers, residual MRSA on carcass surfaces can be transmitted during further processing through human hands, cutting tools, and any surfaces with direct meat contact. Manual handling during processing also can facilitate the entry of human MRSA strains into the production units. Recent surveillance data suggest that 22.5–64.8% of retail beef, pork, chicken, and turkey meats in five different geographical locations in the United States were contaminated with *S. aureus* [148–150]. A Swiss meat-processing plant reported the presence of *S. aureus* on 22.7% of the received chilled pork hind quarters from 18 European suppliers [151]. While investigating German pork processing units, Kastrup [139] determined a MRSA detection frequency of 6% on meat trimmings, 2% on processing equipment, and 5% on employees. Beneke et al. [152] obtained a similar detection rate in the processing area of a German abattoir. In an experimental setting, *S. aureus* at a contamination level of 5–7 log CFU/100 cm² was detectable on dry stainless steel for at least 96 h. In The Netherlands, de Jonge et al. [39] assessed the presence of MRSA in three meat-processing facilities and two institutional kitchens. MRSA was not isolated from any human nose or hand swabs, but 33% of the participants carried MSSA and only 14.3% of the meat samples were contaminated with MRSA. A Dutch study [138] which found that 46% of the retail meat samples, the majority of which came from a single retail shop and contained *S. aureus*, had a high degree of clonal relationship, indicating cross-transmission at some point during processing in the shop.

To pin point the exact source of contamination, it is necessary that the process of slaughtering be analyzed critically. Slaughter and meat processing involve several steps, any of which could introduce contamination with MRSA. Scalding, the first step in the slaughter process, is carried out at 60–62°C for 6–8 min in scalding tanks with rotating bars or through long scalding tanks [153] to loosen the hair from the carcass. An analysis of the effect of scalding on the quantity of coagulase-positive *S. aureus* (CPS) on pig carcasses in two Swiss abattoirs indicated variable data [154]. CPS, isolated from 96 to 100% of all carcasses, was reduced to 18 and 20% along the slaughter line after scalding from one abattoir, but in the second abattoir, it increased to 99% at the end of the line. Dehairing that follows scalding is another critical step that involves mechanical treatment of the carcass with rotating scrapers and rubber flails. This step has a potential to increase dissemination of porcine bacteria from mouth, nose, skin, and intestinal tract due to the accumulation of detritus in the machine. Singeing, which involves the exposure of the carcass for 10–15 s at 900°C, has been reported to decontaminate the surface of pig carcasses and lead to a 2.5- to 3-log reduction in total bacterial counts [153, 155].

Using a probabilistic model, Vossenkuhl et al. [156] found that a high MRSA prevalence at the beginning of the slaughter line was reduced to a low level at the end of the slaughter line (**Figure 1**). However, some investigators reported no effect of singeing on the microflora [157], whereas others have indicated that the reduction achieved by singeing is frequently reversed by polishing, that cleans intensively a skin surface [158–160]. Evisceration of the intestinal tract is another source of contamination with fecal bacteria on the surface of carcasses [154, 161]. To minimize the bacterial contamination, pig carcasses are usually chilled overnight using conventional single-stage chilling regimes, spray chilling, ice bank chilling in humid air at 2°C, and rapid or ultra-rapid chilling [162, 163]. Spescha et al. [154] reported a 77% decrease in the proportion of *S. aureus*-positive carcasses after chilling. Freeze chilling at temperatures of –10 to –25°C for 45–60 min, followed by chilling at 2°C for 23 h reduced *S. aureus* by 1 log CFU cm^{–2} on untrimmed carcasses [151]. It is clear from the published reports that handling of carcasses, proper maintenance of equipment, and personal hygiene play critical roles in the control and spread of *S. aureus* in the final end product.




Figure 1. Change in MRSA prevalence along the slaughter line depending on the variation of the initial MRSA prevalence $P(s_0^+)$ [156]. Reproduced with permission of Elsevier.

4. Molecular epidemiology of *S. aureus* CC398

The emergence of LA-MRSA strains in humans [16, 17, 25, 56] and the presence of an identical MRSA CC398 in pigs, farm workers, veterinarians who attended to the same pig farms and their nonexposed family members [47, 48] suggests animal-to-human or human-to-animal transmission. In an interesting report from the Netherlands, it was shown that farm visitors were positive for CC398 MRSA directly after a farm visit but tested MRSA negative after

24 h [19, 41]. These and other studies indicate that CC398 appears to be frequently shared between animals and humans and is capable of causing infections in both species [25, 70, 164]. Transmission of MRSA between animals and humans is not new, but the MRSA isolates, in most cases, represent an initial human-to-animal transmission [24, 49, 165, 166].

Analysis of MRSA and MSSA from animals and humans spanning 19 countries and four continents indicated that the CC398 lineage originated in humans as an MSSA [167]. The whole-genome sequencing analysis by Price et al. [167] demonstrated that livestock-associated MRSA CC398 lost an immune-evasion cluster (IEC) as it evolved from its human-adapted MSSA. All of the HA-MSSA strains carry ϕ Sa3 prophage in association with human innate immunomodulatory genes that play crucial roles in human niche adaptation (**Figure 2**) [167]. The prophage-associated virulence and adaptation genes are not necessary for nonhuman hosts, therefore, ϕ Sa3 is mostly absent in livestock strains. After their introduction to livestock, MSSA CC398 acquired resistance to methicillin and tetracycline. Since tetracycline is heavily used in animal farming, the tetracycline resistance gene *tet(M)* is nearly universal among livestock-associated MRSA CC398 and MSSA isolates. The MSSA and MRSA CC398 isolates found in humans with direct livestock contact exhibited the same molecular patterns (i.e., ϕ Sa3 prophage negative, *tet(M)*-positive) as the livestock-associated strains, indicating human re-adaptation [167], and were also reported in isolates epidemiologically associated with human-to-human transmission in multiple countries and continents [168]. During the host jump from humans to animals, MRSA CC398 strains also acquired resistance to copper and zinc because of their use in animal feed [130]. The vast majority of LA-MRSA CC398 strains carry *SCCmec* type Vc, which contains the *czrC* gene that confers resistance to copper and zinc.




Figure 2. Gain or loss of genes as *S. aureus* jumps from human to livestock animals.

It is quite possible that LA-MRSA CC398 strains would eventually acquire certain genetic traits (additional antibiotic resistance and virulence factors) that would allow *S. aureus* to colonize both hosts and become a more formidable zoonotic agent [167]. Since ST398 strains are deficient in one or more restriction modification systems [169, 170], this adaptation process may have already occurred, as *pvl*-positive ST398 MRSA strains have been isolated from

severe cases of community-acquired infections [20, 21]. These *pvl*-positive ST398 MRSA strains are also lysogenized with ϕ Sa3 phage coding for the human-specific virulence factors *sak*, *scn*, and *chp*. LA-MRSA *S. aureus* CC398 is not alone in the ability to adapt. A phylogenetic study of *S. aureus* CC97 strains that originated from livestock and caused human infections in Asia, Europe, and North and South America indicated that in the process of adapting to the human host, CC97 acquired either the SCCmec IV or V cassette and the ϕ Sa3 containing an immune evasion cluster [123]. A more recent study in the United States demonstrated that 22% of 30 veterinary students who were initially MRSA negative became positive after visiting MRSA-positive pig farms in Iowa but were negative again by 24 h after the visit [75]. The predominant *spa* type most commonly detected among the students was associated with ST5, suggesting a possible expansion of LA-MRSA to include ST5 [75].

Bayesian phylogenetic analysis of the poultry isolates with the CC5 clade indicated that it arose due to a single human-to-poultry host jump in or near Poland, where CC5 poultry strains acquired an MGE (mobile genetic element), presumably from other resident staphylococcal strains [36]. In addition to gene acquisition, loss of staphylococcal protein A (SpA) has occurred in avian isolates; it encodes virulence factors involved in human disease pathogenesis but is not needed for avian pathogenesis. The lack of protein A expression is a characteristic of the poultry biotype as defined by Devriese et al. [148]. In addition to the CC5 poultry clade, several other poultry isolates had identical or closely related STs to strains commonly associated with humans but had acquired MGE unique to avian strains, indicating that human-to-poultry host switches may be happening relatively frequently. A recent study demonstrated frequent contamination of poultry meat products with *S. aureus* ST5 and ST398 isolates, both of which have human origin [171]. Host adaptation of ST5 from humans to chickens was associated with a loss of genes contributing to human pathogenesis, and this was followed by the acquisition of avian-specific virulence determinants [36, 172]. Further, the presence of a cysteine protease encoded by a plasmid and widely distributed within avian strains suggests its potential role in avian-specific pathogenesis [173]. The adaptation from one host to another appears to be dependent upon the acquisition or loss of MGEs that code for key elements necessary for survival in new host [167, 174]. These cases have not only been associated with livestock [19] but have also occurred in cases with no known livestock contact [53, 54], suggesting a broader transmissibility capacity than originally thought. Further research is required to characterize the full scope of the genetic changes associated with the shift from humans to livestock and vice versa.

5. ST398 evolution and genetic diversity

In spite of similarities between LA- and HA-MRSA isolates, significant amounts of genetic diversity among *spa* and SCCmec cassette types have been documented in ST398 [6, 57, 175]. For instance, ST398 appears to have evolved by multiple acquisitions of the SCCmec elements, such as SCCmec types II, III, IV, IVa, and V [176]. In the Netherlands, two farms were found to have MRSA ST398 with identical *spa* types, but different SCCmec types, suggesting that divergent SCCmec elements were inserted into the clonal MSSA [57]. Similarly, MSSA ST398 (*spa* type t899), MRSA ST398–IVa (*spa* type t899), and MRSA ST398–V (*spa* type t108)

were found in dust samples, nasal swabs, and a blood isolate from workers on the same pig farm [177], suggesting multiple acquisitions of SCC_{mec} cassettes by MSSA precursors. Coagulase-negative staphylococci in the farming environment are suspected as sources of SCC_{mec} [176], and the progeny of emerging MRSA strains are spreading locally rather than globally [178–180]. While SCC_{mec} acquisition seems to be fairly common in MRSA ST398, the transfer of staphylococcal toxin genes, including the Pantone-Valentine leukocidin gene (*pvl*) appears to be rarer [20, 43, 44, 57, 165, 181–184]. Only a handful of studies have found *pvl* positive ST398 [20, 165, 185–187]. Additionally, horizontal transfer of the protein A gene has been suggested, due to the finding of the *spa* type t899 in both ST398 strains and ST9 strains [29, 30, 80, 177].

6. MRSA in companion animals

6.1. Canine and feline MRSA

The first MRSA from pet animals was isolated from dogs in Nigeria in 1972 [188]. *S. intermedius* is the strain most isolated from dogs [189]. However, the predominant canine species among staphylococci was *S. sciuri* in Japan [190]. Various coagulase-positive and -negative staphylococci have been reported in pet animals [191]. Among coagulase-negative staphylococci, *S. felis* was dominant in Brazil [116], and a coagulase-positive *S. intermedius* was dominant in canine species in the UK [192, 193]. The prevalence of canine MRSA was 0.7% in Portugal, 2.3–9% in UK, and ≤20% in Canada [194–197]. Conversely, in cats, it was 1.48% in the UK, ≤4% in Portugal [195, 198] and 21.4% from wounds and skin lesions of cats in the USA [199]. *S. aureus* was found in 8% of dogs with inflammatory skin disease in the USA and one isolate was MRSA [200]. In addition, one MRSA was detected among the *S. aureus* strains isolated from 29% of 48 cats suffering with inflammatory skin disease and two MRSA were found among the *S. aureus* strains from 20% of 50 healthy cats [201]. Previous surgery, hospitalization, antimicrobial agent treatment, contact with humans possessing MRSA, and use of implant devices are regarded as risk factors for MRSA infection in companion animals [195, 202]. Rich and Robert [203] reported that MRSA were isolated from 1.4% of the postoperative and wound infection samples of pet animals in the UK. Lilenbaum et al. found 3% MRSA in Brazilian cats [193]. Southwest Pacific clone-associated community-acquired MRSA (USA1100) and methicillin-resistant *S. pseudintermedius* (MRSP)-associated with the European clone (ST71) were first reported in South America in cats [204].

MRSA isolates from Austria, Belgium, Germany, Ireland, and Portugal were resistant to ciprofloxacin and enrofloxacin, perhaps because of the fluoroquinolone approval for use in companion animals in Europe in the middle of 1990 (Table 2). MRSA ST398 that was identified in dogs and cats in France carried a chloramphenicol acetyltransferase gene, *cat*, that mediates resistance to nonfluorinated phenicols [117]. On the other hand, MRSA isolated in Portugal and Thailand from dogs and Germany from cats possessed a florfenicol-chloramphenicol exporter gene, *fexA*, that mediates resistance to both fluorinated and nonfluorinated phenicols [205–207]. MRSA ST398 isolated from an Austrian dog suffering from vaginitis harbored the *vatC* gene that inactivates streptogramin A [208]. An ABC transporter gene, *lsaE* (responsible

for combined resistance to lincosamides, pleuromutilins, and streptogramin A), and a lincosamide nucleotidyltransferase *lunB* gene (confers resistance only to lincosamides), were present in the MRSA ST45 isolated from dogs in Thailand [205]. In France, an MRSA strain (*agr* III-t008–IV) isolated from a synovial fluid of dog was positive for *pvl* [117]. It carried *sek* and *seq* genes and was suspected to be a USA300 variant that was imported from the USA. *tsst1*-positive MRSA (*agr* II-t002-SCC*mec* I-truncated) was recovered from pet animals in France (Table 3). Leukotoxin genes, most prevalently *lukF* and/or *lukS* followed by *lukD* and/or *lukE*, are reported to be present in most of the MRSA isolated from companion animals, but those from dogs, cats and horses in Austria and the USA did not possess any leukotoxin genes. MRSA isolates CC5-t002–II, CC5-t062 and CC22-t032–IV, isolated in Portugal from a dog, a horse and human, respectively, harbored IEC genes suggesting adaptation to different hosts [206]. The presence of host-adaptive, virulence, and toxin genes makes the companion animal MRSA isolates prime candidates for zoonotic transfer. The first outbreak of human MRSA associated with cats was reported in a rehabilitation geriatric ward in the UK in 1988 [209] but others have since been reported in Australia, Canada, Germany, New Zealand, South America, and the Netherlands [193, 210–212]. Human-associated MRSA, such as EMRSA-15 and CMRSA-2, show a close relationship with pet-associated MRSA [211–214]. Since MRSA ST398 has been isolated from many farm animals, this MLST type found in dogs or cats may have originated from them or people who had contact with them [215]. MRSA SCC*mec* II strains, which are those most frequently affiliated with nosocomial human infections, have been isolated from cats [216].

6.2. Equine MRSA

Since the first report of MRSA from mares with metritis in Japan [217], many isolates have been found in Europe, North America, and Asia (Table 2) [142–144, 218–220]). Haenni et al. [221] identified four *mecC*-positive MRSA from horses in France exhibiting *spa* types, t208, t843, t6220, t11015 and ST types ST49, ST130, and ST1245. Among them, MRSA CC130 (ST130, ST1245) and CC49 (ST49) were documented in animals and humans, respectively. Multidrug-resistant ST8 MRSA was detected in Belgium, Germany, Switzerland, and the USA. A single locus variant of ST8 classified as ST254 MRSA was isolated from horses in Austria, Germany, Ireland, and the UK [183, 208, 222, 223]. Twelve different MLST types have been reported and most of the MRSA strains were grouped into the CC8 or CC398 classes [224]. The CC8-SCC*mec* IV genotype in horses was likely from a contaminated veterinary hospital and later spread to various clinics [219]. This genotype was reported in veterinary hospitals in Canada and the USA [118, 225]. It was first found in infected horses in Ireland and thereafter reported in the Netherlands, Austria, and Germany [226–228]. Many MRSA isolates from Canada showed ST8-SCC*mec* IV-t064 genotype [93]. They were designated as a Canadian epidemic MRSA-5 (CMRSA-5) and were very close to a human clone, USA500. Although the CC8-SCC*mec* IV genotype has been the most frequently found, CC398-SCC*mec* IV has recently become a major genotype. CC398-SCC*mec* IV was first found from infected horses in the Veterinary University of Vienna in Austria [222]. CC398 isolated from nasal samples of horses in the Netherlands and Belgium exhibited high prevalence rates, 9.3 and 10.9%, respectively [227, 229]. More than 25 *spa*-types have been reported, and three types, such as t011, t064 and t451, were the most

Country	Animal	ST	AMP	CEF	CHL	CIP	CLI	ENR	ERY	FLO	FUS	GEN	KAN	LIN	LIZ	QD	SPE	STR	TET	TIA	TOB	TRI	SXT	Genotype	References
Belgium	B [†]	398	*			+	**		+			+	+	+				+	+		+	+			[87]
Belgium	B	398			+	+			+			+	+	+					+		+	+			[133]
Belgium	C [†]	239, 398		+	+			+		+	+	+	+			+		+	+	+		+			[34]
Belgium	C	398				+			+					+					+			+			[133]
Belgium	P [†]	ST9, 80, 239, 398			+	+	+		+		+	+	+		+	+		+	+	+		+	+	<i>aacA-aphD, aadD, aphA3-sat, blaI, blaR, blaZ, cfr, dfrS1, ermB, ermC, fexA, fosB, lnuA, tetK, tetM, vgaA</i>	[121]
Belgium	P	398				+			+			+	+	+					+		+	+			[256]
Belgium	P	398			+	+			+		+	+	+	+					+		+	+			[66]
Belgium	P	398			+	+			+			+	+	+		+			+		+	+			[133]
Brazil	B	398		+		+	+					+						+	+		+			<i>ant4, aac(6')-aph(2''), blaZ, mecA, tetK, tetM</i>	[257]
Central Europe	P	9			+		+		+										+			+			[258]

Country	Animal	ST	AMP	CEF	CHL	CIP	CLI	ENR	ERY	FLO	FUS	GEN	KAN	LIN	LIZ	QD	SPE	STR	TET	TIA	TOB	TRI	SXT	Genotype	References
China	P	6, 9, 63, + 627	+		+		+		+	+						+	+		+	+				<i>aadE</i> , <i>erm33</i> , <i>fexA</i> , <i>lnuB</i> , <i>lsaE</i> , <i>mecA</i> , <i>spw</i> , <i>tetL</i> , <i>vgaA</i>	[259]
China	P	5, 9			+	+	+		+		+								+				+		[260]
China	P	NA	+		+	+	+		+		+								+				+	<i>acc(6')</i> / <i>aph2''</i> , <i>ant(4', 4'')</i> , <i>aph(3')</i> - <i>III</i> , <i>ermC</i> , <i>mrsA</i> , <i>tetK</i> , <i>tetM</i>	[261]
China	P	398	+		+	+	+		+		+					+			+				+		[262]
Finland	P	1, 398	+			+	+		+										+					<i>aadD</i> , <i>blaZ/I/R</i> , <i>ermB</i> , <i>lnuB</i> , <i>mecA</i> , <i>tetK</i> , <i>tetK/M</i>	[120]
Germany	B	398	+		+		+		+		+	+	+	+		+	+		+	+		+	+	<i>aacA</i> - <i>aphD</i> , <i>aadD</i> , <i>aphA3</i> , <i>blaZ-I-R</i> , <i>dfrK</i> , <i>ermA</i> , <i>ermB</i> , <i>ermC</i> , <i>ermT</i> , <i>fex</i> , <i>spc</i> , <i>tetL</i> , <i>tetK</i> , <i>tetM</i> , <i>vgaA</i> , <i>vgaC</i>	[97]

Country	Animal	ST	AMP	CEF	CHL	CIP	CLI	ENR	ERY	FLO	FUS	GEN	KAN	LIN	LIZ	QD	SPE	STR	TET	TIA	TOB	TRI	SXT	Genotype	References
Germany	B	398	+					+	+										+						[102]
Germany	P	398	+	+	+		+	+	+	+		+	+				+		+	+			+	<i>aacA-aphD, aadD, blaZ, dfrG, dfrK, ermA, ermB, ermC, fexA, mecA, qacC, tetK, tetL, tetM, vga vgaA</i>	[44]
Hong Kong	C	9	+	+	+		+	+	+	+		+	+	+		+	+	+	+	+		+	+	<i>aacA-aphD, aadD, aadE, blaZ/I/R, dfrK, ermC, lnuB, lsaE, mecA, spw, tetL</i>	[134]
Hong Kong	P	9			+	+	+		+		+								+					<i>ermB, ermC</i>	[81]
Hong Kong	P	9	+	+			+	+			+	+					+	+	+	+		+		<i>aacA-aphD, aadD, aadE, blaZ/I/R, dfrK, ermC, lnuB, lsaE, mecA, tetL, spw</i>	[134]

Country	Animal	ST	AMP	CEF	CHL	CIP	CLI	ENR	ERY	FLO	FUS	GEN	KAN	LIN	LIZ	QD	SPE	STR	TET	TIA	TOB	TRI	SXT	Genotype	References
Hong Kong	P	9			+	+	+		+		+					+			+			+		<i>ermC</i> , <i>fexA</i> , <i>tetK</i> , <i>tetM</i>	[263]
Hungary	B	1	+						+										+						[92]
Ireland	P	CC398	+		+				+			+	+	+			+		+		+	+		<i>aacA</i> - <i>aphD</i> , <i>aadD</i> , <i>blaZ</i> , <i>dfrG</i> , <i>dfrK</i> , <i>ermA</i> , <i>ermB</i> , <i>fexA</i> , <i>spc</i> , <i>tetK</i> , <i>tetL</i> , <i>tetM</i>	[264]
Italy	B	CC97		+		+	+		+			+	+			+		+	+	+		+		<i>aacA</i> - <i>aphD</i> , <i>aadD</i> , <i>blaI</i> , <i>blaR</i> , <i>blaZ</i> , <i>cat</i> , <i>ermB</i> , <i>ermC</i> , <i>mecA</i> , <i>qacC</i> , <i>sdrM</i> , <i>tetK</i> , <i>tetM</i> , <i>vgaA</i>	[110]
Italy	B	1		+		+			+			+	+					+	+	+		+		<i>aphA3</i> , <i>blaZ</i> , <i>ermC</i> , <i>mecA</i> , <i>sat</i> , <i>tetK</i> , <i>sdrM</i>	[119]
Italy	P	1, 9, 97, 398, 1476			+	+	+		+			+	+					+	+			+			[62]

Country	Animal	ST	AMP	CEF	CHL	CIP	CLI	ENR	ERY	FLO	FUS	GEN	KAN	LIN	LIZ	QD	SPE	STR	TET	TIA	TOB	TRI	SXT	Genotype	References
Italy	P	CC97	+		+	+		+			+	+				+		+	+	+		+		<i>aacA-aphD, aadD, blaI, blaR, blaZ, cat, ermB, ermC, mecA, qacC, sdrM, tetK, tetM, vgaA</i>	[110]
Italy	P	1	+		+	+		+			+	+						+	+	+		+		<i>aacA-aphD, aadD, blaI, blaR, blaZ, cfr, dfrS1, ermA, ermC, ermR, fexA, mecA, qacC, sdrM, tetK, tetM, vgaA</i>	[119]
Italy & Spain	P	1	+		+	+		+			+	+							+	+		+		<i>aacA-aphD, aadD, blaZ, cat, dfrA, ermA, ermC, lnuA, mecA, tetK, tetL, tetM, vgaA, vgbA</i>	[265]

Country	Animal	ST	AMP	CEF	CHL	CIP	CLI	ENR	ERY	FLO	FUS	GEN	KAN	LIN	LIZ	QD	SPE	STR	TET	TIA	TOB	TRI	SXT	Genotype	References
Korea	B	NA	+	+		+	+		+			+	+						+						[89]
Korea	B	1, 72							+			+							+						[108]
Korea	B	NA	+				+		+			+	+						+						[90]
Korea	C	692				+	+		+										+					<i>ermB</i> , <i>ermC</i> , <i>tetL</i> , <i>tetS</i>	[91]
Korea	C	NA	+	+			+		+																[89]
Korea	P	398				+	+		+										+						[85]
Korea	P	5, 72, 398, 541			+	+	+		+		+	+	+			+	+		+	+				<i>aac(6')-le-</i> <i>aph(2'')</i> , <i>ant(3')-</i> <i>IIIa</i> , <i>anat(4')-la</i> , <i>cfr</i> , <i>ermA</i> , <i>ermC</i> , <i>ermT</i> , <i>fexA</i> , <i>lnuB</i> , <i>tetK</i> , <i>tetL</i> , <i>tetM</i>	[91]
Netherland	B	398	+									+						+		+		+			[266]
Netherlands	P	398					+		+			+	+					+		+					[22]
Netherlands	P	398							+			+	+	+				+							[70]
Portugal	B	398			+	+	+	+	+	+			+					+						<i>aphA3</i> , <i>ermC</i> , <i>fexA</i> , <i>tetK</i> , <i>tetM</i>	[206]

Country	Animal	ST	AMP	CEF	CHL	CIP	CLI	ENR	ERY	FLO	FUS	GEN	KAN	LIN	LIZ	QD	SPE	STR	TET	TIA	TOB	TRI	SXT	Genotype	References
Portugal	P	NA				+	+		+			+	+						+	+		+	+	<i>aacA-aphD, aadD, apmA, dfrK, ermC, fexA, qacG, qacJ, tetK, tetM, vgaA</i>	[206]
Spain	P	398, 1379					+		+			+	+					+	+		+	+		<i>aacA-aphD, aadA, aadD, aphA3, dfrA, dfrG, dfrK, ermC, ermT, msrA, str, tetK, tetM, tetL</i>	[255]
Spain	P	398					+		+			+									+		+		[234]
Spain	P	1, 398, 1965, 1966, 1967, 1968, 1969	+	+	+			+		+	+			+		+		+	+			+			[267]
Switzerland	B	1, 398	+	+		+		+											+					<i>aphA, blaI, blaR, blaZ, ermC, mecA, tetM</i>	[42]

Country	Animal	ST	AMP	CEF	CHL	CIP	CLI	ENR	ERY	FLO	FUS	GEN	KAN	LIN	LIZ	QD	SPE	STR	TET	TIA	TOB	TRI	SXT	Genotype	References	
Switzerland	P	398	+	+			+		+			+							+			+	<i>blaI, blaR, blaZ, ermA, mecA, tetK, tetM</i>	[42]		
Switzerland	P	398		+			+		+					+				+	+						[63]	
Taiwan	P	9			+	+	+		+			+							+			+			[268]	
Thailand	P	9		+	+		+					+							+			+			[31]	
Thailand	P	9, 2136		+		+						+					+		+	+		+	+	<i>blaZ, aac-aphD, tetM, vgaA</i>	[269]	
USA	P	398					+		+							+			+						[51]	
USA	P	5, 72, 398, 1340	+				+		+			+				+			+			+			[38]	
B ⁺ : cattle, C ⁺ : chicken, P ⁺ : pigs, *: not tested or sensitive, **: resistant, AMP: ampicillin, CEF: cefoxitin, CHL: chloramphenicol, CIP: ciprofloxacin, CLI: clindamycin, ENR: enrofloxacin, ERY: erythromycin, FLO: florfenicol, FUS: fusidate, GEN: gentamicin, KAN: kanamycin, LIN: lincomycin, LIZ: linezolid, QD: quinupristin/dalfopristin, SPE: spectinomycin, STR: streptomycin, TET: tetracycline, TIA: tiamulin, TOB: tobramycin, TRI: trimethoprim, SXT: trimethoprim-sulfamethoxazole.																										

Table 1. Antimicrobial resistance pattern of bovine, poultry, and porcine MRSA isolates.

Country	Animal	ST	AMP	CEF	CHL	CIP	CLI	ENR	ERY	FLO	FUS	GEN	KAN	LIN	LIZ	QD	SPE	STR	TET	TIA	TOB	TRI	SXT	Genotype	References
Austria	D [†]	22, 254, 398	*			+	**		+			+							+			+		<i>aacA-aphD</i> , [208] <i>dfrA</i> , <i>ermC</i> , <i>tetK</i> , <i>tetM</i> , <i>vatC</i>	
Austria	C [†]	1, 5, 398				+	+		+			+							+					<i>aacA-aphD</i> , [208] <i>ermC</i> , <i>tetK</i> , <i>tetM</i>	
Austria	H [†]	1, 254, 398				+	+		+			+							+			+		<i>aacA-aphD</i> , [208] <i>dfrA</i> , <i>ermC</i> , <i>tetK</i> , <i>tetM</i> , <i>vatC</i>	
Austria	H	1, 254, 398					+		+			+							+					<i>aph2''-aac6'</i> , <i>ermC</i> , <i>mecA</i> , <i>tetM</i>	[222]
Belgium	D	398			+	+			+			+	+	+					+		+	+			[133]
Belgium	C	398			+	+			+					+		+			+						[133]
Belgium	H	8, 398, 2197	+	+			+	+	+			+		+					+			+			[270]
Belgium	H	398					+		+			+		+					+			+			[229]
Brazil	C	30		+																					[204]
Brazil	C	NA	+																						[193]
Canada	D	NA	+			+	+	+	+		+	+							+			+			[271]
Canada & USA	H	NA			+			+	+			+							+			+			[272]
China	C, D	59, 398					+					+							+					<i>aacA-aphD</i> , [273] <i>ermB</i> , <i>mecA</i> , <i>linA</i> , <i>tetK</i>	

Country	Animal	ST	AMP	CEF	CHL	CIP	CLI	ENR	ERY	FLO	FUS	GEN	KAN	LIN	LIZ	QD	SPE	STR	TET	TIA	TOB	TRI	SXT	Genotype	References
Japan	D	5, 30				+			+			+	+					+	+						[279]
Japan	D	NA	+						+			+							+						[280]
Korea	D	72		+																					[281]
Korea	D	NA	+		+	+		+				+									+		+		[282]
Malaysia	D	59		+					+									+	+						[283]
Malaysia	C	55		+					+			+						+	+						[283]
Netherlands	D	NA	+					+	+			+		+											[284]
Nigeria	D	NA		+	+	+	+		+			+							+				+		[285]
Portugal	D	22, 105, 398			+	+	+		+	+			+						+					<i>fexA</i> , <i>ermA</i> , <i>ermC</i> , <i>tetM</i>	[206]
Portugal	C	5, 22				+	+		+		+													<i>ermC</i> , <i>fusC</i> , <i>mphC</i> , <i>msrA</i> ,	[206]
Portugal	D	22				+	+		+		+	+	+			+			+		+			<i>aac(6')-aph(2''), ant(4')-Ia, aph(3')-III, ermB, ermC, msrA, tetM</i>	[286]
Portugal	H	5, 398					+		+		+	+	+					+			+			<i>aacA-aphD, blaZ, dfrK, ermC, fusC, tetM</i>	[287]

[illegible]

Country	Animal	ST	AMP	CEF	CHL	CIP	CLI	ENR	ERY	FLO	FUS	GEN	KAN	LIN	LIZ	QD	SPE	STR	TET	TIA	TOB	TRI	SXT	Genotype	References
USA	C, D	NA			+		+	+	+			+							+				+		[292]
USA	D	NA				+	+		+										+						[293]
USA	C	NA				+	+		+										+						[293]
USA	C	NA			+		+	+	+			+							+				+		[216]
USA & UK	D	NA				+	+	+	+			+		+									+		[294]
USA	H	8, 830	+						+			+							+				+		[118]
C [†] : cats, D [†] : dogs, H [†] : horses, *: not tested or sensitive, **: resistant, AMP: ampicillin, CEF: cefoxitin, CHL: chloramphenicol, CIP: ciprofloxacin, CLI: clindamycin, ENR: enrofloxacin, ERY: erythromycin, FLO: florfenicol, FUS: fusidate, GEN: gentamicin, KAN: kanamycin, LIN: lincomycin, LIZ: linezolid, QD: quinupristin/dalfopristin, SPE: spectinomycin, STR: streptomycin, TET: tetracycline, TIA: tiamulin, TOB: tobramycin, TRI: trimethoprim, SXT: trimethoprim-sulfamethoxazole.																									

Table 2. Antimicrobial resistance pattern of feline, canine and equine MRSA isolates.

Country	Animals	Leukotoxins	Hemolysins	Enterotoxins	Immune-invasion factors	Protease	Superantigen-like proteins	Biofilm-associated	Capsule	Miscellaneous genes	References
Austria	Horses			<i>seb</i>							[208]
	Dogs			<i>sei</i>							
	Cats			<i>sei</i>							
Belgium	Pigs	<i>lukD/E, lukF, lukPV, lukS, lukX, lukY</i>	<i>hla, hlb, hld, hlyIII, hlgA</i>	<i>seg, sei, sem, sen, seo, seu</i>	<i>hysA1, hysA2, isaB, isdA</i>	<i>splA, splB</i>	<i>ssl1, ssl2, ssl3, ssl4, ssl5, ssl6, ssl7, ssl8, ssl9, ssl10, ssl11</i>	<i>bbp, cflA, cflB, cna, ebh, ebpS, eno, fib, fnbA, fnbB, icaACD, map, mprF, sasG, sdrC, sdrD, vwb</i>	<i>capH5, capH8, capI8, capJ5, capJ8, capK5, capK8</i>	<i>edinB, etd, setC</i>	[121]
China	Pigs			<i>sec, seg, sei, sem, sen, seo, seh</i>							[260]
Finland	Pigs			<i>seh</i>				<i>fnbB</i>	<i>cap5, cap8</i>		[120]
France	Cats and dogs	<i>lukF-PV, lukS-PV</i>		<i>sea, sec, sed, seg, sei, sej, sek, sel, sem, sen, seo, seq, ser, seu</i>						<i>tst</i>	[117]
Germany	Pigs			<i>seb, sek, seq</i>							[44]
	Horses	<i>lukD/E, lukF, lukS</i>	<i>hl, hla, hlb, hld, hlyIII-aII, hlgA</i>	<i>sea, seb/k/g, sec/l, sed/j/r, seg/i/m/n/o/u, sep</i>	<i>chp, sak, scn</i>	<i>splAB/E, splE, sspA/B/P</i>					[183]

Country	Animals	Leukotoxins	Hemolysins	Enterotoxins	Immune-invasion factors	Protease	Superantigen-like proteins	Biofilm-associated	Capsule	Miscellaneous genes	References
Italy	Cattle	<i>lukD, lukE, lukF, lukS, lukX, lukY</i>	<i>hl, hIII, hIb, hlgA</i>	<i>seh</i>	<i>hysA1, hysA2, isaB, isdA, sak, scn</i>	<i>aur, splA, splB, sspA, sspB, sspP</i>	<i>ssl1, ssl2, ssl3, ssl4, ssl5, ssl6, ssl7, ssl8, ssl9, ssl10, ssl11</i>	<i>bbp, clfA, clfB, cna, ebh, ebpS, icaA, icaC, icaD, eno, fib, fnbA, fnbB, sasG, sdrC, sdrD, vwb</i>	<i>cap8, capH8, capI8, capJ8, capK8</i>	<i>lmrP, mprF, hsdS2, hsdS3, hsdSx</i>	[119]
	Pigs	<i>lukD, lukE, lukF, lukS, lukX, lukY</i>	<i>hl, hIa, hIII, hIb, hlgA</i>	<i>seh</i>	<i>isaB, isdA, hysA1, hysA2</i>	<i>aur, splA, splB, splE, sspA, sspB, sspP</i>	<i>ssl1, ssl2, ssl3, ssl4, ssl5, ssl6, ssl7, ssl8, ssl9, ssl10, ssl11</i>	<i>bbp, clfA, clfB, cna, ebh, ebpS, icaA, icaC, icaD, eno, fib, fnbA, fnbB, sasG, sdrC, sdrD, vwb</i>	<i>cap8, capH8, capI8, capJ8, capK8</i>	<i>lmrP, mprF, hsdS2, hsdS3, hsdSx</i>	
Italy & Spain	Pigs	<i>lukX</i>	<i>hIb</i>	<i>entH, entU, entX, entY</i>							[265]
Korea	Pigs	<i>lukE/D</i>		<i>sed, seg, sei, sej, sem, sen, seo, sep, ser</i>	<i>chp, sak, scn</i>					<i>tsst1</i>	[91]
	Chicken	<i>lukE/D</i>		–	–						
	Cattle	<i>lukE/D, pvl</i>		<i>seb, sed, seg, sei, sej, sek, sem, sen, seo, seq, ser</i>	<i>chp, sak, scn</i>					<i>tsst1</i>	

Country	Animals	Leukotoxins	Hemolysins	Enterotoxins	Immune-invasion factors	Protease	Superantigen-like proteins	Biofilm-associated	Capsule	Miscellaneous genes	References
Portugal	Pigs	<i>lukF, lukS</i>	<i>hla, hld</i>	–	–						[206]
	Dog	<i>lukD, lukE, lukF, lukS,</i>	<i>hla, hlb, hld</i>	<i>sec, sed, seg, sei, sej, sel, sem, sen, seo, seu, ser</i>	<i>chp, sak, scn</i>						
	Horse	<i>lukF, lukS, lukD, lukE</i>	<i>hla, hlb, hld</i>	<i>seg, sei, sem, sen, seo, seu</i>	<i>chp, sak, scn</i>						
	Calf	<i>lukF, lukS</i>	<i>hla, hld</i>	–	–						
Spain	Pigs	<i>lukD, lukE</i>	<i>hla, hlb, hld, hlg, hlg-v</i>							<i>eta</i>	[255]
Switzerland	Pigs	<i>lukF, lukS, lukY</i>	<i>hla, hld, hlgA</i>	<i>entX, entY</i>							[42]
	Calf	<i>lukF, lukS, lukY</i>	<i>hla, hld, hlgA</i>	<i>entX</i>							
	Cattle	<i>lukD, lukE, lukF, lukY</i>	<i>hla, hld, hlgA</i>	<i>entH, entX, entY</i>							
Thailand	Pigs			<i>entG, entI, entM, entN, entO</i>							[269]
USA	Dogs	<i>lukSF-PV</i>		<i>sea, seb, sec, sed, seg, sei, sej, sem, sen</i>				<i>clfA, clfB, fnbA, fnbB</i>			[118]
	Cats	<i>lukSF-PV</i>		<i>sec, sed, seg, sei, sej, sek, sem, sen</i>				<i>clfA, clfB, fnbA, fnbB</i>			
	Horses			<i>sea, seb, seg, sek, sei, sej, sek</i>				<i>clfA, clfB, fnbA, fnbB</i>			

Table 3. Virulence profiles from food-producing and pet animals.

widespread [230]. In addition, only three SCC_{mec} types (IV, V, VI) have been discovered from horses [218]. Interestingly, no MRSA was isolated from 300 horses on 14 farms in Slovenia, 497 horses on 50 farms in Canada, 87 horses in Austria, and 200 horses in the Netherlands [231–233]. ST22 and ST1117 isolates from horses in Germany had IEC genes, such as *chp*, *sak*, and *scn* [183]. On the other hand, ST8, ST254, and ST398 strains did not carry those genes.

6.3. Antibiotic resistance and enterotoxin genes in LA-MRSA CC398

An analysis of MRSA and MSSA from animals and humans spanning 19 countries and four continents indicated that the CC398 lineage originated in humans as MSSA [167]. After its transmission to livestock, CC398 became resistant to tetracycline, probably because of the heavy tetracycline use in pig production [22]. However, many tetracycline-resistant MRSA strains are found in horses despite the fact that tetracycline is either not used much [229] or sparingly used [93, 229]. Among bovine MRSA isolates tested, most of them were resistant to β -lactam antibiotics [34] as well as tetracycline, erythromycin and gentamicin. CC398 has also been reported to be highly resistant to several other antibiotics, such as ciprofloxacin, tobramycin, clindamycin, and trimethoprim-sulfamethoxazole [234]. Antimicrobial resistance patterns of MRSA and MSSA isolates in Hong Kong were very similar [81]. The only MRSA CC398 isolate that has exhibited resistance against daptomycin and intermediate susceptibility to vancomycin has been described in a case-report from an Italian hospital [235]. Two other isolates, one from a ventilator-associated pneumonia of a farmer and one additional porcine isolate, were also described as resistant to linezolid and possessed the *cfr* gene which is located on transferable plasmids [235]. An outbreak with HA-MRSA ST125 containing *cfr*, reported from a Madrid hospital in 2010 [236], was followed by reports on the emergence of nosocomial coagulase negative staphylococci containing *cfr* [40]. This gene was previously found in staphylococci from animals in Europe [237, 238] but has also been reported in humans in Colombia and the USA [239, 240]. Distribution of the *cfr* gene in MRSA isolates is alarming because, besides linezolid, it confers resistance to oxazolidinones, phenicols, lincosamides, streptogramin A, and pleuromutilins, including the topical antibiotic retapamulin that is used for treatment of human skin infections [239, 241, 242].

About 50% of LA-MRSA CC398 isolates, besides being resistant to antimicrobial agents, also exhibit resistance to copper and zinc mediated by the *czrC* gene [130, 243]. The use of zinc as feed additives may have favored spread of the *czrC* gene in LA-MRSA [130]. So far, this gene has only been found in LA-MRSA [130, 243] but an extended use of copper coating of biomaterials in orthopedic surgery and traumatology, as well as copper surfaces in hospitals, might select for *czrC*-positive HA-MRSA as well. The trimethoprim-resistance gene *dfrK*, located close to *tetL* in LA-MRSA ST398, has also been found on a plasmid [244]. The *tetL* gene was identified in MRSA isolated from diverse livestock animals and meats from different regions of the world [97, 245, 246]. Kadlec and Schwarz demonstrated the presence of plasmid pKKS25-associated resistance genes, *ermT*, *dfrK*, and *tetL*, in MRSA obtained from a nasal swab of a young sow in Germany [247]. A porcine MRSA ST398 was shown to contain a transposon Tn6133 carrying *ant(9)-Ia* and *ermA* genes and a plasmid pKKS825 [248, 249] harboring the resistance genes *vgaC* and *vgaE*, *aadD*, *tetL*, and *dfrK* [248–250]. In Germany, the *vgaE* gene was detected in MRSA ST398 isolated from cattle, turkeys, and chicken and turkey meats

[251]. An apramycin resistance gene, *apmA*, was discovered in a bovine MRSA ST398 [252] and in porcine MRSA ST398 [206, 253]. In Denmark, the quaternary ammonium compound-resistant genes, *qacC* and *qacG* were detected in MRSA CC30 isolates [254]. Wendlandt et al. identified a plasmid pV7037-associated multidrug resistance gene cluster, including the novel resistance genes *lsaE* and *spw*, and other resistance genes, such as *mecA*, *blaZ/I/R*, *tetL*, *dfrK*, *ermC*, and *aadD*, from frozen or chilled chicken carcasses in Hong Kong. The antimicrobial resistance patterns and associated genes from bovine, porcine, and poultry isolates are summarized in **Table 1** and those from companion animals are shown in **Table 2**.

Most of the animal isolates are negative for the *pvl* gene, but Belgian MRSA ST80–IV isolates from healthy pigs were positive for the *pvl* gene and corresponded to the community-acquired CA-MRSA ST80–IV European clone [121]. One Belgian pig CC80 strain contained an exfoliatin (*etd*), an epidermal cell differentiation inhibitor (*edinB*), and staphylococcal exotoxin-like protein (*setC*) and IEC genes (*isaB*, *isdA*, *hysA1*, *hysA2*) [121]. A comparison of the *pvl*-positive MRSA isolates (ST8-t008–IVa) from American pig and pet animals indicated the presence of common virulence profiles (*lukSF-PV*, *clfA*, *clfB*, *fibA*, and *sek*) except for the *fibB* gene of a canine isolate [118]. Spanish ST1379/CC97 porcine isolates carried an exfoliatin (*eta*), a leukotoxin (*lukE/D*), and a gamma-hemolysin (*hlg-2*) but were negative for *etb*, *etc*, *tst*, *pvl*, and enterotoxins [255]. The occurrence and prevalence of enterotoxin genes in MRSA isolates from food-producing or companion animals is summarized in **Table 3**. The presence of multidrug-resistant MRSA in companion and food animals, combined with the enterotoxin genes, will require constant monitoring and evaluation of mitigation strategies.

7. Conclusions

MRSA contamination in food-producing and companion animals poses a serious threat to public health. Incidences of identical LA-MRSA strains in pig farms and persons in close contact with food producing and companion animals suggest a clear link for transmission of these strains between humans and animals. While MRSA isolates from companion and food-producing animals are known to infect humans, the reverse is also true. Studies reviewed in this report indicate an initial transfer of MSSA from humans to animals by deletion of immunomodulatory genes and prophage ØSa3, necessary for human infection but not required for infection in animals, and acquisition of tetracycline and methicillin resistance genes (**Figure 2**). The MRSA that evolved in animals started showing up in humans that were in close contact with them and exhibited traits specifically found in animal isolates, indicating a reverse transmission from animals to humans. Initial reports of MRSA in animals did not indicate the presence of host adaptation, enterotoxin, virulence, and antimicrobial resistance genes in them but they are becoming more prevalent and it is feared that these animals could serve as a reservoir for such strains and play an important role in zoonotic transfers.

Documentation of MRSA isolates (ST59-t437–V) from cattle containing the *pvl* gene and other virulence factors, such as *chp*, *scn*, *seb*, *sek*, and *seq*, toxic shock syndrome toxin 1 (*tsst1* or *tst*) gene and hemolysin, protease, superantigen-like protein, capsule, and biofilm-associated genes make them powerful pathogens that could cause a medical nightmare.

A livestock-associated CC398 lineage MRSA is well known to transfer from animals to humans and other MRSA isolates of different clonal complex groups are also known to be associated with zoonotic transfers. A human pandemic community-associated CC97 lineage MRSA harboring the antimicrobial resistance genes *mecA* and *mecC* has been shown to have originated from animals.

A comprehensive study of the emergence, dissemination, prevention and control of MRSA colonization is required to mitigate the risks to both animal and human health. Rapid advancement of whole genome sequencing technology has the great power of discriminating closely associated MRSA isolates from different sources and could be used for source tracking and differentiating between animal and human origin isolates. In addition, it can be applied to monitor the emergence and dissemination of MRSA isolated from various environments and determine the characteristics of virulence factors and evolution of multi-antimicrobial resistance.

Disclaimer

The views expressed herein do not necessarily reflect those of the US Food and Drug Administration or the US Department of Health and Human Services.

Author details

Jungwhan Chon, Kidon Sung* and Saeed Khan

*Address all correspondence to: kidon.sung@fda.hhs.gov

Division of Microbiology, National Center for Toxicological Research, US Food and Drug Administration, Jefferson, AR, USA

References

- [1] Hennekinne JA, De Buyser ML, Dragacci S. *Staphylococcus aureus* and its food poisoning toxins: characterization and outbreak investigation. FEMS Microbiol Rev. 2012;36(4):815–36.
- [2] Weese JS. Methicillin-resistant *Staphylococcus aureus* in animals. ILAR J. 2010;51(3):233–44.
- [3] Stryjewski ME, Corey GR. Methicillin-resistant *Staphylococcus aureus*: an evolving pathogen. Clin Infect Dis. 2014;58 Suppl 1:S10–9.
- [4] Waness A. Revisiting methicillin-resistant *Staphylococcus aureus* infections. J Glob Infect Dis. 2010;2(1):49–56.
- [5] Taylor AR. Methicillin-resistant *Staphylococcus aureus* infections. Prim Care. 2013;40(3):637–54.

- [6] Pan A, Battisti A, Zoncada A, Bernieri F, Boldini M, Franco A, et al. Community-acquired methicillin-resistant *Staphylococcus aureus* ST398 infection, Italy. *Emerg Infect Dis*. 2009;15(5):845–7.
- [7] McNamee PT, Smyth JA. Bacterial chondronecrosis with osteomyelitis ('femoral head necrosis') of broiler chickens: a review. *Avian Pathol*. 2000;29(5):477–95.
- [8] Eriksen KR. ["Celbenin"-resistant staphylococci]. *Ugeskr Laeger*. 1961;123:384–6.
- [9] Hiramatsu K, Katayama Y, Matsuo M, Sasaki T, Morimoto Y, Sekiguchi A, et al. Multi-drug-resistant *Staphylococcus aureus* and future chemotherapy. *J Infect Chemother*. 2014;20(10):593–601.
- [10] Weber JT. Community-associated methicillin-resistant *Staphylococcus aureus*. *Clin Infect Dis*. 2005;41 Suppl 4:S269–72.
- [11] Vanderhaeghen W, Hermans K, Haesebrouck F, Butaye P. Methicillin-resistant *Staphylococcus aureus* (MRSA) in food production animals. *Epidemiol Infect*. 2010;138(5):606–25.
- [12] Pantosti A. Methicillin-resistant *Staphylococcus aureus* associated with animals and its relevance to human health. *Front Microbiol*. 2012;3:127.
- [13] Bal AM, Coombs GW, Holden MT, Lindsay JA, Nimmo GR, Tattevin P, et al. Genomic insights into the emergence and spread of international clones of healthcare-, community- and livestock-associated methicillin-resistant *Staphylococcus aureus*: blurring of the traditional definitions. *J Glob Antimicrob Resist*. 2016;6:95–101.
- [14] Fitzgerald JR. Livestock-associated *Staphylococcus aureus*: origin, evolution and public health threat. *Trends Microbiol*. 2012;20(4):192–8.
- [15] Fluit AC. Livestock-associated *Staphylococcus aureus*. *Clin Microbiol Infect*. 2012;18(8):735–44.
- [16] Wendlandt S, Kadlec K, Fessler AT, Monecke S, Ehrlich R, van de Giessen AW, et al. Resistance phenotypes and genotypes of methicillin-resistant *Staphylococcus aureus* isolates from broiler chickens at slaughter and abattoir workers. *J Antimicrob Chemother*. 2013;68(11):2458–63.
- [17] Armand-Lefevre L, Ruimy R, Andremon A. Clonal comparison of *Staphylococcus aureus* isolates from healthy pig farmers, human controls, and pigs. *Emerg Infect Dis*. 2005;11(5):711–4.
- [18] Nemati M, Hermans K, Lipinska U, Denis O, Deplano A, Struelens M, et al. Antimicrobial resistance of old and recent *Staphylococcus aureus* isolates from poultry: first detection of livestock-associated methicillin-resistant strain ST398. *Antimicrob Agents Chemother*. 2008;52(10):3817–9.
- [19] Graveland H, Wagenaar JA, Heesterbeek H, Mevius D, van Duinkerken E, Heederik D. Methicillin resistant *Staphylococcus aureus* ST398 in veal calf farming: human MRSA carriage related with animal antimicrobial usage and farm hygiene. *PLoS One*. 2010;5(6):e10990.

- [20] Stegger M, Lindsay JA, Sorum M, Gould KA, Skov R. Genetic diversity in CC398 methicillin-resistant *Staphylococcus aureus* isolates of different geographical origin. Clin Microbiol Infect. 2010;16(7):1017–9.
- [21] Sung JM, Lloyd DH, Lindsay JA. *Staphylococcus aureus* host specificity: comparative genomics of human versus animal isolates by multi-strain microarray. Microbiology. 2008;154(Pt 7):1949–59.
- [22] de Neeling AJ, van den Broek MJ, Spalburg EC, van Santen-Verheувel MG, Dam-Deisz WD, Boshuizen HC, et al. High prevalence of methicillin resistant *Staphylococcus aureus* in pigs. Vet Microbiol. 2007;122(3–4):366–72.
- [23] van Loo I, Huijsdens X, Tiemersma E, de Neeling A, van de Sande-Bruinsma N, Beaujean D, et al. Emergence of methicillin-resistant *Staphylococcus aureus* of animal origin in humans. Emerg Infect Dis. 2007;13(12):1834–9.
- [24] Duquette RA, Nuttall TJ. Methicillin-resistant *Staphylococcus aureus* in dogs and cats: an emerging problem? J Small Anim Pract. 2004;45(12):591–7.
- [25] Smith TC, Pearson N. The emergence of *Staphylococcus aureus* ST398. Vector Borne Zoonotic Dis. 2011;11(4):327–39.
- [26] Johnson AP. Methicillin-resistant *Staphylococcus aureus*: the European landscape. J Antimicrob Chemother. 2011;66 Suppl 4:iv43–iv8.
- [27] Kock R, Mellmann A, Schaumburg F, Friedrich AW, Kipp F, Becker K. The epidemiology of methicillin-resistant *Staphylococcus aureus* (MRSA) in Germany. Dtsch Arztebl Int. 2011;108(45):761–7.
- [28] Larsen J, Imanishi M, Hinjoy S, Tharavichitkul P, Duangsong K, Davis MF, et al. Methicillin-resistant *Staphylococcus aureus* ST9 in pigs in Thailand. PLoS One. 2012;7(2):e31245.
- [29] Neela V, Mohd Zafrul A, Mariana NS, van Belkum A, Liew YK, Rad EG. Prevalence of ST9 methicillin-resistant *Staphylococcus aureus* among pigs and pig handlers in Malaysia. J Clin Microbiol. 2009;47(12):4138–40.
- [30] Wagenaar JA, Yue H, Pritchard J, Broekhuizen-Stins M, Huijsdens X, Mevius DJ, et al. Unexpected sequence types in livestock associated methicillin-resistant *Staphylococcus aureus* (MRSA): MRSA ST9 and a single locus variant of ST9 in pig farming in China. Vet Microbiol. 2009;139(3–4):405–9.
- [31] Patchanee P, Tadee P, Arjkumpa O, Love D, Chanachai K, Alter T, et al. Occurrence and characterization of livestock-associated methicillin-resistant *Staphylococcus aureus* in pig industries of northern Thailand. J Vet Sci. 2014;15(4):529–36.
- [32] Fang HW, Chiang PH, Huang YC. Livestock-associated methicillin-resistant *Staphylococcus aureus* ST9 in pigs and related personnel in Taiwan. PLoS One. 2014;9(2):e88826.

- [33] Argudin MA, Cariou N, Salandre O, Le Guennec J, Nemeghaire S, Butaye P. Genotyping and antimicrobial resistance of *Staphylococcus aureus* isolates from diseased turkeys. *Avian Pathol.* 2013;42(6):572–80.
- [34] Nemeghaire S, Roelandt S, Argudin MA, Haesebrouck F, Butaye P. Characterization of methicillin-resistant *Staphylococcus aureus* from healthy carrier chickens. *Avian Pathol.* 2013;42(4):342–6.
- [35] Buyukcangaz E, Velasco V, Sherwood JS, Stepan RM, Koslofsky RJ, Logue CM. Molecular typing of *Staphylococcus aureus* and methicillin-resistant *S. aureus* (MRSA) isolated from animals and retail meat in North Dakota, United States. *Foodborne Pathog Dis.* 2013;10(7):608–17.
- [36] Lowder BV, Guinane CM, Ben Zakour NL, Weinert LA, Conway-Morris A, Cartwright RA, et al. Recent human-to-poultry host jump, adaptation, and pandemic spread of *Staphylococcus aureus*. *Proc Natl Acad Sci USA.* 2009;106(46):19545–50.
- [37] Argudin MA, Mendoza MC, Rodicio MR. Food poisoning and *Staphylococcus aureus* enterotoxins. *Toxins (Basel).* 2010;2(7):1751–73.
- [38] Molla B, Byrne M, Abley M, Mathews J, Jackson CR, Fedorka-Cray P, et al. Epidemiology and genotypic characteristics of methicillin-resistant *Staphylococcus aureus* strains of porcine origin. *J Clin Microbiol.* 2012;50(11):3687–93.
- [39] de Jonge R, Verdier JE, Havelaar AH. Prevalence of methicillin-resistant *Staphylococcus aureus* amongst professional meat handlers in the Netherlands, March-July 2008. *Euro Surveill.* 2010;15(46):1–5.
- [40] Witte W, Cuny C. Emergence and spread of cfr-mediated multiresistance in staphylococci: an interdisciplinary challenge. *Future Microbiol.* 2011;6(8):925–31.
- [41] van Cleef BA, Graveland H, Haenen AP, van de Giessen AW, Heederik D, Wagenaar JA, et al. Persistence of livestock-associated methicillin-resistant *Staphylococcus aureus* in field workers after short-term occupational exposure to pigs and veal calves. *J Clin Microbiol.* 2011;49(3):1030–3.
- [42] Huber H, Giezendanner N, Stephan R, Zweifel C. Genotypes, antibiotic resistance profiles and microarray-based characterization of methicillin-resistant *Staphylococcus aureus* strains isolated from livestock and veterinarians in Switzerland. *Zoonoses Public Health.* 2011;58(5):343–9.
- [43] Fossum Moen AE, Saltyte Benth J, Alm-Kristiansen K, Bukholm G. Exotoxin-encoding gene content in community-acquired and hospital-acquired methicillin-resistant *Staphylococcus aureus*. *Clin Microbiol Infect.* 2009;15(12):1139–45.
- [44] Kadlec K, Ehricht R, Monecke S, Steinacker U, Kaspar H, Mankertz J, et al. Diversity of antimicrobial resistance pheno- and genotypes of methicillin-resistant *Staphylococcus aureus* ST398 from diseased swine. *J Antimicrob Chemother.* 2009;64(6):1156–64.

- [45] Alt K, Fetsch A, Schroeter A, Guerra B, Hammerl JA, Hertwig S, et al. Factors associated with the occurrence of MRSA CC398 in herds of fattening pigs in Germany. *BMC Vet Res*. 2011;7:69.
- [46] Kock R, Harlizius J, Bressan N, Laerberg R, Wieler LH, Witte W, et al. Prevalence and molecular characteristics of methicillin-resistant *Staphylococcus aureus* (MRSA) among pigs on German farms and import of livestock-related MRSA into hospitals. *Eur J Clin Microbiol Infect Dis*. 2009;28(11):1375–82.
- [47] Garcia-Graells C, Antoine J, Larsen J, Catry B, Skov R, Denis O. Livestock veterinarians at high risk of acquiring methicillin-resistant *Staphylococcus aureus* ST398. *Epidemiol Infect*. 2012;140(3):383–9.
- [48] Cuny C, Nathaus R, Layer F, Strommenger B, Altmann D, Witte W. Nasal colonization of humans with methicillin-resistant *Staphylococcus aureus* (MRSA) CC398 with and without exposure to pigs. *PLoS One*. 2009;4(8):e6800.
- [49] IV VDB, BA VANC, Haenen A, Broens EM, PJ VDW, MJ VDB, et al. Methicillin-resistant *Staphylococcus aureus* in people living and working in pig farms. *Epidemiol Infect*. 2009;137(5):700–8.
- [50] Wulf MW, Sorum M, van Nes A, Skov R, Melchers WJ, Klaassen CH, et al. Prevalence of methicillin-resistant *Staphylococcus aureus* among veterinarians: an international study. *Clin Microbiol Infect*. 2008;14(1):29–34.
- [51] Smith TC, Male MJ, Harper AL, Kroeger JS, Tinkler GP, Moritz ED, et al. Methicillin-resistant *Staphylococcus aureus* (MRSA) strain ST398 is present in midwestern U.S. swine and swine workers. *PLoS One*. 2009;4(1):e4258.
- [52] Hanson BM, Dressler AE, Harper AL, Scheibel RP, Wardyn SE, Roberts LK, et al. Prevalence of *Staphylococcus aureus* and methicillin-resistant *Staphylococcus aureus* (MRSA) on retail meat in Iowa. *J Infect Public Health*. 2011;4(4):169–74.
- [53] Uhlemann AC, Porcella SF, Trivedi S, Sullivan SB, Hafer C, Kennedy AD, et al. Identification of a highly transmissible animal-independent *Staphylococcus aureus* ST398 clone with distinct genomic and cell adhesion properties. *MBio*. 2012;3(2):e00027–12.
- [54] Bhat M, Dumortier C, Taylor BS, Miller M, Vasquez G, Yunen J, et al. *Staphylococcus aureus* ST398, New York City and Dominican Republic. *Emerg Infect Dis*. 2009;15(2):285–7.
- [55] Dressler AE, Scheibel RP, Wardyn S, Harper AL, Hanson BM, Kroeger JS, et al. Prevalence, antibiotic resistance and molecular characterisation of *Staphylococcus aureus* in pigs at agricultural fairs in the USA. *Vet Rec*. 2012;170(19):495.
- [56] Voss A, Loeffen F, Bakker J, Klaassen C, Wulf M. Methicillin-resistant *Staphylococcus aureus* in pig farming. *Emerg Infect Dis*. 2005;11(12):1965–6.

- [57] van Duijkeren E, Ikawaty R, Broekhuizen-Stins MJ, Jansen MD, Spalburg EC, de Neeling AJ, et al. Transmission of methicillin-resistant *Staphylococcus aureus* strains between different kinds of pig farms. *Vet Microbiol.* 2008;126(4):383–9.
- [58] Guardabassi L, Stegger M, Skov R. Retrospective detection of methicillin resistant and susceptible *Staphylococcus aureus* ST398 in Danish slaughter pigs. *Vet Microbiol.* 2007;122(3–4):384–6.
- [59] Meemken D, Cuny C, Witte W, Eichler U, Staudt R, Blaha T. Occurrence of MRSA in pigs and in humans involved in pig production—preliminary results of a study in the northwest of Germany. *Dtsch Tierarztl Wochenschr.* 2008;115(4):132–9.
- [60] Khanna T, Friendship R, Dewey C, Weese JS. Methicillin resistant *Staphylococcus aureus* colonization in pigs and pig farmers. *Vet Microbiol.* 2008;128(3–4):298–303.
- [61] European Food Safety Authority. Analysis of the baseline survey on the prevalence of methicillin-resistant *Staphylococcus aureus* (MRSA) in holdings with breeding pigs in the EU, 2008; Part A: MRSA prevalence estimates. *EFSA J.* 2009;7:1376.
- [62] Battisti A, Franco A, Merialdi G, Hasman H, Iurescia M, Lorenzetti R, et al. Heterogeneity among methicillin-resistant *Staphylococcus aureus* from Italian pig finishing holdings. *Vet Microbiol.* 2010;142(3–4):361–6.
- [63] Oppliger A, Moreillon P, Charriere N, Giddey M, Morisset D, Sakwinska O. Antimicrobial resistance of *Staphylococcus aureus* strains acquired by pig farmers from pigs. *Appl Environ Microbiol.* 2012;78(22):8010–4.
- [64] Broens EM, Graat EA, Van der Wolf PJ, Van de Giessen AW, De Jong MC. Prevalence and risk factor analysis of livestock associated MRSA-positive pig herds in The Netherlands. *Prev Vet Med.* 2011;102(1):41–9.
- [65] European Food Safety Authority. Analysis of the baseline survey on the prevalence of methicillin-resistant *Staphylococcus aureus* (MRSA) in holdings with breeding pigs, in the EU, 2008. Part B. Factors associated with MRSA contamination of holdings. *EFSA J* 2010;8:1597–663.
- [66] Crombe F, Willems G, Dispas M, Hallin M, Denis O, Suetens C, et al. Prevalence and antimicrobial susceptibility of methicillin-resistant *Staphylococcus aureus* among pigs in Belgium. *Microb Drug Resist.* 2012;18(2):125–31.
- [67] Broens EM, Graat EA, Van der Wolf PJ, Van de Giessen AW, De Jong MC. Transmission of methicillin resistant *Staphylococcus aureus* among pigs during transportation from farm to abattoir. *Vet J.* 2011;189(3):302–5.
- [68] Espinosa-Gongora C, Broens EM, Moodley A, Nielsen JP, Guardabassi L. Transmission of MRSA CC398 strains between pig farms related by trade of animals. *Vet Rec.* 2012;170(22):564.

- [69] Weese JS, Zwambag A, Rosendal T, Reid-Smith R, Friendship R. Longitudinal investigation of methicillin-resistant *Staphylococcus aureus* in piglets. *Zoonoses Public Health*. 2011;58(4):238–43.
- [70] van Duijkeren E, Jansen MD, Flemming SC, de Neeling H, Wagenaar JA, Schoormans AH, et al. Methicillin-resistant *Staphylococcus aureus* in pigs with exudative epidermitis. *Emerg Infect Dis*. 2007;13(9):1408–10.
- [71] Schwarz S, Kadlec K, Strommenger B. Methicillin-resistant *Staphylococcus aureus* and *Staphylococcus pseudintermedius* detected in the BfT-GermVet monitoring programme 2004–2006 in Germany. *J Antimicrob Chemother*. 2008;61(2):282–5.
- [72] Weese JS, Reid-Smith R, Rousseau J, Avery B. Methicillin-resistant *Staphylococcus aureus* (MRSA) contamination of retail pork. *Can Vet J*. 2010;51(7):749–52.
- [73] Osadebe LU, Hanson B, Smith TC, Heimer R. Prevalence and characteristics of *Staphylococcus aureus* in Connecticut swine and swine farmers. *Zoonoses Public Health*. 2013;60(3):234–43.
- [74] Gordoncillo MJ, Abdujamilova N, Perri M, Donabedian S, Zervos M, Bartlett P. Detection of methicillin-resistant *Staphylococcus aureus* (MRSA) in backyard pigs and their owners, Michigan, USA. *Zoonoses Public Health*. 2012;59(3):212–6.
- [75] Frana TS, Beahm AR, Hanson BM, Kinyon JM, Layman LL, Karriker LA, et al. Isolation and characterization of methicillin-resistant *Staphylococcus aureus* from pork farms and visiting veterinary students. *PLoS One*. 2013;8(1):e53738.
- [76] Smith TC, Gebreyes WA, Abley MJ, Harper AL, Forshey BM, Male MJ, et al. Methicillin-resistant *Staphylococcus aureus* in pigs and farm workers on conventional and antibiotic-free swine farms in the USA. *PLoS One*. 2013;8(5):e63704.
- [77] Rinsky JL, Nadimpalli M, Wing S, Hall D, Baron D, Price LB, et al. Livestock-associated methicillin and multidrug resistant *Staphylococcus aureus* is present among industrial, not antibiotic-free livestock operation workers in North Carolina. *PLoS One*. 2013;8(7):e67641.
- [78] Neyra RC, Frisancho JA, Rinsky JL, Resnick C, Carroll KC, Rule AM, et al. Multidrug-resistant and methicillin-resistant *Staphylococcus aureus* (MRSA) in hog slaughter and processing plant workers and their community in North Carolina (USA). *Environ Health Perspect*. 2014;122(5):471–7.
- [79] Nadimpalli M, Rinsky JL, Wing S, Hall D, Stewart J, Larsen J, et al. Persistence of livestock-associated antibiotic-resistant *Staphylococcus aureus* among industrial hog operation workers in North Carolina over 14 days. *Occup Environ Med*. 2015;72(2):90–9.
- [80] Cui S, Li J, Hu C, Jin S, Li F, Guo Y, et al. Isolation and characterization of methicillin-resistant *Staphylococcus aureus* from swine and workers in China. *J Antimicrob Chemother*. 2009;64(4):680–3.

- [81] Ho PL, Chow KH, Lai EL, Law PY, Chan PY, Ho AY, et al. Clonality and antimicrobial susceptibility of *Staphylococcus aureus* and methicillin-resistant *S. aureus* isolates from food animals and other animals. J Clin Microbiol. 2012;50(11):3735–7.
- [82] Sergio DM, Koh TH, Hsu LY, Ogden BE, Goh AL, Chow PK. Investigation of methicillin-resistant *Staphylococcus aureus* in pigs used for research. J Med Microbiol. 2007;56(Pt 8):1107–9.
- [83] Hsu LY, Koh TH, Singh K, Kang ML, Kurup A, Tan BH. Dissemination of multisusceptible methicillin-resistant *Staphylococcus aureus* in Singapore. J Clin Microbiol. 2005;43(6):2923–5.
- [84] Tsai HY, Liao CH, Cheng A, Liu CY, Huang YT, Teng LJ, et al. Isolation of methicillin-resistant *Staphylococcus aureus* sequence type 9 in pigs in Taiwan. Int J Antimicrob Agents. 2012;39(5):449–51.
- [85] Lim SK, Nam HM, Jang GC, Lee HS, Jung SC, Kwak HS. The first detection of methicillin-resistant *Staphylococcus aureus* ST398 in pigs in Korea. Vet Microbiol. 2012;155(1):88–92.
- [86] Devriese LA, Van Damme LR, Fameree L. Methicillin (cloxacillin)-resistant *Staphylococcus aureus* strains isolated from bovine mastitis cases. Zentralbl Veterinarmed B. 1972;19(7):598–605.
- [87] Vanderhaeghen W, Cerpentier T, Adriaensen C, Vicca J, Hermans K, Butaye P. Methicillin-resistant *Staphylococcus aureus* (MRSA) ST398 associated with clinical and subclinical mastitis in Belgian cows. Vet Microbiol. 2010;144(1–2):166–71.
- [88] Devriese LA, Hommez J. Epidemiology of methicillin-resistant *Staphylococcus aureus* in dairy herds. Res Vet Sci. 1975;19(1):23–7.
- [89] Lee JH. Methicillin (Oxacillin)-resistant *Staphylococcus aureus* strains isolated from major food animals and their potential transmission to humans. Appl Environ Microbiol. 2003;69(11):6489–94.
- [90] Moon JS, Lee AR, Kang HM, Lee ES, Kim MN, Paik YH, et al. Phenotypic and genetic antibiogram of methicillin-resistant staphylococci isolated from bovine mastitis in Korea. J Dairy Sci. 2007;90(3):1176–85.
- [91] Moon DC, Tamang MD, Nam HM, Jeong JH, Jang GC, Jung SC, et al. Identification of livestock-associated methicillin-resistant *Staphylococcus aureus* isolates in Korea and molecular comparison between isolates from animal carcasses and slaughterhouse workers. Foodborne Pathog Dis. 2015;12(4):327–34.
- [92] Juhasz-Kaszanyitzky E, Janosi S, Somogyi P, Dan A, van der Graaf-van Bloois L, van Duijkeren E, et al. MRSA transmission between cows and humans. Emerg Infect Dis. 2007;13(4):630–2.
- [93] Weese JS, Archambault M, Willey BM, Hearn P, Kreiswirth BN, Said-Salim B, et al. Methicillin-resistant *Staphylococcus aureus* in horses and horse personnel, 2000–2002. Emerg Infect Dis. 2005;11(3):430–5.

- [94] Umoh VJ, Adesiyun AA, Gomwalk NE. Antibigram of staphylococcal strains isolated from milk and milk-products. Zentralbl Veterinarmed B. 1990;37(9):701–6.
- [95] Costa EO, Benites NR, Guerra JL, Melville PA. Antimicrobial susceptibility of *Staphylococcus* spp. isolated from mammary parenchymas of slaughtered dairy cows. J Vet Med B Infect Dis Vet Public Health. 2000;47(2):99–103.
- [96] Erskine RJ, Walker RD, Bolin CA, Bartlett PC, White DG. Trends in antibacterial susceptibility of mastitis pathogens during a seven-year period. J Dairy Sci. 2002;85(5):1111–8.
- [97] Fessler A, Scott C, Kadlec K, Ehricht R, Monecke S, Schwarz S. Characterization of methicillin-resistant *Staphylococcus aureus* ST398 from cases of bovine mastitis. J Antimicrob Chemother. 2010;65(4):619–25.
- [98] Bos ME, Graveland H, Portengen L, Wagenaar JA, Heederik DJ. Livestock-associated MRSA prevalence in veal calf production is associated with farm hygiene, use of antimicrobials, and age of the calves. Prev Vet Med. 2012;105(1–2):155–9.
- [99] Eurosurveillance editorial team. European Union Summary Report on antimicrobial resistance in zoonotic and indicator bacteria from humans, animals and food 2012 published. Euro Surveill. 2014;19(12):20748.
- [100] Graveland H, Wagenaar JA, Bergs K, Heesterbeek H, Heederik D. Persistence of livestock associated MRSA CC398 in humans is dependent on intensity of animal contact. PLoS One. 2011;6(2):e16830.
- [101] Holmes MA, Zadoks RN. Methicillin resistant *S. aureus* in human and bovine mastitis. J Mammary Gland Biol Neoplasia. 2011;16(4):373–82.
- [102] Spohr M, Rau J, Friedrich A, Klittich G, Fetsch A, Guerra B, et al. Methicillin-resistant *Staphylococcus aureus* (MRSA) in three dairy herds in southwest Germany. Zoonoses Public Health. 2011;58(4):252–61.
- [103] Monecke S, Kuhnert P, Hotzel H, Slickers P, Ehricht R. Microarray based study on virulence-associated genes and resistance determinants of *Staphylococcus aureus* isolates from cattle. Vet Microbiol. 2007;125(1–2):128–40.
- [104] Hata E, Katsuda K, Kobayashi H, Uchida I, Tanaka K, Eguchi M. Genetic variation among *Staphylococcus aureus* strains from bovine milk and their relevance to methicillin-resistant isolates from humans. J Clin Microbiol. 2010;48(6):2130–9.
- [105] van Rijen MM, Bosch T, Heck ME, Kluytmans JA. Methicillin-resistant *Staphylococcus aureus* epidemiology and transmission in a Dutch hospital. J Hosp Infect. 2009;72(4):299–306.
- [106] van Rijen MM, Van Keulen PH, Kluytmans JA. Increase in a Dutch hospital of methicillin-resistant *Staphylococcus aureus* related to animal farming. Clin Infect Dis. 2008;46(2):261–3.
- [107] Vandenbroucke-Grauls CM, Beaujean DJ. Methicillin-resistant *Staphylococcus aureus* in pig breeders and cattle breeders. Ned Tijdschr Geneesk. 2006;150(31):1710–2.

- [108] Nam HM, Lee AL, Jung SC, Kim MN, Jang GC, Wee SH, et al. Antimicrobial susceptibility of *Staphylococcus aureus* and characterization of methicillin-resistant *Staphylococcus aureus* isolated from bovine mastitis in Korea. *Foodborne Pathog Dis.* 2011;8(2):231–8.
- [109] Kwon NH, Park KT, Moon JS, Jung WK, Kim SH, Kim JM, et al. Staphylococcal cassette chromosome mec (SCCmec) characterization and molecular analysis for methicillin-resistant *Staphylococcus aureus* and novel SCCmec subtype IVg isolated from bovine milk in Korea. *J Antimicrob Chemother.* 2005;56(4):624–32.
- [110] Feltrin F, Alba P, Kraushaar B, Ianzano A, Argudin MA, Di Matteo P, et al. A livestock-associated, multidrug-resistant, methicillin-resistant *Staphylococcus aureus* clonal complex 97 lineage spreading in dairy cattle and pigs in Italy. *Appl Environ Microbiol.* 2016;82(3):816–21.
- [111] Garcia-Alvarez L, Holden MT, Lindsay H, Webb CR, Brown DF, Curran MD, et al. Methicillin-resistant *Staphylococcus aureus* with a novel mecA homologue in human and bovine populations in the UK and Denmark: a descriptive study. *Lancet Infect Dis.* 2011;11(8):595–603.
- [112] Petersen A, Stegger M, Heltberg O, Christensen J, Zeuthen A, Knudsen LK, et al. Epidemiology of methicillin-resistant *Staphylococcus aureus* carrying the novel mecC gene in Denmark corroborates a zoonotic reservoir with transmission to humans. *Clin Microbiol Infect.* 2013;19(1):E16–22.
- [113] Weese JS, Hannon SJ, Booker CW, Gow S, Avery BP, Reid-Smith RJ. The prevalence of methicillin-resistant *Staphylococcus aureus* colonization in feedlot cattle. *Zoonoses Public Health.* 2012;59(2):144–7.
- [114] Golding GR, Levett PN, McDonald RR, Wylie J, Mulvey MR, editors. Molecular characterization of the first clinical infections of “livestock associated” ST398 methicillin-resistant *Staphylococcus aureus* in Canada. American Society for Microbiology (ASM)-ESCMID Conference on Methicillin-Resistant Staphylococci in Animals; London.
- [115] Lee JH. Occurrence of methicillin-resistant *Staphylococcus aureus* strains from cattle and chicken, and analyses of their mecA, mecR1 and mecI genes. *Vet Microbiol.* 2006;114(1–2):155–9.
- [116] Alves PD, McCulloch JA, Even S, Le Marechal C, Thierry A, Grosset N, et al. Molecular characterisation of *Staphylococcus aureus* strains isolated from small and large ruminants reveals a host rather than tissue specificity. *Vet Microbiol.* 2009;137(1–2):190–5.
- [117] Haenni M, Saras E, Chatre P, Medaille C, Bes M, Madec JY, et al. A USA300 variant and other human-related methicillin-resistant *Staphylococcus aureus* strains infecting cats and dogs in France. *J Antimicrob Chemother.* 2012;67(2):326–9.
- [118] Lin Y, Barker E, Kislow J, Kaldhorne P, Stemper ME, Pantrangi M, et al. Evidence of multiple virulence subtypes in nosocomial and community-associated MRSA genotypes in

- companion animals from the upper midwestern and northeastern United States. Clin Med Res. 2011;9(1):7–16.
- [119] Alba P, Feltrin F, Cordaro G, Porrero MC, Kraushaar B, Argudin MA, et al. Livestock-associated methicillin resistant and methicillin susceptible *Staphylococcus aureus* sequence type (CC)1 in European farmed animals: high genetic relatedness of isolates from Italian cattle herds and humans. PLoS One. 2015;10(8):e0137143.
- [120] Heikinheimo A, Johler S, Karvonen L, Julmi J, Fredriksson-Ahomaa M, Stephan R. New dominant spa type t2741 in livestock-associated MRSA (CC398-MRSA-V) in Finnish fattening pigs at slaughter. Antimicrob Resist Infect Control. 2016;5:6.
- [121] Peeters LE, Argudin MA, Azadikhah S, Butaye P. Antimicrobial resistance and population structure of *Staphylococcus aureus* recovered from pigs farms. Vet Microbiol. 2015;180(1–2):151–6.
- [122] Kitai S, Shimizu A, Kawano J, Sato E, Nakano C, Kitagawa H, et al. Prevalence and characterization of *Staphylococcus aureus* and enterotoxigenic *Staphylococcus aureus* in retail raw chicken meat throughout Japan. J Vet Med Sci. 2005;67(3):269–74.
- [123] Spoor LE, McAdam PR, Weinert LA, Rambaut A, Hasman H, Aarestrup FM, et al. Livestock origin for a human pandemic clone of community-associated methicillin-resistant *Staphylococcus aureus*. MBio. 2013;4(4):e00356–13.
- [124] Pletinckx LJ, Verhegghe M, Dewulf J, Crombe F, De Bleecker Y, Rasschaert G, et al. Screening of poultry-pig farms for methicillin-resistant *Staphylococcus aureus*: sampling methodology and within herd prevalence in broiler flocks and pigs. Infect Genet Evol. 2011;11(8):2133–7.
- [125] Persoons D, Van Hoorebeke S, Hermans K, Butaye P, de Kruif A, Haesebrouck F, et al. Methicillin-resistant *Staphylococcus aureus* in poultry. Emerg Infect Dis. 2009;15(3):452–3.
- [126] Vossenkuhl B, Brandt J, Fetsch A, Kasbohrer A, Kraushaar B, Alt K, et al. Comparison of spa types, SCCmec types and antimicrobial resistance profiles of MRSA isolated from turkeys at farm, slaughter and from retail meat indicates transmission along the production chain. PLoS One. 2014;9(5):e96308.
- [127] Pu S, Han F, Ge B. Isolation and characterization of methicillin-resistant *Staphylococcus aureus* strains from Louisiana retail meats. Appl Environ Microbiol. 2009;75(1):265–7.
- [128] Lin J, Yeh KS, Liu HT, Lin JH. *Staphylococcus aureus* isolated from pork and chicken carcasses in Taiwan: prevalence and antimicrobial susceptibility. J Food Prot. 2009;72(3):608–11.
- [129] Hasman H, Moodley A, Guardabassi L, Stegger M, Skov RL, Aarestrup FM. Spa type distribution in *Staphylococcus aureus* originating from pigs, cattle and poultry. Vet Microbiol. 2010;141(3–4):326–31.

- [130] Cavaco LM, Hasman H, Aarestrup FM. Zinc resistance of *Staphylococcus aureus* of animal origin is strongly associated with methicillin resistance. *Vet Microbiol.* 2011;150(3–4):344–8.
- [131] Smyth DS, Feil EJ, Meaney WJ, Hartigan PJ, Tollersrud T, Fitzgerald JR, et al. Molecular genetic typing reveals further insights into the diversity of animal-associated *Staphylococcus aureus*. *J Med Microbiol.* 2009;58(Pt 10):1343–53.
- [132] Mulders MN, Haenen AP, Geenen PL, Vesseur PC, Poldervaart ES, Bosch T, et al. Prevalence of livestock-associated MRSA in broiler flocks and risk factors for slaughterhouse personnel in The Netherlands. *Epidemiol Infect.* 2010;138(5):743–55.
- [133] Pletinckx LJ, Verheghe M, Crombe F, Dewulf J, De Bleecker Y, Rasschaert G, et al. Evidence of possible methicillin-resistant *Staphylococcus aureus* ST398 spread between pigs and other animals and people residing on the same farm. *Prev Vet Med.* 2013;109(3–4):293–303.
- [134] Wendlandt S, Li J, Ho J, Porta MA, Fessler AT, Wang Y, et al. Enterococcal multiresistance gene cluster in methicillin-resistant *Staphylococcus aureus* from various origins and geographical locations. *J Antimicrob Chemother.* 2014;69(9):2573–5.
- [135] Kwon NH, Park KT, Jung WK, Youn HY, Lee Y, Kim SH, et al. Characteristics of methicillin resistant *Staphylococcus aureus* isolated from chicken meat and hospitalized dogs in Korea and their epidemiological relatedness. *Vet Microbiol.* 2006;117(2–4):304–12.
- [136] Ekkelenkamp MB, Sekkat M, Carpaij N, Troelstra A, Bonten MJ. Endocarditis due to methicillin-resistant *Staphylococcus aureus* originating from pigs. *Ned Tijdschr Geneesk.* 2006;150(44):2442–7.
- [137] de Boer E, Zwartkruis-Nahuis JT, Wit B, Huijsdens XW, de Neeling AJ, Bosch T, et al. Prevalence of methicillin-resistant *Staphylococcus aureus* in meat. *Int J Food Microbiol.* 2009;134(1–2):52–6.
- [138] van Loo IH, Diederiksen BM, Savelkoul PH, Woudenberg JH, Roosendaal R, van Belkum A, et al. Methicillin-resistant *Staphylococcus aureus* in meat products, the Netherlands. *Emerg Infect Dis.* 2007;13(11):1753–5.
- [139] Schraft H, Kleinlein N, Untermann F. Contamination of pig hindquarters with *Staphylococcus aureus*. *Int J Food Microbiol.* 1992;15(1–2):191–4.
- [140] Hendriksen RS, Mevius DJ, Schroeter A, Teale C, Meunier D, Butaye P, et al. Prevalence of antimicrobial resistance among bacterial pathogens isolated from cattle in different European countries: 2002–2004. *Acta Vet Scand.* 2008;50:28.
- [141] Tenhagen BA, Vossenkühl B, Kasbohrer A, Alt K, Kraushaar B, Guerra B, et al. Methicillin-resistant *Staphylococcus aureus* in cattle food chains—prevalence, diversity, and antimicrobial resistance in Germany. *J Anim Sci.* 2014;92(6):2741–51.

- [142] Foster AP. Staphylococcal skin disease in livestock. *Vet Dermatol*. 2012;23(4):342–51, e63.
- [143] Atyah MA, Zamri-Saad M, Siti-Zahrah A. First report of methicillin-resistant *Staphylococcus aureus* from cage-cultured tilapia (*Oreochromis niloticus*). *Vet Microbiol*. 2010;144(3–4):502–4.
- [144] Rhee CH, Woo GJ. Emergence and characterization of foodborne methicillin-resistant *Staphylococcus aureus* in Korea. *J Food Prot*. 2010;73(12):2285–90.
- [145] Hammad AM, Watanabe W, Fujii T, Shimamoto T. Occurrence and characteristics of methicillin-resistant and -susceptible *Staphylococcus aureus* and methicillin-resistant coagulase-negative staphylococci from Japanese retail ready-to-eat raw fish. *Int J Food Microbiol*. 2012;156(3):286–9.
- [146] Sergelidis D, Abraham A, Papadopoulos T, Soultos N, Martziou E, Koulourida V, et al. Isolation of methicillin-resistant *Staphylococcus* spp. from ready-to-eat fish products. *Lett Appl Microbiol*. 2014;59(5):500–6.
- [147] Sugimoto K, Frei R, Graber P. Methicillin-resistant *Staphylococcus aureus* foot infection after fish pedicure. *Infection*. 2013;41(5):1013–5.
- [148] Devriese LA. A simplified system for biotyping *Staphylococcus aureus* strains isolated from animal species. *J Appl Bacteriol*. 1984;56(2):215–20.
- [149] Bhargava K, Wang X, Donabedian S, Zervos M, de Rocha L, Zhang Y. Methicillin-resistant *Staphylococcus aureus* in retail meat, Detroit, Michigan, USA. *Emerg Infect Dis*. 2011;17(6):1135–7.
- [150] Kelman A, Soong YA, Dupuy N, Shafer D, Richbourg W, Johnson K, et al. Antimicrobial susceptibility of *Staphylococcus aureus* from retail ground meats. *J Food Prot*. 2011;74(10):1625–9.
- [151] Carr MA, Thompson LD, Miller MF, Ramsey CB, Kaster CS. Chilling and trimming effects on the microbial populations of pork carcasses. *J Food Prot*. 1998;61(4):487–9.
- [152] Beneke B, Klees S, Stuhrenberg B, Fetsch A, Kraushaar B, Tenhagen BA. Prevalence of methicillin-resistant *Staphylococcus aureus* in a fresh meat pork production chain. *J Food Prot*. 2011;74(1):126–9.
- [153] Borch E, Nesbakken T, Christensen H. Hazard identification in swine slaughter with respect to foodborne bacteria. *Int J Food Microbiol*. 1996;30(1–2):9–25.
- [154] Spescha C, Stephan R, Zweifel C. Microbiological contamination of pig carcasses at different stages of slaughter in two European Union-approved abattoirs. *J Food Prot*. 2006;69(11):2568–75.
- [155] Bolton DJ, Pearce RA, Sheridan JJ, Blair IS, McDowell DA, Harrington D. Washing and chilling as critical control points in pork slaughter hazard analysis and critical control point (HACCP) systems. *J Appl Microbiol*. 2002;92(5):893–902.

- [156] Vossenkuhl B, Sharp H, Brandt J, Fetsch A, Käsböhrer A, Tenhagen BA. Modeling the transmission of livestock associated methicillin-resistant *Staphylococcus aureus* along the pig slaughter line. *Food Control*. 2014;39:17–24.
- [157] Dockerty TR, Ockerman HW, Cahill VR, Kunkle LE, Weiser HH. Microbial level of pork skin as affected by the dressing process. *J Anim Sci* 1970;30:884–90.
- [158] Pearce RA, Bolton DJ, Sheridan JJ, McDowell DA, Blair IS, Harrington D. Studies to determine the critical control points in pork slaughter hazard analysis and critical control point systems. *Int J Food Microbiol*. 2004;90(3):331–9.
- [159] Yu SL, Bolton D, Laubach C, Kline P, Oser A, Palumbo SA. Effect of dehairing operations on microbiological quality of swine carcasses. *J Food Prot*. 1999;62(12):1478–81.
- [160] Saide-Albornoz JJ, Lynn Knipe C, Murano EA, Beran GW. Contamination of pork carcasses during slaughter, fabrication, and chilled storage. *J Food Prot* 1995;58:993–7.
- [161] Rivas T, Vizcaino JA, Herrera FJ. Microbial contamination of carcasses and equipment from an Iberian pig slaughterhouse. *J Food Prot*. 2000;63(12):1670–5.
- [162] Brown T, James SJ. Process design data for pork chilling. *Int J Refrig*. 1992;15:281–9.
- [163] James S. The chill chain “from carcass to consumer”. *Meat Sci*. 1996;43S1:203–16.
- [164] Graveland H, Duim B, van Duijkeren E, Heederik D, Wagenaar JA. Livestock-associated methicillin-resistant *Staphylococcus aureus* in animals and humans. *Int J Med Microbiol*. 2011;301(8):630–4.
- [165] Lewis HC, Molbak K, Reese C, Aarestrup FM, Selchau M, Sorum M, et al. Pigs as source of methicillin-resistant *Staphylococcus aureus* CC398 infections in humans, Denmark. *Emerg Infect Dis*. 2008;14(9):1383–9.
- [166] Cefai C, Ashurst S, Owens C. Human carriage of methicillin-resistant *Staphylococcus aureus* linked with pet dog. *Lancet*. 1994;344(8921):539–40.
- [167] Price LB, Stegger M, Hasman H, Aziz M, Larsen J, Andersen PS, et al. *Staphylococcus aureus* CC398: host adaptation and emergence of methicillin resistance in livestock. *MBio*. 2012;3(1):e00305–11.
- [168] Rooijackers SH, Ruyken M, Roos A, Daha MR, Presanis JS, Sim RB, et al. Immune evasion by a staphylococcal complement inhibitor that acts on C3 convertases. *Nat Immunol*. 2005;6(9):920–7.
- [169] Stegger M, Lindsay JA, Moodley A, Skov R, Broens EM, Guardabassi L. Rapid PCR detection of *Staphylococcus aureus* clonal complex 398 by targeting the restriction-modification system carrying sau1-hsdS1. *J Clin Microbiol*. 2011;49(2):732–4.
- [170] Waldron DE, Lindsay JA. Sau1: a novel lineage-specific type I restriction-modification system that blocks horizontal gene transfer into *Staphylococcus aureus* and between *S. aureus* isolates of different lineages. *J Bacteriol*. 2006;188(15):5578–85.

- [171] Waters AE, Contente-Cuomo T, Buchhagen J, Liu CM, Watson L, Pearce K, et al. Multidrug-resistant *Staphylococcus aureus* in US Meat and Poultry. Clin Infect Dis. 2011;52(10):1227–30.
- [172] Wieler LH, Ewers C, Guenther S, Walther B, Lubke-Becker A. Methicillin-resistant staphylococci (MRS) and extended-spectrum beta-lactamases (ESBL)-producing Enterobacteriaceae in companion animals: nosocomial infections as one reason for the rising prevalence of these potential zoonotic pathogens in clinical samples. Int J Med Microbiol. 2011;301(8):635–41.
- [173] Takeuchi S, Matsunaga K, Inubushi S, Higuchi H, Imaizumi K, Kaidoh T. Structural gene and strain specificity of a novel cysteine protease produced by *Staphylococcus aureus* isolated from a diseased chicken. Vet Microbiol. 2002;89(2–3):201–10.
- [174] Ward MJ, Gibbons CL, McAdam PR, van Bunnik BA, Girvan EK, Edwards GF, et al. Time-scaled evolutionary analysis of the transmission and antibiotic resistance dynamics of *Staphylococcus aureus* clonal complex 398. Appl Environ Microbiol. 2014;80(23):7275–82.
- [175] Jansen MD, Box AT, Fluit AC. SCCmec typing in methicillin-resistant *Staphylococcus aureus* strains of animal origin. Emerg Infect Dis. 2009;15(1):136–7; author reply-7.
- [176] Zhang Y, Agidi S, LeJeune JT. Diversity of staphylococcal cassette chromosome in coagulase-negative staphylococci from animal sources. J Appl Microbiol. 2009;107(4):1375–83.
- [177] Guardabassi L, O'Donoghue M, Moodley A, Ho J, Boost M. Novel lineage of methicillin-resistant *Staphylococcus aureus*, Hong Kong. Emerg Infect Dis. 2009;15(12):1998–2000.
- [178] Fitzgerald JR, Sturdevant DE, Mackie SM, Gill SR, Musser JM. Evolutionary genomics of *Staphylococcus aureus*: insights into the origin of methicillin-resistant strains and the toxic shock syndrome epidemic. Proc Natl Acad Sci USA. 2001;98(15):8821–6.
- [179] Oliveira DC, Tomasz A, de Lencastre H. Secrets of success of a human pathogen: molecular evolution of pandemic clones of methicillin-resistant *Staphylococcus aureus*. Lancet Infect Dis. 2002;2(3):180–9.
- [180] Nubel U, Roumagnac P, Feldkamp M, Song JH, Ko KS, Huang YC, et al. Frequent emergence and limited geographic dispersal of methicillin-resistant *Staphylococcus aureus*. Proc Natl Acad Sci USA. 2008;105(37):14130–5.
- [181] Rijnders MI, Deurenberg RH, Boumans ML, Hoogkamp-Korstanje JA, Beisser PS, Antibiotic Resistance Surveillance Group, et al. Population structure of *Staphylococcus aureus* strains isolated from intensive care unit patients in the Netherlands over an 11-year period (1996 to 2006). J Clin Microbiol. 2009;47(12):4090–5.
- [182] Varshney AK, Mediavilla JR, Robiou N, Guh A, Wang X, Gialanella P, et al. Diverse enterotoxin gene profiles among clonal complexes of *Staphylococcus aureus* isolates from the Bronx, New York. Appl Environ Microbiol. 2009;75(21):6839–49.

- [183] Walther B, Monecke S, Ruscher C, Friedrich AW, Ehricht R, Slickers P, et al. Comparative molecular analysis substantiates zoonotic potential of equine methicillin-resistant *Staphylococcus aureus*. J Clin Microbiol. 2009;47(3):704–10.
- [184] McCarthy AJ, Witney AA, Gould KA, Moodley A, Guardabassi L, Voss A, et al. The distribution of mobile genetic elements (MGEs) in MRSA CC398 is associated with both host and country. Genome Biol Evol. 2011;3:1164–74.
- [185] van Belkum A, Melles DC, Peeters JK, van Leeuwen WB, van Duijkeren E, Huijsdens XW, et al. Methicillin-resistant and -susceptible *Staphylococcus aureus* sequence type 398 in pigs and humans. Emerg Infect Dis. 2008;14(3):479–83.
- [186] Welinder-Olsson C, Floren-Johansson K, Larsson L, Oberg S, Karlsson L, Ahren C. Infection with Panton-Valentine leukocidin-positive methicillin-resistant *Staphylococcus aureus* t034. Emerg Infect Dis. 2008;14(8):1271–2.
- [187] Yu F, Chen Z, Liu C, Zhang X, Lin X, Chi S, et al. Prevalence of *Staphylococcus aureus* carrying Panton-Valentine leukocidin genes among isolates from hospitalised patients in China. Clin Microbiol Infect. 2008;14(4):381–4.
- [188] Ola Ojo M. Bacteriophage types and antibiotic sensitivity of *Staphylococcus aureus* isolated from swabs of the noses and skins of dogs. Vet Rec. 1972;91(6):152–3.
- [189] Wang N, Neilan AM, Klompas M. *Staphylococcus intermedius* infections: case report and literature review. Infect Dis Rep. 2013;5(1):e3.
- [190] Nagase N, Sasaki A, Yamashita K, Shimizu A, Wakita Y, Kitai S, et al. Isolation and species distribution of staphylococci from animal and human skin. J Vet Med Sci. 2002;64(3):245–50.
- [191] Guardabassi L, Schwarz S, Lloyd DH. Pet animals as reservoirs of antimicrobial-resistant bacteria. J Antimicrob Chemother. 2004;54(2):321–32.
- [192] Patel A, Lloyd DH, Howell SA, Noble WC. Investigation into the potential pathogenicity of *Staphylococcus felis* in a cat. Vet Rec. 2002;150(21):668–9.
- [193] Lilenbaum W, Nunes EL, Azeredo MA. Prevalence and antimicrobial susceptibility of staphylococci isolated from the skin surface of clinically normal cats. Lett Appl Microbiol. 1998;27(4):224–8.
- [194] Couto N, Pomba C, Moodley A, Guardabassi L. Prevalence of methicillin-resistant staphylococci among dogs and cats at a veterinary teaching hospital in Portugal. Vet Rec. 2011;169(3):72.
- [195] Loeffler A, Pfeiffer DU, Lindsay JA, Soares Magalhaes RJ, Lloyd DH. Prevalence of and risk factors for MRSA carriage in companion animals: a survey of dogs, cats and horses. Epidemiol Infect. 2011;139(7):1019–28.

- [196] Loeffler A, Boag AK, Sung J, Lindsay JA, Guardabassi L, Dalsgaard A, et al. Prevalence of methicillin-resistant *Staphylococcus aureus* among staff and pets in a small animal referral hospital in the UK. *J Antimicrob Chemother.* 2005;56(4):692–7.
- [197] Weese JS, Faires M, Rousseau J, Bersenas AM, Mathews KA. Cluster of methicillin-resistant *Staphylococcus aureus* colonization in a small animal intensive care unit. *J Am Vet Med Assoc.* 2007;231(9):1361–4.
- [198] Weese JS, van Duijkeren E. Methicillin-resistant *Staphylococcus aureus* and *Staphylococcus pseudintermedius* in veterinary medicine. *Vet Microbiol.* 2010;140(3–4):418–29.
- [199] Gortel K, Campbell KL, Kakoma I, Whitem T, Schaeffer DJ, Weisiger RM. Methicillin resistance among staphylococci isolated from dogs. *Am J Vet Res.* 1999;60(12):1526–30.
- [200] Griffeth GC, Morris DO, Abraham JL, Shofer FS, Rankin SC. Screening for skin carriage of methicillin-resistant coagulase-positive staphylococci and *Staphylococcus schleiferi* in dogs with healthy and inflamed skin. *Vet Dermatol.* 2008;19(3):142–9.
- [201] Abraham JL, Morris DO, Griffeth GC, Shofer FS, Rankin SC. Surveillance of healthy cats and cats with inflammatory skin disease for colonization of the skin by methicillin-resistant coagulase-positive staphylococci and *Staphylococcus schleiferi* ssp. *schleiferi*. *Vet Dermatol.* 2007;18(4):252–9.
- [202] Soares Magalhaes RJ, Loeffler A, Lindsay J, Rich M, Roberts L, Smith H, et al. Risk factors for methicillin-resistant *Staphylococcus aureus* (MRSA) infection in dogs and cats: a case-control study. *Vet Res.* 2010;41(5):55.
- [203] Rich M, Roberts L. Methicillin-resistant *Staphylococcus aureus* isolates from companion animals. *Vet Rec.* 2004;154(10):310.
- [204] Quitoco IM, Ramundo MS, Silva-Carvalho MC, Souza RR, Beltrame CO, de Oliveira TF, et al. First report in South America of companion animal colonization by the USA1100 clone of community-acquired methicillin-resistant *Staphylococcus aureus* (ST30) and by the European clone of methicillin-resistant *Staphylococcus pseudintermedius* (ST71). *BMC Res Notes.* 2013;6:336.
- [205] Chanchaithong P, Perreten V, Schwendener S, Tribuddharat C, Chongthaleong A, Niyomtham W, et al. Strain typing and antimicrobial susceptibility of methicillin-resistant coagulase-positive staphylococcal species in dogs and people associated with dogs in Thailand. *J Appl Microbiol.* 2014;117(2):572–86.
- [206] Couto N, Belas A, Kadlec K, Schwarz S, Pombo C. Clonal diversity, virulence patterns and antimicrobial and biocide susceptibility among human, animal and environmental MRSA in Portugal. *J Antimicrob Chemother.* 2015;70(9):2483–7.
- [207] Weiss S, Kadlec K, Fessler AT, Schwarz S. Identification and characterization of methicillin-resistant *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus haemolyticus* and *Staphylococcus pettenkoferi* from a small animal clinic. *Vet Microbiol.* 2013;167(3–4):680–5.

- [208] Loncaric I, Kunzel F, Licka T, Simhofer H, Spargser J, Rosengarten R. Identification and characterization of methicillin-resistant *Staphylococcus aureus* (MRSA) from Austrian companion animals and horses. *Vet Microbiol*. 2014;168(2–4):381–7.
- [209] Scott GM, Thomson R, Malone-Lee J, Ridgway GL. Cross-infection between animals and man: possible feline transmission of *Staphylococcus aureus* infection in humans? *J Hosp Infect*. 1988;12(1):29–34.
- [210] Strommenger B, Kehrenberg C, Kettlitz C, Cuny C, Verspohl J, Witte W, et al. Molecular characterization of methicillin-resistant *Staphylococcus aureus* strains from pet animals and their relationship to human isolates. *J Antimicrob Chemother*. 2006;57(3):461–5.
- [211] Weese JS, Dick H, Willey BM, McGeer A, Kreiswirth BN, Innis B, et al. Suspected transmission of methicillin-resistant *Staphylococcus aureus* between domestic pets and humans in veterinary clinics and in the household. *Vet Microbiol*. 2006;115(1–3):148–55.
- [212] Malik S, Coombs GW, O'Brien FG, Peng H, Barton MD. Molecular typing of methicillin-resistant staphylococci isolated from cats and dogs. *J Antimicrob Chemother*. 2006;58(2):428–31.
- [213] Grinberg A, Kingsbury DD, Gibson IR, Kirby BM, Mack HJ, Morrison D. Clinically overt infections with methicillin-resistant *Staphylococcus aureus* in animals in New Zealand: a pilot study. *N Z Vet J*. 2008;56(5):237–42.
- [214] Baptiste KE, Williams K, Williams NJ, Wattret A, Clegg PD, Dawson S, et al. Methicillin-resistant staphylococci in companion animals. *Emerg Infect Dis*. 2005;11(12):1942–4.
- [215] Monaco M, Pedroni P, Sanchini A, Bonomini A, Indelicato A, Pantosti A. Livestock-associated methicillin-resistant *Staphylococcus aureus* responsible for human colonization and infection in an area of Italy with high density of pig farming. *BMC Infect Dis*. 2013;13:258.
- [216] Morris DO, Mauldin EA, O'Shea K, Shofer FS, Rankin SC. Clinical, microbiological, and molecular characterization of methicillin-resistant *Staphylococcus aureus* infections of cats. *Am J Vet Res*. 2006;67(8):1421–5.
- [217] Catry B, Van Duijkeren E, Pomba MC, Greko C, Moreno MA, Pyorala S, et al. Reflection paper on MRSA in food-producing and companion animals: epidemiology and control options for human and animal health. *Epidemiol Infect*. 2010;138(5):626–44.
- [218] Maddox TW, Clegg PD, Williams NJ, Pinchbeck GL. Antimicrobial resistance in bacteria from horses: epidemiology of antimicrobial resistance. *Equine Vet J*. 2015;47(6):756–65.
- [219] Tyner GA, Nolen-Walston RD, Hall T, Palmero JP, Couetil L, Javiskas L, et al. A multicenter retrospective study of 151 renal biopsies in horses. *J Vet Intern Med*. 2011;25(3):532–9.

- [220] Weese JS, Rousseau J, Willey BM, Archambault M, McGeer A, Low DE. Methicillin-resistant *Staphylococcus aureus* in horses at a veterinary teaching hospital: frequency, characterization, and association with clinical disease. *J Vet Intern Med*. 2006;20(1):182–6.
- [221] Haenni M, Chatre P, Dupieux C, Metayer V, Maillard K, Bes M, et al. mecC-positive MRSA in horses. *J Antimicrob Chemother*. 2015;70(12):3401–2.
- [222] Cuny C, Strommenger B, Witte W, Stanek C. Clusters of infections in horses with MRSA ST1, ST254, and ST398 in a veterinary hospital. *Microb Drug Resist*. 2008;14(4):307–10.
- [223] Moodley A, Stegger M, Bagcigil AF, Baptiste KE, Loeffler A, Lloyd DH, et al. spa typing of methicillin-resistant *Staphylococcus aureus* isolated from domestic animals and veterinary staff in the UK and Ireland. *J Antimicrob Chemother*. 2006;58(6):1118–23.
- [224] Kock R, Ballhausen B, Bischoff M, Cuny C, Eckmanns T, Fetsch A, et al. The impact of zoonotic MRSA colonization and infection in Germany. *Berl Munch Tierarztl Wochenschr*. 2014;127(9–10):384–98.
- [225] Tokateloff N, Manning ST, Weese JS, Campbell J, Rothenburger J, Stephen C, et al. Prevalence of methicillin-resistant *Staphylococcus aureus* colonization in horses in Saskatchewan, Alberta, and British Columbia. *Can Vet J*. 2009;50(11):1177–80.
- [226] O'Mahony R, Abbott Y, Leonard FC, Markey BK, Quinn PJ, Pollock PJ, et al. Methicillin-resistant *Staphylococcus aureus* (MRSA) isolated from animals and veterinary personnel in Ireland. *Vet Microbiol*. 2005;109(3–4):285–96.
- [227] van Duijkeren E, Moleman M, Sloet van Oldruitenborgh-Oosterbaan MM, Multem J, Troelstra A, Fluit AC, et al. Methicillin-resistant *Staphylococcus aureus* in horses and horse personnel: an investigation of several outbreaks. *Vet Microbiol*. 2010;141(1–2):96–102.
- [228] Axon JE, Carrick JB, Barton MD, Collins NM, Russell CM, Kiehne J, et al. Methicillin-resistant *Staphylococcus aureus* in a population of horses in Australia. *Aust Vet J*. 2011;89(6):221–5.
- [229] Van den Eede A, Martens A, Lipinska U, Struelens M, Deplano A, Denis O, et al. High occurrence of methicillin-resistant *Staphylococcus aureus* ST398 in equine nasal samples. *Vet Microbiol*. 2009;133(1–2):138–44.
- [230] Maddox TW, Clegg PD, Diggle PJ, Wedley AL, Dawson S, Pinchbeck GL, et al. Cross-sectional study of antimicrobial-resistant bacteria in horses. Part 1: prevalence of antimicrobial-resistant *Escherichia coli* and methicillin-resistant *Staphylococcus aureus*. *Equine Vet J*. 2012;44(3):289–96.
- [231] Vengust M, Anderson ME, Rousseau J, Weese JS. Methicillin-resistant staphylococcal colonization in clinically normal dogs and horses in the community. *Lett Appl Microbiol*. 2006;43(6):602–6.

- [232] Springer B, Orendi U, Much P, Hoger G, Ruppitsch W, Krziwanek K, et al. Methicillin-resistant *Staphylococcus aureus*: a new zoonotic agent? Wien Klin Wochenschr. 2009;121(3–4):86–90.
- [233] Burton S, Reid-Smith R, McClure JT, Weese JS. *Staphylococcus aureus* colonization in healthy horses in Atlantic Canada. Can Vet J. 2008;49(8):797–9.
- [234] Morcillo A, Castro B, Rodriguez-Alvarez C, Abreu R, Aguirre-Jaime A, Arias A. Descriptive analysis of antibiotic-resistant patterns of methicillin-resistant *Staphylococcus aureus* (MRSA) st398 isolated from healthy swine. Int J Environ Res Public Health. 2015;12(1):611–22.
- [235] Mammina C, Cala C, Plano MR, Bonura C, Vella A, Monastero R, et al. Ventilator-associated pneumonia and MRSA ST398, Italy. Emerg Infect Dis. 2010;16(4):730–1.
- [236] Morales G, Picazo JJ, Baos E, Candel FJ, Arribi A, Pelaez B, et al. Resistance to linezolid is mediated by the cfr gene in the first report of an outbreak of linezolid-resistant *Staphylococcus aureus*. Clin Infect Dis. 2010;50(6):821–5.
- [237] Kehrenberg C, Cuny C, Strommenger B, Schwarz S, Witte W. Methicillin-resistant and -susceptible *Staphylococcus aureus* strains of clonal lineages ST398 and ST9 from swine carry the multidrug resistance gene cfr. Antimicrob Agents Chemother. 2009;53(2):779–81.
- [238] Gomez-Sanz E, Kadlec K, Fessler AT, Zarazaga M, Torres C, Schwarz S. A novel fexA variant from a canine *Staphylococcus pseudintermedius* isolate that does not confer florfenicol resistance. Antimicrob Agents Chemother. 2013;57(11):5763–6.
- [239] Toh SM, Xiong L, Arias CA, Villegas MV, Lolans K, Quinn J, et al. Acquisition of a natural resistance gene renders a clinical strain of methicillin-resistant *Staphylococcus aureus* resistant to the synthetic antibiotic linezolid. Mol Microbiol. 2007;64(6):1506–14.
- [240] Mendes RE, Deshpande LM, Castanheira M, DiPersio J, Saubolle MA, Jones RN. First report of cfr-mediated resistance to linezolid in human staphylococcal clinical isolates recovered in the United States. Antimicrob Agents Chemother. 2008;52(6):2244–6.
- [241] Kehrenberg C, Schwarz S, Jacobsen L, Hansen LH, Vester B. A new mechanism for chloramphenicol, florfenicol and clindamycin resistance: methylation of 23S ribosomal RNA at A2503. Mol Microbiol. 2005;57(4):1064–73.
- [242] Long KS, Poehlsgaard J, Kehrenberg C, Schwarz S, Vester B. The Cfr rRNA methyltransferase confers resistance to phenicols, lincosamides, oxazolidinones, pleuromutilins, and streptogramin A antibiotics. Antimicrob Agents Chemother. 2006;50(7):2500–5.
- [243] Slifierz MJ, Friendship RM, Weese JS. Methicillin-resistant *Staphylococcus aureus* in commercial swine herds is associated with disinfectant and zinc usage. Appl Environ Microbiol. 2015;81(8):2690–5.

- [244] Schwarz S, Cardoso M, Wegener HC. Nucleotide sequence and phylogeny of the tet(L) tetracycline resistance determinant encoded by plasmid pSTE1 from *Staphylococcus hyicus*. Antimicrob Agents Chemother. 1992;36(3):580–8.
- [245] Fessler AT, Kadlec K, Hassel M, Hauschild T, Eidam C, Ehrlich R, et al. Characterization of methicillin-resistant *Staphylococcus aureus* isolates from food and food products of poultry origin in Germany. Appl Environ Microbiol. 2011;77(20):7151–7.
- [246] Argudin MA, Tenhagen BA, Fetsch A, Sachsenroder J, Kasbohrer A, Schroeter A, et al. Virulence and resistance determinants of German *Staphylococcus aureus* ST398 isolates from nonhuman sources. Appl Environ Microbiol. 2011;77(9):3052–60.
- [247] Kadlec K, Schwarz S. Identification of a plasmid-borne resistance gene cluster comprising the resistance genes erm(T), dfrK, and tet(L) in a porcine methicillin-resistant *Staphylococcus aureus* ST398 strain. Antimicrob Agents Chemother. 2010;54(2):915–8.
- [248] Schwendener S, Perreten V. New transposon Tn6133 in methicillin-resistant *Staphylococcus aureus* ST398 contains vga(E), a novel streptogramin A, pleuromutilin, and lincosamide resistance gene. Antimicrob Agents Chemother. 2011;55(10):4900–4.
- [249] Kadlec K, Schwarz S. Novel ABC transporter gene, vga(C), located on a multiresistance plasmid from a porcine methicillin-resistant *Staphylococcus aureus* ST398 strain. Antimicrob Agents Chemother. 2009;53(8):3589–91.
- [250] Kadlec K, Pomba CF, Couto N, Schwarz S. Small plasmids carrying vga(A) or vga(C) genes mediate resistance to lincosamides, pleuromutilins and streptogramin A antibiotics in methicillin-resistant *Staphylococcus aureus* ST398 from swine. J Antimicrob Chemother. 2010;65(12):2692–3.
- [251] Hauschild T, Fessler AT, Kadlec K, Billerbeck C, Schwarz S. Detection of the novel vga(E) gene in methicillin-resistant *Staphylococcus aureus* CC398 isolates from cattle and poultry. J Antimicrob Chemother. 2012;67(2):503–4.
- [252] Fessler AT, Kadlec K, Schwarz S. Novel apramycin resistance gene apmA in bovine and porcine methicillin-resistant *Staphylococcus aureus* ST398 isolates. Antimicrob Agents Chemother. 2011;55(1):373–5.
- [253] Kadlec K, Fessler AT, Couto N, Pomba CF, Schwarz S. Unusual small plasmids carrying the novel resistance genes dfrK or apmA isolated from methicillin-resistant or -susceptible staphylococci. J Antimicrob Chemother. 2012;67(10):2342–5.
- [254] Seier-Petersen MA, Nielsen LN, Ingmer H, Aarestrup FM, Agerso Y. Biocide susceptibility of *Staphylococcus aureus* CC398 and CC30 isolates from pigs and identification of the biocide resistance genes, qacG and qacC. Microb Drug Resist. 2015;21(5):527–36.
- [255] Gomez-Sanz E, Torres C, Lozano C, Fernandez-Perez R, Aspiroz C, Ruiz-Larrea F, et al. Detection, molecular characterization, and clonal diversity of methicillin-resistant *Staphylococcus aureus* CC398 and CC97 in Spanish slaughter pigs of different age groups. Foodborne Pathog Dis. 2010;7(10):1269–77.

- [256] Verheghe M, Pletinckx LJ, Crombe F, Vandersmissen T, Haesebrouck F, Butaye P, et al. Methicillin-resistant *Staphylococcus aureus* (MRSA) ST398 in pig farms and multispecies farms. *Zoonoses Public Health*. 2013;60(5):366–74.
- [257] Silva NC, Guimaraes FF, Manzi MP, Junior AF, Gomez-Sanz E, Gomez P, et al. Methicillin-resistant *Staphylococcus aureus* of lineage ST398 as cause of mastitis in cows. *Lett Appl Microbiol*. 2014;59(6):665–9.
- [258] Anukool U, O'Neill CE, Butr-Indr B, Hawkey PM, Gaze WH, Wellington EM. Methicillin-resistant *Staphylococcus aureus* in pigs from Thailand. *Int J Antimicrob Agents*. 2011;38(1):86–7.
- [259] Li D, Wu C, Wang Y, Fan R, Schwarz S, Zhang S. Identification of multiresistance gene *cfr* in methicillin-resistant *Staphylococcus aureus* from pigs: plasmid location and integration into a staphylococcal cassette chromosome *mec* complex. *Antimicrob Agents Chemother*. 2015;59(6):3641–4.
- [260] He W, Liu Y, Qi J, Chen H, Zhao C, Zhang F, et al. Food-animal related *Staphylococcus aureus* multidrug-resistant ST9 strains with toxin genes. *Foodborne Pathog Dis*. 2013;10(9):782–8.
- [261] Chao G, Zhang X, Zhang X, Huang Y, Xu L, Zhou L, et al. Phenotypic and genotypic characterization of methicillin-resistant *Staphylococcus aureus* (MRSA) and methicillin-susceptible *Staphylococcus aureus* (MSSA) from different sources in China. *Foodborne Pathog Dis*. 2013;10(3):214–21.
- [262] Yan X, Yu X, Tao X, Zhang J, Zhang B, Dong R, et al. *Staphylococcus aureus* ST398 from slaughter pigs in northeast China. *Int J Med Microbiol*. 2014;304(3–4):379–83.
- [263] Ho J, O'Donoghue M, Guardabassi L, Moodley A, Boost M. Characterization of methicillin-resistant *Staphylococcus aureus* isolates from pig carcasses in Hong Kong. *Zoonoses Public Health*. 2012;59(6):416–23.
- [264] Brennan GI, Abbott Y, Burns A, Leonard F, McManus BA, O'Connell B, et al. The emergence and spread of multiple livestock-associated clonal complex 398 methicillin-resistant and methicillin-susceptible *Staphylococcus aureus* strains among animals and humans in the Republic of Ireland, 2010–2014. *PLoS One*. 2016;11(2):e0149396.
- [265] Franco A, Hasman H, Iurescia M, Lorenzetti R, Stegger M, Pantosti A, et al. Molecular characterization of *spa* type t127, sequence type 1 methicillin-resistant *Staphylococcus aureus* from pigs. *J Antimicrob Chemother*. 2011;66(6):1231–5.
- [266] Tavakol M, Riekerink RG, Sampimon OC, van Wamel WJ, van Belkum A, Lam TJ. Bovine-associated MRSA ST398 in the Netherlands. *Acta Vet Scand*. 2012;54:28.
- [267] Porrero MC, Wassenaar TM, Gomez-Barrero S, Garcia M, Barcena C, Alvarez J, et al. Detection of methicillin-resistant *Staphylococcus aureus* in Iberian pigs. *Lett Appl Microbiol*. 2012;54(4):280–5.

- [268] Lo YP, Wan MT, Chen MM, Su HY, Lauderdale TL, Chou CC. Molecular characterization and clonal genetic diversity of methicillin-resistant *Staphylococcus aureus* of pig origin in Taiwan. *Comp Immunol Microbiol Infect Dis*. 2012;35(6):513–21.
- [269] Vestergaard M, Cavaco LM, Sirichote P, Unahalekhaka A, Dangsakul W, Svendsen CA, et al. SCCmec type IX element in methicillin resistant *Staphylococcus aureus* spa Type t337 (CC9) isolated from pigs and pork in Thailand. *Front Microbiol*. 2012;3:103.
- [270] Van den Eede A, Hermans K, Van den Abeele A, Flore K, Dewulf J, Vanderhaeghen W, et al. The nasal vestibulum is the optimal sampling site for MRSA screening in hospitalised horses. *Vet J*. 2013;197(2):415–9.
- [271] Rubin JE, Chirino-Trejo M. Antimicrobial susceptibility of canine and human *Staphylococcus aureus* collected in Saskatoon, Canada. *Zoonoses Public Health*. 2011;58(7):454–62.
- [272] Anderson ME, Lefebvre SL, Rankin SC, Aceto H, Morley PS, Caron JP, et al. Retrospective multicentre study of methicillin-resistant *Staphylococcus aureus* infections in 115 horses. *Equine Vet J*. 2009;41(4):401–5.
- [273] Zhang W, Hao Z, Wang Y, Cao X, Logue CM, Wang B, et al. Molecular characterization of methicillin-resistant *Staphylococcus aureus* strains from pet animals and veterinary staff in China. *Vet J*. 2011;190(2):e125–9.
- [274] Vincze S, Stamm I, Kopp PA, Hermes J, Adlhoeh C, Semmler T, et al. Alarming proportions of methicillin-resistant *Staphylococcus aureus* (MRSA) in wound samples from companion animals, Germany 2010–2012. *PLoS One*. 2014;9(1):e85656.
- [275] Boost MV, O'Donoghue MM, Siu KH. Characterisation of methicillin-resistant *Staphylococcus aureus* isolates from dogs and their owners. *Clin Microbiol Infect*. 2007;13(7):731–3.
- [276] Leonard FC, Abbott Y, Rossney A, Quinn PJ, O'Mahony R, Markey BK. Methicillin-resistant *Staphylococcus aureus* isolated from a veterinary surgeon and five dogs in one practice. *Vet Rec*. 2006;158(5):155–9.
- [277] Schwaber MJ, Navon-Venezia S, Masarwa S, Tirosh-Levy S, Adler A, Chmelnitsky I, et al. Clonal transmission of a rare methicillin-resistant *Staphylococcus aureus* genotype between horses and staff at a veterinary teaching hospital. *Vet Microbiol*. 2013;162(2–4):907–11.
- [278] Tirosh-Levy S, Steinman A, Carmeli Y, Klement E, Navon-Venezia S. Prevalence and risk factors for colonization with methicillin resistant *Staphylococcus aureus* and other *Staphylococci* species in hospitalized and farm horses in Israel. *Prev Vet Med*. 2015;122(1–2):135–44.
- [279] Ishihara K, Saito M, Shimokubo N, Muramatsu Y, Maetani S, Tamura Y. Methicillin-resistant *Staphylococcus aureus* carriage among veterinary staff and dogs in private veterinary clinics in Hokkaido, Japan. *Microbiol Immunol*. 2014;58(3):149–54.

- [280] Tajima K, Sinjyo A, Ito T, Noda Y, Goto H, Ito N. Methicillin-resistant *Staphylococcus aureus* keratitis in a dog. *Vet Ophthalmol.* 2013;16(3):240–3.
- [281] Jang Y, Bae D, Cho JK, Bahk GJ, Lim SK, Lee YJ. Characterization of methicillin-resistant *Staphylococcus* spp. isolated from dogs in Korea. *Jpn J Vet Res.* 2014;62(4):163–70.
- [282] Pak SI, Han HR, Shimizu A. Characterization of methicillin-resistant *Staphylococcus aureus* isolated from dogs in Korea. *J Vet Med Sci.* 1999;61(9):1013–8.
- [283] Aklilu E, Zakaria Z, Hassan L, Hui Cheng C. Molecular relatedness of methicillin-resistant *S. aureus* isolates from staff, environment and pets at University Veterinary Hospital in Malaysia. *PLoS One.* 2012;7(8):e43329.
- [284] van Duijkeren E, Box AT, Heck ME, Wannet WJ, Fluit AC. Methicillin-resistant staphylococci isolated from animals. *Vet Microbiol.* 2004;103(1–2):91–7.
- [285] Mustapha M, Bukar-Kolo YM, Geidam YA, Gulani IA. Phenotypic and genotypic detection of methicillin-resistant *Staphylococcus aureus* in hunting dogs in Maiduguri metropolitan, Borno State, Nigeria. *Vet World.* 2016;9(5):501–6.
- [286] Coelho C, Torres C, Radhouani H, Pinto L, Lozano C, Gomez-Sanz E, et al. Molecular detection and characterization of methicillin-resistant *Staphylococcus aureus* (MRSA) isolates from dogs in Portugal. *Microb Drug Resist.* 2011;17(2):333–7.
- [287] Couto N, Belas A, Tilley P, Couto I, Gama LT, Kadlec K, et al. Biocide and antimicrobial susceptibility of methicillin-resistant staphylococcal isolates from horses. *Vet Microbiol.* 2013;166(1–2):299–303.
- [288] Sieber S, Gerber V, Jandova V, Rossano A, Evison JM, Perreten V. Evolution of multi-drug-resistant *Staphylococcus aureus* infections in horses and colonized personnel in an equine clinic between 2005 and 2010. *Microb Drug Resist.* 2011;17(3):471–8.
- [289] Davis JA, Jackson CR, Fedorka-Cray PJ, Barrett JB, Brousse JH, Gustafson J, et al. Carriage of methicillin-resistant staphylococci by healthy companion animals in the US. *Lett Appl Microbiol.* 2014;59(1):1–8.
- [290] van Balen J, Kelley C, Nava-Hoet RC, Bateman S, Hillier A, Dyce J, et al. Presence, distribution, and molecular epidemiology of methicillin-resistant *Staphylococcus aureus* in a small animal teaching hospital: a year-long active surveillance targeting dogs and their environment. *Vector Borne Zoonotic Dis.* 2013;13(5):299–311.
- [291] Coughlan K, Olsen KE, Boxrud D, Bender JB. Methicillin-resistant *Staphylococcus aureus* in resident animals of a long-term care facility. *Zoonoses Public Health.* 2010;57(3):220–6.
- [292] Morris DO, Rook KA, Shofer FS, Rankin SC. Screening of *Staphylococcus aureus*, *Staphylococcus intermedius*, and *Staphylococcus schleiferi* isolates obtained from small companion animals for antimicrobial resistance: a retrospective review of 749 isolates (2003–04). *Vet Dermatol.* 2006;17(5):332–7.

- [293] Rankin S, Roberts S, O'Shea K, Maloney D, Lorenzo M, Benson CE. Panton valentine leukocidin (PVL) toxin positive MRSA strains isolated from companion animals. *Vet Microbiol.* 2005;108(1-2):145-8.
- [294] Tomlin J, Pead MJ, Lloyd DH, Howell S, Hartmann F, Jackson HA, et al. Methicillin-resistant *Staphylococcus aureus* infections in 11 dogs. *Vet Rec.* 1999;144(3):60-4.

IntechOpen

IntechOpen