

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

185,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



---

# Effects of Drought and Elevated Atmospheric Carbon Dioxide on Seed Nutrition and $^{15}\text{N}$ and $^{13}\text{C}$ Natural Abundance Isotopes in Soybean Under Controlled Environments

---

Nacer Bellaloui, Alemu Mengistu,  
Hamed K. Abbas and My A. Kassem

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/67511>

---

## Abstract

The objective of the current research was to evaluate the effects of drought and elevated  $\text{CO}_2$  on seed production and seed nutrition under controlled conditions in soybean. Soybean plants were subjected to ambient and elevated  $\text{CO}_2$  and under irrigated and drought conditions. The results showed that drought or drought with elevated  $\text{CO}_2$  resulted in high protein and oleic acid, but low in oil and linoleic and linolenic acids. Significant decrease of sucrose, glucose, and fructose concentrations was noticed, but high content of raffinose and stachyose was observed. Nutrients such as N, P, K, and some micro-nutrients were reduced under drought or drought with normal or elevated  $\text{CO}_2$  concentrations. Seed  $\delta^{15}\text{N}$  ( $^{15}\text{N}/^{14}\text{N}$  ratio) and  $\delta^{13}\text{C}$  ( $^{13}\text{C}/^{12}\text{C}$  ratio) natural abundance isotopes were also altered under drought or drought with ambient or elevated  $\text{CO}_2$  concentrations, reflecting nitrogen and carbon metabolism changes. The current research demonstrated that global climate changes may lead to changes in seed nutrition, and nitrogen and carbon metabolism. Efforts of breeders to select for these traits will sustain food source and food security for humans and livestock as soybean is a major source for protein and oil for human consumption and soy meal for animals.

**Keywords:** elevated carbon dioxide, climate change, seed composition, seed nutrition, drought, soybean

---

## 1. Introduction

Global climate changes due to elevated CO<sub>2</sub> are expected to lead to high heat and drought in some regions, affecting crop production and seed nutrition [1]. Climate change, whether is caused naturally or human-made, threatens crop production and food security. It was reported that food systems that include food availability, food access, and food utilization will suffer when food systems are stressed [1]. It is predicted that climate change will alter the food systems through its effects on crop production and rainfall, leading to drought or flooding or warmer or cooler temperatures [1]. Global warming resulted from climate change leads to high risk of drought and warmer temperatures, resulting in an increase of water demands and evaporation (<http://www.c2es.org/science-impacts/extreme-weather/drought>) [2]. The USA historically has suffered major droughts, and the living memories of the Dust Bowl of the 1930s or the drought of the 1950s and recently the droughts of 2011 in Texas and 2012 in the USA are examples, highlighting our vulnerabilities to drought as we move forward to warmer and drier climate (<http://www.c2es.org/science-impacts/extreme-weather/drought>) [2]. It was also reported that a doubling of CO<sub>2</sub> concentration could result in changes in global climate, including precipitation patterns, resulting in drought conditions in some areas of the world [3]. Generally, elevated CO<sub>2</sub> concentration during crop growth enhances CO<sub>2</sub> exchange rate and final yield [4] as growth at elevated CO<sub>2</sub> stimulates photosynthesis and increases carbon supply in C<sub>3</sub> crops and enhances nutrient supply to match the increase in C acquisition [4]. However, drought stress results in a decrease in CO<sub>2</sub> exchange rate. The mechanisms involved in the reduction of CO<sub>2</sub> exchange rate are still not well understood, although part of the reduction was attributed to stomatal closure [5], leaf water potentials [6], decreases in activities of Photosystem I and Photosystem II [7–9], effects on assimilation products, and photosynthetic enzyme activities [6].

The effects of climate changes on yield were previously reported [1], and it was shown that severe drought stress can lead to crop losses of up to 90% [10–12]. While there has been considerable progress in understanding the sensitivities of crop yield to climate change, evaluation of climate change factors such as CO<sub>2</sub>, elevated temperature, or drought on seed nutrition remains rather limited [13]. Drought resulted from climate change is expected to alter the biochemistry and physiology of crops, impacting the nutritional value of the crop grains. Alteration of carbohydrates including starch and soluble sugars such as glucose, fructose, and sucrose occurs under stress conditions [14]. It is well known that carbohydrates are the major product of photosynthesis, which are transported to stems and leaves (important reserve of photo-assimilates for grain filling) and consequently to the sink (grains) during grain-filling stage [15]. The sugars, sucrose and raffinose, are also known to protect cells against oxidative damage and accumulate later during responses to stress [16, 17]. It was reported that the high levels of raffinose and stachyose were thought to play a role in the acquisition of desiccation tolerance due to over expression of galactinol synthase and accumulation of galactinol and raffinose improving drought tolerance [18, 19]. The mechanisms of involvement of these sugars in drought tolerance are still not fully understood [19–21]. The accumulation of proline, amino acids, and organic acid such as malic acid, fumaric acid, and citric acid is also a common biochemical indicator occurring at high level under drought stress [12]. The accumulation of

these compounds is a part of the mechanism of osmotic adjustment to maintain water potential gradient under drought conditions and to protect subcellular structure from the damaging effects of drought stress [22]. The high level of proline accumulation under drought stress was previously identified as a mechanism for the protection of cytosolic proteins and organelles [12]. Therefore, these biochemical compounds are important to understand the relationship between drought stress and drought tolerance and for decision-making when breeding for drought tolerance [23]. Since drought stress results in limiting the growth and development of the plant and consequently the supply of the photosynthate to grain during grain filling [24], the chemical composition of the grain and its quality can be affected by drought stress [12, 25]. The effect of drought on seed quality could be due to the decrease of total soluble sugars especially in the grain under drought stress due to decreases in the levels of sucrose, stachyose, and verbascose. Sucrose plays an important role in the development of the grain and is sensitive to drought stress and involved in the synthesis of raffinose oligosaccharides including stachyose and verbascose, and these sugars play an important role in seed tolerance to desiccation and against oxidative effects of drought stress [18, 26]. Some sugars, for example, maltose, accumulate in the grain under drought stress, maybe due to starch degradation [12]. For example, during the remobilization phase, starch remobilized was significantly lower under drought stress, but the total amounts of soluble sugars and amino acids remobilized were greater [12], contrasting previous studies that showed higher carbohydrate accumulation under drought stress [27]. Therefore, the role of carbohydrates in drought tolerance is still not well understood, and further investigations are needed to reach the conclusive results.

Research on the possible effects of elevated CO<sub>2</sub> and drought resulted from climate changes on biochemical and chemical composition and mineral nutrition of major crops is limited. We chose soybean as a subject of our research because soybean is a major crop in the world; it is a major source of human nutrition because it contains protein (40%), oil (20%), fatty acids, amino acids, carbohydrates (30%), crude fiber (5%), and ash (5%) and it contains minerals (such as P, K, Ca, Mg, Fe, Cu, Mn, Zn, and Mo), vitamins (B1, B2, and B6), phytoestrogen, such as isoflavones, and phenolics. The objective of the current research was to investigate the possible effects of elevated CO<sub>2</sub> and drought on seed chemical composition (protein, oil, fatty acids, sugars, and minerals). A special attention was given to possible alteration of seed  $\delta^{15}\text{N}$  ( $^{15}\text{N}/^{14}\text{N}$  ratio) and  $\delta^{13}\text{C}$  ( $^{13}\text{C}/^{12}\text{C}$  ratio) natural abundance isotopes as they reflect nitrogen and carbon metabolism.

## 2. Materials and methods

### 2.1. Growth conditions

The experiments were conducted under controlled environments of growth chambers. Soybean seeds were planted in vermiculite and grown under greenhouse conditions until V1 stage where similar plants were transported to pots filled with field soil. The soil texture was 8% sand, 31.6% silt, and 60.4% clay. Soil nutrient concentrations of macro- and

micronutrients were adequate to support the growth and development of plants till maturity. When plants reached R5 (beginning of seed-fill stage) [28], they were transferred to the growth chambers until full maturity. Two soybean cultivars, Freedom and Hutcheson of maturity group V, were used. The plants were subjected to the following four treatments (T): T1 = plants were grown irrigated and subjected to  $360 \mu\text{mol mol}^{-1} \text{CO}_2$  concentration; T2 = plants were grown irrigated and subjected to  $700 \mu\text{mol mol}^{-1} \text{CO}_2$  concentration; T3 = plants were subjected to drought and to  $360 \mu\text{mol mol}^{-1} \text{CO}_2$  concentration; and T4 = plants were subjected to drought and to  $700 \mu\text{mol mol}^{-1} \text{CO}_2$  concentration. Drought stress treatment was imposed by growing the plants at soil water potential of about  $-199 \text{ kPa}$ . For irrigated plants, soil water potential was kept at about  $-15$  to  $-20 \text{ kPa}$ , which was considered to give the field water holding capacity. Soil water potential was monitored by using soil water potential sensors and read by Soil Moisture Meter (Watermark Company, Inc., Wisconsin, USA). For irrigated experiment, the plants were watered as needed. For  $\text{CO}_2$  treatment,  $\text{CO}_2$  concentration was supplied by  $\text{CO}_2$  cylinder located outside the growth chamber, and the rate of  $\text{CO}_2$  was controlled by a regulator, monitoring the  $\text{CO}_2$  concentration flow. The  $\text{CO}_2$  flowed through the tube to the growth chamber. The growth chamber was equipped with  $\text{CO}_2$  sensor to read the concentration of  $\text{CO}_2$  inside the growth chamber. The plants were grown under growth chamber conditions supplied with light source of photon flux density of about  $1000 \mu\text{mol m}^{-2} \text{s}^{-1}$  and provided with a combination of 10,400 W high-pressure sodium and metal halide lights. The experiments were conducted under constant normal temperature of  $26/16^\circ\text{C}$ , day/night.

## 2.2. Determination of seed minerals, N, S, and C

Mature seeds were ground using a Laboratory Mill 3600 (Perten, Springfield, IL, USA), and the ground samples were analyzed for minerals, N, S, and C, by digesting 0.6 g of the ground-dried plant materials in  $\text{HNO}_3$  in a microwave digestion system as detailed elsewhere [29, 30]. Potassium was determined by inductively coupled plasma spectrometry [29, 30]. Each content of N, C, and S was measured in a 0.25 g ground-dried sample which was combusted in atmospheric oxygen of  $1350^\circ\text{C}$ , and then the elements N, S, and C were converted to  $\text{N}_2$ ,  $\text{SO}_2$ , and  $\text{CO}_2$ , respectively. The contents of N, S, and C in seed were measured by an elemental analyzer using thermal conductivity cells (LECO CNS-2000 elemental analyzer, LECO Corporation, St. Joseph, MI, USA) [29, 30].

## 2.3. Determinations of seed protein, oil and fatty acids

The mature seed samples of 25 g were ground using the Laboratory Mill 3600, and seed protein, oil, and fatty acids in the samples were analyzed by near-infrared reflectance [29–31] using a diode array feed analyzer AD 7200 (Perten, Springfield, IL USA). The calibration equation was developed using Preteen's Thermo Galactic Grams PLS IQ software, and the calibration curve was established using Association Official Analytical Chemists (AOAC) methods. The contents of protein and oil were determined based on a seed dry matter [29, 31]. The contents of palmitic, stearic, oleic, linoleic, and linolenic fatty acids were determined based on a total oil basis [29].



## 2.4. Determinations of sucrose, raffinose, stachyose, glucose, and fructose

Seed sucrose, raffinose, and stachyose were determined as detailed elsewhere (Bellaloui et al. [29]) by near-infrared reflectance using the AD 7200 array feed analyzer. Sugars were determined based on a seed dry matter basis [29–31]. The concentration of glucose in mature seed was determined by an enzymatic reaction using a Glucose (HK) Assay Kit, Product Code GAHK-20 (Sigma-Aldrich Co., St Louis, MO, USA) as detailed elsewhere [29]. The concentrations were determined spectrophotometrically by reading the samples at absorbance of 340 nm using the Beckman Coulter DU 800 spectrophotometer. The concentration was expressed as  $\text{mg g}^{-1}$  dry weight. Fructose concentration in mature seed was determined based on an enzymatic reaction using a Fructose Assay Kit, Product Code FA-20 (Sigma-Aldrich Co., St. Louis, MO, USA) as detailed by Bellaloui et al. elsewhere [29]. Fructose concentration was determined spectrophotometrically by the Beckman Coulter DU 800 spectrophotometer by reading the samples at absorbance of 340 nm. Fructose concentration in seed was expressed as  $\text{mg g}^{-1}$  dry weight.

## 2.5. Boron determination

Boron concentrations in mature seeds were determined using the Azomethine-H method [32, 33] as reported elsewhere [29]. Briefly, 1.0 g of ground sample was ashed at  $500^{\circ}\text{C}$ , extracted with 20 ml of 2 M HCl at  $90^{\circ}\text{C}$  for 10 min, and then 4 ml of a buffer solution (containing 25% ammonium acetate, 1.5% EDTA, and 12.5% acetic acid) was added to a filtered 2-ml sample. An amount of 4 ml of fresh azomethine-H solution (0.45% azomethine-H and 1% of ascorbic acid) [34] was added. Boron concentration was determined spectrophotometrically by reading the samples at absorbance of 420 nm using a Beckman Coulter DU 800 spectrophotometer (Beckman Coulter, Inc., Brea, CA, USA). The concentration was expressed as  $\text{mg kg}^{-1}$  dry weight.

## 2.6. Iron determination

Iron concentration in mature seed was determined according to the method described elsewhere [35, 36]. Briefly, 2 g of dry ground samples were acid wet digested, extracted, and reacted with reduced ferrous Fe with 1,10-phenanthroline and as detailed elsewhere [29]. After the extraction, the soluble constituents were dissolved in 2 M HCl, and the phenanthroline solution of 0.25% (w/v) in 25% (v/v) ethanol and quinol solution (1% w/v) was used as a reagent. Standard curve of Fe was created by preparing a range of concentrations from 0.0 to  $4 \mu\text{g ml}^{-1}$  of Fe in 0.4 M HCl. Iron concentration was determined spectrophotometrically by reading the samples at absorbance of 510 nm using the Beckman Coulter DU 800 spectrophotometer. The concentration was expressed as  $\text{mg kg}^{-1}$  dry weight.

## 2.7. Phosphorus determination

Phosphorus content in mature seeds was measured according to [37]. Phosphorus determination was based on the yellow phosphor-vanado-molybdate complex as detailed elsewhere [29]. Briefly, dry ground seed samples of 2 g were ashed. Then, 10 ml of 6M HCl was added,

and phosphorus was extracted using 2 ml of 36% v/v HCl under heat and filtration and 5 ml of 5 M HCl. A volume of 5 ml of reagent (ammonium molybdate-ammonium metavanadate) was added to 5 ml of the filtrate. The reagent ammonium molybdate-ammonium metavanadate was prepared by dissolving 25 g of ammonium molybdate and 1.25 g of ammonium metavanadate in distilled water. The standard curve solutions of P were prepared by using a range of concentrations from 0 to 50  $\mu\text{g ml}^{-1}$  using dihydrogen orthophosphates. The concentration of P was determined spectrophotometrically by reading the absorbance at 400 nm using the Beckman Coulter DU 800 spectrophotometer.

## 2.8. Experimental design and statistical analysis

Two experiments were carried out in a split plot design with arrangement of treatments in a randomized complete block design (RCBD) with four replicates. Main plot was drought/ $\text{CO}_2$  treatment and subplot was cultivar. Each experiment was considered as a replicate for the main plot. One growth chamber was used for each drought/ $\text{CO}_2$  treatment. Treatments in each experiment were the following: T1 = plants were grown irrigated and subjected to 360  $\mu\text{mol mol}^{-1}$   $\text{CO}_2$  concentration; T2 = plants were grown irrigated and subjected to 700  $\mu\text{mol mol}^{-1}$   $\text{CO}_2$  concentration; T3 = plants were grown under drought and subjected to 360  $\mu\text{mol mol}^{-1}$   $\text{CO}_2$  concentration; and T4 = plants were grown under drought and subjected to 700  $\mu\text{mol mol}^{-1}$   $\text{CO}_2$  concentration. Two soybean cultivars were used. The analysis of variance of data was conducted using PROC MIXED in SAS [38]. Experiment (EXP), drought,  $\text{CO}_2$  concentration ( $\text{CO}_2$ ), and their interactions were considered fixed effects, and replication within EXP and replication  $\times$  drought  $\times$   $\text{CO}_2$   $\times$  CV within EXP were considered random effects. The level of significance was  $P \leq 0.05$ . Detailed design of the current experiments was similar to that previously reported by Bellaloui et al. [39].

## 3. Results and discussion

Analysis of variance showed that the main effects of experiment (EXP), drought,  $\text{CO}_2$ , and cultivar (CV) were the most significant factors affecting variability of seed protein, oil, fatty acids, and sugars (**Table 1**). There were no significant effects of the interactions between EXP and other factors, which were expected. Interactions between  $\text{CO}_2$ , CV, and drought showed significant effects for some seed constituents such as protein, oleic, glucose, fructose, and sucrose, indicating the different responses of cultivars to drought and  $\text{CO}_2$  (**Table 1**). Since there were no significant interaction effects between EXP and the others factors, the results were combined across the two experiments. Analysis of variance showed that drought,  $\text{CO}_2$ , and CV were the major factors affecting the variability for macro- and micronutrients in seed (**Table 2**). There were no effects of EXP on nutrients, as expected, because the experiments were conducted under controlled environmental conditions of growth chambers. Both  $\text{CO}_2$  and CV significantly interacted with drought, indicating that the effect of  $\text{CO}_2$  was influenced by drought and the cultivars responded differently due to genotype and genetic background effect (**Table 2**).

Treatments	Protein	Oil	Oleic	Linolenic	Glucose	Fructose	Sucrose	Raffinose	Stachyose
EXP	ns <sup>1</sup>	ns	ns	ns	ns	ns	ns	ns	ns
Drought <sup>2</sup>	*	*	**	**	***	**	**	*	*
CO <sub>2</sub>	**	*	*	*	***	**	***	*	*
CV	*	ns	*	ns	ns	*	*	ns	ns
EXP × drought	ns	ns	ns	ns	ns	ns	ns	ns	ns
EXP × CO <sub>2</sub>	ns	ns	ns	ns	ns	ns	ns	ns	ns
EXP × CV	ns	ns	ns	ns	ns	ns	ns	ns	ns
Drought × CO <sub>2</sub>	ns	ns	ns	ns	ns	ns	ns	ns	ns
Drought × CV	ns	ns	*	ns	*	*	***	ns	ns
CO <sub>2</sub> × CV	*	ns	ns	ns	ns	*	*	ns	ns
EXP × drought × CO <sub>2</sub> × CV	ns	ns	ns	ns	ns	ns	ns	ns	ns

Treatments used were four: T1 = plants were grown irrigated and subjected to 360  $\mu\text{mol mol}^{-1}$  CO<sub>2</sub> concentration; T2 = plants were grown irrigated and subjected to 700  $\mu\text{mol mol}^{-1}$  CO<sub>2</sub> concentration; T3 = plants were grown under drought and subjected to 360  $\mu\text{mol mol}^{-1}$  CO<sub>2</sub> concentration; and T4 = plants were grown under drought and subjected to 700  $\mu\text{mol mol}^{-1}$  CO<sub>2</sub> concentration. Plants were grown in the greenhouse, and they were transferred to growth chambers at the beginning of seed-fill stage (R5) until full maturity (R8). The experiments were conducted under constant normal temperature of 26/16°C, day/night.

<sup>1</sup> \* = significance at  $P \leq 0.05$ ; \*\* = significance at  $P \leq 0.01$ ; \*\*\* = significance at  $P \leq 0.001$ ; ns = not significant.

<sup>2</sup> Drought treatment was imposed by growing the plants at soil water potential of about -199 kPa. For irrigated plant, soil water potential was kept at about -15 to -20 kPa.

**Table 1.** Analysis of variance for soybean cultivars (Freedom and Hutcheson, Maturity group V) for seed protein, oil, fatty acids (%), and sugars (glucose, fructose, sucrose, raffinose, and stachyose, mg g<sup>-1</sup>) and their responses to the main factors of experiment (EXP, two experiments were conducted), drought, carbon dioxide (CO<sub>2</sub>), cultivar (CV), and their interactions.



Treatments	N	P	K	Mg	Fe	B	Cu	Zn	Mn
EXP	ns <sup>1</sup>	ns	ns	ns	ns	ns	ns	ns	ns
Drought <sup>2</sup>	***	***	**	*	*	***	*	*	*
CO <sub>2</sub>	*	*	*	*	*	*	*	**	**
CV	*	**	*	*	***	*	*	*	*
EXP × drought	ns	ns	ns	ns	ns	ns	ns	ns	ns
EXP × CO <sub>2</sub>	ns	ns	ns	ns	ns	ns	ns	ns	ns
EXP × CV	ns	ns	ns	ns	ns	ns	ns	ns	ns
Drought × CO <sub>2</sub>	*	*	*	*	*	***	*	**	*
Drought × CV	**	*	*	*	**	*	*	*	*
CO <sub>2</sub> × CV	**	ns	*	ns	ns	*	ns	*	ns
EXP × drought × CO <sub>2</sub> × CV	ns	ns	ns	ns	ns	ns	ns	ns	ns

Treatments used were four: T1 = plants were grown irrigated and subjected to 360  $\mu\text{mol mol}^{-1}$  CO<sub>2</sub> concentration; T2 = plants were grown irrigated and subjected to 700  $\mu\text{mol mol}^{-1}$  CO<sub>2</sub> concentration; T3 = plants were grown under drought and subjected to 360  $\mu\text{mol mol}^{-1}$  CO<sub>2</sub> concentration; and T4 = plants were grown under drought and subjected to 700  $\mu\text{mol mol}^{-1}$  CO<sub>2</sub> concentration. Plants were grown in the greenhouse, and they were transferred to growth chambers at the beginning of seed-fill stage until maturation (R8). The experiments were conducted under constant normal temperature of 26/16°C, day/night.

\* = significance at  $P \leq 0.05$ ; \*\* = significance at  $P \leq 0.01$ ; \*\*\* = significance at  $P \leq 0.001$ ; ns = not significant

Drought treatment was imposed by growing the plants at soil water potential of about -199 kPa. For irrigated plant, soil water potential was kept at about -15 to -20 kPa.

**Table 2.** Analysis of variance for soybean cultivars (Freedom and Hutcheson, maturity group V) for seed macro- and micronutrients (N, P, K, and Mg: %; Fe, B, Cu, Zn, and Mn: mg kg<sup>-1</sup>) as affected by the main effect factors of experiment (EXP, two experiments were conducted), carbon dioxide (CO<sub>2</sub>), cultivar (CV), and their interactions.

Mean values of seed constituents showed that seed protein, oleic acid, raffinose, and stachyose were higher, and oil, linolenic acid, glucose, fructose, and sucrose were lower under drought or drought with elevated CO<sub>2</sub> conditions (**Table 3**). Seed N was higher under drought or drought with elevated CO<sub>2</sub>. However, seed contents of P, K, Mg, Fe, B, Cu, Zn, and Mn were generally lower under drought or drought with elevated CO<sub>2</sub>. Comparing with ambient CO<sub>2</sub>, elevated CO<sub>2</sub> showed higher contents of seed protein, oil, glucose, sucrose, raffinose, and stachyose and lower contents of seed N, P, B, and Cu in Freedom cultivar only. Cultivar Hutcheson responded differently to ambient CO<sub>2</sub> or elevated CO<sub>2</sub>. The different responses of cultivars to ambient CO<sub>2</sub> or elevated CO<sub>2</sub> are suggested to be due to genotype and/or genetic background (**Tables 1–4**). What is clear is that seed oleic acid was higher and linolenic acid was lower and most nutrients were lower under elevated CO<sub>2</sub> (**Tables 3–6**).  $\delta^{15}\text{N}$  (<sup>15</sup>N/<sup>14</sup>N Ratio) and  $\delta^{13}\text{C}$  (<sup>13</sup>C/<sup>12</sup>C Ratio) natural abundance isotopes showed alterations under drought and drought with elevated CO<sub>2</sub>, indicating that there were changes of nitrogen and carbon metabolism under drought stress (**Figures 1 and 2**).

The higher contents of seed protein, oleic acid, raffinose, and stachyose under drought or drought with elevated CO<sub>2</sub> conditions could be due to drought effects and physiological and biochemical responses to stress. It was reported that drought resulted from climate change is predicted to alter the biochemistry and physiology of crops, impacting the nutritional value of the crop grains [12]. The effects of elevated CO<sub>2</sub> and drought on seed composition (protein, oil, fatty acids, and sugars) and minerals were previously reported. However, results of the effects of elevated CO<sub>2</sub> on seed composition are still conflicting. For example, small effects of elevated CO<sub>2</sub> on seed composition were found [40], whereas others found a significant effect of elevated CO<sub>2</sub> on soybean seed oil in cultivars Essex, Holladay, and NK6955 [41]. The authors also found that oleic fatty acid concentration was positively affected and protein concentration was not affected by CO<sub>2</sub>. Recently, it was found that elevated CO<sub>2</sub> resulted in decreased content of protein and increased contents of oil and oleic acid [39].

It was reported that alteration of carbohydrates such as starch and soluble sugars such as glucose, fructose, and sucrose occurred under stress conditions [14] and sugars of sucrose and raffinose may play a role in protecting cells against oxidative damage and accumulate as a response to stress [16, 17]. Other researchers suggested that high levels of raffinose and stachyose contribute to the acquisition of desiccation tolerance due to overexpression of galactinol synthase and accumulation of galactinol and raffinose improving drought tolerance [18, 19]. Sucrose was reported to be involved in the synthesis of raffinose oligosaccharides including stachyose and verbascose, and these sugars play an important role in seed tolerance to desiccation and drought stress [18, 26]. Some sugars, for example, maltose, accumulate in the grain under drought stress, maybe due to starch degradation [12]. The mechanisms of the involvement of these sugars in drought tolerance are still not fully understood [19–21]. It was concluded that these biochemical compounds can be used as drought indicators and can be used in breeding program to select for drought tolerance [12, 23]. CO<sub>2</sub> elevation resulted in a decrease of seed Na, Ca, Mg, S, Fe, Zn, and Mn [42, 43]. The percentage decrease of nutrients by elevated CO<sub>2</sub> ranged from 0.7 to 19.5%, except for K and P [44]. The decrease of macro- and micronutrients by elevated CO<sub>2</sub> was due to the dilution effect induced by the increase of carbohydrates in seeds [12, 39, 42–44].

Treatments <sup>1</sup>	Protein	Oil	Oleic	Linolenic	Glucose	Fructose	Sucrose	Raffinose	Stachyose
T1	38.5 c	21.3 b	22.2 c	9.6 a	1.4 b	1.01 a	61 b	8.7 d	33.6 d
T2	39.0 c	22.1 a	27.5 b	8.7 b	2.1 a	0.98 a	67 a	9.6 c	34.9 c
T3	41.2 a	19.4 c	29.7 a	6.3 c	0.8 c	0.54 b	32 d	11.5 a	45.3 a
T4	40.3 b	19.6 c	29.5 a	6.8 c	0.9 c	0.52 b	39 c	10.3 b	40.7 b

Treatments used were four: T1 = plants were grown irrigated and subjected to 360  $\mu\text{mol mol}^{-1}$  CO<sub>2</sub> concentration; T2 = plants were grown irrigated and subjected to 700  $\mu\text{mol mol}^{-1}$  CO<sub>2</sub> concentration; T3 = plants were grown under drought and subjected to 360  $\mu\text{mol mol}^{-1}$  CO<sub>2</sub> concentration; and T4 = plants were grown under drought and subjected to 700  $\mu\text{mol mol}^{-1}$  CO<sub>2</sub> concentration. Plants were grown in the greenhouse, and they were transferred to growth chambers at the beginning of seed-fill stage (R5) until full maturity (R8). The experiments were conducted under constant normal temperature of 26/16°C, day/night.

<sup>1</sup>Means within a column of each water treatment followed by the same letter are not significantly different at the 5% level as determined by Fishers' least significant difference (LSD) test. Values are means of four replicates. Drought treatment was imposed by growing the plants at soil water potential of about -199 kPa. For irrigated plant, soil water potential was kept at about -15 to -20 kPa.

**Table 3.** Drought and elevated carbon dioxide effects on soybean (Freedom cultivar, maturity group V) seed protein, oil, fatty acids (%), and sugars (glucose, fructose, sucrose, raffinose, and stachyose, mg g<sup>-1</sup>).

Treatments <sup>1</sup>	N	P	K	Mg	Fe	B	Cu	Zn	Mn
T1	6.5 a	0.64 a	1.9 a	0.34 a	75 a	64 a	12.4 a	51.2 a	42.5 a
T2	5.1 c	0.53 b	1.5 b	0.33 a	72 a	57 b	9.3 b	52.3 a	41.2 a
T3	6.7 a	0.47 c	1.1 c	0.23 b	65 b	41 d	8.2 c	46.5 c	37.6 c
T4	6.1 b	0.46 c	1.2 c	0.34 a	63 b	47 c	8.5 c	49.6 b	40.0 b

Treatments used were four: T1 = plants were grown irrigated and subjected to 360  $\mu\text{mol mol}^{-1}$  CO<sub>2</sub> concentration; T2 = plants were grown irrigated and subjected to 700  $\mu\text{mol mol}^{-1}$  CO<sub>2</sub> concentration; T3 = plants were grown under drought and subjected to 360  $\mu\text{mol mol}^{-1}$  CO<sub>2</sub> concentration; and T4 = plants were grown under drought and subjected to 700  $\mu\text{mol mol}^{-1}$  CO<sub>2</sub> concentration. Plants were grown in the greenhouse, and they were transferred to growth chambers at the beginning of seed-fill stage (R5) until full maturity (R8). The experiments were conducted under constant normal temperature of 26/16°C, day/night.

<sup>1</sup>Means within a column of each water treatment followed by the same letter are not significantly different at the 5% level as determined by Fishers' LSD test. Values are means of four replicates. Drought treatment was imposed by growing the plants at soil water potential of about -199 kPa. For irrigated plant, soil water potential was kept at about -15 to -20 kPa.

**Table 4.** Drought and elevated carbon dioxide effects on soybean (Freedom cultivar, maturity group V) seed macro- and micronutrients (N, P, K, and Mg: %; Fe, B, Cu, Zn, and Mn: mg kg<sup>-1</sup>).

Treatments <sup>1</sup>	Protein	Oil	Oleic	Linolenic	Glucose	Fructose	Sucrose	Raffinose	Stachyose
T1	39.8 c	21.3 a	21.4 c	10.5 a	2.4 a	1.23 a	68 a	11.2 c	41.4 c
T2	40.1 c	21.4 a	29.7 a	9.7 b	2.6 a	1.21 a	66 a	12.5 b	40.5 d
T3	43.5 b	18.5 b	29.6 a	7.5 c	0.7 c	0.73 c	51 c	13.5 a	47.6 a
T4	44.1 a	18.3 b	28.1 b	7.3 c	1.3 b	0.85 b	60 b	13.7 a	46.8 b

Treatments used were four: T1 = plants were grown irrigated and subjected to 360  $\mu\text{mol mol}^{-1}$  CO<sub>2</sub> concentration; T2 = plants were grown irrigated and subjected to 700  $\mu\text{mol mol}^{-1}$  CO<sub>2</sub> concentration; T3 = plants were grown under drought and subjected to 360  $\mu\text{mol mol}^{-1}$  CO<sub>2</sub> concentration; and T4 = plants were grown under drought and subjected to 700  $\mu\text{mol mol}^{-1}$  CO<sub>2</sub> concentration. Plants were grown in the greenhouse, and they were transferred to growth chambers at the beginning of seed-fill stage (R5) until full maturity (R8). The experiments were conducted under constant normal temperature of 26/16°C, day/night.

<sup>1</sup>Means within a column of each water treatment followed by the same letter are not significantly different at the 5% level as determined by Fishers' LSD test. Values are means of four replicates. Drought treatment was imposed by growing the plants at soil water potential of about -199 kPa. For irrigated plant, soil water potential was kept at about -15 to -20 kPa.

**Table 5.** Drought and elevated carbon dioxide effects on soybean (Hutcheson cultivar, maturity group V) seed protein, oil, fatty acids (%), and sugars (glucose, fructose, sucrose, raffinose, and stachyose, mg g<sup>-1</sup>).




Treatments <sup>1</sup>	N	P	K	Mg	Fe	B	Cu	Zn	Mn
T1	6.9 a	0.72 a	2.7 a	0.31 a	76 b	67 a	11.3 a	48 a	43 a
T2	5.3 c	0.59 b	1.7 b	0.26 b	81 a	65 a	10.5 b	43 b	39 b
T3	6.3 b	0.42 c	1.2 c	0.20 c	60 c	51 b	8.1 d	31 d	31 c
T4	6.1 b	0.45 c	1.1 c	0.22 c	63 c	53 b	9.8 c	35 c	33 c


Treatments used were four: T1 = plants were grown irrigated and subjected to 360  $\mu\text{mol mol}^{-1}$  CO<sub>2</sub> concentration; T2 = plants were grown irrigated and subjected to 700  $\mu\text{mol mol}^{-1}$  CO<sub>2</sub> concentration; T3 = plants were grown under drought and subjected to 360  $\mu\text{mol mol}^{-1}$  CO<sub>2</sub> concentration; and T4 = plants were grown under drought and subjected to 700  $\mu\text{mol mol}^{-1}$  CO<sub>2</sub> concentration. Plants were grown in the greenhouse, and they were transferred to growth chambers at the beginning of seed-fill stage (R5) until full maturity (R8). The experiments were conducted under constant normal temperature of 26/16°C, day/night.

<sup>1</sup>Means within a column of each water treatment followed by the same letter are not significantly different at the 5% level as determined by Fishers' LSD test. Values are means of four replicates. Drought treatment was imposed by growing the plants at soil water potential of about -199 kPa. For irrigated plant, soil water potential was kept at about -15 to -20 kPa.


**Table 6.** Drought and elevated carbon dioxide effects on soybean (Hutcheson cultivar, maturity group V) seed macro- and micronutrients (N, P, K, and Mg: %; Fe, B, Cu, Zn, and Mn: mg kg<sup>-1</sup>).



**Figure 1.** Cultivar Freedom (maturity group V).  $\delta^{15}\text{N}$  ( $^{15}\text{N}/^{14}\text{N}$  ratio) (A) and  $\delta^{13}\text{C}$  ( $^{13}\text{C}/^{12}\text{C}$  ratio) (B) natural abundance isotope in soybean seed as influenced by drought and elevated  $\text{CO}_2$ . Treatments used were four: T1 = plants were grown irrigated and subjected to  $360\ \mu\text{mol mol}^{-1}$   $\text{CO}_2$  concentration; T2 = plants were grown irrigated and subjected to  $700\ \mu\text{mol mol}^{-1}$   $\text{CO}_2$  concentration; T3 = plants were grown under drought and subjected to  $360\ \mu\text{mol mol}^{-1}$   $\text{CO}_2$  concentration; and T4 = plants were grown under drought and subjected to  $700\ \mu\text{mol mol}^{-1}$   $\text{CO}_2$  concentration. Plants were grown in the greenhouse, and they were transferred to growth chambers at the beginning of seed-fill stage (R5) until full maturity (R8). The experiments were conducted under constant normal temperature of  $26/16^\circ\text{C}$ , day/night.



Carbon dioxide and drought treatments



Carbon dioxide and drought treatments

**Figure 2.** Cultivar Hutcheson (maturity group V).  $\delta^{15}\text{N}$  ( $^{15}\text{N}/^{14}\text{N}$  ratio) (A) and  $\delta^{13}\text{C}$  ( $^{13}\text{C}/^{12}\text{C}$  ratio) (B) natural abundance isotope in soybean seed as affected by drought and elevated  $\text{CO}_2$ . Treatments used were four: T1 = plants were grown irrigated and subjected to  $360 \mu\text{mol mol}^{-1}$   $\text{CO}_2$  concentration; T2 = plants were grown irrigated and subjected to  $700 \mu\text{mol mol}^{-1}$   $\text{CO}_2$  concentration; T3 = plants were grown under drought and subjected to  $360 \mu\text{mol mol}^{-1}$   $\text{CO}_2$  concentration; and T4 = plants were grown under drought and subjected to  $700 \mu\text{mol mol}^{-1}$   $\text{CO}_2$  concentration. Plants were grown in the greenhouse, and they were transferred to growth chambers at the beginning of seed-fill stage (R5) until full maturity (R8). The experiments were conducted under constant normal temperature of  $26/16^\circ\text{C}$ , day/night.

Our results showed that elevated CO<sub>2</sub> concentration enhanced oleic acid and carbohydrates (mainly sucrose and glucose) and reduced NPK and some micronutrients compared with ambient CO<sub>2</sub>. The increase of carbohydrates due to elevated CO<sub>2</sub> could result from enhanced photosynthesis resulting in higher total carbohydrates [4]. The decrease of NPK and some micronutrients was due to the dilution of concentrations of the nutrients by higher levels of carbohydrates as supported by previous research [12, 39, 42–44]. Higher seed protein, oleic acid, raffinose and stachyose, and lower oil, linolenic acid, glucose, fructose, and sucrose under drought or drought with elevated CO<sub>2</sub> conditions can be used as biochemical indicators/markers to be used for breeding to select for drought tolerance.

Therefore, research is needed to quantify the negative impact of elevated CO<sub>2</sub> and its interactions with biotic and abiotic stresses, including drought, on seed quality to breed for higher seed nutritional qualities and develop appropriate crop management systems [43, 45]. It was reported that the physiological processes influenced by drought can be identified by the biochemical characterization of plant tissues under stress conditions [46]. Limited research was conducted to characterize the biochemical compounds such as protein, amino acids, carbohydrates, and phenolics in the breeding program as a tool for selection, but this tool has promise in selecting for abiotic stress tolerance [24, 47]. The changes of  $\delta^{15}\text{N}$  ( $^{15}\text{N}/^{14}\text{N}$  Ratio) and  $\delta^{13}\text{C}$  ( $^{13}\text{C}/^{12}\text{C}$  Ratio) natural abundance isotopes under drought and drought with elevated CO<sub>2</sub> indicated the increase of  $^{15}\text{N}$  derived from plant gas exchange through stomatal conductance and CO<sub>2</sub> fixation [48, 49]. The alteration of  $^{13}\text{C}/^{12}\text{C}$  ratio indicated a possible shift of carbon metabolism leading to less discrimination against  $\delta^{13}\text{C}$  [50].

## 4. Conclusions

The current research demonstrated that elevated CO<sub>2</sub>, drought, and drought combined with elevated CO<sub>2</sub> altered seed biochemical compounds, including protein, carbohydrates, and minerals. This research increases our knowledge of the interaction between elevated CO<sub>2</sub> and drought resulting from climate changes for seed chemical composition and mineral nutrition.

The high level of oleic acid and low level of linolenic acid are desirable, and they contribute to oil stability and long shelf life of the oil. Sugars of both raffinose and stachyose may play a possible role in drought stress response. Alterations of  $\delta^{15}\text{N}$  and  $\delta^{13}\text{C}$  natural abundance isotopes indicated changes in nitrogen and carbon metabolism. The characterization of these seed biochemical compounds under elevated CO<sub>2</sub> or drought stress can be used as biomarkers in breeding program to select for crop drought tolerance and high seed nutritional qualities, as these traits are related to seed production, quality, and food security. Since limited research was conducted on the effects of climate change on seed quality and mineral nutrition, further research is needed.

## Acknowledgements

We thank Sandra Mosley for lab analysis and greenhouse and growth chamber technical assistance. This work was supported by the US Department of Agriculture, Agricultural Research

Service Project 6402-21220-012-00D. Mentioning trade names or commercial products in this publication is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the US Department of Agriculture.

## Author details

Nacer Bellaloui<sup>1\*</sup>, Alemu Mengistu<sup>2</sup>, Hamed K. Abbas<sup>3</sup> and My A. Kassem<sup>4</sup>

\*Address all correspondence to: [nacer.bellaloui@ars.usda.gov](mailto:nacer.bellaloui@ars.usda.gov)

1 Crop Genetics Research Unit, USDA-ARS, Stoneville, MS, USA

2 Crop Genetics Research Unit, USDA-ARS, Jackson, TN, USA

3 Biological Control of Pests Research Unit, USDA-ARS, Stoneville, MS, USA

4 Department of Biological Sciences, Plant Genomics and Biotechnology Laboratory, Fayetteville State University, Fayetteville, NC, USA

## References

- [1] Gregory PJ, Ingram JSI, Brklacich M: Climate change and food security. *Phil Trans R Soc B*. 2005; **360**: 2139–2148. doi: 10.1098/rstb.2005.1745.
- [2] Center for climate and energy solutions, Drought and climate change <http://www.c2es.org/science-impacts/extreme-weather/drought> Verified August 22, 2016.
- [3] Manabe S, Wetherald RT: On the distribution of climate change resulting from an increase in CO<sub>2</sub>-content of the atmosphere. *J Atmos Sci*. 1980; **37**: 99–118.
- [4] Rogers A, Ainsworth EA, Leakey ADB: Update on legumes and elevated CO<sub>2</sub>: Will elevated carbon dioxide concentration amplify the benefits of nitrogen fixation in legumes? *Plant Physiol*. 2009; **151**: 1009–1016.
- [5] Farquhar GD, Sharkey TD: Stomatal conductance and photosynthesis. *Annu Rev Plant Physiol*. 1982; **33**: 317–345.
- [6] Cu J, Vu V, Allen Jr LH, Bowes G: Drought stress and elevated CO<sub>2</sub> effects on soybean ribulose biphosphate carboxylase activity and canopy photosynthetic rates. *Plant Physiol*. 1987; **83**: 573–578.
- [7] Fellows RJ, Boyer JS: Structure and activity of chloroplasts of sunflower leaves having various water potentials. *Planta*. 1976; **132**: 229–239.
- [8] Keck RW, Boyer JS: Chloroplast response to low leaf water potentials III. Differing inhibition of electron transport and photophosphorylation. *Plant Physiol*. 1974; **53**: 474–479.
- [9] Mayoral ML, Atsmon D, Shimshi D, GRoMET-Elhanan Z: Effect of water stress on enzyme activities in wheat and related wild species: carboxylase activity, electron transport and photophosphorylation in isolated chloroplasts. *Aust J Plant Physiol*. 1981; **8**: 385–393.



- [10] Emam Y, Shekoofa A, Salehi F, Jalali AH: Water stress effects on two common bean cultivars with contrasting growth habits. *Am Eurasian J Agric Environ Sci*. 2010; **9**: 495–499.
- [11] Broughton WJ, Hernández G, Blair M, Beebe S, Gepts P, Vanderleyden J: Beans (*Phaseolus spp.*). *Plant Soil*. 2003; **252**: 55–128.
- [12] Andrade ER, Ribeiro VN, Azevedo CVG, Chiorato AF, Williams TCR, Carbonell SAM: Biochemical indicators of drought tolerance in the common bean (*Phaseolus vulgaris* L.) *Euphytica*. 2016; **210**: 277–289. doi: 10.1007/s10681-016-1720-4
- [13] Caldwell CR, Britz SJ, Mirecki RM: Effect of temperature, elevated carbon dioxide, and drought during seed development on the isoflavone content of dwarf soybean [*Glycine max* (L.) Merrill] grown in controlled environments. *J Agric Food Chem*. 2005; **53**: 1125–1129.
- [14] Nishizawa A, Yabuta Y, Shigeoka S: Galactinol and raffinose as a novel function to protect plants from oxidative damage. *Plant Physiol*. 2008; **23**: 108–122.
- [15] Schnyder H: The role of carbohydrate storage and redistribution in the source-sink relations of wheat and barley during grain filling: A review. *New Phytol*. 1993; **123**: 233–245.
- [16] Cramer GR, Ergul A, Grimpert J, Tillett RL, Tattersall EA, Bohlman MC, Vincent D, Sonderegger J, Evans J, Osborne C, Quilici D, Schlauch KA, Schooley DA, Cushman JC: Water and salinity stress in grapevines: early and late changes in transcript and metabolite profiles. *Funct Integr Genomics*. 2007; **7**: 111–134.
- [17] Rolland F, Baena-Gonzalez E, Sheen J: Sugar sensing and signaling in plants: conserved and novel mechanisms. *Annu Rev Plant Biol*. 2006; **57**: 675–709.
- [18] Taji T, Ohsumi C, Iuchi S, Seki M, Kasuga M, Kobayashi M, Yamaguchi-Shinozaki K, Shinozaki K: Important roles of drought- and cold-inducible genes for galactinol synthase in stress tolerance in *Arabidopsis thaliana*. *Plant J*. 2002; **29**: 417–426.
- [19] Bellaloui N, Gillen AM, Mengistu A, Kebede H, Fisher DK, Smith JR, Reddy KN: Responses of nitrogen metabolism and seed nutrition to drought stress in soybean genotypes differing in slow-wilting phenotype. *Front Plant Sci*. 2013; **4**: 498. doi: 10.3389/fpls.2013.00498. eCollection 2013.
- [20] Chen TH, Murata N: Enhancement of tolerance of abiotic stress by metabolic engineering of betaines and other compatible solutes. *Curr Opin Plant Biol*. 2002; **5**: 250–257. doi: 10.1016/S1369-5266(02)00255-8
- [21] Bartels D, Sunkar R: Drought and salt tolerance in plants. *Crit Rev Plant Sci*. 2005; **24**: 23–58. doi: 10.1080/07352680590910410
- [22] Muller B, Pantin F, Génard M, Turc O, Freixes S, Piques M, Gibon Y: Water deficits uncouple growth from photosynthesis, increase C content, and modify the relationships between C and growth in sink organs. *J Exp Bot*. 2011; **62**: 1715–1729.
- [23] Blum A: Drought resistance, water-use efficiency, and yield potential — are they compatible, dissonant or mutually exclusive? *Aust J Agric Res*. 2005; **56**: 1159–1168.

- [24] Hasanuzzaman M, Nahar K, Gill SS, Gill R, Fujita M: Drought stress responses in plants, oxidative stress, Q1 and antioxidant defense. In: Climate change and plant abiotic stress tolerance, Tuteja N, Gill SS (Eds). 2014; Wiley-Blackwell, Germany, 209–237.
- [25] Wang Y, Frei M: Stressed food—the impact of abiotic environmental stresses on crop quality. *Agric Ecosyst Environ.* 2011; **141**: 271–286.
- [26] Pinheiro C, Rodrigues AP, Carvalho IS, Chaves MM, Ricardo CP: Sugar metabolism in developing lupin seeds is affected by a short-term water deficit. *J Exp Bot.* 2005; **56**: 2705–2712.
- [27] Beebe S, Rao IM, Blair M, Butare L: Breeding for abiotic stress tolerance in common bean: present and future challenges. *CIAT.* 2009: 10-14.
- [28] Fehr WR, Caviness CE: Stages of soybean development. *Iowa Agric. Exp. Stn. Spec. Rep.* 80. 1977; Iowa State University, Ames.
- [29] Bellaloui N, Mengistu A, Walker ER, Young LD: Soybean seed composition as affected by seeding rates and row spacing. *Crop Sci.* 2014; **54**: 1782–1795.
- [30] Bellaloui N, Smith JR, Ray JD, Gillen AM: Effect of maturity on seed composition in the early soybean production system as measured on near-isogenic soybean lines. *Crop Sci.* 2009; **49**, 608–620.
- [31] Wilcox JR, Shibles RM: Interrelationships among seed quality attributes in soybean. *Crop Sci.* 2001; **41**: 11–14.
- [32] Lohse G: Microanalytical azomethine-H method for boron determination in plant tissue. *Commun Soil Sci Plant Anal.* 1982; **13**: 127–134.
- [33] Dordas C, Apostolides G, Goundra O: Boron application affects seed yield and seed quality of sugar beets. *J Agric Sci.* 2007; **145**: 377–384.
- [34] John MK, Chuah HH, Neufeld JH: Application of improved azomethine-H method to the determination of boron in soils and plants. *Anal Lett.* 1975; **8**: 559–568.
- [35] Bandemer SL, Schaible PJ: Determination of iron. A study of the o-phenanthroline method. *Indus Eng Chem, Analytical Edition*, 1944; **16**: 317–319. doi: 10.1021/i560129a013
- [36] Loeppert RL, Inskeep WP: Colorimetric determination of ferrous iron and ferric iron by the 1,10-phenanthroline method. In: *Methods of soil analysis: Part 3, Chemical methods*, Bigham JM (Eds). SSSA; Madison, WI, 1996: 659–661pp.
- [37] Cavell A J: The colorimetric determination of phosphorus in plant materials. *J Sci Food Agric.* 1955; **6**: 479–480.
- [38] SAS. Statistical analysis system. 2002–2010; Cary NC: USA.
- [39] Bellaloui B, Hu Y, Mengistu A, Abbas HK, Kassem MA, Tigabu M: Elevated atmospheric carbon dioxide and temperature affect seed composition, mineral nutrition, and <sup>15</sup>N and <sup>13</sup>C dynamics in soybean genotypes under controlled environments. *A J Plant Biol.* 2016: 56–65. doi: 10.5147/ajpb.2016.0157

- [40] Thomas JMG, Boote KJ, Allen Jr LH, Gallo-Meagher M, Davis JM: Elevated temperature and carbon dioxide effects on soybean seed composition and transcript abundance. *Crop Sci.* 2003; **43**: 1548–1557.
- [41] Heagle AS, Miller JE, Pursley WA: Influence of ozone stress on soybean response to carbon dioxide enrichment: III. Yield and seed quality. *Crop Sci.* 1998; **38**: 128–134.
- [42] Manderscheid R, Bender J, Jager HJ, Weigel HJ: Effects of season long CO<sub>2</sub> enrichment on cereals. II. Nutrient concentrations and grain quality. *Agric Ecosyst Environ.* 1995; **54**: 175–185.
- [43] Upreti DC, Sen S, Dwivedi N: Rising atmospheric carbon dioxide on grain quality in crop plants. *Physiol Mol Biol Plants.* 2010; **16**: 215–227.
- [44] Hogy P, Fangmeier A: Effects of elevated atmospheric CO<sub>2</sub> on grain quality of wheat. *J Cereal Sci.* 2008; **48**: 580–591.
- [45] Prasad P, Vara V, Allen Jr LH, Boote KJ: Crop responses to elevated carbon dioxide and interaction with temperature: grain legumes. *J Crop Improv.* 2005; **13**: 113–155.
- [46] Silvente S, Sobolev AP, Lara M: Metabolite adjustments in drought tolerant and sensitive soybean genotypes in response to water stress. *PLoS One.* 2012; **7**: 1–10.
- [47] Zhang BQ, Yang LT, Li YR: Physiological and biochemical characteristics related to cold resistance in sugar cane. *Sugar Tech.* 2015; **17**: 4.
- [48] Livingston NJ, Guy RD, Sun ZJ, Ethier GJ: The effects of nitrogen stress on the stable carbon isotope composition, productivity and water use efficiency of white spruce (*Picea glauca* (Moench) Voss) seedlings. *Plant Cell Environ.* 1999; **22**: 281–289.
- [49] Matsushima M, Chang SX: Nitrogen and water availabilities and competitiveness of bluejoint: Spruce growth and foliar carbon-13 and nitrogen-15 abundance. *Soil Sci Soc Am J.* 2007; **71**: 1547–1554.
- [50] O'Leary MH: Environmental effects on carbon isotope fractionation in terrestrial plants. In: *Stable isotopes in the biosphere*, Wada E, Yoneyama T, Minagawa M, Ando T, Fry BD (Eds). 1995; Kyoto University Press, Kyoto, Japan, 517–530 pp.