

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

185,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Chitinous Materials for Control of Foodborne Pathogens and Mycotoxins in Poultry

Daniel Hernandez-Patlan, Bruno Solis-Cruz,
Billy M. Hargis and Guillermo Tellez

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/intechopen.76041>

Abstract

Public concern with the incidence of antibiotic-resistant bacteria, particularly among foodborne pathogens has been challenging the poultry industry to find alternative means of control. Chitosan is a modified, natural biopolymer derived by deacetylation of chitin. The antimicrobial activity and film-forming property of chitosan makes it a potential source of food preservative or coating material of natural origin for improvement of quality and shelf life of various foods of agriculture, poultry, beef and seafood origin. In addition to its use as an antimicrobial, it has been shown that it has good properties as a mycotoxin adsorbent. The purposes of the present chapter is to summarize our experience using chitin-chitosan from Deacetylated 95% food grade chitosan (Paragon Specialty Products LLC Rainsville, AL) or *Aspergillus oryzae* meal (Fermacto®, PetAg Inc., Hampshire IL) to control foodborne pathogens, improve performance, biological sanitizer and mycotoxin binder in commercial poultry.

Keywords: chitosan, Fermacto®, *Salmonella*, mycotoxins, gut health

1. Introduction

Chitin ($C_8H_{13}O_5N$)_n is a long-chain polymer of a N-acetylglucosamine (**Figure 1(a)**), a derivative of glucose, and is found in many places globally. It is the main component of the cell walls of fungi, the exoskeletons of arthropods such as crustaceans (e.g., crabs, lobsters and shrimps) and insects, the radula of mollusks, and the beaks of cephalopods, including squid and octopi [1]. In terms of structure, chitin may be compared to the polysaccharide cellulose and, in terms of function, to the protein keratin [2]. Depending on its source, two types of chitin allomorphs




Figure 1. Chemical structure (a) of chitin poly(N-acetyl- β -D-glucosamine) and (b) of chitosan (poly(D-glucosamine) repeat units.

can occur, the α and β forms, which can be differentiated by infrared and solid-state NMR spectroscopy together with X-ray diffraction [3]. Chitosan is a high molecular weight polysaccharide linked by a β -1,4 glycoside and is composed of N-acetyl-glucosamine and glucosamine (**Figure 1(b)**). It is a natural biopolymer derived by deacetylation of chitin and the most widespread polycationic biopolymer [3]. However, although chitosan is obtained from chitin, the applications of the latter compared to chitosan are limited because it is chemically inert [4] and because of its poor solubility [5]. Unlike chitin, chitosan is soluble but in an acidic media since at neutral or alkaline pH it is insoluble. The properties of chitosan can be modified by changing the degree of deacetylation, pH and ionic strength. At neutral pH, most chitosan molecules will lose their charge and precipitate when it is in solution [6].

The application of chitin is focused on obtaining soluble derivatives in aqueous media such as chitosan [3]. Chitosan has several applications in fields such as waste and water treatment, agriculture, fabric and textiles, cosmetics, nutritional enhancement, and food processing. Given its low toxicity and allergenicity, and its biocompatibility, biodegradability and bioactivity, it is a very attractive substance for diverse applications in the pharmaceutical and medical fields, since it has been used for systemic and local delivery of drugs and vaccines [7]. However, one of the most important application is its antimicrobial activity against bacteria, filamentous fungi and yeasts. Chitosan has wide spectrum of activity against Gram-positive and Gram-negative bacteria but it is more effective against Gram-negative bacteria [8, 9]. Furthermore, it has been reported that the antimicrobial activity and film-forming property of chitosan makes it a potential source of food preservative or coating material of natural origin for improvement of quality and shelf life of various foods of agriculture, poultry, beef and seafood origin [3, 10]. The mechanism of the antimicrobial activity of chitosan has not yet been fully elucidated, but several hypotheses have been proposed. The most feasible hypothesis is a change in cell permeability due to interactions between the positively charged chitosan molecules and the negatively charged microbial cell membranes [11, 12]. Other mechanisms include the interaction of diffused hydrolysis products with microbial DNA, which leads to the inhibition of mRNA and protein synthesis and chelation of metals, spore elements, and essential nutrients [5, 13].

The antimicrobial activity of chitosan depends on both intrinsic and extrinsic factors. Among the intrinsic factors are the molecular weight and degree of deacetylation of chitosan. While the extrinsic factors include pH, temperature and reactive time [14]. Moreover, it has been observed that when the chitosan is in nanoparticle form, it has better antimicrobial

properties since its small particle size gives it a greater surface area and high reactivity which could enhance the charge interaction with the bacterial surface and of this way to produce a superior antimicrobial effect [15].

2. Antimicrobial effect of chitosan on *Salmonella* in broiler chickens

Salmonella enterica serovars continue to be among the most important foodborne pathogens worldwide due to the considerable human rates of illness reported, the wide hosts species that are colonized by members of this remarkable pathogen genus, which serve as vectors and reservoirs for spreading these agents to animal and human populations. **Figure 2** shows the distribution of the major serotypes of *Salmonella* with importance in public health. Furthermore, the public concern for the appearance of resistant strains to many antibiotics, particularly among zoonotic pathogens such as common *Salmonella* isolates, is also challenging the poultry industry to find alternative means of control [16]. For these reasons, continued research on sustainable alternatives to antibiotic growth promoters for animal production is needed.

Interest in chitosan, a biodegradable, nontoxic, non-sensitizing, and biocompatible polymer isolated from shellfish, arises from the fact that chitosan is reported to exhibit numerous beneficial effects, including strong antimicrobial and antioxidant activities in foods [18]. Its application in agriculture, horticulture, environmental science, industry, microbiology, and medicine are well reported. A significant interest in the antimicrobial activities of chitosan either as solution, or as powders, edible films and coating against foodborne pathogens, spoilage bacteria, and pathogenic viruses and fungi in several food categories has been extensively investigated [19]. We have evaluated the effect *in vitro* and *in vivo* of chitosan on *Salmonella typhimurium* in broiler chickens [20]. In our *in vitro* crop assay experiments (**Table 1**), 0.2% chitosan significantly




Figure 2. Distribution of the major serotypes of non-typhoidal *Salmonella* associated with human cases (salmonellosis) and poultry meat in EU, 2011 to 2013 [17].

Treatment	Trial 1		Trial 2		Trial 3	
	30 min	6 h	30 min	6 h	30 min	6 h
Control	5.22 ± 0.15 ^a	7.62 ± 0.01 ^a	5.19 ± 0.11 ^a	6.99 ± 0.03 ^a	6.05 ± 0.18 ^a	7.95 ± 0.31 ^a
Chitosan (0.2%)	3.94 ± 0.20 ^b	3.04 ± 0.20 ^b	3.49 ± 0.24 ^b	4.40 ± 0.19 ^b	5.05 ± 0.19 ^b	5.31 ± 0.26 ^b

^{a-b}Values within columns with different lowercase superscripts differ significantly (P < 0.05).

Table 1. Antimicrobial activity of chitosan on *Salmonella typhimurium* in an in vitro crop assay.

Treatment	Trial 1		Trial 2	
	Cecal tonsils (CT)	Log ₁₀ ST/g of CT	Cecal tonsils (CT)	Log ₁₀ ST/g of CT
Control	15/20 (75%)	4.20 ± 0.82 ^a	15/20 (75%)	5.00 ± 0.62 ^a
Chitosan (0.2%)	9/20 (45%)	2.28 ± 0.75 ^b	12/20 (60%)	3.34 ± 0.72 ^b

^{a-b}Values within columns with different lowercase superscripts differ significantly (P < 0.05).

Table 2. Effect of chitosan on *Salmonella enteritidis* cecal tonsils colonization in 7-days-old broiler chickens.

reduced the cfu of ST at 30 min or 6 h compared with control (P < 0.05). In the *in vivo* experiments with 40 day-of-hatch broiler chicks and challenged with 2 × 10⁵ cfu ST, dietary 0.2% chitosan significantly reduce the cfu/g of ST in the ceca in both experiments (**Table 2**). However, no significant reduction in the incidence of ST in cecal tonsils colonization was observed, suggesting that 0.2% chitosan significantly reduced the cfu of ST/gram *in vitro* and *in vivo*.

3. Effect of chitosan as a biological sanitizer on chicken skin

Chickens contain large numbers of microorganisms in their gastrointestinal tract and on their feathers and feet; therefore, storage quality of fresh chicken is partially dependent on the bacteria present on the integument prior to slaughter. Pathogenic microorganisms present in chicken carcasses after processing and throughout scalding and picking can contaminate equipment and other carcasses [21]. Pathogenic bacteria such as *Salmonella spp.* and *Campylobacter spp.* are able to attach to skin and penetrate in skin layers or feather follicles, facilitating their presence on chicken skin and carcass during poultry processing [22]. Critical control point determination at broiler processing has become very important, especially because of the recent attention on

Treatment	Trial 1		Trial 2	
	1 h	24 h	1 h	24 h
Control	6.57 ± 0.11 ^a	6.03 ± 0.02 ^a	6.78 ± 0.06 ^a	7.36 ± 0.06 ^a
Chitosan (0.5%)	6.23 ± 0.03 ^a	5.81 ± 0.06 ^b	7.06 ± 0.08 ^a	6.60 ± 0.17 ^b

^{a-b}Values within columns with different lowercase superscripts differ significantly (P < 0.05).

Table 3. *Salmonella typhimurium* (log₁₀ cfu ± standard error)/square cm of chicken skin treated with 0.5% chitosan solution.

Hazard Analysis and Critical Control Points (HACCP) for reduction of microbial contamination of meat and poultry [23]. For all these reasons, strategies to reduce bacterial contamination on poultry carcasses are important. However, most of the bacterial reduction strategies for poultry comprise the use of antimicrobial chemicals in rinses or washes and their efficacy is reduced by the presence of organic matter. Therefore, it grows the need of biological sanitizers in the processing plant to prevent carcass to carcass cross-contamination by pathogenic bacteria and to lower the potential of foodborne diseases. Interest in chitosan, a biocompatible polymer derived from shellfish, as a biological sanitizer arises from reports showing several beneficial effects such as antimicrobial and antioxidative activities in foods [2]. The use of chitosan in industry, agriculture, and medicine is well described [13]. The antimicrobial activities of chitosan against foodborne pathogens has been broadly investigated in the food industry [24]. Research conducted in our laboratory on the effect of chitosan as a biological sanitizer in chicken skin contaminated with *Salmonella* Typhimurium and aerobic Gram-negative spoilage bacteria present on chicken skin, have revealed that 0.5% chitosan for 30 s dipping ST contaminated skin samples in a solution of 0.5% chitosan reduced ($P < 0.05$) the recovery of ST by 24 h as well as the presence of spoilage-causing psychrotrophic bacteria below detectable levels [19], (Table 3). The antimicrobial activity and film-forming characteristic of chitosan makes it a potential source of food preservative, increasing quality and shelf life of different types of foods [10]. The mechanism of the antimicrobial activity of chitosan has not yet been fully elucidated; nevertheless, different hypotheses have been proposed. The most realistic hypothesis is that chitosan is able to change cell permeability due to interactions between the positive charges of its molecules and the negative charges of the bacterial cell membranes [1]. Other hypotheses include the chelation of metals and essential nutrients, inhibiting bacterial growth had also suggested that high molecular weight chitosan could be able to form a polymer membrane around the bacterial cell, preventing it from receiving nutrients [25].

4. Prebiotic properties of *Aspergillus oryzae* to control foodborne pathogens improve performance and bone mineralization in poultry

Prebiotics are non-digestible food ingredients that are selectively fermented by gut bacteria and are known to have positive effects on gastrointestinal (GI) physiology. Some prebiotics have been shown to selectively stimulate the growth of endogenous lactic acid bacteria in the gut thereby improving the health of the host [26]. Prebiotics selectively modify the colonic microflora and can potentially influence gut metabolism [27]. The commercially available mycelium product of *Aspergillus oryzae*, Fermacto® (PetAg Inc. Hampshire, IL 60140 USA), referred to as *Aspergillus* meal (AM), has no live cells or spores and is proven to enhance the digestive efficiency of the GI tract [28]. *Aspergillus* fiber contains beta-glucans [29], fructooligosaccharides (FOS) [30], chitosan [31], and mannanoligosaccharides (MOS) [32]. Beta-glucan is considered as a powerful immune-enhancing nutritional supplement that affects the intestinal villi and primes the innate immune system to help the body defend itself against viral and bacterial invaders [33]. MOS protect the GI tract from invading toxins and pathogens by binding toxin active sites [34]. FOS and chitosan refer to a class of non-digestible carbohydrates that are readily fermented by beneficial bacteria in the intestine [30]. A healthy population of these beneficial bacteria in the digestive tract enhances the digestion and absorption of nutrients, detoxification and elimination processes,

and helps boost the immune system [35]. With an increase in the dependence on livestock as an important food source, it becomes crucial to achieve good health in order to make rearing of animal food sources safe and beneficial to both animals and humans.

Several studies have demonstrated that prevention of *Salmonella* colonization in chickens can be achieved by feeding prebiotics [36]. According to Lowry et al. [37], dietary beta-glucan reduces SE colonization significantly in chickens. In their experiment, SE from L/S was recovered from 76% of non-treated birds, while only 7% of the birds were positive for SE in the treated group, (Figure 3). Moreover, in the same study, heterophils isolated from birds treated with dietary beta-glucan contained 40% ($p < 0.05$) more SE than heterophils isolated from untreated birds, (Table 4). Heterophils form the first line of defense and killing of *Salmonella* by heterophils is well-described. This corroborates the immunostimulatory effect of beta-glucans and FOS are widely used as prebiotics in a broad range of animal species, and these carbohydrates have been tested with success for protection against *Salmonella* infections in chickens and other avian pathogens [35, 38, 39]. Kim et al. conducted a study where dietary MOS (0.05%) and FOS (0.25%) had an effect on intestinal microflora of broiler chickens, suggesting the use of these prebiotics as an alternative to the use of growth-promoting antibiotics [40]. Finally, chitosan is a modified, natural biopolymer derived by deacetylation of chitin, the main component of the cell walls of fungi and exoskeletons of arthropods. As mention before, chitosan exhibits numerous beneficial effects, including strong antimicrobial and antioxidative activities. Its application in agriculture, horticulture, environmental science, industry, microbiology, and medicine are well reported [10]. According to Huang et al., the use of 0.01 or 0.015% of oligo-chitosan in the diet increased serum levels of immunoglobulins in broiler chickens, suggesting a potential immunomodulatory effect [41]. There have been numerous studies that report the use of chitosan as a mucosal adjuvant, by enhancing IgA levels. It is well known that IgA is active across mucosal surfaces and is the predominant class of antibody against enteric pathogens [42, 43]. The commercial prebiotic supplement derived from *Aspergillus* sp. mycelium is




Figure 3. Effects of dietary β -glucan on SE organ invasion in immature chickens. β -glucan fed as a nutritional supplement to neonatal chickens 3 days prior to SE challenge. (*indicates statistically significant differences, $P < 0.05$).

Treatments	Percent heterophils + SE	Mean #SE/heterophil	Phagocytic index (PI)
Control feed	38.54 ± 0.05 ^b	4.38 ± 1.08 ^b	175.54 ± 44.92 ^b
β-glucan feed	78.84 ± 0.03 ^a	8.20 ± 0.76 ^a	644.10 ± 57.07 ^a

^{a-b}Values within columns with different lowercase superscripts differ significantly (P < 0.05).

Table 4. Effects of feeding β-glucan on chicken heterophil phagocytosis.

unique because it contains all of the above mentioned prebiotic ingredients. Additionally, AM contains 16% protein and 45% fiber and may be used with low levels of protein and amino acid diets to improve performance in commercial poultry [28, 44]. Even though the exact mechanisms of action for prebiotics have not been defined, it may be speculated that the effect is due to changing intestinal flora that promotes the growth of beneficial bacteria. This product has also been shown to benefit poultry through stimulation of growth, most probably by increasing absorption of feed ingredients and improving digestibility [45, 46].

In a recent study conducted in our laboratory we evaluated the effect of 0.2% dietary *Aspergillus* meal (AM) against horizontal transmission of *Salmonella* spp. in turkeys and chickens [36]. The results of this study showed that dietary supplementation with 0.2% *Aspergillus* Meal was able to reduce *Salmonella enteritidis* horizontal transmission in turkeys, (Table 5) and *Salmonella* Typhimurium horizontal transmission in broiler chickens, by reducing the overall colonization levels in birds, (Table 6). Although the mechanism of action is not totally understood, the reduction in *Salmonella* colonization may be related to a synergistic effect between beta-glucan, MOS, chitosan, and FOS present in the *Aspergillus oryzae* mycelium. In a previous work, we showed that dietary AM induces important changes on intestinal morphometry in turkey poult such as increased number of acid mucin cells in the duodenum, neutral mucin cells in the ileum, and

Groups	Day 10 cecal tonsils	Day 20 cecal tonsils	Day 30 cecal tonsils
Control—No AM	20/20 (100%)	18/20 (90%)	15/20 (75%)
AM	15/20 (75%)*	12/20 (60%)*	8/20 (40%)*

* (a) in the Control-No AM group and (b) in the AM group in both tables.

^{a-b}Values within columns with different lowercase superscripts differ significantly (P < 0.05).

Table 5. Effect of dietary *Aspergillus* meal against horizontal transmission of *Salmonella enteritidis* at 10, 20 and 30 days of age in turkeys.

Groups	Trial 1		Trial 2	
	Liver/spleen	Cecal tonsils	Liver/spleen	Cecal tonsils
Control—No AM	18/20 (90%)	20/20 (100%)	19/20 (95%)	18/20 (90%)
AM	6/20 (30%)*	5/20 (25%)*	8/20 (40%)*	6/20 (30%)*

* (a) in the Control-No AM group and (b) in the AM group in both tables.

^{a-b}Values within columns with different lowercase superscripts differ significantly (P < 0.05).

Table 6. Effect of dietary *Aspergillus* meal against horizontal transmission of *Salmonella typhimurium* at 10 days of age in chickens.

	Control	Aspergillus meal
Body weight (Kg)	600.32 ± 52.26 ^b	720.87 ± 63.82 ^a
FC (feed: gain)	1.34 ± 0.03 ^a	1.23 ± 0.02 ^b
Mortality (%)	2.00% ^a	2.50% ^a

^{a-b}Values within columns with different lowercase superscripts differ significantly (P < 0.05).

Table 7. Effect of *Aspergillus* meal on productive parameters in turkey poult at 30 days of ages.

sulphomucin cells in the duodenum and ileum, as well as increased villi height and villi surface area of both duodenum and ileum when compared to control, suggesting that AM prebiotic has an impact on the mucosal architecture and goblet cells proliferation in the duodenum and ileum of neonate poult [45]. Our extended studies using dietary AM prebiotic supplemented for 30 days, have shown significantly increased the body weight of neonate poult and improved feed conversion when compared with poult that received the control basal diet, and interestingly, energy and protein content in the ileum was significantly lower in poult that received dietary AM prebiotic compared with control poult suggesting better digestibility, absorption of those nutrients and bone mineralization [46]. FOS have been shown to stimulate calcium (Ca) and magnesium (Mg) absorption in the intestine and increase bone mineral concentrations in humans and rats as well as stimulate net Ca transport from the epithelium of the small and large intestine [30, 38].

The gastrointestinal tract serves as the interface between diet and the metabolic events that sustain life. Intestinal villi, which play a crucial role in digestion and absorption of nutrients, are underdeveloped at hatch and maximum absorption capacity is attained by 10 days of age [47]. Understanding and optimizing the maturation and development of the intestine in poultry will improve feed efficiency, growth and overall health of the bird. Studies on nutrition and metabolism during the early phase of growth in poult may, therefore, help in optimizing nutritional management for maximum growth. By dietary means it is possible to affect the development of the gut and the competitiveness of both beneficial and harmful bacteria, which can alter not only gut dynamics, but also many physiologic processes due to the end products metabolized by symbiotic gut microflora [48]. Additives such as enzymes, probiotics and prebiotics are now extensively used throughout the world [49–51]. Our studies suggest that the increase in performance and bone parameters in neonatal poult fed with 0.2%AM (Table 7), may be related to a synergistic effect between beta-glucan, MOS, chitosan and FOS from *Aspergillus niger* mycelium [45, 46].

5. Evaluation of chitosan as binding adsorbent material to prevent mycotoxicosis poultry

Mycotoxins are secondary toxic metabolites produced by filamentous fungi which, even at low concentrations, represent an important danger for both animal and human health [52, 53]. Currently, over 300 mycotoxins have been identified worldwide, being aflatoxins, ochratoxins, zearalenone, trichothecenes, and fumonisins, the most frequently found with synergistic toxic effects reported when more than one of these mycotoxins are present in the feed [54, 55]. Mycotoxins are chemically and structurally different, representing serious public health risk

factors since mycotoxins have been shown to have carcinogenic, teratogenic, nephrotoxic, and hepatotoxic effects after the consumption of contaminated grains or animal food products [56]. On the other hand, mycotoxins are equally important in the animal food industry, causing significant economic losses due to diminished performance and productivity, decreased reproductive parameters, and an increased mortality rate associated with the toxicological effects in liver, kidneys, and immune system [52, 57, 58]. Researchers have developed some methods in order to reduce the harmful effects of grains contaminated with mycotoxins. These include physical (thermal and irradiation inactivation); chemical (ozonation and ammoniation); and, biological (bacterial degradation or adsorption [57, 59, 60]. Nevertheless, toxin sequestering agents are the most common and reliable products used for the feed industry due to its economic practicality and aptness for nutritional insight [61, 62]. Several studies have demonstrated that cellulosic materials have adsorption capacities for heavy metal ions and other pollutants [63, 64]. Similarly, some researchers have evaluated the binding activity of chitosan (CS) against several mycotoxins [2, 65]. As a biological polymer, chitosan has been shown to have promising uses as an adsorbent for the removal of various mycotoxins, heavy metal ions, and dyes [65]. Furthermore, it has been tested in the removal of OTA from contaminated drinks, demonstrating that chitosan can reduce the levels of this mycotoxin [1, 66]. On the other hand, some *in vitro* methods have been developed to evaluate the adsorbent capacity of mycotoxin sequestering products [67, 68]. However, these methods may not be directly applicable to poultry diets because they do not use the successive incubation at different pH and enzyme activity conditions similar to the different gastrointestinal compartments of poultry. Recently, we evaluated the adsorption capacity of CS on Aflatoxin B1 (AFB₁); Fumonisin B1 (FUB₁); Ochratoxin (OTA); Trichothecene (T-2); Deoxynivalenol (DON); and, Zearalenone (ZEA), using an *in vitro* digestive model that simulates three gastrointestinal compartment of poultry [69]. In that study, deacetylated 95%, high molecular weight (350 kDa) Chitosan (CS, Paragon Specialty Products, LLC, Rainsville, AL, USA) was tested and acetylated with an aqueous solution of acetic acid 1% (v/v). Then, this solution was dropped into NaOH




Figure 4. Percentage of adsorption of different mycotoxins on chitosan. Bars are the mean values. Error bars displays an interval around each mean, which are based on Fisher’s least significant difference (LSD) procedure.

0.5 M solution and the formed chitosan particles were rinsed three times with pure water and dried [69, 70]. The results showed a moderate adsorbent capacity of CS against five of the six mycotoxins evaluated, except for DON since only 3.5% was adsorbed, (**Figure 4**). Similar results were obtained in another study using non-crosslinked chitosan against different mycotoxins but it is a fact that cross-linking is related to a higher adsorption capacity and pH can affect it [70]. The mycotoxins adsorption capacity of CS is due to the electrostatic interactions. At alkaline pH, the CS is positively charged, while mycotoxins such as AFB1, FUB1, OTA and ZEA are negatively charged [70–72]. In the case of DON and T-2, the interactions appeared to be minor, causing poor adsorption. These results are very similar to those obtained in other studies [70]. Therefore, it could be said that ionic interactions are the main mechanism of mycotoxin adsorption of chitosan.

6. Chitosan nanocarriers: a strategy to improve solubility, permeability and stability of drugs

Another application of chitosan (CS) is its use in nanotechnology for the development of drug delivery systems such as nanoparticles and nanocapsules. These systems emerge as a strategy to improve the dissolution of drugs with low solubility and increase its permeability, which translates into an increase in bioavailability, a greater specificity and also an increase in the stability of drugs against physiological and environmental conditions [73]. In our laboratory, we have developed two nanocapsular systems capable of loading a phytopharmaceutical named Curcumin. This molecule has also been the subject of study in the poultry industry, given its properties, including its antioxidant action, the immunomodulatory, anticoccidial, anti-inflammatory, antimicrobial and growth promotion effects, the latter as an alternative to antibiotic growth promoters in order to maintain the performance and health of the birds [74–76]. In our laboratory, we have already evaluated the antimicrobial activity of curcumin against *Salmonella enteritidis* in an in vitro model that simulates the three compartments of the chicken gastrointestinal tract. The results obtained show that at a dose of 1%, the concentration of *Salmonella enteritidis* decreases slightly but not significantly with respect to the control [77]. However, one of the problems of curcumin, even when it is administered at high doses (12 g/day) is its low bioavailability due to its poor solubility and therefore poor absorption, as well as its rapid metabolism and systemic elimination [76]. In this sense, the development of nanocapsular systems aimed to increase the solubility, permeability and stability of curcumin. Such systems were named chitosan nanocapsules (NC-CS) and Alginate nanocapsules (NC-ALG) and were composed of an oily core of vitamin E surrounded by a biodegradable polymeric shell of either chitosan or alginate respectively.

Both systems were obtained by the a slightly modification of the solvent displacement technique [78]. However, the formation of NC-CS is based on the electrostatic and hydrophobic interactions as well as the hydrogen bonding and van der Waals forces that take place between the chitosan dissolved in an acidic aqueous phase and the lipid cores of Vitamin E formed in the organic phase, causing the polymer to be adsorbed on the surface of the lipid cores [4, 79–81], (**Figure 5(a)**). While NC-ALG were prepared using the “Single-stage procedure” based on the dipolar ionic interactions between the polymer (ALG), which is dissolved in the aqueous phase and the cationic surfactant (CTAB) present in the organic phase which also contains the oil [82], (**Figure 5(b)**).




Figure 5. Structural illustration of NC-CS (a), NC-ALG (b) and its components.

Formulation	Size (nm)	PDI	ζ potential (mV)	%E. E.	pH
NC-NC	116.7 ± 3.2	0.107	24.4 ± 2.1	>98	4.67 ± 0.08
NC-ALG	178 ± 7.9	0.149	-49.0 ± 2.3		6.08 ± 0.06

PDI: polydispersity index; EE: encapsulation efficiency. Values are given as mean ± SD; n = 3.

Table 8. Physicochemical characteristics of the nanocapsules obtained.

The systems were characterized physicochemically in terms of particle size, surface charge, polydispersity index (PDI) and curcumin encapsulation efficiency, (Table 8). Particle size and (PDI) were determined by dynamic light scattering (DLS) using a Malvern Zetasizer Nanoseries 3600 (Worcestershire, UK). The zeta potential values were calculated from the mean electrophoretic mobility values, as determined by Laser Doppler Velocimetry (LDV) using a Malvern Zetasizer Nanoseries 3600 (Worcestershire, UK). The particle size of NC-CS was round 116.7 nm with a PDI of 0.107 and presented positive surface charge (24.4 mV) while NC-ALG was round 178 nm with a PDI of 0.149 and a negative surface charge (-49.0 mV). Curcumin encapsulation efficiency was determined indirectly by Centrifugation-Filtration. Quantification of curcumin was performed by high performance liquid chromatography (HPLC, Merck-Hitachi, Japan) at 425 nm, using a reverse phase Hypersil® Division C8 column (150 × 3 mm, 5 μm; ThermoQuest, Hemel Hempstead, England). Curcumin encapsulation efficiency of both formulations, was >90%, with a final concentration of curcumin around 750 μg/ml [Unpublished work from our laboratory].

The stability to storage conditions is a parameter that must be evaluated in nanoparticulate systems (Table 9). In that study, the storage stability of NC-CS and NC-ALG was around 3 and 2 months respectively. In the case of NC-Cs, after 3 months of storage, the decrease in particle size and the precipitation of CUR were presented with greater magnitude since the chitosan begins to hydrolyze gradually and the viscosity of the formulation based on nanocapsules decreased during the storage period [83]. On the other hand, the results obtained for NC-ALG suggest that the stability of this type of formulation is around 2 months [Unpublished work from our laboratory]. These results are very similar to those reported in other studies, in which they report that the particle size of NC-ALG decreases between month 1 and 5 of storage [84].

Formulation	NC-CS			NC-ALG		
Time (months)	1	2	3	1	2	3
Z-average (nm)	115.2 ± 2.9	101.3 ± 4.1	95.45 ± 3.0	161.9 ± 2.1	157.7 ± 4.77	149.3 ± 2.9
PDI	0.115	0.145	0.196	0.178	0.213	0.245
ζ potential (mV)	23.8 ± 2.1	23.9 ± 3.2	24.5 ± 2.8	-48.7 ± 1.9	-46.8 ± 2.5	-44.3 ± 2.7
Values are given as the mean ± SD; n = 3.						

Table 9. Physicochemical characteristics of the nanocapsules during stability studies under storage conditions (4°C).

An important parameter that was take into account in these nanosystems was the cellular toxicity on caco-2 cells. For this, the conditions for the maintenance of the cell cultures were made according to Déat-Lainé et al. [85] with slight modifications. Before starting the study, the formulations were diluted in cell culture medium (DMEM: Dulbecco’s Modified Eagle Medium) in order to obtain the treatments with different polymer concentrations. Cell viability was determined by MTT assay [86]. In **Figure 6**, the results showed that even at high polymer concentrations (500 µg/mL) the cell viability is above 80%. However, it is a fact that the toxicity increases as so does the polymer concentration. Other studies in Caco-2 cells have shown similar results to those obtained in our laboratory and agree that the toxicity of chitosan nanoparticles is due to their physicochemical properties such as size and surface charge and also to the molecular weight of the chitosan and the concentration at which the cells are exposed [87, 88]. In the case of NC-ALG, the toxicity was lower since the interactions between the carboxyl groups of alginate and cell membranes are weaker because they are of the electrostatic type. The toxicity in these systems is more related to the particle size [89]. From the toxicity study, the polymer concentration to carry out the permeability studies was selected. It should be mentioned that this concentration did not compromise cellular viability.

Permeability studies were carried out on a monolayer of caco-2 cells and the quantification of curcumin was performed by UPLC-TQ-ESI-MS/MS (Waters ACQUITY UPLC system, Milford, MA, USA). Chromatographic analysis was performed on a Waters ACQUITY BEH Shield RP 18 column (2.1 × 100 mm, 1.7 µm). The polymer concentration used was 500 µg/mL of each polymer. Results show that the permeability of curcumin increased 28.6 and 14.6 times when it was in NC-CS and NC-ALG respectively, compared to the dispersion of curcumin in cell culture medium (DMEM: Dulbecco’s Modified Eagle Medium) [Unpublished work from our laboratory]. The increase in the permeability of curcumin in NC-CS is due to the ability of chitosan to temporarily open the tight junctions, which are related to a decrease in the value of transepithelial electrical resistance (TEER, MERSSTX01 electrode, Millicell ERS-2, Millipore, Billerica, MA, USA) (**Table 10**). The mechanism by which chitosan has this capacity is based on the interaction of its protonated amino groups with cell membranes, followed by a reversible structural reorganization of the binding proteins and a specific redistribution of the actin F cytoskeleton and the ZO-1 protein [90, 91]. Furthermore, it has been reported that particles positively charged, with spherical shape and with a monodisperse population have improved cellular uptake through the caveolae-mediated endocytosis and macropinocytosis pathway




Figure 6. Cell viability by the MTT assay on Caco-2 cells 2 h after the addition of NC-CS, NC-ALG and CUR at different concentrations. Values are given as mean \pm SD; n = 3.

Formulation	$P_{APP} \times 10^{-6}$ (cm/s)	R	TEER (%)			
			0 (h)	1 (h)	2(h)	12 (h)
CUR	4.96 ± 0.36	—	100	92 ± 1	93 ± 2	99 ± 4
NC-CS	141.60 ± 37.62	28.6*	100	81 ± 3	82 ± 4	97 ± 6
NC-ALG	72.38 ± 19.33	14.6*	100	90 ± 2	88 ± 6	100 ± 2

Values are given as the mean \pm SD;
n = 3.*p < 0.05 significantly different from CUR.

Table 10. Mean apparent permeability (Papp) and the absorption enhancement ratio (R) of NC-CS, NC-ALG and CUR across Caco-2 cells monolayers after 2 h incubation, as well as the values of transepithelial electrical resistance (TEER) determined at different times.

[65, 92]. Meanwhile, the mechanism of passage of NC-ALG through the monolayer of caco-2 cells depends largely on the particle size mainly. So, the main mechanisms are endocytic such as clathrin-mediated endocytosis, caveolae-mediated endocytosis and micropinocytosis [92, 93]. The results suggest that the use of NC-CS and NC-ALG to improve the bioavailability of curcumin is an interesting strategy to enhance the antimicrobial effect. Previous studies using an *in vitro* digestive model that simulates three gastrointestinal compartments of poultry have demonstrated that raw curcumin does not have good antimicrobial activity against *Salmonella enteritidis* [77]. However, when a solid dispersion of curcumin/PVP K30 was used, it decreased the concentration of *Salmonella enteritidis* more than 3 log in the compartment that simulates the intestine [Unpublished work from our laboratory]. Additional *in vivo* studies in 1-day-old chickens challenged with 10^4 CFU of *Salmonella enteritidis*/bird has shown that the solid dispersion of curcumin/PVP K30 administered in the feed at a concentration of 0.1% decreased more than 2 log the concentration of *Salmonella enteritidis* in ceca-cecal tonsil isolates [Unpublished

work from our laboratory]. Since nanocapsules increased the solubility and permeability of curcumin, the antimicrobial activity of nanocapsules loaded with curcumin developed in our laboratory is being carried out both *in vitro* and *in vivo* against *Salmonella enteritidis*.

7. Conclusion

As seen in this chapter, chitin and its derivatives, such as chitosan, are biopolymers with a wide variety of applications in different areas. Chitosan as a functional biopolymer has different properties. Some of these properties are its intrinsic nutritional value, such as antioxidant properties and health-promoting bioactivities against many chronic diseases, including hypercholesterolemia, hypertension, inflammation and immune diseases. In the case of chitin, its application is more limited given its poor solubility in aqueous medium, however, it has been reported that it has practically the same properties as its derivatives.

Every year millions of people are affected and thousands of them die due to infections and intoxication as a result of foodborne outbreaks, which also cause billions of dollars' worth of damage, public health problems and agricultural product loss. A considerable portion of these outbreaks is related to the consumption of contaminated products with foodborne pathogens and mycotoxins. In this sense, one of the main applications of chitosan is its antimicrobial effect against Gram-positive bacteria such as Gram-negative bacteria, having better activity with the latter due to the ionic interaction that takes place between the positively charged chitosan molecules and the negatively charged microbial cell membranes. Studies conducted on chickens and turkeys challenged with *Salmonella enteritidis* and typhimurium show the antimicrobial capacity of chitosan when it is administered in the feed. Furthermore, *in vitro* studies have demonstrated its properties as an adsorbent, since it can interact ionically with mycotoxins such as AFB1, FUB1, OTA and ZEA given that they are negatively charged, nevertheless, it is a fact that cross-linking is related to a higher adsorption capacity.

Finally, another application of chitosan is its use in nanotechnology for the development of nanoparticles and nanocapsules. These systems are an important strategy to improve the solubility, permeability and stability of molecules that are difficult to formulate. In the case of curcumin, a phytopharmaceutical that has become the subject of study in the poultry industry given its properties, including its antioxidant action, the immunomodulatory, anticoccidial, anti-inflammatory, antimicrobial and growth promotion effects, has problems of solubility and permeability, which causes low bioavailability. However, its association or encapsulation in nanoparticulate systems has shown that the solubility and permeability of this are improved. This suggests that the use of curcumin loaded in chitosan nanocapsules could increase its antimicrobial activity derived from the combination of the effects between chitosan and curcumin on different bacteria.

Acknowledgements

This research was supported by the Arkansas Bioscience Institute under the project: Development of an avian model for evaluation early enteric microbial colonization on the gastrointestinal tract and immune function.

The authors thank the CONACyT for the doctoral scholarship number 447447 and the financial support obtained through the program PAPIIT IN218115 of DGAPA-UNAM.

Author details

Daniel Hernandez-Patlan¹, Bruno Solis-Cruz¹, Billy M. Hargis² and Guillermo Tellez^{2*}

*Address all correspondence to: gtellez@uark.edu

1 Laboratorio 5: LEDEFAR, Unidad de Investigacion Multidisciplinaria, Facultad de Estudios Superiores (FES) Cuautitlan, Universidad Nacional Autonoma de Mexico (UNAM), Cuautitlan Izcalli, Estado de Mexico, Mexico

2 Department of Poultry Science, University of Arkansas, Fayetteville, Arkansas, USA

References

- [1] Bornet A, Teissedre PL. Chitosan, chitin-glucan and chitin effects on minerals (iron, lead, cadmium) and organic (ochratoxin A) contaminants in wines. *European Food Research and Technology*. 2008;**226**:681-689. DOI: 10.1007/s00217-007-0577-0
- [2] Szymczyk P, Filipkowska U, Józwiak T, Kuczajowska-Zadrożna M. Phosphate removal from aqueous solutions by chitin and chitosan in flakes. *Progress on Chemistry and Application of Chitin and its Derivatives*. 2016;**21**:260-272
- [3] Rinaudo M. Chitin and chitosan: Properties and applications. *Progress in Polymer Science*. 2006;**31**:603-632
- [4] Ahmed TA, Aljaeid BM. Preparation, characterization, and potential application of chitosan, chitosan derivatives, and chitosan metal nanoparticles in pharmaceutical drug delivery. *Drug Design, Development and Therapy*. 2016;**10**:483
- [5] Kumar MNVR. A review of chitin and chitosan applications. *Reactive and Functional Polymers*. 2000;**46**:1-27
- [6] Şenel S, McClure SJ. Potential applications of chitosan in veterinary medicine. *Advanced Drug Delivery Reviews*. 2004;**56**:1467-1480
- [7] Kong M, Chen XG, Xing K, Park HJ. Antimicrobial properties of chitosan and mode of action: A state of the art review. *International Journal of Food Microbiology*. 2010;**144**:51-63
- [8] Helander IM, Nurmiäho-Lassila E-L, Ahvenainen R, Rhoades J, Roller S. Chitosan disrupts the barrier properties of the outer membrane of Gram-negative bacteria. *International Journal of Food Microbiology*. 2001;**71**:235-244
- [9] Chung Y-C, Su YP, Chen C-C, Jia G, Wang HL, Wu JCG, Lin JG. Relationship between antibacterial activity of chitosan and surface characteristics of cell wall. *Acta Pharmacologica Sinica*. 2004;**25**:932-936

- [10] Filipkowska U, Józwiak T, Szymczyk P. Application of cross-linked chitosan for phosphate removal from aqueous solutions. *Progress on Chemistry and Application of Chitin and its Derivatives*. 2014;**19**:5-14
- [11] Luo Y, Wang Q. Recent advances of chitosan and its derivatives for novel applications in food science. *Journal of Food Processing & Beverages*. 2013;**1**:1-13
- [12] Rao SB, Sharma CP. Use of chitosan as a biomaterial: Studies on its safety and hemostatic potential. *Journal of Biomedical Materials Research*. 1997;**34**:21-28
- [13] Muzzarelli RAA. Chitins and chitosans as immunoadjuvants and non-allergenic drug carriers. *Marine Drugs*. 2010;**8**:292-312
- [14] Divya K, Vijayan S, George TK, Jisha MS. Antimicrobial properties of chitosan nanoparticles: Mode of action and factors affecting activity. *Fibers and Polymers*. 2017;**18**:221-230
- [15] Romainor ANB, Chin SF, Pang SC, Bilung LM. Preparation and characterization of chitosan nanoparticles-doped cellulose films with antimicrobial property. *Journal of Nanomaterials*. 2014;**2014**:130
- [16] Boyle EC, Bishop JL, Grassl GA, Finlay BB. *Salmonella*: From pathogenesis to therapeutics. *Journal of Bacteriology*. 2007;**189**:1489-1495
- [17] Antunes P, Mourão J, Campos J, Peixe L. Salmonellosis: The role of poultry meat. *Clinical Microbiology and Infection*. 2016;**22**:110-121
- [18] Naksuriya O, Okonogi S. Comparison and combination effects on antioxidant power of curcumin with gallic acid, ascorbic acid, and xanthone. *Drug Discoveries & Therapeutics*. 2015;**9**:136-141
- [19] Menconi A, Hernandez-Velasco X, Latorre JD, Kallapura G, Pumford NR, Morgan MJ, Hargis BM, Tellez G. Effect of chitosan as a biological sanitizer for *Salmonella typhimurium* and aerobic Gram negative spoilage bacteria present on chicken skin. *International Journal of Poultry Science*. 2013;**12**:318
- [20] Menconi A, Pumford NR, Morgan MJ, Bielke LR, Kallapura G, Latorre JD, Wolfenden AD, Hernandez-Velasco X, Hargis BM, Tellez G. Effect of chitosan on *Salmonella typhimurium* in broiler chickens. *Foodborne Pathogens and Disease*. 2014;**11**:165-169
- [21] Corrier DE, Byrd JA, Hargis BM, Hume ME, Bailey RH, Stanker LH. Presence of *Salmonella* in the crop and ceca of broiler chickens before and after preslaughter feed withdrawal. *Poultry Science*. 1999;**78**:45-49
- [22] Byrd JA, Hargis BM, Caldwell DJ, Bailey RH, Herron KL, McReynolds JL, Brewer RL, Anderson RC, Bischoff KM, Callaway TR. Effect of lactic acid administration in the drinking water during preslaughter feed withdrawal on *Salmonella* and *Campylobacter* contamination of broilers. *Poultry Science*. 2001;**80**:278-283
- [23] Cox JM, Pavic A. Advances in enteropathogen control in poultry production. *Journal of Applied Microbiology*. 2010;**108**:745-755

- [24] Choi EH, Yang HP, Chun HS. Chitooligosaccharide ameliorates diet-induced obesity in mice and affects adipose gene expression involved in adipogenesis and inflammation. *Nutrition Research*. 2012;**32**:218-228
- [25] Agnihotri SA, Mallikarjuna NN, Aminabhavi TM. Recent advances on chitosan-based micro- and nanoparticles in drug delivery. *Journal of Controlled Release*. 2004;**100**:5-28
- [26] Gibson GR, Roberfroid MB. Dietary modulation of the human colonic microbiota: Introducing the concept of prebiotics. *The Journal of Nutrition*. 1995;**125**:1401
- [27] Liu X, Cao S, Zhang X. Modulation of gut microbiota-brain axis by probiotics, prebiotics, and diet. *Journal of Agricultural and Food Chemistry*. 2015;**63**:7885-7895
- [28] Torres-Rodriguez A, Sartor C, Higgins SE, Wolfenden AD, Bielke LR, Pixley CM, Sutton L, Tellez G, Hargis BM. Effect of *Aspergillus* meal prebiotic (Fermacto) on performance of broiler chickens in the starter phase and fed low protein diets. *Journal of Applied Poultry Research*. 2005;**14**:665-669
- [29] Mizutani O, Shiina M, Yoshimi A, Sano M, Watanabe T, Yamagata Y, Nakajima T, Gomi K, Abe K. Substantial decrease in cell wall α -1, 3-glucan caused by disruption of the *kexB* gene encoding a subtilisin-like processing protease in *Aspergillus oryzae*. *Bioscience, Biotechnology, and Biochemistry*. 2016;**80**:1781-1791
- [30] Guio F, Rugeles LD, Rojas SE, Palomino MP, Camargo MC, Sánchez OF. Kinetic modeling of fructooligosaccharide production using *Aspergillus oryzae* N74. *Applied Biochemistry and Biotechnology*. 2012;**167**:142-163
- [31] Bays HE, Evans JL, Maki KC, Evans M, Maquet V, Cooper R, Anderson JW. Chitin-glucan fiber effects on oxidized low-density lipoprotein: A randomized controlled trial. *European Journal of Clinical Nutrition*. 2013;**67**:2-7
- [32] Uchima CA, Tokuda G, Watanabe H, Kitamoto K, Arioka M. Heterologous expression and characterization of a glucose-stimulated β -glucosidase from the termite *Neotermes kosshunensis* in *Aspergillus oryzae*. *Applied Microbiology and Biotechnology*. 2011;**89**:1761-1771
- [33] Cox E, Verdonck F, Vanrompay D, Goddeeris B. Adjuvants modulating mucosal immune responses or directing systemic responses towards the mucosa. *Veterinary Research*. 2006;**37**:511-539
- [34] Corrigan A, Horgan K, Clipson N, Murphy RA. Effect of dietary supplementation with a *Saccharomyces cerevisiae* mannan oligosaccharide on the bacterial community structure of broiler cecal contents. *Applied and Environmental Microbiology*. 2011;**77**:6653-6662
- [35] Janssens GP j, Millet S, Van Immerseel F, De Buck J, Hesta M. The impact of prebiotics and salmonellosis on apparent nutrient digestibility and *Salmonella typhimurium* var. Copenhagen excretion in adult pigeons (*Columba livia domestica*). *Poultry Science*. 2004;**83**:1884-1890
- [36] Londero A, Menconi A, Reginatto AR, Bacocina AI, Wolfenden A, Shivaramaiah S, Hargis BM, Tellez G. Effect of an *Aspergillus* meal prebiotic on salmonella infection in turkeys and broiler chickens. *International Journal of Poultry Science*. 2011;**10**:946-951

- [37] Lowry VK, Farnell MB, Ferro PJ, Swaggerty CL, Bahl A, Kogut MH. Purified β -glucan as an abiotic feed additive up-regulates the innate immune response in immature chickens against *Salmonella enterica* serovar Enteritidis. *International Journal of Food Microbiology*. 2005;**98**:309-318
- [38] Xu ZR, Hu CH, Xia MS, Zhan XA, Wang MQ. Effects of dietary fructooligosaccharide on digestive enzyme activities, intestinal microflora and morphology of male broilers. *Poultry Science*. 2003;**82**:1030-1036
- [39] Dahiya JP, Wilkie DC, Van Kessel AG, Drew MD. Potential strategies for controlling necrotic enteritis in broiler chickens in post-antibiotic era. *Animal Feed Science and Technology*. 2006;**129**:60-88
- [40] Kim HG, Lee SY, Kim NR, Lee HY, Ko MY, Jung BJ, Kim CM, Lee JM, Park JH, Han SH. *Lactobacillus plantarum* lipoteichoic acid down-regulated *Shigella flexneri* peptidoglycan-induced inflammation. *Molecular Immunology*. 2011;**48**:382-391
- [41] Huang RL, Yin YL, Wu GY, Zhang YG, Li TJ, Li LL, Li MX, Tang ZR, Zhang J, Wang B. Effect of dietary oligochitosan supplementation on ileal digestibility of nutrients and performance in broilers. *Poultry Science*. 2005;**84**:1383-1388
- [42] Merino-Guzmán R, Latorre JD, Delgado R, Hernandez-Velasco X, Wolfenden AD, Teague KD, Graham LE, Mahaffey BD, Baxter MFA, Hargis BM. Comparison of total immunoglobulin A levels in different samples in leghorn and broiler chickens. *Asian Pacific Journal of Tropical Biomedicine*. 2017;**7**:116-120
- [43] Beal RK, Wigley P, Powers C, Hulme SD, Barrow PA, Smith AL. Age at primary infection with *Salmonella enterica* serovar Typhimurium in the chicken influences persistence of infection and subsequent immunity to re-challenge. *Veterinary Immunology and Immunopathology*. 2004;**100**:151-164
- [44] Harms RH, Miles RD. Research note: Influence of Fermacto® on the performance of laying hens when fed diets with different levels of methionine. *Poultry Science*. 1988;**67**:842-844
- [45] Tellez G, Nava GM, Vincente JL, De Franceschi M, Morales EJ, Prado O, Hargis BM. Evaluation of dietary *Aspergillus* meal on intestinal morphometry in turkey poults. *International Journal of Poultry Science*. 2010;**9**:75-878
- [46] Reginatto AR, Menconi A, Londero A, Lovato M, Rosa AP, Shivaramaiah S, Wolfenden AD, Huff WE, Huff GR, Rath NC. Effects of dietary *Aspergillus* meal prebiotic on turkey poults production parameters and bone qualities. *International Journal of Poultry Science*. 2011;**10**:496-499
- [47] Uni Z, Ganot S, Sklan D. Posthatch development of mucosal function in the broiler small intestine. *Poultry Science*. 1998;**77**:75-82
- [48] Tellez G, Higgins SE, Donoghue AM, Hargis BM. Digestive physiology and the role of microorganisms. *Journal of Applied Poultry Research*. 2006;**15**:136-144
- [49] Latorre JD, Hernandez-Velasco X, Kuttappan VA, Wolfenden RE, Vicente JL, Wolfenden AD, Bielke LR, Prado-Rebolledo OF, Morales E, Hargis BM. Selection of *Bacillus* spp. for

cellulase and xylanase production as direct-fed microbials to reduce digesta viscosity and *Clostridium perfringens* proliferation using an in vitro digestive model in different poultry diets. *Frontiers in Veterinary Science*. 2015;**2**:25

- [50] Latorre JD, Hernandez-Velasco X, Wolfenden RE, Vicente JL, Wolfenden AD, Menconi A, Bielke LR, Hargis BM, Tellez G. Evaluation and selection of *Bacillus* species based on enzyme production, antimicrobial activity, and biofilm synthesis as direct-fed microbial candidates for poultry. *Frontiers in Veterinary Science*. 2016;**3**:95
- [51] Latorre JD, Hernandez-Velasco X, Vicente JL, Wolfenden R, Hargis BM, Tellez G. Effects of the inclusion of a *Bacillus* direct-fed microbial on performance parameters, bone quality, recovered gut microflora, and intestinal morphology in broilers consuming a grower diet containing corn distillers dried grains with solubles. *Poultry Science*. 2017;**96**:2728-2735
- [52] Hussein HS, Brasel JM. Toxicity, metabolism, and impact of mycotoxins on humans and animals. *Toxicology*. 2001;**167**:101-134
- [53] Zain ME. Impact of mycotoxins on humans and animals. *Journal of Saudi Chemical Society*. 2011;**15**:129-144
- [54] Armando MR, Pizzolitto RP, Dogi CA, Cristofolini A, Merkis C, Poloni V, Dalcero AM, Cavaglieri LR. Adsorption of ochratoxin A and zearalenone by potential probiotic *Saccharomyces cerevisiae* strains and its relation with cell wall thickness. *Journal of Applied Microbiology*. 2012;**113**:256-264
- [55] Streit E, Schatzmayr G, Tassis P, Tzika E, Marin D, Taranu I, Tabuc C, Nicolau A, Aprodu I, Puel O. Current situation of mycotoxin contamination and co-occurrence in animal feed – Focus on Europe. *Toxins (Basel)*. 2012;**4**:788-809
- [56] Smith LE, Stoltzfus RJ, Prendergast A. Food chain mycotoxin exposure, gut health, and impaired growth: A conceptual framework. *Advances in Nutrition: An International Review*. 2012;**3**:526-531
- [57] Greco MV, Franchi ML, Rico Golba SL, Pardo AG, Pose GN. Mycotoxins and mycotoxigenic fungi in poultry feed for food-producing animals. *Scientific World Journal*. 2014;**2014**:1-9
- [58] Galarza-Seeber R, Latorre JD, Bielke LR, Kuttappan VA, Wolfenden AD, Hernandez-Velasco X, Merino-Guzman R, Vicente JL, Donoghue A, Cross D. Leaky gut and mycotoxins: Aflatoxin B1 does not increase gut permeability in broiler chickens. *Frontiers in Veterinary Science*. 2016;**3**:10
- [59] Jouany JP. Methods for preventing, decontaminating and minimizing the toxicity of mycotoxins in feeds. *Animal Feed Science and Technology*. 2007;**137**:342-362
- [60] Kolosova A, Stroka J. Evaluation of the effect of mycotoxin binders in animal feed on the analytical performance of standardised methods for the determination of mycotoxins in feed. *Food Additives & Contaminants: Part A*. 2012;**29**:1959-1971
- [61] Avantaggiato G, Solfrizzo M, Visconti A. Recent advances on the use of adsorbent materials for detoxification of *Fusarium* mycotoxins. *Food Additives and Contaminants*. 2005;**22**:379-388

- [62] Di Natale F, Gallo M, Nigro R. Adsorbents selection for aflatoxins removal in bovine milks. *Journal of Food Engineering*. 2009;**95**:186-191
- [63] Hokkanen S, Bhatnagar A, Sillanpää M. A review on modification methods to cellulose-based adsorbents to improve adsorption capacity. *Water Research*. 2016;**91**:156-173
- [64] Tan KB, Abdullah AZ, Horri BA, Salamatinia B. Adsorption mechanism of microcrystalline cellulose as green adsorbent for the removal of cationic methylene blue dye. *Journal of the Chemical Society of Pakistan*. 2016;**38**:651-664
- [65] Zhao L, Yang G, Shi Y, Su C, Chang J. Co-delivery of Gefitinib and chloroquine by chitosan nanoparticles for overcoming the drug acquired resistance. *Journal of Nanobiotechnology*. 2015;**13**:57. DOI: 10.1186/s12951-015-0121-5
- [66] Mine Kurtbay H, Bekçi Z, Merdivan M, Yurdakoç K. Reduction of ochratoxin A levels in red wine by bentonite, modified bentonites, and chitosan. *Journal of Agricultural and Food Chemistry*. 2008;**56**:2541-2545
- [67] Ledoux DR, Rottinghaus GE. In vitro and in vivo testing of adsorbents for detoxifying mycotoxins in contaminated feedstuffs. *Biotechnology Feed Industry*. Nottingham, UK: Nottingham University Press; 1999. pp. 369-379
- [68] Kong C, Shin SY, Kim BG. Evaluation of mycotoxin sequestering agents for aflatoxin and deoxynivalenol: An in vitro approach. *Spring*. 2014;**3**:346
- [69] Solís-Cruz B, Hernández-Patlán D, Beyssac E, Latorre JD, Hernandez-Velasco X, Merino-Guzman R, Tellez G, López-Arellano R. Evaluation of chitosan and cellulosic polymers as binding adsorbent materials to prevent aflatoxin B1, fumonisin B1, ochratoxin, trichothecene, deoxynivalenol, and zearalenone mycotoxicoses through an in vitro gastrointestinal model for poultry. *Polymers (Basel)*. 2017;**9**:529
- [70] Zhao Z, Liu N, Yang L, Wang J, Song S, Nie D, Yang X, Hou J, Wu A. Cross-linked chitosan polymers as generic adsorbents for simultaneous adsorption of multiple mycotoxins. *Food Control*. 2015;**57**:362-369
- [71] Daković A, Tomašević-Čanović M, Rottinghaus GE, Matijašević S, Sekulić Ž. Fumonisin B 1 adsorption to octadecyldimethylbenzyl ammonium-modified clinoptilolite-rich zeolitic tuff. *Microporous and Mesoporous Materials*. 2007;**105**:285-290
- [72] Bazin I, Faucet-Marquis V, Monje M-C, El Khoury M, Marty J-L, Pfohl-Leszkowicz A. Impact of pH on the stability and the cross-reactivity of ochratoxin A and citrinin. *Toxins (Basel)*. 2013;**5**:2324-2340
- [73] Ghadi A, Mahjoub S, Tabandeh F, Talebnia F. Synthesis and optimization of chitosan nanoparticles: Potential applications in nanomedicine and biomedical engineering. *Caspian Journal of Internal Medicine*. 2014;**5**:156
- [74] Khan RU, Naz S, Javdani M, Nikousefat Z, Selvaggi M, Tufarelli V, Laudadio V. The use of turmeric (*Curcuma longa*) in poultry feed. *World's Poultry Science Journal*. 2012;**68**:97-103
- [75] Khalafalla RE, Müller U, Shahiduzzaman M, Dyachenko V, Desouky AY, Alber G, Daugschies A. Effects of curcumin (diferuloylmethane) on *Eimeria tenella* sporozoites in vitro. *Parasitology Research*. 2011;**108**:879-886

- [76] Anand P, Kunnumakkara AB, Newman RA, Aggarwal BB. Bioavailability of curcumin: Problems and promises. *Molecular Pharmaceutics*. 2007;**4**:807-818
- [77] Hernandez-Patlan D, Solis-Cruz B, Méndez-Albores A, Latorre JD, Hernandez-Velasco X, Tellez G, López-Arellano R. Comparison of PrestoBlue® and plating method to evaluate antimicrobial activity of ascorbic acid, boric acid and curcumin in an in vitro gastrointestinal model. *Journal of Applied Microbiology*. 2017;**124**:423-430. DOI: 10.1111/jam.13659
- [78] Rivera-Rodriguez GR, Lollo G, Montier T, Benoit JP, Passirani C, Alonso MJ, Torres D. In vivo evaluation of poly-L-asparagine nanocapsules as carriers for anti-cancer drug delivery. *International Journal of Pharmaceutics*. 2013;**458**:83-89. DOI: 10.1016/j.ijpharm.2013.09.038
- [79] Alishahi A, Mirvaghefi A, Tehrani MR, Farahmand H, Shojaosadati SA, Dorkoosh FA, Elsabee MZ. Shelf life and delivery enhancement of vitamin C using chitosan nanoparticles. *Food Chemistry*. 2011;**126**:935-940
- [80] Luo Y, Teng Z, Wang Q. Development of zein nanoparticles coated with carboxymethyl chitosan for encapsulation and controlled release of vitamin D3. *Journal of Agricultural and Food Chemistry*. 2012;**60**:836-843
- [81] Dimzon IKD, Ebert J, Knepper TP. The interaction of chitosan and olive oil: Effects of degree of deacetylation and degree of polymerization. *Carbohydrate Polymers*. 2013;**92**:564-570
- [82] Oyarzun-Ampuero FA, Rivera-Rodríguez GR, Alonso MJ, Torres D. Hyaluronan nanocapsules as a new vehicle for intracellular drug delivery. *European Journal of Pharmaceutical Sciences*. 2013;**49**:483-490
- [83] Rokhati N, Widjajanti P, Pramudono B, Susanto H. Performance comparison of α - and β -amylases on chitosan hydrolysis. *ISRN Chemical Engineering*. 2013;**2013**:186159
- [84] Ma H, Qi X, Maitani Y, Nagai T. Preparation and characterization of superparamagnetic iron oxide nanoparticles stabilized by alginate. *International Journal of Pharmaceutics*. 2007;**333**:177-186. DOI: 10.1016/j.ijpharm.2006.10.006
- [85] Déat-Lainé E, Hoffart V, Garrait G, Beyssac E. Whey protein and alginate hydrogel microparticles for insulin intestinal absorption: Evaluation of permeability enhancement properties on Caco-2 cells. *International Journal of Pharmaceutics*. 2013;**453**:336-342
- [86] Vázquez-Durán A, Díaz-Torres R, Ramírez-Noguera P, Moreno-Martínez E, Méndez-Albores A. Cytotoxic and genotoxic evaluation of tortillas produced by microwave heating during alkaline-cooking of aflatoxin-contaminated maize. *Journal of Food Science*. 2014;**79**:T1024-T1029
- [87] Schipper NG, Varum KM, Artursson P. Chitosans as absorption enhancers for poorly absorbable drugs. 1: Influence of molecular weight and degree of acetylation on drug transport across human intestinal epithelial (Caco-2) cells. *Pharmaceutical Research*. 1996;**13**:1686-1692

- [88] Prego C, Torres D, Fernandez-Megia E, Novoa-Carballal R, Quiñoá E, Alonso MJ. Chitosan-PEG nanocapsules as new carriers for oral peptide delivery: Effect of chitosan pegylation degree. *Journal of Controlled Release*. 2006;**111**:299-308
- [89] Xiang Y, Liu Y, Mi B, Leng Y. Hydrated polyamide membrane and its interaction with alginate: A molecular dynamics study. *Langmuir*. 2013;**29**:11600-11608. DOI: 10.1021/la401442r
- [90] Van der Lubben IM, Verhoef JC, Borchard G, Junginger HE. Chitosan and its derivatives in mucosal drug and vaccine delivery. *European Journal of Pharmaceutical Sciences*. 2001;**14**:201-207. DOI: 10.1016/S0928-0987(01)00172-5
- [91] Amidi M, Mastrobattista E, Jiskoot W, Hennink WE. Chitosan-based delivery systems for protein therapeutics and antigens. *Advanced Drug Delivery Reviews*. 2010;**62**:59-82. DOI: 10.1016/j.addr.2009.11.009
- [92] Salatin S, Yari Khosroushahi A. Overviews on the cellular uptake mechanism of polysaccharide colloidal nanoparticles. *Journal of Cellular and Molecular Medicine*. 2017;**21**:1668-1686. DOI: 10.1111/jcmm.13110
- [93] Li Q, Liu C-G, Yu Y. Separation of monodisperse alginate nanoparticles and effect of particle size on transport of vitamin E. *Carbohydrate Polymers*. 2015;**124**:274-279. DOI: 10.1016/j.carbpol.2015.02.007

IntechOpen