

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

185,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



New Approaches for Chagas' Disease Chemotherapy

Guadalupe García Liñares

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/intechopen.77235>

Abstract

The latest advances concerning drug design and chemotherapy development to combat the Chagas' disease are discussed. This chapter is based on the metabolic differences between the pathogenic parasite and mammal hosts that led to the progress in the search for novel metabolic pathways in parasites that may be essential for parasite's survival but with no counterpart in the host. There is a considerable amount of work in the search of more promising molecular targets for drug design. However, the chemotherapy for this disease remains unsolved. It is based on old and fairly not specific drugs associated with long-term treatments, severe side effects, drug resistance, and different strains' susceptibility. Herein, a thorough analysis of selected molecular targets is described in terms of their potential usefulness for drug design. Therefore, rational approaches to the chemotherapeutic control of American trypanosomiasis describing some useful metabolic pathways are covered. Enzymes involved in ergosterol biosynthesis (squalene synthase, HMG-CoA reductase, farnesyl diphosphate synthase (FPPS), sterol 24-methyltransferase, and sterol 14 α -demethylase), trypanothione system (glutathionyl-spermidine synthetase, trypanothione synthetase, and trypanothione reductase), cysteine proteases, trans-sialidase, and so on are discussed. The design of specific inhibitors of these metabolic activities as possible means of controlling the parasites without damaging the hosts is presented.

Keywords: *Trypanosoma cruzi*, drug development, molecular targets

1. Introduction

Chagas' disease or American trypanosomiasis is among the most prevalent parasitic diseases worldwide [1–3]. It has been estimated that around 20 million people are infected and over 40 million individuals are facing the risk of infection by the hemoflagellates protozoan *Trypanosoma cruzi*, the responsible agent of Chagas' disease. This disease is endemic in Latin

American countries, but the migration of individuals and blood transfusions have made possible the occurrence of the Chagas' disease in developed countries. Currently, no vaccine is available, so the control of the disease is limited to its detection, vector control, screening of blood banks and organ donors, and case finding of infected pregnant women. Moreover, the development of a safer and more effective chemotherapeutic intervention is crucial for the treatment of Chagas disease.

This disease is characterized by three phases: (1) an early acute phase in which trypomastigotes circulate in blood and infect cells transforming into the asexually-multiplying amastigotes; (2) a subsequent intermediate phase where the cells are broken, parasites are released to the blood and infect other cells (in this phase there are unspecific symptoms like fever, allergic reactions, acute heart failure, or meningoencephalitis); (3) a late chronic phase, a prolonged, and asymptomatic indeterminate phase, where parasites establish in their target organs (during this stage patients may have nonspecific clinical manifestations or present major complications such as cardiomyopathy and/or megaesophagus and megacolon syndromes [4]. The absence of adequate treatment in the acute phase results in the development of the above-mentioned stages of the disease.

The existing chemotherapy remains deficient; it is based on two old drugs empirically discovered, (1) nifurtimox, actually discontinued, and (2) benznidazole (**Figure 1**). Although both of these compounds are able to cure at least 50% of recent infections, they present important drawbacks such as selective drug sensitivity on different *T. cruzi* strains, serious side effects including vomiting, anorexia, peripheral neuropathy, allergic dermatopathy, and long-term treatment [2, 5, 6]. Moreover, these compounds are not effective in the chronic stage of the disease. Consequently, the development of novel, safe, and affordable compounds with potent antiparasitic activity is urgently needed [7, 8]. The existence of *T. cruzi* populations naturally resistant to benznidazole and nifurtimox led to the search for compounds with a different mechanism of action [2, 8]. The development of new drugs that are more effective and safer than those currently available is urgently necessary.

There is a considerable amount of work in the search of unique aspects of the biochemistry and physiology of *T. cruzi* intending to find specific molecular targets for drug design [9]. It can be thought that the selective inhibition of a biosynthetic pathway that leads to a crucial metabolite for parasite survival would not have any significant toxic effect on the host. Based on these facts, this chapter will discuss the search for new approaches based on metabolic differences between the pathogenic parasite and mammal hosts and the development of new potential antiparasitic drugs in the last years.



Figure 1. Drugs currently used for Chagas' disease treatment.

2. Molecular targets

There are many metabolic pathways and enzymes unique to *Trypanosoma cruzi* that constitute excellent molecular targets for drug development. However, despite the specificity of new compounds targeting parasite molecules, the effect of these drugs on mammalian metabolism must be carefully evaluated. A few of the most studied compounds targeting specific *T. cruzi* enzymes will be reviewed here.

2.1. Sterol biosynthesis

Isoprenoid biosynthetic pathway constitutes one of the most important metabolic pathways of all organisms because isoprenoids are essential for numerous biochemical functions. In trypanosomatids diverse enzymes of this biosynthetic pathway are involved in key process. Sterol biosynthesis in parasites differs from that in mammalian hosts since the final product is ergosterol instead cholesterol, the main sterol present in the mammals. As *T. cruzi* is entirely dependent on endogenously produced sterols for survival and proliferation, the sterol biosynthetic pathway constitutes an attractive target for drug development.

Reduction of endogenous sterols induces inhibition of the multiplication of *T. cruzi*. Then, the restriction of an enzyme of this biosynthetic pathway will inhibit the growth of the parasite [10, 11]. Sterol composition in *T. cruzi* is very similar to fungi with ergosterol and 24-ethylergosterol being the primary mature sterols in the epimastigote stage. Fungisterol and 24-ethylfungisterol are the major sterols produced by the amastigote stage. In consequence, antifungal drugs are potentially capable of decreasing pathogen growth.

There are several interesting enzymes in the pathway as potential targets for anti-trypanosomal chemotherapy. For example, sterol 14 α -demethylase, sterol 24-methyltransferase, farnesyl diphosphate synthase, squalene synthase, and HMG-CoA reductase.

2.1.1. Sterol 14 α -demethylase (CYP51)

Sterol 14 α -demethylase is a CYP monooxygenase that catalyzes the removal of the 14 α -methyl group from eburicol. Unlike other hemoproteins, the hemo cofactor in CYP51 is coordinated to cysteine residue instead of histidine. Since it is an essential enzyme in sterol biosynthesis, the activity inhibition could be lethal in organisms requiring sterols for membrane function [12]. This enzyme constitutes an interesting target as it has the advantage of being inhibited by antifungal agents currently in clinical use. Azoles are the most efficient antifungal drugs and there have been reported numerous examples of its antiparasitic effects [13–15].

The first azoles such as ketoconazole (3), miconazole (4), or fluconazole (5) (**Figure 2**) were found to have potent *in vitro* activity but did not cure the *T. cruzi* infection. The mechanism of action involves binding to *T. cruzi* CYP51 and a disruption of sterol biosynthesis resulting in accumulation of 14-methylated sterols [16]. New azole drugs developed to combat fungal infections have been evaluated for the activity against *T. cruzi*. The experimental azole drug D0870 (6) was the first to cure the chronic infection in mice [17]. Unfortunately, D0870 was discontinued as an antifungal agent due to undesired side effects.




Figure 2. Structures of sterol 14 α -demethylase (CYP51) inhibitors.

Among second-generation azoles, posaconazole (7) is the most potent drug against *T. cruzi*; because of its broad-spectrum antifungal activity, it could potentially be repurposed for use in Chagas' disease (**Figure 2**) [14]. It has been demonstrated that posaconazole is curative in the chronic murine model. In addition, it has potent activity against benznidazole and nifurtimox resistant *T. cruzi* strains [18]. Recently, a case was reported in Spain, describing the cure of chronic infection by treatment with posaconazole in an immunosuppressed patient. This compound is currently in phase II clinical trial for Chagas' disease. [19, 20] Unfortunately, posaconazole is a very expensive drug, so its application becomes impractical for patients with limited resources.

Ravuconazole (8) is another azole antifungal drug that has been evaluated as an antiparasitic agent and it appeared to be very efficient in restraining the parasitemia *in vitro* against *T. cruzi* in murine models [21]. However, since the half-life is much longer in humans than in mice, it is possible that this drug has curative effects. In fact, a prodrug of ravuconazole (E1224 9) is in a phase II trial [20]. In addition, ravuconazole has a simpler chemical structure than posaconazole, so the cost might be lower (**Figure 2**).

Tipifarnib (10) is an antitumor agent inhibiting human protein farnesyl transferase (**Figure 2**). It has considerable *in vitro* activity against *T. cruzi*. It was determined that the drug produces restriction of sterol biosynthesis by inhibition of CYP51 [22]. Analogs of tipifarnib have been designed and synthesized with improved CYP51 inhibitory activity and excellent pharmacokinetic properties [23].

2.1.2. Sterol 24-methyltransferase (24-SMT)

This enzyme catalyzes the methenylation of zymosterol, an important step for the biosynthesis of ergosterol and other related 24-alkylated sterols, which are the main sterols

2.1.3. 3-Hydroxy-3-methyl-glutaryl-coenzyme a reductase (HMG-CoA reductase)

Current cardiovascular drugs have been tested for the treatment of Chagas' disease but new therapeutic drugs based on statins with a new anti-inflammatory approach have arisen as potential antiparasitic agents. Statins are thought to be associated with their ability to reduce cholesterol synthesis [28]. It was observed that lovastatin (**14**) and simvastatin (**15**) (**Figure 4**) have inhibited the growth of epimastigotes of *T. cruzi* and simvastatin could potentially inhibit HMG-CoA reductase both in epimastigotes and trypomastigotes [29]. Moreover, the combination of lovastatin with ketoconazole allowed the elimination of the presence of parasites into the blood flow and, in this way, prevented host death [30].

The enzyme farnesyl pyrophosphate synthase, also known as farnesyl diphosphate synthase (FPPS) belongs to the E-family of the prenyltransferases and it has a key role in the isoprenoid biosynthetic pathway. Isopentenyl diphosphate (IPP) and its isomer dimethylallyl diphosphate (DMAPP) are synthesized via mevalonate pathway from acetyl-CoA [31]. DMAPP is the precursor for the biosynthesis of different and very important isoprenoids like sterols, ubiquinones, triterpenoids, and prenylated proteins.




Figure 3. General structures of inhibitors of 24-SMT.




Figure 4. Chemical structures of inhibitors of the enzymatic activity of HMG-CoA reductase.

FPPS catalyzes two sequential steps: the addition of DMAPP to isopentenyl diphosphate (IPP) to form geranyl diphosphate (GPP) and the addition of DMAPP to geranyl diphosphate to produce farnesyl pyrophosphate (FPP). The inhibition or alteration of FPPS activity can regulate the isoprenoid metabolism. Therefore, FPPS has been selected as an excellent target for different disorders and anticancer, antibacterial, and antiparasitic drugs design among others [32–34].

Bisphosphonates act as inhibitors of bone resorption binding to the bone mineral. They are currently used for the treatment of several bone disorders like osteoporosis, Paget's disease, and hypercalcemia [35]. Bisphosphonates were the first FPPS inhibitors known and were reported as potent antiparasitic agents. Nitrogen-containing bisphosphonates like pamidronate (**16**), alendronate (**17**), risedronate (**18**), and ibandronate (**19**) were originally found to be effective against *T. cruzi* without toxicity to the host cells (**Figure 5**) [36]. Risedronate has shown a significantly increased survival of *T. cruzi*-infected mice *in vivo*. It was also found that diverse bisphosphonate derivatives were effective growth inhibitors of other pathogenic trypanosomatids and apicomplexan parasites [32, 37, 38].

Bisphosphonates have the disadvantage that they are highly polar and are rapidly removed from the circulatory system. Therefore, more lipophilic derivatives were developed (**20**) [38–40]. Bisphosphonates have a great potentiality as antiparasitic agents with characterized mechanisms of action involving the inhibition of FPPS, being very proper candidates to control and treat American trypanosomiasis. In addition, bisphosphonates have the advantage of being inexpensively synthesized and many compounds are FDA-approved drugs for the long-term treatment of several diseases.

2.1.5. Squalene synthase (SQS)

Squalene synthase (SQS) is also a key enzyme of ergosterol biosynthesis, which catalyzes the condensation of two farnesyl pyrophosphate molecules to form presqualene diphosphate and the subsequent loss of diphosphate, rearrangement, and reduction by NADPH to form squalene. Then, squalene epoxidase catalyzes the epoxidation of squalene affording oxidosqualene, which is cyclized by oxidosqualene cyclase to form lanosterol [41, 42]. Therefore,




Figure 5. Chemical structures of representative bisphosphonates targeting FPPS.

any of these enzymes constitute an excellent molecular target for antitrypanosomal agent's development. SQS is also under intense study as a possible target for cholesterol-lowering drugs in humans [41].

Quinuclidines were developed as cholesterol-decreasing agents but actually, they also turned out to be potent SQS inhibitors. However, they showed poor SQS selectivity. It has also been reported that this class of compounds eliminated the parasite both *in vitro* and *in vivo* [43, 44].

SQ109 (**21**, **Figure 6**), an ethylenediamine currently in phase II clinical trials for the treatment of tuberculosis, is of great interest for the etiological treatment of Chagas' disease [45]. Studies have conveyed that SQ109 was an inhibitor of dehydrosqualene synthase from *Staphylococcus aureus*, a protein very similar to squalene synthase, suggesting that SQ109 might also inhibit *T. cruzi* SQS [46]. In fact, recently, it was determined that SQ109 is active against all life cycle stages of *T. cruzi*, detecting the most potent activity against the highly infective trypomastigote form. Furthermore SQ109 showed synergism with posaconazole [47].

Other very interesting SQS inhibitors that have been discovered include thiocyanates like WC-9 (**22**), which proved to be a potent inhibitor of this enzyme [48]. Fluorine-containing thiocyanate derivatives exhibited higher efficacy as inhibitors of *T. cruzi* proliferation (**23,24**) (**Figure 4**) [49]. Recently, the structures of human SQS and *T. cruzi* SQS bound to a substrate-like inhibitor were reported suggesting an interesting alternative for the development of selective drugs [50].




Figure 6. Chemical structures of inhibitors of the enzymatic activity of squalene synthase.

2.2. Cruzipain

Cruzipain is the main cysteine proteinase of *T. cruzi*, which is essential for the survival of the parasite. This enzyme is involved in different cellular functions such as nutrition, penetration into the host cell, defense, and differentiation processes [51]. It has been extensively studied as a valid target for new drug development [52, 53]. There are several three-dimensional structures of cruzipain with different inhibitors allowing the identification of structural regions of this enzyme that will enable the design of new agents [54, 55].

Numerous structurally varied compounds that inhibit proliferation of *T. cruzi* by inhibiting the enzymatic activity of cruzipain have been reported. Among the compounds tested, K777 (**25**), a vinyl sulfone derivative was active against a wide range of susceptible and resistant strains (**Figure 7**). Moreover, it was able to cure *T. cruzi* acute and non-acute infection in mice, showing also synergistic activity with benznidazole [56, 57].




Figure 7. Structures of inhibitors of cruzipain activity.

Molecules containing thiosemicarbazones have been extensively explored [58–60]. Among these, 3, 4-dichlorophenyl thiosemicarbazone (**26**) is one of the most potent cruzipain inhibitor [61]. Recently, thiazolidinone derivatives (**27**) were identified as strong antiparasitic compounds (**Figure 7**) [62, 63].

Benzimidazoles have also been studied as cruzipain inhibitors and various derivatives have been synthesized and evaluated. *N*-(2-(1H-benzo[d]imidazol-2-yl) ethyl)-2-(1-bromonaphthalene-2-yloxy) acetamide (**28**) was the most potent enzyme inhibitor but it showed moderate trypanocidal activity [64, 65].

Other very potent cruzipain inhibitors have been developed such as oxyguanidine derivatives WRR-483, WRR-669, and WRR-676 (**29–31**). Some of these compounds showed suitable metabolic stability and a remarkable trypanocidal activity [66].

Very recently a series of peptidyl nitroalkenes was designed, synthesized, and evaluated as cruzipain inhibitors. Several compounds showed high activity against the enzyme observing the peptidic nature to be the determinant for their inhibitory activity [67].

Odanacatib (**32**), clofazimine (**33**), and benidipine (**34**) are examples of drug repurposing for the development of novel therapeutics for the Chagas' diseases (**Figure 5**). Odanacatib is a cathepsin K inhibitor used for the treatment of postmenopausal osteoporosis; it is a potent cysteine protease inhibitor, which is in Phase III clinical trials [68]. Clofazimine is an antibiotic applied to the treatment of leprosy [69] and benidipine is a calcium channel-blocking agent employed in the treatment of hypertension [70]. Clofazimine and benidipine were recently tested in a murine model of chronic Chagas' disease infection. Both compounds have reduced the parasitemia and the inflammatory effects have been well tolerated [71].

2.3. *Trans*-sialidase

The enzyme *trans*-sialidase (TS) belongs to the group of enzymes that are secreted by the parasite involved in the processes of cell invasion and immune evasion [72]. *T. cruzi* incorporates sialic acid (**35**) from exogenous sialoglycoconjugates by a *trans*-glycosylation reaction [73]. TS catalyze this transfer from host sialoglycoconjugates to glycoconjugates or mucins, which are attached to the cell membrane of *T. cruzi* via glycosphosphatidylinositol anchors [74, 75]. Added to its important role, the absence of this enzyme in mammalian organisms makes it an excellent molecular target.

Within TS' inhibitors it can be mentioned fluorinated compounds like 9-benzoyl-3-fluoro-*N*-acetylneuraminic acid (**36**), and 3-fluorosialyl fluoride (**37**) selectively bind to the active site of the enzyme [76]. Sulfonamide-containing hydroxylated chalcones (**38**) and quinolones (**39**) are also specific inhibitors of *T. cruzi* TS, being dihydroxylated more potent than monohydroxylated derivatives (**Figure 8**) [77].

Some approved FDA drugs were evaluated on trypomastigotes deriving out of a computational screening protocol. The anti-inflammatory sulfasalazine (**40**) showed potent antiparasitic effects in *in vivo* assays, but with moderate TS inhibition. However, this drug and sulfonamide-containing compounds could be used as leading drugs in the development of new TS inhibitors [78].




Figure 8. Representative inhibitors of *trans*-sialidase.

2.4. Trypanothione system

Trypanothione ($T[SH]_2$ or N^1, N^8 -bis-(glutathionyl)spermidine, (41)) is a dithiol, which is responsible for the thiol metabolism in trypanosomatids by maintaining the intracellular redox balance [79]. Trypanothione is responsible for the protection against oxidative stress in *T. cruzi* trapping reactive oxygen species [80].

The protective reactions involve two key enzymes, which keep the trypanothione system operating: trypanothione reductase (TryR), homologous to mammal glutathione reductase and trypanothione synthetase (TryS) [81]. TryR catalyzes the reduction of trypanothione disulfide ($T[S]_2$, 42) to $T[SH]_2$ (Figure 9); TryS catalyzes the synthesis of $T[SH]_2$ from glutathione and spermidine, keeping the amount of total trypanothione constant [82]. Taking into account the fact that trypanothione system is essential to the survival of parasites, these enzymes emerge as valid molecular targets for the search of new chemotherapeutic agents. In addition, TryS has no counterpart in mammals, making even more interesting to the design more specific and safer inhibitors [2].

Various reports have conveyed that different classes of compounds have selective enzyme inhibition activity such as tricyclic ring structures, bicyclic, and heterocyclic compounds or polyamines among others.




Figure 9. Structures of the oxidized and reduced forms of trypanothione.

Mepacrine (**43**) is a tricyclic antimalarial, which showed TryR inhibition without affecting human glutathione reductase and several mepacrine derivatives have been prepared. Although these derivatives presented greater potency, they turned out to be toxic to human cells [83].

Tricyclic phenothiazine-containing drugs are currently used as an antidepressant and have additionally exhibited antimicrobial activity [84]. Some phenothiazines demonstrated inhibitory activity against TryR [85]. Within them, thioridazine (**44**) is one of the most potent TryR inhibitors as investigations have suggested [86]. Clomipramine (**45**) [87] is another psychiatric drug with inhibiting action towards TryR. Both thioridazine and clomipramine seem to have *in vivo* effects against *T. cruzi* (Figure 10).

Library screening has allowed the identification of other classes of inhibitors that could be useful for the development of more potent TryR inhibitors. Some examples of these compounds are indatraline (**46**), a monoamine transporter inhibitor, 1-(2-(benzhydryloxy)ethyl)-4-(3-phenylpropyl)piperazine (GBR-12935, **47**) and a benzothiophene-piperidine derivative (BTCP **48**) [88]. Other or additional, studies allowed finding analogs of these compounds, which have increased the potency against TryR exhibiting higher selectivity [89]. The most interesting compounds were a piperazine-phenothiazine derivative of GBR-12935 (**49**) and a diaryl sulfide BTCP derivative [50, 88, 90].




Figure 10. Chemical structures of relevant compounds targeting trypanothione reductase.

Different compounds can behave as TryR inhibitors when turning into reactive radical species through the reduction single-electron step [83, 91]. Within these structures, 1, 4 naphthoquinones and nitrofurans have been largely studied [92, 93]. The most potent derivative of 1,4 naphthoquinone was a quinone–coumarin hybrid [51] despite showing toxicity against rat skeletal myoblasts [94].

Between nitrofurans compounds, several 5-nitrofuroic acid derivatives have been synthesized and evaluated against *T. cruzi*. The best compound was 5-nitro-furan-2-carboxylic acid dibenzyl amide [52], which it significantly increased the trypanocidal nifurtimox activity being TryTR your molecular target [95].

3. Conclusions

Currently, the chemotherapy of American trypanosomiasis remains a serious problem in the field of neglected tropical diseases. There are no vaccines, and chemotherapy is limited to old drugs, which present important drawbacks. Taking into account that this disease is associated with poor populations and bad housing conditions, pharmaceutical companies have no economic motivations. Therefore, all efforts to the development of new drugs must be made by academic and/or governmental institutions and new chemotherapies are needed urgently. In order to search new, safer and efficient drugs for the Chagas' treatment, an overview of possible molecular targets based on specific features of the biochemistry of *Trypanosoma cruzi* was given. Although there are numerous potentially valid targets, only the more representative ones were discussed here. Furthermore, some of the new potential antiparasitic drugs as well as drugs applied to other human illness were described in this work. However, despite numerous efforts and progress in the searching of new or repositioned compounds for American trypanosomiasis chemotherapy, no ideal drugs are yet available for human treatment

Author details

Guadalupe García Liñares

Address all correspondence to: linares@qo.fcen.uba.ar

Departamento de Química Orgánica y UMYMFOR, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Buenos Aires, Argentina

References

- [1] Second W. Report on Neglected Tropical Diseases. Sustaining the Drive to Overcome the Global Impact of Neglected Tropical Diseases; 2013
- [2] Linares GE, Ravaschino EL, Rodriguez JB. Progresses in the field of drug design to combat tropical protozoan parasitic diseases. *Current Medicinal Chemistry*. 2006;**13**(3):335-360

- [3] Pérez-Molina JA, Molina I. Chagas' disease. The Lancet. 2017;**391**(10115):82-94
- [4] Coura JR, Borges-Pereira J. Chronic phase of Chagas' disease: Why should it be treated? A comprehensive review. Memórias do Instituto Oswaldo Cruz. 2011;**106**(6):641-645
- [5] Bermudez J, Davies C, Simonazzi A, Real JP, Palma S. Current drug therapy and pharmaceutical challenges for Chagas' disease. Acta Tropica. 2016;**156**:1-16
- [6] Morillo CA, Marin-Neto JA, Avezum A, Sosa-Estani S, Rassi A Jr, Rosas F, et al. Randomized trial of benznidazole for chronic Chagas' cardiomyopathy. The New England Journal of Medicine. 2015;**373**(14):1295-1306
- [7] Moraes CB, Franco CH. Novel drug discovery for Chagas' disease. Expert Opinion on Drug Discovery. 2016;**11**(5):447-455
- [8] Urbina JA, McKerrow JH. Drug susceptibility of genetically engineered *Trypanosoma cruzi* strains and sterile cure in animal models as a criterion for potential clinical efficacy of anti-*T. cruzi* drugs. Antimicrobial Agents and Chemotherapy. 2015;**59**(12):7923-7924
- [9] Urbina J. New insights in Chagas' disease treatment. Drugs of the Future. 2010;**35**(5):409-419
- [10] Buckner FS, Urbina JA. Recent developments in sterol 14-demethylase inhibitors for Chagas' disease. International Journal for Parasitology: Drugs and Drug Resistance. 2012;**2**:236-242
- [11] Lepesheva GI, Villalta F, Waterman MR. Targeting *Trypanosoma cruzi* sterol 14 α -demethylase (CYP51). Advances in Parasitology. 2011;**75**:65
- [12] Lepesheva GI, Ott RD, Hargrove TY, Kleshchenko YY, Schuster I, Nes WD, et al. Sterol 14 α -demethylase as a potential target for antitrypanosomal therapy: Enzyme inhibition and parasite cell growth. Chemistry & Biology. 2007;**14**(11):1283-1293
- [13] Araújo M, Martins-Filho O, Pereira M, Brener Z. A combination of benznidazole and ketoconazole enhances efficacy of chemotherapy of experimental Chagas' disease. The Journal of Antimicrobial Chemotherapy. 2000;**45**(6):819-824
- [14] Molina J, Martins-Filho O, Brener Z, Romanha AJ, Loebenberg D, Urbina JA. Activities of the triazole derivative SCH 56592 (posaconazole) against drug-resistant strains of the protozoan parasite *Trypanosoma* (Schizotrypanum) *cruzi* in immunocompetent and immunosuppressed murine hosts. Antimicrobial Agents and Chemotherapy. 2000;**44**(1):150-155
- [15] Urbina JA, Lira R, Visbal G, Bartolí J. *In vitro* antiproliferative effects and mechanism of action of the new triazole derivative UR-9825 against the protozoan parasite *Trypanosoma* (Schizotrypanum) *cruzi*. Antimicrobial Agents and Chemotherapy. 2000;**44**(9):2498-2502
- [16] Urbina JA, Payares G, Contreras LM, Liendo A, Sanoja C, Molina J, et al. Antiproliferative effects and mechanism of action of SCH 56592 against *Trypanosoma* (Schizotrypanum) *cruzi*: *in vitro* and *in vivo* studies. Antimicrobial Agents and Chemotherapy. 1998;**42**(7):1771-1777

- [17] Urbina JA, Payares G, Molina J, Sanoja C, Liendo A, Lazardí K, et al. Cure of short- and long-term experimental Chagas' disease using D0870. *Science-New York then Washington*. 1996;**273**:969-970
- [18] Lepesheva GI, Hargrove TY, Anderson S, Kleshchenko Y, Furtak V, Wawrzak Z, et al. Structural insights into inhibition of sterol 14 α -demethylase in the human pathogen *Trypanosoma cruzi*. *The Journal of Biological Chemistry*. 2010;**285**(33):25582-25590
- [19] Pinazo M-J, Espinosa G, Gállego M, López-Chejade PL, Urbina JA, Gascón J. Successful treatment with posaconazole of a patient with chronic Chagas' disease and systemic lupus erythematosus. *The American Journal of Tropical Medicine and Hygiene*. 2010;**82**(4):583-587
- [20] Clayton J. Chagas' disease: Pushing through the pipeline. *Nature*. 2010:S12
- [21] de Figueiredo Diniz L, Caldas IS, da Matta Guedes PM, Crepalde G, de Lana M, Carneiro CM, et al. Effects of ravuconazole treatment on parasite load and immune response in dogs experimentally infected with *Trypanosoma cruzi*. *Antimicrobial Agents and Chemotherapy*. 2010;**54**(7):2979-2986
- [22] Hucke O, Gelb MH, Verlinde CL, Buckner FS. The protein farnesyltransferase inhibitor Tipifarnib as a new lead for the development of drugs against Chagas' disease. *Journal of Medicinal Chemistry*. 2005;**48**(17):5415-5418
- [23] Kraus JM, Tatipaka HB, McGuffin SA, Chennamaneni NK, Karimi M, Arif J, et al. Second generation analogues of the cancer drug clinical candidate tipifarnib for anti-Chagas' disease drug discovery. *Journal of Medicinal Chemistry*. 2010;**53**(10):3887-3898
- [24] Urbina JA, Vivas J, Lazardí K, Molina J, Payares G, Pirns MM, et al. Antiproliferative effects of $\Delta 24$ (25) sterol methyl Transferase inhibitors on *Trypanosoma* (Schizotrypanum) *cruzi*: *In vitro* and *in vivo* studies. *Chemotherapy*. 1996;**42**(4):294-307
- [25] Vivas J, Urbina JA, de Souza W. Ultrastructural alterations in *Trypanosoma* (Schizotrypanum) *cruzi* induced by $\Delta 24$ (25) sterol methyl transferase inhibitors and their combinations with ketoconazole. *International Journal of Antimicrobial Agents*. 1996;**7**(4):235-240
- [26] Gros L, Castillo-Acosta VM, Jiménez CJ, Sealey-Cardona M, Vargas S, Estévez AM, et al. New azasterols against *Trypanosoma brucei*: Role of 24-sterol methyltransferase in inhibitor action. *Antimicrobial Agents and Chemotherapy*. 2006;**50**(8):2595-2601
- [27] Gigante F, Kaiser M, Brun R, Gilbert IH. Design and preparation of sterol mimetics as potential antiparasitics. *Bioorganic & Medicinal Chemistry*. 2010;**18**(20):7291-7301
- [28] Quist-Paulsen P. Statins and inflammation: An update. *Current Opinion in Cardiology*. 2010;**25**(4):399-405
- [29] Silva RR, Shrestha-Bajracharya D, Almeida-Leite CM, Leite R, Bahia MT, Talvani A. Short-term therapy with simvastatin reduces inflammatory mediators and heart inflammation during the acute phase of experimental Chagas' disease. *Memórias do Instituto Oswaldo Cruz*. 2012;**107**(4):513-521

- [30] Urbina J, Lazardi K, Marchan E, Visbal G, Aguirre T, Piras M, et al. Mevinolin (lovastatin) potentiates the antiproliferative effects of ketoconazole and terbinafine against *Trypanosoma* (Schizotrypanum) *cruzi*: *in vitro* and *in vivo* studies. Antimicrobial Agents and Chemotherapy. 1993;**37**(3):580-591
- [31] Lombard J, Moreira D. Origins and early evolution of the mevalonate pathway of isoprenoid biosynthesis in the three domains of life. Molecular Biology and Evolution. 2010;**28**(1):87-99
- [32] Martin MB, Grimley JS, Lewis JC, Heath HT, Bailey BN, Kendrick H, et al. Bisphosphonates inhibit the growth of *Trypanosoma brucei*, *Trypanosoma cruzi*, *Leishmania donovani*, *Toxoplasma gondii*, and *Plasmodium falciparum*: A potential route to chemotherapy. Journal of Medicinal Chemistry. 2001;**44**(6):909-916
- [33] Reddy R, Dietrich E, Lafontaine Y, Houghton TJ, Belanger O, Dubois A, et al. Bisphosphonated benzoxazinorifamycin prodrugs for the prevention and treatment of osteomyelitis. ChemMedChem. 2008;**3**(12):1863-1868
- [34] Miller K, Erez R, Segal E, Shabat D, Satchi-Fainaro R. Targeting bone metastases with a bispecific anticancer and antiangiogenic polymer-alendronate-taxane conjugate. Angewandte Chemie, International Edition. 2009;**48**(16):2949-2954
- [35] Reszka AA, Rodan GA. Nitrogen-containing bisphosphonate mechanism of action. Mini Reviews in Medicinal Chemistry. 2004;**4**(7):711-719
- [36] Urbina JA, Moreno B, Vierkoter S, Oldfield E, Payares G, Sanoja C, et al. *Trypanosoma cruzi* contains major pyrophosphate stores, and its growth *in vitro* and *in vivo* is blocked by pyrophosphate analogs. The Journal of Biological Chemistry. 1999;**274**(47):33609-33615
- [37] Ghosh S, Chan JM, Lea CR, Meints GA, Lewis JC, Tovian ZS, et al. Effects of bisphosphonates on the growth of *Entamoeba histolytica* and *Plasmodium* species *in vitro* and *in vivo*. Journal of Medicinal Chemistry. 2004;**47**(1):175-187
- [38] Szajnman SH, Garcia GE, Li Z-H, Jiang C, Galizzi M, Bontempi EJ, et al. Synthesis and biological evaluation of 2-alkylaminoethyl-1, 1-bisphosphonic acids against *Trypanosoma cruzi* and *Toxoplasma gondii* targeting farnesyl diphosphate synthase. Bioorganic & Medicinal Chemistry. 2008;**16**(6):3283-3290
- [39] Recher M, Barboza AP, Li Z-H, Galizzi M, Ferrer-Casal M, Szajnman SH, et al. Design, synthesis and biological evaluation of sulfur-containing 1, 1-bisphosphonic acids as antiparasitic agents. European Journal of Medicinal Chemistry. 2013;**60**:431-440
- [40] Zhang Y, Cao R, Yin F, Hudock MP, Guo R-T, Krysiak K, et al. Lipophilic bisphosphonates as dual farnesyl/geranylgeranyl diphosphate synthase inhibitors: An X-ray and NMR investigation. Journal of the American Chemical Society. 2009;**131**(14):5153-5162
- [41] Urbina JA, Concepcion JL, Rangel S, Visbal G, Lira R. Squalene synthase as a chemotherapeutic target in *Trypanosoma cruzi* and *Leishmania mexicana*. Molecular and Biochemical Parasitology. 2002;**125**(1):35-45

- [42] Oldfield E. Targeting isoprenoid biosynthesis for drug discovery: Bench to bedside. *Accounts of Chemical Research*. 2010;**43**(9):1216-1226
- [43] Urbina JA, Concepcion JL, Caldera A, Payares G, Sanoja C, Otomo T, et al. *In vitro* and *in vivo* activities of E5700 and ER-119884, two novel orally active squalene synthase inhibitors, against *Trypanosoma cruzi*. *Antimicrobial Agents and Chemotherapy*. 2004;**48**(7):2379-2387
- [44] Sealey-Cardona M, Cammerer S, Jones S, Ruiz-Pérez LM, Brun R, Gilbert IH, et al. Kinetic characterization of squalene synthase from *Trypanosoma cruzi*: Selective inhibition by quinuclidine derivatives. *Antimicrobial Agents and Chemotherapy*. 2007;**51**(6):2123-2129
- [45] Sacksteder KA, Protopopova M, Barry CE, Andries K, Nacy CA. Discovery and development of SQ109: A new antitubercular drug with a novel mechanism of action. *Future Microbiology*. 2012;**7**(7):823-837
- [46] Lin F-Y, Liu Y-L, Li K, Cao R, Zhu W, Axelson J, et al. Head-to-head prenyl transferases: Anti-infective drug targets. *Journal of Medicinal Chemistry*. 2012;**55**(9):4367-4372
- [47] Veiga-Santos P, Li K, Lameira L, de Carvalho TMU, Huang G, Galizzi M, et al. SQ109, a new drug lead for Chagas' disease. *Antimicrobial Agents and Chemotherapy*. 2015;**59**(4):1950-1961
- [48] Urbina JA, Concepcion JL, Montalvetti A, Rodriguez JB, Docampo R. Mechanism of action of 4-phenoxyphenoxyethyl thiocyanate (WC-9) against *Trypanosoma cruzi*, the causative agent of Chagas' disease. *Antimicrobial Agents and Chemotherapy*. 2003;**47**(6):2047-2050
- [49] Liñares GG, Gismondi S, Codesido NO, Moreno SN, Docampo R, Rodriguez JB. Fluorine-containing aryloxyethyl thiocyanate derivatives are potent inhibitors of *Trypanosoma cruzi* and *Toxoplasma gondii* proliferation. *Bioorganic & Medicinal Chemistry Letters*. 2007;**17**(18):5068-5071
- [50] Shang N, Li Q, Ko T-P, Chan H-C, Li J, Zheng Y, et al. Squalene synthase as a target for Chagas' disease therapeutics. *PLoS Pathogens*. 2014;**10**(5):e1004114
- [51] Sajid M, McKerrow JH. Cysteine proteases of parasitic organisms. *Molecular and Biochemical Parasitology*. 2002;**120**(1):1-21
- [52] G Duschak V. A decade of targets and patented drugs for chemotherapy of Chagas' disease. *Recent Patents on Anti-Infective Drug Discovery*. 2011;**6**(3):216-259
- [53] Martinez-Mayorga K, Byler KG, Ramirez-Hernandez AI, Terrazas-Alvares DE. Cruzain inhibitors: Efforts made, current leads and a structural outlook of new hits. *Drug Discovery Today*. 2015;**20**(7):890-898
- [54] Brak K, Kerr ID, Barrett KT, Fuchi N, Debnath M, Ang K, et al. Nonpeptidic tetrafluorophenoxymethyl ketone cruzain inhibitors as promising new leads for Chagas' disease chemotherapy. *Journal of Medicinal Chemistry*. 2010;**53**(4):1763-1773

- [55] Chen YT, Brinen LS, Kerr ID, Hansell E, Doyle PS, McKerrow JH, et al. *In vitro* and *in vivo* studies of the trypanocidal properties of WRR-483 against *Trypanosoma cruzi*. PLoS Neglected Tropical Diseases. 2010;4(9):e825
- [56] McKerrow JH, Doyle P, Engel J, Podust L, Robertson S, Ferreira R, et al. Two approaches to discovering and developing new drugs for Chagas' disease. Memórias do Instituto Oswaldo Cruz. 2009;104:263-269
- [57] Zingales B, Miles MA, Moraes CB, Luquetti A, Guhl F, Schijman AG, et al. Drug discovery for Chagas' disease should consider *Trypanosoma cruzi* strain diversity. Memórias do Instituto Oswaldo Cruz. 2014;109(6):828-833
- [58] Espíndola JWP, de Oliveira Cardoso MV, de Oliveira Filho GB, e Silva DAO, Moreira DRM, Bastos TM, et al. Synthesis and structure–activity relationship study of a new series of antiparasitic aryloxyl thiosemicarbazones inhibiting *Trypanosoma cruzi* cruzain. European Journal of Medicinal Chemistry. 2015;101:818-835
- [59] Fonseca NC, da Cruz LF, da Silva Villela F, do Nascimento Pereira GA, de Siqueira-Neto JL, Kellar D, et al. Synthesis of a sugar-based thiosemicarbazone series and structure-activity relationship versus the parasite cysteine proteases rhodesain, cruzain, and schistosoma mansonii cathepsin B1. Antimicrobial Agents and Chemotherapy. 2015; 59(5):2666-2677
- [60] Moreira DRM, Lima Leite AC, Cardoso MVO, Srivastava RM, Hernandez MZ, Rabello MM, et al. Structural design, synthesis and structure–activity relationships of Thiazolidinones with enhanced anti-*Trypanosoma cruzi* activity. ChemMedChem. 2014; 9(1):177-188
- [61] Carvalho L, Gomes M, Rocha B, de Oliveira R, Maria E, de Melo E. Anti-parasite effects of new thiosemicarbazones and their products thiazolidinone including cellular aspects of intracellular elimination of *Trypanosoma cruzi* *in vitro*. Journal of Developing Drugs 2014;3(126):10.4172
- [62] Hernandez MZ, Rabello MM, Leite ACL, Cardoso MVO, Moreira DRM, Brondani DJ, et al. Studies toward the structural optimization of novel thiazolylhydrazone-based potent antitrypanosomal agents. Bioorganic & Medicinal Chemistry. 2010;18(22):7826-7835
- [63] de Oliveira Filho GB, de Oliveira Cardoso MV, Espíndola JWP, Ferreira LFGR, de Simone CA, Ferreira RS, et al. Structural design, synthesis and pharmacological evaluation of 4-thiazolidinones against *Trypanosoma cruzi*. Bioorganic & Medicinal Chemistry. 2015;23(23):7478-7486
- [64] Ferreira RS, Dessoy MA, Pauli I, Souza ML, Krogh R, Sales AI, et al. Synthesis, biological evaluation, and structure–activity relationships of potent noncovalent and Nonpeptidic Cruzain inhibitors as anti-*Trypanosoma cruzi* agents. Journal of Medicinal Chemistry. 2014;57(6):2380-2392
- [65] Ferreira RS, Simeonov A, Jadhav A, Eidam O, Mott BT, Keiser MJ, et al. Complementarity between a docking and a high-throughput screen in discovering new cruzain inhibitors. Journal of Medicinal Chemistry. 2010;53(13):4891-4905

- [66] Jones BD, Tochowicz A, Tang Y, Cameron MD, McCall L-I, Hirata K, et al. Synthesis and evaluation of oxyguanidine analogues of the cysteine protease inhibitor WRR-483 against Cruzain. *ACS Medicinal Chemistry Letters*. 2015;7(1):77-82
- [67] Latorre A, Schirmeister T, Kesselring J, Jung S, Johé P, Hellmich UA, et al. Dipeptidyl nitroalkenes as potent reversible inhibitors of cysteine proteases Rhodesain and Cruzain. *ACS Medicinal Chemistry Letters*. 2016;7(12):1073-1076
- [68] Ndao M, Beaulieu C, Black WC, Isabel E, Vasquez-Camargo F, Nath-Chowdhury M, et al. Reversible cysteine protease inhibitors show promise for a Chagas' disease cure. *Antimicrobial Agents and Chemotherapy*. 2014;58(2):1167-1178
- [69] Cholo MC, Steel HC, Fourie PB, Germishuizen WA, Anderson R. Clofazimine: Current status and future prospects. *The Journal of Antimicrobial Chemotherapy*. 2011;67(2):290-298
- [70] Yao K, Nagashima K, Miki H. Pharmacological, pharmacokinetic, and clinical properties of benidipine hydrochloride, a novel, long-acting calcium channel blocker. *Journal of Pharmacological Sciences*. 2006;100(4):243-261
- [71] Sbaraglini ML, Bellera CL, Fraccaroli L, Larocca L, Carrillo C, Talevi A, et al. Novel cruzipain inhibitors for the chemotherapy of chronic Chagas' disease. *International Journal of Antimicrobial Agents*. 2016;48(1):91-95
- [72] Buschiazzo A, Amaya MF, Cremona ML, Frasch AC, Alzari PM. The crystal structure and mode of action of trans-sialidase, a key enzyme in *Trypanosoma cruzi* pathogenesis. *Molecular Cell*. 2002;10(4):757-768
- [73] Previato J, Andrade A, Vermelho A, Firmino J, Mendonça-Previato L. Evidence for N-glycolylneuraminic acid incorporation by *Trypanosoma cruzi* from infected animal. *Memórias do Instituto Oswaldo Cruz*. 1990;85:38-39
- [74] Cross GA, Takle GB. The surface trans-sialidase family of *Trypanosoma cruzi*. *Annual Reviews in Microbiology*. 1993;47(1):385-411
- [75] Ferguson M, Brimacombe J, Cottaz S, Field R, Güther L, Homans S, et al. Glycosylphosphatidylinositol molecules of the parasite and the host. *Parasitology*. 1994;108(S1):S45-S54
- [76] Buchini S, Buschiazzo A, Withers SG. A new generation of specific *Trypanosoma cruzi* trans-sialidase inhibitors. *Angewandte Chemie, International Edition*. 2008;47(14):2700-2703
- [77] Kim JH, Ryu HW, Shim JH, Park KH, Withers SG. Development of new and selective *Trypanosoma cruzi* trans-Sialidase inhibitors from sulfonamide Chalcones and their derivatives. *Chembiochem*. 2009;10(15):2475-2479
- [78] Lara-Ramirez EE, López-Cedillo JC, Noguera-Torres B, Kashif M, Garcia-Perez C, Bocanegra-Garcia V, et al. An *in vitro* and *in vivo* evaluation of new potential trans-sialidase inhibitors of *Trypanosoma cruzi* predicted by a computational drug repositioning method. *European Journal of Medicinal Chemistry*. 2017;132:249-261

- [79] Fairlamb AH, Cerami A. Identification of a novel, thiol-containing co-factor essential for glutathione reductase enzyme activity in trypanosomatids. *Molecular and Biochemical Parasitology*. 1985;**14**(2):187-198
- [80] Brigelius-Flohé R, Flohé L. Basic principles and emerging concepts in the redox control of transcription factors. *Antioxidants & Redox Signaling*. 2011;**15**(8):2335-2381
- [81] Flohé L. The trypanothione system and its implications in the therapy of trypanosomatid diseases. *International Journal of Medical Microbiology*. 2012;**302**(4):216-220
- [82] Oza SL, Tetaud E, Ariyanayagam MR, Warnon SS, Fairlamb AH. A single enzyme catalyses formation of trypanothione from glutathione and spermidine in *Trypanosoma cruzi*. *The Journal of Biological Chemistry*. 2002;**277**(39):35853-35861
- [83] Krauth-Siegel RL, Bauer H, Schirmer RH. Dithiol proteins as guardians of the intracellular redox milieu in parasites: Old and new drug targets in trypanosomes and malaria-causing plasmodia. *Angewandte Chemie, International Edition*. 2005;**44**(5):690-715
- [84] Amaral L, Viveiros M, Kristiansen JE. "non-antibiotics": Alternative therapy for the management of MDRTB and MRSA in economically disadvantaged countries. *Current Drug Targets*. 2006;**7**(7):887-891
- [85] Bernardes L, Zani C, Carvalho I. Trypanosomatidae diseases: From the current therapy to the efficacious role of trypanothione reductase in drug discovery. *Current Medicinal Chemistry*. 2013;**20**(21):2673-2696
- [86] Presti MSL, Bazán PC, Strauss M, Báez AL, Rivarola HW, Paglini-Oliva PA. Trypanothione reductase inhibitors: Overview of the action of thioridazine in different stages of Chagas' disease. *Acta Tropica*. 2015;**145**:79-87
- [87] Fauro R, Presti SL, Bazan C, Baez A, Strauss M, Triquell F, et al. Use of clomipramine as chemotherapy of the chronic phase of Chagas' disease. *Parasitology*. 2013;**140**(7):917-927
- [88] Richardson JL, Nett IR, Jones DC, Abdille MH, Gilbert IH, Fairlamb AH. Improved tricyclic inhibitors of trypanothione reductase by screening and chemical synthesis. *ChemMedChem*. 2009;**4**(8):1333-1340
- [89] Walton JG, Jones DC, Kiuru P, Durie AJ, Westwood NJ, Fairlamb AH. Synthesis and evaluation of Indatraline-based inhibitors for Trypanothione Reductase. *ChemMedChem*. 2011;**6**(2):321-328
- [90] Eberle C, Lauber BS, Fankhauser D, Kaiser M, Brun R, Krauth-Siegel RL, et al. Improved inhibitors of trypanothione reductase by combination of motifs: Synthesis, inhibitory potency, binding mode, and antiprotozoal activities. *ChemMedChem*. 2011;**6**(2):292-301
- [91] Gendron T, Lanfranchi DA, Davioud-Charvet E. Redox-active agents in reactions involving the Trypanothione/Trypanothione Reductase-based system to fight Kinetoplastid parasites. *Trypanosomatid Diseases: Molecular Routes to Drug Discovery*. 2013:405-428
- [92] Leroux AE, Krauth-Siegel RL. Thiol redox biology of trypanosomatids and potential targets for chemotherapy. *Molecular and Biochemical Parasitology*. 2016;**206**(1):67-74

- [93] R Wilkinson S, Bot C, M Kelly J, S Hall B. Trypanocidal activity of nitroaromatic prodrugs: Current treatments and future perspectives. *Current Topics in Medicinal Chemistry*. 2011;**11**(16):2072-2084
- [94] Belluti F, Uliassi E, Veronesi G, Bergamini C, Kaiser M, Brun R, et al. Toward the development of dual-targeted Glyceraldehyde-3-phosphate dehydrogenase/Trypanothione Reductase inhibitors against *Trypanosoma brucei* and *Trypanosoma cruzi*. *ChemMedChem*. 2014;**9**(2):371-382
- [95] Arias D, Herrera F, Garay A, Rodrigues D, Forastieri P, Luna L, et al. Rational design of nitrofurantoin derivatives: Synthesis and valuation as inhibitors of *Trypanosoma cruzi* trypanothione reductase. *European Journal of Medicinal Chemistry*. 2017;**125**:1088-1097