

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Skeletal Muscle as a Therapeutic Target for Natural Products to Reverse Metabolic Syndrome

Sithandiwe Eunice Mazibuko-Mbeje,
Phiwayinkosi V. Dludla, Bongani B. Nkambule,
Nnini Obonye and Johan Louw

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/intechopen.78687>

Abstract

Natural compounds, especially polyphenols have become a popular area of research mainly due to their apparent health benefits. Increasing the phenolic content of a diet, apart from its antioxidant benefit, has a beneficial effect on signaling molecules involved in carbohydrate and lipid metabolism. These effects could potentially protect against metabolic syndrome, a cluster of metabolic complications such as obesity, insulin resistance and type 2 diabetes that is characterized by a dysregulated carbohydrate, and lipid metabolism. Research continues to investigate various natural compounds for their amelioration of impaired signaling mechanisms that may lead to dysregulated metabolism to find means to improve the life expectancy of patients with metabolic syndrome. In this chapter, a systematic search through major databases such as MEDLINE/PubMed, EMBASE, and Google Scholar of literature reporting on the ameliorative potential of commonly investigated natural products that target skeletal muscle to ameliorate metabolic syndrome associated complications was conducted. The selected natural products that are discussed include apigenin, aspalathin, berberine, curcumin, epigallocatechin gallate, hesperidin, luteolin, naringenin, quercetin, resveratrol, rutin, and sulforaphane.

Keywords: skeletal muscle, metabolic syndrome, insulin resistance, type 2 diabetes, natural products

1. Introduction

A considerable amount of interest has been placed on the discovery of novel naturally occurring plant-derived compounds for the treatment and prevention of various diseases. Bioactive

compounds of plant origin have long been shown to possess strong ameliorative properties against various communicable and noncommunicable diseases [1, 2]. For example, since its traditional use during the 1950s, artemisinin, an antimalarial *qinghao* derived lactone, has been the leading therapy for the treatment of *Plasmodium falciparum* malaria worldwide [3]. Similarly, the traditional use of galegine, an alkaloid isolated from *Galega officinalis*, led to the discovery of biguanide class of antidiabetic medications such as metformin [4]. Agents such as metformin are effective at lowering blood glucose levels and combating complications associated with insulin resistance (IR), the major characteristic of the metabolic syndrome [5]. However, the continued rise in the mortality of diabetic patients warrants an investigation into alternative therapies to reduce the burden of noncommunicable diseases. Naturally derived compounds such as polyphenols are increasingly explored for their therapeutic potential to reverse IR and thus decrease the risk of developing the metabolic syndrome. This may eventually lead to an increased life expectancy of diabetic individuals [6]. Thus, due to its modulatory effect of glucose and lipid metabolism, skeletal muscle has been a target to a growing number of therapeutic interventions in an effort to reverse IR and improve the management of metabolic syndrome [7, 8]. Here, we systematically assessed the available literature on the ameliorative potential of some of the prominent natural products against IR associated complications. A systematic search was conducted on all major databases such as MEDLINE/PubMed, EMBASE, and Google Scholar, for available literature reporting on the ameliorative properties of some of the prominent natural compounds including apigenin, aspalathin, berberine, curcumin, epigallocatechin gallate, hesperidin, luteolin, naringenin, quercetin, resveratrol, rutin, and sulforaphane against IR related to the development of metabolic syndrome. The search was conducted from inception until the end of January 2018, gray literature such as abstract proceedings and pre-prints were also included. There were no language restrictions implemented while review articles were screened for primary findings.

2. Apigenin

Apigenin (PubChem CID: 5280443) is a natural flavone (4',5,7-trihydroxyflavone) with the molecular formula $C_{15}H_{10}O_5$ (MW 270.24 g/mol) that is abundantly present in fruits and vegetables, including parsley, chamomile, and celery (**Figure 1**) [9]. Apigenin was identified as the main yellow dye compound in the flowers of *Delphinium Zalil* as early as the 1890s [10], and its bioavailability and metabolism profile has been studied as far back as the 1970s [11].




Figure 1. The chemical structure of apigenin (4',5,7-trihydroxyflavone).

Although pharmacokinetic studies show that apigenin has low bioavailability [12, 13], this compound has been detected in rat plasma after intravenous bolus administration [14], and it was demonstrated that human intestinal microbiota might contribute to its metabolism [15]. The known metabolites of apigenin detected in the urine of rats consist of p-hydroxyphenylpropionic acid, p-hydroxycinnamic acid, and p-hydroxybenzoic acid metabolites [11] while it is known glucosides include apiin, apigenin, vitexin, isovitexin, and rhoifolin.

In relation to its biological activities, increasing studies have demonstrated that apigenin displays a broad spectrum of anticarcinogenic properties as reviewed by Sung et al. [16]. Some of the well-studied mechanisms associated with the chemo-preventative capabilities of apigenin include its anti-inflammatory activity, its ability to suppress cell proliferation and oxidative stress, as well as its modulatory effect of autophagy and apoptosis [16, 17]. Interestingly, similar mechanisms have also been implicated in the development and aggravation of IR and its related complications. In a recent study, Jung et al. [18] showed that in addition to reducing circulating free fatty acids (FFAs), total cholesterol, and apolipoprotein B levels, apigenin modulated transcriptional factors linked with the development of obesity and related metabolic disturbances in high fat diet (HFD)-induced mice. This study showed that apigenin upregulated the expression of genes responsible for the regulation of beta-oxidation, oxidative phosphorylation, as well as electron transport chain and cholesterol homeostasis, which are all essential target sites for the control of substrate usage in cells. Although limited studies are reporting on its effect on skeletal muscle, two recent studies have shown that apigenin can regulate skeletal muscle function. For instance, Choi et al. [19] showed that this flavone improved mitochondrial function and exercise capacity by reducing the expression of atrophic genes such as RING-finger protein-1 and Atrogin 1 in mice fed HFD. Jang et al. [20] demonstrated that in C2C12 cells and skeletal muscle of C57BL/6 mice, this flavone promoted hypertrophy and myogenic differentiation by regulating protein arginine methyltransferase 7 (Prmt7)-peroxisome proliferator-activated receptor gamma coactivator 1-alpha (PGC-1α)-G protein-coupled receptor 56 (GPR56) pathway, as well as the Prmt7-p38-myoD pathway. Although additional studies are required to further assess the impact of apigenin in the modulation of metabolic disease-related complications through the regulation of skeletal muscle function, the two aforementioned studies suggest that this flavone has a potential to protect against skeletal muscle weakness associated with metabolic complications.

3. Aspalathin

Aspalathin (PubChem CID: 11282394) is a natural C-glucosyl dihydrochalcone (3'-β-D-glucopyranosyl-2',3,4,4',6'- pentahydroxydihydrochalcone) with the molecular formula C₂₁H₂₄O₁₁ (MW 452.412 g/mol) (**Figure 2**) [21]. Although aspalathin was known to be uniquely found in rooibos [22], recent evidence has shown that this C-linked dihydrochalcone glucoside can be detected in trace amounts in two other species of *Acacia pendula* [23]. Aspalathin is considered to have a poor bioavailability profile in different experimental settings as reviewed by Muller et al. [24] and Johnson et al. [25]. While Stalmach et al. [26], using high-performance liquid chromatography-mass spectrometry method, showed that O-methyl-aspalathin-O-glucuronide




Figure 2. The chemical structure of aspalathin (3'-β-D-glucopyranosyl-2',3,4,4',6'-pentahydroxydihydrochalcone).

and eriodictyol-*O*-sulfate were the main metabolites excreted following ingestion of rooibos extract containing 10-fold higher levels of aspalathin in human subjects. In addition, a recent study by Bowles et al. [27] showed that aspalathin can be absorbed and metabolized to mostly sulfate conjugates detected in the urine of mice. However, additional evidence is required to establish the pharmacokinetic profile of aspalathin.

Relevant to its biological activity, the initial evidence demonstrated that aspalathin possess strong antioxidant properties by scavenging 2,2-diphenyl-β-picrylhydrazyl (DPPH) radical in vitro [28]. This effect was important since experimental and clinical studies support the notion that drug compounds that enhance intracellular antioxidant properties can further exhibit a wide range of beneficial effects against the development of metabolic syndrome [29]. In addition to its robust antioxidant activity [28, 30–34], aspalathin can ameliorate inflammation [35–39], protect cardiac cells exposed to high glucose concentrations [40–44], and also display glucose lowering properties [45–50]. In addition to work by our group [46, 48], studies conducted by Kawano et al. [51] and Son et al. [50] have reported on the effect of pure aspalathin or an aspalathin rich green rooibos extract on the signaling mechanisms that regulate glucose and lipid metabolism in skeletal muscle. Activation of 5' AMP-activated protein kinase (AMPK), an important kinase in the regulation of energy production, as well as increasing the expression and translocation of glucose transporter (GLUT) 4 have been the key molecular targets by aspalathin in the skeletal muscle. Thus, although additional evidence such as assessing the therapeutic effect of this dihydrochalcone on skeletal muscle biopsies of insulin-resistant human subjects is still necessary, its aforementioned potential to target AMPK, and improve glucose uptake is of major importance for future therapeutic development.

4. Berberine

Berberine (PubChem CID: 2353) is a quaternary alkaloid (5,6-Dihydro-9,10-dimethoxybenzo[g]-1,3-benzodioxolo[5,6-a]quinolizinium) with the molecular formula $C_{20}H_{18}NO_4^+$ (MW 336.37 g/mol) that is present in several plants including *Hydrastis canadensis*, *Xanthorrhiza simplicissima*, *Phellodendron amurense*, and *Berberis aristata* (**Figure 3**) [52]. Berberrubine, thalifendine, demethyleberberine, and jatrorrhizine are some of the major metabolites detected in plasma following the administration of berberine in rats, with the liver and intestinal bacteria identified to participate in the metabolism, and disposition of this compound in vivo [53]. Although a number of factors, including it being hydrophilic in nature and its containment of quaternary ammonium groups contribute to the low bioavailability of berberine [54]. Interestingly, the absorption of berberine in the small intestine can be enhanced by d-α-tocopheryl polyethylene glycol 1000 succinate [55].




Figure 3. The chemical structure of berberine (5,6-dihydro-9,10-dimethoxybenzo[g]-1,3-benzodioxolo[5,6-a]quinolizinium).

Therefore, further research is required to better understand and inform on mechanisms that can add to our current knowledge on the bioavailability of berberine, which is crucial in improving its efficacy *in vivo*.

Berberine has a long history of medicinal use in traditional Chinese and Native American medicine [56] and has demonstrated a number of beneficial effects against metabolic complications, including amelioration of IR. Berberine demonstrated an enhanced effect to reduce body weight and raise plasma triglyceride levels while improving glucose tolerance and insulin action in both type 2 diabetic (*db/db*) mice and in FHD fed rats [57]. Interestingly, similar to aspalathin, an increase of glucose uptake through activation of AMPK as well as enhanced translocation of GLUT4 in skeletal muscle remains important in the ameliorative potential of berberine against IR [58–62]. However, it has been reported that berberine can alter muscle metabolism by altering mitochondrial function, resulting in the development of muscle atrophy in normal, and diabetic (*db/db*) mice [63]. Although the results were not in human subjects, these findings remain relevant since loss of muscle mass is an important feature that occurs in type 2 diabetic patients, especially in older individuals [64]. These results suggest that precaution should be taken when using these quaternary alkaloids, especially considering the toxicity of high doses [65]. In addition to acting by targeting the mitochondria [65], another mechanism by which berberine can reverse IR include downregulating toll-like receptor 4 (TLR4)/inhibitor of nuclear factor kappa-B kinase subunit beta (IKK β)/nuclear factor kappa-light-chain-enhancer of activated B cells (NF- κ B) inflammation signaling pathway, leading to reduced inflammation [66].

5. Curcumin

Curcumin, also known as diferuloylmethane (PubChem CID: 969516; (1E,6E)-1,7-Bis (4-hydroxy-3-methoxyphenyl) hepta-1,6-diene-3,5-dione), is a major polyphenolic derivative of turmeric (*Curcuma longa*) with the molecular formula C₂₁H₂₀O₆ (MW 368.39 g/mol) (**Figure 4**) [67]. A single oral dose administration of curcumin can lead to the detection of its metabolites, glucuronide, and sulfate conjugates in plasma of human subjects [68]. Although is considered to have a safety profile, curcumin displays poor bioavailability profile that is coupled with quick metabolism and systemic removal [69]. However, recent developments such as blocking of metabolic pathways by




Figure 4. The chemical structure of curcumin ((1E,6E)-1,7-bis(4-hydroxy-3-methoxyphenyl)hepta-1,6-diene-3,5-dione).

concomitant administration with other agents, conjugation, and modification of structure, as well as modulation of route and medium of administration are some of the explored approaches to improve the bioavailability of curcumin as reviewed by Prasad et al. [70]. Indeed, increasing research over the past 30 years has focused on exploring the pharmacokinetics, safety profile, and efficacy of this natural product in order to enhance its therapeutic profile in humans [71].

An increasing number of reviews has been published to keep track of the cumulative literature informing on the therapeutic potential of curcumin, including anticancer, antioxidant, anti-inflammatory, and antibacterial activities [70–72]. Relevant to its effect on skeletal muscle function. A study published in 2005 by Farid et al. [73] showed that curcumin failed to inhibit NF- κ B activity, leading to its inability to ameliorate loss of muscle mass in the soleus. However, in a follow-up study published in 2008, curcumin presented enhanced effect in blocking sepsis-induced muscle proteolysis, at least in part by inhibiting NF- κ B, and p38 activities in rats [74]. In L6 or C2C12 myotubes exposed to high palmitate concentrations as a model of IR, curcumin reversed IR by increasing glucose and FFA oxidation, at least in part by mediating LKB1-AMPK pathways, as well as suppressing insulin receptor substrate 1 (IRS-1) Ser³⁰⁷ and protein kinase B (AKT) phosphorylation [75–77]. Although similar evidence has been supported by in vivo experiments on skeletal muscle tissue of either diabetic or nondiabetic rodents [75, 77], curcumin displays an enhanced capacity to protect against oxidative stress associated complications by improving mitochondrial biogenesis, and other antioxidant mechanisms [78–81]. This involves activation of the nuclear factor (erythroid-derived 2)-like 2 (NRF2) [82], an essential intracellular antioxidant response element that is a target of various natural products aiming to reduce metabolic disease-associated complications.

6. Epigallocatechin gallate

Epigallocatechin gallate (PubChem CID: 65064) is an ester of epigallocatechin and gallic acid ([(2R,3R)-5,7-dihydroxy-2-(3,4,5-trihydroxyphenyl)chroman-3-yl] 3,4,5-trihydroxybenzoate, with the molecular formula C₂₂H₁₈O₁₁ (MW 458.375 g/mol), that is abundantly found in tea (Figure 5) [83]. Due to the popularity of green tea and as one of its major components, epigallocatechin gallate remains one of the highly consumed polyphenolic compounds [84]. Although it is detectable in its original form in human plasma after oral administration [85], epigallocatechin gallate is considered to have very low oral bioavailability profile as reviewed by Mereles and colleagues [86]. Although additional evidence is required to improve its




Figure 5. The chemical structure of epigallocatechin gallate ([[(2R,3R)-5,7-dihydroxy-2-(3,4,5-trihydroxyphenyl)chroman-3-yl] 3,4,5-trihydroxybenzoate).

bioavailability, there has been an extensive exploration of this polyphenolic compound for its chemopreventive properties. Among the 10 polyphenols present in green tea, epigallocatechin gallate was found to exhibit the most antiproliferative and antiapoptotic effects [87].

It has already been established that epigallocatechin gallate can ameliorate complications linked with the development of the metabolic syndrome, by improving insulin sensitivity in both obese rodents and patients [88–90]. The enhanced therapeutic effect of this catechin has been associated with the modulation of various signaling pathways, including targeting of genes involved in cell survival, FFA regulation, mitochondrial energetics, intracellular antioxidant response, and others as reviewed by Singh and colleagues [91]. A number of studies have demonstrated several mechanisms associated with the ameliorative effect of epigallocatechin gallate on IR and associated complications in skeletal muscle. In addition to strengthening muscle integrity [92–94], accumulative data has been presented that this catechin can improve insulin sensitivity by enhancing glucose uptake, reduce lactate concentrations, enhancing mitochondrial capacity and stimulating beta-oxidation in cultured cells, or rodents as well as obese human subjects [95–100]. Inhibition of oxidative stress, activation of AMPK, increased expression of PGC-1 α , NAD-dependent protein deacetylase sirtuin-1 (SIRT1), nuclear respiratory factor 1, medium chain acyl coA decarboxylase, uncoupling protein 3 (UCP3), AKT, and peroxisome proliferator-activated receptor alpha (PPAR α) are some of the mechanisms targeted by epigallocatechin to enhance skeletal muscle function in a diseased state [101–104].

7. Hesperidin

Hesperidin (PubChem CID: 10621) is a flavanone glycoside ((2S)-5-hydroxy-2-(3-hydroxy-4-methoxyphenyl)-7-[(2S,3R,4S,5S,6R)-3,4,5-trihydroxy-6-[[[(2R,3R,4R,5R,6S)-3,4,5-trihydroxy-6-methyloxan-2-yl]oxymethyl]oxan-2-yl]oxy-2,3-dihydrochromen-4-one) with the molecular formula C₂₈H₃₄O₁₅ (MW 610.565 g/mol) that is present in high amounts in citrus fruits (**Figure 6**)




Figure 6. The chemical structure of hesperidin ((2S)-5-hydroxy-2-(3-hydroxy-4-methoxyphenyl)-7-[(2S,3R,4S,5S,6R)-3,4,5-trihydroxy-6-[[[(2R,3R,4R,5R,6S)-3,4,5-trihydroxy-6-methyloxan-2-yl]oxymethyl]oxan-2-yl]oxy-2,3-dihydrochromen-4-one).

[105]. Although it has a low bioavailability due to the rutinose moiety attached to the flavonoid [106], hesperidin can be converted to glucuronides and sulfoglucuronides, which have been shown to be excreted in urine nearly 24 hours after the orange juice ingestion [107]. In a randomized controlled trial, Nielsen et al. [108] demonstrated that removal of the rhamnose group to yield hesperetin-7-glucoside improved the bioavailability of the aglycone hesperetin. Suggesting that additional interventions are required to improve the bioavailability of citrus flavonoids such as hesperidin.

Increasing data has supported the notion that hesperidin possesses increased potential to lower raised blood glucose and lipid levels in various models of type 2 diabetes [109–111]. When administered in rats subjected to swimming exercise, this citrus flavonoid improved the biochemical and antioxidant profile of the animals [112]. This compound may induce its therapeutic effect through the regulation of genes implicated in insulin signaling such as insulin receptor substrate 1, GLUT2/4, and those linked with lipid metabolism, including sterol regulatory element-binding protein 1c (SREBP-1c), fatty acid synthase (FAS) and acetyl-CoA carboxylase [113]. Although data on its effect on skeletal muscle is currently limited, it can reverse IR by reducing muscle glycogen content and ischemia-reperfusion injury while promoting myogenic differentiation through the activation of MyoD-mediated myogenin expression in cultured cells and animals [109, 114, 115].

8. Luteolin

Luteolin (PubChem CID: 5280445) is a flavone glycoside (2-(3,4-Dihydroxyphenyl)- 5,7-dihydroxy-4-chromenone) with the molecular formula $C_{15}H_{10}O_6$ (MW 286.239 g/mol) that is rich in various dietary sources such as fruits, vegetables, and teas (**Figure 7**) [116]. As with most flavonoids, during its metabolism luteolin is broken down to its glucuronides, which can eventually pass through intestinal mucosa as shown by Yasuda and colleagues [117]. Although studies reporting on the pharmacokinetic profile of luteolin in human subjects are limited, this flavone is quickly absorbed in rats and can be detected in urine and feces while showing a slow elimination rate [118]. Furthermore, luteolin from peanut hull extract can be easily absorbed compared to the pure compound, with its absorption more efficient in the




Figure 7. The chemical structure of luteolin (2-(3,4-dihydroxyphenyl)-5,7-dihydroxy-4-chromenone).

jejunum and duodenum than in the colon and ileum [119]. Alternatively, luteolin-loaded solid lipid nanoparticles prepared by hot microemulsion ultrasonic technique can also improve the solubility and increase the compound concentration in plasma of rats [120].

In addition to its strong antioxidant effects [121], *in vitro* experiments have provided evidence that luteolin possesses chemopreventive and anti-inflammatory properties [122, 123]. Hydroxyl groups and 2–3 double bond remain key structural features of luteolin that are linked to its enhanced therapeutic effect [124]. Recent studies show that this flavone attenuates hepatic steatosis and IR by upregulating PPAR γ protein expression and activating AMPK α 1 signaling, which may be linked to the improvement in circulating FFA levels in diet-induced obese mice [125, 126]. However, only a few studies have reported on the effect of luteolin on the skeletal muscle. Available literature has reported on its effect in preventing lipopolysaccharide-induced muscle atrophy, oxidative stress-induced tissue injury and inflammation, partly through regulation of atrogen-1/MAFbx expression, and c-Jun N-terminal kinases (JNK) phosphorylation reported on [127–129].

9. Naringenin

Naringenin (PubChem CID: 932) is a flavanone (5,7-dihydroxy-2-(4-hydroxyphenyl)chroman-4-one) with the molecular formula $C_{15}H_{12}O_5$ (MW 272.256 g/mol) that is also predominantly found in citrus fruits (**Figure 8**) [130]. The chemical structure of naringenin comprises three hydroxy groups at the 4', 5, and 7 carbons while its glycoside, naringin contains an additional disaccharide neohesperidose that is linked via its carbon end. Although naringenin can be




Figure 8. The chemical structure of naringenin (5,7-dihydroxy-2-(4-hydroxyphenyl)chroman-4-one).

detected as monoglucuronides in plasma and urine after ingestion of orange fruit juice in human subjects [131], the bioavailability of naringenin can be influenced by its glycosidic moiety. Felgines et al. [132] demonstrated that kinetics of absorption of naringenin and naringenin-7-glucoside was similar. In addition, naringenin-7-rhamnoglucoside exhibited a delay in its intestinal absorption, resulting in decreased bioavailability after ingestion in rats. On the other hand, complexation of naringenin with hydroxypropoyl- β -cyclodextrin has been another viable alternative to improve the bioavailability of naringenin, which is important to enhance its therapeutic potential [133].

Naringenin is among the well-studied citrus flavonoids shown to prevent complications associated with IR and the metabolic syndrome. Its role in preventing the deterioration in skeletal muscle mass and protecting against metabolic associated complication is summarized. In low-density lipoprotein (LDL) receptor-null ($Ldlr^{-/-}$) mice fed HFD, this flavanone reduced fasting hyperinsulinemia, improved glucose utilization and increased insulin sensitivity through regulation of SREBP-1c-mediated lipogenesis [134]. It stimulated glucose uptake but failed to have a significant effect on basal or insulin-stimulated AKT phosphorylation while significantly increasing AMPK phosphorylation/activation in cultured L6 myotubes [135]. Bhattacharya and colleagues showed that naringenin stimulates glucose uptake, indicating a dependence on GLUT4 activity as well as phosphatidylinositol-4,5-bisphosphate 3-kinase (PI3K) and/or p38MAPK activity [136]. Maintenance of muscle mass by reducing muscle diacylglycerol content, improving hyperinsulinemia, promoting phosphorylation of p38/MAPK via estrogen receptor beta ($ER\beta$), lowering reactive oxygen species (ROS) production, and enhancing tyrosine phosphorylation are other mechanisms associated with protective effect of naringenin in either cultured cells or in vivo animal models [137–140].

10. Quercetin

Quercetin (PubChem CID: 5280343) is classified as a flavonol (2-(3,4-dihydroxyphenyl)-3,5,7-trihydroxy-4H-chromen-4-one) with the molecular formula $C_{15}H_{10}O_7$ (MW 302.238 g/mol) that is abundantly found in various fruits and vegetables (**Figure 9**) [141]. Quercetin is one of the most abundant dietary flavonoids that is rapidly metabolized to glucuronides and sulfates that can be detected in plasma and urine [142]. Although oral bioavailability of quercetin remains




Figure 9. The chemical structure of quercetin (2-(3,4-dihydroxyphenyl)-3,5,7-trihydroxy-4H-chromen-4-one).

low, the type of sugar moiety attached to its structure may affect its absorption. This has been as demonstrated with quercetin glycosides from onion which have a higher absorption rate compared to apple-derived quercetin [143, 144]. Quercetin-4'-O-glucoside and quercetin-3-O-rutinoside (rutin) are one the accomplished glycosides of quercetin, and their absorption rate and extent can be influenced by plant matrix as demonstrated by Graefe and colleagues [145]. However, it is clear that further investigations into improvement strategies for pure quercetin aglycone are required to improve the therapeutic potential of this flavonol.

Quercetin exhibits a wide range of biological functions. Although Stewart et al. [146] failed to show any beneficial effect of quercetin against IR in diet induced-obese mice, other researchers have shown that this flavonol plays a major role in modulating several signaling pathways to reverse metabolic syndrome and improve skeletal muscle function, either in vitro on cultured cells or in vivo in animals and samples from human subjects [148–168]. In L6 myotubes and skeletal muscle of genetical modified (*ob/ob*) mice, quercetin improved insulin sensitivity by increasing GLUT4 expression [147]. Several studies using different experimental models have also demonstrated the positive effect of quercetin in improving skeletal muscle insulin sensitivity through enhanced uptake of glucose, and reducing oxidative stress or inflammation-induced damage, with modulation of tumor necrosis alpha (TNF- α), AKT, peroxisome proliferator-activated receptor gamma coactivator 1-alpha (PGC-1 α) and AMPK as prime pathways involved in the process [148–158]. The therapeutic potential of quercetin extends to its preventative effect against ischemia–reperfusion injury, as well as strengthening muscle fibers through the modulation of calcium homeostasis, and enhancing intracellular antioxidants [159–168].

11. Resveratrol

Resveratrol (PubChem CID: 445154) is a phytoalexin stilbenoid (3,5,4'-trihydroxy-trans-stilbene) with the molecular formula $C_{14}H_{12}O_3$ (MW 228.247 g/mol) that is present in abundant amounts in various food sources such as grapes, blueberries, and red wine (**Figure 10**) [169]. Upon ingestion, resveratrol can be metabolized to form conjugated sulfates and glucuronides, namely resveratrol monosulfate, monosulfate dihydroresveratrol, and monoglucuronide dihydroresveratrol, as reviewed by Gambini and colleagues [170]. Although the bioavailability of resveratrol is considered low, it can vary depending on the method of administration and type of dietary source ingested [171]. The dimethyl ether analog of resveratrol, pterostilbene,




Figure 10. The chemical structure of resveratrol (3,5,4'-trihydroxy-trans-stilbene).

has been shown to exhibit a higher bioavailability, in terms of total plasma levels of both the parent compound and metabolites than does resveratrol [172]. However, Li et al. [173] showed that intravenous and oral pharmacokinetic characteristics of trans-resveratrol can be improved through encapsulating with PP123 self-assembling lecithin-based mixed polymeric micelles. Suggesting that alternative methods to improve the bioavailability of resveratrol are required, which may translate to enhanced therapeutic potential *in vivo*.

Resveratrol has displayed a variety of antidiabetic effects in rodent models. In addition, resveratrol attenuates thermal hyperalgesia, cold allodynia, as well as raised serum lipid levels [174–176]. In diabetic individuals, resveratrol administration is associated with significantly improved glucose and insulin control [177]. The systematic search of evidence linking resveratrol and IR in skeletal muscle revealed up to 18 studies published between 2007 and 2017, with 9 papers produced between 2016 and 2017, suggesting that this phytoalexin stilbenoid is increasingly explored for therapeutic effect against metabolic associated complications. Although Williams and colleagues showed no effect on insulin signaling pathways [178], stimulation of glucose uptake by resveratrol in cultured C2C12 cells or skeletal muscle has been linked with activation of extracellular signal-related kinase/p38/PI3K [179]. Its effect in promoting glucose uptake and improving insulin sensitivity was also associated with increased NAD-dependent protein deacetylase sirtuin-1 (SIRT1) expression, activation of AMPK while abolishing phosphorylation of extracellular signal-regulated kinase 1/2 (ERK1/2), JNK, and I κ B kinase α/β (IKK α /IKK β) [180–190]. Other documented beneficial effect of resveratrol includes inhibiting ischemia–reperfusion injury through its potent antioxidant properties [191], reducing cell proliferation through upregulating PGC-1 α [192], promoting muscle regeneration and attenuating the impact of ROS [193], and elevated forearm skeletal muscle mitochondrial capacity [194].

12. Rutin

Rutin (PubChem CID: 5280805) is a glycoside combining the flavonol quercetin and the disaccharide rutinose (2-(3,4-dihydroxyphenyl)-5,7-dihydroxy-3-[α -L-rhamnopyranosyl-(1 \rightarrow 6)- β -D-glucopyranosyloxy]-4H-chromen-4-one) with the molecular formula C₂₇H₃₀O₁₆ (MW 610.521 g/mol) that is found in many plants and fruits, as well as tea infusions (**Figure 11**) [195]. Upon oral administration, rutin can be metabolized into sulfates and glucuronides of quercetin that are detected in blood, whereas unchanged forms of rutin and quercetin were not detected [142, 196]. Although quercetin glycosides from onions demonstrate an enhanced absorptive capacity than pure aglycones [143, 144], some studies have showed that rutin has a lower oral absorption rate than quercetin [142, 197]. However, as with the use of natural deep eutectic solvents [198], alternative methods to improve the absorptive capacity of rutin is tested to improve therapeutic effect *in vivo*.

Like quercetin, rutin exhibits a wide variety of biological properties, mostly attributed to its strong antioxidant properties [199, 200]. It is accomplished that rutin displays enhanced potential




Figure 11. The chemical structure of rutin (2-(3,4-dihydroxyphenyl)-5,7-dihydroxy-3-[α -L-rhamnopyranosyl-(1 \rightarrow 6)- β -D-glucopyranosyloxy]-4H-chromen-4-one).

to improve insulin sensitivity by regulating genes involved in glucose and lipid metabolism such as GLUT4, PPAR γ , and tyrosine phosphatase 1B in cultured cells or skeletal muscle of rodents [201–204]. However, from the study by Zyma et al. [205], that demonstrated that rutin induces conformational changes in the myosin structure of skeletal muscle of rabbits accompanied by an increase in ATPase activity, accumulative evidence has supported muscle strengthening capacity of this polyphenol. For example, Su et al. [206] presented data showing that rutin promoted skeletal muscle endurance capacity by modulating markers of mitochondrial biogenesis such as PGC-1 α and SIRT1 expression in ICR mice subjected to a weight-loaded forced swim test. These findings were further supported by data showing that rutin increased the mitochondrial size and mitochondrial DNA content as well as gene expression related to mitochondrial biogenesis, such as PGC1- α , NRF-1, transcription factor A, and SIRT1 [207, 208].

13. Sulforaphane

Sulforaphane (PubChem CID: 5350) is an isothiocyanate (1-isothiocyanato-4-methylsulfinylbutane) with the molecular formula $C_6H_{11}NOS_2$ (MW 177.28 g/mol) that is found in cruciferous vegetables such as cabbages, broccoli, and brussels sprouts (**Figure 12**) [209]. Although sulforaphane displays a dose-dependent pharmacokinetic behavior, as higher doses show reduced absorptive potential, lower doses of the compound can be rapidly absorbed in rats following intravenous administration, with the absolute bioavailability being able to reach 82% [210]. In human subjects consuming fresh broccoli sprouts or the broccoli sprout extract, with each estimated to provide 200 μ mol sulforaphane daily, the




Figure 12. The chemical structure of sulforaphane (1-isothiocyanato-4-methylsulfinylbutane).

compound metabolites were found to be three times higher in plasma and urine of sprout consumers, suggesting enhanced sulforaphane absorption from sprouts [211]. Therefore, dietary form and dosing schedule of sulforaphane may influence impact absorption and therapeutic potential in human subjects.

Sulforaphane has received a considerable interest due to its ability to simultaneously control multiple cellular targets involved in various metabolic complications. For instance, in rats fed HFD, this isothiocyanate has displayed an enhanced hypoglycemic potential as well as the elevation of GLUT3 expression in the cerebral cortex and hypothalamus, leading to improved glucose tolerance [212]. Other studies [213, 214] have supported the beneficial effect of sulforaphane or its stable precursor glucoraphanin, to reverse IR, mostly through its robust antioxidant properties. In skeletal muscle, sulforaphane has exhaustive exercise-induced muscle damage, reducing muscle glycogen content, and enhanced exercise endurance capacity through inhibition of pro-inflammatory response and enhancing antioxidant response by upregulating NRF2 expression [215–220].

14. Conclusions

Natural compounds have gained popularity for their potential beneficial effect to fight metabolic diseases due to their less adverse effect compared to synthetic drugs. Furthermore, natural compounds serve as a valuable source for the discovery of new drugs. Currently, knowledge shows that natural compounds can ameliorate IR, however, the gap in scientific evidence of plant-derived therapeutic benefits still exist due to the slow rate of translation of animal studies findings into human clinical trials. In this chapter, evidently reported the great potential and the future promise of natural compounds for the management and treatment of metabolic disorders, specifically IR, obesity, and T2D. Therefore, further research is required to assess the use of natural compounds alone or in combination with well know antidiabetic drugs might result in synergistic and enhanced effects in combating metabolic diseases.

Acknowledgements

This work was supported by the Biomedical Research and Innovation Platform of the South African Medical Research Council (SAMRC) baseline funding and the South African National Research Foundation (NRF; grant number 87836 to SE Mazibuko-Mbeje). The grant holders acknowledge that opinions, findings, and conclusions or recommendations expressed in any publication generated by the SAMRC or NRF supported research are those of the authors, and that the SAMRC and NRF accept no liability whatsoever in this regard. PVD was partially supported as a Post-Doctoral Fellow by funding from the SAMRC.

Abbreviations

AKT	protein kinase B
AMPK	5' AMP-activated protein kinase
ATP	adenosine triphosphate
CD36	cluster of differentiation 36
DPPH	2,2-diphenyl- β -picrylhydrazyl
ER β	estrogen receptor beta
FAS	fatty acid synthase
FFA	free fatty acid
GLUT	glucose transporter
HFD	high fat diet
IR	insulin resistance
IRS-1	suppressing insulin receptor substrate 1
JNK	c-Jun N-terminal kinases
LDL	low density lipoprotein
LKB1	serine/threonine kinase 11
MAPK	mitogen-activated protein kinase
MW	molecular weight
NF- κ B	nuclear factor kappa-light-chain-enhancer of activated B cells
NRF2	nuclear factor (erythroid-derived 2)-like 2
PGC-1 α	peroxisome proliferator-activated receptor gamma coactivator 1-alpha
PI3K	phosphatidylinositol-4,5-bisphosphate 3-kinase
PPAR	peroxisome proliferator-activated receptor
Prmt7	protein arginine methyltransferase 7
SIRT1	NAD-dependent protein deacetylase sirtuin-1
SREBP-1c	sterol regulatory element-binding protein 1c
T2D	type 2 diabetes mellitus
UCP	uncoupling protein

Author details

Sithandiwe Eunice Mazibuko-Mbeje^{1,2*}, Phiwayinkosi V. Dlodla¹, Bongani B. Nkambule³, Nnini Obonye¹ and Johan Louw^{1,4}

*Address all correspondence to: sithandiwe.mazibuko@mrc.ac.za

1 Biomedical Research and Innovation Platform, South African Medical Research Council, Tygerberg, South Africa

2 Division of Medical Physiology, Faculty of Health Sciences, Stellenbosch University, Tygerberg, South Africa

3 School of Laboratory Medicine and Medical Sciences (SLMMS), College of Health Sciences, University of KwaZulu-Natal, Durban, South Africa

4 Department of Biochemistry and Microbiology, University of Zululand, South Africa

References

- [1] Garg N et al. Natural products as mediators of disease. *Natural Product Reports*. 2017; **34**(2):194-219
- [2] Ansari N, Khodagholi F. Natural products as promising drug candidates for the treatment of Alzheimer's disease: Molecular mechanism aspect. *Current Neuropharmacology*. 2013;**11**(4):414-429
- [3] World Health Organization. Overview of Malaria Treatment. 2017. Available from: <http://www.who.int/malaria/areas/treatment/overview/en/>. [Accessed: 01 February 2018]
- [4] Bailey CJ. Metformin: Historical overview. *Diabetologia*. 2017;**60**(9):1566-1576
- [5] Bodmer M et al. Metformin, sulfonylureas, or other antidiabetes drugs and the risk of lactic acidosis or hypoglycemia: A nested case-control analysis. *Diabetes Care*. 2008; **31**(11):2086-2091
- [6] Rios JL, Francini F, Schinella GR. Natural products for the treatment of type 2 diabetes mellitus. *Planta Medica*. 2015;**81**(12-13):975-994
- [7] Joya-Galeana J et al. Effects of insulin and oral anti-diabetic agents on glucose metabolism, vascular dysfunction and skeletal muscle inflammation in type 2 diabetic subjects. *Diabetes/Metabolism Research and Reviews*. 2011;**27**(4):373-382
- [8] Cetrone M, Mele A, Tricarico D. Effects of the antidiabetic drugs on the age-related atrophy and sarcopenia associated with diabetes type II. *Current Diabetes Reviews*. 2014;**10**(4):231-237
- [9] PubChem Compound Database CID: 5280443. National Center for Biotechnology Information. Available from: <https://pubchem.ncbi.nlm.nih.gov/compound/5280443>. [Accessed: 05 January 2018]

- [10] Perkin AGPJ. The colouring matters of the Indian dye stuff asbarg, *Delphinium zalil*. Journal of the Chemical Society, Transactions. 1898;**73**(267)
- [11] Griffiths LA, Smith GE. Metabolism of apigenin and related compounds in the rat. Metabolite formation in vivo and by the intestinal microflora in vitro. The Biochemical Journal. 1972;**128**(4):901-911
- [12] Gradolatto A et al. Pharmacokinetics and metabolism of apigenin in female and male rats after a single oral administration. Drug Metabolism and Disposition. 2005;**33**(1):49-54
- [13] Gradolatto A et al. Metabolism of apigenin by rat liver phase I and phase II enzymes and by isolated perfused rat liver. Drug Metabolism and Disposition. 2004;**32**(1):58-65
- [14] Wan L et al. Quantitative determination of apigenin and its metabolism in rat plasma after intravenous bolus administration by HPLC coupled with tandem mass spectrometry. Journal of Chromatography. B, Analytical Technologies in the Biomedical and Life Sciences. 2007;**855**(2):286-289
- [15] Hanske L et al. The bioavailability of apigenin-7-glucoside is influenced by human intestinal microbiota in rats. The Journal of Nutrition. 2009;**139**(6):1095-1102
- [16] Sung B, Chung HY, Kim ND. Role of apigenin in cancer prevention via the induction of apoptosis and autophagy. Journal of Cancer Prevention. 2016;**21**(4):216-226
- [17] Wang IK, Lin-Shiau SY, Lin JK. Induction of apoptosis by apigenin and related flavonoids through cytochrome c release and activation of caspase-9 and caspase-3 in leukaemia HL-60 cells. European Journal of Cancer. 1999;**35**(10):1517-1525
- [18] Jung UJ, Cho YY, Choi MS. Apigenin ameliorates dyslipidemia, hepatic steatosis and insulin resistance by modulating metabolic and transcriptional profiles in the liver of high-fat diet-induced obese mice. Nutrients. 2016;**8**(5)
- [19] Choi WH et al. Apigenin ameliorates the obesity-induced skeletal muscle atrophy by attenuating mitochondrial dysfunction in the muscle of obese mice. Molecular Nutrition & Food Research. 2017;**61**(12)
- [20] Jang YJ et al. Apigenin enhances skeletal muscle hypertrophy and myoblast differentiation by regulating Prmt7. Oncotarget. 2017;**8**(45):78300-78311
- [21] PubChem Compound Database CID: 11282394. National Center for Biotechnology Information. Available from: <https://pubchem.ncbi.nlm.nih.gov/compound/11282394>. [Accessed: 05 January 2018]
- [22] Bramati L et al. Quantitative characterization of flavonoid compounds in rooibos tea (*Aspalathus linearis*) by LC-UV/DAD. Journal of Agricultural and Food Chemistry. 2002;**50**(20):5513-5519
- [23] Stander MA. Analysis of phenolic compounds in rooibos tea (*Aspalathus linearis*) with a comparison of flavonoid-based compounds in natural populations of plants from different regions. Journal of Agricultural and Food Chemistry. 2017;**65**(47):10270-10281

- [24] Muller CJ et al. Potential of rooibos, its major C-glucosyl flavonoids and Z-2-(beta-D-glucopyranoxy)-3-phenylpropenoic acid in prevention of metabolic syndrome. *Critical Reviews in Food Science and Nutrition*. 2016;227-246
- [25] Johnson R et al. Aspalathin from rooibos (*Aspalathus linearis*): A bioactive C-glucosyl dihydrochalcone with potential to target the metabolic syndrome. *Planta Medica*. 2018; **84**(9-10):568-583. DOI: 10.1055/s-0044-100622
- [26] Stalmach A et al. Bioavailability of C-linked dihydrochalcone and flavanone glucosides in humans following ingestion of unfermented and fermented rooibos teas. *Journal of Agricultural and Food Chemistry*. 2009;**57**(15):7104-7111
- [27] Bowles S et al. Intestinal transport characteristics and metabolism of C-glucosyl dihydrochalcone, aspalathin. *Molecules*. 2017;**22**(4)
- [28] Von Gadow A, Joubert E, Hansmann CF. Comparison of the antioxidant activity of aspalathin with that of other plant phenols of rooibos tea (*Aspalathus linearis*), α -tocopherol, BHT, and BHA. *Journal of Agricultural and Food Chemistry*. 1997;**45**:632-638
- [29] Gregorio BM et al. The potential role of antioxidants in metabolic syndrome. *Current Pharmaceutical Design*. 2016;**22**(7):859-869
- [30] Joubert E et al. Superoxide anion and α,α -diphenyl- β -picrylhydrazyl radical scavenging capacity of rooibos (*Aspalathus linearis*) aqueous extracts, crude phenolic fractions, tannin and flavonoids. *Food Research International*. 2004;**37**(2):133-138
- [31] Kondo M et al. Xanthine oxidase inhibitory activity and hypouricemic effect of aspalathin from unfermented rooibos. *Journal of Food Science*. 2013;**78**(12):H1935-H1939
- [32] Snijman PW et al. Antioxidant activity of the dihydrochalcones aspalathin and nothofagin and their corresponding flavones in relation to other rooibos (*Aspalathus linearis*) flavonoids, epigallocatechin gallate, and Trolox. *Journal of Agricultural and Food Chemistry*. 2009;**57**(15):6678-6684
- [33] Chen W et al. Ameliorative effect of aspalathin from rooibos (*Aspalathus linearis*) on acute oxidative stress in *Caenorhabditis elegans*. *Phytomedicine*. 2013;**20**(3-4):380-386
- [34] Van der Merwe JD et al. In vitro hepatic biotransformation of aspalathin and nothofagin, dihydrochalcones of rooibos (*Aspalathus linearis*), and assessment of metabolite antioxidant activity. *Journal of Agricultural and Food Chemistry*. 2010;**58**(4):2214-2220
- [35] Ku SK et al. Aspalathin and nothofagin from rooibos (*Aspalathus linearis*) inhibits high glucose-induced inflammation in vitro and in vivo. *Inflammation*. 2015;**38**(1):445-455
- [36] Ku SK et al. Antithrombotic activities of aspalathin and nothofagin via inhibiting platelet aggregation and FIIa/FXa. *Archives of Pharmacal Research*. 2015;**38**(6):1080-1089
- [37] Kwak S, Han MS, Bae JS. Aspalathin and nothofagin from rooibos (*Aspalathus linearis*) inhibit endothelial protein C receptor shedding in vitro and in vivo. *Fitoterapia*. 2015; **100**:179-186

- [38] Lee W, Bae JS. Anti-inflammatory effects of aspalathin and nothofagin from rooibos (*Aspalathus linearis*) in vitro and in vivo. *Inflammation*. 2015;**38**(4):1502-1516
- [39] Lee W, Kim KM, Bae JS. Ameliorative effect of aspalathin and nothofagin from rooibos (*Aspalathus linearis*) on HMGB1-induced septic responses in vitro and in vivo. *The American Journal of Chinese Medicine*. 2015;**43**:991-1012
- [40] Dlodla PV et al. Aspalathin protects the heart against hyperglycemia-induced oxidative damage by up-regulating Nrf2 expression. *Molecules*. 2017;**22**(1)
- [41] Johnson R et al. Aspalathin, a dihydrochalcone C-glucoside, protects H9c2 cardiomyocytes against high glucose induced shifts in substrate preference and apoptosis. *Molecular Nutrition & Food Research*. 2016;**60**(4):922-934
- [42] Johnson R et al. The transcription profile unveils the cardioprotective effect of aspalathin against lipid toxicity in an in vitro H9c2 model. *Molecules*. 2017;**22**(2)
- [43] Smit SE et al. Myocardial glucose clearance by aspalathin treatment in young, mature, and obese insulin-resistant rats. *Planta Medica*. 2017;**84**(2):75-82
- [44] Johnson R et al. Aspalathin reverts doxorubicin-induced cardiotoxicity through increased autophagy and decreased expression of p53/mTOR/p62 signaling. *Molecules*. 2017;**22**(10)
- [45] Mazibuko SE et al. Aspalathin improves glucose and lipid metabolism in 3T3-L1 adipocytes exposed to palmitate. *Molecular Nutrition & Food Research*. 2015;**59**(11):2199-2208
- [46] Mazibuko SE et al. Amelioration of palmitate-induced insulin resistance in C2C12 muscle cells by rooibos (*Aspalathus linearis*). *Phytomedicine*. 2013;**20**(10):813-819
- [47] Mikami NTJ et al. Green rooibos extract from *Aspalathus linearis*, and its component, aspalathin, suppress elevation of blood glucose levels in mice and inhibit α -amylase and α -glucosidase activities in vitro. *Food Science and Technology Research*. 2015;**21**(2):231-240
- [48] Muller CJ et al. Acute assessment of an aspalathin-enriched green rooibos (*Aspalathus linearis*) extract with hypoglycemic potential. *Phytomedicine*. 2012;**20**(1):32-39
- [49] Najafian M, Najafian B, Najafian Z. The effect of aspalathin on levels of sugar and lipids in streptozotocin-induced diabetic and normal rats. *Zahedan Journal of Research in Medical Sciences*. 2016;**18**(11):e4963
- [50] Son MJ et al. Aspalathin improves hyperglycemia and glucose intolerance in obese diabetic Ob/Ob mice. *European Journal of Nutrition*. 2013;**52**(6):1607-1619
- [51] Kawano A et al. Hypoglycemic effect of aspalathin, a rooibos tea component from *Aspalathus linearis*, in type 2 diabetic model db/db mice. *Phytomedicine*. 2009;**16**(5):437-443
- [52] PubChem Compound Database CID: 2353. National Center for Biotechnology Information. Available from: <https://pubchem.ncbi.nlm.nih.gov/compound/2353#section=Top>. [Accessed: 05 January 2018]

- [53] Zuo F et al. Pharmacokinetics of berberine and its main metabolites in conventional and pseudo germ-free rats determined by liquid chromatography/ion trap mass spectrometry. *Drug Metabolism and Disposition*. 2006;**34**(12):2064-2072
- [54] Hua W et al. Determination of berberine in human plasma by liquid chromatography-electrospray ionization-mass spectrometry. *Journal of Pharmaceutical and Biomedical Analysis*. 2007;**44**(4):931-937
- [55] Chen W et al. Bioavailability study of berberine and the enhancing effects of TPGS on intestinal absorption in rats. *AAPS PharmSciTech*. 2011;**12**(2):705-711
- [56] Kumar A et al. Current knowledge and pharmacological profile of berberine: An update. *European Journal of Pharmacology*. 2015;**761**:288-297
- [57] Lee YS et al. Berberine, a natural plant product, activates AMP-activated protein kinase with beneficial metabolic effects in diabetic and insulin-resistant states. *Diabetes*. 2006;**55**(8):2256-2264
- [58] Turner N et al. Berberine and its more biologically available derivative, dihydroberberine, inhibit mitochondrial respiratory complex I. *Diabetes*. 2008;**57**(5):1414-1418
- [59] Ma X et al. Berberine-induced activation of 5'-adenosine monophosphate-activated protein kinase and glucose transport in rat skeletal muscles. *Metabolism*. 2010;**59**(11):1619-1627
- [60] Zhang Q et al. Berberine moderates glucose and lipid metabolism through multipathway mechanism. *Evidence-Based Complementary and Alternative Medicine*. 2011;**2011**:924851
- [61] Gomes AP et al. Berberine protects against high fat diet-induced dysfunction in muscle mitochondria by inducing SIRT1-dependent mitochondrial biogenesis. *Biochimica et Biophysica Acta*. 2012;**1822**(2):185-195
- [62] Liu C et al. Effects of berberine on amelioration of hyperglycemia and oxidative stress in high glucose and high fat diet-induced diabetic hamsters in vivo. *BioMed Research International*. 2015;**2015**:313808
- [63] Wang H et al. Atrogin-1 affects muscle protein synthesis and degradation when energy metabolism is impaired by the antidiabetes drug berberine. *Diabetes*. 2010;**59**(8):1879-1889
- [64] Perry BD et al. Muscle atrophy in patients with type 2 diabetes mellitus: Roles of inflammatory pathways, physical activity and exercise. *Exercise Immunology Review*. 2016;**22**:94-109
- [65] Pereira GC et al. Mitochondrially targeted effects of berberine [natural yellow 18, 5,6-dihydro-9,10-dimethoxybenzo(g)-1,3-benzodioxolo(5,6-a) quinolizinium] on K1735-M2 mouse melanoma cells: Comparison with direct effects on isolated mitochondrial fractions. *The Journal of Pharmacology and Experimental Therapeutics*. 2007;**323**(2):636-649
- [66] Zhang DS et al. Effect of berberine on the insulin resistance and TLR4/IKKbeta/NF-kappaB signaling pathways in skeletal muscle of obese rats with insulin resistance. *Sichuan Da Xue Xue Bao. Yi Xue Ban*. 2015;**46**(6):827-831

- [67] PubChem Compound Database CID: 969516. National Center for Biotechnology Information. Available from: <https://pubchem.ncbi.nlm.nih.gov/compound/969516>. [Accessed: 06 January 2018]
- [68] Vareed SK et al. Pharmacokinetics of curcumin conjugate metabolites in healthy human subjects. *Cancer Epidemiology, Biomarkers & Prevention*. 2008;**17**(6):1411-1417
- [69] Anand P et al. Bioavailability of curcumin: Problems and promises. *Molecular Pharmaceutics*. 2007;**4**(6):807-818
- [70] Prasad S, Tyagi AK, Aggarwal BB. Recent developments in delivery, bioavailability, absorption and metabolism of curcumin: The golden pigment from golden spice. *Cancer Research and Treatment*. 2014;**46**(1):2-18
- [71] Gupta SC, Patchva S, Aggarwal BB. Therapeutic roles of curcumin: Lessons learned from clinical trials. *The AAPS Journal*. 2013;**15**(1):195-218
- [72] Sharma RA, Steward WP, Gescher AJ. Pharmacokinetics and pharmacodynamics of curcumin. *Advances in Experimental Medicine and Biology*. 2007;**595**:453-470
- [73] Farid M et al. Effects of dietary curcumin or N-acetylcysteine on NF-kappaB activity and contractile performance in ambulatory and unloaded murine soleus. *Nutrition & Metabolism (London)*. 2005;**(2)**:20
- [74] Poylin V et al. The NF-kappaB inhibitor curcumin blocks sepsis-induced muscle proteolysis. *Mediators of Inflammation*. 2008;**2008**:317851
- [75] Na LX et al. Curcumin improves insulin resistance in skeletal muscle of rats. *Nutrition, Metabolism, and Cardiovascular Diseases*. 2011;**21**(7):526-533
- [76] Deng YT et al. Suppression of free fatty acid-induced insulin resistance by phytopolyphenols in C2C12 mouse skeletal muscle cells. *Journal of Agricultural and Food Chemistry*. 2012;**60**(4):1059-1066
- [77] Panzhinskiy E et al. Novel curcumin derivative CNB-001 mitigates obesity-associated insulin resistance. *The Journal of Pharmacology and Experimental Therapeutics*. 2014;**349**(2):248-257
- [78] Ray Hamidie RD et al. Curcumin treatment enhances the effect of exercise on mitochondrial biogenesis in skeletal muscle by increasing cAMP levels. *Metabolism*. 2015;**64**(10):1334-1347
- [79] Receno CN et al. Curcumin supplementation effects on aging skeletal muscle. *The FASEB Journal*. 2017;**31**(Suppl 1)
- [80] Maithilikarpagaselvi N et al. Preventive effect of curcumin on inflammation, oxidative stress and insulin resistance in high-fat fed obese rats. *Journal of Complementary and Integrative Medicine*. 2016;**13**(2):137-143
- [81] Ono T et al. Curcumin ameliorates skeletal muscle atrophy in type 1 diabetic mice by inhibiting protein ubiquitination. *Experimental Physiology*. 2015;**100**(9):1052-1063

- [82] He HJ et al. Curcumin attenuates Nrf2 signaling defect, oxidative stress in muscle and glucose intolerance in high fat diet-fed mice. *World Journal of Diabetes*. 2012;**3**(5):94-104
- [83] PubChem Compound Database CID: 65064. National Center for Biotechnology Information. Available from: <https://pubchem.ncbi.nlm.nih.gov/compound/65064>. [Accessed: 06 January 2018]
- [84] Graham HN. Green tea composition, consumption, and polyphenol chemistry. *Preventive Medicine*. 1992;**21**(3):334-350
- [85] Lee MJ et al. Pharmacokinetics of tea catechins after ingestion of green tea and (–)-epigallocatechin-3-gallate by humans: Formation of different metabolites and individual variability. *Cancer Epidemiology, Biomarkers & Prevention*. 2002;**11**(10 Pt 1):1025-1032
- [86] Mereles D, Hunstein W. Epigallocatechin-3-gallate (EGCG) for clinical trials: More pitfalls than promises? *International Journal of Molecular Sciences*. 2011;**12**(9):5592-5603
- [87] Du GJ et al. Epigallocatechin gallate (EGCG) is the most effective cancer chemopreventive polyphenol in green tea. *Nutrients*. 2012;**4**(11):1679-1691
- [88] Bose M et al. The major green tea polyphenol, (–)-epigallocatechin-3-gallate, inhibits obesity, metabolic syndrome, and fatty liver disease in high-fat-fed mice. *The Journal of Nutrition*. 2008;**138**(9):1677-1683
- [89] Gan L et al. Green tea polyphenol epigallocatechin-3-gallate ameliorates insulin resistance in non-alcoholic fatty liver disease mice. *Acta Pharmacologica Sinica*. 2015;**36**(5):597-605
- [90] Pezeshki A et al. The effect of green tea extract supplementation on liver enzymes in patients with nonalcoholic fatty liver disease. *International Journal of Preventive Medicine*. 2016;**7**:28
- [91] Singh BN, Shankar S, Srivastava RK. Green tea catechin, epigallocatechin-3-gallate (EGCG): Mechanisms, perspectives and clinical applications. *Biochemical Pharmacology*. 2011;**82**(12):1807-1821
- [92] Nagasawa T et al. Induction of oxidatively modified proteins in skeletal muscle by electrical stimulation and its suppression by dietary supplementation of (–)-epigallocatechin gallate. *Bioscience, Biotechnology, and Biochemistry*. 2000;**64**(5):1004-1010
- [93] Dorchies OM et al. Green tea extract and its major polyphenol (–)-epigallocatechin gallate improve muscle function in a mouse model for Duchenne muscular dystrophy. *American Journal of Physiology. Cell Physiology*. 2006;**290**(2):C616-C625
- [94] Meador BM et al. The green tea polyphenol epigallocatechin-3-gallate (EGCG) attenuates skeletal muscle atrophy in a rat model of sarcopenia. *The Journal of Frailty & Aging*. 2015;**4**(4):209-215
- [95] Jung KH et al. Epigallocatechin gallate stimulates glucose uptake through the phosphatidylinositol 3-kinase-mediated pathway in L6 rat skeletal muscle cells. *Journal of Medicinal Food*. 2008;**11**(3):429-434

- [96] Ueda M et al. Epigallocatechin gallate promotes GLUT4 translocation in skeletal muscle. *Biochemical and Biophysical Research Communications*. 2008;**377**(1):286-290
- [97] Liu HW et al. Dietary (–)-Epigallocatechin-3-gallate supplementation counteracts aging-associated skeletal muscle insulin resistance and fatty liver in senescence-accelerated mouse. *Journal of Agricultural and Food Chemistry*. 2015;**63**(38):8407-8417
- [98] Most J et al. Combined epigallocatechin-3-gallate and resveratrol supplementation for 12 wk increases mitochondrial capacity and fat oxidation, but not insulin sensitivity, in obese humans: A randomized controlled trial. *The American Journal of Clinical Nutrition*. 2016;**104**(1):215-227
- [99] Most J et al. A 3-day EGCG-supplementation reduces interstitial lactate concentration in skeletal muscle of overweight subjects. *Scientific Reports*. 2015;**5**:17896
- [100] Ng HLH et al. Acute vascular and metabolic actions of the green tea polyphenol epigallocatechin 3-gallate in rat skeletal muscle. *The Journal of Nutritional Biochemistry*. 2017;**40**:23-31
- [101] Pence BD et al. Effects of exercise and dietary epigallocatechin gallate and beta-alanine on skeletal muscle in aged mice. *Applied Physiology, Nutrition, and Metabolism*. 2016;**41**(2):181-190
- [102] Wang L et al. Epigallocatechin gallate reduces slow-twitch muscle fiber formation and mitochondrial biosynthesis in C2C12 cells by repressing AMPK activity and PGC-1 α expression. *Journal of Agricultural and Food Chemistry*. 2016;**64**(34):6517-6523
- [103] Sae-Tan S et al. (–)-Epigallocatechin-3-gallate increases the expression of genes related to fat oxidation in the skeletal muscle of high fat-fed mice. *Food & Function*. 2011;**2**(2):111-116
- [104] Mirza KA et al. Attenuation of muscle wasting in murine C2C 12 myotubes by epigallocatechin-3-gallate. *Journal of Cachexia, Sarcopenia and Muscle*. 2014;**5**(4):339-345
- [105] PubChem Compound Database CID: 10621. National Center for Biotechnology Information. Available from: <https://pubchem.ncbi.nlm.nih.gov/compound/10621>. [Accessed: 06 January 2018]
- [106] Ameer B et al. Flavanone absorption after naringin, hesperidin, and citrus administration. *Clinical Pharmacology and Therapeutics*. 1996;**60**(1):34-40
- [107] Manach C et al. Bioavailability in humans of the flavanones hesperidin and narirutin after the ingestion of two doses of orange juice. *European Journal of Clinical Nutrition*. 2003;**57**(2):235-242
- [108] Nielsen IL et al. Bioavailability is improved by enzymatic modification of the citrus flavonoid hesperidin in humans: A randomized, double-blind, crossover trial. *The Journal of Nutrition*. 2006;**136**(2):404-408
- [109] Ahmed OM et al. Antidiabetic effects of hesperidin and naringenin in type 2 diabetic rats. *Diabetologia Croatica*. 2012;**41**(2):53-67
- [110] Amiot MJ, Riva C, Vinet A. Effects of dietary polyphenols on metabolic syndrome features in humans: A systematic review. *Obesity Reviews*. 2016;**17**(7):573-586

- [111] Jung UJ et al. The hypoglycemic effects of hesperidin and naringin are partly mediated by hepatic glucose-regulating enzymes in C57BL/KsJ-db/db mice. *The Journal of Nutrition*. 2004;**134**(10):2499-2503
- [112] De Oliveira DM, Dourado GK, Cesar TB. Hesperidin associated with continuous and interval swimming improved biochemical and oxidative biomarkers in rats. *Journal of the International Society of Sports Nutrition*. 2013;**10**:27
- [113] Pu P. Protection mechanisms of hesperidin on mouse with insulin resistance. *Zhongguo Zhong Yao Za Zhi*. 2016;**41**(17):3290-3295
- [114] Jeong H et al. Hesperidin promotes MyoD-induced myogenic differentiation in vitro and in vivo. *British Journal of Pharmacology*. 2011;**163**(3):598-608
- [115] Ekinici Akdemir FN et al. A comparative study on the antioxidant effects of hesperidin and ellagic acid against skeletal muscle ischemia/reperfusion injury. *Journal of Enzyme Inhibition and Medicinal Chemistry*. 2016;**31**(Suppl 4):114-118
- [116] PubChem CID: 5280445. 2017. PubChem Compound Database CID: 10621. National Center for Biotechnology Information. Available from: <https://pubchem.ncbi.nlm.nih.gov/compound/5280445>. [Accessed: 07 January 2018]
- [117] Yasuda MT et al. Absorption and metabolism of luteolin and its glycosides from the extract of *Chrysanthemum morifolium* flowers in rats and Caco-2 cells. *Journal of Agricultural and Food Chemistry*. 2015;**63**(35):7693-7699
- [118] Chen T et al. Absorption and excretion of luteolin and apigenin in rats after oral administration of *Chrysanthemum morifolium* extract. *Journal of Agricultural and Food Chemistry*. 2007;**55**(2):273-277
- [119] Zhou P et al. Intestinal absorption of luteolin from peanut hull extract is more efficient than that from individual pure luteolin. *Journal of Agricultural and Food Chemistry*. 2008;**56**(1):296-300
- [120] Dang H et al. Luteolin-loaded solid lipid nanoparticles synthesis, characterization, & improvement of bioavailability, pharmacokinetics in vitro and vivo studies. *Journal of Nanoparticle Research*. 2014;**16**:2347
- [121] Zhang YC et al. Antioxidant and Nrf2 inducing activities of luteolin, a flavonoid constituent in *Ixeris sonchifolia* Hance, provide neuroprotective effects against ischemia-induced cellular injury. *Food and Chemical Toxicology*. 2013;**59**:272-280
- [122] Lin Y et al. Luteolin, a flavonoid with potential for cancer prevention and therapy. *Current Cancer Drug Targets*. 2008;**8**(7):634-646
- [123] López-Lázaro M. Distribution and biological activities of the flavonoid luteolin. *Mini Reviews in Medicinal Chemistry*. 2009;**9**(1):31-59
- [124] Wang T, Li Q, Bi K. Bioactive flavonoids in medicinal plants: Structure, activity and biological fate. *Asian Journal of Pharmaceutics*. 2018;**13**(1):12-23

- [125] Kwon EY et al. Luteolin attenuates hepatic steatosis and insulin resistance through the interplay between the liver and adipose tissue in mice with diet-induced obesity. *Diabetes*. 2015;**64**(5):1658-1669
- [126] Zhang L et al. Luteolin reduces obesity-associated insulin resistance in mice by activating AMPK α 1 signalling in adipose tissue macrophages. *Diabetologia*. 2016;**59**(10):2219-2228
- [127] Rostoka E et al. Effects of lycopene, indole-3-carbinol, and luteolin on nitric oxide production and iNOS expression are organ-specific in rats. *Arhiv za Higijenu Rada i Toksikologiju*. 2010;**61**(3):275-285
- [128] Shiota C et al. Flavones inhibit LPS-induced atrogen-1/MAFbx expression in mouse C2C12 skeletal myotubes. *Journal of Nutritional Science and Vitaminology (Tokyo)*. 2015;**61**(2):188-194
- [129] Duan FF et al. Antifatigue effect of luteolin-6-C-neohesperidoside on oxidative stress injury induced by forced swimming of rats through modulation of Nrf2/ARE signaling pathways. *Oxidative Medicine and Cellular Longevity*. 2017;**2017**:3159358
- [130] PubChem CID: 932. PubChem Compound Database CID: 10621. National Center for Biotechnology Information. Available from: <https://pubchem.ncbi.nlm.nih.gov/compound/932>. [Accessed: 08 January 2018]
- [131] Brett GM et al. Absorption, metabolism and excretion of flavanones from single portions of orange fruit and juice and effects of anthropometric variables and contraceptive pill use on flavanone excretion. *The British Journal of Nutrition*. 2009;**101**(5):664-675
- [132] Felgines C et al. Bioavailability of the flavanone naringenin and its glycosides in rats. *American Journal of Physiology. Gastrointestinal and Liver Physiology*. 2000;**279**(6):G1148-G1154
- [133] Shulman M et al. Enhancement of naringenin bioavailability by complexation with hydroxypropyl-beta-cyclodextrin. [corrected]. *PLoS One*. 2011;**6**(4):e18033
- [134] Mulvihill EE et al. Naringenin prevents dyslipidemia, apolipoprotein B overproduction, and hyperinsulinemia in LDL receptor-null mice with diet-induced insulin resistance. *Diabetes*. 2009;**58**(10):2198-2210
- [135] Zygmunt K et al. Naringenin, a citrus flavonoid, increases muscle cell glucose uptake via AMPK. *Biochemical and Biophysical Research Communications*. 2010;**398**(2):178-183
- [136] Bhattacharya S et al. Naringenin and falcarinol stimulate glucose uptake and TBC1D1 phosphorylation in porcine myotube cultures. *Journal of Biomedical and Pharmaceutical Research*. 2014;**2**(2):91-98
- [137] Kannappan S, Anuradha CV. Naringenin enhances insulin-stimulated tyrosine phosphorylation and improves the cellular actions of insulin in a dietary model of metabolic syndrome. *European Journal of Nutrition*. 2010;**49**(2):101-109
- [138] Pellegrini M et al. Naringenin modulates skeletal muscle differentiation via estrogen receptor alpha and beta signal pathway regulation. *Genes & Nutrition*. 2014;**9**(5):425

- [139] Assini JM et al. Naringenin prevents obesity, hepatic steatosis, and glucose intolerance in male mice independent of fibroblast growth factor 21. *Endocrinology*. 2015;**156**(6):2087-2102
- [140] Ke JY et al. Citrus flavonoid, naringenin, increases locomotor activity and reduces diacylglycerol accumulation in skeletal muscle of obese ovariectomized mice. *Molecular Nutrition & Food Research*. 2016;**60**(2):313-324
- [141] PubChem CID: 5280343. PubChem Compound Database CID: 10621. National Center for Biotechnology Information. Available from: <https://pubchem.ncbi.nlm.nih.gov/compound/5280343>. [Accessed: 09 January 2018]
- [142] Erlund I et al. Pharmacokinetics of quercetin from quercetin aglycone and rutin in healthy volunteers. *European Journal of Clinical Pharmacology*. 2000;**56**(8):545-553
- [143] Lee J, Mitchell AE. Pharmacokinetics of quercetin absorption from apples and onions in healthy humans. *Journal of Agricultural and Food Chemistry*. 2012;**60**(15):3874-3881
- [144] Hollman PC et al. Absorption of dietary quercetin glycosides and quercetin in healthy ileostomy volunteers. *The American Journal of Clinical Nutrition*. 1995;**62**(6):1276-1282
- [145] Graefe EU et al. Pharmacokinetics and bioavailability of quercetin glycosides in humans. *Journal of Clinical Pharmacology*. 2001;**41**(5):492-499
- [146] Stewart LK et al. Failure of dietary quercetin to alter the temporal progression of insulin resistance among tissues of C57BL/6J mice during the development of diet-induced obesity. *Diabetologia*. 2009;**52**(3):514-523
- [147] Anhe GF et al. Quercetin decreases inflammatory response and increases insulin action in skeletal muscle of Ob/Ob mice and in L6 myotubes. *European Journal of Pharmacology*. 2012;**689**(1-3):285-293
- [148] Dai X et al. Quercetin but not quercitrin ameliorates tumor necrosis factor- α -induced insulin resistance in C2C12 skeletal muscle cells. *Biological & Pharmaceutical Bulletin*. 2013;**36**(5):788-795
- [149] Arias N et al. Quercetin can reduce insulin resistance without decreasing adipose tissue and skeletal muscle fat accumulation. *Genes & Nutrition*. 2014;**9**(1):361
- [150] Henagan TM et al. Dietary quercetin supplementation in mice increases skeletal muscle PGC1 α expression, improves mitochondrial function and attenuates insulin resistance in a time-specific manner. *PLoS One*. 2014;**9**(2):e89365
- [151] Le NH et al. Quercetin protects against obesity-induced skeletal muscle inflammation and atrophy. *Mediators of Inflammation*. 2014;**2014**:834294
- [152] Eid HM et al. The molecular basis of the antidiabetic action of quercetin in cultured skeletal muscle cells and hepatocytes. *Pharmacognosy Magazine*. 2015;**11**(41):74-81
- [153] Dhanya R et al. Quercetin, a lead compound against type 2 diabetes ameliorates glucose uptake via AMPK pathway in skeletal muscle cell line. *Frontiers in Pharmacology*. 2017;**8**:336

- [154] Henagan T et al. Quercetin and red onion extract attenuate high fat diet-induced insulin resistance while increasing energy expenditure and skeletal muscle mitochondrial number despite a decrease in Pgc1a. *The FASEB Journal*. 2014;**28**(Suppl 1)
- [155] Liu K et al. Quercetin oppositely regulates insulin-mediated glucose disposal in skeletal muscle under normal and inflammatory conditions: The dual roles of AMPK activation. *Molecular Nutrition & Food Research*. 2016;**60**(3):551-565
- [156] Kannappan S, Anuradha CV. Insulin sensitizing actions of fenugreek seed polyphenols, quercetin & metformin in a rat model. *The Indian Journal of Medical Research*. 2009;**129**(4):401-408
- [157] Davis JM et al. Quercetin increases brain and muscle mitochondrial biogenesis and exercise tolerance. *American Journal of Physiology. Regulatory, Integrative and Comparative Physiology*. 2009;**296**(4):R1071-R1077
- [158] Casuso RA et al. Oral quercetin supplementation hampers skeletal muscle adaptations in response to exercise training. *Scandinavian Journal of Medicine & Science in Sports*. 2014;**24**(6):920-927
- [159] Shoshan V et al. Quercetin inhibits Ca^{2+} uptake but not Ca^{2+} release by sarcoplasmic reticulum in skinned muscle fibers. *Proceedings of the National Academy of Sciences of the United States of America*. 1980;**77**(8):4435-4438
- [160] McKenna E et al. Dissociation of phospholamban regulation of cardiac sarcoplasmic reticulum Ca^{2+} ATPase by quercetin. *The Journal of Biological Chemistry*. 1996;**271**(40):24517-24525
- [161] Orzechowski A et al. Effect of quercetin and DMSO on skeletal myogenesis from C2C12 skeletal muscle cells with special reference to PKB/Akt activity, myogenin and Bcl-2 expression. *Basic and Applied Myology*. 2001;**11**(1):31-44
- [162] Lee EH, Meissner G, Kim DH. Effects of quercetin on single Ca^{2+} release channel behavior of skeletal muscle. *Biophysical Journal*. 2002;**82**(3):1266-1277
- [163] Mukai R et al. Quercetin prevents unloading-derived disused muscle atrophy by attenuating the induction of ubiquitin ligases in tail-suspension mice. *Journal of Natural Products*. 2010;**73**(10):1708-1710
- [164] Bouitbir J et al. Opposite effects of statins on mitochondria of cardiac and skeletal muscles: A 'mitohormesis' mechanism involving reactive oxygen species and PGC-1. *European Heart Journal*. 2012;**33**(11):1397-1407
- [165] Leelayuwat N et al. Quercetin enhances endurance capacity via antioxidant activity and size of muscle fibre type. *Journal of Pharmacy and Nutrition Sciences*. 2012;**2012**(2):160-164
- [166] Ekinci Akdemir FN et al. Quercetin protects rat skeletal muscle from ischemia reperfusion injury. *Journal of Enzyme Inhibition and Medicinal Chemistry*. 2016;**31**(Suppl 2):162-166

- [167] Spaulding HR et al. Long-term quercetin dietary enrichment partially protects dystrophic skeletal muscle. *PLoS One*. 2016;**11**(12):e0168293
- [168] Kohara A et al. Enzymatically modified isoquercitrin supplementation intensifies plantaris muscle fiber hypertrophy in functionally overloaded mice. *Journal of the International Society of Sports Nutrition*. 2017;**(32)**:14
- [169] PubChem Compound Database CID: 445154. National Center for Biotechnology Information. Available from: <https://pubchem.ncbi.nlm.nih.gov/compound/445154>. [Accessed: 10 January 2018]
- [170] Gambini J et al. Properties of resveratrol: In vitro and in vivo studies about metabolism, bioavailability, and biological effects in animal models and humans. *Oxidative Medicine and Cellular Longevity*. 2015;**2015**:837042
- [171] Boocock DJ et al. Phase I dose escalation pharmacokinetic study in healthy volunteers of resveratrol, a potential cancer chemopreventive agent. *Cancer Epidemiology, Biomarkers & Prevention*. 2007;**16**(6):1246-1252
- [172] Kapetanovic IM et al. Pharmacokinetics, oral bioavailability, and metabolic profile of resveratrol and its dimethylether analog, pterostilbene, in rats. *Cancer Chemotherapy and Pharmacology*. 2011;**68**(3):593-601
- [173] Li TP et al. Physical and pharmacokinetic characterizations of trans-resveratrol (t-rev) encapsulated with self-assembling lecithin-based mixed polymeric micelles (saLMPMs). *Scientific Reports*. 2017;**7**(1):10674
- [174] Su HC, Hung LM, Chen JK. Resveratrol, a red wine antioxidant, possesses an insulin-like effect in streptozotocin-induced diabetic rats. *American Journal of Physiology. Endocrinology and Metabolism*. 2006;**290**(6):E1339-E1346
- [175] Kwon YI, Vatter DA, Shetty K. Evaluation of clonal herbs of *Lamiaceae* species for management of diabetes and hypertension. *Asia Pacific Journal of Clinical Nutrition*. 2006;**15**(1):107-118
- [176] Sharma S, Kulkarni SK, Chopra K. Resveratrol, a polyphenolic phytoalexin attenuates thermal hyperalgesia and cold allodynia in STZ-induced diabetic rats. *Indian Journal of Experimental Biology*. 2006;**44**(7):566-569
- [177] Zhu X et al. Effects of resveratrol on glucose control and insulin sensitivity in subjects with type 2 diabetes: Systematic review and meta-analysis. *Nutrition & Metabolism (London)*. 2017;**14**:60
- [178] Williams CB et al. An examination of resveratrol's mechanisms of action in human tissue: Impact of a single dose in vivo and dose responses in skeletal muscle ex vivo. *PLoS One*. 2014;**9**(7):e102406
- [179] Deng JY et al. Activation of estrogen receptor is crucial for resveratrol-stimulating muscular glucose uptake via both insulin-dependent and -independent pathways. *Diabetes*. 2008;**57**(7):1814-1823

- [180] Bruckbauer A et al. Synergistic effects of leucine and resveratrol on insulin sensitivity and fat metabolism in adipocytes and mice. *Nutrition & Metabolism (London)*. 2012;**9**(1):77
- [181] Goh KP et al. Effects of resveratrol in patients with type 2 diabetes mellitus on skeletal muscle SIRT1 expression and energy expenditure. *International Journal of Sport Nutrition and Exercise Metabolism*. 2014;**24**(1):2-13
- [182] Sin TK et al. SIRT1-dependent myoprotective effects of resveratrol on muscle injury induced by compression. *Frontiers in Physiology*. 2015;**6**:293
- [183] Frendo-Cumbo S, MacPherson RE, Wright DC. Beneficial effects of combined resveratrol and metformin therapy in treating diet-induced insulin resistance. *Physiological Reports*. 2016;**4**(15):e12877
- [184] Gospin R et al. Resveratrol improves insulin resistance with anti-inflammatory and “browning” effect in adipose tissue of overweight humans. *JIM*. 2016;**64**:814-815
- [185] Dugdale HF et al. The role of resveratrol on skeletal muscle cell differentiation and myotube hypertrophy during glucose restriction. *Molecular and Cellular Biochemistry*. 2018;**444**(1-2):109-123
- [186] Godínez Salas ET et al. Effect of the bioactive compounds genistein and resveratrol on insulin resistance in patients with metabolic syndrome. *The FASEB Journal*. 2017;**31**(1)
- [187] Sadeghi A et al. Resveratrol ameliorates palmitate-induced inflammation in skeletal muscle cells by attenuating oxidative stress and JNK/NF-kappaB pathway in a SIRT1-independent mechanism. *Journal of Cellular Biochemistry*. 2017;**118**(9):2654-2663
- [188] Sung MM et al. Resveratrol improves exercise performance and skeletal muscle oxidative capacity in heart failure. *American Journal of Physiology. Heart and Circulatory Physiology*. 2017;**312**(4):H842-h853
- [189] Tran HT et al. Resveratrol ameliorates the chemical and microbial induction of inflammation and insulin resistance in human placenta, adipose tissue and skeletal muscle. *PLoS One*. 2017;**12**(3):e0173373
- [190] Zheng J et al. Resveratrol improves insulin resistance of catch-up growth by increasing mitochondrial complexes and antioxidant function in skeletal muscle. *Metabolism*. 2012;**61**(7):954-965
- [191] Elmali N et al. Effects of resveratrol on skeletal muscle in ischemia-reperfusion injury. *Ulusal Travma ve Acil Cerrahi Dergisi*. 2007;**13**(4):274-280
- [192] Kaminski J et al. Dietary resveratrol modulates metabolic functions in skeletal muscle cells. *Journal of Food and Drug Analysis*. 2012;**20**(Suppl 1):398-401
- [193] Bosutti A, Degens H. The impact of resveratrol and hydrogen peroxide on muscle cell plasticity shows a dose-dependent interaction. *Scientific Reports*. 2015;**5**:8093
- [194] Polley KR et al. Influence of exercise training with resveratrol supplementation on skeletal muscle mitochondrial capacity. *Applied Physiology, Nutrition, and Metabolism*. 2016;**41**(1):26-32

- [195] PubChem Compound Database CID: 5280805. National Center for Biotechnology Information. Available from: <https://pubchem.ncbi.nlm.nih.gov/compound/5280805>. [Accessed: 11 January 2018]
- [196] Yang C et al. Bioavailability and metabolic pharmacokinetics of rutin and quercetin in rats. *Journal of Food and Drug Analysis*. 2005;**13**(3):244-250
- [197] Manach C et al. Bioavailability of rutin and quercetin in rats. *FEBS Letters*. 1997;**409**(1):12-16
- [198] Faggian M et al. Natural deep eutectic solvents (NADES) as a tool for bioavailability improvement: Pharmacokinetics of rutin dissolved in proline/glycine after oral administration in rats: Possible application in nutraceuticals. *Molecules*. 2016;**21**(11):1531
- [199] Yang J, Guo J, Yuan J. In vitro antioxidant properties of rutin. *LWT - Food Science and Technology*. 2008;**41**(6):1060-1066
- [200] Gatineau E et al. Chronic intake of sucrose accelerates sarcopenia in older male rats through alterations in insulin sensitivity and muscle protein synthesis. *The Journal of Nutrition*. 2015;**145**(5):923-930
- [201] Cai Y et al. Effects of rutin on the expression of PPARgamma in skeletal muscles of db/db mice. *Planta Medica*. 2012;**78**(9):861-865
- [202] Lee DG et al. Effect of rutin from tartary buckwheat sprout on serum glucose-lowering in animal model of type 2 diabetes. *Acta Pharmaceutica*. 2016;**66**(2):297-302
- [203] Hsu CY et al. Rutin potentiates insulin receptor kinase to enhance insulin-dependent glucose transporter 4 translocation. *Molecular Nutrition & Food Research*. 2014;**58**(6):1168-1176
- [204] Kappel VD et al. Rutin potentiates calcium uptake via voltage-dependent calcium channel associated with stimulation of glucose uptake in skeletal muscle. *Archives of Biochemistry and Biophysics*. 2013;**532**(2):55-60
- [205] Zyma VL et al. Interaction of flavonoid compounds with contractile proteins of skeletal muscle. *General Physiology and Biophysics*. 1988;**7**(2):165-175
- [206] Su KY et al. Rutin, a flavonoid and principal component of *Saussurea involucreata*, attenuates physical fatigue in a forced swimming mouse model. *International Journal of Medical Sciences*. 2014;**11**:528-537
- [207] Seo S et al. Rutin increases muscle mitochondrial biogenesis with AMPK activation in high-fat diet-induced obese rats. *Nutrients*. 2015;**7**(9):8152-8169
- [208] Nieman DC et al. Quercetin's influence on exercise performance and muscle mitochondrial biogenesis. *Medicine and Science in Sports and Exercise*. 2010;**42**(2):338-345
- [209] PubChem Compound Database CID: 5350. National Center for Biotechnology Information. Available from: <https://pubchem.ncbi.nlm.nih.gov/compound/5350>. [Accessed: 12 January 2018]

- [210] Hanlon N et al. Absolute bioavailability and dose-dependent pharmacokinetic behaviour of dietary doses of the chemopreventive isothiocyanate sulforaphane in rat. *The British Journal of Nutrition*. 2008;**99**(3):559-564
- [211] Atwell LL et al. Absorption and chemopreventive targets of sulforaphane in humans following consumption of broccoli sprouts or a myrosinase-treated broccoli sprout extract. *Molecular Nutrition & Food Research*. 2015;**59**(3):424-433
- [212] Souza CG et al. Chronic sulforaphane oral treatment accentuates blood glucose impairment and may affect GLUT3 expression in the cerebral cortex and hypothalamus of rats fed with a highly palatable diet. *Food & Function*. 2013;**4**(8):1271-1276
- [213] Nagata N et al. Glucoraphanin ameliorates obesity and insulin resistance through adipose tissue browning and reduction of metabolic endotoxemia in mice. *Diabetes*. 2017;**66**(5):1222-1236
- [214] Fu J et al. Divergent effects of sulforaphane on basal and glucose-stimulated insulin secretion in beta-cells: Role of reactive oxygen species and induction of endogenous antioxidants. *Pharmaceutical Research*. 2013;**30**(9):2248-2259
- [215] Malaguti M et al. Sulforaphane treatment protects skeletal muscle against damage induced by exhaustive exercise in rats. *Journal of Applied Physiology*. 2009;**107**(4):1028-1036
- [216] Fan H et al. Sulforaphane causes a major epigenetic repression of myostatin in porcine satellite cells. *Epigenetics*. 2012;**7**(12):1379-1390
- [217] Whitman SA et al. Nrf2 modulates contractile and metabolic properties of skeletal muscle in streptozotocin-induced diabetic atrophy. *Experimental Cell Research*. 2013;**319**(17):2673-2683
- [218] Sun CC et al. Sulforaphane attenuates muscle inflammation in dystrophin-deficient mdx mice via NF-E2-related factor 2 (Nrf2)-mediated inhibition of NF-kappaB signaling pathway. *The Journal of Biological Chemistry*. 2015;**290**(29):17784-17795
- [219] Oh S et al. Nuclear factor (erythroid derived 2)-like 2 activation increases exercise endurance capacity via redox modulation in skeletal muscles. *Scientific Reports*. 2017;**7**
- [220] Uruno A et al. Nrf2-mediated regulation of skeletal muscle glycogen metabolism. *Molecular and Cellular Biology*. 2016;**36**(11):1655-1672

