We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists



186,000

200M



Our authors are among the

TOP 1% most cited scientists





WEB OF SCIENCE

Selection of our books indexed in the Book Citation Index in Web of Science™ Core Collection (BKCI)

Interested in publishing with us? Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected. For more information visit www.intechopen.com



Chapter

Decadal Pollution Assessment and Monitoring along the Kenya Coast

Eric Ochieng Okuku, Kiteresi Linet Imbayi, Owato Gilbert Omondi, Wanjeri Veronica Ogolla Wayayi, Mwalugha Catherine Sezi, Kombo Mokeira Maureen, Stephen Mwangi and Nancy Oduor

Abstract

Marine contamination arising from land-based sources is on the rise along the Kenyan Coast. We carried out a decadal pollution survey between 2008 and 2018 to determine the levels of various pollutants (nutrients, trace metals, persistent organic pollutants, and ²¹⁰Po) in water, sediment, and biota collected from selected locations in Kenya. Nutrient levels in water ranged between <0.10 and 1560.00, <0.10 and 1320.00, and <0.10 and 3280.00 μ g/L for PO₄³⁻-P, (NO₂⁻ + NO₃⁻)-N, and NH₄⁺-N, respectively, while Chl-a values ranged between 0.02 and 119.37 mg/L. Total PAH, PCBs, and OCPs in sediment from the studied locations ranged from BDL-37800, 0.012–7.99 and BDL-6.10 ng/g. High level of PAH in Kilindini port was primarily from petroleum sources. DDD + DDE/DDT ratio was above 0.5 suggesting historical input. Sediment trace metal concentration from selected locations in Kenyan estuaries had various ranges, that is, Al ($0.06-9804284.00 \mu g/g$), Zn (3.82–367.20 μg/g), Cu (7.5–169.60), Cd (DL –2.40 μg/g), Mn (BDL-169.60 μg/g), Cr (2.55–239.10 µg/g), and Pb (BDL-135.60) µg/g dw. Surface sediment ²¹⁰Po activities ranged between 20.29 and 43.44 Bq kg⁻¹ dw. Chl-a and PO₄³⁻-P data revealed enhance primary productivity in Mombasa peri-urban creeks and estuarine areas. Although the reported concentrations of trace metals and POPs are low in most locations from Kenya, there is a potential risk of bioaccumulation of these contaminants in marine biota; thus, there is a need for continuous monitoring to protect both ecosystem and human health.

Keywords: marine pollution, nutrients, heavy metal

1. Introduction

Kenya is experiencing an increase in human population, urbanization, and industrialization which is characterized by massive utilization of natural resources coupled with an increase in waste production [1]. UNEP GEF-funded WIOLaB project identified that over 70% of pollutants entering marine and coastal ecosystems originate from the land [2]. Many pollutants from these land-based sources (such as sewage, oil hydrocarbons, sediments, nutrients, pesticides, litter, and plastics and toxic wastes) enter the sea via rivers and surface runoff [3]. For instance, sewage is still being indiscriminately discharged into peri-urban creeks of Mombasa [4].

Kenya's coastal waters are thus continuously enriched with large amounts of nutrients, particularly nitrogen (N) and phosphorus (P), silica (Si) and sulfur (S) from anthropogenic activities and wastewater from households and industries [4] despite the fact that effluent pollution had been identified as the most serious of all land-based threats to the marine environment and an area which requires more attention [5]. Excessive nutrient enrichment results in nutrient pollution and eutrophication with severe economic, environmental, and ecological implications to the marine environment due to the shift in the Redfield ratio [6]. Some of the impacts of nutrient enrichment include increased abundance and biomass of phytoplankton and macroalgae [7]; reduced light penetration; increased development of hypoxic, anoxic, and dead zones [8, 9]; changes in phytoplankton community structure and alteration of food chains and webs [10]; development of harmful algal blooms that may produce substances toxic to aquatic organisms or humans [11]; dieback of sea-grasses, algal beds, and corals [12, 13]; loss of biodiversity; and increased incidences and duration of harmful algal blooms [14].

The other class of contaminants that are discharged through sewage, surface runoff, and industrial discharge includes heavy metals and persistent organic pollutants (POPs). Trace metals such as lead (Pb), cadmium (Cd), copper (Cu), chromium (Cr), nickel (Ni), mercury (Hg), arsenic (As), and zinc (Zn) are not easily degraded in the natural environment and are bioaccumulated along the food chain [15]. They are also known to affect productivity, reproduction, and survival of marine organisms and can be hazardous to human health at elevated concentrations [15].

Persistent contamination of organic pollutants (POPs), such as organochlorine pesticides (OCPs), polycyclic aromatic hydrocarbons (PAHs), and polychlorinated biphenyls (PCBs) in the aquatic environment is a worldwide problem [16]. The anthropogenic input of these POPs in the Indian Ocean includes industrial and sewage effluents, storm water drains, shipping activities, spillage, rivers, atmospheric fallout, coastal activities, and natural oil seeps. For instance, these compounds have been used extensively in Kenya for various applications in industries and agriculture [17]. They are characterized by long-term stability (not easily degradable in the environment and can persist in sediments for decades or even centuries), bioaccumulative nature, long-range transport capability, and may have high toxic effects on aquatic living organisms [18, 19]. These chemicals have since been replaced with organophosphate and carbamate, which are less persistent in the environment under the Stockholm Convention on Persistent Organic Pollutants [20, 21]. However, their residues are still present in the marine environment.

Among the natural radionuclides, alpha emitters (e.g. ²¹⁰Po) have been reported to have significant radiological effects resulting from their accumulation in organisms [22]. ²¹⁰Po is a naturally occurring radioactive material (NORM) alpha-emitter occurring in very low concentrations in the environment as a part of the uranium decay chain. ²¹⁰Po has a radioactive half-life of 138.4 days and is produced in the marine environment from the decay of ²¹⁰Pb (which is a daughter isotope derived from ²²⁶Ra dissolved in seawater). ²¹⁰Pb can also be introduced into marine environment directly from the atmosphere from the decay of ²²²Rn [22, 23]. It is ubiquitously distributed in rocks, soils, making up earth's crust, in the atmosphere and in natural waters [24]. It can also be derived in insignificant quantities from lead-containing wastes from uranium, vanadium, and radium refining operations [25]. The activity of ²¹⁰Po in the environment has increased in the past due to human activity such as fossil fuel combustion, use of phosphate fertilizers in agriculture, and discharge of domestic and industrial sewage.

UNEP-GEF WIO-LaB Project identified municipal and industrial effluents, contaminated surface runoff including groundwater and agricultural runoff [2] as the major sources of land-based pollution and microbial contamination, suspended solids, chemical pollution, marine litter (including debris), and eutrophication

(harmful/nuisance algal blooms) as priority pollution categories in the WIO region. In Kenya, limited action has been taken to control or manage pollution. For instance, only a couple of studies have been carried out to determine long-term pollution status or their impacts on marine and coastal ecosystems and the potential human health.

During the assessment and monitoring program, water matrix was analyzed for nutrient pollution whereas sediments were analyzed for trace metals, radionuclides, and POP contamination. Sediment was the preferred matrix for persistent pollutants given that it a known sink and an archive of pollutants brought into the aquatic environment from direct discharges, surface runoff, atmospheric fallout, and nonpoint source [19, 26, 27]. The distribution of these contaminants in sediments can serve as a useful index of pollution and potential environmental risks [17, 28].

2. Materials and methods

Water and sediment samples were collected from the estuarine areas of Ramisi-Vanga system in the Kenyan South Coast; peri-urban creeks of Mombasa; and estuarine areas of the North Coast (Figure 1). Ramisi-Vanga system (Ramisi, Mwena, and Umba) is a low-lying coastal plain submergent complex (below 30 m contour) with extensive cover of mangrove forest, intertidal areas covered with seagrass beds and shallow water lagoons harboring the coral reefs. These critical systems are interlinked through the exchange of water, nutrients, and carbon by the tidally controlled circulation and river discharge [29]. The Kisite-Mpunguti Marine Protected Area (KMMPA) and Shimoni are situated approximately 90 km south of Mombasa with very few scattered fringe mangrove cover on the shoreline though there is no river input into this system apart from sewage and runoff [30]. The area constitutes an important tourist attraction and important resource for the surrounding communities [31]. The main characterizing features of interest around Mombasa peri-urban creeks (Makupa, Kilindini, Mtwapa, and Tudor creeks) include the Kilindini port and the Mombasa Municipal refuse dumpsite, (Kibarani), both of which are found in the vicinity of the Makupa creek stations and the Kenya Meat Commission (KMC) beef factory located within the vicinity of the Tudor creek [32]. Mtwapa creek is situated 25 km north of Mombasa. It is a tidal creek lined by mangrove forests and extensive mud banks. The creek receives freshwater input through seasonal rivers and it is reported to be relatively eutrophic due to direct release of raw sewage into the creek at the vicinity of Shimo la Tewa government prison [30]. These peri-urban creeks were chosen due to anthropogenic activities in the adjacent areas while riverine estuarine systems were chosen given that river-derived material like water, sediments, and pollutants have a tremendous ecological and toxicological influence on coastal zones [35]. North Coast sites comprised of estuarine areas of River Tana which arises from Aberdare range and Mount Kenya passing through the arid and semiarid areas of the eastern part of the country and finally drain into the Indian Ocean in a fan-shaped delta [34] while Sabaki river originates from Athi river where it joins Tsavo river and forms river Galana where it also discharges into the sea. These two rivers affect the marine ecological conditions of Watamu, Malindi, and Lamu [33].

Nutrient sample collection, preservation, shipment, storage, and analysis were done following procedures described by Parsons [36], APHA [37] and measured using Genesys 10S Vis spectroscopy (Thermo Scientific[™]). Sediment samples were collected using Uwitec corer (fitted with a 50-cm plastic sampler, Ø 8 cm) following the procedures provided in IAEA TECDOC-1415. The samples were transported to the laboratory awaiting analysis.



Figure 1.

Map showing sampling sites: Estuaries (Umba, Ramisi, Sabaki, Mwena, and Tana); north coast (Kilifi, Malindi, and Lamu); south coast (Kibuyuni, Shimoni, Wasini, Sii Island, Gazi, and Kisite); and peri-urban creeks of Mombasa (Mtwapa, Tudor, and Makupa).

For analysis of trace metals and alpha-emitter ²¹⁰Po radionuclide, top 5 cm of sediment for each site were pooled, mixed by hand, and freeze-dried, homogenized sieved, dry-weighed, and then digested in a microwave (Microwave Accelerated Reaction system). Elemental analysis of Cu, Zn, Mn, Pb, Cd, and Cr was carried out using inductively coupled plasma mass spectrometer (ICP-MS) (Varian, Australia) while ²¹⁰Po activity was measured using alpha-spectrometry with silicon surface barrier detectors (EG and G) coupled to a PC running Maestro TM data acquisition software [38].

PCBs, OCPs, and PAH were analyzed as described by Thompson et al. [39] where wet samples were freeze-dried using Heto powerdry LL3000 freeze dryer (Thermo

Scientific), then known weight of the sediment sample was spiked with internal standard for PCBs (PCB 30, PCB 103, PCB 155, PCB 198 from Promochem, Dr. Ehrenstorfer GmbH, France), OCPs (d8 4,4′ DDT from Cambridge Isotope Laboratory, France), and PAH (per-deuterated PAHs; Phenanthrene-d10, benzo [a]pyrene-d12, benzo [e] pyrene d12 and benzo [g, h, i]perylene-d12 from Cambridge Isotope Laboratories, Andover, USA; and Fluoranthene-d10, chrysene-d12 and pyrene-d10 from MSD isotopes, Division of Merck Frost Canada INC, Montreal, CND) to quantify the recoveries. The samples were then extracted using START E microwave-assisted extraction system (Milestone, Italy). The extract was concentrated using Rapidvap LABCONCO (Serlabo Technologies, France) and then concentrated samples were subjected to the clean-up process. Thereafter, the extracts were finally concentrated under nitrogen, transferred to 100-µl container and analyzed using GC-MS.

3. Pollutants' source identification and risk assessment

To determine the possible pollution sources of DDT, the approach by Kilunga et al. and Mohammed et al. [18, 40] (that classified sediments with (DDD +DDE)/ DDT ratio greater than 0.5 as historic and ratio less than 0.5 as recent input) was applied. The ratio of DDE to DDD was used in this study to give an idea of the degradation pathway of DDTs in the sediment [41]. DDE/DDD ratio greater than 1 was indicative of anaerobic degradation of DDT into DDD while the ratio less than 1 implies DDT conversion into DDE via aerobic degradation [42].

The pollution data obtained compared with effects range-low (ER-L) and effects range-median (ER-M) values (sediment quality guidelines, SQGs) provided by MacDonald et al. and MacDonald and Ingersoll [43, 44] were used to evaluate the ecotoxicological significance/potential impacts of trace metals, PAH, PCBs, and DDT on benthic organism. ER-L represents the value at which toxicity may be observed in sensitive marine species (i.e., pollution is at a level that may not cause harm to aquatic environment) and ER-M represents the concentration below which adverse effects are expected to occur.

4. Results and discussion

4.1 Nutrient pollution

Results of this study showed high values of nutrient concentration in stations within estuaries compared to oceanic water stations (Figure 2a-d). The highest value of phosphate concentration was recorded in Kisite (304.82 μ g/L) followed by River Sabaki (205.79 μ g/L) and the least was in Kilifi (7.52 μ g/L) whereas the highest values of ammonium were, however, recorded in Mombasa stations of Tudor creek $(81.71 \,\mu g/L)$. The stations in Mombasa were also observed to have high nutrient levels compared to Malindi and Lamu (Figure 2d and b), a condition that can be attributed to the high population associated with municipal and industrial wastes in Mombasa compared to the other towns. It was observed that the nutrient levels for stations in Tana river estuary, which was hypothetically thought to be receiving higher nutrient input from the agricultural and residential population in the river catchment, were strikingly low (Figure 2a). This observation of low nutrient and chlorophyll-a concentration can be explained by the presence of several dams, which form the Seven Folks Dams Scheme. These dams are known to settle down suspended materials with adsorbed nutrients [45], therefore reducing the transportation of these sediments and nutrients downstream. Nutrient ranges recorded



Figure 2.

Showing nutrient phosphates ($\mu g/L$), nitrates ($\mu g/L$), ammonia ($\mu g/L$), and chlorophyll-a ($\mu g/L$) distribution pattern observed in: (a) south coast estuaries: Umba, Ramisi, Sabaki, Mwena, and Tana; (b) north coast open water, stations: Kilifi, Malindi, and Lamu; (c) south coast shoreline area: Kibuy (Kibuyuni), Shimoni, Wasini, Sii Island, Gazi, and Kisite; and (d) peri-urban stations of Mombasa: Mtwapa, Tudor, and Makupa.

in this study are similar to those realized by [46, 47] in the Indian Mangroves of 14–600 and 1.00–80 μ g/L for nitrates and phosphates respectively. Chlorophyll-a showed a distribution pattern almost similar to that of nutrients with low values recorded in Kilifi (0.03 μ g/L) while the highest ranges were observed in River Sabaki (8.19 μ g/L).

4.2 Spatial distribution of heavy metals

The aluminum (Al) levels in the peri-urban creeks had a higher concentration range: $0.06-17483.66 \ \mu g/g$) compared to estuaries (range: 719484.00– 9804284.00 $\ \mu g/g$) and pristine area Gazi Bay (335.00 $\ \mu g/g$). The metal concentration seemed normal since aluminum concentrations are not likely to be significantly affected by anthropogenic aluminum sources [47].

Cr concentration levels in estuaries were higher (range: $101.10-239.10 \mu g/g$) compared to Gazi Bay ($1.58 \mu g/g$), Lamu port (range: $2.55-7.58 \mu g/g$), and periurban creeks (range: $11.53-24.43 \mu g/g$) with an exception of Kilindini harbor that had Cr concentration of 42,338.00 $\mu g/g$. The relatively high concentration of Cr in the river could have been caused by mining activities upstream whereas the high level observed in one of the stations in Kilindini could have resulted from the cement bulk storage at Kilindini harbor which is adjacent to the station [48].

Pb concentration in estuaries was higher (range: $45.70-135.60 \ \mu g/g$) compared to the peri-urban creeks (range: $2.75-40.65 \ \mu g/g$). These high concentrations could be attributed to both natural and anthropogenic Pb point-source discharges, primarily from its use as a fuel additive, which has made it a pervasive and persistent pollutant worldwide [49]. The concentration reported in this study was, however, lower compared to Pattani Bay (range: $79.00-97.00 \ \mu g/g$), Thailand Laptev, Russia (range: $16.00-22.00 \ \mu g/g$; Nolting in press) and the levels previously reported for the Kenyan Coastal Zone (range: $0.13-0.56 \ \mu g/g$; by [50].

The concentration of Zn was higher in the peri-urban creeks (range: 23.36–264.30 μ g/g) compared to the estuaries (range: 144.20–367.20 μ g/g) and Gazi Bay (3.82 μ g/g). High levels of Zn are associated to the dissolution of sacrificial zinc anodes used in marinas, ships, and on leisure boats at the Kenya ports and water sports areas; the adjacent Kibarani dumpsite is also a contributor since most types of wastes in the dumpsite find their way into Makupa creek [51].

Mn level in estuaries had higher concentrations (range: $3809.00-6781.00 \ \mu g/g$) compared to peri-urban creeks (range: $16.10-58.50 \ \mu g/g$) and Gazi Bay ($0.77 \ \mu g/g$). The same trend was observed in Cu levels in which estuaries had the highest concentration (range: $63.30-169.60 \ \mu g/g$) compared to peri-urban creeks (range: $(0.51-52.16 \ \mu g/g)$ and Gazi Bay which was BDL. Cd levels in peri-urban creeks were higher (range: BDL-2.40 \ \mu g/g), compared to the pristine area Gazi Bay ($0.03 \ \mu g/g$) (**Table 1**). These values may be attributed to historical industrial activity around the estuaries, that is, gypsum mining in Tana River County; five cement manufacturing companies located in Athi River: Athi River Mining, Bamburi Portland Cement, East African Portland Cement, Savanna Cement, and Simba Cement; and small landfill along Sabaki river [52, 53].

The mean ²¹⁰Po activities determined in the surface sediment from the three peri-urban creeks ranged between 20.29 and 43.44 Bq kg⁻¹ dw. Mtwapa creek had the highest mean (37.56 \pm 2.14 Bq kg⁻¹ dw) compared to Makupa creek (30.42 \pm 2.16 Bq kg⁻¹ dw) and Tudor creek (28.64 \pm 2.86 Bq kg⁻¹ dw). These results are comparable to those of surface sediments from the Venice lagoon ecosystem in Italy in the range of 26–45 Bq kg⁻¹ [54].

The high levels of ²¹⁰Po activity in Mtwapa creek could have originated from the seasonal river (River Mto Mkuu) which flows into the creek discharging finer sediment from the mainland agriculture activities. In addition, high levels of silt and organic matter were found in Mtwapa creek and this may have resulted in high levels of ²¹⁰Po activity since ²¹⁰Po tends to adsorb on a finer particle of silt (<63 μ m) due to its specific surface area compared to coarse particles (>63 μ m) [49, 50]. The same observation was made by Aközcan and Uğur Görgün [57] in Izmir Bay and Didim who attributed high levels of ²¹⁰Po to Buyuk Menderes river.

4.3 Organic pollutant pollution

PAH concentration in the surface sediment from Ramisi, Sabaki, and Tana rivers was in the range of BDL-0.16, 0.06–7.98, and 0.78–13.20 ng/g, respectively. Higher concentration of PAH in Tana river could be as a result of rapid urbanization, industrialization, gasoline or diesel fuel from fishing boats, and high erosion activities in their catchment areas. The concentrations were, however, lower compared to those reported by Chen and coworkers [58] who recorded concentration ranges of 362.00–15, 667.00 ng/g (dw) for Weihe river, a river draining arid and semiarid regions as Tana river.

The surveyed sites in Lamu, namely, Wange Fish Port, proposed Lamu Port, and Mokowe Fish port, had PAH concentration of less than 10.00 ng/g (dw). This is because the port is still under construction and un-operational. The PAH levels were lower than the levels observed in Kilindini port (range: <50.00–37,800.00 ng/g dw) with a mean concentration of 6086.00 ng/g dw. The PAH level in Kilindini was, however, higher than levels reported by Saleem et al. [59, 53] in Karachi harbor (mean: 27.52 ng/g dw) and [54, 60] in Kaohsiung harbor, Taiwan (range: 472.00–16,201.00 ng/g; mean 5764.00 ng/g). High level of PAH was observed in Kilindini due leakage of combusted and un-combusted fuel from intense shipping activities, oil spills experienced in Makupa creeks, and oil terminal activities in Kilindini

	Pristine environment	Estuaries			Peri-urban		SQG	
Sampling site	Gazi Bay	Sabaki mouth	Tana mouth	Lamu port	Makupa creek	Kilindini harbor	ER-L	ERM-
Al	335.00	9804284.00	719484.00	659.50–1674.90	17483.66	0.06–2.85		NA
As	_		_	BDL-3.54	_	<0.01–2.99	33	NG
Cd	0.027	(- +)	_	BDL-0.13	2.4	<0.02–0.11	5	9.6
Со	0.072	SP	_	_	_	42	_	_
Cr	1.576	101.10	239.10	2.55–7.58		11.53-42,338	80	370.00
Cu	BDL	63.30	169.60	0.51–2.98	52.16	6.57–15.3	70	270.00
Hg	4.623		_	BDL	_	<0.01	0.15	_
Mn	0.757	3809.00	6781.00	16.10–58.5	90.47	7	\	NG
Ni	_	4.90	13.10	_	_	8.47–16.22	30	_
Pb	_	45.70	135.600	BDL	40.65	2.75–11.46	35	218.00
Zn	3.82	144.20	367.20	_	264.3	23.36–38.45	120	410.00

 Table 1.

 Decadal comparison of trace metals of the Kenyan coastal sediment with the set international standards for sediment quality.

port as well as ferry services. This port serves many shipping vessels involved in cargo importation for Kenya as well as the landlocked countries in East Africa. The total OCPs namely HCN, Heptachlor, Hepoxide, 2,4' DDE, Cis chlordane, Trans Nonachlor, 4,4'DDE, 2,4' DDD, 4,4' DDD, 2,4' DDT, 4,4' DDT, and Mirex in sediments from Ramisi, Sabaki, and Tana estuarine areas ranged between ND-0.13, 0.37–1.81, and 0.03–3.94 ng/g respectively. The concentration of OCP reported by this study (range: 0.03–3.94 ng/g) was lower compared to the levels reported for Tana estuary by Lalah et al. [61, 55] (range: <0.003–108.51 ng/g). This confirms the reduction in utilization of the products containing OCPs in Kenya.

The DDT and its metabolites 2,4′ DDE, 4,4′ DDE, 2,4′ DDD, 4,4′ DDD, 2,4′ DDT, and 4,4′ DDT were detected in the surface sediment in both Sabaki and Tana river estuaries. The highest concentrations of $\sum 6$ DDTs were recorded in Tana river (11.57 ng/g dw) compared to Sabaki (2.34 ng/g dw) and Ramisi (0.23 ng/g dw). This suggests that even though DDT was banned in Kenya in 1986, there could still be illegal use of DDT in Kenya [56].

The DDE-to-DDD ratio for Ramisi, Sabaki, and Tana river estuaries were 0.20, 0.16, and 0.17 respectively, suggesting an aerobic degradation of DDT to DDD in the areas via dehydrochlorination and oxidation processes [18, 62]. The study revealed high ratios (above 0.5) for Ramisi (3.38), Sabaki (2.82), and Tana rivers (2.10), indicating a historical input of DDT into the estuaries. The historical input of DDT into marine environment has similarly been reported by Aly Salem et al. [63] for Egyptian Mediterranean Coast, as well as Lu et al. [28] for Poyang Lake, China. This shows the reduced application of DDT in some part of the world.

High concentration of PCBs was observed in River Tana estuary (range: 4.94–7.99 ng/g dw) compared to concentration in Ramisi (ND-0.15 ng/g dw) and Sabaki (0.58–2.40 ng/g dw) which could be attributed to leakage or inadvertent disposal of materials used in transformers and capacitors along the Tana river channel since Tana river holds the Seven Folks hydropower dams which generate approximately 49% of Kenyan electricity.

4.4 Ecological risk assessment of pollution along the Kenyan Coast

According to [64], some of the estuaries and peri-urban stations (Mwena, Ramisi, Sabaki, Umba, Lamu, Kibuyuni, Sii Island, and Makupa) could be classified as eutrophic given that they have phosphates levels >0.021 mg/L whereas Shimoni and Gazi could be classified as oligotrophic. Trace metal concentrations in estuaries (River Sabaki and Tana mouth) except for Pb and Cr (**Table 1**) in the sediments were below the effect range-low (ER-L)/effect range-median (ER-M). This indicates that high concentration may pose adverse effects [65].

Sediment quality criteria and concentration ranges of PAHs, PCBs, and DDT contaminants are summarized in (**Table 2**). The \sum PAHs, \sum PCBs, and \sum DDT concentration in sediment from Ramisi, Sabaki, and Kilindini were below ER-L and ER-M suggesting the low ecological risk of these compounds to sediment-dwelling

	Ramisi	Sabaki	Tana	Kilindini	Lamu	ER-L	ER-M
∑РАН	0.01–0.16	0.79–3.75	0.78–13.2	<50-37,800	<10	4000	35,000
∑PCB	0.12-0.15	0.58–2.40	4.99–7.99	<10	_	50	400
∑DDT	0.01–0.128	0.22–1.57	0.25–5.23	_	_	3	350

Table 2.

Comparison of PAH, DDT, and PCB (ng/g dw) concentrations in sediments of the study area with the sediment quality guidelines (SQG).

organisms. However, the concentration level of \sum DDT was higher than ER-L in Tana estuary but was significantly lower than the ER-M values suggesting that the accumulated DDT in sediments in Tana estuary could have a potential ecotoxicological impact on the benthic fauna.

5. Conclusion

From the study, it is evident that human influences like urbanization and agricultural activities are the major nutrient contributors to the marine environment through continuous discharge of untreated sewage effluents and agricultural runoffs delivered by rivers. Trace metal concentrations in estuaries, peri-urban creeks, and Lamu port were below the ER-L and ER-M indicating that there are no potential adverse effects on marine biota with an exception of Kilindini harbor that had a higher Cr concentration that may pose adverse effects. The concentrations of PAH, PCB, and OCP were generally high in Tana estuary as compared to the other estuaries in this study; however, the concentrations were lower than ER-L suggesting no potential impacts on organisms. It was also observed that the concentration level of \sum DDT was higher in Tana estuary than ER-L and lower for ER-M values suggesting that the accumulated persistent organic pollutants in sediments from Tana estuary could have a potential ecotoxicological impact on the benthic fauna. It is therefore important to put in place effective sewage and wastewater management strategies and implement good agricultural practices to manage the emerging pollution problem along the Kenyan Coast. A monitoring program for these pollutants is also important so as to always have an early warning.

Acknowledgements

Funding for this work was provided by Kenya Marine and Fisheries Research Institute, KMFRI) and RAF 7008, 7009, 7015 Project and CRP-K41016 projects jointly funded by International Atomic Energy Agency, IAEA and Government of Kenya. We are greatly indebted to the Directors of these Institutions for supporting this work. We also appreciate the efforts of KMFRI staff that assisted in field samples collection and analysis in one way or the other.

Author details

Eric Ochieng Okuku^{*}, Kiteresi Linet Imbayi, Owato Gilbert Omondi, Wanjeri Veronica Ogolla Wayayi, Mwalugha Catherine Sezi, Kombo Mokeira Maureen, Stephen Mwangi and Nancy Oduor Kenya Marine and Fisheries Research Institute, Mombasa, Kenya

*Address all correspondence to: ochiengokuku2003@yahoo.com

IntechOpen

© 2019 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

References

[1] Okuku EO, Mubiana VK, Hagos KG, Peter HK, Blust R. Bioavailability of sediment-bound heavy metals on the East African Coast. West Indian Ocean Journal of Marine Science. 2010;**9**(1):31-42

[2] UNEP-GEF. Project Document for the UNEP-GEF WIO-LaB Project entitled 'Addressing Land-based Activities in the Western Indian Ocean. 2004

[3] Daoji L, Daler D. Ocean pollution from land-based sources: East China Sea, China. AMBIO: A Journal of the Human Environment. 2004;**33**(1):107-113. DOI: 10.1579/0044-7447-33.1.107

[4] Ohowa B, Munga D, Kiteresi LI, Wanjeri VO, Okumu S, Kilonzo J, et al. Sewage pollution in the Coastal waters of Mombasa City, Kenya: A norm rather than an exception. Int J Environ Res. 2011;(4)

[5] Centre UWCM, Group C of ML on S (Programme). DAW. Seamounts, deepsea corals and fisheries: Vulnerability of deep-sea corals to fishing on seamounts beyond areas of national jurisdiction. UNEP/Earthprint; 2006

[6] Redfield AC. The biological control of chemical factors in the environment. American Scientist. 1958;**46**(3):230A-2221A

[7] van der Struijk LF, Kroeze C. Future trends in nutrient export to the coastal waters of South America: Implications for occurrence of eutrophication. Global Biogeochem Cycles. 2010;**24**(4). DOI: 10.1029/2009GB003572

[8] Diaz RJ, Rosenberg R. Spreading dead zones and consequences for marine ecosystems. Science (80-). 2008;**321**(5891):926-929

[9] Selman M, Greenhalgh S, Diaz R, Sugg Z. Eutrophication and hypoxia in coastal areas: A global assessment of the state of knowledge. World Resour Inst. 2008;**284**:1-6

[10] Howarth RW. In: Visgilio GR, Whitelaw DM, editors. Atmospheric Deposition and Nitrogen Pollution in Coastal Marine Ecosystems BT-Acid in the Environment: Lessons Learned and Future Prospects. Boston, MA: Springer US; 2007. pp. 97-116. DOI: 10.1007/978-0-387-37562-5_6

[11] SAB EPA. Reactive Nitrogen in the United States: An Analysis of Inputs, Flows, Consequences, and Management Options. EPA-SAB-11-013.
Washington, DC: Science Advisory Board, US Environmental Protection Agency; 2011. Available from: http:// yosemite.epa.gov/sab/sabproduct.nsf/ WebBOARD/INCSupplemental

[12] Burkholder JM, Mason KM, Glasgow HB Jr. Water-column nitrate enrichment promotes decline of eelgrass Zostera marina: Evidence from seasonal mesocosm experiments. Marine Ecology Progress Series. 1992;**81**(2):163-178

[13] Gappa JJL, Tablado NHM. Influence of sewage pollution on a rocky intertidal community dominated by the mytilid Brachidontes. Marine Ecology Progress Series. 1990;**63**:163-175

[14] Howarth RW, Marino R, Swaney DP, Boyer EW. Wastewater and watershed influences on primary productivity and oxygen dynamics in the lower Hudson River estuary. The Hudson River Estuary. 2006;**136**

[15] Okuku EO. Bioavailability of Sediment-Bound Heavy Metals along the Coast of Eastern Africa and Evaluation of Metal Concentrations along the Trophic Hierarchy. 2007

[16] Alonso-Hernandez CM, Mesa-Albernas M, Tolosa I. Organochlorine pesticides (OCPs) and polychlorinated biphenyls (PCBs) in sediments from the Gulf of Batabanó, Cuba. Chemosphere. 2014;**94**:36-41. Available from: http:// www.sciencedirect.com/science/article/ pii/S0045653513012368

[17] Barakat AO, Mostafa A, Wade TL,
Sweet ST, El Sayed NB. Distribution and ecological risk of organochlorine pesticides and polychlorinated biphenyls in sediments from the Mediterranean coastal environment of Egypt. Chemosphere. 2013;93(3):
545-554. Available from: http://www. sciencedirect.com/science/article/pii/ S0045653513009065

[18] Kilunga PI, Sivalingam P, Laffite A, Grandjean D, Mulaji CK, de Alencastro LF, et al. Accumulation of toxic metals and organic micro-pollutants in sediments from tropical urban rivers, Kinshasa, Democratic Republic of the Congo. Chemosphere. 2017;**179**:37-48. Available from: http://www. sciencedirect.com/science/article/pii/ S0045653517304538

[19] Miglioranza KSB, Gonzalez M, Ondarza PM, Shimabukuro VM, Isla FI, Fillmann G, et al. Assessment of Argentinean Patagonia pollution: PBDEs, OCPs and PCBs in different matrices from the Río Negro basin. Sci Total Environ. 2013;**452-453**:275-285. Available from: http://www. sciencedirect.com/science/article/pii/ S0048969713002386

[20] Barasa MW, Lalah JO, Wandiga SO. Seasonal variability of persistent organochlorine pesticide residues in marine fish along the Indian Ocean coast of Kenya. Toxicol Environ Chem. 2008;**90**(3):535-547. DOI: 10.1080/02772240701556312

[21] Sun H, Qi Y, Zhang D, Li QX, Wang J. Concentrations, distribution, sources and risk assessment of organohalogenated contaminants in soils from Kenya, Eastern Africa. Environ Pollut. 2016;**209**:177-185. Available from: http://www. sciencedirect.com/science/article/pii/ S0269749115302001

[22] Skwarzec B, Jahnz A. The inflow of polonium ²¹⁰Po from Vistula river catchments area. J Environ Sci Heal -Part A Toxic/Hazardous Subst Environ Eng. 2007;**42**(14):2117-2122

[23] Thi Van T, Duc Tam N, Duc Nhan D, Quang Long N, Van Thang D, Duy Cam B. Water-sediment distribution and behaviour of Polonium (²¹⁰Po) in a shallow coastal area with high concentration of dissolved organic matters in water, North Vietnam. Nucl Sci Technol. 2016;**6**(2):1-14

[24] Matthews KM, Kim C-K, Martin P. Determination of ²¹⁰Po in environmental materials: A review of analytical methodology. Appl Radiat Isot. 2007;**65**(3):267-279. Available from: http://www.sciencedirect.com/science/ article/pii/S0969804306003290

[25] Hains A. Lead Poisoning: Triggers and Thresholds. Buffalo: NY Partnersh Public Good; 2017

[26] Arrebola JP, Cuellar M, Claure E, Quevedo M, Antelo SR, Mutch E, et al. Concentrations of organochlorine pesticides and polychlorinated biphenyls in human serum and adipose tissue from Bolivia. Environ Res. 2012;**112**:40-47. Available from: http:// www.sciencedirect.com/science/article/ pii/S0013935111002507

[27] Commendatore MG, Franco MA, Gomes Costa P, Castro IB, Fillmann G, Bigatti G, et al. Butyltins, polyaromatic hydrocarbons, organochlorine pesticides, and polychlorinated biphenyls in sediments and bivalve mollusks in a mid-latitude environment from the Patagonian coastal zone. Environmental Toxicology and Chemistry. 2015;**34**(12):2750-2763. DOI: 10.1002/etc.3134

[28] Lu M, Zeng D-C, Liao Y, Tong B. Distribution and characterization of organochlorine pesticides and polycyclic aromatic hydrocarbons in surface sediment from Poyang Lake, China. Sci Total Environ. 2012;**433**:491-497. Available from: http://www. sciencedirect.com/science/article/pii/ S0048969712009412

[29] Kiteresi LI, Okuku EO, Mwangi SN, Ohowa B, Wanjeri VO, Okumu S, et al. The influence of land based activities on the phytoplankton communities of Shimoni-Vanga system, Kenya. International Journal of Environmental Research. 2012;**6**(1):151-162

[30] Nyunja JA, Mavuti KM, Wakwabi EO. Trophic ecology of *Sardinella gibbosa* (Pisces: Clupeidae) and *Atherinomorous lacunosus* (Pisces: Atherinidae) in Mtwapa Creek and Wasini Channel, Kenya. Western Indian Ocean Journal of Marine Science. 2002;1(2):181-189. Available at: http:// www.oceandocs.net/handle/1834/40

[31] Meyler S, Felix H, Crouthers R. Abundance and distribution of Indo-Pacific humpback dolphins (*Sousa chinensis*) in the Shimoni Archipelago, Kenya. Western Indian Ocean Journal of Marine Science. 2011;**10**(2):201-209. Available at: http://41.215.122.106/ dspace/bitstream/0/5262/1/Meyler2012. pdf

[32] Mwashote BM. Cadmium and lead levels in Mombasa, Kenya levels of cadmium and lead in water, sediments and selected fish species in Mombasa, Kenya. Western Indian Ocean Journal of Marine Science. 2003;**2**(1):25-34

[33] Hoorweg JC, Versleijen N, Wangila B, Degen A. Income diversification and fishing practices among artisanal fishers on the Malindi-Kilifi coast. 2006

[34] Okuku EO, Bouillon S, Ochiewo JO, Munyi F, Kiteresi LI, Tole M. The impacts of hydropower development on rural livelihood sustenance. International Journal of Water Resources Development. 2016;**32**(2):267-285

[35] Wu Y, Wang X, Ya M, Li Y, Hong H. Distributions of organochlorine compounds in sediments from Jiulong River Estuary and adjacent Western Taiwan Strait: Implications of transport, sources and inventories. Environ Pollut. 2016;**219**:519-527. Available from: http:// www.sciencedirect.com/science/article/ pii/S0269749116304699

[36] Parsons TR. A Manual of Chemical & Biological Methods for Seawater Analysis. Elsevier; 2013

[37] APHA. Standard Methods for the Examination of Water and Wastewater. 9th ed. Washington, DC: American Public Health Association, American Water Works Association and Water Environmental Federation; 1998

[38] Benedik L, Vrecek P. Determination of Pd-210 and Po-210 in environmental samples. Acta Chimica Slovenica. 2001;**48**(2):199-213

[39] Thompson S, Budzinski H, LeMenach K, Letellier M, Garrigues P. Multi-residue analysis of polycyclic aromatic hydrocarbons, polychlorobiphenyls, and organochlorine pesticides in marine sediments. Analytical and Bioanalytical Chemistry. 2002;**372**(1):196-204. DOI: 10.1007/s00216-001-1170-1

[40] Mohammed A, Peterman P, Echols K, Feltz K, Tegerdine G, Manoo A, et al. Polychlorinated biphenyls (PCBs) and organochlorine pesticides (OCPs) in harbor sediments from Sea Lots, Portof-Spain, Trinidad and Tobago. Marine Pollution Bulletin. 2011;**62**(6): 1324-1332. Available from: http://www. sciencedirect.com/science/article/pii/ S0025326X1100186X

[41] Wu Q, Leung JYS, Yuan X, Huang X, Li H, Huang Z, et al. Biological

risk, source and pollution history of organochlorine pesticides (OCPs) in the sediment in Nansha mangrove, South China. Marine Pollution Bulletin. 2015;**96**(1):57-64. Available from: http:// www.sciencedirect.com/science/article/ pii/S0025326X15003276

[42] Ali U, Li J, Zhang G, Mahmood A, Jones KC, Malik RN. Presence, deposition flux and mass burden of persistent organic pollutants (POPs) from Mehmood Booti Drain sediments, Lahore. Ecotoxicology and Environmental Safety. 2016;**125**:9-15. Available from: http://www. sciencedirect.com/science/article/pii/ S0147651315301664

[43] MacDonald DD, IngersollCG, Smorong DE, LindskoogR. Development and Evaluationof Numerical Sediment QualityAssessment Guidelines for FloridaInland Waters. 2003

[44] MacDonald DD, Ingersoll CG. A guidance manual to support the assessment of contaminated sediments in freshwater ecosystems. US Environmental Protection Agency, Great Lakes National Program Office; 2002

[45] Rosenberg DM, McCully P, Pringle CM. Global-scale environmental effects of hydrological alterations: introduction. AIBS Bulletin.2000;50(9):746-751

[46] Bala Krishna Prasad M. Nutrient stoichiometry and eutrophication in Indian mangroves. Environmental Earth Sciences. 2012;**67**(1):293-299. DOI: 10.1007/s12665-011-1508-8

[47] Zhang W, Feng H, Chang J, Qu J, Xie H, Yu L. Heavy metal contamination in surface sediments of Yangtze River intertidal zone: An assessment from different indexes. Environmental Pollution. 2009;**157**(5):1533-1543. Available from: http://www. sciencedirect.com/science/article/pii/ S0269749109000219 [48] Kamau JN, Kuschk P, Machiwa J, Macia A, Mothes S, Mwangi S, et al. Investigating the distribution and fate of Al, Cd, Cr, Cu, Mn, Ni, Pb and Zn in sewage-impacted mangrovefringed creeks of Kenya, Tanzania and Mozambique. Journal of Soils and Sediments. 2015;**15**(12):2453-2465. DOI: 10.1007/s11368-015-1214-3

[49] Mager EM, Esbaugh AJ, Brix KV, Ryan AC, Grosell M. Influences of water chemistry on the acute toxicity of lead to *Pimephales promelas* and *Ceriodaphnia dubia*. Comparative Biochemistry and Physiology, Part C: Toxicology & Pharmacology. 2011;**153**(1):82-90

[50] Everaarts JM, Swennen C, Cheewasedtham W. Heavy metals (Cu, Zn, Cd, Pb) in surface sediment and organisms in a short food chain from the intertidal zone of Pattani Bay, Thailand–A preliminary study. Wallaceana. 1994;**72**:17-24

[51] Obhođaš J, Valković V. Contamination of the coastal sea sediments by heavy metals. Applied Radiation and Isotopes. 2010;**68**(4): 807-811. Available from: http://www. sciencedirect.com/science/article/pii/ S0969804309008100

[52] Davies TC, Osano O. Sustainable mineral development: Case study from Kenya. Geological Society of London, Special Publication. 2005;**250**(1):87-93. Available from: http://sp.lyellcollection. org/content/250/1/87.abstract

[53] Johnson UK. River sediment supply, sedimentation and transport of the highly turbid sediment plume in Malindi Bay, Kenya. Journal of Geographical Sciences. 2013;**23**(3): 465-489. DOI: 10.1007/s11442-013-1022-x

[54] Jia G, Belli M, Sansone U, Rosamilia S, Blasi M. ²¹⁰Pb and ²¹⁰Po concentrations in the Venice lagoon ecosystem (Italy) and the potential radiological impact to the local

public and environment. Journal of Radioanalytical and Nuclear Chemistry. 2003;**256**(3):513-528

[55] Mahmood ZUW, Mohamed CAR, Ishak AK, Bakar NSA, Ishak K. Vertical distribution of ²¹⁰Pb and ²²⁶Ra and their activity ratio in marine sediment core of the East Malaysia coastal waters. Journal of Radioanalytical and Nuclear Chemistry. 2011;**289**(3):953-959. Available from: http://link.springer. com/10.1007/s10967-011-1206-8

[56] Venunathan N, Narayana Y. Activity of ²¹⁰Po and ²¹⁰Pb in the riverine environs of coastal Kerala on the southwest coast of India. Journal of Radiation Research and Applied Science. 2016:392-399

[57] Aközcan S, Uğur Görgün A. Trace metal and radionuclide pollution in marine sediments of the Aegean Sea (Izmir Bay and Didim). Environment and Earth Science. 1 Aug 2013;**69**(7):2351-2355

[58] Chen Y, Jia R, Yang S. Distribution and source of polycyclic aromatic hydrocarbons (PAHs) in water dissolved phase, suspended particulate matter and sediment from Weihe river in Northwest China. International Journal of Environmental Research and Public Health. 2015;**12**

[59] Saleem M, Ali Khan A, Ahmed K, Qamaruddin M, Kahkashan S, Imran Hasany S, et al. Polycyclic Aromatic Hydrocarbons (PAHs) in Surface Sediments from the Karachi Harbor and off-Clifton Coast, Karachi: Spatial Distribution, Composition and Ecological Risk Assessment. Vol. 112017. pp. 2222-3045

[60] Chen C-W, Chen C-F. Distribution, origin, and potential toxicological significance of polycyclic aromatic hydrocarbons (PAHs) in sediments of Kaohsiung Harbor, Taiwan. Marine Pollution Bulletin. 2011;**63**(5):417-423. Available from: http://www. sciencedirect.com/science/article/pii/ S0025326X11002505

[61] Lalah JO, Yugi PO, Jumba IO,
Wandiga SO. Organochlorine pesticide residues in Tana and Sabaki Rivers in Kenya. Bulletin of Environmental Contamination and Toxicology.
2003;71(2):298-307

[62] Cheng H, Lin T, Zhang G, Liu G, Zhang W, Qi S, et al. DDTs and HCHs in sediment cores from the Tibetan Plateau. Chemosphere. 2014;**94**:183-189. Available from: http://www. sciencedirect.com/science/article/pii/ S0045653513013982

[63] Aly Salem DMS, Khaled A, El
Nemr A. Assessment of pesticides
and polychlorinated biphenyls
(PCBs) in sediments of the Egyptian
Mediterranean Coast. Egyptian Journal
of Aquatic Research. 2013;39(3):
141-152. Available from: http://www.
sciencedirect.com/science/article/pii/
S1687428513001131

[64] Siokou-Frangou I, Pagou K. Assessment of the trophic conditions and ecological status in the Inner Saronikos Gulf. InTechnical Report for the Ministry of Environment, Planning and Public Works. NCMR: Athens. Mar 2000. p. 43

[65] Benson NU, Anake WU, Essien JP, Enyong P, Olajire AA. Distribution and risk assessment of trace metals in Leptodius exarata, surface water and sediments from Douglas Creek in the Qua Iboe Estuary. Journal of Taibah University For Science. 1 May 2017;**11**(3):434-439