

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Cyclotomic and Littlewood Polynomials Associated to Algebras

José-Antonio de la Peña

Abstract

Let A be a finite dimensional algebra over an algebraically closed field k . Assume A is a basic connected and triangular algebra with n pairwise non-isomorphic simple modules. We consider the Coxeter transformation $\phi_A(T)$ as the automorphism of the Grothendieck group $K_0(A)$ induced by the Auslander-Reiten translation τ in the derived category $D^b(\text{mod}_A)$ of the module category mod_A of finite dimensional left A -modules. In this paper we study the Mahler measure $\mathbb{M}(\chi_A)$ of the Coxeter polynomial χ_A of certain algebras A . We consider in more detail two cases: (a) A is said to be *cyclotomic* if all eigenvalues of χ_A are roots of unity; (b) A is said to be of *Littlewood type* if all coefficients of χ_A are $-1, 0$ or 1 . We find criteria in order that A is of one of those types. In particular, we establish new records according to Mossingshoff's list of *Record Mahler measures* of polynomials q with $1 < \mathbb{M}(q)$ as small as possible, ordered by their number of roots outside the unit circle.

Keywords: finite dimensional algebra, coxeter transformation, derived category, accessible algebra, characteristic polynomial, cyclotomic polynomial, littlewood type

1. Introduction


Assume throughout the paper that K is an algebraically closed field. We assume that A is a triangular finite dimensional basic K -algebra, that is, of the form $A = KQ/I$, where I is an ideal of the path algebra KQ for Q a quiver without oriented cycles. In particular, A has finite global dimension. The Coxeter transformation ϕ_A is the automorphism of the Grothendieck group $K_0(A)$ induced by the Auslander-Reiten translation τ in the derived category $D^b(A)$ see [1]. The characteristic polynomial χ_A of ϕ_A is called the *Coxeter polynomial* χ_A of A , or simply χ_A see [15, 17]. It is a monic self-reciprocal polynomial, therefore it is $\chi_A = a_0 + a_1T + a_2T^2 + \dots + a_{n-2}T^{n-2} + a_{n-1}T^{n-1} + a_nT^n \in \mathbb{Z}[T]$, with $a_i = a_{n-i}$ for $0 \leq i \leq n$, and $a_0 = 1 = a_n$.

Consider the roots $\lambda_1, \dots, \lambda_n$ of χ_A , the so called *spectrum* of A . There is a number of measures associated to the absolute values $|\lambda|$ for λ in the spectrum $\text{Spec}(\phi_A)$ of A . For instance, the *spectral radius* of A is defined as $\rho_A = \max\{|\lambda| : \lambda \in \text{Spec}(\phi_A)\}$ and the *Mahler measure* of χ_A defined as $\mathbb{M}(\chi_A) = \max\left\{1, \prod_{|\lambda| > 1} |\lambda|\right\}$. Recently, some explorations on the relations of the Mahler measure $\mathbb{M}(\chi_A)$ and properties of the algebra A have been initiated.

For a one-point extension $A = B[N]$, we show that $\mathbb{M}(\chi_B) \leq \mathbb{M}(\chi_A)$. The strongest statements and examples will be given for the class of accessible algebras. We say

that an algebra A is *accessible from* B if there is a sequence $B = B_1, B_2, \dots, B_s = A$ of algebras such that each B_{i+1} is a one-point extension (resp. coextension) of B_i for some exceptional B_i -module M_i . As a special case, a K -algebra A is called *accessible* if A is accessible from the one vertex algebra K .

We say that A is of *cyclotomic type* if the eigenvalues of ϕ_A lie on the unit circle. Many important finite dimensional algebras are known to be of cyclotomic type: hereditary algebras of finite or tame representation type, canonical algebras, some extended canonical algebras and many others. On the other hand, there are well-known classes of algebras with a mixed behavior with respect to cyclotomicity. For instance, in Section 6 below we consider the class of *Nakayama algebras*. Let $N(n, r)$ be the quotient obtained from the linear quiver with n vertices



with relations $x^r = 0$. The Nakayama algebras $N(n, 2)$ are easily proven to be of cyclotomic type, while those of the form $N(n, 3)$ are of cyclotomic type as consequence of lengthly considerations in [18]. The case $r = 4$ is more representative: $N(n, 4)$ is of cyclotomic type for all $0 \leq n \leq 100$ except for $n = 10, 22, 30, 42, 50, 62, 70, 82$ and 90 . Clearly, if A is of cyclotomic type then $|\text{Tr}(\phi_A)^k| \leq n$, for $k \geq 0$. We show the following theorem.

Theorem 1: *Let M be an unimodular $n \times n$ -matrix. The following are equivalent:*

- a. M is of cyclotomic type;
- b. for every positive integer $0 \leq k \leq n$, we have $|\text{Tr}(M^k)| \leq n$.

We also consider algebras A of *Littlewood type* where χ_A has all its coefficients in the set $\{-1, 0, 1\}$. Among other structure results, we prove.

Proposition. *The closure \bar{P} of the set P of roots of Littlewood polynomials, equals the set R of roots of Littlewood series.*

Our results make use of well established techniques in the *representation theory of algebras*, as well as results from the *theory of polynomials* and *transcendental number theory*, where Mahler measure has its usual habitat. We stress here the natural context of these investigations on the largely unexplored overlapping area of these important subjects. Hence, rather than a comprehensive study we understand our work as a preliminary exploration where examples are most valuable.

2. Measures for polynomials

2.1 Self-reciprocal polynomials

A polynomial $p(z)$ of degree n is said to be self-reciprocal if $p(z) = z^n p(1/z)$. The following table displays the number $a(n)$ of polynomials p of degree n (for small n) with $p(0)$ non-zero, $b(n)$ is the number of such polynomials which are additionally self-reciprocal, and $c(n)$ is the number of those which are self-reciprocal and where $p(-1)$ is the square of an integer.

n	1	2	3	4	5	6	7	8	9	10	11	12	15	20	25
$a(n)$	2	6	10	24	38	78	118	224	330	584	838	1420	4514	30,532	152,170
$b(n)$	1	5	5	19	19	59	59	165	165	419	419	1001	2257	20,399	76,085
$c(n)$	1	3	5	12	19	34	59	99	165	244	419	598	2257	12,526	76,085

Indeed, there is an efficient algorithm to determine such polynomials of given degree n , based on a quadratic bound for $n \leq 4f(n)^2$ in terms of Euler totient function, $f(n)$.

Cyclotomic polynomials Φ_n and their products are a natural source for self-reciprocal polynomials. Clearly, $\Phi_1(z) = z - 1$ is not self-reciprocal, but all remaining Φ_n (with $n \geq 2$) are. Hence, exactly the polynomials $(z - 1)^{2k} \prod_{n \geq 2} \Phi_n^{e_n}$ with natural numbers k and e_n are self-reciprocal with spectral radio one and without eigenvalue zero.

It is not a coincidence that in the above tables we have $b(n) = c(n + 1)$ for n even and $b(n) = c(n)$ for n odd. Indeed, if p is self-reciprocal of odd degree then $p(-1) = 0$, hence $p(z) = (z + 1)q(z)$ where q is also self-reciprocal.

2.2 Mahler measure

Let A be a finite dimensional K -algebra with finite global dimension. The Grothendieck group $K_0(A)$ of the category mod_A of finite dimensional (right) A -modules, formed with respect to short exact sequences, is naturally isomorphic to the Grothendieck group of the derived category, formed with respect to exact triangles.

The Coxeter transformation ϕ_A is the automorphism of the Grothendieck group $K_0(A)$ induced by the Auslander-Reiten translation τ . The characteristic polynomial $\chi_A(T)$ of ϕ_A is called the Coxeter polynomial $\chi_A(T)$ of A , or simply χ_A . It is a monic self-reciprocal polynomial, therefore it is $\chi_A(T) = a_0 + a_1T + a_2T^2 + \dots + a_{n-2}T^{n-2} + a_{n-1}T^{n-1} + a_nT^n \in \mathbb{Z}[T]$, with $a_i = a_{n-i}$ for $0 \leq i \leq n$, and $a_0 = 1 = a_n$.

Consider the roots $\lambda_1(A), \dots, \lambda_n(A)$ of χ_A , the so called *spectrum* of A . In [15], a measure for polynomials was introduced. Namely, the *Mahler measure* of χ_A is $\mathbb{M}(\chi_A) = \max\{1, \prod_{i=1}^n |\lambda_i|\}$. By a celebrated result of Kronecker [9], see also [7, Prop. 1.2.1], a monic integral polynomial p , with $p(0) \neq 0$, has $\mathbb{M}(p) = 1$ if and only if p factorizes as product of cyclotomic polynomials. As observed in [18], A is of *cyclotomic type* if and only if $\mathbb{M}(\chi_A) = 1$, that is, $\chi_A(T)$ factorizes as product of cyclotomic polynomials.

2.3 Spectral radius one, periodicity

If the spectrum of A lies in the unit disk, then all roots of χ_A lie on the unit circle, hence A has spectral radius $\rho_A = 1$. Clearly, for fixed degree there are only finitely many monic integral polynomials with this property.

The following finite dimensional algebras are known to produce Coxeter polynomials of spectral radius one:

1. hereditary algebras of finite or tame representation type;
2. all canonical algebras;
3. (some) extended canonical algebras;
4. generalizing (2), (some) algebras which are derived equivalent to categories of coherent sheaves.

We put $v_n = 1 + x + x^2 + \dots + x^{n-1}$. Note that v_n has degree $n - 1$. There are several reasons for this choice: first of all $v_n(1) = n$, second this normalization yields convincing formulas for the Coxeter polynomials of canonical algebras and

hereditary stars, third representing a Coxeter polynomial — for spectral radius one — as a rational function in the v_n ’s relates to a Poincaré series, naturally attached to the setting.

Dynkin type	Star symbol	v -factorization	Cyclotomic factorization	Coxeter number
A_n	$[n]$	v_{n+1}	$\prod_{d n, d>1} \Phi_d$	$n+1$
D_n	$[2, 2, n-2]$	$\frac{v_2(v_2v_{n-2})}{(v_2v_{n-2})v_{n-1}} v_{2(n-1)}$	$\Phi_2 \prod_{\substack{d 2(n-1) \\ d \neq 1, d \neq n-1}} \Phi_d$	$2(n-1)$
E_6	$[2, 3, 3]$	$\frac{v_2v_3(v_3)}{(v_3)v_4v_6} v_{12}$	$\Phi_3\Phi_{12}$	12
E_7	$[2, 3, 4]$	$\frac{v_2v_3(v_4)}{(v_4)v_6v_9} v_{18}$	$\Phi_2\Phi_{18}$	18
E_8	$[2, 3, 5]$	$\frac{v_2v_3v_5}{v_6v_{10}v_{15}} v_{30}$	Φ_{30}	30

In the column ‘ v -factorization’, we have added some extra terms in the nominator and denominator which obviously cancel.

Inspection of the table shows the following result:

Proposition. *Let k be an algebraically closed field and A be a connected, hereditary k -algebra which is representation-finite. Then the Coxeter polynomial χ_A determines A up to derived equivalence.* □

2.4 Triangular algebras

Nearly all algebras considered in this survey are triangular. By definition, a finite dimensional algebra is called *triangular* if it has triangular matrix shape

$$\begin{bmatrix} A_1 & M_{12} & \cdots & M_{1n} \\ 0 & A_2 & \cdots & M_{2n} \\ & & \ddots & \vdots \\ 0 & 0 & \cdots & A_n \end{bmatrix}$$

where the diagonal entries A_i are skew-fields and the off-diagonal entries M_{ij} , $j > i$, are A_i , A_j -bimodules. Each triangular algebra has finite global dimension.

Proposition. *Let A be a triangular algebra over an algebraically closed field K . Then $\chi_A(-1)$ is the square of an integer.*

Proof. Let C be the Cartan matrix of A with respect to the basis of indecomposable projectives. Since A is triangular and K is algebraically closed, we get $\det C = 1$, yielding

$$\chi_A = |xI + C^{-1}C^t| = |C^{-1}| \cdot |xC + C^t| = |C^t + xC|.$$

Hence $\chi_A(-1)$ is the determinant of the skew-symmetric matrix $S = C^t - C$. Using the skew-normal form of S , see [16, Theorem IV.1], we obtain $S' = U^t S U$ for some $U \in GL_n(\mathbb{Z})$, where S' is a block-diagonal matrix whose first block is the zero matrix of a certain size and where the remaining blocks have the shape $\begin{bmatrix} 0 & m_i \\ -m_i & 0 \end{bmatrix}$ with integers m_i . The claim follows. □

Which self-reciprocal polynomials of spectral radius one are Coxeter polynomials? The answer is not known. If arbitrary base fields are allowed, we conjecture that all self-reciprocal polynomials are realizable as Coxeter polynomials of triangular

algebras. Restricting to algebraically closed fields, already the request that $\chi_A(-1)$ is a square discards many self-reciprocal polynomials, for instance the cyclotomic polynomials $\Phi_4, \Phi_6, \Phi_8, \Phi_{10}$. Moreover, the polynomial $f = x^3 + 1$, which is the Coxeter polynomial of the non simply-laced Dynkin diagram \mathbb{B}_3 , does not appear as the Coxeter polynomial of a triangular algebra over an algebraically closed field, despite of the fact that $f(-1) = 0$ is a square. Indeed, the Cartan matrix

$$\begin{bmatrix} 1 & a & b \\ 0 & 1 & c \\ 0 & 0 & 1 \end{bmatrix}$$

yields the Coxeter polynomial $f = x^3 + \alpha x^2 + \alpha x + 1$, where $\alpha = abc - a^2 - b^2 - c^2 + 3$. The equation $a^2 + b^2 + c^2 - abc = 3$ of Hurwitz-Markov type does not have an integral solution. (Use that reduction modulo 3 only yields the trivial solution in \mathbb{F}_3 .)

2.5 Relationship with graph theory

Given a (non-oriented) graph Δ , its *characteristic polynomial* κ_Δ is defined as the characteristic polynomial of the adjacency matrix M_Δ of Δ . Observe that, since M_Δ is symmetric, all its eigenvalues are real numbers. For general results on graph theory and spectra of graphs see [4].

There are important interactions between the theory of graph spectra and the representation theory of algebras, due to the fact that if C is the Cartan matrix of $A = K[\vec{\Delta}]$, then M_Δ is determined by the symmetrization $C + C^t$ of C , since $M_\Delta = C + C^t - 2I$. We shall see that information on the spectra of M_Δ provides fundamental insights into the spectral analysis of the Coxeter matrix Φ_A and the structure of the algebra A .

A fundamental fact for a hereditary algebra $A = K[\vec{\Delta}]$, when $\vec{\Delta}$ is a *bipartite quiver*, that is, every vertex is a sink or source, is that $\text{Spec}(\Phi_A) \subset \mathbb{S}^1 \cup \mathbb{R}^+$. This was shown as a consequence of the following important identity.

Proposition. [2] *Let $A = K[\vec{\Delta}]$ be a hereditary algebra with $\vec{\Delta}$ a bipartite quiver without oriented cycles. Then $\chi_A(x^2) = x^n \kappa_\Delta(x + x^{-1})$, where n is the number of vertices of $\vec{\Delta}$ and κ_Δ is the characteristic polynomial of the underlying graph Δ of $\vec{\Delta}$.*

Proof. Since $\vec{\Delta}$ is bipartite, we may assume that the first m vertices are sources and the last $n - m$ vertices are sinks. Then the adjacency matrix A of Δ and the Cartan matrix C of A , in the basis of simple modules, take the form: $A = N + N^t$, $C = I_n - N$, where

$$N = \begin{pmatrix} 0 & D \\ 0 & 0 \end{pmatrix}$$

for certain $m \times m$ -matrix D . Since $N^2 = 0$, then $C^{-1} = I_n + N$. Therefore

$$\begin{aligned} \det(x^2 I_n - \Phi_A) &= \det(x^2 I_n + (I_n - N)(I_n + N)^t) \det(I_n - N^t) \\ &= \det(x^2 I_n - x^2 N^t + (I_n - N)) \\ &= x^n \det((x + x^{-1}) I_n - x N^t - x^{-1} N) \\ &= x^n \det((x + x^{-1}) I_n - A). \end{aligned}$$

□

The above result is important since it makes the spectral analysis of bipartite quivers and their underlying graphs almost equivalent. Note, however, that the representation theoretic context is much richer, given the categorical context behind the spectral analysis of quivers. The representation theory of bipartite quivers may thus be seen as a categorification of the class of graphs, allowing a bipartite structure.

Constructions in graph theory. Several simple constructions in graph theory provide tools to obtain in practice the characteristic polynomial of a graph. We recall two of them (see [4] for related results):

- a. Assume that a is a vertex in the graph Δ with a unique neighbor b and Δ' (resp. Δ'') is the full subgraph of Δ with vertices $\Delta_0 \setminus \{a\}$ (resp. $\Delta_0 \setminus \{a, b\}$), then

$$\kappa_{\Delta} = x\kappa_{\Delta'} - \kappa_{\Delta''}$$

- b. Let Δ_i be the graph obtained by deleting the vertex i in Δ . Then the first derivative of κ_{Δ} is given by

$$\kappa'_{\Delta} = \sum_i \kappa_{\Delta_i}$$

The above formulas can be used inductively to calculate the characteristic polynomial of trees and other graphs. They immediately imply the following result that will be used often to calculate Coxeter polynomials of algebras.

Proposition. Let $A = K[\vec{\Delta}]$ be a bipartite hereditary algebra. The following holds:

- i. Let a be a vertex in the graph Δ with a unique neighbor b . Consider the algebras B and C obtained as quotients of A modulo the ideal generated by the vertices a and a, b , respectively. Then

$$\chi_A = (x + 1)\chi_B - x\chi_C$$

- ii. The first derivative of the Coxeter polynomial satisfies:

$$2x\chi'_A = n\chi_A + (x - 1) \sum_i \chi_{A^{(i)}}$$

where $A^{(i)} = K[\vec{\Delta} \setminus \{i\}]$ is an algebra obtained from A by ‘killing’ a vertex i .

Proof. Use the corresponding results for graphs and A’Campo’s formula for the algebras A and its quotients $A^{(i)}$. \square

3. Important classes of algebras

In this section we give the definitions and main properties of such classes of finite dimensional algebras where information on their spectral properties is available.

3.1 Hereditary algebras


Let A be a finite dimensional K -algebra. For simplicity we assume $A = K[\vec{\Delta}]/I$

for a quiver $\vec{\Delta}$ without oriented cycles and I an ideal of the path algebra. The following facts about the Coxeter transformation Φ_A of A are fundamental:

- i. Let S_1, \dots, S_n be a complete system of pairwise non-isomorphic simple A -modules, P_1, \dots, P_n the corresponding projective covers and I_1, \dots, I_n the injective envelopes. Then ϕ_A is the automorphism of $K_0(A)$ defined by $\Phi_A[P_i] = -[I_i]$, where $[X]$ denotes the class of a module X in $K_0(A)$.
- ii. For a hereditary algebra $A = K[\vec{\Delta}]$, the spectral radius $\rho_A = \rho_{\Phi_A}$ determines the representation type of A in the following manner:
 - a. A is representation-finite if $1 = \rho_A$ is not a root of the Coxeter polynomial χ_A .
 - b. A is tame if $1 = \rho_A \in \text{Roots}(\chi_A)$.
 - c. A is wild if $1 < \rho_A$. Moreover, if A is wild connected, Ringel [20] shows that the spectral radius ρ_A is a simple root of χ_A . Then Perron-Frobenius theory yields a vector $y^+ \in K_0(A) \otimes_{\mathbb{Z}} \mathbb{R}$ with positive coordinates such that $\Phi_A y^+ = \rho_A y^+$. Since χ_A is self reciprocal, there is a vector $y^- \in K_0(A) \otimes_{\mathbb{Z}} \mathbb{R}$ with positive coordinates such that $\Phi_A y^- = \rho_A^{-1} y^-$. The vectors y^+, y^- play an important role in the representation theory of $A = K[\vec{\Delta}]$, see [5, 17].

Explicit formulas, special values. We are discussing various instances where an explicit formula for the Coxeter polynomial is known.

star quivers. Let A be the path algebra of a hereditary star $[p_1, \dots, p_t]$ with respect to the standard orientation, see



Since the Coxeter polynomial χ_A does not depend on the orientation of A we will denote it by $\chi_{[p_1, \dots, p_t]}$. It follows from [11, prop. 9.1] or [2] that

$$\chi_{[p_1, \dots, p_t]} = \prod_{i=1}^t v_{p_i} \left((x+1) - x \sum_{i=1}^t \frac{v_{p_i-1}}{v_{p_i}} \right). \quad (1)$$

In particular, we have an explicit formula for the sum of coefficients of $\chi = \chi_{[p_1, \dots, p_t]}$ as follows:

$$\chi(1) = \prod_{i=1}^t p_i \left(2 - \sum_{i=1}^t \left(1 - \frac{1}{p_i} \right) \right). \quad (2)$$

This special value of χ has a specific mathematical meaning: up to the factor $\prod_{i=1}^t p_i$ this is just the orbifold-Euler characteristic of a weighted projective line \mathbb{X} of weight type (p_1, \dots, p_t) . Moreover,

1. $\chi(1) > 0$ if and only if the star $[p_1, \dots, p_t]$ is of Dynkin type, correspondingly the algebra A is representation-finite.

- 2. $\chi(1) = 0$ if and only if the star $[p_1, \dots, p_t]$ is of extended Dynkin type, correspondingly the algebra A is of tame (domestic) type.
- 3. $\chi(1) < 0$ if and only if $[p_1, \dots, p_t]$ is not Dynkin or extended Dynkin, correspondingly the algebra A is of wild representation type.

The above deals with all the Dynkin types and with the extended Dynkin diagrams of type $\tilde{\mathbb{D}}_n, n \geq 4$, and $\tilde{\mathbb{E}}_n, n = 6, 7, 8$. To complete the picture, we also consider the extended Dynkin quivers of type $\tilde{\mathbb{A}}_n (n \geq 2)$ restricting, of course, to quivers without oriented cycles. Here, the Coxeter polynomial depends on the orientation: If p (resp. q) denotes the number of arrows in clockwise (resp. anti-clockwise) orientation ($p, q \geq 1, p + q = n + 1$), that is, the quiver has type $\mathbb{A}(p, q)$, the Coxeter polynomial χ is given by

$$\chi_{(p,q)} = (x - 1)^2 v_p v_q. \tag{3}$$

Hence $\chi(1) = 0$, fitting into the above picture.
The next table displays the v -factorization of extended Dynkin quivers.

Extended Dynkin type	Star symbol	Weight symbol	Coxeter polynomial
$\tilde{\mathbb{A}}_{p,q}$	—	(p, q)	$(x - 1)^2 v_p v_q$
$\tilde{\mathbb{D}}_n, n \geq 4$	$[2, 2, n-2]$	$(2, 2, n - 2)$	$(x - 1)^2 v_2^2 v_{n-2}$
$\tilde{\mathbb{E}}_6$	$[3, 3, 3]$	$(2, 3, 3)$	$(x - 1)^2 v_2 v_3^2$
$\tilde{\mathbb{E}}_7$	$[2, 4, 4]$	$(2, 3, 4)$	$(x - 1)^2 v_2 v_3 v_4$
$\tilde{\mathbb{E}}_8$	$[2, 3, 6]$	$(2, 3, 5)$	$(x - 1)^2 v_2 v_3 v_5$

Remark: As is shown by the above table, proposition 2.3 extends to the tame hereditary case. That is, the Coxeter polynomial of a connected, tame hereditary K -algebra A (remember, K is algebraically closed) determines the algebra A up to derived equivalence. This is no longer true for wild hereditary algebras, not even for trees.

3.2 Canonical algebras

Canonical algebras were introduced by Ringel [19]. They form a key class to study important features of representation theory. In the form of tubular canonical algebras they provide the standard examples of tame algebras of linear growth. Up to tilting canonical algebras are characterized as the connected K -algebras with a separating exact subcategory or a separating tubular one-parameter family (see [12]). That is, the module category $\text{mod} - \Lambda$ accepts a *separating tubular family* $\mathcal{T} = (T_\lambda)_{\lambda \in P_1 K}$, where T_λ is a homogeneous tube for all λ with the exception of t tubes $T_{\lambda_1}, \dots, T_{\lambda_t}$ with T_{λ_i} of rank p_i ($1 \leq i \leq t$).

Canonical algebras constitute an instance, where the explicit form of the Coxeter polynomial is known, see [11] or [10].

Proposition. *Let Λ be a canonical algebra with weight and parameter data (p, λ) . Then the Coxeter polynomial of Λ is given by*

$$\chi_\Lambda = (x - 1)^2 \prod_{i=1}^t v_{p_i}. \tag{4} \square$$

The Coxeter polynomial therefore only depends on the weight sequence \mathbf{p} . Conversely, the Coxeter polynomial determines the weight sequence — up to ordering.

3.3 Incidence algebras of posets

Let X be a finite partially ordered set (poset). The incidence algebra KX is the K -algebra spanned by elements e_{xy} for the pairs $x \leq y$ in X , with multiplication defined by $e_{xy}e_{zw} = \delta_{yz}e_{xw}$. Finite dimensional right modules over KX can be identified with commutative diagrams of finite dimensional K -vector spaces over the Hasse diagram of X , which is the directed graph whose vertices are the points of X , with an arrow from x to y if $x < y$ and there is no $z \in X$ with $x < z < y$.

We recollect the basic facts on the Euler form of posets and refer the reader to [6] for details. The algebra KX is of finite global dimension, hence its Euler form is well-defined and non-degenerate. Denote by C_X , Φ_X the matrices of the bilinear form and the corresponding Coxeter transformation with respect to the basis of the simple KX -modules.

The incidence matrix of X , denoted 1_X , is the $X \times X$ matrix defined by $(1_X)_{xy} = 1$ if $x \leq y$ and otherwise $(1_X)_{xy} = 0$. By extending the partial order on X to a linear order, we can always arrange the elements of X such that the incidence matrix is uni-triangular. In particular, 1_X is invertible over \mathbb{Z} . Recall that the Möbius function $\mu_X : X \times X \rightarrow \mathbb{Z}$ is defined by $\mu_X(x, y) = (1_X)_{xy}^{-1}$.

Lemma. *a. $C_X = 1_X^{-1}$.*

b. Let $x, y \in X$. Then $(\Phi_X)_{xy} = -\sum_{x \leq z < y} \mu_X(y, z)$.

Proposition. *If X and Y are posets, then $C_{X \times Y} = C_X \otimes C_Y$ and $\Phi_{X \times Y} = -\Phi_X \otimes \Phi_Y$.*

4. Cyclotomic polynomials and polynomials of Littlewood type

4.1 Cyclotomic polynomials

We recall some facts about *cyclotomic polynomials*.

The n -cyclotomic polynomial $\Phi_n(T)$ is inductively defined by the formula

$$T^n - 1 = \prod_{d|n} \Phi_d(T). \quad (5)$$

The *Möbius function* is defined as follows:

$$\mu(n) = \begin{cases} 0 & \text{if } n \text{ is divisible by a square} \\ (-1)^r & \text{if } n = p_1 p_2 \dots p_r \text{ is a factorization into distinct primes.} \end{cases}$$

A more explicit expression for the cyclotomic polynomials is given by

$$\Phi_n(T) = \prod_{\substack{1 \leq d < n \\ d|n}} v_{n/d}(T)^{\mu(d)} \quad (6)$$


for $n \geq 2$, where $v_n = 1 + T + T^2 + \dots + T^{n-1}$.

4.2 Hereditary stars

A path algebra $K\Delta$ is said to be of *Dynkin type* if the underlying graph $|\Delta|$ of Δ is one of the *ADE-series*, that is, of type, \mathbb{A}_n , \mathbb{D}_n , for some $n \geq 1$ or \mathbb{E}_k , for $k = 6, 7, 8$.

There are various instances where an explicit formula for the Coxeter polynomial is known.

Let A be the path algebra of a hereditary star $[p_1, \dots, p_t]$ with respect to the standard orientation, see [13].



Since the Coxeter polynomial χ_A does not depend on the orientation of A we will denote it by $\chi_{[p_1, \dots, p_t]}$. It follows that


$$\chi_{[p_1, \dots, p_t]} = \prod_{i=1}^t v_{p_i} \left((T+1) - T \sum_{j=1}^t \frac{v_{p_j-1}}{v_{p_j}} \right).$$

In particular, we have an explicit formula for the sum of coefficients of $\chi_{[p_1, \dots, p_t]}$ as follows:

$$\sum_{i=0}^n a_i = \chi_{[p_1, \dots, p_t]}(1) = \prod_{i=1}^t p_i \left(2 - \sum_{i=1}^t \left(1 - \frac{1}{p_i} \right) \right).$$

4.3 Wild algebras

Let c be the real root of the polynomial $T^3 - T - 1$, approximately $c = 1.325$. As observed in [21], a wild hereditary algebra A associated to a graph Δ without multiple arrows has spectral radius $\rho_A > c$ unless Δ is one of the following graphs:



In these cases, for $m \geq 8$

$$c > \rho_{[2,4,5]} > \rho_{[2,3,m]} > \rho_{[2,3,7]} = \mu_0$$

where $\mu_0 = 1.176280\dots$ is the real root of the Coxeter polynomial

$$T^{10} + T^9 - T^7 - T^6 - T^5 - T^4 - T^3 + T + 1$$


associated to any hereditary algebra whose underlying graph is $[2, 3, 7]$. Observe that in these cases, the Mahler measure of the algebra equals the spectral radius.

4.4 Lehmer polynomial

In 1933, D. H. Lehmer found that the polynomial

$$T^{10} + T^9 - T^7 - T^6 - T^5 - T^4 - T^3 + T + 1$$

has Mahler measure $\mu_0 = 1.176280\dots$, and he asked if there exist any smaller values exceeding 1. In fact, the polynomial above is the Coxeter polynomial of the hereditary algebra whose underlying graph [2, 3, 7] is depicted below.



We say that a matrix M is of *Mahler type* (resp. *strictly Mahler type*) if either $\mathbb{M}(M) = 1$ or $\mathbb{M}(M) \geq \mu_0$ (resp. $\mathbb{M}(M) > \mu_0$). Earlier this year, Jean-Louis Verger-Gaugry announced a proof of Lehmer’s conjecture, see <https://arxiv.org/pdf/1709.03771.pdf>. The key result (Theorem 5.28, p. 122) is a Dobrowolski type minoration of the Mahler Measure $\mathbb{M}(\beta)$. Experts are still reading the arguments, but there is no conclusive opinion.

4.5 Happel’s trace formula

In [8], Happel shows that the trace of the Coxeter matrix can be expressed as follows:

$$-\mathrm{Tr}(\phi_A) = \sum_{k=0}^{\infty} (-1)^k \dim_K H^k(A) \tag{7}$$

where $H^k(A)$ denotes the k -th Hochschild cohomology group. In particular, if the Hochschild cohomology ring $H^*(A)$ is trivial, that is, $H^i(A) = 0$ for $i > 0$ and $H^0(A) = K$, then $\mathrm{Tr}(\phi_A) = -1$.

For an algebra A and a left A -module N we call

$$A[N] = \begin{bmatrix} A & 0 \\ N & K \end{bmatrix}$$

the *one-point extension* of A by N . This construction provides an order of vertices to deal with *triangular algebras*, that is, algebras KQ/I , where I is an ideal of the path algebra KQ for Q a quiver without oriented cycles.

4.6 One-point extensions

Let B be an algebra and M a B -module. Consider the one-point extension $A = B[N]$. In [19] it is shown the Coxeter transformations of A and B are related by

$$\phi_A = \begin{pmatrix} \phi_B & -C_B^T n^T \\ -n\phi_B & nC_B^T n^T - 1 \end{pmatrix} \tag{8}$$

where C_B is the *Cartan matrix* of B which satisfies $\phi_B = -C_B^{-T}C_B$ and n is the class of N in the Grothendieck group $K_0(B)$. In case $A = B[N]$ with N an *exceptional* module, it follows that

$$\mathrm{Tr}(\phi_A) = \mathrm{Tr}(\phi_B)$$

We recall that the *Euler quadratic form* is defined as $q_A(x) = xC_A^t x^t$. Assume that $A = B[M]$ for an algebra B and an indecomposable module M . In many cases, we get

that $q_A(m) > 0$, for m the dimension vector of M (for instance, if M is preprojective, or if q_A coincides with the Tits form of A ...)

Proposition. *Let A be an accessible algebra, such that $q_A(m) > 0$ for m the dimension vector of M , where $A = B[M]$ for certain algebra B and an indecomposable module M . Then the following happens:*

a. $\text{Tr}(\phi_A) \geq -1$;

b. if $\text{Tr}(\phi_B) = -1$ and $q_B(m) = 1$, then $\text{Tr}(\phi_A) = -1$.

Proof. Assume that $A = B[M]$ for an algebra B and an indecomposable module M such that $q_A(m) > 0$ for m the dimension vector of M . Then B is also accessible. By induction hypothesis, $\text{Tr}(\phi_B) \geq -1$. Then

$$\text{Tr}(\phi_A) = \text{Tr}(\phi_B) + (mC_B^T m^T - 1) \geq -1 + (mC_B^T m^T - 1) = -1 + (q_B(m) - 1) \geq -1$$

This shows (a).

For (b) assume that $\text{Tr}(\phi_B) = -1$ and $q_B(m) = 1$, then

$$\text{Tr}(\phi_A) = \text{Tr}(\phi_B) + (mC_B^T m^T - 1) = -1 + (mC_B^T m^T - 1) = -1 + (q_B(m) - 1) = -1$$

□

4.7 Strongly accessible algebras

Theorem: *A finite dimensional accessible algebra A then it is strongly accessible if and only if $\text{Tr}(\phi_A) = -1$.*

Proof. Assume A is strongly accessible from A_0 . Since $q_A(m) \geq 1$, for $A = B[M]$ a one-point extension of the subcategory B of A by the exceptional module M (since then $q_A(m) = \dim_K \text{End}_A(M)$). By the Proposition above

$$\text{Tr}(\phi_A) = \text{Tr}(\phi_{A_{n-1}}) = \dots = \text{Tr}(\phi_{A_0}) = -1$$

Conversely, assume that $\text{Tr}(\phi_A) = -1$ and write $A = B[M]$ as a one-point extension of the subcategory B of A by the module M . We shall prove that M is exceptional.

$$-1 = \text{Tr}(\phi_A) = \text{Tr}(\phi_B) + (mC_B^T m^T - 1) \geq -1 + (mC_B^T m^T - 1) = -1 + (q_B(m) - 1) \geq -1$$

Equality holds and $q_B(m) = 1$, since M is indecomposable, it follows that the extension ring of M is trivial. □

4.8 Stable matrices

The following statement is Theorem 1 for stable matrices.

Proposition. *Suppose M is a stable unimodular $n \times n$ -matrix. Let $\chi_M = c_0 + c_1 T + c_2 T^2 + \dots + c_{n-2} T^{n-2} + c_{n-1} T^{n-1} + c_n T^n$ be its characteristic polynomial.*

Suppose that $0 < \text{Tr} M^k \leq m$ for $p \leq k \leq p + n - 1$ and certain integers $1 \leq p$ and m .

Then $0 < \text{Tr} M^k \leq m$ for all integers $p \leq k$.

In particular, M is of cyclotomic type.

Proof. Consider the coefficients c_0, c_1, \dots, c_n of χ_M . Since M is stable then $c_n = 1, c_{n-1} < 0, c_{n-2} > 0$ and the signs alternate until we meet a j with $c_j c_0 < 0$. Cayley-Hamilton theorem states that $\chi_M(M) = 0$. Then

$$0 = c_0 1_n + c_1 M + c_2 M^2 + \dots + c_{n-1} M^{n-1} + c_n M^n$$

Then

$$c_0 1_n + c_2 M^2 + \dots + c_{2m} M^{2m} = c_1 M + c_3 M^3 + \dots + c_{2m-1} M^{2m-1} + c_{2(m+r)-1} M^{2(m+r)-1}$$

Let $c > 0$ be the common value of the trace of this matrix.

Write $n = 2m + r$ for $r = 0$ or 1 . Consider the matrices

$$P = \frac{1}{c} (c_0 1_n + c_2 M^2 + \dots + c_{2m} M^{2m})$$

$$Q = -\frac{1}{c} \left((c_1 M + c_3 M^3 + \dots + c_{2m-1} M^{2m-1} + c_{2(m+r)-1} M^{2(m+r)-1}) \right)$$

so that we get two expressions of P as positive linear combinations of powers of M .

Suppose that $n = 2m + 1$. By hypothesis we have $\text{Tr}(P) \leq n$. Moreover, since $c_n = 1$ then

$$\text{Tr}(M^n) \leq \text{Tr}(Q) = \text{Tr}(P) \leq n$$

The claim follows by induction.

Otherwise, $n = 2m$. The claim follows similarly. \square

4.9 Theorem 1

Proof of Theorem 1. Observe that $M = \phi_A$ is a real unimodular matrix. One implication of the Theorem was shown before. Suppose that $|\text{Tr}(M^k)| \leq n$ or equivalently, $-n \leq \text{Tr}(M^k) \leq n$ for $0 \leq k \leq n$. The Proposition above yields that M is cyclotomic. \square

4.10 Polynomials of Littlewood type

An integral self-reciprocal polynomial $p(t) = p_0 + p_1 t + \dots + p_{n-1} t^{n-1} + p_n t^n$ is of Littlewood type if every coefficient non-zero p_i has modulus 1. A polynomial $p(t)$ of Littlewood type with all $p_i \neq 0$, for $i = 0, 1, \dots, n$, is said to be Littlewood.

Lemma. *If z is a root of a polynomial of Littlewood type, then*

$$1/2 < |z| < 2$$

Proof. Suppose z is a root of a polynomial of Littlewood type. Then

$$1 = \epsilon_1 z + \epsilon_2 z^2 + \dots + \epsilon_n z^n$$

for some $\epsilon_i \in \{-1, 0, 1\}$.

If $|z| < 1$ then $1 \leq |z| + |z|^2 + \dots + |z|^n < |z|/(1 - |z|)$ so $|z| > 1/2$. Since z is the root of a polynomial of Littlewood type if and only if z^{-1} is, then $1/2 < |z| < 2$.

Moreover, if $|z| > 1$, then $1/|z| < 1$ and $1/2 < 1/|z| < 2$. Hence $1/2 < |z| < 2$. \square

4.11 Littlewood series

Definition. A *Littlewood series* is a power series all of whose coefficients are 1, 0 or -1 .

Let $P = \{z \in \mathbb{C} : z \text{ is the root of some Littlewood polynomial}\}$.

Remarks:

- a. Littlewood series converge for $|z| < 1$.
- b. A point $z \in \mathbb{C}$ with $|z| < 1$ lies in P if and only if some Littlewood series vanishes at this point.
- c. A Littlewood polynomial is not a Littlewood series. But any Littlewood polynomial, say $p(z) = a_0 + \dots + a_d z^d$ yields a Littlewood series having the same roots z with $|z| < 1$: indeed, consider the series

$$P(z) = p(z)/(1 - z^{d+1}) = a_0 + \dots + a_d z^d + a_0 z^{d+1} + \dots + a_d z^{2d+1} + a_0 z^{2d+2} + \dots$$

Thus $P \subset R$, where R is the set of roots of Littlewood series. We shall show the Proposition at the Introduction.

Proof. Let \mathcal{L} be the set of Littlewood series. Then $\mathcal{L} = \{-1, 0, 1\}^{\mathbb{N}}$, so with the product topology it is homeomorphic to the Cantor set. Choose $0 < r < 1$. Let \mathbb{F} be the space of finite multisets of points z with $|z| < r$, modulo the equivalence relation generated by $S \cong S \cup X$ when $|X| = r$.


Claim. Any Littlewood series has finitely many roots in the disc $|z| \leq r$. The map $f : \mathcal{L} \rightarrow \mathbb{F}$ sending a Littlewood series to its multiset of roots in this disc is continuous.

Since \mathcal{L} is compact, the image of f is closed. From this we can show that R , the set of roots of Littlewood series, is closed. Since Littlewood polynomials are densely included in \mathcal{L} and f is continuous, we get that P , the set of roots of Littlewood polynomials, is dense in R . It follows that $\bar{P} = R$, as we wanted to show. \square

5. An example

5.1 Construction

For m a natural number and let $n = 3 + 6m$. Let R_n be an algebra formed by n commutative squares. Consider the one-point extension $A_m = R_n[P_n]$ with P_n the unique indecomposable projective R_n -module of K -dimension 2. Observe that A_m (resp. C_{n-1}) is given by the following quiver with $n + 1$ vertices and commutative relations (resp. $n - 1$ vertices and relations):



We claim:

- a. $\chi_{A_m} = T^n + T^{n-1} - T^3 \chi_{A_{m-1}} + T + 1$, for all $n \geq 1$. As consequence, the algebras A_m and C_n are of Littlewood type;
- b. the number of eigenvalues of ϕ_{A_m} not lying in the unit disk is at least m ;
- c. $\mathbb{M}(\chi_{A_m}) \leq 8$.

Proof. (a): Consider $m \geq 1$, $n = 3 + 6m$ and the algebra $B_n = R_{3+6m}$ such that $A_m = B_n[P_n]$ and the perpendicular category P_n^\perp in $D^b(B_n)$ is derived equivalent to $\text{mod}(C_{n-1})$ where C_{n-1} is a proper quotient of an algebra derived equivalent to R_{2+6m} . Therefore

$$\begin{aligned}\chi_{A_{m+1}} &= (T+1)\chi_{R_{n+6}} - T\chi_{C_{n+5}} \\ &= (T+1)(T^{n+6} + T^{n+5} + T+1) - T^3(T+1)\chi_{R_n} - T\chi_{C_{n+5}}\end{aligned}$$

We shall calculate $\chi_{C_{2+6m}}$. Observe that C_{2+6m} is tilting equivalent to the one-point extension $R_{1+6m}[P_1]$. Hence

$$\begin{aligned}\chi_{C_{2+6m}} &= (T+1)\chi_{R_{1+6m}} - T\chi_{R_{6m}} = T^{2+6m} + T^{1+6m} - T^3\left\{(T+1)\chi_{R_{1+6(m-1)}} - T\chi_{R_{6(m-1)}}\right\} \\ &\quad + T+1 = T^{2+6m} + T^{1+6m} - T^3\chi_{C_{2+6(m-1)}} + T+1\end{aligned}$$

which implies

$$\begin{aligned}\chi_{A_{m+1}} &= (T+1)(T^{n+6} + T^{n+5} + T+1) - T^3(T+1)\chi_{R_n} - T(T^{n+5} + T^{n+4} + T+1) \\ &\quad - T^3T\chi_{C_{n-1}} = T^{n+7} + T^{n+6} - T^3\chi_{A_m} + T+1\end{aligned}$$

as claimed.

As consequence of formula (a) we observe the following:

$$(a') \quad L(\chi_{A_m}) = 4m + 5.$$

(b) By induction, we shall construct polynomials r_m representing χ_{A_m} .

For $m = 0$, we have $\chi_{A_0} = T^4 + T^3 + T^2 + T + 1$, which is represented by the polynomial $r_0 = T^4 - 3T^2 + 1$.

Observe that $(T^{n-1} + 1) = v_n - Tv_{n-2}$ then $T^n + T^{n-1} + T + 1 = (T+1)(T^{n-1} + 1)$ is represented by $w_n = T(u_{n-1} - u_{n-3})$.

For $n = 4 + 6m$, we define $r_m = w_n - T^3r_{m-1}$. We verify by induction on m that r_m represents χ_{A_m} :

$$\begin{aligned}\chi_{A_m}(T^2) &= (T^2+1)(T^{2n-2}+1) - T^6\chi_{A_{m-1}}(T^2) \\ &= T^n w_n(T+T^{-1}) - T^6T^{n-6}r_{m-1}(T+T^{-1}) = T^n r_m(T+T^{-1})\end{aligned}$$

For instance,

$$\begin{aligned}r_1 &= w_{10} - T^3r_0 = T\{(T^9 - 8T^7 + 21T^5 - 20T^3 + 5T) - (T^7 - 6T^5 + 10T^3 - 4T)\} \\ &\quad - T^3\{T^4 - 3T^2 + 1\} \\ &= T^{10} - 9T^8 - T^7 + 27T^6 + 3T^5 - 30T^4 - T^3 + 9T^2\end{aligned}$$

which has $\xi(r_1) = 4$ changes of sign in the sequence of coefficients. According to *Descartes rule of signs*, r_1 has at most $\xi(r_1) = 4$ positive real roots. Since r_1 represents χ_{A_1} , then χ_{A_1} has at most $2\xi(r_1) = 8$ roots in the unit circle. That is, χ_{A_1} has at least 2 roots z with $|z| \neq 1$.

We shall prove, by induction, that r_m has at most $\xi(r_m) = 2(m+1)$ positive real roots. Indeed, write

$$r_m = T^n - (n-1)T^{n-2} - T^3q_m + (n-1)T^2$$

for some polynomial q_m of degree $n - 6$ with signs of its coefficients $+ - - + + - - \dots \pm$ so that $\xi(q_m) = 2m$. Then

$$r_{m+1} = w_{n+6} - T^3 r_m = Tu_{n+5} - Tu_{n+3} - T^3 r_m$$

an addition of three polynomials with signs of coefficients given as follows:

$$\begin{array}{cccccccc} + & 0 & - & 0 & + & 0 & - & 0 & \dots & + & 0 & 0 \\ & & - & 0 & + & 0 & - & 0 & \dots & + & 0 & 0 \\ & & & - & + & + & - & - & \dots & 0 & 0 & 0 \end{array}$$

Hence $r_{m+1} = T^{n+6} - (n+5)T^{n+4} - T^3 q_{m+1} + (n+5)T^2$ where the polynomial q_{m+1} of degree n has signs of its coefficients $+ - - + + - - \dots \pm$ so that $\xi(q_{m+1}) = \xi(q_m) + 2 = 2(m+1)$. Hence $\xi(r_m) = 2 + \xi(q_m) = 2(m+1)$.

By the Lemma below, χ_{A_m} has at most $4(m+1)$ roots in the unit circle. Equivalently, χ_{A_m} has at least $4 + 6m - 4(m+1) = 2m$ roots outside the unit circle. Hence χ_{A_m} has at least m roots z satisfying $|z| > 1$.

Lemma. Let q be a polynomial representing the polynomial p . Assume q accepts at most s positive real roots, then p has at most $2s$ roots in the unit circle.

Proof. Let μ_1, \dots, μ_s be the positive real roots of q . Let $z = a + ib$ be a root of p with $a^2 + b^2 = 1$. Consider $w = c + id$ a complex number with $w^2 = z$. Then $0 = p(z) = w^n q(w + w^{-1})$ where $w + w^{-1} = (c + id) + (c - id) = 2c$. Then $2c = \epsilon \lambda_j$ for some $\epsilon \in \{1, -1\}$ and $1 \leq j \leq s$. Hence

$$z = w^2 = \left(\frac{1}{2} \lambda_j^2 - 1 \right) + i \left(2\epsilon \lambda_j \sqrt{1 - \lambda_j^2} \right)$$

can be selected in two different ways. □

(c) For $n = 6m + 4$ we have $\chi_{A_m} = T^n + T^{n-1} - T^3 \chi_{A_{m-1}} + T + 1$. Then

$$\chi_{A_m} = \xi_m + (-1)^{m-1} T^{2m+4} \chi_{10}, \text{ where } \xi_m = T^n + T^{n-1} - T^3 \xi_{m-1} + T + 1$$

for $m \geq 2$ and $\xi_1 = 0$.

We observe that ξ_m is a product of cyclotomic polynomials. Indeed, since $\xi_m(-1) = 0$ we can write

$$\xi_m = (T + 1) \sigma_m \text{ and } \sigma_m = T^{n-1} - T^3 \sigma_{m-1} + 1$$

for $m \geq 2$ and $\sigma_1 = 0$.

Recall $\Phi_{2^s-1} = T^{s-1} + T^{s-2} + \dots + T + 1$ and $\Phi_{2^s}(T) = \Phi_s(-T)$. Moreover, $\Phi_{3p}(T) = \Phi_p(T^3)$, if p is a power of 2. Altogether this yields

$$\begin{aligned} \Phi_{6(2^{2(m+1)}-1)}(T) &= \Phi_{2(2^{2(m+1)}-1)}(T^3) = \Phi_{2^{2(m+1)}-1}(-T^3) \\ &= T^{6m+3} - T^{6m} + \dots - T^3 + 1 = \sigma_m \end{aligned}$$

hence

$$\xi_m = \Phi_2 \Phi_{6(2^{2(m+1)}-1)}$$

confirming the claim.

We estimate the Mahler measure of $\chi_{A_m} = \xi_m + (-1)^{m-1} T^{2m+4} \chi_{A_{10}}$. Write $\chi_{A_m} = f_m + g_m$, where f_m is the cyclotomic summand. Observe that $L(g_m) = L(\chi_{A_{10}}) = 8$ and apply Lemma (3.4) with $\mathbb{M}(f_m) = 1$ to get

$$\mathbb{M}(\chi_{A_m}) \leq \mathbb{M}(f_m)L(g_m) = 8$$

With the help of computer programs we calculate more accurate values of the Mahler measure of some of the above examples:

No. vertices	No. roots outside unit disk	Mahler measure
178	29	1.28368024451292
184	30	1.28327850483340
190	31	1.28386917621114
196	32	1.28395305512596

Comparing with the list of *Record Mahler measures by roots outside the unit circle* in Mossinghoff's web page we see:

- i. for the entry 29 the Mahler measure is the same in both tables;
- ii. the entries 30 and 31 have a smaller Mahler measure in our table, establishing new records;
- iii. the entry 32 of our table seems to be new. Further entries could be calculated.

6. Coefficients of Coxeter polynomials

6.1 Derived tubular algebras


There are interesting invariants associated to the Coxeter polynomial of a triangular algebra $A = k[\Delta]/I$. For instance, the evaluation of the Coxeter polynomial $\chi_A(-1) = m^2$ for some integer m . Clearly, this number is a derived invariant. A simple argument yields that $m = 0$ in case Δ has an odd number of vertices. In [14], it was shown that for a representation-finite accessible algebra A with $\text{gl.dim } A \leq 2$ the invariant $\chi_A(-1)$ equals zero or one. The criterion was applied to show that a canonical algebra is derived equivalent to a representation-finite algebra if and only if it has weight type $(2, p, p + k)$, where $p \geq 2$ and $k \geq 0$. In particular, the tubular canonical algebra of type $(3, 3, 3)$ is not derived equivalent to a representation-finite algebra, while the tubular algebras of type $(2, 4, 4)$ or $(2, 3, 6)$ are.

6.2 Strong towers

Recall from [14] that a *strong tower* $\mathbb{T} = (A_0 = k, A_1, \dots, A_n = A)$ of access to A satisfies that $A_{i+1} = A_i[M_i]$ or $[M_i]A_i$ for some exceptional module M_i in such a way that, in case $A_{i+1} = A_i[M_i]$ (resp. $A_{i+1} = [M_i]A_i$), the perpendicular category $\underline{M_i}^\perp$ (resp. ${}^\perp \underline{M_i}$) of M_i in mod_{A_i} is equivalent to $\text{mod}_{C_{i-1}}$ for some accessible algebra C_{i-1} , $i = 1, \dots, n - 1$. In the extension situation the perpendicular category M_i^\perp (resp. ${}^\perp M_i$ in the coextension situation) in $D^b(\text{mod}_{A_i})$ is equivalent to $D^b(\text{mod}_{C_{i-1}})$ and B_i is derived equivalent to a one-point (co-)extension of C_{i-1} . An algebra C_i as above is called an *i-th perpendicular restriction* of the tower \mathbb{T} , observe that it is well-defined only up to derived equivalence. We denote by s_i the number of connected components of the algebra C_i ; in particular, $s_1 = 1$.

There are many examples of *strongly accessible algebras*, that is, algebras derived equivalent to algebras with a strong tower of access. The following are some instances:

- a. A canonical algebra C of weight (p_1, \dots, p_t) is strongly accessible if and only if $t = 3$, in that case, C is derived-equivalent to a representation-finite algebra if and only if the weight type does not dominate $(3, 3, 3)$.
- b. The following sequence of poset algebras defines strong towers of access:



6.3 Towering numbers

Consider a strong tower $\mathbb{T} = (A_0 = k, A_1, \dots, A_n = A)$ of access to A such that A_{i+1} is an one-point (co)extension of A_i by M_i and C_{i-1} the corresponding i -th perpendicular restriction of \mathbb{T} . Let C_{i-1} have s_{i-1} connected components, $i = 2, \dots, n-1$. Define the *first towering number* of \mathbb{T} as the sum $s_{\mathbb{T}}(A) = \sum_{i=1}^{n-2} s_i$.


Theorem. Let A be a strongly accessible algebra with n vertices, then the first towering number $s_{\mathbb{T}}(A) = \sum_{i=1}^{n-2} s_i$ of \mathbb{T} is a derived invariant, that is, depends only on the derived class of A . It is $s_{\mathbb{T}}(A) = n - 1 - a_2$, where a_2 is the coefficient of the quadratic term in the Coxeter polynomial of A .

Proof. Assume $A = A_n$ and $B = A_{n-1}$ such that $A = B[M]$ for M an exceptional B -module and let $C = C_{n-2}$ be the algebra such that mod_C is derived equivalent to the perpendicular category M^\perp formed in $D^b(\text{mod}_B)$. Then $\chi_A(t) = (1+t)\chi_B(t) - t\chi_C(t)$. Write $\chi_B(t) = 1 + t + \sum_{i=2}^{n-3} b_i t^i + t^{n-2} + t^{n-1}$ and $\chi_C(t) = 1 + \sum_{i=1}^{n-3} c_i t^i + t^{n-2}$. By induction hypothesis we may assume that $s(B) = n - 2 - b_2$. Then $a_2 = b_2 + 1 - c_1$. Moreover, since C is a direct sum accessible algebras, then $c_1 = \sum_{i=0}^{n-2} (-1)^i \dim_k H^i(C) = \dim_k H^0(C) = s_{n-2}$. Hence $a_2 = n - 1 - s(B) - s_{n-2} = n - 1 - s(A)$. \square

Corollary. Let $\mathbb{T} = (A_1 = k, \dots, A_n = A)$ be a strong tower of access to A . Let $A = B[M]$ for $B = A_{n-1}$ with M exceptional and C a perpendicular restriction of B via M . Consider the Coxeter polynomials $\chi_A(t) = 1 + t + a_2 t^2 + \dots + a_{n-2} t^{n-2} + t^{n-1} + t^n$ and $\chi_B(t) = 1 + t + b_2 t^2 + \dots + b_{n-3} t^{n-3} + t^{n-2} + t^{n-1}$, then $a_2 \leq b_2$, with equality if and only if C is connected. In particular, $a_2 \leq 1$.

Proof. First recall that for a connected accessible algebra the linear term of the Coxeter polynomial has coefficient 1. Let $\chi_C(t) = 1 + c_1 t + c_2 t^2 + \dots + c_{n-4} t^{n-4} + c_{n-3} t^{n-3} + t^{n-2}$ be the Coxeter polynomial of C . If C is the direct sum of connected accessible algebras C_1, \dots, C_s , then $c_1 = s$. Therefore, $a_2 = b_2 + b_1 - c_1 = b_2 - (s - 1) \leq b_2$. By induction hypothesis, we get $a_2 \leq 1$. \square

Let A be the algebra given by the following quiver with relation $\gamma\beta\alpha = 0$:



which is derived equivalent to the quiver algebra B with the zero relation as depicted in the second diagram. Clearly, $A = A'[M]$, where A' is a quiver algebra of type \mathbb{A}_4 and M is an indecomposable module with M^\perp the category of modules of the disconnected quiver $\bullet \rightarrow \bullet \rightarrow \bullet$, that is $s_3(A) = 2$. Moreover $s_2(A) = s_2(A') = 1$ and $s(A) = 4$. On the other hand $B = [N]B'$ such that B' is not hereditary. A calculation yields $s_3(B) = 1$ and $s_2(B) = s_2(B') = 2$, obviously implying that $s(B) = 4$.

Some properties of the invariant s :

- i. Let A and B be accessible algebras and A be accessible from B , then $s(B) \leq s(A)$. Equality holds exactly when $A = B$.
- ii. Let A be an accessible schurian algebra (that is for every couple of vertices i, j , $\dim_k A(i, j) \leq 1$), then for every convex subcategory B we have $s(B) \leq s(A)$.

6.4 Totally accessible algebras

An accessible algebra A with $n = 2r + r_0$ vertices, and $r_0 \in \{0, 1\}$, is said to be *totally accessible* if there is a family of (not necessarily connected) algebras $C^{(n)} = A', C^{(n-2)}, C^{(n-4)}, \dots, C^{(r_0)}$ satisfying:

- a. A is derived equivalent to A' ;
- b. for each $0 \leq i = n - 2j \leq n$, there is a strong tower $\mathbb{T}^{(j)} = (C^{(j,1)} = k, \dots, C^{(j,i)} = C^{(i)})$ of access to $C^{(i)}$;
- c. $C^{(i-2)}$ is an $i - 1$ -th perpendicular restriction of $\mathbb{T}^{(j)}$, that is, $C^{(i)}$ is a one-point (co)extension of $C^{(j,i-1)}$ by a module N_{i-1} and $C^{(i-2)}$ is a perpendicular restriction of $C^{(j,i-1)}$ via N_{i-1} .

The tower $\mathbb{T}^{(j)}$ is said to be a j -th derivative of the tower $\mathbb{T}^{(0)}$.

Examples that we have encountered of totally accessible algebras are:

- i. Hereditary tree algebras: since for any connected hereditary tree algebra A with at least 3 vertices, there is an arrow $a \rightarrow b$ with a a source (or dually a sink) and $A = B[P_b]$ such that the perpendicular restriction of B via P_b is the algebra hereditary tree algebra C obtained from A by deleting the vertices a, b .
- ii. Accessible representation-finite algebras A with $\text{gl.dim } A \leq 2$, since then the perpendicular restrictions of any strong tower (which exists by [14]) satisfy the same set of conditions.
- iii. Certain canonical algebras: for instance the tame canonical algebra A of weight type $(2, 4, 4)$ is an extension $A = B[M]$ of a hereditary algebra B of extended Dynkin type $[2, 4, 4]$ by a module M in a tube of rank 4, then the perpendicular restriction of B via M is the hereditary algebra C of extended Dynkin type $[3, 3, 3]$, see for example [?](10.1). Since C is totally accessible, so A is. Moreover $s(A) = 8$.
- iv. Let A be an accessible algebra of the form $A = B[M]$ for an algebra B and an exceptional module M and let C the perpendicular restriction of B via M . If A is totally accessible, then B and C are totally accessible.

The following results extend some of the features observed in the examples above.

Proposition. *a. Assume that A is a totally accessible algebra, then $\chi_A(-1) \in \{0, 1\}$.*

b. Assume that A is an accessible but not totally accessible algebra with $\text{gl.dim } A \leq 2$, then one of the following conditions hold:

i. for every exceptional B -module such that $A = B[M]$ and any perpendicular restriction C of B via M , then C is not accessible;

ii. there exists a homological epimorphism $\phi : A \rightarrow B$ such that $\chi_B(-1) > 1$.

Proof. (a): Consider the perpendicular restriction C of B via M , such that $\chi_A(t) = (1+t)\chi_B(t) - t\chi_C(t)$. Therefore $\chi_A(-1) = \chi_C(-1)$ and moreover, C is totally accessible. Then by induction hypothesis, $\chi_A(-1) = \chi_{C^{(m)}}(-1)$ for a totally accessible algebra $C^{(m)}$ with number of vertices $m = 1$ or $m = 2$. Clearly, $C^{(m)}$ is either k , $k \oplus k$ or hereditary of type \mathbb{A}_2 , which yields the desired result.

(b): Assume A is an accessible algebra with $\text{gl.dim } A \leq 2$ and such that for every homological epimorphism $\phi : A \rightarrow B$ we have $\chi_B(-1) \in \{0, 1\}$. Let $A = B[M]$ for an accessible algebra B and an exceptional B -module M such that C is a perpendicular restriction of B via M . Since $\text{gl.dim } A \leq 2$ then there is a homological epimorphism $A \rightarrow C$ and $\text{gl.dim } C \leq 2$. Observe that for every homological epimorphism $\psi : B \rightarrow B'$ (resp. $\psi : C \rightarrow C'$) there is a homological epimorphism $\phi : A \rightarrow B'$ (resp. $\phi : A \rightarrow C'$), hence $\chi_{B'}(-1)$ (resp. $\chi_{C'}(-1)$) is 0 or 1. By induction hypothesis, B is totally accessible. Moreover if C is accessible, then the induction hypothesis yields that C is totally accessible and also A is totally accessible, a contradiction. Therefore C is not accessible. \square

7. On the quadratic coefficient of the Coxeter polynomial of a totally accessible algebra

7.1 Derived algebras of linear type

Recall that an *extended canonical* algebra of weight type $\langle p_1, \dots, p_t \rangle$ is a one-point extension of the canonical algebra of weight type $[p_1, \dots, p_t]$ by an indecomposable projective module. As in (1.3), the extended canonical algebras of type $\langle p_1, p_2, p_3 \rangle$ is strongly accessible. Moreover, the extended canonical algebra A of type $\langle 3, 4, 5 \rangle$ (with 12 points) has Coxeter polynomial $1 + t + t^2 + \dots + t^{12}$ which is also the Coxeter polynomial of a linear hereditary algebra H with 12 vertices. Clearly A and H are not derived equivalent.

The following generalizes a result of Happel who considers the case of Coxeter polynomials associated to hereditary algebras [8].

Theorem 1. *Let A be a totally accessible algebra with n vertices and let $\chi_A(t) = \sum_{i=0}^n a_i t^i$ be the Coxeter polynomial of A . The following are equivalent:*

i. $a_2 = 1$;

ii. let $\mathbb{T} = (A_1 = k, \dots, A_{n-1}, A_n = A)$ be a strong tower of access to A and C_i the i -th perpendicular restriction of \mathbb{T} , for all $1 \leq i \leq n - 2$, then the algebras C_i are connected;

iii. A is derived equivalent to a quiver algebra of type \mathbb{A}_n .

Proof. (i) \Leftrightarrow (ii): Let $\mathbb{T} = (A_1 = k, \dots, A_n = A)$ be a strong tower of access to A . In case each C_i is connected, then $s(A) = n - 2$, that is $a_2 = 1$. If $a_2 = 1$, then $n - 2 = s_{\mathbb{T}}(A) = \sum_{i=1}^{n-2} s_i$ with each $s_i \geq 1$. (i) \Leftrightarrow (iii): We know that an algebra A derived equivalent to a quiver algebra of type \mathbb{A}_n has $\chi_A(t) = \sum_{i=0}^n t^i$, in particular, $a_2 = 1$. Assume that an accessible algebra A has the quadratic coefficient of its Coxeter polynomial $a_2 = 1$. Let $A = B[M]$ for an accessible algebra $B = A_{n-1}$ and an exceptional module M . Since B is also totally accessible with a tower $\mathbb{T}' = (A_1 = k, \dots, A_{n-1} = B)$ satisfying (ii), then the quadratic coefficient of the Coxeter polynomial of B is $b_2 = 1$ and we may assume that B is derived equivalent to a quiver algebra of type \mathbb{A}_{n-1} . In particular, B is representation-finite with a preprojective component \mathcal{P} such that the orbit graph $\mathcal{O}(\mathcal{P})^\tau$ is of type \mathbb{A}_{n-1} (recall that the orbit graph has vertices the τ -orbits in the quiver \mathcal{P} with Auslander-Reiten translation τ and there is an edge between the orbit of X and the orbit of Y if there is some numbers a, b and an irreducible morphism $\tau^a X \rightarrow \tau^b Y$). Observe that for any X in $D^b(\text{mod}_A)$ not in the orbit of M , there is some translation $\tau^a X$ belonging to M^\perp , implying that in case M^τ has two neighbors in the orbit graph then M^\perp is not connected, that is $s_{n-2} > 1$ and $a_2 = n - 1 - s(A) \leq 0$, a contradiction. Therefore, M^τ has just one neighbor in $\mathcal{O}(\mathcal{P})^\tau$, hence A is derived of type \mathbb{A}_n . \square

7.2 Theorem 2

Consider a tower $A_1, \dots, A_n = A$ of accessible algebras where A_{i+1} is a one-point (co)extension of A_i by the indecomposable M_i and C_i is such that M_i^\perp is derived equivalent to $D^b(\text{mod}_{C_i})$. Assume that $C_i^{(j)}$, for $1 \leq j \leq s_i$, are the connected components of the category C_i . Consider the corresponding Coxeter polynomials:

$$\begin{aligned}\chi_{A_i}(t) &= 1 + t + \sum_{j=2}^{i-2} a_j^{(i)} t^j + t^{i-1} + t^i, \\ \chi_{C_i}(t) &= 1 + s_i t + \sum_{r=2}^{n_i-2} c_{i,r} t^r + s_i t^{n_i-1} + t^{n_i}, \\ \chi_{C_i^{(j)}}(t) &= 1 + t + \sum_{s=2}^{n_{i,j}-2} c_{i,s}^{(j)} t^s + t^{n_{i,j}-1} + t^{n_{i,j}},\end{aligned}$$

where clearly, $\sum_{j=1}^{s_i} n_{i,j} = n_i$.

Lemma. (α) For every $1 \leq j \leq i - 2$, we have $a_j^{(i)} \leq 1$.

($\alpha\alpha$) For every $1 \leq j \leq i - 2$, we have $a_j^{(i)} \leq c_{i,j}$ and $a_j^{(i)} \leq a_j^{(i-1)}$.

Proof. We shall check that (α) implies ($\alpha\alpha$), then we show that (α') holds by induction on j .

Indeed, assume that (α) holds and proceed to show ($\alpha\alpha$) by induction on j . If $j = 0, 1$, then $a_j^{(i)} = 1 = a_{i-j}^{(i)}$. Assume that $2 \leq j \leq i - 2$ and $a_j^{(i)} \leq c_{i,j}$ and $a_j^{(i)} \leq a_j^{(i-1)}$. Then

$$a_{j+1}^{(i)} = a_{j+1}^{(i-1)} + \left(a_j^{(i-1)} - c_{j,i-1} \right) \leq a_{j+1}^{(i-1)} \leq \dots \leq a_{j+1}^{(j+1)} = 1.$$

Let $0 \leq j \leq i - 2$. If $j = 0, 1$ we have $a_0^{(i)} = 1 = c_{i,0}$ and $a_1^{(i)} \leq s_1(A) = c_{i,1}$. Moreover $a_1^{(i)} = a_1^{(i-1)}$. Assume (α) holds for $j \geq 2$, then.

$$a_{j+1}^{(i)} = a_{j+1}^{(i-1)} + \left(a_j^{(i-1)} - c_{j,i-1} \right) \leq a_{j+1}^{(i-1)},$$

$$a_{j+1}^{(i)} - c_{i,j+1} = a_{j+2}^{(i)} - a_{j+2}^{(i-1)} \leq 0. \quad \square$$

Theorem 2. Let A be a totally accessible algebra with Coxeter polynomial $\chi_A(t) = 1 + t + a_2 t^2 + \dots + a_{n-2} t^{n-2} + t^{n-1} + t^n$, then:

a. $a_j \leq 1$, for every $2 \leq j \leq n-2$;

b. if for some $2 \leq j \leq n-2$, we have $a_j = 1$ then A is derived equivalent to a hereditary algebra of type \mathbb{A}_n .

Proof. Keep the notation as in (4.1). Then (a) is the case $i = n$ of the Lemma above.

We shall prove (b) by induction on n the number of vertices of A . Let $j = 2$ and assume $a_2 = 1$, then (3.1) implies that A is derived equivalent to \mathbb{A}_n . Consider now $2 < j < n-2$ and assume that $a_j = 1$, we get:

$$1 = a_j^{(n)} = a_j^{(n-1)} + \left(a_{j-1}^{(n-1)} - c_{n-1,j-1} \right) \leq a_j^{(n-1)} \leq 1$$

The last inequality due to (a), hence $a_j^{(n-1)} = 1$. Induction hypothesis yields that A_{n-1} is derived equivalent to \mathbb{A}_{n-1} and its Auslander-Reiten quiver consists of a preprojective component \mathcal{P} . In particular, $a_2^{(n-1)} = 1$, which implies that $s_{n-3}(A_{n-1}) = 1$, that is, $A = A_{n-1}[M]$ for some exceptional module M such that M^\perp is derived equivalent to mod_C for a connected algebra C , that is, $s(A) = n-2$ and by (3.1), $A = B[M]$ is derived equivalent to a hereditary algebra of type \mathbb{A}_n . \square

7.3 Examples

If A is a representation-finite accessible algebra with $\text{gl.dim } A \leq 2$, then A is totally accessible. On the other hand the algebra B with quiver:

$$1 \xrightarrow{x} 2 \xrightarrow{x} 3 \xrightarrow{x} 4 \dots \xrightarrow{x} 11 \xrightarrow{x} 12$$

and $x^3 = 0$ is representation-finite and accessible (but not $\text{gl.dim } B \leq 2$). The Coxeter polynomial of B is:

$$\chi_B(t) = 1 + t - t^3 - t^4 + t^6 - t^8 - t^9 + t^{11} + t^{12}.$$

Then observe that the 6-th coefficient is 1 but the algebra B is not derived equivalent to Dynkin type \mathbb{A}_{12} .

8. On the traces of Coxeter matrices

Let A be an algebra such that not all roots of χ_A are roots of unity. By the result of Kronecker [36], not all of the spectrum of A lies in the unit disk. Equivalently, the spectral radius $\rho_A = \max\{|\lambda| : \lambda \text{ eigenvalue of } \phi_A\} > 1$. Arrange the eigenvalues of ϕ_A so that $\mu_1, \mu_2, \dots, \mu_n$ have absolute values $\rho_A = r_1 > r_2 > \dots > r_s$ and multiplicities m_1, \dots, m_s , respectively. Therefore $s \geq 2$ and

$$|\det \phi_A| = r_1^{m_1} r_2^{m_2} \dots r_s^{m_s} = 1.$$

We define the *critical power* $\kappa(A)$ as the minimal k such that

$$|\text{Tr}(\phi_A^k)| > n$$

Since r_1 is a simple eigenvalue of ϕ_A , then it follows that $\kappa(A)$ is well defined due to the existence of k satisfying the following chain of inequalities:

$$|\text{Tr}(\phi_A^k)| = \left| \sum_{j=1}^n \mu_j^k \right| \geq r_1^{km_1} - \sum_{j=2}^s r_j^{km_j} \geq r_1^k - (n-1)r_2^k > n.$$

The following is a reformulation of Theorem 2.

Theorem. Let A be an algebra such that not all roots of χ_A are roots of unity. We have $\kappa(A) \leq n$.

Proof. Indeed, suppose that A is not of cyclotomic type and $\kappa(A) > n$, that is, $|\text{Tr}(\phi_A^k)| \leq n$ for all $0 \leq k \leq n$. Observe that $M = \phi_A$ is a unimodular matrix and therefore, Theorem 2 implies that M is of cyclotomic type, which yields a contradiction. \square

Remark: We consider explicitly the case $n = 2$ in the above Theorem. Obviously, the Cartan matrix of A is of the form

$$C = \begin{pmatrix} 1 & a \\ 0 & 1 \end{pmatrix} \quad \phi_A = -C^{-1}C^T = \begin{pmatrix} a^2 - 1 & a \\ -a & -1 \end{pmatrix}$$

for some $a \geq 1$. Then ϕ_A has the indicated shape. If A is not cyclotomic, then $a \geq 3$ and $\text{Tr}(\phi_A^2) = (a^2 - 2)^2 - 2 > 2$.

9. Stability of a real matrix

9.1 Stability of matrices and the Lyapunov criterion

Let M be a real invertible $n \times n$ -matrix with eigenvalues $\lambda_j = r_j e^{i\theta_j}$, for some numbers $\theta_j \in [0, 2\pi)$ and $j = 1, \dots, n$. We will say that M is *stable* (resp. *semi-stable*) if the real part $\text{Re}(e^{i\theta_j}) = \cos \theta_j$ of the argument of the eigenvalue λ_j is positive (resp. non-negative), for every $j = 1, \dots, n$. The following is well-known, we sketch a proof for the sake of completeness.

Proposition. Let M be a stable (resp. semi-stable) $n \times n$ -matrix. Then the characteristic polynomial $\chi_M = T^n + a_{n-1}T^{n-1} + \dots + a_1T + a_0$ has coefficients satisfying $(-1)^{n-j}a_j > 0$ (resp. ≥ 0), for $j = 0, 1, \dots, n$;

Proof. Observe that $(-1)^n p(-T)$ is the product of polynomials $T - \alpha$ with $\alpha \in \mathbb{R}$ and $(T - (\alpha + i\beta))(T - (\alpha - i\beta)) = T^2 - 2\alpha T + (\alpha^2 + \beta^2)$ with $0 \neq \beta, \alpha \in \mathbb{R}$. Stability (resp. semi-stability) implies that $\alpha < 0$ (resp. $\alpha \leq 0$) above. Therefore, $(-1)^n p(-T)$ is product of polynomials with positive coefficients. \square

Remark: In most of the literature the stability concept we use goes by the name of *positive stability*, while the *stability* name is used also as *Hurwitz stability*, or *Lyapunov stability*.

The system of differential equations

$$y'(t) = -My(t)$$

is said to be *stable* if for every vector $d = (d_1, \dots, d_n)$, the solution $v(t) = e^{-tM}d$ of the above system has the property that $\lim_{t \rightarrow \infty} v(t) = 0$.

We recall here the celebrated.

Lyapunov criterion: The system $y'(t) = -My(t)$ is stable if and only if M is a stable matrix, equivalently there is a real positive definite matrix P such that

$$M^T P + PM = I_n.$$

It is not hard to see that given M , the corresponding P is unique. A proof of the criterion and its equivalence to other stability conditions are considered in [13].

9.2 Semi-stable powers

Let μ_1, \dots, μ_n be the eigenvalues of the real matrix M with $\mu_j = \rho_j e^{2\pi i \theta_j}$ in polar form. Observe that μ_j^k , for $j = 1, \dots, n$, are the eigenvalues of M^k and

$$\text{Tr} M^k = \sum_{j=1}^n \rho_j^k \cos(k\theta_j) \leq \sum_{j=1}^n |\mu_j^k| |\cos(k\theta_j)|$$

Lemma. For a positive integer $k \geq 1$ the following assertions are equivalent:

a. M^k is a semi-stable matrix;

b. $\text{Tr}(M^k) = \sum_{j=1}^n |\mu_j|^k |\cos(k\theta_j)|$.

Proof. If M^k is a semi-stable matrix, then $\mu_j^k = \rho_j^k (\cos(k\theta_j) + i \sin(k\theta_j))$ has $\cos(k\theta_j) \geq 0$. Since M is a real matrix then $\text{Tr}(M^k) = \sum_{j=1}^n \rho_j^k \cos(k\theta_j) \geq 0$. Therefore

$$\text{Tr}(M^k) = \sum_{j=1}^n \rho_j^k |\cos(k\theta_j)|.$$

Assume that $\text{Tr}(M^k) = \sum_{j=1}^n |\lambda_j|^k |\cos(k\theta_j)|$. Since $|\lambda_j^k| \geq \rho_j^k \cos(k\theta_j)$ for $j = 1, \dots, n$, adding up, we get

$$\text{Tr}(M^k) \geq \sum_{j=1}^n \rho_j^k \cos(k\theta_j) = \text{Tr}(M^k)$$

Hence we have equalities $|\lambda_j^k| |\cos(k\theta_j)| = \rho_j^k \cos(k\theta_j)$ for $j = 1, \dots, n$. Then M^k is semi-stable. \square

We say that k is a *stable power* (resp. *semi-stable power*) of M if M^k is a stable (resp. semi-stable) matrix.

10. Nakayama algebras

10.1 Cyclotomic Nakayama algebras

As a well-understood example the representation theory of the *Nakayama algebras* stands apart. Let $N(n, r)$ be the quotient obtained from the linear quiver with n vertices with radical rad_A of nilpotency index r .

For instance, for $A = N(6, 3)$ the Cartan matrix C and Coxeter matrix ϕ are:

$$C = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 & 1 \end{pmatrix} \text{ and } \phi = \begin{pmatrix} -1 & 1 & 0 & -1 & 1 & 1 \\ -1 & 0 & 1 & -1 & 0 & 1 \\ -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 1 & 0 \end{pmatrix}$$

whose characteristic polynomial is cyclotomic as we know from [18] or might be verified calculating $\text{Tr}(\phi_B^k) \leq n$, for $1 \leq k \leq 72$ and applying the criterion of Theorem 1. Indeed, for.

$k =$	$\text{Tr} \chi_A^k =$
11	-1
1, 2, 5, 7, 9, 10, 13, 14, 17	$= 1$
3, 6, 15	$= 2$
4, 8, 16	$= 3$
12	$= 6$

Starting with $k = 17$ the sequence of traces repeats cyclically. Therefore, $\text{Tr}(\chi_A^k) \leq 6$ for all $0 \leq k$. Then $N(6, 3)$ is of cyclotomic type.

10.2 An example

We recall in some length the argument given in [18] for the cyclotomicity of $N(n, 3)$, for all $n \geq 1$.

Consider the algebra R_{2n} with $2n$ vertices and whose quiver is given as

$$\begin{array}{ccccccc} 1 & \rightarrow & 2 & \rightarrow & 3 & \rightarrow & \cdots \rightarrow n \\ \downarrow & \searrow & \downarrow & \searrow & \downarrow & & \downarrow \\ 1' & \rightarrow & 2' & \rightarrow & 3' & \rightarrow & \cdots \rightarrow n' \end{array}$$

with all commutative relations. The corresponding Coxeter polynomial

$$\chi_{R_{2n}} = \chi_{\mathbb{A}_n} \otimes \chi_{\mathbb{A}_2} = v_{n+1} \otimes v_3$$

is a product of cyclotomic polynomials, therefore $\chi_{R_{2n}}$ is a cyclotomic polynomial. In fact $R_{2n} = \mathbb{A}_n \otimes \mathbb{A}_2$, where \mathbb{A}_s is the hereditary algebra associated to the linear quiver $1 \rightarrow 2 \rightarrow \cdots \rightarrow s$.

For $2m + 1$ odd, we consider.

$$R_{2m+1} \quad \begin{array}{ccccccc} \circ & \rightarrow & \circ & \rightarrow & \cdots & \rightarrow & \circ \rightarrow \circ \\ \downarrow & \searrow & \downarrow & & & & \downarrow \searrow \\ \circ & \rightarrow & \circ & \rightarrow & \cdots & \rightarrow & \circ \rightarrow \circ \end{array}$$

The following holds for the sequence of algebras R_n and its Coxeter polynomials χ_{R_n} :

- a. R_n is derived equivalent to $N(n, 3)$.

$$b. \chi_{R_n} = T^n + T^{n-1} - T^3 \chi_{R_{n-6}} + T + 1, \text{ for all } n \geq 6;$$

$$c. \mathbb{M}(\chi_{R_n}) = 1.$$

Observe that the sequence of algebras (R_n) forms an *interlaced tower of algebras*, that is, it is a sequence of triangular algebras R_1, \dots, R_n , such that R_s is a basic algebra with s simple modules and, among others, the condition

$$\chi_{R_{s+1}} = (T + 1)\chi_{R_s} - T\chi_{R_{s-1}}$$

is satisfied for $s = 1, \dots, n - 1$. Moreover, A_{s+1} is a one-point extension (or coextension) of an accessible algebra A_s by an exceptional A_s - module M_s such that the perpendicular category M_s^\perp formed in the derived category is triangular equivalent to $\text{mod}(A_{s-1})$, for $s = m + 1, \dots, n - 1$.

The following was shown in [18]: *Consider an interlaced tower of algebras A_m, \dots, A_n with $m \leq n - 2$. If $\text{Spec } \phi_{A_n}$ is contained in the union of the unit circle and the semi-ray of positive real numbers then either all A_i are of cyclotomic type or $M(\chi_{A_m}) < M(\chi_{A_n})$. In the latter case, $M(\chi_{A_n}) < \prod_{s=m}^{n-1} M(\chi_{A_s})$.*

Since we know that $M(\chi_{R_{2n}}) = 1$, for all $n \geq 0$, we conclude that $M(\chi_{R_n}) = 1$, for all $n \geq 0$. That is the Nakayama algebras of the form $N(n, 3)$ are of cyclotomic type.

10.3 Non-cyclotomic Nakayama algebras

Calculation of $\text{Tr } \phi_A^k$ for $A = N(n, r)$ and k in intervals, for data sets (n, r, k) , yield interesting information. Namely,

- a. Many Nakayama algebras are of cyclotomic type;
- b. Not all Nakayama algebras are of cyclotomic type. The case $r = 4$ illustrates this claim:

$N(n, 4)$ is of cyclotomic type for all $0 \leq n \leq 100$ except for $n = 10, 22, 30, 42, 50, 62, 70, 82$ and 90

- c. A canonical algebra C of weight (p_1, \dots, p_t) is strongly accessible if and only if $t = 3$, in that case, C is derived-equivalent to a representation-finite algebra if and only if the weight type does not dominate $(3, 3, 3)$.

IntechOpen

Author details

José-Antonio de la Peña^{1,2}

1 Instituto de Matemáticas, Universidad Nacional Autónoma de México, México

2 El Colegio Nacional, México

*Address all correspondence to: jantdelap@gmail.com

IntechOpen

© 2019 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

References

- [1] Assem I, Simson D, Skowroński A. Elements of the representation theory of associative algebras 1: Techniques of representation theory. In: London Mathematical Society Student Texts 65. Cambridge University Press; 2006
- [2] Boldt A. Methods to determine Coxeter polynomials. Linear Algebra and its Applications. 1995;**230**: 151-164
- [3] Cross GW. Three types of matrix stability. Linear Algebra and its Applications. 1978;**20**:253-263
- [4] Cvetković DM, Doob M, Sachs H. Spectra of Graphs, Theory and Application. Academic Press; 1979
- [5] Dlab V, Ringel CM. Eigenvalues of Coxeter transformations and the Gelfand-Kirillov dimension of the preprojective algebras. Proceedings of the American Mathematical Society. 1981;**83**:228-232
- [6] Ladkani S. On the periodicity of Coxeter transformations and the non-negativity of their Euler forms. Linear Algebra and its Applications. 2008; **428**(4):742-753
- [7] Goodman FM, de la Harpe P, Jones VFR. Coxeter Graphs and Towers of Algebras. New York, Berlin, Heidelberg: Springer-Verlag; 1989
- [8] Happel D. Hochschild Cohomology of Finite Dimensional Algebras. Lect. Notes Math. Vol. 1404. Springer Verlag; 1989. pp. 108-126
- [9] Kronecker L. Zwei Sätze über Gleichungen mit ganzzahligen Coefficienten. Crelle's Journal. 1857; **Ouvres I**:105-108
- [10] Lenzing H. A K-theoretic study of canonical algebras. Conference Proceedings, Canadian Mathematical Society. 1996;**18**:433-454
- [11] Lenzing H, de la Peña JA. Wild canonical algebras. Mathematische Zeitschrift. 1997;**224**:403-425
- [12] Lenzing H, de la Peña JA. Concealed-canonical algebras and separating tubular families. Proceedings of the London Mathematical Society. 1999;**78**:513-540
- [13] Lenzing H, de la Peña JA. Spectral analysis of finite dimensional algebras and singularities. In: Skowroński A, editor. Trends in Representation Theory of Algebras and Related Topics. Zürich: EMS Publishing House; 2008. pp. 541-588
- [14] Lenzing H, de la Peña JA. Extended canonical algebras and Fuchsian singularities. Mathematische Zeitschrift. 2011;**268**(1-2):143-167
- [15] Mahler K. Lectures on transcendental Number Theory. Lecture Notes in Mathematics. Vol. 546. Berlin: Springer Verlag; 1976
- [16] Newman M. Integral Matrices. New York: Academic Press; 1972
- [17] de la Peña JA, Takane M. Spectral properties of Coxeter transformations and applications. Archiv der Mathematik. 1990;**55**:120-134
- [18] de la Peña JA. Algebras whose Coxeter polynomials are product of cyclotomic polynomials. Algebras and Representation Theory. 2014;**17**(3): 905-930
- [19] Ringel CM. Tame Algebras and Integral Quadratic Forms. Lect. Notes Math. Vol. 1099. Berlin-Heidelberg-New York: Springer; 1984
- [20] Ringel CM. The spectral radius of the Coxeter transformations for a

generalized Cartan matrix.
Mathematische Annalen. 1994;**300**:
331-339

[21] Xi C. On wild hereditary algebras
with small growth numbers.
Communications in Algebra. 1990;
18(10):3413-3420

IntechOpen

IntechOpen