

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

185,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Solid-State Fermentation of Cassava Products for Degradation of Anti-Nutritional Value and Enrichment of Nutritional Value

Mohamed Hawashi, Tri Widjaja and Setiyo Gunawan

Abstract

The cassava plant is grown in tropical and subtropical countries, which represents, alongside with its by-products, an important source of food and feed. Hence, this plant has the capacity to promote the economic development of those countries and provide food security. However, cassava has some disadvantages due to the antinutrient compounds produced in its tissues. In addition, the cassava roots have a low protein content. Due to the economic and practical advantages, the solid-state fermentation (SSF) has been used as a cost-effective and efficient processing method to detoxify the cassava products and enrich them in nutrients. This chapter reviews the solid-state fermentation technique of cassava products for the production of valuable components for food and feed applications, microorganisms involved in this process, and key factors used to optimize the SSF process.

Keywords: anti-nutritional value, cassava, nutritional value, processing variables, solid-state fermentation

1. Introduction

Cassava (*Manihot esculenta* Crantz) is grown in tropical and subtropical countries. It is a vital source of food and feed and it can promote economic development and provide food security [1]. Cassava production has been promoted globally by the International Fund for Agricultural Development (IFAD) and the United Nations Food and Agriculture Organization (FAO) to develop cassava strategies [2]. Reports indicate that production rates will reach 300 million tons per year by 2020 [3]. Due to its high drought tolerance, cassava plant cultivation can take place even under critical environmental conditions, with an ideal high yield of approximately 50% for leaves and 6% for roots at plant maturity [4]. Its peel may make up 10–20% of the roots' wet weight [5]. However, cassava has some disadvantages; its tissues contain anti-nutritional compounds and very low protein content [6, 7].

Among all the antinutrients, hydrogen cyanide (HCN) is of great concern, the concentration of which is in cassava and its by-products are much higher than the World Health Organization (WHO) safe limit for human consumption (10 ppm) [8, 9]. Konzo is an irreversible neurological disease associated with intake of HCN [10]. Therefore, a detoxification process is needed to reduce anti-nutritional levels

in order to consume cassava safely. Solid-state fermentation (SSF) has been used as an economical and efficient processing method for enriching and detoxifying cassava and its by-products [11, 12]. Various process parameters such as particle size, moisture content, water activity, pH, the inoculum size, incubation time, concentration of nutrient supplementation, and temperature can affect the microbial growth, enzyme production, and formation of the product during the SSF process [13].

This chapter discusses fermented cassava products through solid-state fermentation for food and feed applications, as well as microorganisms involved in solid-state fermentation and the essential processing variables used to optimize the process.

2. Fermentation processes

Fermentation has been one of the most used technologies to improve the taste and sensory properties of food and continues to be one of the most widely used methods of preserving the food for a length of time [14, 15]. The cassava fermentation process is a strategy to improve nutritional value by enriching protein and detoxifying toxic and anti-nutritional compounds, in particular by reducing toxic cyanogenic glycosides to a safe level of consumption in cassava products as well as reducing post-harvest losses [16–18].

There are two kinds of fermentation, i.e., spontaneous (natural) fermentation and controlled fermentation. For the natural fermentation, the conditions are selected so that to produce the most suitable microorganisms for the production of growth by-products characteristic of a particular type of fermentation [19]. The controlled fermentation is generally used when the natural fermentation is unstable or the bacteria are not able to grow. In this case, specific microbial strains, such as lactic acid bacteria (LAB), yeast, and fungal are isolated, characterized, and preserved for later use as starter cultures [20]. Under optimal growth conditions, these cultures can be used as single or combined starter cultures. As a result, the quality of products and their organoleptic characteristics are well controlled and predictable [20, 21].

However, the fermentation process can be broadly categorized into submerged fermentation (involving soaking in water) and solid-state fermentation (without soaking in water) [22]. The solid-state fermentation (SSF) technique has several advantages over submerged fermentation (SmF). However, the SSF has some constraints. **Table 1** illustrates the advantages and disadvantages of SSF over SmF [23].

2.1 Solid-state fermentation and its application in cassava products

In recent years, the cassava population has developed numerous processing methods (soaking, boiling, drying, and fermentation) [24–26]. SSF is one of the promising processes of enriching protein and detoxifying of cassava products [27–29].

Fermented cassava products by SSF, such as flour, gari, starch, bread, and biomass contain high protein content that can either be consumed by humans or animals, replacing expensive, conventional protein sources in different parts of Latin America, Africa, and Asia [30]. The major fermented cassava products by SSF can be derived from different parts of the cassava plant, such as roots, peels, and leaves.

2.1.1 Cassava roots

Cassava is grown in many developing countries for its roots as a primary source of carbohydrates and ranks third in the developing countries as the leading source

Parameter	Solid-state fermentation	Submerged fermentation
Substrates	Insoluble substrates (starch, cellulose, pectins, lignin)	Soluble substrates (sugars)
Aseptic techniques	Sterilization of steam and non-sterile conditions	Sterilization of heat and aseptic control
Temperature	Difficult temperature control	Easy temperature control
Water	Low water consumption	High water consumption
pH control	Difficult pH control	Easy control of pH
Industrial level	Relatively small scale, newly designed equipment is needed	The industrial level is available
Inoculation	Spore inoculation, batch process	Easy inoculation, continuous process
Contamination	Contamination risk of low-growth fungi	Contamination risk of single strain bacteria
Energy	Low consumption of energy	High consumption of energy
Equipment volumes	Low volumes and low equipment costs	High volumes and high equipment costs
Pollution (effluents)	No volumes of effluents	High volumes of effluents
Concentration/products	100/300 g/L	30–80 g/L

Table 1.
Comparative characteristics of solid-state and submerged fermentations.

of energy in human diets along with rice and wheat [31]. World production of cassava is estimated at 277 million tons of fresh root in 2017 [32]. Cassava root has several advantages compared to other crop roots, including high productivity, resistance to droughts and pests, flexible harvesting age, and it can be kept in the ground until they are needed [33]. However, cassava root also has certain disadvantages; its tissues contain toxic compounds (a cyanogenic glycoside), low protein content (1% fresh root weight), and short shelf life of 1–3 days [34].

Food processing techniques have been used to convert cassava tubers into flour as an alternative way to preserve the roots after harvesting and then further use it for industrial and traditional purposes [35, 36]. Gari and flour are the most popular fermented food products from cassava roots by SSF. In West Africa, approximately 200 million people consume gari [37, 38]. **Figure 1** shows the production of flour and gari under the solid-state fermentation [11].

The purpose of cassava root fermentation is to increase the low protein content from 2% to about 7% or more than the critical crude protein content [39]. To achieve this goal, several solid-state fermentation techniques have been used. Raimbault et al. [40] reported the principle underlying the SSF procedure for the enrichment of cassava flour. This procedure led to the enrichment of crude protein from 1 to 18–20%, which improved between 1700 and 1900% after 30 h of fermentation. Oboh and Elusiyan [41] studied the effect of solid-state fermentation by *R. oryzae* and *S. cerevisiae* on the improvement of nutritional values of cassava flour produced from two different varieties of cassava root. The nutritional contents of cassava flour were assayed before and after 72 h of fermentation. This study has observed that *S. cerevisiae* was more effective than *R. oryzae* in the nutrient enrichment of cassava flour. The results of this study are presented in **Figures 2–5**. Essers et al. [42] investigated the effect of SSF on the degradation of hydrogen cyanide level in cassava root using six fungal strains, namely *Rhizopus stolonifer*, *Rhizopus*




Figure 1.
The production chart of cassava products (flour and gari) under SSF.




Figure 2.
Proximate composition of the cassava flour obtained from cassava varieties of low HCN subjected to SSF.

oryzae, *Mucor racemosus*, *Bacillus sp.* *Geotrichum candidum*, and *Neurospora sitophila*. The reduction in cyanide content was more than 60% after 72 h of fermentation.

In addition, Oboh and Akindahunsi [11] investigated the effect of solid-state fermentation with *S. cerevisiae* on the nutritional and antinutrient contents of cassava products (flour and gari). After 72 h of fermentation, the results revealed that the content of protein and fats in cassava flour increased by 10.9 and 4.5%, respectively. The protein and fat content of fermented gari also improved by 6.3% and 3.0%. In contrast, the content of cyanide in flour and gari decreased to 9.5 and 9.1 (mg/kg),




Figure 3.
Proximate composition of the cassava flour obtained from cassava varieties of medium HCN subjected to SSF.




Figure 4.
Mineral contents of the cassava flour obtained from cassava varieties of low HCN subjected to SSF.

respectively. However, the tannin content, crude fiber, and ash content of the cassava products did not change significantly under SSF.

2.1.2 Cassava peels

Cassava wastes, such as peels and leaves and starch residues make up 25% of the total cassava plant [43]. Cassava peel is the leading waste from the cassava plant, but its use is limited due to the high content of cyanide and fiber as well as low protein and therefore disposed of it after cassava processing into food or other industrial products [44, 45]. Many efforts have been made using SSF techniques to enrich the protein content and degrade the cyanide level of cassava peels for animal feed.

Bayitse et al. [12] studied protein enrichment of cassava residue using *Trichoderma pseudokoningii* under solid-state fermentation for 12 days, urea, and ammonium sulfate was used as a nitrogen source, and the moisture content ranged from 60 to 70%. The result showed an improvement in crude protein content of 12.5% using urea as a nitrogen source, and a moisture content of 70%, as compared




Figure 5.
Mineral contents of the cassava flour obtained from cassava varieties of medium HCN subjected to SSF.

to 8.89 and 6.37% improvement observed with ammonium sulfate as a nitrogen source, and without using nitrogen source. The study observed a decrease in cyanide content, but it did not attribute it to the fermentation effect of *Trichoderma pseudokoningii*, rather it stated that the reduction could have been as a result of the pre-processing of cassava peels.

Iyayi and Losel [43] also evaluated protein improvement of cassava peels using different types of microorganisms and fermentation time (*Saccharomyces cerevisiae*, *Aspergillus niger*, *Rhizomucor miehei*, and *Mucor strictus*). The solid-state fermentation of cassava peels by *S. cerevisiae* produced the highest protein content from 5.6 to 16.74% for 21 days. Also, they reported the maximum fermentation period for the protein enrichment of cassava peel to be from 12 to 15 days, after which no significant change was observed, which is in line with the work reported by Bayitse et al. [12].

Ezekiel and Aworh [13] evaluated the effectiveness of SSF with *Trichoderma viride* on the reduction of cyanide content and enrichment of the crude protein content of cassava peel by optimizing the fermentation conditions such as moisture content, pH, particle size, nitrogen source, and incubation temperature. The optimum SSF conditions were found at the initial moisture content of 60% (v/w), the particle size of 4.00 mm, a pH of 6.0, 30°C of temperature, and ammonium sulfate (10 g N/kg substrate) as nitrogen sources. After 8 days of fermentation, the cyanide content was reduced by 71% and improved the crude protein content from 4.2 to 10.43% at optimized conditions.

In another study by Ruqayyah et al. [45], the application of response surface methodology was used to optimize SSF conditions (moisture content, inoculum size, and pH) with *P. tigrinus* to enrich the crude protein content of cassava peel. A maximum protein content of 89.58 (mg/g) was obtained at 75% (v/w) moisture content, 7% (v/w) inoculum size, and pH of 5.3 with a fermentation time of 15 days. The optimum level resulted in a significant enrichment of the protein content by 55.16%.

Oboh [46] investigated the effect of solid-state fermentation of cassava peel with a mixture of *Saccharomyces cerevisiae* and two strains of lactic acid bacteria, *Lactobacillus delbrueckii* and *Lactobacillus coryniformis* to improve the nutritional value and detoxification of cassava peel. The chemical composition of cassava peel has been analyzed before and after fermentation. The results showed the effective

performance of the SSF technique in removing cyanide by 86% after 7 days of fermentation. On the other hand, the mineral composition of the cassava peel did not change during the fermentation. The results of this study are presented in **Table 2**.

2.1.3 *Cassava leaves*

Cassava leaves are an extremely rich source of proteins, vitamins, and minerals that exceed some of the other green vegetables [47, 48]. The production of cassava leaves is estimated at 10 tons of dry leaves per hectare, which has a similar yield with the roots [49]. Cassava leaves are consumed in most Southeast Asian and African societies, such as Indonesia, Malaysia, Congo, Madagascar, and Nigeria [50, 51]. However, cassava leaves contain both nutritive (33.8–37.4% protein content) and anti-nutritional compounds [301.04–192.47 (mg/100 g) HCN content] [52]. Boiling, soaking, steaming, drying the sun, drying the oven, and cooking are the most common methods for processing cassava leaves in African and Asian countries [53].

The origin of HCN in the cassava leaves is a two-step process [54, 55]. First, the linamarin, a cyanogenic glycoside, which represent 93% of cyanogenic glycosides found in cassava (7% is lotaustralin), is hydrolyzed by linamarase (a beta-glycosidase) into glucose and cyanohydrin. Then, in the second step, the cyanohydrin is decomposed, either enzymatically or not, to HCN and acetone. The nonenzymatic pathway depends on pH. At pH > 6, the HCN is liberated, but at an acidic pH (~5), the process is much lower, and the resulting HCN is therefore relatively lower in concentration. However, this approach did not assure full hydrolysis of cyanogens. The partial breakdown of the leaf cells only partially releases linamarase resulting in only a certain proportion of the cyanogenic compounds being converted to HCN. This implies that a proportion of the cyanogens remain present in the leaves after processing and resulting in the release of HCN directly into the human body upon consumption.

The conventional methods have been proven to be ineffective for lowering the cyanide content in cassava leaves to the safe limit, at the same time causing a significant loss of protein and essential nutrients, which is highly desired from the cassava leaves [56–60]. Hence, the establishment of a universally acceptable method that produces edible leaves with low cyanide level while maintaining

Composition	Fresh	Naturally fermented	Fermented with a mixed culture
Crude protein (%)	8.2 ± 0.1	11.1 ± 0.3	21.5 ± 1.2
Crude fiber (%)	11.7 ± 0.5	6.5 ± 0.5	11.7 ± 0.5
Fat (%)	3.1 ± 0.4	3.5 ± 0.2	2.1 ± 0.1
Ash (%)	6.4 ± 0.4	6.0 ± 0.2	7.2 ± 0.2
Carbohydrate (%)	64.6 ± 0.2	67.3 ± 0.4	51.1 ± 0.4
Moisture (%)	5.1 ± 0.3	5.7 ± 0.2	6.4 ± 0.4
Ca (ppm)	0.03 ± 0.00	0.03 ± 0.00	0.03 ± 0.00
Na (ppm)	00.04 ± 0.00	0.04 ± 0.00	0.04 ± 0.00
Zn (ppm)	0.01 ± 0.00	0.01 ± 0.00	0.01 ± 0.00
K (ppm)	0.05 ± 0.00	0.05 ± 0.00	0.05 ± 0.00
HCN (mg/kg)	45 ± 0.3	24 ± 0.2	6.1 ± 0.4

Table 2.
The effect of fermentation on the chemical composition of cassava peels.

maximum nutritional content is challenging and still far away from being established. Among the efforts made so far, Morales et al. [61] proposed a solid-state fermentation of cassava leaves, reducing the cyanide content while improving the nutritional value of the processed leaves. SSF was performed using *Rhizopus oligosporus*, and babassu mesocarp flour was the substrate used, supplemented by cassava leaf flour. The solid-state fermentation decreased the total cyanide content of the cassava leaves by 94.18%, also SSF increased the quantity and quality of crude protein content by 15%, resulting in the relative nutritional value of 98.18% for food, which is equivalent to casein (100%). Furthermore, Kobawila et al. [62] investigated the effect of alkaline fermentation on the reduction of cyanide level in cassava leaves to produce ntoba mbod. The dominant microflora in the fermentation of the cassava leaves was *Bacillus subtilis*, *Bacillus macerans*, and *Bacillus pumilus*. These bacteria can utilize cyanide acid for their nutrition [63]. Thus, they are responsible for the reduction of the cyanide content in the medium of fermentation (~70% removal). However, the report did not provide the effect of the fermentation process on the protein content of cassava leaves.

One of the essential criteria for the solid-state fermentation is the selection of an appropriate microorganism [64]. Several research works have explored different types of microorganisms mainly fungi, yeasts, and bacteria, as well as different substrates to favor the metabolism of the microorganisms in SSF of cassava products. Examples of microorganisms associated with solid-state fermentation of cassava products for food and feed applications are summarized in **Table 3**.

Microorganism	Substrate	Product	References
<i>Rhizopus oryzae</i>	Cassava root	Gari	[29]
<i>Rhizopus oryzae</i> (TISTR 3052), <i>Rhizopus oryzae</i> (TISTR 3058), <i>Rhizopus delemar</i> (TISTR 3534), and <i>Rhizopus delemar</i> (TISTR 3190)	Cassava flour	Bread	[65]
<i>Panus tigrinus</i> (M609RQY)	Cassava peels	Animal feed	[66]
<i>Aspergillus niger</i> and <i>Panus tigrinus</i>	Cassava peels	Poultry feed	[67]
<i>Rhizopus oryzae</i> and <i>Saccharomyces cerevisiae</i>	Cassava pulp	Animal feed	[68]
<i>Saccharomyces cerevisiae</i> , <i>Aspergillus niger</i> , <i>Rhizomucor miehei</i> , and <i>Mucor strictus</i>	Cassava leaves	Animal feed	[43]
<i>Rhizopus oryzae</i>	Cassava pulp	Lactic acid production	[69]
<i>Lactobacillus plantarum</i> and <i>Rhizopus oryzae</i>	Cassava root	Cellulase production	[70]
<i>Bacillus</i> sp., <i>Mucor racemosus</i> , <i>R. oryzae</i> , <i>Neurospora sitophila</i> <i>R. stolonifer</i> and <i>Geotrichum candidum</i>	Cassava root	Cassava flour	[71]
<i>Rhizopus stolonifer</i> LAU 07	Cassava peel	Feed supplements	[72]
<i>Rhizopus</i> sp.	Cassava starch and leaves	Lactic acid and ethanol productions	[73]

Table 3.
Examples of microorganisms associated with the SSF of cassava products.

3. Environmental factors

The process control of the solid-state fermentation parameters is closely related to the metabolic regulation of microorganisms [74]. Based on the metabolic needs of the fermentation microorganisms, the control of water activity, oxygen content, temperature, and pH are the main solid-state fermentation parameters [23]. In the solid-state fermentation process, the water, gas, and heat caused by the growth microbes are the dominant factors that determine the environmental changes. The environmental factors can affect the microbial growth and formation of the product during the SSF process [75, 13]. Therefore, the physical-chemical parameters must be controlled.

3.1 Water activity and moisture content

The unique feature of solid-state fermentation is that there is almost no free water in the substrate [76]. However, microorganisms can grow depending upon the water activity of the substrate [64, 75]. The growth of fungi and some yeast usually requires a water activity value between 0.6 and 0.7 [77]. In addition to meet the microbial physiological requirements, the water content level plays a decisive role in the variation of the three-phase structure relating to water retention, permeability, and thermal conductivity. The degree of swelling in the SSF system was low at a lower moisture level and hence increased water stress reduces nutrient solubility. On the contrary, the higher level of humidity results in changes in substrates that reduce porosity, thus contributing to stickiness and reduced gas exchange [78, 79]. According to Grover et al. [80], the required moisture content should range between 60 and 80% for an efficient SSF system.

3.2 Temperature

The fermentation temperature affects microbial growth, spore germination, and the formation of product [81]. Heat generation in solid-state fermentation system is more problematic than in liquid fermentation. Due to poor heat conductivity and accumulation of metabolic heat in the material combined with substrate shrinkage and decreased porosity, gas convection is severely impeded. Previous studies showed that the significant resistance to heat transfer in solid-state fermentation was low conduction efficiency [82, 83].

Therefore, moisturizing is a common measure of temperature control. In addition, routine operations (e.g., forced ventilation and jacket cooling) all can solve these problems [84]. The evaporative cooling is one of the main solid-state fermentation temperature control measures [85, 86]. In general, the aeration could reduce the temperature gradient of the medium [23]. The forced ventilation can take away more than 80% of the heat generated from the substrate [84]. From the current investigation, it is difficult to maintain the temperature at an ideal range in SSF system. To reach this aim, the main strategy used in large-scale solid-state fermentation is to combine ventilation and humidity [77].

3.3 Oxygen concentration

The gas environment is a critical factor that significantly affects the relative levels of biomass and the production of an enzyme [23]. Oxygen uptake rate (OUR) and carbon dioxide production (CDPR) can be used to assess the state of the solid-state fermentation process. However, different microorganisms cause these

assessments to vary. Ghildyal et al. [87] studied the impact of the gas concentration gradient on product yield in a tray solid-state fermentation bioreactor. The results showed that the variations of O₂ and CO₂ concentration gradients were visible, which severely affected product yield. The yield decreased when gradient increases. Gowthaman et al. [88] also studied the impact of gas concentration gradient on the product in a packing bed bioreactor. The results showed that the gas concentration gradient could be eliminated and the ability of mass transfer can be enhanced by forced ventilation, which increased enzyme activity.

3.4 pH value

In general, if the initial pH value of the medium is adjusted, the variations of pH value during the solid-state fermentation process need to be considered [89]. During the fermentation process, the pH values change drastically. The reason is that organic acids including citric and lactic are secreted during the fermentation process, which decreases the pH [23]. While the increase in pH was rationalized in terms of organic acid decomposition and protein degradation in the raw materials into amino acids and peptide fractions [90]. The pH values are difficult to determine by conventional detection in SSF due to the low water content of the substrate. Nitrogen-containing inorganic salts (such as urea) are often used as sources of nitrogen to offset the pH variation in the fermentation process [91, 92].

In the study conducted by Ezekiel and Aworh [13] to evaluate the effect of pH on protein enrichment and soluble sugars of cassava peel by *Trichoderma viride*, the fungus was grown in a controlled pH medium of 4.– 6.0 with an incubation time of 8 days. The optimal growth condition was observed at pH 6.0. The protein increased in cassava peels from 230 at pH 4.0 to 270 (mg/gm) at pH 6.0. Also, the sugars yield at pH 5.0 and 6.0 was five times higher compared to pH 4.0. According to the study, the growth rate of the fungi at pH below five was affected by high acidity, leading to reduced bio-conversion of sugars into protein.

4. Conclusions

The results discussed in this chapter highlighted the importance of the SSF technique applied to cassava to improve its nutritional value. The solid-state fermentation using microbial protein is beneficial for the reduction of cyanide contents while the content of protein and other nutrients is increased compared to those obtained by the conventional approaches, i.e., soaking, boiling, and drying. Thus, the SSF technique for processing cassava products is better suited for developing societies and rural communities in the African and Asian countries that do not have easy access to available protein sources.

Acknowledgements

The authors are thankful to the Ministry of Research, Technology and Higher Education of the Republic of Indonesia for its financial support to this project through the grant no. 849/PKS/ITS/2018.

Conflict of interest

The authors declare no conflict of interest.

IntechOpen


IntechOpen

Author details

Mohamed Hawashi, Tri Widjaja and Setiyo Gunawan*
Department of Chemical Engineering, Institut Teknologi Sepuluh Nopember (ITS),
Surabaya, Indonesia

*Address all correspondence to: gunawan@chem-eng.its.ac.id

IntechOpen

© 2019 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

References

- [1] FAO, IFAD. The global cassava development strategy and implementation plan. In: Proceedings of the FAO and IFAD Validation Forum on the Global Cassava Development Strategy; 26-28 April 2000; Rome. Rome: FAO and IFAD; 2001. pp. 13-15
- [2] Howeler RH. Endorsement of the global cassava development strategy. In: Proceedings of the FAO and IFAD Validation Forum on the Global Cassava Development Strategy; 26-28 April 2000; Rome. Rome: FAO and IFAD; 2001. p. 57
- [3] Agustian A. Bioenergy development in the agricultural sector: Potential and constraints of cassava bioenergy development. Analisis Kebijakan Pertanian. 2015;**13**(1):19-38
- [4] Tewe OO, Lutaladio N. Cassava for Livestock Feed in Sub-Saharan Africa. Rome: FAO; 2004. p. 64
- [5] Obadina AO, Oyewole OB, Sanni LO, Abiola SS. Fungal enrichment of cassava peels proteins. African Journal of Biotechnology. 2006;**5**(3):302-304. DOI: 10.5897/AJB05.360
- [6] Gunawan S, Widjaja T, Zullaikah S, Istianah N, Aparamarta HW, Prasetyoko D, et al. Effect of fermenting cassava with *Lactobacillus plantarum*, *Saccharomyces cerevisiae*, and *Rhizopus oryzae* on the chemical composition of their flour. International Food Research Journal. 2015;**22**(3):1280-1287
- [7] Hawashi M, Ningsih TS, Cahyani SBT, Widjaja KT, Gunawan S. Optimization of the fermentation time and bacteria cell concentration in the starter culture for cyanide acid removal from wild cassava (*Manihot glaziovii*). MATEC Web of Conferences. 2018;**156**:01004. DOI: 10.1051/mateconf/201815601004
- [8] Codex Alimentarius Commission. Codex Alimentarius. Rome: Food and Agriculture Organization; 1992
- [9] Hadiyat MA, Wahyudi RD. Integrating steepest ascent for the Taguchi experiment: A simulation study. International Journal of Technology. 2013;**3**:280-287. DOI: 10.14716/ijtech.v4i3.132
- [10] Bradbury JH. Simple wetting method to reduce cyanogen content of cassava flour. Journal of Food Composition and Analysis. 2006;**19**(4):388-393. DOI: 10.1016/J.JFCA.2005.04.012
- [11] Oboh G, Akindahunsi AA. Biochemical changes in cassava products (flour & gari) subjected to *Saccharomyces cerevisiae* solid media fermentation. Food Chemistry. 2003;**82**(4):599-602. DOI: 10.1016/S0308-8146(03)00016-5
- [12] Bayitse R, Hou X, Laryea G, Bjerre AB. Protein enrichment of cassava residue using *Trichoderma pseudokoningii* (ATCC 26801). AMB Express. 2015;**5**(1):80. DOI: 10.1186/s13568-015-0166-8
- [13] Ezekiel OO, Aworh OC. Solid state fermentation of cassava peel with *Trichoderma viride* (ATCC 36316) for protein enrichment. World Academy of Science, Engineering and Technology. 2013;**7**(3):6892-6991
- [14] Motarjemi Y. Impact of small scale fermentation technology on food safety in developing countries. International Journal of Food Microbiology. 2002;**75**(3):213-229. DOI: 10.1016/S0168-1605(01)00709-7
- [15] Smid EJ, Hugenholtz J. Functional genomics for food fermentation processes. Annual Review of

Food Science and Technology.
 2010;**1**:497-519. DOI: 10.1146/annurev.
 food.102308.124143

[16] Caplice E, Fitzgerald GF. Food fermentations: Role of microorganisms in food production and preservation. *International Journal of Food Microbiology*. 1999;**50**(1-2):131-149. DOI: 10.1016/S0168-1605(99)00082-3

[17] Kostinek M, Specht I, Edward VA, Schillinger U, Hertel C, Holzapfel WH, et al. Diversity and technological properties of predominant lactic acid bacteria from fermented cassava used for the preparation of Gari, a traditional African food. *Systematic and Applied Microbiology*. 2005;**28**(6):527-540. DOI: 10.1016/j.syapm.2005.03.001

[18] Achi OK, Akomas NS. Comparative assessment of fermentation techniques in the processing of fufu, a traditional fermented cassava product. *Pakistan Journal of Nutrition*. 2006;**5**(3):224-229

[19] Stiles ME, Holzapfel WH. Lactic acid bacteria of foods and their current taxonomy. *International Journal of Food Microbiology*. 1997;**36**(1):1-29. DOI: 10.1016/S0168-1605(96)01233-0

[20] Zulu RM, Dillon VM, Owens JD. Munkoyo beverage, a traditional Zambian fermented maize gruel using *Rhynchosia* root as amylase source. *International Journal of Food Microbiology*. 1997;**34**(3):249-258. DOI: 10.1016/S0168-1605(96)01195-6

[21] Oguntoyinbo FA, Cho GS, Trierweiler B, Kabisch J, Rösch N, Neve H, et al. Fermentation of African kale (*Brassica carinata*) using *L. plantarum* BFE 5092 and *L. fermentum* BFE 6620 starter strains. *International Journal of Food Microbiology*. 2016;**238**:103-112. DOI: 10.1016/j.ijfoodmicro.2016.08.030

[22] Ray RC, Swain MR. Bio-ethanol, bioplastics and other fermented industrial products from cassava

starch and flour. In: Colleen MP, editor. *Cassava: Farming, Uses and Economic Impact*. Hauppauge: Nova; 2011. pp. 1-32

[23] Raimbault M. General and microbiological aspects of solid substrate fermentation. *Electronic Journal of Biotechnology*. 1998;**1**(3):26-27. DOI: 10.4067/S0717-34581998000300007

[24] Nambisan B, Sundaresan S. Effect of processing on the cyanoglucoside content of cassava. *Journal of the Science of Food and Agriculture*. 1985;**36**(11):1197-1203. DOI: 10.1002/jsfa.2740361126

[25] Bradbury JH, Denton IC. Rapid wetting method to reduce cyanogen content of cassava flour. *Food Chemistry*. 2010;**121**(2):591-594. DOI: 10.1016/j.foodchem.2009.12.053

[26] Ezekiel OO, Aworh OC, Blaschek HP, Ezeji TC. Protein enrichment of cassava peel by submerged fermentation with *Trichoderma viride* (ATCC 36316). *African Journal of Biotechnology*. 2010;**9**(2):187-194. DOI: 10.5897/AJB09.620

[27] Reade AE, Gregory KF. High-temperature production of protein-enriched feed from cassava by fungi. *Applied and Environmental Microbiology*. 1975;**30**(6):897-904

[28] Vlavonou BM. Cassava processing technologies in Africa. In: *Proceedings of the Interregional Experts' Group Meeting on the Exchange of Technologies for Cassava Processing Equipment and Food Products*; 13-19 April 1988; Ibadan, Nigeria. New York: UNICEF House; 1988. pp. 19-25

[29] Akindahunsi AA, Oboh G, Oshodi AA. Effect of fermenting cassava with *Rhizopus oryzae* on the chemical

composition of its flour and Gari products. *Rivista Italiana delle Sostanze Grasse*. 1999;**76**:437-440

[30] Behera SS, Ray RC. Microbial linamarase in cassava fermentation. In: Ramesh RC, Christina MS, editors. *Microbial Enzyme Technology in Food Applications*. Boca Raton: CRC Press; 2017. pp. 337-346

[31] Food and Agricultural Organization. *Food Outlook: Global Market Analysis*. Rome: FAO; 2009. pp. 23-27

[32] Food and Agricultural Organization. *Food Outlook: Biannual Report on Global Food Markets*. Rome: FAO; 2018

[33] Gunawan S, Istighfarah Z, Aparamarta HW, Syarifah F, Dwitasari I. Utilization of modified cassava flour and its by-products. In: Klein C, editor. *Handbook on Cassava*. New York: Nova Science Publisher; 2017. pp. 271-295

[34] Westby A. Cassava utilization, storage and small-scale processing. In: Hillocks RJ, Thresh JM, Bellotti AC, editors. *Cassava: Biology, Production and Utilization*. New York: CABI Publishing; 2002. pp. 281-300

[35] Defloor I, Nys M, Delcour JA. Wheat starch, cassava starch, and cassava flour impairment of the breadmaking potential of wheat flour. *Cereal Chemistry*. 1993;**70**(5):526-530

[36] Dakwa S, Sakyi-Dawson E, Diako C, Annan NT, Amoa-Awua WK. Effect of boiling and roasting on the fermentation of soybeans into dawadawa (*soy-dawadawa*). *International Journal of Food Microbiology*. 2005;**104**(1):69-82. DOI: 10.1016/j.ijfoodmicro.2005.02.006

[37] Yao AA, Dortu C, Egounlety M, Pinto C, Edward VA, Huch M, et al. Production of freeze-dried lactic acid bacteria starter culture

for cassava fermentation into gari. *African Journal of Biotechnology*. 2009;**8**(19):4996-5004

[38] Udoro EO, Kehinde AT, Olanukanmi SG, Charles TA. Studies on the physicochemical, functional and sensory properties of gari processed from dried cassava chips. *Journal of Food Processing & Technology*. 2014;**5**(1):293. DOI: 10.4172/2157-7110.1000293

[39] Aro SO. Improvement in the nutritive quality of cassava and its by-products through microbial fermentation. *African Journal of Biotechnology*. 2008;**7**(25):4789-4797. DOI: 10.5897/AJB08.1005

[40] Raimbault M, Deschamps F, Meyer F, Senez JC. Direct protein enrichment of starchy products by fungal solid fermentation. In: *Proceedings of the 5th International Conference on Global Impacts of Applied Microbiology*; 21-26 November 1977. Bangkok; 1977

[41] Oboh G, Elusiyan CA. Changes in the nutrient and anti-nutrient content of micro-fungi fermented cassava flour produced from low-and medium-cyanide variety of cassava tubers. *African Journal of Biotechnology*. 2007;**6**(18):2150-2157. DOI: 10.5897/AJB2007.000-2336

[42] Essers AA, Jurgens CM, Nout MR. Contribution of selected fungi to the reduction of cyanogen levels during solid substrate fermentation of cassava. *International Journal of Food Microbiology*. 1995;**26**(2):251-257. DOI: 10.1016/0168-1605(94)00116-N

[43] Iyayi EA, Losel DM. Protein enrichment of cassava by-products through solid state fermentation by fungi. *Journal of Food Technology in Africa*. 2001;**6**(4):116-118. DOI: 10.4314/jfta.v6i4.19301

[44] Iyayi EA, Tewe OO. Effect of protein deficiency on utilization

of cassava peel by growing pigs.
 In: Proceedings of the IITA/ILCA/
 University of Ibadan Workshop on
 the Potential Utilisation of Cassava
 as Livestock Feed in Africa; 14-18
 November 1988. Ibadan: IITA; 1988.
 pp. 54-57

[45] Ruqayyah TI, Jamal P, Alam
 MZ, Mirghani ME, Jaswir I, Ramli
 N. Application of response surface
 methodology for protein enrichment
 of cassava peel as animal feed by
 the white-rot fungus *Panus tigrinus*
 M609RQY. Food Hydrocolloids.
 2014;**42**(15):298-303. DOI: 10.1016/j.
 foodhyd.2014.04.027

[46] Oboh G. Nutrient enrichment of
 cassava peels using a mixed culture of
Saccharomyces cerevisiae and *Lactobacillus*
spp solid media fermentation
 techniques. Electronic Journal of
 Biotechnology. 2006;**9**(1):46-49. DOI:
 10.4067/S0717-34582006000100007

[47] Montagnac JA, Davis CR,
 Tanumihardjo SA. Nutritional value
 of cassava for use as a staple food and
 recent advances for improvement.
 Comprehensive Reviews in Food Science
 and Food Safety. 2009;**8**(3):181-194.
 DOI: 10.1111/j.1541-4337.2009.00077.x

[48] Wargiono J, Richana N, Hidajat A.
 Contribution of cassava leaves used
 as a vegetable to improved human
 nutrition in Indonesia. In: Proceedings
 of the Seventh Regional Workshop on
 Cassava Research and Development
 in Asia: Exploring New Opportunities
 for an Acent Crop; 28 October – 01
 November 2002. Bangkok: CIAT; 2007.
 pp. 466-471

[49] Morgan NK, Choct M. Cassava:
 Nutrient composition and nutritive
 value in poultry diets. Animal Nutrition.
 2016;**2**(4):253-261. DOI: 10.1016/j.
 aninu.2016.08.010

[50] Gidamis AB, O'Brien GM,
 Poulter NH. Cassava detoxification of

traditional Tanzanian cassava foods.
 International Journal of Food Science
 and Technology. 1993;**28**(2):211-218.
 DOI: 10.1111/j.1365-2621.1993.
 tb01266.x

[51] Balagopalan C. Cassava utilization in
 food, feed and industry. In: Hillocks RJ,
 Thresh JM, Bellotti AC, editors. Cassava:
 Biology, Production and Utilization.
 New York: CABI Publishing; 2002.
 pp. 301-318

[52] Achidi AU, Ajayi OA, Maziya-
 Dixon BU, Bokanga M. The effect of
 processing on the nutrient content of
 cassava (*Manihot esculenta* Crantz)
 leaves. Journal of Food Processing &
 Preservation. 2008;**32**(3):486-502. DOI:
 10.1111/j.1745-4549.2007.00165.x

[53] Fasuyi AO. Nutrient composition
 and processing effects on cassava
 leaf (*Manihot esculenta*, Crantz)
 antinutrients. Pakistan Journal of
 Nutrition. 2005;**4**(1):37-42

[54] Vetter J. Plant cyanogenic
 glycosides. Toxicon. 2000;**38**(1):11-36.
 DOI: 10.1016/S0041-0101(99)00128-2

[55] Montagnac JA, Davis CR,
 Tanumihardjo SA. Processing
 techniques to reduce toxicity and
 antinutrients of cassava for use
 as a staple food. Comprehensive
 Reviews in Food Science and Food
 Safety. 2009;**8**(1):17-27. DOI:
 10.1111/j.1541-4337.2008.00064.x

[56] Padmaja G, Steinkraus
 KH. Cyanide detoxification in
 cassava for food and feed uses.
 Critical Reviews in Food Science and
 Nutrition. 1995;**35**(4):299-339. DOI:
 10.1080/10408399509527703

[57] Ngudi DD, Kuo YH, Lambein F.
 Amino acid profiles and protein quality
 of cooked cassava leaves or 'saka-saka'.
 Journal of the Science of Food and
 Agriculture. 2003;**83**(6):529-534. DOI:
 10.1002/jsfa.1373

- [58] Ngudi DD, Kuo YH, Lambein F. Cassava cyanogens and free amino acids in raw and cooked leaves. *Food and Chemical Toxicology*. 2003;**41**(8):1193-1197. DOI: 10.1016/S0278-6915(03)00111-X
- [59] Bradbury JH, Denton IC. Mild method for removal of cyanogens from cassava leaves with retention of vitamins and protein. *Food Chemistry*. 2014;**1**(158):417-420. DOI: 10.1016/j.foodchem.2014.02.132
- [60] Latif S, Müller J. Potential of cassava leaves in human nutrition: A review. *Trends in Food Science and Technology*. 2015;**44**(2):147-158. DOI: 10.1016/j.tifs.2015.04.006
- [61] Morales EM, Domingos RN, Angelis DF. Improvement of protein bioavailability by solid-state fermentation of babassu mesocarp flour and cassava leaves. *Waste and Biomass Valorization*. 2018;**9**(4):581-590. DOI: 10.1007/s12649-016-9759-y
- [62] Kobawila SC, Louembe D, Keleke S, Hounhouigan J, Gamba C. Reduction of the cyanide content during fermentation of cassava roots and leaves to produce bikedi and ntoba mbodi, two food products from Congo. *African Journal of Biotechnology*. 2005;**4**(7):689-696. DOI: 10.5897/AJB2005.000-3128
- [63] Knowles CJ. Microorganisms and cyanide. *Bacteriological Reviews*. 1976;**40**(3):652-680
- [64] Pandey A. Recent process developments in solid-state fermentation. *Process Biochemistry*. 1992;**27**(2):109-117. DOI: 10.1016/0032-9592(92)80017-W
- [65] Begum R, Rakshit SK, Rahman SM. Protein fortification and use of cassava flour for bread formulation. *International Journal of Food Properties*. 2011;**14**(1):185-198. DOI: 10.1080/10942910903160406
- [66] Jamal P, Tijani RI, Alam MZ, Mirghani ME. Effect of operational parameters on solid-state fermentation of cassava peel to an enriched animal feed. *Journal of Applied Sciences*. 2012;**12**(11):1166-1170. DOI: 10.3923/jas.2012.1166.1170
- [67] Purwadaria T. Solid substrate fermentation of cassava Peel for poultry feed ingredient. *WARTAZOA. Indonesian Bulletin of Animal and Veterinary Sciences*. 2013;**23**(1):15-22. DOI: 10.14334/wartazoa.v23i1.955
- [68] Thongkratok R, Khempaka S, Molee W. Protein enrichment of cassava pulp using microorganisms' fermentation techniques for use as an alternative animal feedstuff. *Journal of Animal and Veterinary Advances*. 2010;**9**(22):2859-2862. DOI: 10.3923/javaa.2010.2859.2862
- [69] Phrueksawan P, Kulpreecha S, Sooksai S, Thongchul N. Direct fermentation of L (+)-lactic acid from cassava pulp by solid-state culture of *Rhizopus oryzae*. *Bioprocess and Biosystems Engineering*. 2012;**35**(8):1429-1436. DOI: 10.1007/s00449-012-0731-3
- [70] Roger DD, Jean-Justin EN, Francois-Xavier ET. Cassava solid-state fermentation with a starter culture of *Lactobacillus plantarum* and *Rhizopus oryzae* for cellulase production. *African Journal of Microbiology Research*. 2011;**5**(27):4866-4872. DOI: 10.5897/AJMR11.790
- [71] Essers AJ, Bennik MH, Nout MJ. Mechanisms of increased linamarin degradation during solid-substrate fermentation of cassava. *World Journal of Microbiology and Biotechnology*. 1995;**11**(3):266-270. DOI: 10.1007/BF00367096
- [72] Lateef A, Oloke JK, Kana EG, Oyeniyi SO, Onifade OR, Oyeleye AO, et al. Improving the quality of

agro-wastes by solid-state fermentation: Enhanced antioxidant activities and nutritional qualities. *World Journal of Microbiology and Biotechnology*. 2008;**24**(10):2369-2374. DOI: 10.1007/s11274-008-9749-8

[73] Azmi AS, Yusuf N, Jimat DN, Puad NI. Co-production of lactic acid and ethanol using *Rhizopus Sp.* from hydrolyzed inedible cassava starch and leaves. *IIUM Engineering Journal*. 2016;**17**(2):1-10. DOI: 10.31436/iiumej.v17i2.610

[74] Chen HZ, Li ZH. Bioreactor engineering. *Chinese Journal of Process Biotechnology*. 1998;**18**:46-49

[75] Nagel FJ. Process Control of Solid-State Fermentation: Simultaneous Control of Temperature and Moisture Content [Thesis]. Wageningen: Wageningen University; 2002

[76] Pandey A. Solid-state fermentation. *Biochemical Engineering Journal*. 2003;**13**(2-3):81-84. DOI: 10.1016/S1369-703X(02)00121-3

[77] Gervais P, Molin P. The role of water in solid-state fermentation. *Biochemical Engineering Journal*. 2003;**13**(2-3):85-101. DOI: 10.1016/S1369-703X(02)00122-5

[78] Mahanta N, Gupta A, Khare SK. Production of protease and lipase by solvent tolerant *Pseudomonas aeruginosa* PseA in solid-state fermentation using *Jatropha curcas* seed cake as substrate. *Bioresource Technology*. 2008;**99**(6):1729-1735. DOI: 10.1016/j.biortech.2007.03.046

[79] Mustafa SR, Husaini A, Hipolito CN, Hussain H, Suhaili N, Roslan HA. Application of response surface methodology for optimizing process parameters in the production of amylase by *Aspergillus flavus* NSH9 under solid state fermentation. *Brazilian Archives of Biology and Technology*. 2016;**59**. DOI: 10.1590/1678-4324-2016150632

[80] Grover A, Maninder A, Sarao LK. Production of fungal amylase and cellulase enzymes via solid state fermentation using *Aspergillus oryzae* and *Trichoderma reesei*. *International Journal of Advancements in Research & Technology*. 2013;**2**(8):108-124

[81] Lonsane BK, Ghildyal NP, Budiatman S, Ramakrishna SV. Engineering aspects of solid-state fermentation. *Enzyme and Microbial Technology*. 1985;**7**(6):258-265. DOI: 10.1016/0141-0229(85)90083-3

[82] Saucedo-Castañeda G, Gutierrez-Rojas M, Bacquet G, Raimbault M, Viniegra-González G. Heat transfer simulation in solid substrate fermentation. *Biotechnology and Bioengineering*. 1990;**35**(8):802-808. DOI: 10.1002/bit.260350808

[83] González-Blanco P, Saucedo-Castañeda G, Viniegra-González G. Protein enrichment of sugar cane by-products using solid-state cultures of *Aspergillus terreus*. *Journal of Fermentation and Bioengineering*. 1990;**70**(5):351-354. DOI: 10.1016/0922-338X(90)90150-U

[84] Manpreet S, Sawraj S, Sachin D, Pankaj S, Banerjee UC. Influence of process parameters on the production of metabolites in solid-state fermentation. *Malaysian Journal of Microbiology*. 2005;**1**(2):1-9

[85] Durand A, Arnous P, de Chardin OT, Chereau D, Boquien C. Protein enrichment of sugar beet pulp by solid-state fermentation. In: Ferranti MP, Fiechter A, editors. *Production and Feeding of Single-Cell Protein*. London: Applied Science Publisher; 1983. pp. 120-123

[86] Grajek W. Cooling aspects of solid-state cultures of mesophilic and thermophilic fungi. *Journal of Fermentation Technology*.

1988;**66**(6):675-679. DOI:
10.1016/0385-6380(88)90072-6

[87] Ghildyal NP, Gowthaman MK, Rao KR, Karanth NG. Interaction of transport resistances with biochemical reaction in packed-bed solid-state fermentors: Effect of temperature gradients. *Enzyme and Microbial Technology*. 1994;**16**(3):253-257. DOI: 10.1016/0141-0229(94)90051-5

[88] Gowthaman MK, Ghildyal NP, Rao KR, Karanth NG. Interaction of transport resistances with biochemical reaction in packed bed solid state fermenters: The effect of gaseous concentration gradients. *Journal of Chemical Technology and Biotechnology*. 1993;**56**(3):233-239. DOI: 10.1002/jctb.280560303

[89] Mitchell DA, Do DD, Greenfield PF, Doelle HW. A semi-mechanistic mathematical model for growth of *Rhizopus oligosporus* in a model solid-state fermentation system. *Biotechnology and Bioengineering*. 1991;**38**(4):353-362. DOI: 10.1002/bit.260380405

[90] Awasthi MK, Pandey AK, Bundela PS, Khan J. Co-composting of organic fraction of municipal solid waste mixed with different bulking waste: Characterization of physicochemical parameters and microbial enzymatic dynamic. *Bioresource Technology*. 2015;**182**:200-207. DOI: 10.1016/j.biortech.2015.01.104

[91] Raimbault M. Fermentation en milieu solid: Croissance de champignons filamentous sure substrate amylacé. Paris: ORSTOM; 1981. p. 291

[92] Correia R, Magalhaes M, Macêdo G. Protein enrichment of pineapple waste with *Saccharomyces cerevisiae* by solid state bioprocessing. *Journal of Scientific and Industrial Research*. 2007;**66**(3):259-262