We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

ALAD (δ -aminolevulinic Acid Dehydratase) as Biosensor for Pb Contamination

Muhsin Konuk, İbrahim Hakkı Cigerci and Safiye Elif Korcan,
Afyon Kocatepe University, Faculty of Science & Literatures, Biology Department,
Afyonkarahisar
Turkey

1. Introduction

Heavy metals are defined as those having a specific density of more than 5 g/cm³. The main threats to human health are associated with exposure to lead, cadmium, mercury and arsenic (arsenic is a metalloid, but is usually classified as a heavy metal) (Järup 2003). Heavy metals have been used in many different areas for thousands of years. Lead has been used for at least 5000 years, early applications including building materials, pigments for glazing ceramics, and pipes for transporting water. In ancient Rome, lead acetate was used to sweeten old wine, and some Romans might have consumed as much as a gram of lead a day. Mercury was allegedly used by the Romans as a salve to alleviate teething pain in infants, and was later (from the 1300s to the late 1800s) employed as a remedy for syphilis. Although adverse health effects of heavy metals have been known for a long time, exposure to heavy metals continues and is even increasing in some areas. Since the middle of the 19th century, production of heavy metals increased steeply for more than 100 years, with concomitant emissions to the environment. Emissions of heavy metals to the environment occur *via* a wide range of processes and pathways, including to the air, waters, and soil (Järup 2003).

Lead, mercury, cadmium and arsenic form a significant potential threat to human health, both occupational and environmental (Hu 2000). Over the past few decades, heavy metal contamination of aquatic system has attracted the attention of a number of researchers in all over the world. Many industrial and agricultural processes have contributed to the contamination of fresh water systems thereby causing adverse effects on aquatic biota and human health (Wang 2002, Dautremepuits 2004). The fact that heavy metals cannot be destroyed through biological degradation and have the ability to accumulate in the environment make these toxicants deleterious to the aquatic environment and consequently to humans who depend on aquatic products as sources of food. Heavy metals do mainly accumulate in the tissues of aquatic animals and can be of public health concern to human beings (Kalay 1999, Ashraf 2005).

In fact, there are several heavy metals known to be carcinogens, including arsenic, chromium and nickel. Many of the toxic effects of metals, including carcinogenicity, can be modified by concurrent exposure to other metals. The emphasis now is on testing the effects of mixtures of chemicals to simulate actual environmental conditions. Researches have already demonstrated links between lead and polychlorinated biphenyl exposure and

Source: Intelligent and Biosensors, Book edited by: Vernon S. Somerset,
 ISBN 978-953-7619-58-9, pp. 386, January 2010, INTECH, Croatia, downloaded from SCIYO.COM

neurodevelopmental effects. It is now frightened that neurobehavioural and neurodevelopmental deterioration may take place in the next generations (Tang et al. 1999, Winneke et al. 2002, Stein et al. 2002, Yang et al. 2003).

Reports which link autoimmune diseases to environmental factors have appeared in recent literatures (Molina & Ehrenfeld 2003, Dooley & Hogan 2003). In many exposed populations, some individuals are extra sensitive while some extra tolerant. Especially, children are more susceptible to mercury and lead exposures (Needleman et al. 1990). In order to comprehend the basis of these special high or low risk groups, regarding comparative anatomy, physiology, metabolism and genetics, more in depth studies in toxicodynamics and toxicokinetics are needed. It is assumed that state-of-the-art technology where occupational safety, product safety, environmental quality maintenance and accident prevention are in-built under good manufacturing practices. As a result of these studies, conventional problems in occupational toxicology are expected to decrease. Exposure to chemicals already loaded in the environment will still be responsible for delayed effects and indirect exposure. Molecular epidemiology, armed with designed molecular probes and noninvasive diagnostics will become a leading component in risk assessment and health management in future.

2. Lead and lead pollution

Lead (Pb) is one of the most widely used metals in industries and almost in all over the world exposure to Pb continues to be a common problem. Batteries, paints and pigments, plastic, ceramic, secondary foundries and welding are the most important occupational settings. The general population may get exposed to Pb due to food and water contamination, and air pollution caused by industrial emission and fuel containing Pb compounds.

Due to environmental ubiquity and persistence of Pb, its accumulation in organisms and biomass throughout the trophic chain, imply a continuous exposure. This metal can cause mortality in cases of acute poisoning or can indirectly affect the populations by altering reproductive success, behavior, immune response, and physiology in cases of chronic exposure (Mazliah et al. 1989, Burger 1995, Burger & Gochfeld 2000b, Fair & Ricklefs 2002).

Sometimes wild birds might be exposed to very high metal levels, for example, at waste disposal sites or through the ingestion of lead-shot pellets. Such acute poisonings are easily diagnosed, although longer-term effects are difficult to assess. Lead in blood is a good indicator of newly exposure, while chronic exposure can be estimated when concentrations in accumulator tissue(s) are available. Although studies on dead animals provide useful information, ethical, legal, and scientific reasons indicate the need for other types of more easily available samples (feathers, eggs, excrements, regurgitated food, etc), which enable us to estimate exposure conditions (Burger & Gochfeld 2000a, Dauwe et al. 2000). In recent years the usefulness of feathers as a biomarker of heavy metal exposure has been investigated. Results were very satisfying to monitor mercury and lead levels, while contradictory results have been obtained for cadmium (Furness 1993).

In spite of significant reductions in use, most notably in paint production and as a fuel additive, Pb continues to enter the environment primarily by anthropogenic resources, retaining its status as a priority pollutant (USEPA 2006).

As the focus has turned towards remediation concerning to prevent the human exposure, much is still needed in the way of determining appropriate measures to monitor and protect

the aquatic environment. Usually, water quality criteria (WQC) continue to rely principally on water hardness (i.e. Ca^{2+}) despite growing evidence that other chemical parameters [e.g. pH, salinity and dissolved organic carbon (DOC)], which may vary greatly on a local basis, also strongly influence Pb toxicity (Macdonald et al. 2002, Grosell et al. 2006).

Efforts to improve WQC for metals have resulted in several toxicity models which designed to encircle the influences of all major water chemistry parameters. The most widely accepted model, the biotic ligand model (BLM), is currently used by the USEPA to set WQC for copper. In core, the BLM is responsible for site-specific water conditions considering the competitive effects from other cations and complexity with organic/inorganic agents that prevent the metal from interacting with the site of toxic action (Paquin et al. 2002).

There has been no demonstrated biological need for Pb. Therefore, its uptake and toxicity is likely mediated through imitating of other cations (Ballatori 2002). The most reasonable candidate is Ca^{2+} . There are strong evidences that Pb acts as a Ca^{2+} antagonist (Busselberg et al. 1991, Rogers & Wood 2004). However, the identification of a specific ligand for Pb remains elusive. As in mammals, the principal effects of chronic Pb exposure to fish are presumably hematological (Hodson et al. 1978), neurological (Davies et al. 1976) and renal (Patel et al. 2006) defects. Some studies have also examined reproduction and behavioral effects (Holcombe et al. 1976, Weber 1993).

These metals and other toxicants are commonly present as mixtures in the environment. Genomic approaches are well suited to locate such problems by filling in where more conventional methods prove insufficient to precise key environmental stressors or elucidate the contributions and additive effects from multiple toxicants. Additionally, microarrays give opportunities not only for establishing the molecular basis of toxicity, but potential for gaining insights into modes of action and higher order effects. Thus, defining toxicant-specific mechanisms that link signature gene transcript profiles to chronic effects would greatly help in monitoring and diagnosing water quality and also prioritizing higher ranked tests in ecological risk assessment. The significance of genomics in this regard was recently referred by the USEPA (Dix et al. 2006).

3. Lead toxicity and δ -ALAD as biosensor for lead toxicity

Pb is a natural component of ecosystems with no known biological role and is highly toxic. Its toxicity originates from its ability to mimic biologically important metals and to produce membrane damage through lipid peroxidation. Most Pb poisoning symptoms are thought to occur by interfering with an essential enzyme, δ -aminolevulinic acid dehydratase (ALAD), the activity of which is markedly inhibited by Pb. This is in total agreement with almost all studies and confirms the toxic effects of Pb for the taxa including bacteria, fishes, amphibians, reptiles, birds, mammals and humanbeings.

A biosensor is a device for the detection of an analyte that combines a biological component with a physicochemical detector component (<http://en.wikipedia.org/wiki/Biosensor>). It consists of 3 parts:

- The *sensitive biological element* biological material (eg. tissue, microorganisms, organelles, cell receptors, enzymes, antibodies, nucleic acids, etc.), a biologically derived material or biomimic] the sensitive elements can be created by biological engineering.
- the *transducer* or the *detector element* (works in a physicochemical way; optical, piezoelectric, electrochemical, etc.) that transforms the signal resulting from the

interaction of the analyte with the biological element into another signal (i.e., transducers) that can be more easily measured and quantified;

- Associated electronics or signal processors that are primarily responsible for the display of the results in a user-friendly way (Cavalcanti 2008).

An enzyme, an analytical device, can be used as a biosensor. This combines an enzyme with a transducer to produce a signal proportional to target analyte concentration. This signal can result from a change in proton concentration, release or uptake of gases, such as ammonia or oxygen, light emission, absorption or reflectance, heat emission, and so forth, brought about by the reaction catalyzed by the enzyme used. The transducer converts this signal into a measurable response, such as current, potential, temperature change, or absorption of light through electrochemical, thermal, or optical means. This signal can be further amplified, processed, or stored for later analysis. Because of their specificity and catalytic properties, enzymes have found widespread use as sensing elements in biosensors. Since the development of the first enzyme-based sensor by Clark & Lyons (1962), who immobilized glucose oxidase on an oxygen-sensing electrode to measure glucose, there has been an impressive proliferation of applications involving a wide variety of substrates. A variety of the enzymes belonging to classes of oxido-reductases, hydrolases, and lyases have been integrated with different transducers to construct biosensors for applications in health care, veterinary medicine, food industry, environmental monitoring, and defense (Guilbault 1984).

Enzyme biosensors have been widely used in clinical and food analysis, where analytes represent natural substrates of the enzymes employed. At difference, in environmental analysis, pollutants (e.g. pesticides, heavy metals, etc.) are generally detected as monitoring the inhibition of enzymatic activity caused by those toxic materials. The reduced specificity of inhibition phenomenon makes only possible the determination of such parameters, i.e. total concentration of the substances belonging to a certain class. On the other hand, it is expected that biosensor detects only contaminants, which are actually harmful to life.

Enzyme biosensor has got some advantages and same disadvantages. Their advantages are: being more specific than cell based sensors; faster responds due to shorter diffusion a path (no cell walls). Disadvantages: being more expensive to produce; and enzymes are often unstable. Additionally, many enzymes need cofactors for the detection of substances ([http://www.rpi.edu/dept/chem-eng/Biotech Environ/BIOSEN/enzbio.htm](http://www.rpi.edu/dept/chem-eng/Biotech%20Environ/BIOSEN/enzbio.htm)).

Biosensors for heavy metals have been mainly developed in environmental analysis in water (Evtugyn et al. 1999). As far as enzyme biosensors for heavy metal determination are concerned, a certain number of papers have appeared, reporting the use of different enzymes and biosensor configurations/transducers (Ciucu et al. 2001, Compagnone et al. 1995 and Donlan et al. 1989b, Starodub et al. 1999, Vel Krawczyc et al. 2000, Pirvutoiu et al. 2001, and Dzyadevych et al. 2003). In several cases the inhibition of enzymes by heavy metals is reversible, even if for rapid restoration of enzymatic activity the use of strong ligands, like EDTA, is required.

Most of the heavy metals bind to the sulfhydryl groups, thus inhibiting enzymic activity, disrupting cellular transport and causing changes in protein functions. The toxicity of heavy metals includes the blocking of functional groups of important molecules, e.g. enzymes, polynucleotides, transport systems for essential nutrients and ions, and substitution of essential ions from cellular sites.

During the last decades there was an increasing interest to investigate other sublethal endpoints, especially in relation to those biochemical responses that may be considered as

early biosensors of contamination (Huggett et al. 1992) Among them, the inhibition of the enzyme δ -aminolevulinic acid dehydratase (ALAD, E.C. 4.2.1.24) is recognized as a useful biomarker of Pb exposure and effect, both in humans and other animal species (Rand 1995, Timbrell 2000).

Endogenous metals are essential components of many enzyme systems, for instance, δ -aminolevulinic acid dehydratase (δ -ALAD or called PBGS, EC 4.2.1.24) is a metalloenzyme requiring zinc ions for activity (Jaffe et al. 1995). δ -ALAD catalyses the asymmetric condensation of two aminolevulinic acid (ALA) molecules to form porphobilinogen (PBG) in heme biosynthesis (Gibson et al. 1955) (Figure 1-2) pathway. The pyrrole is common precursor of the tetrapyrrole pigments such as heme, chlorophyll, and cobalamin, corrins, and its biosynthesis pathways are similar in all organisms (Senior et al. 1996, Shoolingin-Jordan 2003). PBGS is very highly conserved in sequence and structure but contains a remarkable phylogenetic variation in metal ion usage for catalytic and allosteric functions

Fig. 1. The heme biosynthetic pathway. Mitochondrial enzymes are depicted in green and cytosolic enzymes in red (Richard et al., 2006).

Fig. 2. Synthesis of porphobilinogen (PBG). Two molecules of ALA (blue and orange) are condensed to form PBG, a monopyrrole, by the cytosolic enzyme aminolevulinic acid dehydratase (ALAD) (Richard et al. 2006).

(Jaffe 2000, 2003). As of 2003, approximately one-half of the ~130 PBGS sequences available contained the binding determinants for a catalytic zinc ion, and about one-half did not (Jaffe 2003). On the other hand, approximately 90% of the known PBGS sequences contain the binding determinants for allosteric magnesium. The only known PBGS sequences that lack the binding determinants for both the catalytic zinc and the allosteric magnesium are in the bacterial genus *Rhodobacter* (Jaffe 2003).

δ -ALAD is a sulfhydryl containing enzyme (Gibson et al. 1955, Barnard et al. 1977) and numerous metals such as lead (Rodrigues et al. 1989, 1996, and Goering 1993), mercury (Rocha et al., 1993, 1995), and other compounds that oxidize sulfhydryl groups modified its activity (Emanuelli et al. 1996, Barbosa et al. 1998, Flora et al. 1998). Therefore, δ -ALAD is inhibited by substances that compete with zinc and/or that oxidize the -SH groups (Farina et al. 2002, Nogueira et al. 2003a-b, Santos et al. 2004) and is linked to situations associated with oxidative stress (Folmer et al. 2002, Pande et al. 2001, Pande & Flora 2002, Tandon et al. 2002, Soares et al. 2003). In addition, human exposure to Pb^{2+} causes an important inhibition of blood δ -ALAD (Meredith et al. 1979, Fujita et al. 1981, Pappas et al. 1995, Polo et al. 1995, Pires et al. 2002) and is associated with an intense anemia accompanied by an increase in urinary δ -ALA excretion (Oskarsson 1989). Therefore, δ -ALAD activity is used as one of the most reliable indicators of Pb^{2+} intoxication in humans and other animals (Meredith et al. 1979, Pappas et al. 1995).

ALADs have been purified from a wide variety of sources, including bovine liver (Gibson et al. 1955), human erythrocytes (Anderson & Desnick 1979), *Rhodopseudomonas capsulatus*, *Rhodobacter sphaeroides*, (Nandi & Shemin 1973; Nandi et al. 1968), *Escherichia coli* (Spencer & Jordan 1993) and spinach, *Spinacia oleracea* (Liedgens et al. 1983) during the time. Although the fundamental catalytic properties of all ALADs are similar, differences in enzyme primary structure, metal ion requirement and thiol sensitivity have been observed between the various purified enzymes. Metal dependency allows ALADs to be divided into two main categories, the Zn^{2+} -dependent and the Mg^{2+} -dependent dehydratases. The Zn^{2+} -dependent enzymes include the ALADs from mammalian sources, which have 'pH optima'

of between 6.3 and 7.1 and have been shown to require Zn^{2+} for maximal catalytic activity (Shemin 1972, Cheh & Neilands 1976). The yeast and *E. coli* enzymes can also be included in this class, requiring Zn^{2+} for activity but with more alkaline pH optima than the animal counterparts: 9.8 for the yeast and 8.5 for the enzyme from *E. coli* (Borralho et al. 1990). The animal, yeast and *E. coli* ALADs have a homo-octameric structure and have thiol groups that are extremely sensitive to oxidation. The oxidation of the thiol groups has been shown to be accompanied by a decrease in catalytic activity and a stoichiometric loss of bound metal ions (Tsukamoto et al. 1979), thereby demonstrating that the cysteine residues are required for Zn^{2+} binding. It has been established that ALADs from this class contain both catalytic and non-catalytic Zn^{2+} (Dent et al. 1990). Techniques such as EXAFS predict that the non-catalytic Zn^{2+} has a tetrahedral co-ordination of at least two and often four cysteine residues. The catalytic Zn^{2+} can be bound in either a tetrahedral or pentaco-ordinate fashion with cysteine, histidine and often water as ligands (Jaffe 1993). The Mg^{2+} -dependent class of dehydratases includes the plant ALADs, which have been reported to have alkaline pH optima of c.a. 8.0-8.5 (Liedgens et al. 1983), but again these values were determined by measurement of an average rate of reaction as described above. They have an absolute dependence on Mg^{2+} as well as subtle differences in their primary structure, especially in the putative metal-binding domains. In addition, some of the plant enzymes seem to be homohexameric and to be less sensitive to oxidation than their animal counterparts; consequently a minor role has been postulated for their thiol groups (Liedgens et al. 1983). This may be due to the fact that the cysteine residues are not involved in metal chelation in the Mg^{2+} -dependent enzymes and their oxidation therefore does not lead to the loss of metal ions. In an attempt to show conclusive differences and/or similarities in the ALADs, a detailed study of the properties of the ALADs from *E. coli*, yeast and pea was conducted. Evidences were presented supporting this hypothesis that the variances in metal binding between the enzymes are a reflection of significant biochemical differences that affect substrate recognition and binding and can therefore be used in the design of specific inhibitors (Senior et al. 1996). It was also revealed that the yeast enzyme, previously assumed to be only Zn^{2+} -binding, is similar to the *E. coli* ALAD in that Mg^{2+} can be substituted at the catalytic site to restore enzyme activity although there is no stimulation of activity. Finally, the crystallization of the yeast ALAD is reported, which will permit the determination of the structure of the enzyme by X-ray diffraction methods.

The yeast ALAD has been overexpressed, purified and found to be a Zn^{2+} -dependent, Mg^{2+} -binding enzyme that is similar in behaviour to, but not identical with, the ALAD from *E. coli*.

Comparative studies with the ALADs from three different sources have given an insight into some of the features required for molecular recognition, demonstrating that there are real differences both between and within the different classes of ALADs. These are sufficient to enable the selective inhibition of the enzymes. It will not be possible to rationalize all of the inhibition results collected until the three-dimensional structure of the yeast enzyme has been solved and substantial progress has been made towards this goal with the reported crystallization of the yeast enzyme.

More specific biochemical screening methods are being used by toxicologists such as, protein kinase variants, nitric oxide, interleukin 4 (IL4) and auto-antibodies in plumbism apart from gross changes such as stippling of erythrocytes or inhibition of ALAD (Nag et al. 1996).

Not only δ -ALAD is inhibited but also a number of other enzymes in heme biosynthesis pathway, including coproporphyrinogen oxidase and ferrochelatase are affected by lead.

Inhibitions of ALAD are most profound, and the degree of erythrocyte ALAD inhibition has been used clinically to estimate the degree of Pb poisoning in humans. ALA has neurotoxic activity and may contribute to Pb-induced neurotoxicity (Sithisarankul et al. 1997). At the molecular level, Pb displaces a zinc ion at the metal binding site, not the active site, producing inhibition through a change in the enzyme quaternary structure. ALAD is the second enzyme in the heme biosynthetic pathway, which is cytosolic and non-limiting in heme synthesis in healthy cells. Heme plays important roles in oxygen transport, electron transport systems, detoxification, and transcriptional regulations.

Porphobilinogen is the pyrrole precursor utilized by all living systems for the biosynthesis of tetrapyrroles, including hemes and chlorophylls (Jordan 1991). The ALAD polymorphism has not been established, but it is clear that geographic and strainspecific factors define the distribution of the two recognized ALAD alleles (Fleming et al. 1998). It has also been shown that organisms bred in environments containing high levels of Pb are endowed with multiple copies of the ALAD gene (Bishop et al. 1998).

4. Result

Determination of the blood Pb level alone cannot indicate the toxicity of Pb, since each individual has different degrees of tolerance of Pb (Marcus 1985). As known, atomic absorption spectrophotometry (AAS) or ICP is required for the determination of Pb, and both are expensive. These instruments are available only in specialist laboratories, and can be operated only by a well-trained technician. For these reasons same Pb affected enzymes have been employed as biosensors in monitoring Pb toxicity. Among these, ALAD is most popular one, and its results showed good combination with the blood Pb level determined by atomic absorption spectrophotometry or ICP. There are some mathematical equations devised by several authors to give the Pb concentration by looking at the activity of ALAD (Ogunseitan 1999, Ogunseitan et al. 2000, Korcan et al. 2007, Cigerci et al. 2008, Konuk et al. 2008).

The expression of ALAD activity gives us a clear indication of the severity of the effect of Pb pollution along the pollution gradient. That is why it is an important biosensor for Pb contamination and pollution. Further studies should be focused on the determination of molecular basis of its effect on ALADs in different organisms and these studies should be strengthened by immobilization studies.

5. References

- Anderson, P.M., Desnick, R.J., (1979). Purification and Properties of δ -Aminolevulinic Acid Dehydratase from Human Erythrocytes, *J. Biol. Chem.*, 254, 6924-6930.
- Ashraf, W., (2005). Accumulation of heavy metals in kidney and heart tissues of *Epinephelus microdon* fish from the Arabian Gulf. *Environ Monit Assess.* 101, 311.
- Ballatori, N., (2002). Transport of toxic metals by molecular mimicry. *Environ. Health Perspect.* 110 (Suppl. 5), 689-694.
- Barbosa, N.B.V., Rocha, J.B.T., Zeni, G., Emanuelli, T., Beque, M.C., Braga, A.L., (1998). Effect of organic selenium on δ -aminolevulinate dehydratase from liver, kidney, and brain of adults rats. *Toxicol. Appl. Pharmacol.* 149, 243-253.
- Barnard, G.F., Itoh, R., Hohberger, L.H., Shemin, D., (1977). Mechanism of porphobilinogen synthase – possible role of essential thiol-groups. *J. Biol. Chem.*, 252, 8965-8974.

- Bishop, T. R., Miller, M. W., Wang, A., and Dierks, P. M., (1998). Multiple copies of the ALAD gene are located at the *Lv* locus in *Mus domesticus* mice. *Genomics*, 48, 221–231.
- Borralho, L.M., Ortiz, C.H.D., Panek, A.D. and Mattoon, J.R. (1990). Purification of δ -aminolevulinic acid dehydratase from genetically engineered yeast. *Yeast* 6, 319–330.
- Burger, J., (1995). A risk assessment for lead in birds. *J Toxicol Environ Health* 45:369–396.
- Burger, J., Gochfeld, M., (2000a). Metals levels in feathers of 12 species of seabirds from Midway Atoll in the northern Pacific Ocean. *Sci Total Environ* 257:37–52.
- Burger, J., Gochfeld, M., (2000b). Effects of lead on birds (Laridae): A review of laboratory and field studies. *J Toxicol Environ Health B Crit. Rev.* 3(2):59–78.
- Busselberg, D., Evans, M.L., Rahmann, H., Carpenter, D.O., (1991). Effects of inorganic and triethyl lead and inorganic mercury on the voltage activated calcium channel of *Aplysia* neurons. *Neurotoxicol.*, 12, 733–744.
- Cavalcanti, A., Shirinzadeh, B., Zhang, M., Kretly, L.C., (2008). "Nanorobot hardware architecture for medical defense". *Sensors* 8 (5): 2932–2958.
- Cheh, A. and Neilands, J. B., (1976). The aminolevulinic acid dehydratase: Molecular and environmental properties, *Struct. Bond.*, 29, 123–170.
- Ciğerci, İ.H., Korcan, S.E., Konuk, M. and Öztürk, S., (2008). Comparison of ALAD activities of *Citrobacter* and *Pseudomonas* strains and their usage as biomarker for Pb contamination. *Environ. Monitor. Assess.*, 139(1–3), 41–48.
- Ciucu, A., Lupu, A., Pirvutoi, S. and Palleschi, G., (2001). Biosensors for heavy metals determination based on enzyme inhibition. *Scientific Bulletin–University "Politehnica" of Bucharest, Series B, Chem. Mater. Sci.* 63 (4), pp. 33–44.
- Clark, L.C. and Lyons, C., (1962). Electrode system for continuous monitoring of cardiovascular surgery. *Ann. NY Acad. Sci.*, 102, 2945.
- Compagnone, D., Palleschi, G., Varallo, G. and Imperiali, P.L., (1995). Amperometric biosensors for the determination of heavy metals In: T. Vo-Dinh, Editors, *Proceedings of SPIE*, vol. 2504: Environmental Monitoring and Hazardous Waste Site Remediation, pp. 141–152.
- Dautremepuits, C., Paris-Palacios, S., Betoulle, S., Vernet, G., (2004). Modulation in hepatic and head kidney parameters of carp (*Cyprinus carpio* L.) induced by copper and chitosan. *Comp. Biochem. Physiol. C Toxicol. Pharmacol.*, 137, 325–33.
- Dauwe, T., Bervoets L., Blust, R., Pinxten, R., Eens, M., (2000). Can excrement and feathers of nestlings songbirds be used as biomonitors for heavy metal pollution? *Arch. Environ. Contam. Toxicol.*, 39:541–546.
- Davis, P.H., Goettl, J.P., Sinley, J.R., Smith, N.F., (1976). Acute and chronic toxicity of lead to rainbow trout *Salmo gairdneri*, in hard and soft water. *Water Res.*, 10, 199–206.
- Dent, A.J., Beyersmann, D., Block, C. and Hasnain, S.S., (1990). Two different zinc sites in bovine δ -aminolevulinic acid dehydratase distinguished by EXAFS. *Biochemistry*, 29, 7822–7828.
- Dix, D.J., Gallagher, K., Benson, W.H., Groskinsky, B.L., McClintock, J.T., Dearfield, K.L., Farland, W.H., (2006). A framework for the use of genomics data at the EPA. *Nat. Biotechnol.*, 24, 1108–1111.
- Donlan, A. M., Moody, G.J. and Thomas, J.D.R., (1989). The amperometric detection of some enzyme inhibitors, *Anal. Proceed.* 26, pp. 369–371.

- Dooley, M.A., Hogan, S.L., (2003). Environmental epidemiology and risk factors for autoimmune disease. *Curr. Opin. Rheumatol.*, 15, 99–103.
- Dzyadevych, V., A.P. Soldatkin, Y.I. Korpan, V.N. Arkhypova, A.V. El'skaya, J. Chovelon, C. Martelet and N. Jaffrezic-Renault, (2003), Biosensors based on enzyme field-effect transistors for determination of some substrates and inhibitors, *Anal. Bioanal. Chem.*, 377, 496–506.
- Emanuelli, T., Rocha, J.B.T., Pereira, M.E., Porciuncula, L.O., Morsch, V.M., Martins, A.F., Souza, D.O., (1996). Effect of mercuric chloride intoxication and dimercaprol treatment on aminolevulinate dehydratase from brain, liver and kidney of adult mice. *Pharmacol. Toxicol.*, 79, 138–143.
- Evtugyn, G.A., Budnikov, H.C. and Nikolskaya, E.B., (1999). Biosensors for the determination of environmental inhibitors of enzymes, *Russ. Chem. Rev.*, 68, 1041–1064.
- Fair, J.M., Ricklefs, R.E., (2002). Physiological, growth, and immune responses of Japanese quail chicks to the multiple stressors of immunological challenge and lead shot. *Arch. Environ. Contam. Toxicol.*, 42:77–87.
- Farina, M., Barbosa, N.B.V., Nogueira, C.W., Folmer, V., Zeni, G., Andrade, L.H., Braga, L.A., Rocha, J.B.T., (2002). Reaction of diphenyl diselenide with hydrogen peroxide and inhibition of daminolevulinate dehydratase from rat liver and cucumber leaves. *Braz. J. Med. Biol. Res.*, 35, 623–631.
- Fleming, D. E. B., Chettle, D. R., Wetmur, J. G., Desnick, R. J., Robin, J. P., Boulay, D., (1998). Effect of the deltaaminolevulinatedehydratase polymorphism on the accumulation of lead in bone and blood in lead smelter workers. *Environ. Res.*, 77, 49–61.
- Flora, S.J.S., Gubrelay, U., Kannan, G.M., Mathur, R., (1998). Effects of zinc supplementation during chelating agent administration in cadmium intoxication in rats. *J. Appl. Toxicol.*, 18, 357–362.
- Folmer, V., Soares, J.C.M., Rocha, J.B.T., (2002). Oxidative stress in mice is dependent on the free glucose content of the diet. *Int. J. Biochem. Cell Biol.*, 34, 1279–1286.
- Furness, R.W., (1993). Birds as monitors of pollutants. In: Furness RW, Greenwood JJD (eds) *Birds as monitors of environmental change*. Chapman & Hall, London, pp 86–143.
- Gibson, K.D., Neureberger, A., Scott, J.J., (1955). The purification and properties of delta-aminolevulinic acid dehydratase. *Biochem. J.*, 61, 618–629.
- Grosell, M., Gerdes, R., Brix, K.V., (2006). Influence of Ca, humic acid and Ph on lead accumulation and toxicity in the fathead minnow during prolonged water-borne lead exposure. *Comp. Biochem. Physiol. C-Toxicol. Pharmacol.* 143, 473–483.
- Guilbault, G.G., (1984). *Analytical Uses of Immobilized Enzymes* Marcel Dekker, New York.
- Hodson, P.V., Blunt, B.R., Spry, D.J., (1978). Chronic toxicity of water-borne and dietary lead to rainbow trout (*Salmo gairdneri*) in Lake Ontario water. *Water Res.*, 12, 869–878.
- Holcombe, G.W., Benoit, D.A., Leonard, E.N., McKim, J.M., (1976). Long-term effects of lead exposure on three generations of brook trout (*Salvelinus fontinalis*). *J. Fish. Res. Board Can.*, 33, 1731–1741.
- <http://en.wikipedia.org/wiki/Biosensor>
- <http://www.rpi.edu/dept/chem-eng/Biotech Environ/BIOSSEN/enzbio.htm>
- Hu, H., (2000). Exposure to metals. *Prim. Care.*, 2, 983–996.

- Huggett, R.J., Kimerle, R.A., Mehrie P.M. and Bergman, H.L., (1992). Biomarkers: Biochemical Physiological and Histological Markers of Anthropogenic Stress, Lewis Publishers, London.
- Jaffe, E.K., (2000). The porphobilinogen synthase family of metalloenzymes. *Acta Crystallog. D Biol Crystallogr.*, 56:115-128.
- Jaffe, E.K., (2003). An unusual phylogenetic variation in the metal ion binding sites of porphobilinogen synthase. *Chem Biol.*, 10:25-34.
- Jaffe, E.K., Ali, S., Mitchell, L.W., Taylor, K.M., Volin, M., Markham, G.D., (1995). Characterization of the role of the stimulatory magnesium of *Escherichia coli* porphobilinogen synthase. *Biochem.*, 34, 244-251.
- Järup, L., (2003). Hazards of heavy metal contamination, *Brit. Med. Bull.*, 68:167-182.
- Jordan, P. M., (1991). New comprehensive biochemistry. Amsterdam, The Netherlands: Elsevier.
- Kalay, M., Ay, P., Canil, M., (1999). Heavy metal concentration in fish tissues from the northeast Mediterranean. *Bull Environ. Contam. Toxicol.*, 63, 673-671.
- Konuk, M., Cigerci, İ.H., Aksan, Ş. and Korcan, S.E., (2008). Isolation and biochemical characterization of δ -aminolevulinic acid dehydratase from *Streptomyces yokosukanensis* ATCC 25520. *Appl. Biochem. Microbiol.*, 44, 356-360.
- Korcan, S.E., Cigerci, İ.H. and Konuk, M., (2007). Screening of δ -Aminolevulinic Acid Dehydratase from *Pseudomonas* strains as biosensor for Lead and some other metals contamination. *Environ. Monitor. Asses.*, 134, 263-269.
- Liedgens, W., C. Lutz, and H. A. W. Schneider., (1983). Molecular properties of 5-aminolevulinic acid dehydratase from *Spinacia oleracea*. *Eur. J. Biochem.*, 135:75-79.
- Macdonald, A., Silk, L., Schwartz, M., Playle, R.C., (2002). A lead-gill binding model to predict acute lead toxicity to rainbowtrout (*Oncorhynchus mykiss*). *Comp. Biochem. Physiol. C-Toxicol. Pharmacol.*, 133, 227-242.
- Marcus, A.H., (1985). Multicompartment Kinetic Models for Lead: Part1, B one Kinetic and Variable Absorption in Human without excessive lead exposure. *Env. Res.*, 3, 459-479.
- Mazliah, J., Barron, S., Bental, E., Reznik, I., (1989). The effect of chronic lead intoxication in mature chickens. *Avian Dis.*, 33:566-570.
- Meredith, P.A., Moore, M.R., Goldberg, A., (1979). Erythrocyte ALA dehydratase activity and blood protoporphyrin concentrations as indices of lead exposure and altered haem biosynthesis. *Cli. Sci. Mol. Med.*, 56, 61-69.
- Molina, V., Ehrenfeld, M., (2003). Environmental factors in autoimmune diseases, 4-5 February 2003. Durham, NC, USA. *Autoimm. Rev.*, 2, 284-287.
- Nag, D., Jaffery, F.N., Viswanathan, P.N., (1996). Clinical and biochemical screening tests for identification of high risk groups. In: Richardson, M.L. (Ed.), Risk Reduction. Chemicals and Energy into the 21st Century. Taylor & Francis, London, UK, pp. 285-302.
- Nandi, D.L. and Shemin, D., (1968). Delta-aminolevulinic acid dehydratase of *Rhodospseudomonas spheroides* II. Association to polymers and dissociation to subunits. *J. Biol. Chem.*, 243, 1231-1235.
- Nandi, D.L. and Shemin, D., (1973). ALAD of *Rhodopseudomonas capsulatus*, *Arch. Biochem. Biophys.*, 158, 305-311.

- Needleman, H.L., Schell, A., Bellinger, D., Leviton, A., Allred, E.N., (1990). The long-term effects of exposure to low doses of lead in childhood: an 11 year followup report. *N. Engl. J. Med.* 32, 83–88.
- Nogueira, C.W., Borges, V.C., Zeni, G., Rocha, J.B.T., (2003b). Organochalcogens effects on aminolevulinate dehydratase activity from human erythrocytic cells in vitro. *Toxicol.*, 191, 169–178.
- Nogueira, C.W., Soares, F.A., Nascimento, P.C., Muller, D., Rocha, J.B.T., (2003a). 2,3-Dimercaptopropane-1-sulfonic acid and meso- 2,3-dimercaptosuccinic acid increase mercury- and cadmiuminduced inhibition of δ -aminolevulinate dehydratase. *Toxicol.*, 184, 85–95.
- Ogunseitan, O. A., (1999). Microbial proteins as biomarkers for ecosystem health. In K. Scow, G Fogg, D. Hinton, & M. Johnson (Eds.), *Integrated assessment of ecosystem healthy* (pp. 207–222). Boca Raton, FL: Lewis.
- Ogunseitan, O.A., Yang, S. and Ericson, J., (2000). Mikrobial δ -aminolevulinate dehydratase as a biosensor of lead bioavailability in contaminated environments. *Soil Biol. Biochem.*, 32, 1899–1906.
- Oskarsson, A., (1989). Effects of perinatal treatment with lead and dissulfiram on ALA-D activity in blood, liver and kidney and urinary ALA excretion in rats. *Pharm. Toxicol.*, 64, 344–348.
- Pande, M., Flora, S.J.S., (2002). Lead induced oxidative damage and its response to combined administration of α -lipoic acid and succimers in rats. *Toxicol.*, 177, 187–196.
- Pande, M., Mehta, A., Pant, B.P., Flora, S.J.S., (2001). Combined administration of a chelating agent and an antioxidant in the prevention and treatment of acute lead intoxication in rats. *Environ.Toxicol. Pharmacol.*, 9, 173–184.
- Pappas, J.B., Ahlquist, J.T., Allen, E.M., Banner Jr., W., (1995). Oral dimercaptosuccinic acid and ongoing exposure to lead: effects on heme synthesis and lead distribution in a rat model. *Toxicol.Appl. Pharm.*, 133, 121–129.
- Paquin, P.R., Gorsuch, J.W., Apte, S., Batley, G.E., Bowles, K.C., Campbell, P.G., Delos, C.G., Di Toro, D.M., Dwyer, R.L., Galvez, F., Gensemer, R.W., Goss, G.G., Hostrand, C., Janssen, C.R., McGeer, J.C., Naddy, R.B., Playle, R.C., Santore, R.C., Schneider, U., Stubblefield, W.A., Wood, C.M., Wu, K.B., (2002). The biotic ligand model: a historical overview. *Comp. Biochem. Physiol. C-Toxicol. Pharmacol.*, 133, 3–35.
- Patel, M., Rogers, J.T., Pane, E.F., Wood, C.M., (2006). Renal responses to acute lead waterborne exposure in the freshwater rainbow trout (*Oncorhynchus mykiss*). *Aquat. Toxicol.*, 80, 362–371.
- Pires, J.B., Miekeley, N., Donangelo, C.M., (2002). Calcium supplementation during lactation blunts erythrocyte lead levels and deltaaminolevulinic acid dehydratase zinc-reactivation in women nonexposed to lead and with marginal calcium intakes. *Toxicol.*, 175, 247–255.
- Pirvutoiu, I. Surugiu, E.S. Dei, A. Ciucu, V. Magearu and B. Danielsson, (2001), Flow injection analysis of mercury (II) based on enzyme inhibition and thermometric detection, *Analyst* 126, 1612–1616.
- Rand, G.M., (1995). *Fundamentals of Aquatic Toxicology: Effects, Environmental Fate and Risk Assessment*, Taylor & Francis, London.

- Richard, S. A., Phillips, J. D., Kushner, J. P., (2006). Biosynthesis of heme in mammals. *Biochim. Biophys. Acta.*, 1763, 723–736.
- Rocha, J.B.T., Freitas, A.J., Marques, M.B., Pereira, M.E., Emanuelli, T., Souza, D.O., (1993). Effects of methylmercury exposure during the second stage of rapid postnatal brain growth on negative geotaxis and on delta-aminolevulinic acid dehydratase of suckling rats. *Braz. J. Med. Biol. Res.*, 26, 1077–1083.
- Rocha, J.B.T., Pereira, M.E., Emanuelli, T., Christofari, R.S., Souza, D.O., (1995). Effect of treatment with mercury chloride and lead acetate during the second stage of rapid postnatal brain growth on delta-aminolevulinic acid dehydratase (ALA-D) activity in brain, liver kidney and blood of suckling rats. *Toxicol.*, 100, 27–37.
- Rodrigues, A.L., Bellinaso, M.L., Dick, T., (1989). Effect of some metalions on blood and liver delta-aminolevulinate dehydratase of *Pimelodus maculatus* (Pisces, Pimelodidae). *Comp. Biochem. Physiol.*, 94, 65–69.
- Rogers, J.T., Wood, C.M., (2004). Characterization of branchial lead-calcium interaction in the freshwater rainbow trout *Oncorhynchus mykiss*. *J. Exp. Biol.*, 207, 813–825.
- Santos, F.W., Oro, T., Zeni, G., Rocha, J.B.T., do Nascimento, P.C., Nogueira, C.W., (2004). Cadmium induced testicular damage and its response to administration of succimer and diphenyl diselenide in mice. *Toxicol. Lett.*, 152, 255–263.
- Senior, N.M., Brocklehurst, K., Cooper, J.B., Wood, S.P., Erksine, P., Shoolingin- Jordan, P., Thomas, P.G., Warren, M.J., (1996). Comparative studies on the 5- aminolevulinic acid dehydratase from *Pisum sativum*, *Escherichia coli* and *Saccharomyces cerevisiae*. *Biochem J.* 320:410-412
- Shemin, D., (1972). Aminolevulinic acid dehydratase, *Enzymes*, 7, 323-337.
- Shoolingin-Jordan, P.M., (2003). The biosynthesis of Coproporphyrinogen III. In: Kadish KM, Smith KM and Guillard R. , editor. *The Porphyrin Handbook*. Vol. 12. Amsterdam, Elsevier Science; pp. 33–74.
- Sithisarankul, P., Schwartz, B.S., Lee, B.K., Kalsey, K.T., Strickland, P.T., (1997). Aminolevulinic acid (ALA) dehydratase genotype mediates plasma levels of the neurotoxin, 5-aminolevulinic acid, in lead exposed workers. *Am. J. Ind. Med.*, 32, 15–28.
- Soares, J.C.M., Folmer, V., Rocha, J.B.T., (2003). Influence of dietary selenium supplementation and exercise on thiol-containing enzymes in mice. *Nutrition* 19, 627–632.
- Spencer, P., Jordan, P.M., (1993). Purification and characterization of 5-aminolaevulinic acid dehydratase from *Escherichia coli* and a study of the reactive thiols at the metal-binding domain. *Biochem. J.*, 290, 279-287.
- Starodub, N.F., Kanjuk, N.I., Kukla, A.L., Shirshov and Yu, M., (1999). Multi-enzymatic electrochemical sensor: field measurements and their optimization, *Anal. Chim. Acta*, 385, pp. 461–466.
- Stein, J., Schettler, T., Wallinga, D., Vallenti, M., (2002). In harm's way: toxic threats to child development. *J. Dev. Behav. Pediatr. (Suppl. 1)*, 13–22.
- Tandon, S.K., Singh, S., Prasad, S., Srivastava, S., Siddiqui, M.K.J., (2002). Reversal of lead-induced oxidative stress by chelating agent, antioxidant, or their combination in the rat. *Environ. Res.*, 90, 61–66.
- Tang, H.W., Huel, G., Compagna, D., Hellier, G., Boissinot, C., Blot, P., (1999). Neurodevelopment evaluation of 9-month old infants exposed to low levels of Pb

- in vitro: involvement of monoamine neurotransmitters. *J. Appl. Toxicol.*, 19, 167–172.
- Timbrell, J.A., (2000). *Principles of Biochemical Toxicology*, Taylor & Francis, London.
- Tsukamoto, I., Yoshinaga, T., Sano, S., (1979). The role of zinc with special reference to the essential thiol groups in delta-aminolevulinic acid dehydratase of bovine liver. *Biochim. Biophys. Acta.*, 12;570(1):167–178.
- USEPA, (2006). *National Recommended Water Quality Criteria*. Office of Water, Washington, DC.
- Vel Krawczyc, K., Moszczynska, M. and Trojanowicz, M., (2000). Inhibitive determination of mercury and other metal ions by potentiometric urea biosensor, *Biosens. Bioelectron.*, 15, pp. 681–691.
- Wang, W. X., (2002). Interaction of trace metals and different marine food chains. *Mar. Ecol. Prog. Ser.* 243, 295–309.
- Weber, D.N., (1993). Exposure to sublethal levels of waterborne lead alters reproductive behavior patterns in fathead minnows (*Pimephales promelas*). *Neurotoxicol.*, 14, 347–358.
- Winneke, G., Walkowiak, J., Lilienthal, H., (2002). PCB induced neurodevelopmental toxicity in human infants and its potential mediation by endocrine dysfunction. *Toxicol.*, 181–182, 161–165.
- Yang, J.H., Derr-Yellin, E.C., Kodavanti, P.R., (2003). Alterations in brain protein kinase c isoforms following developmental exposure to a polychlorinated biphenyl mixture. *Brain Res. Mol. Brain Res.*, 123–135.

IntechOpen

Intelligent and Biosensors

Edited by Vernon S. Somerset

ISBN 978-953-7619-58-9

Hard cover, 386 pages

Publisher InTech

Published online 01, January, 2010

Published in print edition January, 2010

The use of intelligent sensors have revolutionized the way in which we gather data from the world around us, how we extract useful information from that data, and the manner in which we use the newly obtained information for various operations and decision making. This book is an attempt to highlight the current research in the field of Intelligent and Biosensors, thereby describing state-of-the-art techniques in the field and emerging new technologies, also showcasing some examples and applications.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Muhsin Konuk, İbrahim Hakkı Ciğerci and Safiye Elif Korcan (2010). ALAD (δ -aminolevulinic Acid Dehydratase) as Biosensor for Pb Contamination, Intelligent and Biosensors, Vernon S. Somerset (Ed.), ISBN: 978-953-7619-58-9, InTech, Available from: <http://www.intechopen.com/books/intelligent-and-biosensors/alad-aminolevulinic-acid-dehydratase-as-biosensor-for-pb-contamination>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2010 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike-3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/), which permits use, distribution and reproduction for non-commercial purposes, provided the original is properly cited and derivative works building on this content are distributed under the same license.

IntechOpen

IntechOpen