

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Microalgae and Its Use in Nutraceuticals and Food Supplements

Joshi Nilesh Hemantkumar and Mor Ilza Rahimbhai

Abstract

Microalgae are a large diverse group of microorganisms comprising photoautotrophic protists and prokaryotic cyanobacteria—also called as blue-green algae. These microalgae form the source of the food chain for more than 70% of the world's biomass. It contains higher nutritional values, with rapid growth characteristics. Microalgae are autotrophic organisms and extensively desired for use in nutraceuticals and as supplement in diet. Many microalgal species are documented for health benefits, by strengthening immune system and by increasing the nutritional constitution of body. In this chapter the major economically important species like *Spirulina*, *Chlorella*, *Haematococcus*, and *Aphanizomenon* are described with reference to its importance as nutraceuticals and food.

Keywords: microalgae, *Spirulina*, *Chlorella*, nutraceuticals, food supplements

1. Introduction

Microalgae are a large diverse group of microorganisms comprising photoautotrophic protists and prokaryotic cyanobacteria—also called as blue-green algae. These microalgae form the source of the food chain for more than 70% of the world's biomass [1]. Microalgae are single-celled, microscopic photosynthetic organisms, found in freshwater and marine environment. They produce compounds such as protein, carbohydrates, and lipids. Mostly, microalgae are photosynthetic microorganisms; it does not contain cell organelles unlike land plants. They use the carbon from air for energy production.

Microalgae can be cultivated photosynthetically using CO₂, solar energy, and water. It can be cultivated in shallow lagoons, marginal ponds, raceway ponds, or artificial tanks. The use of plastic tubes/reactors in pond system can achieve up to seven times the production efficiency compared to open culture system [2].

There are more than 300,000 species of microalgae, out of which around 30,000 are documented. They live in complex natural habitats and can adapt rapidly in extreme conditions (in variation of extreme weather conditions). This ability makes them capable to produce secondary metabolites, with novel structure and biologically active functions.

Microalgae produce some useful bio-products including β -carotene, astaxanthin, docosahexaenoic acid (DHA), eicosapentaenoic acid (EPA), bioactive and functional pigments, natural dyes, polysaccharides, antioxidants, and algal extracts.




Figure 1.
Application of microalgae in various fields.

The first commercial cultivation of *Chlorella* was started in 1960 in Japan for nutraceuticals. Microalgae grow fast and produce large biomass with high protein content and consist the source of “single cell protein” [3] (**Figure 1**).

Algae are classified into many major groups, based on pigment composition, storage compound, and diversity in features of its ultrastructure. However, advance molecular biology-based techniques are nowadays used to check the relation between taxonomic groups and families of the specific class.

The global market value of microalgae is estimated to be around US\$ 6.5 billion, out of which about US\$ 2.5 billion is generated by the health food sector, US\$ 1.5 billion by the production of DHA, and US\$ 700 million by aquaculture. The annual production of microalgae is approximately 7.5 million tons.

2. Microalgal diversity

The diversity of microalgae is vast and represents an intact resource. The scientific literature indicates the existence of 200,000 to several million species of microalgae when compared to about 250,000 species of higher plants [4].

Green microalgae usually grow in freshwater and seawater, whereas several other species of microalgae grow in extremely saline environments, such as the Great Salt Lake in UT, USA, and the Dead Sea in Israel. Within these aqueous habitats, some algae grow inside the deeper waters, others populate the subsurface water column, and a few grow at the limits of the photic zone, 200–300 m below the water surface [5–8]. The microalgae are small in size (mostly 5–50 μm) and characterized by a simple morphology, usually unicellular. Accordingly, most of the species are not observed as an individual cell/specimen but become noticeable only when it generates large colony, specially in the form of green, black, red, or brown patches on the water surface. Coastlines between 45 and 30°N are suitable regions for algal farming, in particular in those territories at the south of the Mediterranean that experience warmer climates and whose temperature does not go too much below 15°C throughout the year [2]. This type of warmer climate of the Mediterranean region can facilitate the algal growth in the open or closed pond system.

The environmental parameters favorable for mass scale culture are explored by many countries. For example, Israel and few Mediterranean countries have explored specific parameters for each economically important alga and cultivating it by maintaining them artificially for mass scale production. Few advance countries have started culturing several microalgal strains of biofuel production, while countries like Libya, Cyprus, and Turkey also have plenty of marginal lands to harvest algae. For these countries a limited water resource is not a constraint as they are using recycled brackish or saline water. With the high temperatures in the Mediterranean region, the open or closed pond system would probably be the most efficient and suitable to grow algae.

Terrestrial microalgae belong primarily to three diverse evolutionary pedigrees: the blue-green algae (Cyanobacteria), the green algae (Chlorophyta and Streptophyta), and the diatoms (Bacillariophyceae, Ochrophyta) [9]. However, the species of green and blue-green algal group are the majorly studied group, taxonomically as well as with economical perspective. Nevertheless, the understanding of the patterns of geographical distribution in terrestrial algae is inadequate, mainly due to poor understanding of the diversity of these organisms [10].

3. Microalgae: uses as nutraceuticals and food

Microalgae have a wide range of industrial applications, in food industries, wastewater purification, and pharmaceutical formulations [11]. Microalgae can also be used for high-value food, health food for human, polysaccharides, food and fodder additives, cosmetics, antioxidants, anti-inflammatory objects, dyes and feed for aquaculture, and preparation of biofilms [3, 12, 13].

The most widely used microalgae include Cyanophyceae (blue-green algae), Chlorophyceae (green algae), Bacillariophyceae (including diatoms), and Chrysophyceae (including golden algae). **Table 1** highlights some major microalgal species, products, and their application.

Since the last 20 years, biotechnological and nutraceutical application of microalgae has focused specifically on four major microalgae: (a) *Spirulina* (*Arthrospira*), (b) *Chlorella*, (c) *Dunaliella salina*, and (d) *Haematococcus pluvialis*.

3.1 *Spirulina*

Spirulina is a prokaryotic cyanobacterium that has been commercially produced for over 30 years for uses including fish food, vitamin supplements, food dyes, aquaculture, pharmaceuticals, and nutraceuticals [15, 16]. *Spirulina* is manufactured by many pharmaceutical companies. This alga is thought of as a super food and is widely cultured, primarily in specifically designed raceway ponds and photobioreactors, to meet the current demand.

Spirulina is one of the algae studied for large-scale commercial culture. It grows best at a high pH (9–11) and high bicarbonate concentrations. Generally raceway ponds are used to culture *Spirulina*. The water depth in the pond is generally 300–500 mm depending on the physical–chemical parameters and density of microalgae. The depth is also dependent on pond size, water flow velocity, and light absorption by the algal culture. Water temperature and pH have a large effect on the productivity of this species. It grows well between the temperatures of 35 and 37°C [17]. *Spirulina* is a filamentous microalga; hence, its harvesting is relatively easy.

Spirulina contains 60–70% protein by weight (including many amino acids) and contains up to 10 times more beta-carotene than carrots per unit mass [18]. *Spirulina* is rich in nutrients such as B vitamins, phycocyanin, chlorophyll, vitamin E,

Group/species	Extract	Use/application
<i>Arthrospira (Spirulina) platensis</i>	Phycocyanin, biomass	Health food, cosmetics
<i>Arthrospira (Spirulina)</i>	Protein, vitamin B12	Antioxidant capsule, immune system
<i>Aphanizomenon flos-aquae</i>	Protein, essential fatty acids, β -carotene	Health food, food supplement
<i>Chlorella</i> spp.	Biomass, carbohydrate extract	Animal nutrition, health drinks, food supplement
<i>Dunaliella salina</i>	Carotenoids, β -carotene	Health food, food supplement, feeds
<i>Haematococcus pluvialis</i>	Carotenoids, astaxanthin	Health food, food supplement, feeds
<i>Odontella aurita</i>	Fatty acids, EPA	Pharmaceuticals, cosmetics, anti-inflammatory
<i>Porphyridium cruentum</i>	Polysaccharides	Pharmaceuticals, cosmetics
<i>Isochrysis galbana</i>	Fatty acids	Animal nutrition
<i>Phaeodactylum tricornutum</i>	Lipids, fatty acids	Nutrition, fuel production
<i>Lyngbya majuscula</i>	Immune modulators	Pharmaceuticals, nutrition
<i>Schizochytrium</i> sp.	DHA and EPA	Food, beverage, and food supplement
<i>Cryptocodinium cohnii</i>	DHA	Brain development, infant health and nutrition
<i>Nannochloropsis oculata</i>	Biomass	Food for larval and juvenile marine fish

Adopted from [14].

Table 1.
The major microalgal species, products, and application.

omega 6 fatty acids, and many minerals [19]. *Spirulina* is used for weight loss [20], diabetes [21], high blood pressure, and hypertension [22]. It has documented antiviral [23, 24] and anticancer properties [25]. *Spirulina* positively affects cholesterol metabolism by increasing HDL levels, which can lead to healthy cardiovascular functions [26]. Romay et al. [27] described the antioxidant and anti-inflammatory properties of C-phycocyanin, which is a prevalent pigment in *Spirulina*. Essential amino acids like leucine, isoleucine, and valine are significantly present in *Spirulina*. *Spirulina* contains fatty acids like linolenic and γ -linolenic acid and ω -3 and ω -6 polyunsaturated fatty acids. *Spirulina platensis* is a natural source of DHA accounting up to 9.1% of the total fatty acids. The mineral content of *Spirulina* depends on the water in which it was grown, and its content of iron, calcium, and magnesium provides high nutritional value [16]. Belay et al. [18] reported that *Spirulina* powder contains provitamin A (2.330×10^3 IU/kg), β -carotene (140 mg 100/g), vitamin E (100 mg 100/g), thiamin B1 (3.5 mg 100/g), riboflavin B2 (4.0 mg 100/g), niacin B3 (14.0 mg 100/g), vitamin B6 (0.8 mg 100/g), inositol (64 mg 100/g), vitamin B12 (0.32 mg 100/g), biotin (0.005 mg 100/g), folic acid (0.01 mg 100/g), pantothenic acid (0.1 mg 100/g), and vitamin K (2.2 mg 100/g). The amino acid composition of *Spirulina* is given in **Table 2**.

3.2 *Chlorella*

Chlorella is a single-cell, spherical shaped (2–10 μ m in diameter), and photoautotrophic green microalga with no flagella. It multiplies rapidly requiring only CO₂, water, sunlight, and a small amount of minerals. *Chlorella* has been grown commercially cultured in photobioreactors [28] and harvested by centrifugation or

Amino acids (g kg ⁻¹)	<i>Spirulina</i>	<i>Chlorella</i>
Alanine	47	48
Arginine	43	36
Aspartic acid	61	52
Cysteine	6	4
Glutamic acid	91	63
Glycine	32	34
Histidine	10	13
Isoleucine	35	26
Leucine	54	53
Lysine	29	35
Methionine	14	15
Phenylalanine	28	31
Proline	27	29
Serine	32	28
Threonine	32	27
Tryptophan	9	6
Tyrosine	30	21
Valine	40	36

Adapted from [16].

Table 2.
Amino acid profile of Spirulina and Chlorella microalgae (in mg kg⁻¹).

autoflocculation. After harvesting the biomass is spray-dried, and the cell powder is sold directly. *Chlorella* contains 11–58% protein, 12–28% carbohydrate, and 2–46% lipids of its dry weight [29]. It also contains various vitamins such as β -carotene (180 mg 100/g), provitamin A (55,500 IU/kg), thiamin B1 (1.5 mg 100/g), vitamin E (<1 mg 100/g), riboflavin B2 (4.8 mg 100/g), niacin B3 (23.8 mg 100/g), vitamin B6 (1.7 mg 100/g), inositol (165.0 mg 100/g), vitamin B12 (125.9 mg 100/g), biotin (191.6 mg 100/g), folic acid (26.9 mg 100/g), and pantothenic acid (1.3 mg 100/g) [18, 30]. The amino acid composition of *Chlorella* is shown in Table 2.

Chlorella is able to decrease blood pressure, lower cholesterol levels, and enhance the immune system [16]. It also has the potential to relieve fibromyalgia, hypertension, or ulcerative colitis [31, 32]. The presence of aortic atheromatous lesions was significantly inhibited, and low-density lipoprotein (LDL) cholesterol levels were greatly suppressed upon consumption of *Chlorella* [33]. Some *Chlorella* consumers have mentioned a potential correlation between some brands of *Chlorella* tablets and nausea, vomiting, and other gastrointestinal issues. *Chlorella* has been labeled as a weak allergen and may be of clinical significance to certain types of people [34].

3.3 *Dunaliella*

Dunaliella (*D. salina*) is a unicellular green alga which contains large amounts of β -carotene, glycerol, and protein that can easily be extracted through its thin cell wall. *Dunaliella* does not require waters appropriate for agricultural and domestic

uses and can be cultured in brackish water, marine water, and highly saline water. Global production of *Dunaliella* is estimated to be 1200 tons dry weight per year [16]. The dominant companies that produce *Dunaliella*, mainly for beta-carotene production, are located in Israel, China, the USA, and Australia and include Betatene, Western Biotechnology, AquaCarotene Ltd., Cyanotech Corp., and Nature Beta Technologies [35].

Dunaliella produces many carotenoid pigments with the dominant being beta-carotene and smaller amounts of lutein and lycopene [36]. Some strains of *Dunaliella* contain up to 14% of beta-carotene on dry weight basis. The total carotenoid content of *Dunaliella* varies with the physicochemical parameters and growth conditions. In optimal environmental condition, it can yield around 400 mg beta-carotene/m² of cultivation area [37]. Carotenoids from *Dunaliella* are potent free radical scavengers that reduce levels of lipid peroxidation and enzyme inactivation, thereby restoring enzyme activity. Research has shown beta-carotene to prevent cancer of various organs like the lungs, cervix, pancreas, colon, rectum, breast, prostate, and ovary by means of antioxidant activity [36]. It has also been shown to promote regression of certain types of cancer. Supplements of *Dunaliella* have also shown excellent hepatoprotective effects and reduced the occurrence of liver lesions [38].

3.4 *Haematococcus pluvialis*

Haematococcus pluvialis (*H. pluvialis*) is unicellular biflagellate freshwater green microalga. This species is known for its ability to accumulate large quantities of

Composition content (% of DW)	Green stage	Red stage
Proteins	29–45	17–25
Lipids (% of total)	20–25	32–37
Neutral lipids	59	51.9–53.5
Phospholipids	23.7	20.6–21.1
Glycolipids	11.5	25.7–26.5
Carbohydrates	15–17	36–40
Carotenoids (% of total)	0.5	2–5
Neoxanthin	8.3	n.d
Violaxanthin	12.5	n.d
β-carotene	16.7	1
Lutein	56.3	0.5
Zeaxanthin	6.3	n.d
Astaxanthin (including esters)	n.d	81.2
Adonixanthin	n.d	0.4
Adonirubin	n.d	0.6
Canthaxanthin	n.d	5.1
Echinenone	n.d	0.2
Chlorophylls	1.5	2 0

Adapted from [16].

Table 3.
Composition of H. pluvialis biomass in green and red cultivation stages.

strong antioxidant astaxanthin (up to 2–3% on dry weight) under any conditions. The principal commercial astaxanthin-producing microalga is *H. pluvialis* [9, 26]. Astaxanthin is used as a nutritional supplement and anti-inflammatory and anti-cancer agent for cardiovascular diseases and is recently recorded to prevent diabetes and neurodegenerative disorders and stimulates immunization. It also has anti-inflammatory properties and is used for various commercial applications in the dosage forms as biomass, capsules, creams, granulated powders, oils, soft gels, syrups, and tablets [39].

Photoautotrophic culture of *H. pluvialis* is mainly carried out in open raceway ponds or closed photobioreactors. The accumulation of astaxanthin is affected by environmental factors such as light, temperature, pH, salt concentration, and nutritional stresses. The cellular composition of *H. pluvialis* varies notably between its “green” and “red” stages of cultivation [40]. Specific biochemical characters of green and red stage of *H. pluvialis* are described in **Table 3**. The table shows that carbohydrate content in the green stage is approximately half of the red stage. *H. pluvialis* can accumulate approximately 5% DW of astaxanthin which is considered as a natural source of this high-value carotenoid protein [37].

3.5 *Aphanizomenon*

Aphanizomenon is a prokaryotic cyanobacterium commonly found in freshwater systems. There are approximately 500 tons of dried *Aphanizomenon* produced annually for use in food and pharmaceutical industries [41]. The dominant production source of *Aphanizomenon* in North America is Upper Klamath Lake and Klamath Falls, Oregon, and currently constitutes a significant part of the health food supplement industry throughout North America. *Aphanizomenon* contains a significant amount of C-phyococyanin, a light-harvesting pigment. It has antioxidant and anti-inflammatory properties [42]. *Aphanizomenon* also exhibits high

Component	<i>Spirulina</i>	<i>Dunaliella</i>	<i>Haematococcus</i>	<i>Chlorella</i>	<i>Aphanizomenon</i>
Protein	63	7.4	23.6	64.5	1.0
Fat	4.3	7.0	13.8	10.0	3.0
Carbohydrate	17.8	29.7	38.0	15.0	23.0
Chlorophyll	1.15	2.2	0.4 (red) 1.1 (green)	5.0	1.8
Magnesium	0.319	4.59	1.14	0.264	0.2
B-carotene	0.12	1.6	0.054	0.086	0.42
Vitamin B1 (thiamin)	0.001	0.0009	0.00047	0.0023	0.004
Vitamin B2 (riboflavin)	0.0045	0.0009	0.0017	0.005	0.0006
Vitamin B3 (niacin)	0.0149	0.001	0.0066	0.025	0.025
Vitamin B5 (pantothenic acid)	0.0013	0.0005	0.0014	0.0019	0.0008
Vitamin B6 (pyridoxine)	0.00096	0.0004	0.00036	0.0025	0.0013
Vitamin B9 (folic acid)	0.000027	0.00004	0.00029	0.0006	0.0001
Vitamin B12 (cobalamine)	0.00016	0.000004	0.00012	0.000008	0.0006

Adopted from [16].

Table 4.
Summary of referenced biochemical constitutions of average nutritional compositions (g per 100 g DW).

hypo-cholesterolemic activity, significantly greater than soybean oil, which decreases blood cholesterol and triglyceride levels [43–45]. It also produces polyunsaturated fatty acids (i.e., omega 3 and omega 6), a deficiency of which has been linked to immunosuppression, arthritis, cardiovascular diseases, mental health issues, and dermatological problems [16]. A summary on biochemical characters of all these economically important species is described in **Table 4**.

4. Summary

As the human population continues to increase, the demand for nutritive food and health products increases concomitantly. The sources of nutritive biomass that can meet this demand are pursued rampantly. Their wide diversity, fast growth, and diverse uses make them easily accepted for commercial culture. Microalgae require much fewer resources as compared to other crops. The role of algae in human health and nutrition will continually increase with additional research in the areas of health benefits and culturing. The usage of currently produced algae primarily includes food, food additives, aquaculture, colorants, cosmetics, pharmaceuticals, and nutraceuticals. Very few algal species are being cultivated for human use. There are likely more species of algae that have not been identified than ones that have and those still numbers in the thousands. Therefore, the potential for algal use in the realms of food consumption, health supplements, energy production, and many more is likely to intensify in the years to come.

Author details


Joshi Nilesh Hemantkumar^{1*} and Mor Ilza Rahimbhai²

¹ College of Fisheries Science, Junagadh Agricultural University, Veraval, Gujarat, India

² Department of Botany, H & H. B College of Science, Saurashtra University, Rajkot, Gujarat, India

*Address all correspondence to: nileshhj@gmail.com

IntechOpen

© 2019 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

References

- [1] Wiessner W, Schnepf E, Starr RC. Algae, Environment and Human Affairs. Bristol: Biopress Ltd.; 1995
- [2] Singh J, Gu S. Commercialization potential of microalgae for production of biofuels. *Renewable and Sustainable Energy Reviews*. 2010;**14**:2596-2610
- [3] Varfolomeev SD, Wasserman LA. Microalgae as source of biofuel, food, fodder and medicine. *Applied Biochemistry and Microbiology*. 2011; **47**:789-807
- [4] Norton TA, Melkonian M, Andersen RA. Algal biodiversity. *Phycologia*. 1996;**35**:308-326
- [5] Rindi F, McIvor L, Sherwood AR, Friedl T, Guiry MD, Sheath RG. Molecular phylogeny of the green algal order Prasiolales (Trebouxiophyceae, Chlorophyta). *Journal of Phycology*. 2007;**43**:811-822
- [6] Lewis LA, Flechtner VR. Green algae (*Chlorophyta*) of desert microbiotic crusts: Diversity of North American taxa. *Taxon*. 2002;**51**:443-451
- [7] Broady PA. Diversity, distribution and dispersal of Antarctic terrestrial algae. *Biodiversity and Conservation*. 1996;**5**:1307-1335
- [8] Sharma NK, Rai AK, Singh S, Brown RM. Airborne algae: Their present status and relevance. *Journal of Phycology*. 2007;**43**:615-627
- [9] Fritsch F. The subaerial and freshwater algal flora of the tropics. A phyto-geographical and ecological study. *Annals of Botany*. 1907;**21**: 235-275
- [10] Lewis LA, McCourt RM. Green algae and the origin of land plants. *American Journal of Botany*. 2004;**91**: 1535-1556
- [11] Mobin S, Alam F. Biofuel production from algae utilizing wastewater. In: *Proceedings of 19th Australasian Fluid Mechanics Conference*; Article No 27; 2014
- [12] Borowitzka MA. Algae as food. In: BJB W, editor. *Microbiology of Fermented Foods*. Thomson Science; 1998. pp. 585-602
- [13] Hudek K, Davis LC, Ibbini J, Erickson L. Commercial products from algae. In: Bajpai et al., editors. *Algal Biorefineries*. Dordrecht: Springer Science+Business Media; 2014. pp. 275-295
- [14] Mobin S, Alam F. Some promising microalgal species for commercial applications: A review. *Energy Procedia*. 2017;**110**:510-517
- [15] Ciferri O, Tiboni O. The biochemistry and industrial potential of *Spirulina*. *Annual Review of Microbiology*. 1985;**39**: 503-526
- [16] Bishop WM, Zubeck HM. Evaluation of microalgae for use as nutraceuticals and nutritional supplements. *Journal of Nutrition & Food Sciences*. 2012;**2**:147
- [17] Richmond A. *Spirulina*. In: Borowitzka MA, Borowitzka LJ, editors. *Micro-Algal Biotechnology*. Cambridge: Cambridge University Press; 1988. pp. 85-121
- [18] Belay A, Ota Y, Miyakawa K, Shimamatsu H. Current knowledge on potential health benefits of *Spirulina*. *Journal of Applied Phycology*. 1993;**5**: 235-241
- [19] Gershwin ME, Belay A. *Spirulina in Human Nutrition and Health*. Boca Raton, FL, USA: CRC Press; 2008

- [20] Becker EW, Jakover B, Luft D, Schmuelling RM. Clinical and biochemical evaluations of the alga *Spirulina* with regard to its application in the treatment of obesity: A double-blind cross-over study. *Nutrition Reports International*. 1986; **33**:565-574
- [21] Takai Y, Hosoyamada Y, Kato T. Effect of water-soluble and water insoluble fractions of *Spirulina* over serum lipids and glucose resistance of rats. *Journal of Japan Society of Nutrition and Food Sciences*. 1991; **44**: 273-277
- [22] Iwata K, Inayama T, Kato T. Effects of *Spirulina platensis* on plasma lipoprotein lipase activity in fructose-induced hyperlipidemic rats. *Journal of Nutritional Science and Vitaminology (Tokyo)*. 1990; **36**:165-171
- [23] Kromhout D. Fish oil consumption and coronary heart disease. In: *Dietary LW3 and LW6 Fatty Acids: Biological Effects and Nutritional Effects and Nutritional Essentiality*. New York, USA: Plenum Publishing; 1989
- [24] Simopoulos AP. Omega-3 fatty acids in health and disease and in growth and development. *The American Journal of Clinical Nutrition*. 1991; **54**:438-463
- [25] Tang G, Suter PM. Vitamin A, nutrition, and health values of algae: *Spirulina*, *Chlorella*, and *Dunaliella*. *Journal of Pharmacy and Nutrition Sciences*. 2011; **1**:111-118
- [26] de Caire GZ, de Cano MS, de Mule CZ, Steyerthal N, Piantanida M. Effect of *Spirulina platensis* on glucose, uric acid and cholesterol levels in the blood of rodents. *International Journal of Experimental Botany*. 1995; **57**:93-96
- [27] Romay C, Armesto J, Ramirez D, González R, Ledon N. Antioxidant and anti-inflammatory properties of C-phycocyanin from blue-green algae. *Inflammation Research*. 1998; **47**:36-41
- [28] Borowitzka MA. Commercial production of microalgae: Ponds, tanks, tubes and fermenters. *Journal of Biotechnology*. 1999; **70**:313-321
- [29] Zhu LD, Hiltunen E, Antila E, Zhong JJ, Yuan ZH, Wang ZM. Microalgal biofuels: Flexible bioenergies for sustainable development. *Renewable and Sustainable Energy Reviews*. 2014; **30**:1035-1046
- [30] Belay A, Ota Y, Miyakawa K, Shimamatsu H. Production of high quality *Spirulina* at Earthrise Farms. In: Phang SM, Lee K, Borowitzka MA, Whitton B, editors. *Algal Biotechnology in the Asia-Pacific Region*. University of Kuala Lumpur; 1994. pp. 92-102
- [31] Merchant RE, Andre CA. A review of recent clinical trials of the nutritional supplement *Chlorella pyrenoidosa* in the treatment of fibromyalgia, hypertension, and ulcerative colitis. *Alternative Therapies in Health and Medicine*. 2001; **7**:79-91
- [32] Sansawa H, Takahashi M, Tsuchikura S, Endo H. Effect of *Chlorella* and its fractions on blood pressure, cerebral stroke lesions, and life-span in strokeprone spontaneously hypertensive rats. *Journal of Nutritional Science and Vitaminology (Tokyo)*. 2006; **52**:457-466
- [33] Sano T, Tanaka Y. Effect of dried, powdered *Chlorella vulgaris* on experimental atherosclerosis and alimentary hypercholesterolemia in cholesterol-fed rabbits. *Artery*. 1987; **14**: 76-84
- [34] Tiberg E, Dreborg S, Björkstén B. Allergy to green algae (*Chlorella*) among children. *The Journal of Allergy and Clinical Immunology*. 1995; **96**:257-259

- [35] Del Campo JA, García-González M, Guerrero MG. Outdoor cultivation of microalgae for carotenoid production: Current state and perspectives. *Applied Microbiology and Biotechnology*. 2007; **74**:1163-1174
- [36] Chidambara Murthy KN, Vanitha A, Rajesha J, Mahadeva Swamy M, Sowmya PR. In vivo antioxidant activity of carotenoids from *Dunaliella salina*—A green microalga. *Life Sciences*. 2005; **76**:1381-1390
- [37] Finney KF, Pomeranz Y, Bruinsma BL. Use of algae *Dunaliella* as a protein supplement in bread. *Cereal Chemistry*. 1984; **61**:402-406
- [38] Hsu YW, Tsai CF, Chang WH, Ho YC, Chen WK. Protective effects of *Dunaliella salina*—A carotenoids-rich alga, against carbon tetrachloride-induced hepatotoxicity in mice. *Food and Chemical Toxicology*. 2008; **46**:3311-3317
- [39] Guerin M, Huntley ME, Olaizola M. *Haematococcus* astaxanthin: Applications for human health and nutrition. *Trends in Biotechnology*. 2003; **21**: 210-216
- [40] Shah MMR, Liang Y, Cheng JJ, Daroch M. Astaxanthin-producing green microalga *Haematococcus pluvialis*: From single cell to high value commercial products. *Frontiers in Plant Science*. 2016; **7**:531
- [41] Dashwood R, Guo D. Protective properties of chlorophylls against the covalent binding of heterocyclic amines to DNA in vitro and in vivo. *Princess Takamatsu Symposia*. 1995; **23**:181-189
- [42] Kushak RI, Drapeau C, Van Cott EM, Winter HH. Favorable effect of blue-green algae *Aphanizomenon flos-aquae* on rat plasma lipids. *Journal of the American Nutraceutical Association*. 2000; **2**:59-65
- [43] Miyamoto E, Tanioka Y, Nakao T, Barla F, Inui H. Purification and characterization of a corrinoid-compound in an edible cyanobacterium *Aphanizomenon flos-aquae* as a nutritional supplementary food. *Journal of Agricultural and Food Chemistry*. 2006; **54**:9604-9607
- [44] Liang S, Liu X, Chen F, Chen Z. Current microalgal health food R & D activities in China. *Hydrobiologia*. 2004; **512**:45-48
- [45] Pulz O, Gross W. Valuable products from biotechnology of microalgae. *Applied Microbiology and Biotechnology*. 2004; **65**:635-648