

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Polymerase Chain Reaction

*Shaheen Shahzad, Mohammad Afzal, Shomaila Sikandar
and Imran Afzal*

Abstract

Polymerase chain reaction (PCR) is an efficient and one of the most common methods used in biological sciences for in vitro multiplication of a target DNA molecule. The technique has significantly contributed in changing and developing different fields of biological sciences since 1980s. PCR has a vital role in supporting the processes involved in genetic engineering, particularly the cloning of DNA fragments used to modify the genomes of microorganisms, animals, and plants. Consequently, the technique has numerous applications in fundamental and applied research in medicine agriculture, environment, and bio-industry. The main focus of this chapter is to describe briefly the principles, methodology, various types, and applications of PCR in different fields. Besides, different components of PCR, trouble shooting during the execution, and limitations of the techniques are also outlined.

Keywords: PCR, primer, DNA template, nucleotides, sequence, polymerase

1. Introduction

Polymerase chain reaction (PCR) is one of the most commonly used method in modern molecular biology. The technique was developed by the Nobel laureate, Kary Mullis, in 1984. It is an in vitro process to multiply a target molecule of DNA with extreme precision, making it easy to be handled and examined by routine molecular biological methods [1–4]. Since its inception, PCR has significantly contributed in changing and developing biological sciences. The first PCR machine was introduced in market in 1988. The Human Genome Project has been result of PCR based approaches [5, 6]. Owing to its wide range of applications, numerous variants of PCR techniques have emerged over the past few decades [2–4].

PCR begins with the separation (denaturation) of the strands of a target DNA molecule (known as template) followed by annealing (hybridization) of oligonucleotide primers to the target template. The annealed primers provide a start for the DNA polymerase point to add new nucleotides (deoxynucleoside triphosphates or dNTPs). The sequence of nucleotides to be added is determined by the template. This entire process of the amplification of template, i.e., separation, annealing, and polymerization, is accomplished in vitro by cyclical alterations of temperature [2, 4, 7–10]. DNA polymerases used in PCR originate in thermophilic microorganisms, largely archaea, thriving temperature between 41 and 122°C. This ability to withstand high temperature is required in PCR to melt or separate the double-stranded DNA. Today, PCR has become a main stay in biotechnology, genomics, diagnostics, systematics, and many more areas [2–6].

2. Principles of PCR

PCR typically involves a series of 20–40 repeated temperature changes, called cycles, with each cycle usually comprising three discrete temperature shifts (Figure 3) [2, 4, 7–10].

1. Denaturation: 94–96°C
2. Primer annealing (depending on the primer): 45–60°C
3. Primer extension: usually 72°C

The cycling steps often start and end with a temperature step called “hold” where product extension is performed at (>90) and (~72°C), respectively. The final product is kept at 4°C before its analysis or storage. The most of PCR methods typically amplify DNA fragments of up to ~10 kilo base pairs (kb). However, some techniques can amplify up to 40 kb.

3. The components of PCR

Setting up a basic PCR requires many ingredients, reagents, and conditions which are described below (Figure 1) [2, 4, 7–10].




Figure 1.
The component of PCR.

3.1 DNA template

The double-stranded DNA molecule amplified by PCR is called the target or template DNA. The template defines the sequence in which new nucleotides are added during the PCR process [11]. This process is carried out in vitro by cyclically varying temperature, enabling separation of DNA strands, hybridization of primers, and polymerization.

DNA isolated from any source can be used as a template for PCR provided that it contains the target sequence. The DNA used in PCR can be isolated from blood, tissue, forensics specimens, paleontological samples, or microbial/tissue cells grown in the lab. Whatever the source, we need to have some information of the target DNA sequence, so that primers for PCR can be designed [2, 12]. The PCR primers can be designed very easily nowadays owing to the plethora of software tools that only requires target sequence information. If the sequence information is not known, the designing of primers becomes very challenging. This problem can be circumvented by using degenerate primers [2].

3.2 Primers

Primers are single-stranded DNA molecules usually synthesized commercially, i.e., polynucleotides of variable sizes [2, 7–10]. These short polynucleotide DNA strands have a free 3' hydroxyl group, also called as 3' end. The free 3' hydroxyl group on the primer is needed by the DNA polymerase to add new nucleotides during the polymerization process, thereby synthesizing a new complementary strand [12, 13]. The binding of DNA primer to the target requires the separation of two complementary DNA strands (Denaturation) which is generally achieved by heating process. To perform PCR, two primers are needed to enhance both the strands of the template: a primer for one strand (or sense strand), called the “forward primer,” which is the beginning of the template, and another primer for the complementary strand (or the antisense strand) called the “reverse primer.” Thus, both the primers bind to 5' ends of the sense and antisense strand.

The length of primers plays an important role in correctly identifying their designated target complementary regions. Increasing the length of primers improves their chances of matching the target (specificity). These primers are usually commercially synthesized with their size ranging between 18 and 25 nucleotides.

Primers should bind (anneal or hybridize) to the template with good specificity and strength to ensure amplification of the correct sequence. The specific temperature that is needed for primer annealing also depends on the primer sequences, e.g., the longer the primer, the higher the annealing temperature. Therefore, the maximum specificity and efficiency of PCR depends on optimal primer sequences and appropriate primer concentrations [2, 7, 8]. This in return depends on the way primers are designed and used. Improper primers may amplify undesired DNA segments (non-specific products), lower the yield of specific products, or completely fail the results of PCR. These undesired outcomes can be circumvented by designing and validating primers that preferentially bind to their target sequences. The online IDT Sci Tools Software Oligo Analyzer 3.1 and Primer Quest are invaluable aids both in primer design and validation [14]. These software tools also ensure that the two primers do not contain sequences that are complementary to each other. If primers contain self-complementary sequences, then hybridization will occur to each other, and they form “primer-dimmers.” Consequently, the primers will fail to bind to their target template, leading to a compromised PCR efficiency. In addition, presence of complementary sequences within a primer leads to the formation of hairpin loop structures [15].

As the bonding of guanine and cytosine bases (GC) is stronger than that between adenine and thymine (AT) bases, primers having GC at 3' end should be preferred for a strong bonding with the template [16]. However, the primers should not contain runs of three or more C or G bases, as this may lead to nonspecific binding to G- or C-rich sequences (mispriming) in the DNA which is not the target sequence [17].

3.3 DNA polymerase

Discovery of DNA polymerase in 1955 was the onset of PCR technology, which exploits the ability of bacterial DNA polymerase to make a complementary strand of a target DNA [2, 4, 7, 8, 18]. DNA polymerase starts making a new DNA from the 3' end of the template. The 3' end of the two template strands is where the primers bind which are then extended by the DNA polymerase. The most commonly used DNA polymerase is Taq DNA polymerase isolated from *Thermus aquaticus*, a thermophilic bacterium. Taq polymerase extends the DNA chain by adding ~1.0 kb per min with the enzymatic half-life achieved at 95°C in 40 minutes. Alternatively, the DNA polymerase from *Pyrococcus furiosus*, called Pfu, is also used widely due to its 3'–5' exonuclease activity (proofreading) which is not present in Taq DNA polymerase. Proofreading allows Pfu to remove incorrectly added nucleotide during polymerization and therefore to synthesize new DNA with minimum errors. A recombinant DNA polymerase, KOD DNA polymerase, derived from the thermophilic solfatara bacterium *Thermococcus kodakarensis* KOD1 type strain, functions optimally at 85°C with 3'–5' exonuclease proofreading activity, resulting in blunt-ended DNA products [19, 20]. KOD DNA polymerase exhibits high fidelity and processivity for small amplicons. However, for the amplicons over 5 kb, the amplification is lowered due to strong 3'–5' exonuclease activity of the enzyme [5]. This problem can be solved by mixing wild type with the mutant form of the enzyme (with lower 3'–5' exonuclease activity), which can result in more correct amplification of the amplicons between 5 and 15 kb [21]. Other sources of DNA polymerases used in PCR include thermophilic species like *Thermus thermophilus* (Tth) and *Thermus flavus* (Tfl) [18].

3.4 Nucleotides

PCR requires four different deoxynucleoside triphosphates or dNTPs to synthesize new DNA strands: adenine(A), guanine(G), cytosine(C), thymine(T). The dNTPs are usually provided at a concentration of 200 µM in the reaction mixture [22]. The concentration of these four dNTPs must be equal in the reaction mixture, as unequal concentration of even a single dNTPs leads to misincorporation of nucleotides by the DNA polymerase.

3.5 Buffer solution

The function of PCR buffer solution is to provide suitable conditions and chemicals to the DNA polymerase for optimal activity and stability [23]. The buffers often contain Tris-HCl, KCl, and sometimes MgCl₂. PCR buffers are often available in 10× concentration and are sometimes Taq formulation-specific including the compounds shown in **Table 1**.

3.6 Monovalent cations

Potassium chloride (KCl) is normally used in a PCR amplification of DNA fragments at a final concentration of 50 mM [24].

| Component | Function |
|-----------------------------------|--|
| 100 mM Tris-HCl (pH 8.8 at 25 °C) | Maintains reaction pH |
| 500 mM KCl | Stabilizes primer-template annealing |
| 15 mM MgCl ₂ | Cofactor for DNA polymerase |
| 0.8% (v/v) Nonidet P40 (Optional) | Suppresses secondary structure formation |

(Thermo Fisher Scientific™B16: <https://www.thermofisher.com/order/catalog/product/B16>)

Table 1.
Concentrations of PCR buffers.

3.7 Divalent cations

Magnesium ions are needed by the DNA polymerase enzyme as a cofactor. The divalent cations may include magnesium or manganese ions; generally, Mg²⁺ is used, but Mn²⁺ can be utilized for PCR-mediated DNA mutagenesis, as higher Mn²⁺ concentration increases the error rate during DNA synthesis [25].

3.8 PCR tube

PCR is performed in a small, thin-walled plastic tube called PCR tube. The tube is specifically designed to permit favorable thermal conductivity equilibration during thermal cycling.

3.9 Thermal cycler

A thermal cycler or thermocycler is a device used to rapidly heat and cool the reaction mixtures and cycle them between the three PCR temperature steps [26]. Many modern thermocyclers employ the Peltier effect to achieve this temperature ramping, which is done by reversing the electric current [11]. Modern thermocyclers are also provided with heated lids to prevent condensation of reaction mixture during PCR operation. Older thermocyclers lacked this feature, and the evaporation was prevented by applying oil or wax balls on the surface of PCR mixture.

4. Procedure

Each cycle or round of PCR comprises three major steps, viz., denaturation, annealing, and extension, repeated for 30 or 40 cycles on a thermocycler (**Figure 2**) [2, 4, 7–10]. A number of parameters determine the range of temperature and the duration of each cycle step (**Figure 3**), e.g., the polymerase used for DNA synthesis; melting temperature (T_m) of the primers; and the concentration of reagents used, i.e., divalent ions and dNTPs. The melting temperature depends on the length and specific nucleotide sequence of a primer. At T_m, half of the DNA molecules are in the single-stranded form.

4.1 Denaturation

It is the first cycling step that involves heating the reaction mixture to 94–98°C for 20–30 seconds. Such higher temperature disrupts the hydrogen bonding of the two complementary strands to produce the single-stranded DNA templates. Thus, denaturation prepares the DNA template for the binding of primers.




Figure 2.
A basic PCR protocol—DNA synthesis cycle.




Figure 3.
PCR—a simple thermocycling protocol.

4.2 Annealing

After denaturation, the next cycling step is annealing, in which the temperature of the PCR reaction is decreased to 50–65°C and kept for 20–60 seconds. This promotes hybridization between primers and single-stranded templates. Optimal annealing occurs at temperatures that are 3–5°C less than the primer T_m . The primers should have sufficient length and GC content to strongly bind to their target template during annealing.

4.3 Extension

DNA polymerase synthesizes (polymerizes) new DNA molecule by adding dNTPs complementary to the template bases in a 5′–3′ direction. The temperature and extension time depend on the type of DNA polymerase used: Taq polymerase performs optimally between 75 and 80°C. However, the enzyme is routinely used at 72°C. The extension time also depends on the length of the template.

A single cycle of PCR comprises the entire processes of denaturation, annealing, and extension/elongation. Under ideal conditions (optimal temperature ramping, presence of substrates/reagents, absence of inhibitors), the quantity of target DNA is doubled at the end of each cycle, resulting in exponential amplification of the specific DNA segment.

Final elongation: This single step is optional and performed at 70–74°C. The final elongation may take for 5–15 minutes after the last PCR cycle and allows any remaining single-stranded DNA to be fully extended.

Final hold: The final step may be employed for short-term storage of the reaction by cooling the reaction chamber to 4–15°C for an indefinite time.

5. Magnitude of amplification

Multiple cycling steps of a PCR can exponentially increase the copies of target DNA template to millions. The number of DNA copies produced by a PCR can be calculated using the formula “ 2^n ”, where n is the number of cycles [27, 28]. For example, a PCR set for 36 cycles results in 68 billion copies of the template. Under optimal conditions, even with minimal efficiency, a PCR in 50 ml volume may produce 0.2 mg of 150 bp DNA from 100 template molecules after 35–40 cycles, with the molar weight of the fragment equal to 99,000 Da.

6. Validating PCR

The degree of a PCR success can be determined in many ways [3, 29–36]:

1. Ethidium bromide (EtBr) can be used for the staining of amplified DNA product [31, 32]. It has UV absorbance maxima at 300 and 360 nm, and an emission maximum at 590 nm, and being a DNA intercalator, EtBr inserts itself between the base pairs in the double helix. The detection limit of DNA bound to ethidium bromide is 0.5–5.0 ng/band.
2. A three primer combination approach can provide a more cost-effective end-labeling of PCR products: (i) fluorescently labeled universal primer, (ii) modified locus-specific primers, and (iii) 5' universal primer sequence tails [34].
3. Agarose gel electrophoresis: The most commonly employed validating method, gel electrophoresis, makes use of electric current to separate charged molecules like DNA using gel as molecular sieve. Gelling agents can be agarose (for DNA >500 bp) or polyacrylamide (<500 bp). Different DNA sequences are separated out based on their sizes. DNA staining dyes (like EtBr) are applied to the gel to help visualize the DNA bands using UV transilluminator [34, 36]. The presence of a correct size DNA band (confirmed using a DNA ladder) indicates that the target sequence was present and that the PCR has amplified a correct product. Absence of any DNA band indicates that the target DNA was absent, while the presence of incorrect size DNA band indicates production of a spurious product [37].
4. The direct sequencing is often not practiced due to inaccessibility or cost of DNA sequencer or even the time needed to undertake such an analysis. However, restriction enzyme digestion can also be used to assess the sequence of an amplicon indirectly [29].

7. Types of PCR

Owing to ever-growing applications, a wide variety of PCR techniques have emerged over the past few decades [2–4]. Some of the variants are mere optimization close to the basic PCR to fulfill the specific needs. Others have undergone massive modifications to suit novel applications in different biological, biomedical, agricultural, and environmental fields [6].

7.1 Conventional PCR

This is a standard PCR in which a single-primer pair is used to bind to the two separated target strands. The primers also define the target sequences that will be copied. The PCR generates millions of copies of the target DNA sequences [2–4].

7.2 Multiplex-PCR

It is a special type of PCR for the detection of pathogenic microorganisms by using several pairs of primers annealing to different target sequences in a single sample [2–4, 38]. The multiplex-PCR is mainly used to identify exonic or intronic sequences to detect mutations, deletions, insertions, and rearrangements in pathogenic specimens.

7.3 Nested-PCR

It is used to increase the specificity of DNA amplification by reducing the nonspecific amplification [2–4, 39]. The two sets of primer pairs are used for a single locus point in two successive PCR reactions. The first round of PCR is performed with a primer pair that anneals to the sequence that flanks the target region. This generates a much larger DNA product that includes the target sequence. The second PCR is performed with a primer pair that precisely anneals to the target sequence, internal to the product of first PCR. This ensures that only the correct product is amplified in the second PCR [7]. Although Nested PCR improves specificity of amplification, it has disadvantage like primer-dimer formations [40].

7.4 Real-time PCR/quantitative PCR (qPCR)

A qPCR is a technique used to quantify the amplification of a template DNA in real time during the PCR reaction. This type of PCR is commonly employed to estimate the number of DNA targets present in a sample or to study and compare the gene expression [7, 37]. When real-time PCR is used quantitatively (qPCR), the amount of amplification is measured either by using a nonspecific fluorescent dyes or sequence-specific DNA oligonucleotide fluorescent probes [4, 41]. When quantitative PCR is used above/below a certain amount of DNA molecules, it is called semi quantitative real-time PCR. Although the quantitative real-time PCR has many applications, it is more frequently used in basic research and diagnostic purposes. There is a growing industrial use of the technique, e.g., quantification of microbial load in processed foods, detection of GMOs, quantification of pathogenic viruses, etc. [42–44].

7.5 Hot start/cold finish PCR

This technique reduces nonspecific amplification during the initial stages of a PCR [4, 7, 45]. To prevent nonspecific amplification at lower temperatures, hybrid polymerases are used which remain inactive at ambient temperature and is only activated at higher temperatures. Inhibition of the polymerase activity at ambient temperature is done by using an antibody or covalently bound inhibitors. Simply, in this technique the reaction components are heated to the DNA melting temperature (e.g., 95°C) before adding the polymerase.

7.6 Touchdown PCR (step-down PCR)

This type of PCR is designed to minimize nonspecific amplification by gradually decreasing the primer annealing temperature in the successive cycles. PCR is started with initial cycles having an annealing temperature 3–5°C higher than the primer T_m . The annealing temperature is then gradually decreased to 3–5°C lower below the T_m . The higher annealing temperature increases the specificity of the primers at initial stages of the reaction, while the lower temperature permits more efficient amplification later at the end [4, 7, 46].

7.7 Assembly PCR or polymerase cycling assembly (PCA)

This technique is used for the synthesis of long DNA molecules from long oligonucleotides with short overlapping segments, alternating between sense and antisense directions. The process begins with an initial PCR with primers that have an overlap, followed by a second PCR using the products of the first PCR as the template to generate the final full-length DNA structure [4, 7, 47].

7.8 Colony PCR

It is a convenient high-throughput technique used to confirm the addition of DNA insert in the recombinant clones and their uptake by the bacterial cell. A single set of insert specific primers are designed for the areas of the vector flanking the site where target DNA fragments are already inserted. This results in the amplification of the inserted sequences. The technique is used for the screening of bacterial colonies transformed with the recombinant vectors and to perform PCR without initially extracting the bacterial genomic DNA [3, 4, 7, 48].

7.9 Methylation-specific PCR (MSP)

It is a variant of PCR used to identify promoter hyper-methylation at CpG islands in cell lines and clinical samples, including fresh/frozen tissues. The target DNA is first treated with sodium bisulfite, which transforms the unmethylated cytosine bases in to uracil, which pair with adenosine of the PCR primers. The modified DNA is then amplified using two types of primers that only differ at their CpG islands. One primer set anneals to DNA with cytosine (corresponding to methylated cytosine), while the other anneals to DNA with uracil (corresponding to unmethylated cytosine). The MSP technique provides quantitative information about the methylation when used in quantitative PCR [3, 4, 7, 49, 50].

7.10 Inverse PCR

This type of PCR is used to detect the sequences that surround the target DNA (flanking sequences). It involves a series of restriction enzyme digestions and self-ligation. The primers amplify sequences at either end of the target by extending outward from the known DNA segment [4, 7, 51].

7.11 Reverse transcription-PCR (RTP)

In this technique, the PCR is preceded by a reaction converting RNA into cDNA using viral reverse transcriptase. The resulting cDNA is used as a template for a second conventional PCR. The technique is widely used in the detection of RNA viruses and to

study gene expression [7, 52, 53]. A variant of the RTP, called differential-display reverse transcription-PCR or RNA arbitrarily primed PCR (RAP-PCR), is used to study and compare the gene expression of organism grown under different conditions. The variant employs the use of short and random 10-mer or 11-mer radio-labeled primers that are annealed at low stringency conditions to promote the extension of random sequences during the first PCR cycle. This is followed by high-stringency cycles to extend the products of first cycle. The resulting products are analyzed using standard sequencing gels, and RAP-PCR fingerprints are visualized by autoradiography. The technique is extremely useful in studying tissue-specific and condition-specific gene expressions [54, 55].

8. Variants of PCR

In addition to the above mentioned techniques, numerous other variants of PCR are in use to serve a wide variety of research, diagnostic, and industrial needs, e.g., after exponential PCR, allele specific PCR, asymmetric PCR, arbitrary PCR, core sample PCR, degenerate PCR, dial-out PCR, digital PCR, high-fidelity PCR, hot start PCR, in silico PCR, inter-sequence PCR, ligation-mediated PCR, mini primer PCR, nanoparticle-PCR, overlap-extension PCR, solid-phase PCR, splicing by overlap/overhang extension PCR, suicide PCR, thermal asymmetric interlaced PCR, etc. Some of the important variants of PCR are described below:

8.1 Extreme PCR

In extreme PCR the concentration of primers and polymerase is increased 10–20 times; the amplification rate of instrument reaches about 0.4–2.0 s/. When the primers' concentration is more than 10 mol/L, the polymerase concentration is 1 mol/L, and the extreme PCR is suitable for rapid detection of virulent infectious and bioterrorism pathogens [56].

8.2 Photonic PCR

It is achieved by fast heating and based on energy conversion, thus shortening the PCR time. The specific process is carried out by using electronic resonance light emitting diode. The energy conversion process is more rapid than the conventional cooling process, causing amplification of target DNA within 5 min and thus making the PCR detection more convenient and fast [57, 58].

8.3 COLD-PCR

It is a low denatured temperature-PCR for enriching mutant genes by reducing the reactive temperature of PCR. The basic principle is founded on the base mismatch in any strand of DNA affecting the denaturation temperature. Therefore, the denaturation temperature of the mutant DNA is often lower than that of wild type DNA. The assay is often used for viral gene mutation [59] detection, cancer associated gene mutations (p53) [60, 61], EGFR, KRAS, etc.), beta globulin (HBB) mutations that cause beta thalassemia [62], etc.

8.4 Nanoparticle-PCR

Gold nanoparticles have superior electrical, optical, thermal, and catalytic activities and have the same properties as single-stranded binding proteins (ssb), which bind to single-stranded DNA and do not interact with double-stranded DNA. Therefore, the

amplification effect of high GC template can be significantly improved by adding gold nanoparticles as additives to slowdown or touchdown PCR reaction systems [63].

8.5 HPE-PCR

It is an amplification technique for templates with long DNA chains and large numbers of CTG repeats involving the increase in the denaturation temperature of PCR to solve the problem of high content of DNA (G+C) [64].

8.6 LATE-PCR

It generates high concentrations of single-stranded DNA that can be analyzed at the end point using probes which hybridize over a wide temperature range [65].

8.7 Digital PCR

Digital PCR (dPCR) enables precise and sensitive quantification of nucleic acids in a wide range of applications in both healthcare and environmental analysis. It is based on detection in two discrete optical channels, focused on the quantification of one or two targets within a single reaction [66]. The technique has become a promising quantification strategy that combines absolute quantification with high sensitivity.

9. Applications of PCR

The PCR technique and its several advanced variants act as powerful tools with specialized applications which were once impossible by the scientific world [67, 68]. This versatile technique brought enormous benefits and scientific developments such as genome sequencing, gene expressions in recombinant systems, and the study of molecular genetic analysis, including the rapid determination of both paternity and the diagnosis of infectious disease [69, 70]. It enables the *in vitro* synthesis of nucleic acids through which a DNA segment can be specifically replicated in a semiconservative way. It generally exhibits excellent detection limits [71, 72]. It has significantly transformed

| Application | Description | Reference |
|------------------------------------|---|-----------|
| Diagnosis of infections | PCR approaches are used to specifically and sensitively diagnose infections (bacterial, viral, protozoan, fungal). They are routinely used in clinical laboratories to confirm and quantify these infectious agents | [71, 74] |
| Diagnosis of genetic defects | PCR-based detection systems are used to accurately detect (before disease onset) and confirm (after the onset) many genetic disorders | [74–76] |
| Diagnosis and prognosis of cancers | PCR-based approaches can identify cancer genes and analyze their expression to determine genetic predisposition to certain cancers, confirmation of cancer type, their prognosis, and treatment | [74, 77] |
| Phylogenetics | Phylogenetic analysis of organisms routinely relies on PCR amplification of phylogenetic markers to identify and classify them | [78, 79] |
| Archeology | Ancient DNA (aDNA) recovered from archeological remains are usually degraded and are in low amounts. Such miniscule quantities of aDNA are amplified using PCR techniques to improve their quality and quantity to make them analyzable for archeological study | [80–82] |
| Recombinant DNA technology | PCR techniques are used to generate hybrid DNA with ease and precision. The techniques are also employed to clone DNA in to specific vectors to get protein expression | [83, 84] |

| Application | Description | Reference |
|------------------------------|--|--------------|
| Metagenomics | Gene-targeted metagenomics combines PCR with metagenomics to identify rarest members of a sampled community and rare genes in the community members | [85, 86] |
| Site-directed mutagenesis | PCR-based approaches are commonly used to insert mutations (deletions, additions, and substitutions) at specific locations in a gene to study role of specific amino acids in the structure and function of proteins | [87, 88] |
| Personalized medicine | PCR technologies are employed in pharmacogenomics and pharmacogenetics to track genetic markers that determine the response of individuals to treatments and are used to design tailor-made drugs and to prescribe drugs in effective doses | [74, 89] |
| Forensics sciences | The power of PCR is employed to amplify poor quality and quantity DNA samples from crime scenes and make them reliably analyzable. | [67, 68, 90] |
| DNA profiling | DNA profiling methods utilize PCR-based approaches to exploit the polymorphic nature of DNA (SNPs, DNA repeats, etc.) to study the structure and diversity ecological communities, phylogeny, and population genetics | [90] |
| Gene expression profiling | Reverse-transcriptase PCR and qPCR are routinely employed to profile the expression of genes and to validate transcriptome profiles generated through techniques like microarray and RNA-seq | [91–93] |
| Identifying medicinal plants | PCR-based DNA barcoding is a tool that utilizes specific DNA sequences to rapidly and accurately identify medicinal plants species from other morphologically similar plants. This approach is also used by ecologists and conservation biologists to identifying endangered and new species | [94] |
| Detecting GMO | PCR techniques are used to quickly and reliably track the presence of genetically modified organism in food and feed to ensure their regulation and protection of consumer rights | [95, 96] |
| Meat traceability | PCR methods are used to identifying and quantifying adulteration of meat in raw and processed food. | [97–99] |

Table 2.
The most important applications of PCR.

the scientific research and diagnostic medicine. Over the years, it has become a vital of clinical and diagnostic research. It has a wide range of applications in almost every field of science, for example, clinicians widely use the technique for disease diagnosis. Biologists, including agriculturists, clone and sequence genes using PCR and rapidly carry out sophisticated quantitative and genomic studies. Now for criminal identification, PCR assays are commonly employed. DNA fingerprinting is also used in paternity testing, where the DNA from an individual is matched with that of his possible children, siblings, or parents [67, 68]. Besides, PCR has enormous role in diagnosing genetic disease, whether inherited genetic changes or as a result of spontaneous genetic mutations, is becoming more common. Diseases can be diagnosed even before birth. Even PCR can also be employed with significant precision to predict cure of diseases [73]. The most important applications of PCR are summarized in **Table 2**.

10. Limitations of PCR

Since its discovery in 1980s, the PCR technique has brought about significant changes in biological sciences. Huge scientific undertakings like the Human Genome Project have been possible due to PCR-based approaches [5, 6]. It is a very sensitive

and flexible technique to amplify DNA of interest. A very small amount of the target DNA can be used as a starting material. Even old or degraded DNA samples may yield successful amplification. However, there is also a long list of PCR limitations. High-quality DNA amplification needs information about target DNA sequence. The sensitivity of PCR is also its major disadvantage since the very end result of a PCR is highly susceptible to contamination or false amplification. Therefore, amplification of DNA by PCR may not be 100% specific. Moreover, the specificity of amplification is dependent on physicochemical parameter, such as temperature and Mg^{++} concentration. The PCR is also inhibited by the presence of certain chemicals such as ethanol, phenol, isopropanol, detergent compounds like sodium dodecyl sulfate (SDS), high salt concentration, chelators, etc. There is an upper limit to the size of DNA that can be synthesized by PCR. Additionally, analysis and product detection usually take much longer time than the PCR reaction itself.

Author details

Shaheen Shahzad^{1*}, Mohammad Afzal², Shomaila Sikandar³ and Imran Afzal³

1 Genomics Research Lab, Department of Bioinformatics and Biotechnology, Faculty of Basic and Applied Sciences, International Islamic University, Islamabad, Pakistan

2 Consultant for Health Research and M&E, Pakistan

3 Department of Biology, Lahore Garrison University, Lahore, Pakistan

*Address all correspondence to: drshaheen@iiu.edu.pk

IntechOpen

© 2020 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

References

- [1] Sárosi I, Gerald E, Girish VN. Polymerase chain reaction and its application. *Orvosi Hetilap*. 1992;**133**:16-21
- [2] Bermingham N, Luettich K. Polymerase chain reaction and its applications. *Current Diagnostic Pathology*. 2003;**9**(3):159-164
- [3] Rajalakshmi S. Different types of PCR techniques and its applications. *International Journal of Pharmaceutical, Chemical and Biological Sciences*. 2017;**7**(3):285-292
- [4] Patel SV, Bosamia TC, Bhalani HN, Singh P, Kumar A. Polymerase chain reaction (PCR). *Agrbios*. 2015;**13**(9):10-12
- [5] Mullis KB. The unusual origin of the polymerase chain reaction. *Scientific American*. 1990;**4**:56-65
- [6] Garibyan L, Avashia N. Research techniques made simple: Polymerase chain reaction (PCR). *The Journal of Investigative Dermatology*. 2013;**133**(3):e6. DOI: 10.1038/jid.2013.1
- [7] Zádor E. The polymerase chain reaction. Theoretical course: Basic biochemical methods and ischemic heart models. Supported by: HURO/0901/069/2.3.1 HU-RO-DOCS. 2011. Available from: http://www.academia.edu/35361310/The_polymerase_chain_reaction_Theoretical_course_Basic_biochemical_methods_and_ischemic_heart_models_Supported_by_HURO_0901_069_2.3.1_HU-RO-DOCS
- [8] Dubey VK. Proteomics and Genomics. Lecture 37: Polymerase Chain Reaction. Available from: http://nptel.ac.in/courses/102103017/module37/lec37_slide1.htm [Accessed: 24 May 2018]
- [9] Saiki RK, Scharf S, Faloona F, Mullis K, Hoorn GT, Arnheim N. Polymerase chain reaction. *Science*. 1985;**230**:1350-1354
- [10] Joshi M, Deshpande JD. Polymerase chain reaction: Methods, principles and application. *International Journal of Biomedical Research*. 2010;**2**(1):81-97
- [11] Bell LE. Cooling, heating, generating power, and recovering waste heat with thermoelectric systems. *Science*. 12 Sep 2008;**321**(5895):1457-1461
- [12] Ye J, Coulouris G, Zaretskaya I, Cutcutache I, Rozen S, Madden TL. Primer-BLAST: A tool to design target-specific primers for polymerase chain reaction. *BMC Bioinformatics*. 2012;**13**(1):134
- [13] Olerup O, Zetterquist H. HLA-DR typing by PCR amplification with sequence-specific primers (PCR-SSP) in 2 hours: An alternative to serological DR typing in clinical practice including donor-recipient matching in cadaveric transplantation. *HLA*. 1992;**39**(5):225-235
- [14] Owczarzy R, Tataurov AV, Wu Y, Manthey JA, McQuisten KA, Almabrazi HG, et al. IDT SciTools: A suite for analysis and design of nucleic acid oligomers. *Nucleic Acids Research*. 2008;**36**(Suppl_2):W163-W169
- [15] Vallone PM, Butler JM. Auto dimer: A screening tool for primer-dimer and hairpin structures. *BioTechniques*. 2004;**37**(2):226-231
- [16] Li LC, Dahiya R. MethPrimer: Designing primers for methylation PCRs. *Bioinformatics*. 1 Nov 2002;**18**(11):1427-1431
- [17] Lexow P, Ragnhildstveit E, LingVitae AS. Method for characterising polynucleotides. United States Patent Application US 10/553,113 [Accessed: 02 April 2009]

- [18] Cline J, Braman JC, Hogrefe HH. PCR fidelity of Pfu DNA polymerase and other thermostable DNA polymerases. *Nucleic Acids Research*. 1 Sep 1996;**24**(18):3546-3551
- [19] Benson LM, Null AP, Muddiman DC. Advantages of *Thermococcus kodakaraensis* (KOD) DNA polymerase for PCR-mass spectrometry based analyses. *Journal of the American Society for Mass Spectrometry*. 1 Jun 2003;**14**(6):601-604
- [20] Morikawa M, Izawa Y, Rashid N, Hoaki T, Imanaka T. Purification and characterization of a thermostable thiol protease from a newly isolated hyperthermophilic *Pyrococcus* sp. *Applied and Environmental Microbiology*. 1 Dec 1994;**60**(12):4559-4566
- [21] Nishioka M, Mizuguchi H, Fujiwara S, Komatsubara S, Kitabayashi M, Uemura H, et al. Long and accurate PCR with a mixture of KOD DNA polymerase and its exonuclease deficient mutant enzyme. *Journal of Biotechnology*. 15 Jun 2001;**88**(2):141-149
- [22] Lorenz TC. Polymerase chain reaction: Basic protocol plus troubleshooting and optimization strategies. *Journal of Visualized Experiments*. 2012;**63**:3998. DOI: 10.3791/3998
- [23] Hwang IT, Kim YJ, Kim SH, Kwak CI, Gu YY, Chun JY. Annealing control primer system for improving specificity of PCR amplification. *BioTechniques*. 2003;**35**(6):1180-1191
- [24] Huggett J, O'Grady J, editors. *Molecular Diagnostics: Current Research and Applications*. Caister: Academic Press; 2014
- [25] Innis M, Gelfand D. Optimization of PCR: Conversations between Michael and David. In: *PCR Applications*. 1999. pp. 3-22
- [26] Lund K, Rothenberg BE. Applied Chemical, Engineering Systems Inc. Thermal Cycler. United States patent US 6,558,947 [Accessed: 06 May 2003]
- [27] Vos P, Hogers R, Bleeker M, Reijans M, Lee TV, Hornes M, et al. AFLP: A new technique for DNA fingerprinting. *Nucleic Acids Research*. 1995;**23**(21):4407-4414
- [28] Barnes WM. PCR amplification of up to 35-kb DNA with high fidelity and high yield from lambda bacteriophage templates. *Proceedings of the National Academy of Sciences*. 15 Mar 1994;**91**(6):2216-2220
- [29] Telenti A, Marchesi F, Balz M, Bally F, Böttger EC, Bodmer T. Rapid identification of mycobacteria to the species level by polymerase chain reaction and restriction enzyme analysis. *Journal of Clinical Microbiology*. 1993;**31**(2):175-178
- [30] Millipore Sigma. Technical Guide to PCR Technologies—Assay Optimization and Validation. Darmstadt, Germany: Merck KGaA; 2018. Available from: <https://www.sigmaaldrich.com/technical-documents/articles/biology/assay-optimization-and-validation.html> [Accessed: 30 May 2018]
- [31] Mariani BD, Martin DS, Chen AF, Yagi H, Lin SS, Taun RS. PCR—Molecular diagnostic technology for monitoring chronic osteomyelitis. *Journal of Experimental Orthopaedics*. 2014;**19**
- [32] Yap EP, McGee JO. Non isotopic SSCP detection in PCR products by ethidium bromide staining. *Trends in Genetics*. 1992;**8**(2):49
- [33] Nakayama H, Arakaki A, Maruyama K, Takeyama H, Matsunaga T. Single-nucleotide polymorphism analysis using fluorescence resonance energy transfer between DNA-labeling fluorophore, fluorescein isothiocyanate,

and DNA intercalator, POPO-3, on bacterial magnetic particles. *Biotechnology and Bioengineering*. 2003;**84**(1):96-102

[34] Blacket MJ, Robin C, Good RT, Lee SF, Miller AD. Universal primers for fluorescent labelling of PCR fragments—An efficient and cost-effective approach to genotyping by fluorescence. *Molecular Ecology Resources*. 2012;**12**(3):456-463

[35] Gariyan L, Avashia N. Research Techniques made simple-PCR. *The Journal of Investigative Dermatology*. 2013;**133**(3):6

[36] Querci M, Jermini M, van den Eede G, editors. Training Course on the Analysis of Food Samples for the Presence of Genetically Modified Organisms. JRC-European Commission, World Health Organization—Regional Office for Europe; 2006

[37] Huang T, Yu X, Gelbič I, Guan X. RAP-PCR fingerprinting reveals time-dependent expression of development-related genes following differentiation process of *Bacillus thuringiensis*. *Canadian Journal of Microbiology*. 19 Jun 2015;**61**(9):683-690

[38] Ibrahim WA, Ghany A, Nasef SA, Hatem ME. Comparative study on the use of RT-PCR and standard isolation techniques for the detection of *Salmonella* in broiler chicks. *International Journal of Veterinary Science and Medicine*. 2014;**2**:67

[39] Kawasaki ES. Amplification of RNA. In: Innis MA, Gelfand DH, Stunski JJ, editors. *PCR Protocols, a Guide to Methods and Applications*. New York: Academic Press; 1990. pp. 21-27

[40] Docker MF, Delvin RH, Richard J, Khattra J. Sensitive and specific PCR-assay detection of *Loma salmonae* (microsporea). *Diseases of Aquatic Organisms*. 1997;**29**:41-48

[41] Logan J, Edwards K, Saunders N. *Real-Time PCR: Current Technology and Applications*. Caister: Academic Press. 2009. ISBN: 978-1-904455-39-4

[42] Espy MJ, Uhl JR, Sloan LM, Buckwalter SP, Jones MF, Vetter EA, et al. Real-time PCR in clinical microbiology: Applications for routine laboratory testing. *Clinical Microbiology Reviews*. 2006;**19**(1):165-256

[43] Fillion M. *Quantitative Real-time PCR in Applied Microbiology*. Caister: Academic Press; 2012. ISBN: 978-1-908230-01-0

[44] Baldwin BG. Phylogenetic utility of the internal transcribed spacers of nuclear ribosomal DNA in plants: An example from the compositae. *Molecular Phylogenetics and Evolution*. 1992;**1**(1):3-16

[45] Chou Q, Russell M, Birch DE, Raymond J, Bloch W. Prevention of pre-PCR mis-priming and primer dimerization improves low-copy-number amplifications. *Nucleic Acids Research*. 1992;**20**(7):1717-1723

[46] Don RH, Cox PT, Wainwright BJ, Baker K, Mattick JS. "Touchdown" PCR to circumvent spurious priming during gene amplification. *Nucleic Acids Research*. 1991;**19**(14):4008

[47] Stemmer WP, Cramer A, Ha KD, Brennan TM, Heyneker HL. Single-step assembly of a gene and entire plasmid from large numbers of oligodeoxyribonucleotides. *Gene*. 1995;**164**(1):49-53

[48] Pavlov AR, Pavlova NV, Kozyavkin SA, Slesarev AI. In: Kieleczawa J, editor. *DNA Sequencing II: Optimizing Preparation and Cleanup*. Jones and Bartlett; 2006. pp. 241-257

[49] Herman JG, Graff JR, Myöhänen S, Nelkin BD, Baylin SB.

Methylation-specific PCR: A novel PCR assay for methylation status of CpG islands. Proceedings of the National Academy of Sciences of the United States of America. 1996;**93**(18):9821-9826

[50] Licchesi JD, Herman JG. Methylation-specific PCR. In: DNA Methylation. Humana Press; 2009. pp. 305-323

[51] Ochman H, Gerber AS, Hartl DL. Genetic applications of an inverse polymerase chain reaction. Genetics. 1988;**120**(3):621-623

[52] Morris T, Robertson B, Gallagher M. Rapid reverse transcription-PCR detection of hepatitis C virus RNA in serum by using the Taq Man fluorogenic detection system. Journal of Clinical Microbiology. 1996;**34**(12):2933-2936

[53] Sheridan GE, Masters CI, Shallcross JA, Mackey BM. Detection of mRNA by reverse transcription-PCR as an indicator of viability in *Escherichia coli* cells. Applied and Environmental Microbiology. 1998;**64**(4):1313-1318

[54] Ruijter JM, Ramakers C, Hoogaars WM, Karlen Y, Bakker O, Van den Hoff MJ, et al. Amplification efficiency: Linking baseline and bias in the analysis of quantitative PCR data. Nucleic Acids Research. 22 Feb 2009;**37**(6):e45

[55] Welsh J, Chada K, Dalal SS, Cheng R, Ralph D, McClelland M. Arbitrarily primed PCR fingerprinting of RNA. Nucleic Acids Research. 11 Sep 1992;**20**(19):4965-4970

[56] Farrar JS, Wittwer CT. Extreme PCR: Efficient and specific DNA amplification in 15-60 seconds. Clinical Chemistry. 1 Jan 2014;**61**(1):145-153

[57] Yu M, Cao Y, Ji Y. The principle and application of new PCR Technologies. In: IOP Conference Series: Earth and

Environmental Science; Decemver 2017 (Vol. 100, No. 1, p. 012065). IOP Publishing

[58] Son JH, Cho B, Hong S, Lee SH, Hoxha O, Haack AJ, et al. Ultrafast photonic PCR. Light: Science & Applications. Jul 2015;**4**(7):e280

[59] Wong DK, Tsoi O, Huang FY, Seto WK, Fung J, Lai CL, et al. Application of co-amplification at lower denaturation temperature-PCR (COLD-PCR) sequencing for the early detection of antiviral drug resistance mutations of hepatitis B virus. Journal of Clinical Microbiology. 20 Jun 2014;**52**(9):3209-3215

[60] Liu C, Lin J, Chen H, Shang H, Jiang L, Chen J, et al. Detection of HBV genotypic resistance mutations by coamplification at lower denaturation temperature PCR coupled with Sanger sequencing. Journal of Clinical Microbiology. 4 Jun 2014;**52**(8):2933-2939

[61] Lewandowska MA, Jóźwicki W, Jochymski C, Kowalewski J. Application of PCR methods to evaluate EGFR, KRAS and BRAF mutations in a small number of tumor cells in cytological material from lung cancer patients. Oncology Reports. 1 Sep 2013;**30**(3):1045-1052

[62] Galbiati S, Brisci A, Lalatta F, Seia M, Makrigiorgos GM, Ferrari M, et al. Full COLD-PCR protocol for noninvasive prenatal diagnosis of genetic diseases. Clinical Chemistry. 1 Jan 2011;**57**(1):136-138

[63] Li H, Huang J, Lv J, An H, Zhang X, Zhang Z, et al. Nanoparticle PCR: Nanogold-assisted PCR with enhanced specificity. Angewandte Chemie. 12 Aug 2005;**117**(32):5230-5233

[64] Orpana AK, Ho TH, Alagrund K, Ridanpää M, Aittomäki K, Stenman J. Novel heat pulse extension-PCR-based

method for detection of large CTG-repeat expansions in myotonic dystrophy type 1. *The Journal of Molecular Diagnostics*. 1 Jan 2013;**15**(1):110-115

[65] Pierce KE, Wangh LJ. Rapid detection and identification of hepatitis C virus (HCV) sequences using mismatch-tolerant hybridization probes: A general method for analysis of sequence variation. *BioTechniques*. Sep 2013;**55**(3):125-132

[66] Hindson BJ, Ness KD, Masquelier DA, Belgrader P, Heredia NJ, Makarewicz AJ, et al. High-throughput droplet digital PCR system for absolute quantitation of DNA copy number. *Analytical Chemistry*. 28 Oct 2011;**83**(22):8604-8610

[67] Green MR, Sambrook J, Sambrook J. *Molecular Cloning: A Laboratory Manual*. 4th ed. New York: Cold Spring Harbor Laboratory Press; 2012

[68] Lakshmi V, Sudha T, Rakhi D, Anilkumar G, Dandona L. Application of polymerase chain reaction to detect HIV-1 DNA in pools of dried blood spots. *Indian Journal of Microbiology*. 2011;**51**(2):147-152. DOI: 10.1007/s12088-011-0135-0

[69] Valones MAA, Guimarães RL, Brandão LAC, de Souza PRE, de Albuquerque Tavares Carvalho A, Crovela S. Principles and applications of polymerase chain reaction in medical diagnostic fields: A review. *Brazilian Journal of Microbiology*. 2009;**40**(1):1-11. DOI: 10.1590/S1517-83822009000100001

[70] Novais CM, Pires-Alves M, Silva FF. PCR in real time. *Revista Biotecnologia Ciência & Desenvolvimento - Edição nº*. 2004;**33**:11

[71] Speers DJ, Ryan S, Harnett G, Chidlow G. Diagnosis of malaria aided by polymerase chain reaction in two

cases with low-level parasitaemia. *Internal Medicine Journal*. Dec 2003;**33**(12):613-615

[72] Estrela C. *Metodologia científica: ciência, ensino, pesquisa*. Artes Médicas; 2005

[73] Cortelli S, Jorge AO, Querido SM, Cortelli JR. PCR and culture in the subgingival detection of *Actinobacillus actinomycetemcomitans*: A comparative study. *Brazilian Dental Science*. 12 Aug 2010;**6**(2)

[74] Dwivedi S, Purohit P, Misra R, Pareek P, Goel A, Khattri S, et al. Diseases and molecular diagnostics: A step closer to precision medicine. *Indian Journal of Clinical Biochemistry*. 1 Oct 2017;**32**(4):374-398

[75] Mahdieh N, Rabbani B. An overview of mutation detection methods in genetic disorders. *Iranian Journal of Pediatrics*. Aug 2013;**23**(4):375

[76] <http://www.genelink.com/gds/GDSintro.asp>

[77] Kondo M, Hoshi SL, Yamanaka T, Ishiguro H, Toi M. Economic evaluation of the 21-gene signature (Oncotype DX®) in lymph node-negative/positive, hormone receptor-positive early-stage breast cancer based on Japanese validation study (JBCRG-TR03). *Breast Cancer Research and Treatment*. 1 Jun 2011;**127**(3):739-749

[78] Klindworth A, Pruesse E, Schweer T, Peplies J, Quast C, Horn M, et al. Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation sequencing-based diversity studies. *Nucleic Acids Research*. 1 Jan 2013;**41**(1):e1

[79] Marthaler D, Raymond L, Jiang Y, Collins J, Rossow K, Rovira A. Rapid detection, complete genome sequencing, and phylogenetic

analysis of porcine deltacoronavirus. *Emerging Infectious Diseases*. Aug 2014;**20**(8):1347

[80] Malainey ME. Ancient DNA and the polymerase chain reaction. In: *A Consumer's Guide to Archaeological Science*. New York, NY: Springer; 2011. pp. 237-253

[81] Wales N, Andersen K, Cappellini E, Ávila-Arcos MC, Gilbert MT. Optimization of DNA recovery and amplification from non-carbonized archaeobotanical remains. *PLoS ONE*. 27 Jan 2014;**9**(1):e86827

[82] Del Gaudio S, Cirillo A, Di Bernardo G, Galderisi U, Thanassoulas T, Pitsios T, et al. Preamplification procedure for the analysis of ancient DNA samples. *The Scientific World Journal*. 2013;**2013**

[83] Elion EA, Marina P, Yu L. Constructing recombinant DNA molecules by PCR. *Current Protocols in Molecular Biology*. 2007 Apr;**78**(1):3-17

[84] Dubey AA, Singh MI, Jain V. Rapid and robust PCR-based all-recombinant cloning methodology. *PLoS ONE*. 23 Mar 2016;**11**(3):e0152106

[85] Rosen MJ, Callahan BJ, Fisher DS, Holmes SP. Denoising PCR-amplified metagenome data. *BMC Bioinformatics*. Dec 2012;**13**(1):283

[86] Lee S, Cantarel B, Henrissat B, Gevers D, Birren BW, Huttenhower C, et al. Gene-targeted metagenomic analysis of glucan-branching enzyme gene profiles among human and animal fecal microbiota. *The ISME Journal*. Mar 2014;**8**(3):493

[87] Rouached H. Efficient procedure for site-directed mutagenesis mediated by PCR insertion of a novel restriction site. *Plant Signaling & Behavior*. 1 Dec 2010;**5**(12):1547-1548

[88] Edelheit O, Hanukoglu A, Hanukoglu I. Simple and efficient site-directed mutagenesis using two single-primer reactions in parallel to generate mutants for protein structure-function studies. *BMC Biotechnology*. Dec 2009;**9**(1):61

[89] Cho SH, Jeon J, Kim SI. Personalized medicine in breast cancer: A systematic review. *Journal of Breast Cancer*. 1 Sep 2012;**15**(3):265-272

[90] Eppelen J, Lubjuhn T. *DNA Profiling and DNA Fingerprinting*. Springer Science & Business Media; 11 Dec 2012

[91] Mooney C, Raouf R, El-Naggar H, Sanz-Rodriguez A, Jimenez-Mateos EM, Henshall DC. High throughput qPCR expression profiling of circulating microRNAs reveals minimal sex- and sample timing-related variation in plasma of healthy volunteers. *PLoS ONE*. 23 Dec 2015;**10**(12):e0145316

[92] Sheibani-Tezerji R, Rattei T, Sessitsch A, Trognitz F, Mitter B. Transcriptome profiling of the endophyte *Burkholderia phytofirmans* PsJN indicates sensing of the plant environment and drought stress. *MBio*. 30 Oct 2015;**6**(5):e00621-e00615

[93] Condomines M, Hose D, Rème T, Requirand G, Hundemer M, Schoenhals M, et al. Gene expression profiling and real-time PCR analyses identify novel potential cancer-testis antigens in multiple myeloma. *The Journal of Immunology*. 1 Jan 2009;**183**(2):832-840

[94] Kress WJ, Erickson DL. DNA barcodes: genes, genomics, and bioinformatics. *Proceedings of the National Academy of Sciences*. 26 Feb 2008;**105**(8):2761-2762

[95] Fraiture MA, Herman P, Taverniers I, De Loose M, Deforce D, Roosens NH. Current and new

approaches in GMO detection:
Challenges and solutions. *BioMed
Research International*. 2015;2015

[96] Rabiei M, Mehdizadeh M,
Rastegar H, Vahidi H, Alebouyeh M.
Detection of genetically modified maize
in processed foods sold commercially
in Iran by qualitative PCR. *Iranian
Journal of Pharmaceutical Research*.
2013;12(1):25

[97] Han SH, Oh HS, Cho IC. Identifying
the species of origin in commercial
sausages in South Korea. *Journal
of Applied Animal Research*. 1 Jan
2017;45(1):179-184

[98] Ren J, Deng T, Huang W,
Chen Y, Ge Y. A digital PCR method
for identifying and quantifying
adulteration of meat species in raw and
processed food. *PLoS ONE*. 20 Mar
2017;12(3):e0173567

[99] Fumière O, Marien A,
Fernández Pierna JA, Baeten V,
Berben G. Development of a real-time
PCR protocol for the species origin
confirmation of isolated animal
particles detected by NIRM. *Food
Additives and Contaminants*. 1 Aug
2010;27(8):1118-1127