

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

185,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



The Monte Carlo Techniques and the Complex Probability Paradigm

Abdo Abou Jaoude

Abstract

The concept of mathematical probability was established in 1933 by Andrey Nikolaevich Kolmogorov by defining a system of five axioms. This system can be enhanced to encompass the imaginary numbers set after the addition of three novel axioms. As a result, any random experiment can be executed in the complex probabilities set \mathcal{C} which is the sum of the real probabilities set \mathcal{R} and the imaginary probabilities set \mathcal{M} . We aim here to incorporate supplementary imaginary dimensions to the random experiment occurring in the “real” laboratory in \mathcal{R} and therefore to compute all the probabilities in the sets \mathcal{R} , \mathcal{M} , and \mathcal{C} . Accordingly, the probability in the whole set $\mathcal{C} = \mathcal{R} + \mathcal{M}$ is constantly equivalent to one independently of the distribution of the input random variable in \mathcal{R} , and subsequently the output of the stochastic experiment in \mathcal{R} can be determined absolutely in \mathcal{C} . This is the consequence of the fact that the probability in \mathcal{C} is computed after the subtraction of the chaotic factor from the degree of our knowledge of the nondeterministic experiment. We will apply this innovative paradigm to the well-known Monte Carlo techniques and to their random algorithms and procedures in a novel way.

Keywords: degree of our knowledge, chaotic factor, complex probability set, probability norm, complex random vector, convergence probability, divergence probability, simulation

1. Introduction

“Thus, joining the rigor of the demonstrations of science to the uncertainty of fate, and reconciling these two seemingly contradictory things, it can, taking its name from both, appropriately arrogate to itself this astonishing title: the geometry of chance.”

Blaise Pascal

“You believe in the God who plays dice, and I in complete law and order.”

Albert Einstein, Letter to Max Born

“Chance is the pseudonym of God when He did not want to sign.”

Anatole France

“There is a certain Eternal Law, to wit, Reason, existing in the mind of God and governing the whole universe.”

Saint Thomas Aquinas

“An equation has no meaning for me unless it expresses a thought of God.”

Srinivasa Ramanujan

Calculating probabilities is the crucial task of classical probability theory. Adding supplementary dimensions to nondeterministic experiments will yield a deterministic expression of the theory of probability. This is the novel and original idea at the foundations of my complex probability paradigm. As a matter of fact, probability theory is a stochastic system of axioms in its essence; that means that the phenomena outputs are due to randomness and chance. Adding new imaginary dimensions to the nondeterministic phenomenon happening in the set \mathcal{R} will lead to a deterministic phenomenon, and thus, a probabilistic experiment will have a certain output in the set \mathcal{C} of complex probabilities. If the chaotic experiment becomes fully predictable, then we will be completely capable to foretell the output of random events that occur in the real world in all probabilistic processes. Accordingly, the task that has been achieved here was to extend the set \mathcal{R} of random real probabilities to the deterministic set $\mathcal{C} = \mathcal{R} + \mathcal{M}$ of complex probabilities and this by incorporating the contributions of the set \mathcal{M} which is the set of complementary imaginary probabilities to the set \mathcal{R} . Consequently, since this extension reveals to be successful, an innovative paradigm of stochastic sciences and prognostic was put forward in which all nondeterministic phenomena in \mathcal{R} was expressed deterministically in \mathcal{C} . I coined this novel model by the term “the complex probability paradigm” that was initiated and established in my 14 earlier research works [1–14].

2. The purpose and the advantages of the current chapter

The advantages and the purpose of the present chapter are to [15–39]:

1. Extend the theory of classical probability to cover the complex numbers set, hence to connect the probability theory to the field of complex analysis and variables. This task was initiated and developed in my earlier 14 works.
2. Apply the novel paradigm and its original probability axioms to Monte Carlo techniques.
3. Prove that all phenomena that are nondeterministic can be transformed to deterministic phenomena in the complex probabilities set which is \mathcal{C} .
4. Compute and quantify both the chaotic factor and the degree of our knowledge of Monte Carlo procedures.
5. Represent and show the graphs of the functions and parameters of the innovative model related to Monte Carlo algorithms.
6. Demonstrate that the classical probability concept is permanently equal to 1 in the set of complex probabilities; thus, no chaos, no randomness, no ignorance, no uncertainty, no unpredictability, no nondeterminism, and no disorder exist in

$$\mathcal{C} \text{ (complex set)} = \mathcal{R} \text{ (real set)} + \mathcal{M} \text{ (imaginary set)}.$$

7. Prepare to apply this inventive paradigm to other topics in prognostics and to the field of stochastic processes. These will be the goals of my future research publications.




Figure 1.
 The diagram of the major aims of the complex probability paradigm.

Regarding some applications of the novel established model and as a subsequent work, it can be applied to any nondeterministic experiments using Monte Carlo algorithms whether in the continuous or in the discrete cases.

Moreover, compared with existing literature, the major contribution of the current chapter is to apply the innovative complex probability paradigm to the techniques and concepts of the probabilistic Monte Carlo simulations and algorithms.

The next figure displays the major aims and purposes of the complex probability paradigm (CPP) (**Figure 1**).

3. The complex probability paradigm

3.1 The original Andrey Nikolaevich Kolmogorov system of axioms

The simplicity of Kolmogorov’s system of axioms may be surprising [1–14]. Let E be a collection of elements $\{E_1, E_2, \dots\}$ called elementary events and let F be a set of subsets of E called random events. The five axioms for a finite set E are:

Axiom 1: F is a field of sets.

Axiom 2: F contains the set E .

Axiom 3: A nonnegative real number $P_{rob}(A)$, called the probability of A , is assigned to each set A in F . We have always $0 \leq P_{rob}(A) \leq 1$.

Axiom 4: $P_{rob}(E)$ equals 1.

Axiom 5: If A and B have no elements in common, the number assigned to their union is

$$P_{rob}(A \cup B) = P_{rob}(A) + P_{rob}(B)$$

hence, we say that A and B are disjoint; otherwise, we have

$$P_{rob}(A \cup B) = P_{rob}(A) + P_{rob}(B) - P_{rob}(A \cap B)$$

And we say also that $P_{rob}(A \cap B) = P_{rob}(A) \times P_{rob}(B/A) = P_{rob}(B) \times P_{rob}(A/B)$ which is the conditional probability. If both A and B are independent then $P_{rob}(A \cap B) = P_{rob}(A) \times P_{rob}(B)$.

Moreover, we can generalize and say that for N disjoint (mutually exclusive) events $A_1, A_2, \dots, A_j, \dots, A_N$ (for $1 \leq j \leq N$), we have the following additivity rule:

$$P_{rob}\left(\bigcup_{j=1}^N A_j\right) = \sum_{j=1}^N P_{rob}(A_j)$$

And we say also that for N independent events $A_1, A_2, \dots, A_j, \dots, A_N$ (for $1 \leq j \leq N$), we have the following product rule

$$P_{rob}\left(\bigcap_{j=1}^N A_j\right) = \prod_{j=1}^N P_{rob}(A_j)$$

3.2 Adding the imaginary part \mathcal{M}

Now, we can add to this system of axioms an imaginary part such that:

Axiom 6: Let $P_m = i \times (1 - P_r)$ be the probability of an associated complementary event in \mathcal{M} (the imaginary part) to the event A in \mathcal{R} (the real part). It follows that $P_r + P_m/i = 1$ where i is the imaginary number with $i = \sqrt{-1}$ or $i^2 = -1$.

Axiom 7: We construct the complex number or vector $Z = P_r + P_m = P_r + i(1 - P_r)$ having a norm $|Z|$ such that

$$|Z|^2 = P_r^2 + (P_m/i)^2.$$

Axiom 8: Let P_c denotes the probability of an event in the complex probability universe \mathcal{C} where $\mathcal{C} = \mathcal{R} + \mathcal{M}$. We say that P_c is the probability of an event A in \mathcal{R} with its associated event in \mathcal{M} such that

$$P_c^2 = (P_r + P_m/i)^2 = |Z|^2 - 2iP_rP_m \text{ and is always equal to 1.}$$

We can see that by taking into consideration the set of imaginary probabilities, we added three new and original axioms, and consequently the system of axioms




Figure 2.
The EKA or the CPP diagram.

defined by Kolmogorov was hence expanded to encompass the set of imaginary numbers.

3.3 A brief interpretation of the novel paradigm

To summarize the novel paradigm, we state that in the real probability universe \mathcal{R} , our degree of our certain knowledge is undesirably imperfect and hence unsatisfactory; thus, we extend our analysis to the set of complex numbers \mathcal{C} which incorporates the contributions of both the set of real probabilities which is \mathcal{R} and the complementary set of imaginary probabilities which is \mathcal{M} . Afterward, this will yield an absolute and perfect degree of our knowledge in the probability universe $\mathcal{C} = \mathcal{R} + \mathcal{M}$ because $P_c = 1$ constantly. As a matter of fact, the work in the universe \mathcal{C} of complex probabilities gives way to a sure forecast of any stochastic experiment, since in \mathcal{C} we remove and subtract from the computed degree of our knowledge the measured chaotic factor. This will generate in the universe \mathcal{C} a probability equal to 1 ($P_c^2 = DOK - Chf = DOK + MChf = 1 = P_c$). Many applications taking into consideration numerous continuous and discrete probability distributions in my 14 previous research papers confirm this hypothesis and innovative paradigm. The extended Kolmogorov axioms (EKA) or the complex probability paradigm (CPP) can be shown and summarized in the next illustration (Figure 2).

4. The Monte Carlo techniques and the complex probability paradigm parameters

4.1 The divergence and convergence probabilities

Let R_E be the exact result of the stochastic phenomenon or of a multidimensional or simple integral that are not always possible to compute by probability theory ordinary procedures or by deterministic numerical means or by calculus [1–14]. And let R_A be the phenomenon and integrals approximate results calculated by the techniques of Monte Carlo:

The relative error in the Monte Carlo methods is $\text{Rel.Error} = \left| \frac{R_E - R_A}{R_E} \right| = \left| 1 - \frac{R_A}{R_E} \right|$.

Additionally, the percent relative error is $= 100\% \times \left| \frac{R_E - R_A}{R_E} \right|$ and is always between 0% and 100%. Therefore, the relative error is always between 0 and 1. Hence

$$0 \leq \left| \frac{R_E - R_A}{R_E} \right| \leq 1 \Leftrightarrow \begin{cases} 0 \leq \left(\frac{R_E - R_A}{R_E} \right) \leq 1 & \text{if } R_A \leq R_E \\ 0 \leq - \left(\frac{R_E - R_A}{R_E} \right) \leq 1 & \text{if } R_A \geq R_E \end{cases} \Leftrightarrow \begin{cases} 0 \leq R_A \leq R_E \\ R_E \leq R_A \leq 2R_E \end{cases}$$

Moreover, we define the real probability in the set \mathcal{R} by

$$P_r = 1 - \left| \frac{R_E - R_A}{R_E} \right| = 1 - \left| 1 - \frac{R_A}{R_E} \right| = \begin{cases} 1 - \left(1 - \frac{R_A}{R_E} \right) & \text{if } 0 \leq R_A \leq R_E \\ 1 + \left(1 - \frac{R_A}{R_E} \right) & \text{if } R_E \leq R_A \leq 2R_E \end{cases}$$

$$= \begin{cases} \frac{R_A}{R_E} & \text{if } 0 \leq R_A \leq R_E \\ 2 - \frac{R_A}{R_E} & \text{if } R_E \leq R_A \leq 2R_E \end{cases}$$

= 1 – the relative error in the Monte Carlo method.

= probability of Monte Carlo method convergence in \mathcal{R} .

And therefore,

$$P_m = i(1 - P_r) = i \left\{ 1 - \left[1 - \left| \frac{R_E - R_A}{R_E} \right| \right] \right\} = i \left\{ 1 - \left[1 - \left| 1 - \frac{R_A}{R_E} \right| \right] \right\} = i \left| 1 - \frac{R_A}{R_E} \right|$$

$$= \begin{cases} i \left(1 - \frac{R_A}{R_E} \right) & \text{if } 0 \leq R_A \leq R_E \\ -i \left(1 - \frac{R_A}{R_E} \right) & \text{if } R_E \leq R_A \leq 2R_E \end{cases} = \begin{cases} i \left(1 - \frac{R_A}{R_E} \right) & \text{if } 0 \leq R_A \leq R_E \\ i \left(\frac{R_A}{R_E} - 1 \right) & \text{if } R_E \leq R_A \leq 2R_E \end{cases}$$

= probability of Monte Carlo method divergence in the imaginary complementary probability set \mathcal{M} since it is the imaginary complement of P_r .

Consequently,

$$P_m/i = 1 - P_r = \left| 1 - \frac{R_A}{R_E} \right| = \begin{cases} 1 - \frac{R_A}{R_E} & \text{if } 0 \leq R_A \leq R_E \\ \frac{R_A}{R_E} - 1 & \text{if } R_E \leq R_A \leq 2R_E \end{cases}$$

= the relative error in the Monte Carlo method.

= probability of Monte Carlo method divergence in \mathcal{R} since it is the real complement of P_r .

In the case where $0 \leq R_A \leq R_E \Rightarrow 0 \leq \frac{R_A}{R_E} \leq 1 \Rightarrow 0 \leq P_r \leq 1$ and we deduce also that $0 \leq \left(1 - \frac{R_A}{R_E} \right) \leq 1 \Rightarrow 0 \leq P_m/i \leq 1$ and $\Rightarrow 0 \leq P_m \leq i$.

And in the case where $R_E \leq R_A \leq 2R_E \Rightarrow 1 \leq \frac{R_A}{R_E} \leq 2 \Rightarrow 0 \leq \left(2 - \frac{R_A}{R_E} \right) \leq 1 \Rightarrow 0 \leq P_r \leq 1$ and we deduce also that $0 \leq \left(\frac{R_A}{R_E} - 1 \right) \leq 1 \Rightarrow 0 \leq P_m/i \leq 1$ and $\Rightarrow 0 \leq P_m \leq i$.

Consequently, if $R_A = 0$ or $R_A = 2R_E$ that means before the beginning of the simulation, then

$$P_r = P_{rob} \text{ (convergence) in } \mathcal{R} = 0$$

$$P_m = P_{rob} \text{ (divergence) in } \mathcal{M} = i$$

$$P_m/i = P_{rob} \text{ (divergence) in } \mathcal{R} = 1$$

And if $R_A = R_E$ that means at the end of Monte Carlo simulation, then

$$P_r = P_{rob} \text{ (convergence) in } \mathcal{R} = 1$$

$$P_m = P_{rob} \text{ (divergence) in } \mathcal{M} = 0$$

$$P_m/i = P_{rob} \text{ (divergence) in } \mathcal{R} = 0$$

4.2 The complex random vector Z in $\mathcal{C} = \mathcal{R} + \mathcal{M}$

$$\text{We have } Z = P_r + P_m = \begin{cases} \frac{R_A}{R_E} + i \left(1 - \frac{R_A}{R_E} \right) & \text{if } 0 \leq R_A \leq R_E \\ \left(2 - \frac{R_A}{R_E} \right) + i \left(\frac{R_A}{R_E} - 1 \right) & \text{if } R_E \leq R_A \leq 2R_E \end{cases} .$$

$$= \text{Re}(Z) + i\text{Im}(Z)$$

where

$$\operatorname{Re}(Z) = P_r = \begin{cases} \frac{R_A}{R_E} & \text{if } 0 \leq R_A \leq R_E \\ 2 - \frac{R_A}{R_E} & \text{if } R_E \leq R_A \leq 2R_E \end{cases} = \text{the real part of } Z$$

and

$$\operatorname{Im}(Z) = P_m/i = \begin{cases} 1 - \frac{R_A}{R_E} & \text{if } 0 \leq R_A \leq R_E \\ \frac{R_A}{R_E} - 1 & \text{if } R_E \leq R_A \leq 2R_E \end{cases} = \text{the imaginary part of } Z.$$

That means that the complex random vector Z is the sum in \mathcal{C} of the real probability of convergence in \mathcal{R} and of the imaginary probability of divergence in \mathcal{M} .

If $R_A = 0$ or $R_A = 2R_E$ (before the simulation begins), then

$$P_r = \frac{R_A}{R_E} = \frac{0}{R_E} = 0 \text{ or } P_r = 2 - \frac{R_A}{R_E} = 2 - \frac{2R_E}{R_E} = 2 - 2 = 0.$$

and

$$P_m = i\left(1 - \frac{R_A}{R_E}\right) = i\left(1 - \frac{0}{R_E}\right) = i(1 - 0) = i \text{ or } P_m = i\left(\frac{R_A}{R_E} - 1\right) = i\left(\frac{2R_E}{R_E} - 1\right) = i(2 - 1) = i$$

therefore $Z = 0 + i = i$.

If $R_A = \frac{R_E}{2}$ or $R_A = \frac{3R_E}{2}$ (at the middle of the simulation), then

$$P_r = \begin{cases} \frac{R_A}{R_E} & \text{if } 0 \leq R_A \leq R_E \\ 2 - \frac{R_A}{R_E} & \text{if } R_E \leq R_A \leq 2R_E \end{cases} = \begin{cases} \frac{R_E}{2R_E} = 0.5 & \text{if } 0 \leq R_A \leq R_E \\ 2 - \frac{3R_E}{2R_E} = 0.5 & \text{if } R_E \leq R_A \leq 2R_E \end{cases} \Leftrightarrow P_r = 0.5$$

and

$$P_m = \begin{cases} i\left(1 - \frac{R_A}{R_E}\right) & \text{if } 0 \leq R_A \leq R_E \\ i\left(\frac{R_A}{R_E} - 1\right) & \text{if } R_E \leq R_A \leq 2R_E \end{cases} = \begin{cases} i\left(1 - \frac{R_E}{2R_E}\right) = 0.5i & \text{if } 0 \leq R_A \leq R_E \\ i\left(\frac{3R_E}{2R_E} - 1\right) = 0.5i & \text{if } R_E \leq R_A \leq 2R_E \end{cases} \Leftrightarrow P_m = 0.5i$$

therefore $Z = 0.5 + 0.5i$.

If $R_A = R_E$ (at the simulation end), then

$$P_r = \begin{cases} \frac{R_A}{R_E} = \frac{R_E}{R_E} = 1 & \text{if } 0 \leq R_A \leq R_E \\ 2 - \frac{R_A}{R_E} = 2 - \frac{R_E}{R_E} = 2 - 1 = 1 & \text{if } R_E \leq R_A \leq 2R_E \end{cases} \Leftrightarrow P_r = 1$$

and

$$\begin{aligned}
 P_m &= \begin{cases} i \left(1 - \frac{R_A}{R_E}\right) & \text{if } 0 \leq R_A \leq R_E \\ i \left(\frac{R_A}{R_E} - 1\right) & \text{if } R_E \leq R_A \leq 2R_E \end{cases} = \begin{cases} i \left(1 - \frac{R_E}{R_E}\right) & \text{if } 0 \leq R_A \leq R_E \\ i \left(\frac{R_E}{R_E} - 1\right) & \text{if } R_E \leq R_A \leq 2R_E \end{cases} \\
 &= \begin{cases} 0 & \text{if } 0 \leq R_A \leq R_E \\ 0 & \text{if } R_E \leq R_A \leq 2R_E \end{cases} \\
 &\Leftrightarrow P_m = 0
 \end{aligned}$$

therefore $Z = 1 + 0i = 1$.

4.3 The degree of our knowledge, DOK

We have

$$\begin{aligned}
 DOK = |Z|^2 &= P_r^2 + (P_m/i)^2 = \begin{cases} \left(\frac{R_A}{R_E}\right)^2 & \text{if } 0 \leq R_A \leq R_E \\ \left(2 - \frac{R_A}{R_E}\right)^2 & \text{if } R_E \leq R_A \leq 2R_E \end{cases} + \begin{cases} \left(1 - \frac{R_A}{R_E}\right)^2 & \text{if } 0 \leq R_A \leq R_E \\ \left(\frac{R_A}{R_E} - 1\right)^2 & \text{if } R_E \leq R_A \leq 2R_E \end{cases} \\
 &= \begin{cases} \left(\frac{R_A}{R_E}\right)^2 + \left(1 - \frac{R_A}{R_E}\right)^2 & \text{if } 0 \leq R_A \leq R_E \\ \left(2 - \frac{R_A}{R_E}\right)^2 + \left(\frac{R_A}{R_E} - 1\right)^2 & \text{if } R_E \leq R_A \leq 2R_E \end{cases} = \begin{cases} 2\left(\frac{R_A}{R_E}\right)^2 - 2\left(\frac{R_A}{R_E}\right) + 1 & \text{if } 0 \leq R_A \leq R_E \\ 2\left(\frac{R_A}{R_E}\right)^2 - 6\left(\frac{R_A}{R_E}\right) + 5 & \text{if } R_E \leq R_A \leq 2R_E \end{cases}
 \end{aligned}$$

From CPP we have that $0.5 \leq DOK \leq 1$, then if $DOK = 0.5$

$$\Leftrightarrow \begin{cases} 2\left(\frac{R_A}{R_E}\right)^2 - 2\left(\frac{R_A}{R_E}\right) + 1 = 0.5 & \text{if } 0 \leq R_A \leq R_E \\ 2\left(\frac{R_A}{R_E}\right)^2 - 6\left(\frac{R_A}{R_E}\right) + 5 = 0.5 & \text{if } R_E \leq R_A \leq 2R_E \end{cases}$$

then solving the second-degree equations for $\frac{R_A}{R_E}$ gives

$$\begin{cases} \frac{R_A}{R_E} = 1/2 & \text{if } 0 \leq R_A \leq R_E \\ \frac{R_A}{R_E} = 3/2 & \text{if } R_E \leq R_A \leq 2R_E \end{cases} \Leftrightarrow \begin{cases} R_A = R_E/2 & \text{if } 0 \leq R_A \leq R_E \\ R_A = 3R_E/2 & \text{if } R_E \leq R_A \leq 2R_E \end{cases} \text{ and vice versa.}$$

That means that DOK is minimum when the approximate result R_A is equal to half of the exact result R_E if $0 \leq R_A \leq R_E$ or when the approximate result is equal to three times the half of the exact result if $R_E \leq R_A \leq 2R_E$, which means at the middle of the simulation.

In addition, if $DOK = 1$, then

$$\Leftrightarrow \begin{cases} 2\left(\frac{R_A}{R_E}\right)^2 - 2\left(\frac{R_A}{R_E}\right) + 1 = 1 & \text{if } 0 \leq R_A \leq R_E \\ 2\left(\frac{R_A}{R_E}\right)^2 - 6\left(\frac{R_A}{R_E}\right) + 5 = 1 & \text{if } R_E \leq R_A \leq 2R_E \end{cases} \Leftrightarrow \begin{cases} \left(\frac{R_A}{R_E}\right)^2 - \left(\frac{R_A}{R_E}\right) = 0 & \text{if } 0 \leq R_A \leq R_E \\ 2\left(\frac{R_A}{R_E}\right)^2 - 6\left(\frac{R_A}{R_E}\right) + 4 = 0 & \text{if } R_E \leq R_A \leq 2R_E \end{cases}$$

$$\Leftrightarrow \begin{cases} R_A = 0 \text{ OR } R_A = R_E & \text{if } 0 \leq R_A \leq R_E \\ R_A = 2R_E \text{ OR } R_A = R_E & \text{if } R_E \leq R_A \leq 2R_E \end{cases} \text{ and vice versa.}$$

That means that DOK is maximum when the approximate result R_A is equal to 0 or $2R_E$ (before the beginning of the simulation) and when it is equal to the exact result R_E (at the end of the simulation). We can deduce that we have perfect and total knowledge of the stochastic experiment before the beginning of Monte Carlo simulation since no randomness was introduced yet, as well as at the end of the simulation after the convergence of the method to the exact result.

4.4 The chaotic factor, Chf

We have

$$\begin{aligned} Chf &= 2iP_rP_m \\ &= 2i \times \begin{cases} \frac{R_A}{R_E} & \text{if } 0 \leq R_A \leq R_E \\ 2 - \frac{R_A}{R_E} & \text{if } R_E \leq R_A \leq 2R_E \end{cases} \times \begin{cases} i\left(1 - \frac{R_A}{R_E}\right) & \text{if } 0 \leq R_A \leq R_E \\ i\left(\frac{R_A}{R_E} - 1\right) & \text{if } R_E \leq R_A \leq 2R_E \end{cases} \end{aligned}$$

Since $i^2 = -1$ then

$$Chf = \begin{cases} -2\left(\frac{R_A}{R_E}\right)\left(1 - \frac{R_A}{R_E}\right) & \text{if } 0 \leq R_A \leq R_E \\ -2\left(2 - \frac{R_A}{R_E}\right)\left(\frac{R_A}{R_E} - 1\right) & \text{if } R_E \leq R_A \leq 2R_E \end{cases}$$

From CPP we have that $-0.5 \leq Chf \leq 0$, and then if $Chf = -0.5$

$$\Leftrightarrow \begin{cases} -2\left(\frac{R_A}{R_E}\right)\left(1 - \frac{R_A}{R_E}\right) = -0.5 & \text{if } 0 \leq R_A \leq R_E \\ -2\left(2 - \frac{R_A}{R_E}\right)\left(\frac{R_A}{R_E} - 1\right) = -0.5 & \text{if } R_E \leq R_A \leq 2R_E \end{cases} \Leftrightarrow \begin{cases} R_A = R_E/2 & \text{if } 0 \leq R_A \leq R_E \\ R_A = 3R_E/2 & \text{if } R_E \leq R_A \leq 2R_E \end{cases}$$

and vice versa.

That means that Chf is minimum when the approximate result R_A is equal to half of the exact result R_E if $0 \leq R_A \leq R_E$ or when the approximate result is equal to three times the half of the exact result if $R_E \leq R_A \leq 2R_E$, which means at the middle of the simulation.

In addition, if $Chf = 0$ then

$$\Leftrightarrow \begin{cases} -2\left(\frac{R_A}{R_E}\right)\left(1 - \frac{R_A}{R_E}\right) = 0 & \text{if } 0 \leq R_A \leq R_E \\ -2\left(2 - \frac{R_A}{R_E}\right)\left(\frac{R_A}{R_E} - 1\right) = 0 & \text{if } R_E \leq R_A \leq 2R_E \end{cases} \Leftrightarrow \begin{cases} R_A = 0 \text{ OR } R_A = R_E & \text{if } 0 \leq R_A \leq R_E \\ R_A = 2R_E \text{ OR } R_A = R_E & \text{if } R_E \leq R_A \leq 2R_E \end{cases}$$

And, conversely, if $\begin{cases} R_A = 0 \text{ OR } R_A = R_E & \text{if } 0 \leq R_A \leq R_E \\ R_A = 2R_E \text{ OR } R_A = R_E & \text{if } R_E \leq R_A \leq 2R_E \end{cases}$, then $Chf = 0$.

That means that Chf is equal to 0 when the approximate result R_A is equal to 0 or $2R_E$ (before the beginning of the simulation) and when it is equal to the exact result R_E (at the end of the simulation).

4.5 The magnitude of the chaotic factor, $MChf$

We have

$$MChf = |Chf| = -2iP_rP_m = -2i \times \begin{cases} \frac{R_A}{R_E} & \text{if } 0 \leq R_A \leq R_E \\ 2 - \frac{R_A}{R_E} & \text{if } R_E \leq R_A \leq 2R_E \end{cases} \times \begin{cases} i\left(1 - \frac{R_A}{R_E}\right) & \text{if } 0 \leq R_A \leq R_E \\ i\left(\frac{R_A}{R_E} - 1\right) & \text{if } R_E \leq R_A \leq 2R_E \end{cases}$$

Since $i^2 = -1$ then

$$MChf = \begin{cases} 2\left(\frac{R_A}{R_E}\right)\left(1 - \frac{R_A}{R_E}\right) & \text{if } 0 \leq R_A \leq R_E \\ 2\left(2 - \frac{R_A}{R_E}\right)\left(\frac{R_A}{R_E} - 1\right) & \text{if } R_E \leq R_A \leq 2R_E \end{cases}$$

From CPP we have that $0 \leq MChf \leq 0.5$, and then if $MChf = 0.5$

$$\Leftrightarrow \begin{cases} 2\left(\frac{R_A}{R_E}\right)\left(1 - \frac{R_A}{R_E}\right) = 0.5 & \text{if } 0 \leq R_A \leq R_E \\ 2\left(2 - \frac{R_A}{R_E}\right)\left(\frac{R_A}{R_E} - 1\right) = 0.5 & \text{if } R_E \leq R_A \leq 2R_E \end{cases} \Leftrightarrow \begin{cases} R_A = R_E/2 & \text{if } 0 \leq R_A \leq R_E \\ R_A = 3R_E/2 & \text{if } R_E \leq R_A \leq 2R_E \end{cases}$$

and vice versa.

That means that $MChf$ is maximum when the approximate result R_A is equal to half of the exact result R_E if $0 \leq R_A \leq R_E$ or when the approximate result is equal to three times the half of the exact result if $R_E \leq R_A \leq 2R_E$, which means at the middle of the simulation. This implies that the magnitude of the chaos ($MChf$) introduced by the random variables used in Monte Carlo method is maximum at the halfway of the simulation.

In addition, if $MChf = 0$, then

$$\Leftrightarrow \begin{cases} 2\left(\frac{R_A}{R_E}\right)\left(1 - \frac{R_A}{R_E}\right) = 0 & \text{if } 0 \leq R_A \leq R_E \\ 2\left(2 - \frac{R_A}{R_E}\right)\left(\frac{R_A}{R_E} - 1\right) = 0 & \text{if } R_E \leq R_A \leq 2R_E \end{cases} \Leftrightarrow \begin{cases} R_A = 0 \text{ OR } R_A = R_E & \text{if } 0 \leq R_A \leq R_E \\ R_A = 2R_E \text{ OR } R_A = R_E & \text{if } R_E \leq R_A \leq 2R_E \end{cases}$$

And, conversely, if $\begin{cases} R_A = 0 \text{ OR } R_A = R_E & \text{if } 0 \leq R_A \leq R_E \\ R_A = 2R_E \text{ OR } R_A = R_E & \text{if } R_E \leq R_A \leq 2R_E \end{cases}$, then $MChf = 0$.

That means that $MChf$ is minimum and is equal to 0 when the approximate result R_A is equal to 0 or $2R_E$ (before the beginning of the simulation) and when it is

equal to the exact result R_E (at the end of the simulation). We can deduce that the magnitude of the chaos in the stochastic experiment is null before the beginning of Monte Carlo simulation since no randomness was introduced yet, as well as at the end of the simulation after the convergence of the method to the exact result when randomness has finished its task in the stochastic Monte Carlo method and experiment.

4.6 The probability P_c in the probability set $\mathcal{C} = \mathcal{R} + \mathcal{M}$

We have

$$\begin{aligned} P_c^2 &= DOK - Chf = DOK + MChf \\ &= \begin{cases} 2\left(\frac{R_A}{R_E}\right)^2 - 2\left(\frac{R_A}{R_E}\right) + 1 & \text{if } 0 \leq R_A \leq R_E \\ 2\left(\frac{R_A}{R_E}\right)^2 - 6\left(\frac{R_A}{R_E}\right) + 5 & \text{if } R_E \leq R_A \leq 2R_E \end{cases} - \begin{cases} -2\left(\frac{R_A}{R_E}\right)\left(1 - \frac{R_A}{R_E}\right) & \text{if } 0 \leq R_A \leq R_E \\ -2\left(2 - \frac{R_A}{R_E}\right)\left(\frac{R_A}{R_E} - 1\right) & \text{if } R_E \leq R_A \leq 2R_E \end{cases} \\ &= \begin{cases} 1 & \text{if } 0 \leq R_A \leq R_E \\ 1 & \text{if } R_E \leq R_A \leq 2R_E \end{cases} \Leftrightarrow P_c^2 = 1 \text{ for } 0 \leq \forall R_A \leq 2R_E \end{aligned}$$

$\Leftrightarrow P_c = 1$ = probability of convergence in \mathcal{C} ; therefore,

$$P_c = \begin{cases} \frac{R_A}{R_E} = 1 & \text{if } 0 \leq R_A \leq R_E \\ 2 - \frac{R_A}{R_E} = 1 & \text{if } R_E \leq R_A \leq 2R_E \end{cases} \Leftrightarrow \begin{cases} R_A = R_E & \text{if } 0 \leq R_A \leq R_E \\ R_A = R_E & \text{if } R_E \leq R_A \leq 2R_E \end{cases}$$

$\Leftrightarrow R_A = R_E$ for $0 \leq \forall R_A \leq 2R_E$ continuously in the probability set $\mathcal{C} = \mathcal{R} + \mathcal{M}$. This is due to the fact that in \mathcal{C} , we have subtracted in the equation above the chaotic factor Chf from our knowledge DOK , and therefore we have eliminated chaos caused and introduced by all the random variables and the stochastic fluctuations that lead to approximate results in the Monte Carlo simulation in \mathcal{R} . Therefore, since in \mathcal{C} we have always $R_A = R_E$, then the Monte Carlo simulation which is a stochastic method by nature in \mathcal{R} becomes after applying the CPP a deterministic method in \mathcal{C} since the probability of convergence of any random experiment in \mathcal{C} is constantly and permanently equal to 1 for any iterations number N .

4.7 The rates of change of the probabilities in \mathcal{R} , \mathcal{M} , and \mathcal{C}

Since

$$Z = P_r + P_m = \begin{cases} \frac{R_A}{R_E} + i\left(1 - \frac{R_A}{R_E}\right) & \text{if } 0 \leq R_A \leq R_E \\ \left(2 - \frac{R_A}{R_E}\right) + i\left(\frac{R_A}{R_E} - 1\right) & \text{if } R_E \leq R_A \leq 2R_E \end{cases} = \text{Re}(Z) + i\text{Im}(Z)$$

Then

$$\begin{aligned}
\frac{dZ}{dR_A} &= \frac{dP_r}{dR_A} + \frac{dP_m}{dR_A} = \begin{cases} \frac{d}{dR_A} \left[\frac{R_A}{R_E} + i \left(1 - \frac{R_A}{R_E} \right) \right] & \text{if } 0 \leq R_A \leq R_E \\ \frac{d}{dR_A} \left[\left(2 - \frac{R_A}{R_E} \right) + i \left(\frac{R_A}{R_E} - 1 \right) \right] & \text{if } R_E \leq R_A \leq 2R_E \end{cases} \\
&= \begin{cases} \frac{d}{dR_A} \left[\frac{R_A}{R_E} \right] + \frac{d}{dR_A} \left[i \left(1 - \frac{R_A}{R_E} \right) \right] & \text{if } 0 \leq R_A \leq R_E \\ \frac{d}{dR_A} \left[2 - \frac{R_A}{R_E} \right] + \frac{d}{dR_A} \left[i \left(\frac{R_A}{R_E} - 1 \right) \right] & \text{if } R_E \leq R_A \leq 2R_E \end{cases} \\
&= \begin{cases} \frac{1}{R_E} - \frac{i}{R_E} = \frac{1}{R_E} (1 - i) & \text{if } 0 \leq R_A \leq R_E \\ -\frac{1}{R_E} + \frac{i}{R_E} = \frac{1}{R_E} (i - 1) & \text{if } R_E \leq R_A \leq 2R_E \end{cases}
\end{aligned}$$

Therefore,

$$\begin{aligned}
\operatorname{Re} \left[\frac{dZ}{dR_A} \right] &= \frac{dP_r}{dR_A} = \begin{cases} +\frac{1}{R_E} & \text{if } 0 \leq R_A \leq R_E \\ -\frac{1}{R_E} & \text{if } R_E \leq R_A \leq 2R_E \end{cases} \\
&= \begin{cases} \text{constant} > 0 & \text{if } 0 \leq R_A \leq R_E \text{ and } R_E > 0 \\ \text{constant} < 0 & \text{if } R_E \leq R_A \leq 2R_E \text{ and } R_E > 0 \end{cases}
\end{aligned}$$

That means that the slope of the probability of convergence in \mathcal{R} or its rate of change is constant and positive if $0 \leq R_A \leq R_E$, and constant and negative if $R_E \leq R_A \leq 2R_E$, and it depends only on R_E ; hence, we have a constant increase in P_r (the convergence probability) as a function of the iterations number N as R_A increases from 0 to R_E and as R_A decreases from $2R_E$ to R_E till P_r reaches the value 1 that means till the random experiment converges to R_E :

$$\begin{aligned}
\operatorname{Im} \left[\frac{dZ}{dR_A} \right] &= \frac{1}{i} \frac{dP_m}{dR_A} = \frac{d(P_m/i)}{dR_A} = \begin{cases} -\frac{1}{R_E} & \text{if } 0 \leq R_A \leq R_E \\ +\frac{1}{R_E} & \text{if } R_E \leq R_A \leq 2R_E \end{cases} \\
&= \begin{cases} \text{constant} < 0 & \text{if } 0 \leq R_A \leq R_E \text{ and } R_E > 0 \\ \text{constant} > 0 & \text{if } R_E \leq R_A \leq 2R_E \text{ and } R_E > 0 \end{cases}
\end{aligned}$$

That means that the slopes of the probabilities of divergence in \mathcal{R} and \mathcal{M} or their rates of change are constant and negative if $0 \leq R_A \leq R_E$ and constant and positive if $R_E \leq R_A \leq 2R_E$ and they depend only on R_E ; hence, we have a constant decrease in P_m/i and P_m (the divergence probabilities) as functions of the iterations number N as R_A increases from 0 to R_E and as R_A decreases from $2R_E$ to R_E till P_m/i and P_m reach the value 0 that means till the random experiment converges to R_E .

Additionally,

$$\begin{aligned} \left| \frac{dZ}{dR_A} \right|^2 &= \left[\frac{dP_r}{dR_A} \right]^2 + \left[\frac{1}{i} \frac{dP_m}{dR_A} \right]^2 = \left[\frac{dP_r}{dR_A} \right]^2 + \left[\frac{d(P_m/i)}{dR_A} \right]^2 \\ &= \begin{cases} \left(\frac{1}{R_E} \right)^2 + \left(-\frac{1}{R_E} \right)^2 & \text{if } 0 \leq R_A \leq R_E \\ \left(-\frac{1}{R_E} \right)^2 + \left(\frac{1}{R_E} \right)^2 & \text{if } R_E \leq R_A \leq 2R_E \end{cases} \\ \Leftrightarrow \left| \frac{dZ}{dR_A} \right|^2 &= \frac{1}{R_E^2} + \frac{1}{R_E^2} = \frac{2}{R_E^2} \quad \text{for } 0 \leq \forall R_A \leq 2R_E \\ \Leftrightarrow \left| \frac{dZ}{dR_A} \right| &= \frac{\sqrt{2}}{R_E} = \text{constant} > 0 \quad \text{if } R_E > 0; \end{aligned}$$

that means that the module of the slope of the complex probability vector Z in \mathcal{C} or of its rate of change is constant and positive and it depends only on R_E ; hence, we have a constant increase in $\text{Re}(Z)$ and a constant decrease in $\text{Im}(Z)$ as functions of the iterations number N and as Z goes from $(0, i)$ at $N = 0$ till $(1, 0)$ at the simulation end; hence, till $\text{Re}(Z) = P_r$ reaches the value 1 that means till the random experiment converges to R_E .

Furthermore, since $Pc^2 = DOK - Chf = DOK + MChf = 1$, then $Pc = 1 = \text{probability of convergence in } \mathcal{C}$, and consequently

$$\frac{d(Pc)}{dR_A} = \frac{d(1)}{dR_A} = 0,$$

which means that Pc is constantly equal to 1 for every value of R_A , of R_E , and of the iterations number N , which means for any stochastic experiment and for any simulation of Monte Carlo method. So, we conclude that in $\mathcal{C} = \mathcal{R} + \mathcal{M}$, we have complete and perfect knowledge of the random experiment which has become now a deterministic one since the extension in the complex probability plane \mathcal{C} defined by the CPP axioms has changed all stochastic variables to deterministic variables.

5. The new paradigm parameter evaluation

We can infer from what has been developed earlier the following:

The real probability of convergence $P_r(N) = 1 - \left| \frac{R_E - R_A(N)}{R_E} \right|$.

We have $0 \leq N \leq N_C$ where $N = 0$ corresponds to the instant before the beginning of the random experiment when $R_A(N = 0) = 0$ or $= 2R_E$ and where $N = N_C$ (iterations number needed for the method convergence) corresponds to the instant at the end of the random experiments and Monte Carlo methods when $R_A(N = N_C) \rightarrow R_E$.

The imaginary complementary probability of divergence $P_m(N) = i \left| \frac{R_E - R_A(N)}{R_E} \right|$.

The real complementary probability of divergence $P_m(N)/i = \left| \frac{R_E - R_A(N)}{R_E} \right|$.

The random vector of complex probability

$$Z(N) = P_r(N) + P_m(N) = \left[1 - \left| \frac{R_E - R_A(N)}{R_E} \right| \right] + i \left| \frac{R_E - R_A(N)}{R_E} \right|$$

The degree of our knowledge

$$\begin{aligned} DOK(N) &= |Z(N)|^2 = P_r^2(N) + [P_m(N)/i]^2 = \left[1 - \left|\frac{R_E - R_A(N)}{R_E}\right|\right]^2 + \left[\left|\frac{R_E - R_A(N)}{R_E}\right|\right]^2 \\ &= 1 + 2iP_r(N)P_m(N) = 1 - 2P_r(N)[1 - P_r(N)] = 1 - 2P_r(N) + 2P_r^2(N) \\ &= 1 - 2\left|\frac{R_E - R_A(N)}{R_E}\right| + 2\left[\frac{R_E - R_A(N)}{R_E}\right]^2. \end{aligned}$$

$DOK(N)$ is equal to 1 when $P_r(N) = P_r(0) = 0$ and when $P_r(N) = P_r(N_C) = 1$.
The Chaotic factor

$$\begin{aligned} Chf(N) &= 2iP_r(N)P_m(N) = -2P_r(N)[1 - P_r(N)] = -2P_r(N) + 2P_r^2(N) \\ &= -2\left|\frac{R_E - R_A(N)}{R_E}\right| + 2\left[\frac{R_E - R_A(N)}{R_E}\right]^2 \end{aligned}$$

$Chf(N)$ is null when $P_r(N) = P_r(0) = 0$ and when $P_r(N) = P_r(N_C) = 1$.
The magnitude of the chaotic factor $MChf$

$$\begin{aligned} MChf(N) &= |Chf(N)| = -2iP_r(N)P_m(N) = 2P_r(N)[1 - P_r(N)] = 2P_r(N) - 2P_r^2(N) \\ &= 2\left|\frac{R_E - R_A(N)}{R_E}\right| - 2\left[\frac{R_E - R_A(N)}{R_E}\right]^2 \end{aligned}$$

$MChf(N)$ is null when $P_r(N) = P_r(0) = 0$ and when $P_r(N) = P_r(N_C) = 1$.

At any iteration number N $0 \leq \forall N \leq N_C$, the probability calculated in the set \mathcal{C} of complex probabilities is as follows:

$$\begin{aligned} Pc^2(N) &= [P_r(N) + P_m(N)/i]^2 = |Z(N)|^2 - 2iP_r(N)P_m(N) = DOK(N) - Chf(N) \\ &= DOK(N) + MChf(N) = 1 \end{aligned}$$

then


$$Pc^2(N) = [P_r(N) + P_m(N)/i]^2 = \{P_r(N) + [1 - P_r(N)]\}^2 = 1^2 = 1 \Leftrightarrow Pc(N) = 1(\text{continuously}).$$

Thus, the prediction in the set \mathcal{C} of the probabilities of convergence of the random Monte Carlo methods is always certain.

Let us consider afterward a multidimensional integral and a stochastic experiment to simulate the Monte Carlo procedures and to quantify, to draw, as well as to visualize all the prognostic and *CPP* parameters.

6. The flowchart of the prognostic model of Monte Carlo techniques and *CPP*

The flowchart that follows illustrates all the procedures of the elaborated prognostic model of *CPP*.



7. Simulation of the new paradigm

Note that all the numerical values found in the simulations of the new paradigm for any iteration cycles N were computed using the 64-bit MATLAB version 2020 software and compared to the values found by Microsoft Visual C++ programs. Additionally, the reader should be careful of the truncation and rounding errors since we represent all numerical values by at most five significant digits and since we are using Monte Carlo techniques of simulation and integration which yield approximate results under the influence of stochastic aspects and variations. We have considered for this purpose a high-capacity computer system: a workstation computer with parallel microprocessors, a 64-bit operating system, and a 64-GB RAM.

7.1 The continuous random case: a four-dimensional multiple integral

The Monte Carlo technique of integration can be summarized by the following equation:

$$\int_{a_1}^{b_1} \int_{a_2}^{b_2} \dots \int_{a_n}^{b_n} f(x_1, x_2, \dots, x_n).dx_1 dx_2 \dots dx_n \cong \frac{[(b_1 - a_1) \times (b_2 - a_2) \times \dots \times (b_n - a_n)]}{N} \sum_{j=1}^N f(x_{1j}, x_{2j}, \dots, x_{nj})$$

Let us consider here the multidimensional integral of the following function:

$$\begin{aligned} \int_0^{4/3} \int_0^{4/3} \int_0^{4/3} \int_0^{4/3} xyzw .dx dy dz dw &= \int_0^{4/3} \int_0^{4/3} \int_0^{4/3} \left[\frac{x^2}{2} \right]_0^{4/3} yzw .dy dz dw = \int_0^{4/3} \int_0^{4/3} \int_0^{4/3} \frac{16}{18} yzw .dy dz dw \\ &= \frac{8}{9} \int_0^{4/3} \int_0^{4/3} \left[\frac{y^2}{2} \right]_0^{4/3} zw .dz dw = \frac{8}{9} \int_0^{4/3} \int_0^{4/3} \frac{16}{18} zw .dz dw = \frac{64}{81} \int_0^{4/3} \left[\frac{z^2}{2} \right]_0^{4/3} w .dw = \frac{64}{81} \int_0^{4/3} \frac{16}{18} w .dw \\ &= \frac{512}{729} \left[\frac{w^2}{2} \right]_0^{4/3} = \frac{512}{729} \times \frac{16}{18} = \frac{512}{729} \times \frac{8}{9} = \frac{4,096}{6,561} = 0.62429507696997411 \dots \end{aligned}$$

$\Leftrightarrow R_E = 0.62429507696997411 \dots$ by the deterministic methods of calculus.

$\Leftrightarrow f(x, y, z, w) = xyzw$, where x, y, z , and w follow a discrete uniform distribution U such that

$$x \mapsto U(0, 4/3), y \mapsto U(0, 4/3), z \mapsto U(0, 4/3), w \mapsto U(0, 4/3)$$




Figure 3.
The increasing convergence of the Monte Carlo method up to N = 50 iterations.

$$\Leftrightarrow \int_0^{4/3} \int_0^{4/3} \int_0^{4/3} \int_0^{4/3} xyzw .dxdydzdw \cong \frac{[(4/3 - 0) \times (4/3 - 0) \times (4/3 - 0) \times (4/3 - 0)]}{N} \sum_{j=1}^N x_j y_j z_j w_j$$
$$= \frac{256/81}{N} \sum_{j=1}^N x_j y_j z_j w_j = R_A$$

with $1 \leq N \leq N_C$ after applying Monte Carlo method.

Furthermore, the four figures (**Figures 3–6**) illustrate and prove the increasing convergence of Monte Carlo simulation and technique to the exact result




Figure 4.
The increasing convergence of the Monte Carlo method up to N = 100 iterations.




Figure 5.
The increasing convergence of the Monte Carlo method up to N = 500 iterations.




Figure 6.
The increasing convergence of the Monte Carlo method up to $N = 100,000$ iterations.




Figure 7.
The CPP parameters and the Monte Carlo method for a multiple integral.

$R_E = 0.62429507696997411 \dots$ for $N = 50, 100, 500$, and $N = N_C = 100,000$ iterations. Consequently, we have $\lim_{N \rightarrow +\infty} P_r(N) = \lim_{N \rightarrow +\infty} \left\{ 1 - \left| \frac{R_E - R_A(N)}{R_E} \right| \right\} = 1 - \left| \frac{R_E - R_E}{R_E} \right| = 1 - 0 = 1$ which is equal to the probability of convergence of Monte Carlo technique as $N \rightarrow +\infty$.

Moreover, **Figure 7** shows undoubtedly and graphically the relation of all the parameters of the complex probability paradigm ($Chf, R_A, P_r, MChf, R_E, DOK, P_m/i, Pc$) to the Monte Carlo technique after applying CPP to this four-dimensional integral.

7.2 The discrete random case: the matching birthday problem

An interesting problem that can be solved using simulation is *the famous birthday problem*. Suppose that in a room of n persons, each of the 365 days of the year (not a leap year) is equally likely to be someone's birthday. It can be proved from the theory of probability and contrary to intuition that only 23 persons need to be present for the probability to be better than fifty-fifty that at least two of them will have the same birthday.

Many people are interested in checking the theoretical proof of this statement, so we will demonstrate it briefly before doing the problem simulation. After someone is asked about his or her birthday, the probability that the next person asked will not have the same birthday is $364/365$. The probability that the third person's birthday will not match those of the first two people are $363/365$. It is well-known that the probability of two independent and successive events happening is the product of the probability of the separate events. In general, the probability that the n th person asked will have a birthday different from that of anyone already asked is

$$P(\text{all } n \text{ birthdays are different}) = \left(\frac{365}{365}\right) \times \left(\frac{364}{365}\right) \times \left(\frac{363}{365}\right) \times \dots \\ \times \left(\frac{365 - (n - 1)}{365}\right)$$

The probability that the n th person asked will provide a match is 1 minus this value:

$$P(\text{matching birthdays}) = \\ 1 - \left(\frac{365}{365}\right) \times \left(\frac{364}{365}\right) \times \left(\frac{363}{365}\right) \times \dots \times \left(\frac{365 - (n - 1)}{365}\right) \\ = 1 - \frac{(365) \times (364) \times (363) \times \dots \times [365 - (n - 1)]}{365^n} = R_E$$

which shows that with 23 persons, the chances are 50.7%; with 55 persons, the chances are 98.6% or almost theoretically certain that at least two out of 55 people will have the same birthday. The table gives the theoretical probabilities of matching birthdays for a selected number of people n (**Table 1**).

Without using the probability theory, we can write a routine that uses the random number generator to compute the approximate chances for groups of n persons. Obviously, what is needed here is to choose n random integers from the set of integers $\{1, 2, 3, \dots, 365\}$ and to check whether there is a match. When we repeat this experiment a large number of times, we can calculate afterward the probability of at least one match in any gathering of n persons. Note that if $n \geq 366$, then $P(\text{matching birthdays}) = 1$ by the famous pigeonhole principle.

Furthermore, the four figures (**Figures 8–11**) illustrate and prove the increasing convergence of Monte Carlo simulation and technique to the exact result $R_E = 0.706316242719 \dots$ for $n = 30$ and for $N = 50, 100, 500$, and $N = N_C = 750, 000$ iterations. Consequently, we have $\lim_{N \rightarrow +\infty} P_r(N) = \lim_{N \rightarrow +\infty} \left\{ 1 - \left| \frac{R_E - R_A(N)}{R_E} \right| \right\} = 1 - \left| \frac{R_E - R_E}{R_E} \right| = 1 - 0 = 1$ which is equal to the probability of convergence of Monte Carlo technique as $N \rightarrow +\infty$.

Moreover, **Figure 12** shows undoubtedly and graphically the relation of all the parameters of the complex probability paradigm ($Chf, R_A, P_r, MChf, R_E, DOK, P_m/i, Pc$) to the Monte Carlo technique after applying CPP to this problem of matching birthday.

Number of people n	Theoretical probability = R_E
$n = 5$	$P = 0.027135573700$
$n = 10$	$P = 0.116948177711$
$n = 15$	$P = 0.252901319764$
$n = 20$	$P = 0.411438383581$
$n = 22$	$P = 0.475695307663$
$n = 23$	$P = 0.507297234324$
$n = 25$	$P = 0.568699703969$
$n = 30$	$P = 0.706316242719$
$n = 35$	$P = 0.814383238875$
$n = 40$	$P = 0.891231809818$
$n = 45$	$P = 0.940975899466$
$n = 50$	$P = 0.970373579578$
$n = 55$	$P = 0.986262288816$
$n = 100$	$P = 0.999999692751$
$n = 133$	$P = 0.999999999999$
$n = 365$	$P = 1.000000000000$

Table 1.
Some theoretical probabilities of matching birthdays for n people where $1 \leq n \leq 365$.




Figure 8.
The increasing convergence of the Monte Carlo method up to $N = 50$ iterations.

7.2.1 The cubes of complex probability

In **Figure 13** and in the first cube, the simulation of Chf and DOK as functions of the iterations N and of each other is executed for the problem of matching birthday. If we project $Pc^2(N) = DOK(N) - Chf(N) = 1 = Pc(N)$ on the plane $N = 0$ iterations,




Figure 9.
The increasing convergence of the Monte Carlo method up to $N = 100$ iterations.




Figure 10.
The increasing convergence of the Monte Carlo method up to $N = 500$ iterations.

we will get the line in cyan. The starting point of this line is point J ($DOK = 1$, $Chf = 0$) when $N = 0$ iterations, and then the line gets to point ($DOK = 0.5$, $Chf = -0.5$) when $N = 375,000$ iterations and joins finally and again point J ($DOK = 1$, $Chf = 0$) when $N = N_C = 750,000$ iterations. The graphs of $Chf(N)$ (pink, green, blue) in different planes and $DOK(N)$ (red) represent the other curves. We can notice that point K ($DOK = 0.5$, $Chf = -0.5$, $N = 375,000$ iterations) is the minimum of all these curves. We can notice also that point L has the coordinates ($DOK = 1$, $Chf = 0$, $N = N_C = 750,000$ iterations). Additionally, the three points J, K, and L correspond to the same points that exist in **Figure 12**.




Figure 11.
The increasing convergence of the Monte Carlo method up to $N = 750,000$ iterations.




Figure 12.
The CPP parameters and the Monte Carlo techniques for the matching birthday problem.

In **Figure 14** and in the second cube, we simulate the probability of convergence $P_r(N)$ and its complementary real probability of divergence $P_m(N)/i$ as functions of the iterations N for the problem of matching birthday. If we project $Pc^2(N) = P_r(N) + P_m(N)/i = 1 = Pc(N)$ on the plane $N = 0$ iterations, we will get the line in cyan. The starting point of this line is point $(P_r = 0, P_m/i = 1)$, and the final point is

The Matching Birthday Problem: DOK and Chf in terms of N and of each other




Figure 13.
Chf and DOK in terms of each other and of N for the problem of matching birthday.

The Matching Birthday Problem: The Probabilities P_r and P_m / i in terms of N




Figure 14.
 P_m / i and P_r in terms of each other and of N for the problem of matching birthday.




Figure 15.

The vector of complex probability Z in terms of N for the problem of matching birthday.

point $(P_r = 1, P_m/i = 0)$. The graph of $P_r(N)$ in the plane $P_r(N) = P_m(N)/i$ is represented by the red curve. The starting point of this graph is point J ($P_r = 0, P_m/i = 1, N = 0$ iterations), and then it gets to point K ($P_r = 0.5, P_m/i = 0.5, N = 375,000$ iterations) and joins finally point L ($P_r = 1, P_m/i = 0, N = N_C = 750,000$ iterations). The graph of $P_m(N)/i$ in the plane $P_r(N) + P_m(N)/i = 1$ is represented by the blue curve. We can notice how much point K is important and which is the intersection of the blue and red graphs when $P_r(N) = P_m(N)/i = 0.5$ at $N = 375,000$ iterations. Additionally, the three points J, K, and L correspond to the same points that exist in **Figure 12**.

In **Figure 15** and in the third cube, we simulate the vector of complex probabilities $Z(N)$ in \mathcal{C} as a function of the real probability of convergence $P_r(N) = \text{Re}(Z)$ in \mathcal{R} and of its complementary imaginary probability of divergence $P_m(N) = i \times \text{Im}(Z)$ in \mathcal{M} , and as a function of the iterations N for the problem of matching birthday. The graph of $P_r(N)$ in the plane $P_m(N) = 0$ is represented by the red curve, and the graph of $P_m(N)$ in the plane $P_r(N) = 0$ is represented by the blue curve. The graph of the vector of complex probabilities $Z(N) = P_r(N) + P_m(N) = \text{Re}(Z) + i \times \text{Im}(Z)$ in the plane $P_r(N) = iP_m(N) + 1$ is represented by the green curve. The graph of $Z(N)$ has point J ($P_r = 0, P_m = i, N = 0$ iterations) as the starting point and point L ($P_r = 1, P_m = 0, N = N_C = 750,000$ iterations) as the end point. If we project $Z(N)$ curve on the plane of complex probabilities whose equation is $N = 0$ iterations, we get the line in cyan which is $P_r(0) = iP_m(0) + 1$. This projected line has point J ($P_r = 0, P_m = i, N = 0$ iterations) as the starting point and point ($P_r = 1, P_m = 0, N = 0$ iterations) as the end point. We can notice how much point K is important, and it corresponds to $P_r = 0.5$ and

$P_m = 0.5i$ when $N = 375,000$ iterations. Additionally, the three points J, K, and L correspond to the same points that exist in **Figure 12**.

8. Perspectives and conclusion

In the current chapter, the extended and original Kolmogorov model of eight axioms (*EKA*) was connected and applied to the random and classical Monte Carlo techniques. Thus, we have bonded Monte Carlo algorithms to the novel *CPP* paradigm. Accordingly, the paradigm of “complex probability” was more expanded beyond the scope of my 14 earlier studies on this topic.

Also, as it was proved and demonstrated in the original paradigm, when $N = 0$ (before the beginning of the random simulation) and when $N = N_C$ (after the convergence of Monte Carlo algorithm to the exact result), then the chaotic factor (*Chf* and *MChf*) is 0, and the degree of our knowledge (*DOK*) is 1 since the stochastic aspects and variations have either not commenced yet or they have terminated their job on the random phenomenon. During the course of the nondeterministic phenomenon ($N > 0$), we have $0 < MChf \leq 0.5$, $0.5 \leq DOK < 1$, and $-0.5 \leq Chf < 0$, and it can be noticed that throughout this entire process, we have continually and incessantly $Pc^2 = DOK - Chf = DOK + MChf = 1 = Pc$, which means that the simulation which seemed to be random and nondeterministic in the set \mathcal{R} is now deterministic and certain in the set $\mathcal{C} = \mathcal{R} + \mathcal{M}$, and this after adding the contributions of \mathcal{M} to the experiment happening in \mathcal{R} and thus after removing and subtracting the chaotic factor from the degree of our knowledge. Additionally, the probabilities of convergence and divergence of the random Monte Carlo procedure that correspond to each iteration cycle N have been determined in the three sets of probabilities which are \mathcal{C} , \mathcal{M} , and \mathcal{R} by P_c , P_m , and P_r , respectively. Subsequently, at each instance of N , the novel Monte Carlo techniques and *CPP* parameters *DOK*, *Chf*, *MChf*, R_E , R_A , P_r , P_m , P_m/i , P_c , and Z are perfectly and surely predicted in the set of complex probabilities \mathcal{C} with P_c kept as equal to 1 continuously and forever. Also, referring to all these shown simulations and obtained graphs all over the entire chapter, we can visualize and quantify both the system chaos and stochastic influences and aspects (expressed by *Chf* and *MChf*) and the certain knowledge (expressed by *DOK* and P_c) of Monte Carlo algorithms. This is definitely very wonderful, fruitful, and fascinating and demonstrates once again the advantages of extending the five axioms of probability of Kolmogorov and thus the benefits and novelty of this original theory in applied mathematics and prognostics that can be called verily: “the complex probability paradigm.”

Moreover, it is important to mention here that one essential and very well-known probability distribution was taken into consideration in the current chapter which is the uniform and discrete probability distribution as well as a specific generator of uniform random numbers, knowing that the original *CPP* model can be applied to any generator of uniform random numbers that exists in literature. This will yield certainly analogous results and conclusions and will confirm without any doubt the success of my innovative theory.

As a prospective and future challenges and research, we intend to more develop the novel conceived prognostic paradigm and to apply it to a diverse set of nondeterministic events like for other stochastic phenomena as in the classical theory of probability and in stochastic processes. Additionally, we will implement *CPP* to the first-order reliability method (FORM) in the field of prognostic in engineering and also to the problems of random walk which have huge consequences when applied to economics, to chemistry, to physics, and to pure and applied mathematics.

Nomenclature

\mathcal{R}	the events real set
\mathcal{M}	the events imaginary set
\mathcal{C}	the events complex set
i	the imaginary number with $i^2 = -1$ or $i = \sqrt{-1}$
EKA	extended Kolmogorov axioms
CPP	complex probability paradigm
P_{rob}	any event probability
P_r	the probability in the real set \mathcal{R} = the probability of convergence in \mathcal{R}
P_m	the probability in the complementary imaginary set \mathcal{M} that corresponds to the real probability set in \mathcal{R} = the probability of divergence in \mathcal{M}
P_c	the probability in \mathcal{R} of the event with its associated event in \mathcal{M} = the probability in the set $\mathcal{C} = \mathcal{R} + \mathcal{M}$ of complex probabilities
R_E	the exact result of the random experiment
R_A	the approximate result of the random experiment
Z	complex probability number = complex random vector = sum of P_r and P_m
$DOK = Z ^2$	the degree of our knowledge of the stochastic experiment or system, it is the square of the norm of Z
Chf	the chaotic factor of Z
$MChf$	the magnitude of the chaotic factor of Z
N	the number of iterations cycles = number of random vectors
N_C	the number of iterations cycles till the convergence of Monte Carlo method to R_E = the number of random vectors till convergence.

Author details

Abdo Abou Jaoude
Department of Mathematics and Statistics, Faculty of Natural and Applied Sciences,
Notre Dame University - Louaize, Lebanon

*Address all correspondence to: abdoaj@idm.net.lb

IntechOpen

© 2020 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

References

- [1] Abou Jaoude A, El-Tawil K, Kadry S. Prediction in complex dimension using Kolmogorov's set of axioms. *Journal of Mathematics and Statistics, Science Publications*. 2010;**6**(2):116-124
- [2] Abou Jaoude A. The complex statistics paradigm and the law of large numbers. *Journal of Mathematics and Statistics, Science Publications*. 2013;**9**(4):289-304
- [3] Abou Jaoude A. The theory of complex probability and the first order reliability method. *Journal of Mathematics and Statistics, Science Publications*. 2013;**9**(4):310-324
- [4] Abou Jaoude A. Complex probability theory and prognostic. *Journal of Mathematics and Statistics, Science Publications*. 2014;**10**(1):1-24
- [5] Abou Jaoude A. The complex probability paradigm and analytic linear prognostic for vehicle suspension systems. *American Journal of Engineering and Applied Sciences, Science Publications*. 2015;**8**(1):147-175
- [6] Abou Jaoude A. The paradigm of complex probability and the Brownian motion. *Systems Science and Control Engineering, Taylor and Francis Publishers*. 2015;**3**(1):478-503
- [7] Abou Jaoude A. The paradigm of complex probability and Chebyshev's inequality. *Systems Science and Control Engineering, Taylor and Francis Publishers*. 2016;**4**(1):99-137
- [8] Abou Jaoude A. The paradigm of complex probability and analytic nonlinear prognostic for vehicle suspension systems. *Systems Science and Control Engineering, Taylor and Francis Publishers*. 2016;**4**(1):99-137
- [9] Abou Jaoude A. The paradigm of complex probability and analytic linear prognostic for unburied petrochemical pipelines. *Systems Science and Control Engineering, Taylor and Francis Publishers*. 2017;**5**(1):178-214
- [10] Abou Jaoude A. The paradigm of complex probability and Claude Shannon's information theory. *Systems Science and Control Engineering, Taylor and Francis Publishers*. 2017;**5**(1):380-425
- [11] Abou Jaoude A. The paradigm of complex probability and analytic nonlinear prognostic for unburied petrochemical pipelines. *Systems Science and Control Engineering, Taylor and Francis Publishers*. 2017;**5**(1):495-534
- [12] Abou Jaoude A. The paradigm of complex probability and Ludwig Boltzmann's entropy. *Systems Science and Control Engineering, Taylor and Francis Publishers*. 2018;**6**(1):108-149
- [13] Abou Jaoude A. The paradigm of complex probability and Monte Carlo methods. *Systems Science and Control Engineering, Taylor and Francis Publishers*. 2019;**7**(1):407-451
- [14] Abou Jaoude A. Analytic prognostic in the linear damage case applied to buried petrochemical pipelines and the complex probability paradigm. In: *Fault Detection, Diagnosis and Prognosis*. London, UK: IntechOpen; 2020. DOI: 10.5772/intechopen.90157
- [15] Abou Jaoude A. *The Computer Simulation of Monté Carlo Methods and Random Phenomena*. United Kingdom: Cambridge Scholars Publishing; 2019
- [16] Abou Jaoude A. *The Analysis of Selected Algorithms for the Stochastic Paradigm*. United Kingdom: Cambridge Scholars Publishing; 2019
- [17] Abou Jaoude A. *Applied mathematics: Numerical methods and*

- algorithms for applied mathematicians [PhD thesis]. Spain: Bircham International University. 2004. Available from: <http://www.bircham.edu>
- [18] Abou Jaoude A. Computer science: Computer simulation of Monté Carlo methods and random phenomena [PhD thesis]. Spain: Bircham International University. 2005. Available from: <http://www.bircham.edu>
- [19] Abou Jaoude A. Applied statistics and probability: Analysis and algorithms for the statistical and stochastic paradigm [PhD thesis]. Spain: Bircham International University. 2007. Available from: <http://www.bircham.edu>
- [20] Metropolis N. The Beginning of the Monte Carlo Method. Los Alamos Science (1987 Special Issue dedicated to Stanislaw Ulam); 1987. pp. 125-130
- [21] Eckhardt R. Stan Ulam, John von Neumann, and the Monte Carlo Method. Los Alamos Science, Special Issue (15); 1987. pp. 131-137
- [22] Mazhdrakov M, Benov D, Valkanov N. The Monte Carlo Method. Engineering Applications. Cambridge: ACMO Academic Press; 2018. p. 250. ISBN: 978-619-90684-3-4
- [23] Peragine M. The Universal Mind: The Evolution of Machine Intelligence and Human Psychology. San Diego, CA: Xiphias Press; 2013. [Retrieved: 17 December 2018]
- [24] McKean HP. Propagation of chaos for a class of non-linear parabolic equations. In: Lecture Series in Differential Equations, Session 7. Arlington, VA: Catholic University; 1967. pp. 41-57. Bibcode: 1966PNAS...56.1907M. DOI: 10.1073/pnas.56.6.1907. PMC: 220210. PMID: 16591437
- [25] Herman K, Theodore HE. Estimation of particle transmission by random sampling. National Bureau of Standards: Applied Mathematics Series. 1951;12:27-30
- [26] Turing AM. Computing machinery and intelligence. Mind. LIX. 1950;238: 433-460. DOI: 10.1093/mind/LIX.236.433
- [27] Barricelli NA. Symbiogenetic evolution processes realized by artificial methods. Methodos. 1957:143-182
- [28] Del Moral P. Feynman-Kac Formulae. Genealogical and Interacting Particle Approximations. Series: Probability and Applications. Berlin: Springer; 2004. p. 575
- [29] Assaraf R, Caffarel M, Khelif A. Diffusion Monte Carlo methods with a fixed number of walkers. Physical Review E. 2000;61(4):4566-4575. Bibcode: 2000PhRvE...61.4566A. DOI: 10.1103/physreve.61.4566. Archived from the original (PDF) on 2014-11-07
- [30] Caffarel M, Ceperley D, Kalos M. Comment on Feynman-Kac path-integral calculation of the ground-state energies of atoms. Physical Review Letters. 1993;71(13):2159. Bibcode: 1993PhRvL...71.2159C. DOI: 10.1103/physrevlett.71.2159. PMID: 10054598
- [31] Hetherington JH. Observations on the statistical iteration of matrices. Physical Review A. 1984;30(2713): 2713-2719. Bibcode: 1984PhRvA...30.2713H. DOI: 10.1103/PhysRevA.30.2713
- [32] Fermi E, Richtmyer RD. Note on Census-Taking in Monte Carlo Calculations (PDF). LAM. 805 (A). Declassified Report Los Alamos Archive; 1948
- [33] Rosenbluth MN, Rosenbluth AW. Monte-Carlo calculations of the average extension of macromolecular chains. The Journal of Chemical Physics. 1955; 23(2):356-359. Bibcode: 1955JChPh...23...356R. DOI: 10.1063/1.1741967

[34] Gordon NJ, Salmond DJ, Smith AFM. Novel approach to nonlinear/non-Gaussian Bayesian state estimation. *Radar and Signal Processing, IEE Proceedings F*. April 1993;**140**(2): 107-113. DOI: 10.1049/ip-f-2.1993.0015. ISSN 0956-375X

[35] Kitagawa G. Monte Carlo filter and smoother for non-Gaussian nonlinear state space models. *Journal of Computational and Graphical Statistics*. 1996;**5**(1):1-25. DOI: 10.2307/1390750. JSTOR 1390750

[36] Carvalho H, Del Moral P, Monin A, Salut G. Optimal non-linear filtering in GPS/INS integration. *IEEE Transactions on Aerospace and Electronic Systems*. 1997;**33**(3):835-850

[37] Del Moral P, Rigal G, Salut G. Nonlinear and Non-Gaussian Particle Filters Applied to Inertial Platform Repositioning. LAAS-CNRS, Toulouse, Research Report No. 92207, STCAN/DIGILOG-LAAS/CNRS Convention STCAN No. A.91.77.013. 1991. p. 94

[38] Crisan D, Gaines J, Lyons T. Convergence of a branching particle method to the solution of the Zakai. *SIAM Journal on Applied Mathematics*. 1998;**58**(5):1568-1590. DOI: 10.1137/s0036139996307371

[39] Crisan D, Lyons T. Nonlinear filtering and measure-valued processes. *Probability Theory and Related Fields*. 1997;**109**(2):217-244. DOI: 10.1007/s004400050131