

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

185,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Aptamers: Magic Bullet for Theranostic Applications

Arghya Sett

Abstract

Aptamers are a short polymer of oligonucleotides (natural or modified) that can bind to its cognate target (small molecules to large macromolecules like proteins, cells, microorganisms etc.) with high affinity and selectivity. They can fold into unique secondary and tertiary conformation in solution (pH, ionic concentration) and bind to their targets in a specific manner (binding constants in sub-nano to pico molar range). They rival the monoclonal antibodies and other specific biological ligands with respect to affinity, stability, robustness, non-immunogenicity and facile to synthesis. Nucleic acid aptamers are selected from an oligonucleotide library by an iterative process called SELEX (Systematic Evolution of Ligands by Exponential Enrichment Analysis). These aptamers are compatible to any kind of chemical modification, conjugation and functionalization. Briefly, this chapter discusses about the diagnostic and therapeutic application of aptamers.

Keywords: aptamers, SELEX, theranostics, chemical antibodies

1. Introduction

Aptamers (Latin word *aptus* means 'to fit' and Greek meros meaning 'part') are single stranded oligonucleotides, which act as synthetic ligands for its cognate target molecules. [1, 2] These molecules show high target specificity, selectivity and affinity, which resemble 'monoclonal antibody'. Similar to antibodies, aptamers also have immense potential to interact with their targets by structural recognition and thus they are termed as 'chemical antibodies'. [3] Different conformations allow aptamers to bind specifically with their target by 'lock and key' model. This hypothesis of binding mechanism is driven by the secondary and tertiary structure of aptamers in-bound state with their targets. Adopting various structures like hairpin loops, bulges, stem-loop, quartets, G-quadruplex, pseudo knots, aptamers can fit into the binding region of the target. [4] Intra and inter molecular interactions like hydrogen bonding, Vander Waals force, hydrophobic interaction, electrostatic forces play major crucial role in aptamer-target interaction. [5] However, the aptamers are primarily synthetic molecules and naturally occurring ribozymes are single stranded RNA molecules, which also have similar recognition domain acting in a similar manner. [6, 7] Aptamers are capable of forming various stable three-dimensional structures in physiological solution. The folding process in solution and the ligand-induced conformational switch is strongly dependent on the presence of divalent cations (magnesium, potassium etc.). [8] There are plethora of computer algorithms

enable sequence based modeling of secondary structure of the oligonucleotide aptamers which actually strengthen the predictability of strongest binders with lowest free energy. [9, 10] Aptamers fold into tertiary conformations and bind to their targets through shape complementarity at the aptamer-target interface. [11] An aptamer binds to a protein can modulate protein functions by interfering with protein interaction with natural partners. Similar to antibodies, aptamers can enter to specific target cells via receptor-mediated endocytosis upon binding to cell surface ligands. [12] Importantly, aptamers can penetrate into tumor cores much more efficiently than antibodies due to their ~20–25-fold smaller sizes compared with full sized monoclonal antibodies. [13, 14]

Monoclonal Antibody	Aptamer
Stability: Monoclonal antibodies require refrigeration to avoid denaturation. Limited shelf life. [15]	Stability: Aptamers do not require refrigeration. Indefinite shelf life. [16]
Immunogenicity: They can cause immunogenic response. [17]	Immunogenicity: Aptamers are non-immunogenic. [18]
Production: laborious, expensive, high batch-to-batch variation.	Production: simpler and controlled chemical reactions, little to no variation, automated, chemical synthesis, no contamination.
Size: Larger in size, they can resist filtration by the kidneys, long half-lives. However, their size prevents access to smaller areas. [19]	Size: Aptamers are small molecules. They are especially subject to kidney filtration, resulting in short half-lives. Compared to antibodies, aptamers can bind to smaller targets. [20]
Ability to modification: Antibodies cannot accommodate conjugates without negative consequences such as reduced activity.	Ability to modification: Easy to modify, modifications can also be incorporated during synthesis to prevent kidney filtration. [21]

Table 1.
Comparison between Monoclonal antibody vs. Aptamer:




Figure 1.
Publication trend for Search strings: “Aptamers as diagnostics” and “Aptamers as therapeutics” (Source: Scopus).




Figure 2.
 Publication trend for Search strings: “Aptamers as theranostics” (Source: Scopus).

Compared to antibodies, aptamers can be produced using cell-free chemical synthesis and are therefore less expensive for large-scale manufacture. Aptamers exhibit extremely low variability between batches and have better controlled post-production modification, they are minimally immunogenic, and are small in size. (Table 1) The rapidly growing aptamer industry was predicted to reach US \$244.93 million by 2020. [22] Presently more than 40 companies are actively engaged in diagnostics and therapeutics research to commercialize these “magic bullets” globally (EU countries, Asia, USA, UK etc.). [23] The largest company is “SomaLogic” (company based on SOMAmer- a patented “Slow Off-rate Modified Aptamer) founded by Prof. Larry Gold at Colorado, USA. Since the advent of aptamers scientists and researchers exploit different applications of aptamers that reflects the following trends in the publications. (Figures 1 and 2).

2. Basics of SELEX screening process

Back in 1990, two individual groups Prof. Larry Gold and Craig Tuerk from University of Boulder, USA and Prof. Jack Szostak and his student A.D. Ellington from Havard University, USA discovered the evolution process to obtain the oligonucleotide binders and they coined the term ‘Aptamer’ and the process as ‘SELEX’. [24, 25] Systematic Evolution of Ligands by Exponential Enrichment (SELEX) is a common screening process by which aptamers can be selected from an aptamer library which consist of 10^{24-25} number of various sequences. The method attempts to isolate an aptamer of interest from a pool of randomized library by an iterative cycle of incubation with the target, partitioning and amplification, until the pool of aptamers enriched enough to fit with the target. The SELEX procedure iterates over five basic steps- incubation of aptamer pools with the target, binding, partitioning and washing (to get rid of non-binders which are loosely bound with the target), then elution of positive target-bound aptamers and amplification of enriched pools. Traditionally, the positive pool eluted from last round is being analysed, and high-throughput sequencing is performed.

An array of different RNA and DNA aptamers were isolated against a vast array of targets: ions, [26] low molecular weight metabolites, [27, 28] proteins, [29–31] sugar moieties [32] lipids, [33] and even whole cells. [34, 35]

3. Library selection

To select highly selective, specific aptamers, design of the initial aptamer library is the first and foremost step. In case of determination of the length of the random region researchers should consider the sequence space and structural diversity. The complexity of the initial aptamer library depends on the length of the random window of the aptamer library (If the random window is 40 and if we consider DNA aptamer library, so the complexity of the library is: 4^{40} that equals to 10^{24-25}). [36]

Special libraries would consist of specifically designed flanking sequences directing the aptamers to form a specific secondary structure, or include modified nucleotides. In capture SELEX, there is unique docking sequence (12–14 nucleotides long) which enables the library in such a way, that highly sensitive aptamers can be fished out against small molecules. [37, 38] The extended genetic alphabets or combination of artificial xeno nucleic acids (XNA) greatly broaden the diversity of sequences and can influence the properties of the aptamers, such as their in vivo stability or nuclease resistance. [39–42] Modified nucleotides can be introduced either during the library synthesis or in the post-selection optimization.

In a review article, Maria *et al.* summarized all key features of designing nucleic acid libraries for SELEX like nature, composition of the library (RNA, DNA or modified nucleotides), the length of a randomized region and the presence of fixed sequences. Different randomization strategies and computer algorithms of library designs were also discussed. [43]

4. Various SELEX processes

Specific aptamers are screened by the iterative process of SELEX from a highly diverse pool of oligonucleotides. [44–46] After the incubation of the random aptamer pool with the target followed by the removal of non-binding aptamers, the bound aptamer species are recovered. These recovered nucleic acid sequences are amplified with PCR (in the case of DNA aptamer) or RT-PCR (for RNA aptamers). In addition to selection against a purified target molecule, SELEX process can be performed against live bacterial cells and even in mammalian cell lines to isolate cancer cell specific aptamers and furthermore it can lead to the identification of novel biomarkers. [47, 48]




Figure 3.
Typical schema of Cross-over SELEX process.

A giant advancement of SELEX technology has been made since its discovery in 1990. Conventional SELEX is a well-established and effective method but due to its immense time- and labor-consumption, continuous improvement of alternative methods for aptamer selection has been inevitable.

High throughput SELEX (HT-SELEX), Functional screening (Microfluidics or Flow cytometry based SELEX), Cross-over SELEX (where the target is alternatively changing from proteins and cells), (**Figure 3**) *in-vivo* SELEX, Spiegelmers selection, de-convolution SELEX are few examples of modern-era screening strategy of aptamers. [49] Cutting-edge functional screening process, the chemical modifications, Next-generation sequencing (NGS technology) enable SELEX more efficient, cost-effective and considerably less time-consuming.

5. Modifications of naturally occurring aptamers

DNA is the backbone of central dogma of our life cycle. Moreover, any form of nucleic acids play a crucial role in our genetic codon. DNA/RNA is an essential biomacromolecule consist of nucleotide bases such as adenine (A), thymine (T), uracil (U), guanine (G), cytosine (C).

There are various types of modifications (nucleotide base modifications, phosphate backbone modifications, peptide mimic oligonucleotides PNA etc.) available which can prevent aptamers from nuclease degradation. Locked nucleic acid (LNA) is one among them where 2'-oxygen has been linked to the 4'-carbon of the ribose sugar by a methylene bridge, thus completely locking the sugar into a 3'-endo conformation. LNAs increase the thermodynamic stability, binding affinity, and enable the oligonucleotides to prevent serum degradation. [50–52] These modifications enable the aptamers for biological applications.

Compared to LNAs, the unlocked nucleic acid (UNA) is an acyclic ribose derivative that has increased flexibility. UNAs do not consist the C2'-C3' bond, which confers the flexibility observed in this modified nucleotide. [53] LNAs increase the melting temperature of the nucleotide by 1–10°C per LNA insertion but UNAs reduce the melting temperature by 5–10°C retaining the nuclear resistance. In case of, Peptide nucleic acid (PNA) in which sugar-phosphate backbone is modified by short stretch of N-(2-aminoethyl)-glycine units connected by peptide bonds, enhances biostability of the modified candidates. [54]

5.1 Aptamers in Drug development pipeline

Aptamers have been incorporated in drug development pipeline as they have the capacity to block the downstream signalling (phosphorylation of kinases etc.) of different biomolecules. They can play an important role to regulate various cellular crosstalks. To screen therapeutic aptamers either DNA aptamers or 2'-fluoro modified RNA, a combination of 2'-fluoro pyrimidines and 2'-hydroxyl purines (fYrR) are of major interest. fYrR is the “nuclease stable RNA” and can be easily generated by Y639F modified T7 RNA polymerase. Fovista, an anti-platelet derived growth factor (PDGF) aptamer, was previously DNA aptamer but later modified to augment the stability with the addition of backbone modifications. [55] As with the 2'-fluoro modification, the 2'-OMe modifications adopt a C3'-endo conformation. US FDA approved the first aptamer (Macugen®, pegaptanib sodium) in 2004 against vascular endothelial growth factor for the treatment of age-related macular degeneration. [56] This aptamer was modified with 2'-fluoro-pyrimidines and 2'-O-methyl-purines. The stability of the small aptamer was a critical factor but later which can be circumvented with a 3'-cap and a polyethylene glycol

molecule, the half-life of Macugen® was extended to 131 hours at max. [57, 58] Anti-vascular endothelial growth factor (VEGF 165) aptamer Macugen, and an anti-Factor IXa aptamer REG1 were both selected from fYrR libraries, and subsequently 2'-O-methyl nucleosides have been incorporated in order to increase serum stability. [57]

There is a plethora of polymerase enzymes like KOD, Pwo, Phusion, Superscript III, vent (exo-), T7 polymerase have all been shown to be capable of incorporating modified triphosphates into DNA and RNA strands, which open up a new opportunities in aptamer selection strategies. [59] The use of Pfx DNA polymerase allows amplification of Ds-Px base pair in Ex-SELEX protocol where extended genetic alphabets were included in complexity of nucleic acid library. [60]

Several limitations of aptamers should be considered in the process of *in-vivo* applications of nucleic acid aptamers. Being polynucleotides, nucleic acid aptamers are naturally susceptible to enzymes degradation by exo and/or endo-nucleases, leading to a reduced *in vivo* circulatory half-life. This drawback can be alleviated by side chain chemical modifications to aptamers, incorporating unnatural nucleotide bases (locked and unlocked nucleic acids) and capping the aptamer ends, thus minimizing the susceptibility to endonuclease and exonuclease attack. [50, 51] Short blood residence time is another challenge with *in vivo* aptamer applications, which is due to fast removal of aptamer from the circulation by renal filtration as most aptamers have a size smaller than the renal filtration threshold of 40 kDa. [31] To achieve desired serum half-life, aptamers can be engineered by conjugation with a terminal polyethylene glycol (PEG), although this may compromise the extent of tumor penetration [61]. In some cases, post-SELEX modifications following the selection of aptamers may alter the 3-D structure of the aptamers, leading to the lost or altered binding affinity and specificity. Such problems can be prevented by using random aptamer pools containing modified nucleotides during the SELEX process. [62, 63] In addition, the ability of aptamers to interact with cells may decrease due to repulsion of nucleic acids by negatively-charged cell membranes. This can be refuted by increasing the binding affinity and specificity of aptamers toward their cell surface receptors to trigger receptor-mediated endocytosis.

In the field of oncology, two aptamers, namely, AS1411 and NOX-A12, have entered clinical trials. [45, 64] AS1411 (formerly ARGO100; Antisoma) is a guanine quadruplex aptamer obtained from a guanine-rich oligonucleotide library in the anti-proliferation screen, which is not a typical SELEX process. [65] The guanine quadruplex structure benefits AS1411 because it is resistant to nuclease degradation and enhances cell uptake. In *in-vitro* validations, AS1411 inhibits more than 80 types of cancer cell lines. In addition, the target of AS1411 has been verified to be nucleolin, and the relevant mechanism and specificity against cancer cells have also been described. In preclinical tests, AS1411 shows efficacy toward xenograft models, including non-small cell lung, renal, and breast cancers. It entered clinical trials in 2003 and passed phase II trials for acute myeloid leukemia. A subsequent phase II trial for renal cell carcinoma started in 2008 (clinical trial ID NCT00740441). [66] NOXA12 (Olaptesed pegol; Noxxon) is an L-form RNA aptamer known as Spiegelmer and is used for cancer therapy. NOX-A12 can bind to its target chemokine CXCL-12 and blocks its interaction with its receptor. [67] This disrupts the tissue gradient of CXCL-12 and reduces the cancer cell homing that might lead to tumor metastasis and drug resistance. [68] Currently, phase II clinical trials for NOX-A12 are underway for the treatment of chronic lymphocytic leukemia and refractory multiple myeloma (clinical trial IDs NCT01486797 and NCT01521533). [67] Aptamer based cancer therapeutics have immense potential for precise and less toxic treatment for cancer patients. [46]

6. Aptamers as diagnostic agents

Aptamers can be used *in-vitro* and *in-vivo* as well. [69] In terms of *in vivo* diagnostics, ‘escort’ aptamers can be implied as vehicles for a detectable molecules, such as radionuclides, fluorophores etc. [70–72] The development of new agents like radio-pharmaceuticals is challenging. There are some important factors such as efficiency of the radiolabeling process, specific activity (radioactivity per moles e.g. Ci/ μmol), chemical purity, radiochemical and chemical stability and shelf life of the final product. [73] Mostly, radiolabeling strategies for aptamers are similar as for proteins, or antibodies. Aptamers can be easily chemically modified at its 5’ or 3’ end with a desired functional group for radiolabeling (**Figure 4**).

Radiohalogens (fluorine-18, bromine-76, iodine-125 etc.) are the most commonly used for radiolabelling oligonucleotides which are often accompanied with prosthetic groups. Recently, click-chemistry for radiofluorination was demonstrated on antisense oligonucleotides and siRNAs. [74, 75] Another report used photoconjugation as strategy for the radiofluorination of an aptamer. [76] Oligonucleotides have also been radiolabeled with the radiohalogens such as bromine-76 for PET imaging and iodine-123 for SPECT (Single photon emission computed tomography) imaging. In addition, iodine-125 has been used to radiolabel antisense oligonucleotides, aptamers and spiegelmers for theranostic applications. Due to the harsh and non-aqueous reaction conditions usually needed to radiolabel prosthetic groups, it is performed before the conjugation process to the oligonucleotide. [73]




Figure 4.

Aptamers modified with radiolabelled molecules for disease diagnosis (Figure adapted from Gijs et al) [73].

Till date, a plethora of aptamers have been modified or labelled with radioactive molecules. Aptamers against several important biomarkers like PMSA, Tenascin C, thrombin, MUC1 were already exploited for radiolabelling. Aptamer-based radiopharmaceuticals were primarily developed for imaging and therapy of cancer diseases, metabolic disorders and others. The aptamers are mainly radiolabeled with technetium-99 m for SPECT (Single photon emission computed tomography), PET (Positron emission tomography) imaging. Very few aptamers were published related to PET imaging, and there is only one study of radiolabeled aptamers for therapy by Bandekar *et al.* [77] Other radiolabeled aptamers have only been tested for preclinical applications or in the course of preclinical assesement.

Molecular nuclear imaging technique is a diagnostic process of non-invasive visualization of any disease *in-vivo* at molecular level with high precision. For nuclear imaging, the probes used for radiolabelling has to be modified accordingly. Aptamers are the most promising candidates with versatile modification capibility, can be easily engineered for various imaging and other diagnostic purposes.

The first radiolabeled aptamer for nuclear imaging was discovered by Charlton *et al.* A DNA aptamer, NX21909, was selected against human neutrophil elastase, an enzyme which is secreted by neutrophils and macrophages during inflammation to kill pathogens. [78]

Aptamer TTA1, an RNA aptamer targeting the extracellular matrix protein tenascin C (TN-C), was the first radiolabeled aptamer which was used as molecular cancer imaging agent. Aptamer TTA1 was generated by a cross-over SELEX involving the purified recombinant TN-C protein and TN-C-positive U251 glioblastoma cells. [79, 80]

7. Lightup aptasensors for diagnostic applications

There are a unique group of aptamers (generally RNA aptamers) which can bind specifically with their cognate fluorogen molecules like DFHBI, thiazole orange, thioflavin T etc. [49, 81, 82] The non-fluorescent moelcules (native unbound state) become fluorescent (bound state) after binding to the aptamers and these “light-up” aptamers generate fluorescence signal. In the omni-presence of target molecules (small pre-miRNAs) and malachite green (fluorogen) light up aptasensors ‘malaswitch’ exhibit fluorescence enhancement. [83] We can engineer the small-molecule specific aptamers (like aptamers for some pesticides, toxins, small metaboltes) in such a way, that combined with light-up aptamers, they can generate a detectable signal. Light up aptasensors are promising alternative biosensor for label free sensitive detection of small molecules. [84]

8. Future perspectives

With more focus on *in vivo* studies for potential clinical applications, aptamers can be developed in combination with DNA nanostructures, nanomaterial, or microfluidic devices as diagnostic probes or therapeutic agents for cancers, infectious diseases, genetic, metabolic, neurological disorders, lifestyle diseases and several others. The use of aptamers as targeting agents in drug delivery can also be explored. Aptamers might be exploited to develop portable, low-cost and robust diagnostic kit using simple devices for real-time and on-site POC (point-of-care) detection and monitoring, instead of the laborious and time-consuming diagnostic tests currently available only in clinical labs. Regarding therapeutics approach, there is still untapped potential in the combination of the target recognition and strong


binding property of aptamers with exquisitely designed nanomaterials. It can be used as an effective alternative drug delivery platform. Variety of materials such as liposomes, polymer vesicles and silica nanoparticles, combined with DNA/RNA aptamers, has shown feasibility for use in *in vivo* targeted drug delivery. [85, 86] The integration of diagnostic capability with therapeutic interventions termed, as “Theranostics” is critical to address the challenges of disease heterogeneity and adaptation. Although aptamers have immense potential as theranostic agents, tailor-made modifications, validation of experiments need to be executed before aptamer-based drug delivery can reach clinical trials and eventually the patient management system.

Author details

Arghya Sett
Institute of Organic Chemistry and Biochemistry, Academy of Sciences of the
Czech Republic, Prague, Czech Republic

*Address all correspondence to: arghya.sett@gmail.com

IntechOpen

© 2020 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

References

- [1] Famulok M, Mayer G. Aptamers and SELEX in chemistry & biology. Vol. 21, Chemistry and Biology. 2014.
- [2] Nimjee SM, White RR, Becker RC, Sullenger BA. Aptamers as Therapeutics. *Annu Rev Pharmacol Toxicol* [Internet]. 2017 Jan 6;57:61-79. Available from: <https://pubmed.ncbi.nlm.nih.gov/28061688>
- [3] Bauer M, Strom M, Hammond DS, Shigdar S. Anything You Can Do, I Can Do Better: Can Aptamers Replace Antibodies in Clinical Diagnostic Applications? *Molecules*. 2019;24(23):4377.
- [4] Cai S, Yan J, Xiong H, Liu Y, Peng D, Liu Z. Investigations on the interface of nucleic acid aptamers and binding targets. *Analyst* [Internet]. 2018;143(22):5317-38. Available from: <http://dx.doi.org/10.1039/C8AN01467A>
- [5] Elskens JP, Elskens JM, Madder A. Chemical Modification of Aptamers for Increased Binding Affinity in Diagnostic Applications: Current Status and Future Prospects. *Int J Mol Sci*. 2020;21(12):4522.
- [6] Breaker RR, Joyce GF. The Expanding View of RNA and DNA Function. *Chem Biol* [Internet]. 2014;21(9):1059-65. Available from: <http://linkinghub.elsevier.com/retrieve/pii/S1074552114002373>
- [7] Doudna JA, Cech TR. The chemical repertoire of natural ribozymes. *Nature*. 2002;418(6894):222-8.
- [8] Horiya S, Li X, Kawai G, Saito R, Katoh A, Kobayashi K, et al. RNA LEGO: magnesium-dependent formation of specific RNA assemblies through kissing interactions. *Chem Biol*. 2003;10(7):645-54.
- [9] Zuker M. Mfold web server for nucleic acid folding and hybridization prediction. *Nucleic Acids Res* [Internet]. 2003 Jul 1;31(13):3406-15. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC169194/>
- [10] Wolfe BR, Porubsky NJ, Zadeh JN, Dirks RM, Pierce NA. Constrained multistate sequence design for nucleic acid reaction pathway engineering. *J Am Chem Soc*. 2017;139(8):3134-44.
- [11] Khati M. The Future of Aptamers in Medicine. *J Clin Pathol* [Internet]. 2010;63(6):480-7. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/20360137>
- [12] Cibiel A, Dupont DM, Ducongé F. Methods to identify aptamers against cell surface biomarkers. *Pharmaceuticals*. 2011;4(9):1216-35.
- [13] Soper SA, Brown K, Ellington A, Frazier B, Garcia-Manero G, Gau V, et al. Point-of-care biosensor systems for cancer diagnostics/prognostics. *Biosens Bioelectron*. 2006;21(10):1932-42.
- [14] Brody EN, Willis MC, Smith JD, Jayasena S, Zichi D, Gold L. The use of aptamers in large arrays for molecular diagnostics. Vol. 4, *Molecular Diagnosis*. 1999. p. 381-8.
- [15] Le Basle Y, Chennell P, Tokhadze N, Astier A, Sautou V. Physicochemical stability of monoclonal antibodies: a review. *J Pharm Sci*. 2020;109(1):169-90.
- [16] Bruno JG. Long shelf life of a lyophilized DNA aptamer beacon assay. *J Fluoresc*. 2017;27(2):439-41.
- [17] Kozlowski S, Swann P. Current and future issues in the manufacturing and development of monoclonal antibodies. *Adv Drug Deliv Rev*. 2006;58(5-6):707-22.

- [18] Esposito CL, Passaro D, Longobardo I, Condorelli G, Marotta P, Affuso A, et al. A Neutralizing RNA Aptamer against EGFR Causes Selective Apoptotic Cell Death. *PLoS One* [Internet]. 2011;6(9):e24071. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3167817&tool=pmcentrez&rendertype=abstract>
- [19] Šmuc T, Ahn I-Y, Ulrich H. Nucleic acid aptamers as high affinity ligands in biotechnology and biosensorics. *J Pharm Biomed Anal*. 2013;81:210-7.
- [20] Majumder P, Gomes KN, Ulrich H. Aptamers: from bench side research towards patented molecules with therapeutic applications. *Expert Opin Ther Pat* [Internet]. 2009 Nov 1;19(11):1603-13. Available from: <https://doi.org/10.1517/13543770903313746>
- [21] Meek KN, Rangel AE, Heemstra JM. Enhancing aptamer function and stability via in vitro selection using modified nucleic acids. *Methods*. 2016 Aug;106:29-36.
- [22] Kedzierski S, Khoshnejad M, Caltagirone GT. Synthetic antibodies: the emerging field of aptamers. *Bioprocess J*. 2012;11:46-9.
- [23] Kaur H, Bruno JG, Kumar A, Sharma TK. Aptamers in the Therapeutics and Diagnostics Pipelines. *Theranostics* [Internet]. 2018 Jul 1;8(15):4016-32. Available from: <https://pubmed.ncbi.nlm.nih.gov/30128033>
- [24] Ellington AD, Szostak JW. In vitro selection of RNA molecules that bind specific ligands. *Nature*. 1990;346(6287):818.
- [25] Tuerk C, Gold L. Systematic evolution of ligands by exponential enrichment: RNA ligands to bacteriophage T4 DNA polymerase. *Science*. 1990;249(4968):505-10.
- [26] Hofmann H-P, Limmer S, Hornung V, Sprinzl M. Ni²⁺-binding RNA motifs with an asymmetric purine-rich internal loop and a GA base pair. *Rna*. 1997;3(11):1289-300.
- [27] Huizenga DE, Szostak JW. A DNA aptamer that binds adenosine and ATP. *Biochemistry*. 1995;34(2):656-65.
- [28] Miyachi Y, Shimizu N, Ogino C, Fukuda H, Kondo A. Selection of a DNA aptamer that binds 8-OHdG using GMP-agarose. *Bioorg Med Chem Lett*. 2009;19(13):3619-22.
- [29] Sett A, Borthakur BB, Sharma JD, Kataki AC, Bora U. DNA aptamer probes for detection of estrogen receptor α positive carcinomas. *Transl Res*. 2017;183:104-20.
- [30] Sett A, Borthakur BB, Bora U. Selection of DNA aptamers for extra cellular domain of human epidermal growth factor receptor 2 to detect HER2 positive carcinomas. *Clin Transl Oncol*. 2017;19(8).
- [31] Ahirwar R, Vellarikkal SK, Sett A, Sivasubbu S, Scaria V, Bora U, et al. Aptamer-Assisted Detection of the Altered Expression of Estrogen Receptor Alpha in Human Breast Cancer. *PLoS One* [Internet]. 2016;11(4):e0153001. Available from: <http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0153001>
- [32] Boese BJ, Breaker RR. In vitro selection and characterization of cellulose-binding DNA aptamers. *Nucleic Acids Res*. 2007;35(19):6378-88.
- [33] Betat H, Vogel S, Struhalla M, Förster H-H, Famulok M, Welzel P, et al. Aptamers that recognize the lipid moiety of the antibiotic moenomycin A. *Biol Chem*. 2003;384(10-11):1497-500.
- [34] Dwivedi HP, Smiley RD, Jaykus L-A. Selection and characterization of DNA aptamers with binding selectivity to

Campylobacter jejuni using whole-cell SELEX. *Appl Microbiol Biotechnol*. 2010;87(6):2323-34.

[35] Wang G, Liu J, Chen K, Xu Y, Liu B, Liao J, et al. Selection and characterization of DNA aptamer against glucagon receptor by cell-SELEX. *Sci Rep*. 2017;7(1):1-10.

[36] Pobanz K, Lupták A. Improving the odds: Influence of starting pools on in vitro selection outcomes. *Methods*. 2016;106:14-20.

[37] Stoltenburg R, Nikolaus N, Strehlitz B. Capture-SELEX: selection of DNA aptamers for aminoglycoside antibiotics. *J Anal Methods Chem*. 2012;2012.

[38] Paniel N, Istamboulié G, Triki A, Lozano C, Barthelmebs L, Noguér T. Selection of DNA aptamers against penicillin G using Capture-SELEX for the development of an impedimetric sensor. *Talanta*. 2017;162:232-40.

[39] Taylor AI, Houlihan G, Holliger P. Beyond DNA and RNA: The Expanding Toolbox of Synthetic Genetics. *Cold Spring Harb Perspect Biol*. 2019 Jun;11(6).

[40] Lee J, Schwieter KE, Watkins AM, Kim DS, Yu H, Schwarz KJ, et al. Expanding the limits of the second genetic code with ribozymes. *Nat Commun*. 2019 Nov;10(1):5097.

[41] Morihiro K, Kasahara Y, Obika S. Biological applications of xeno nucleic acids. *Mol Biosyst*. 2017 Jan;13(2):235-45.

[42] Rangel AE, Chen Z, Ayele TM, Heemstra JM. In vitro selection of an XNA aptamer capable of small-molecule recognition. *Nucleic Acids Res*. 2018;46(16).

[43] Vorobyeva MA, Davydova AS, Vorobjev PE, Venyaminova AG. Key

Aspects of Nucleic Acid Library Design for in Vitro Selection. *Int J Mol Sci* [Internet]. 2018 Feb 5;19(2):470. Available from: <https://pubmed.ncbi.nlm.nih.gov/29401748>

[44] Jayasena SD. Aptamers: An emerging class of molecules that rival antibodies in diagnostics. *Clin Chem*. 1999;45(9):1628-50.

[45] Keefe AD, Pai S, Ellington A. Aptamers as therapeutics. *Nat Rev Drug Discov* [Internet]. 2010;9(7):537-50. Available from: <http://dx.doi.org/10.1038/nrd3141>

[46] Xiang D, Shigdar S, Qiao G, Wang T, Kouzani AZ, Zhou S-F, et al. Nucleic Acid Aptamer-Guided Cancer Therapeutics and Diagnostics: the Next Generation of Cancer Medicine. *Theranostics* [Internet]. 2015 Jan 1;5(1):23-42. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4265746/>

[47] Cho EJ, Lee J-W, Ellington AD. Applications of aptamers as sensors. *Annu Rev Anal Chem (Palo Alto Calif)*. 2009;2:241-64.

[48] Li S, Xu H, Ding H, Huang Y, Cao X, Yang G, et al. Identification of an aptamer targeting hnRNP A1 by tissue slide-based SELEX. 2009;(February):327-36.

[49] Autour A, Bouhedda F, Cubi R, Ryckelynck M. Optimization of fluorogenic RNA-based biosensors using droplet-based microfluidic ultrahigh-throughput screening. *Methods*. 2019;161:46-53.

[50] Barciszewski J, Medgaard M, Koch T, Kurreck J, Erdmann VA. Locked Nucleic Acid Aptamers. In: *Nucleic Acid and Peptide Aptamers: Methods and Protocols* [Internet]. 2009. p. 165-86. Available from: <http://www.springerlink.com/index/10.1007/978-1-59745-557-2>

- [51] Veedu RN, Wengel J. Locked nucleic acids: promising nucleic acid analogs for therapeutic applications. *Chem Biodivers*. 2010;7(3):536-42.
- [52] Darfeuille F, Hansen JB, Orum H, Di Primo C, Toulmé J-J. LNA/DNA chimeric oligomers mimic RNA aptamers targeted to the TAR RNA element of HIV-1. *Nucleic Acids Res* [Internet]. 2004 Jun 4;32(10):3101-7. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/15181175>
- [53] Langkjær N, Pasternak A, Wengel J. UNA (unlocked nucleic acid): a flexible RNA mimic that allows engineering of nucleic acid duplex stability. *Bioorg Med Chem*. 2009;17(15):5420-5.
- [54] Wittung P, Nielsen PE, Buchardt O, Egholm M, Norde B. DNA-like double helix formed by peptide nucleic acid. *Nature*. 1994;368(6471):561-3.
- [55] Floege J, Ostendorf T, Janssen U, Burg M, Radeke HH, Vargeese C, et al. Novel approach to specific growth factor inhibition in vivo: antagonism of platelet-derived growth factor in glomerulonephritis by aptamers. *Am J Pathol*. 1999;154(1):169-79.
- [56] Santosh B, Yadava PK. Nucleic Acid Aptamers: Research Tools in Disease Diagnostics and Therapeutics. *Biomed Res Int* [Internet]. 2014;2014:1-13. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4090538&tool=pmcentrez&rendertype=abstract>
- [57] Ruckman J, Green LS, Beeson J, Waugh S, Gillette WL, Henninger DD, et al. 2'-Fluoropyrimidine RNA-based aptamers to the 165-amino acid form of vascular endothelial growth factor (VEGF165) inhibition of receptor binding and VEGF-induced vascular permeability through interactions requiring the exon 7-encoded domain. *J Biol Chem*. 1998;273(32):20556-67.
- [58] Lee J-H, Canny MD, De Erkenez A, Krilleke D, Ng Y-S, Shima DT, et al. A therapeutic aptamer inhibits angiogenesis by specifically targeting the heparin binding domain of VEGF165. *Proc Natl Acad Sci*. 2005;102(52):18902-7.
- [59] Lipi F, Chen S, Chakravarthy M, Rakesh S, Veedu RN. In vitro evolution of chemically-modified nucleic acid aptamers: Pros and cons, and comprehensive selection strategies. *RNA Biol* [Internet]. 2016 Dec 1;13(12):1232-45. Available from: <https://doi.org/10.1080/15476286.2016.1236173>
- [60] Kimoto M, Matsunaga K, Hirao I. Evolving Aptamers with Unnatural Base Pairs. *Curr Protoc Chem Biol*. 2017;9(4):315-39.
- [61] Ng EWM, Shima DT, Calias P, Cunningham ET, Guyer DR, Adamis AP. Pegaptanib, a targeted anti-VEGF aptamer for ocular vascular disease. *Nat Rev Drug Discov*. 2006;5(2):123-32.
- [62] Sampson T. Aptamers and SELEX: The technology. *World Pat Inf*. 2003;25(2):123-9.
- [63] Shamah SM, Healy JM, Cload ST. Complex target SELEX. *Acc Chem Res*. 2008;41(1):130-8.
- [64] Lao Y-H, Phua KKL, Leong KW. Aptamer nanomedicine for cancer therapeutics: barriers and potential for translation. *ACS Nano*. 2015;9(3):2235-54.
- [65] Carvalho J, Paiva A, Campello MPC, Paulo A, Mergny J-L, Salgado GF, et al. Aptamer-based targeted Delivery of a G-quadruplex Ligand in Cervical Cancer Cells. *Sci Rep*. 2019;9(1):1-12.
- [66] Rosenberg JE, Bambury RM, Van Allen EM, Drabkin HA, Lara PN, Harzstark AL, et al. A phase II trial of AS1411 (a novel nucleolin-targeted DNA aptamer) in metastatic renal

cell carcinoma. *Invest New Drugs*. 2014;32(1):178-87.

[67] Hoellenriegel J, Zboralski D, Maasch C, Rosin NY, Wierda WG, Keating MJ, et al. The Spiegelmer NOX-A12, a novel CXCL12 inhibitor, interferes with chronic lymphocytic leukemia cell motility and causes chemosensitization. *Blood*. 2014;123(7):1032-9.

[68] Kruschinski MGSC-CMJTDZZR. Anti-CXCL12/SDF-1 Spiegelmer® Nox-A12 alone and in combination with bendamustine and rituximab in patients with relapsed chronic lymphocytic leukemia (CLL): results from a phase IIa study [Abstract 1635]. 55th American Society of Hematology Annual Meeting and Exhibition. 2013.

[69] Pestourie C, Tavitian B, Duconge F. Aptamers against extracellular targets for in vivo applications. *Biochimie*. 2005;87(9-10):921-30.

[70] Zhou J, Rossi JJ. Cell-specific aptamer-mediated targeted drug delivery. *Oligonucleotides*. 2011;21(1):1-10.

[71] Li N, Nguyen HH, Byrom M, Ellington AD. Inhibition of cell proliferation by an anti-EGFR aptamer. *PLoS One*. 2011;6(6):e20299.

[72] Hicke BJ, Stephens AW. Escort aptamers: a delivery service for diagnosis and therapy. *J Clin Invest*. 2000;106(8):923-8.

[73] Gijs M, Aerts A, Impens N, Baatout S, Luxen A. Aptamers as radiopharmaceuticals for nuclear imaging and therapy. *Nucl Med Biol*. 2016;43(4):253-71.

[74] Inkster JAH, Adam MJ, Storr T, Ruth TJ. Labeling of an antisense oligonucleotide with [18F] FPy5yne. *Nucleosides, Nucleotides and Nucleic Acids*. 2009;28(11-12):1131-43.

[75] Mercier F, Paris J, Kaisin G, Thonon D, Flagothier J, Teller N, et al. General method for labeling siRNA by click chemistry with fluorine-18 for the purpose of PET imaging. *Bioconjug Chem*. 2011;22(1):108-14.

[76] Lange CW, VanBrocklin HF, Taylor SE. Photoconjugation of 3-azido-5-nitrobenzyl-[18F] fluoride to an oligonucleotide aptamer. *J Label Compd Radiopharm*. 2002;45(3):257-68.

[77] Bandekar A, Zhu C, Jindal R, Bruchertseifer F, Morgenstern A, Sofou S. Anti-prostate-specific membrane antigen liposomes loaded with 225Ac for potential targeted antivasculature α -particle therapy of cancer. *J Nucl Med*. 2014;55(1):107-14.

[78] Charlton J, Sennello J, Smith D. In vivo imaging of inflammation using an aptamer inhibitor of human neutrophil elastase. *Chem Biol*. 1997;4(11):809-16.

[79] Hicke BJ, Marion C, Chang Y-F, Gould T, Lynott CK, Parma D, et al. Tenascin-C aptamers are generated using tumor cells and purified protein. *J Biol Chem*. 2001;276(52):48644-54.

[80] Hicke BJ, Stephens AW, Gould T, Chang Y-F, Lynott CK, Heil J, et al. Tumor targeting by an aptamer. *J Nucl Med*. 2006;47(4):668-78.

[81] Ouellet J. RNA Fluorescence with Light-Up Aptamers. *Front Chem* [Internet]. 2016;4(June):1-12.

[82] Bouhedda F, Autour A, Ryckelynck M. Light-up RNA aptamers and their cognate fluorogens: from their development to their applications. *Int J Mol Sci*. 2018;19(1):44.

[83] Sett A, Zara L, Dausse E, Toulme J-J. Engineering light-up aptamers for the detection of RNA hairpins through kissing interaction. *Anal Chem*. 2020 Jun 4;0(ja).

[84] Durand G, Lisi S, Ravelet C, Dausse E, Peyrin E, Toulmé JJ. Riboswitches based on kissing complexes for the detection of small ligands. *Angew Chemie - Int Ed.* 2014;53(27):6942-5.

[85] Germer K, Leonard M, Zhang X. RNA aptamers and their therapeutic and diagnostic applications. Vol. 4, *International Journal of Biochemistry and Molecular Biology*. 2013. p. 27-40.

[86] Tan W, Wang H, Chen Y, Zhang X, Zhu H, Yang C, et al. Molecular aptamers for drug delivery. *Trends Biotechnol.* 2011;29(12):634-40.