

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

185,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



# Chronic Obstructive Pulmonary Disease Related to Wood and Other Biomass Smoke: A Different Phenotype or Specific Diseases?

*Carlos A. Torres-Duque, Felipe Severiche-Bueno  
and Mauricio González-García*

## Abstract

Around 41% of the world's population continue using solid fuels, including wood and other types of biomass, for cooking or heating their homes. Long-term indoor exposure to wood smoke, and biomass smoke in general, is a risk factor for developing chronic obstructive pulmonary disease (COPD). In some regions of the world, biomass exposure is a more frequent cause of COPD than exposure to cigarette smoke. Recently it has been described notable differences between COPD associated with wood smoke (WS-COPD) and that caused by tobacco smoking (TS-COPD): significantly less emphysema and more airway inflammation in WS-COPD. Recognizing these differences, some authors have suggested that WS-COPD should be considered a new COPD phenotype. This chapter summarizes the differences between WS-COPD and TS-COPD. The information about the characteristics of COPD caused by other types of biomass fuels, different from wood, is very scarce. Accepting that the smoke derived from wood burning and tobacco smoking have some differences (etiology), the inhalation patterns are different (pathogenesis) and the physiopathological mechanisms they induce may also differ, we analyze if the disease caused by indoor chronic exposure to wood smoke should be considered as another COPD phenotype or a distinct nosological entity.

**Keywords:** chronic obstructive pulmonary disease (COPD), wood smoke, biomass, tobacco, phenotype, emphysema, chronic bronchitis, bronchial anthracofibrosis, indoor air pollution

## 1. Introduction

***Solid and biomass fuels are the most important global environmental risk factor.*** Around 41% of the world's population, over 2.8 billion people, particularly in developing countries, still use solid fuels, whether coal or biomass (wood, vegetable remains and dung), for cooking or heating their homes [1, 2]. In some countries, these fuels are the main source of energy for over 70% of the rural population. In countries where migration from rural areas to cities is high, the population of urban dwellers over the age of 40 years frequently has a significant history of exposure

to biomass fuels. One example is Colombia, where 39% of the population over 40 years of age living in the five main cities had cooked using wood as fuel for more than 10 years before relocating [3].

Between 1980 and 2010, the population exposed to household air pollution (HAP) increased from 333 million to 646 million in sub-Saharan Africa and from 162 million to 190 million in the eastern Mediterranean. In south-east Asia, it remained stable during the same period at around 1 billion people [1].

Biomass fuels are usually burnt in open fires and inefficient traditional cook-stoves, often in poorly ventilated cooking spaces, resulting in indoor high levels of air pollutants including carbon monoxide (CO) and particulate matter (PM). The people most exposed are women who are routinely responsible for cooking and their young children [4].

HAP is responsible for nearly 5% of the global disease burden, making it globally the single most important environmental risk factor [5]. In 2017, it has been estimated that HAP contributed to 1.8 million (95% CI, 1.1–2.7) deaths and 60.9 million (95% CI, 34.6–93.3) disability-adjusted life-years (DALYs) globally [6]. Respiratory disease was the leading cause of these deaths and DALYs attributable to HAP accounting for 38% of all deaths (0.7 million [0.4–1.0]) and 75% of all DALYs (45.7 million [26.8–68.8]) [6]. Among the premature deaths related to HAP, 20% are due to chronic obstructive pulmonary disease (COPD) [2].

**Biomass fuels and COPD.** Several systematic reviews, meta-analyses and reviews of evidence confirm that individuals chronically exposed to solid fuels at home have a higher risk of developing COPD [6–13]. In addition, people chronically exposed to biomass smoke also have a high risk of chronic bronchitis [10, 14, 15]. The pooled analysis of the Pathak's study [7] showed that, globally, exposure to indoor air pollution due to solid biomass fuels increased risk of COPD by 2.65 (95% CI, 2.13–3.31) and chronic bronchitis by 2.89 (95% CI, 2.18–3.82) times more compared to non-biomass fuels. The risk of COPD was higher in Africa region (odds ratio [OR]: 3.19), Asia (OR: 2.88), South America (OR: 2.15), Europe (OR: 2.30) and North America (OR: 2.14). This distribution confirms that although the risk is higher in developing regions [16], the high risk is also present in developed countries, as some studies have shown [17, 18].

In some highly populated countries, like India and China, the exposure to biomass smoke is a significant risk factor for COPD, mainly in women living in rural zones [15, 19–24]. In some areas of India and China, this exposure is the most important risk factor of COPD [20–22, 25–30]. In Latin America, the PREPOCOL [3], the CRONICAS [31] and the PUMA [32] studies have confirmed that the use of biomass fuels, frequently wood, for cooking is a significant and independent risk factor for COPD, stronger in women from rural areas.

Although the risk of COPD from long-term indoor exposure to biomass fuels is particularly high in women [23, 33–36], a population study (n = 5539) showed that, after adjusting for age, smoking, educational level and occupational exposure, men exposed to wood smoke for more than 10 years had a higher risk of COPD (OR: 1.50) [37]. The risk of COPD increases significantly with the length of exposure to wood smoke and with simultaneous exposure to tobacco smoke [37].

This evidence supports that HAP from burning solid fuels, including biomass, is the biggest worldwide risk factor for COPD [38–40]. However, the prevalence of biomass-related COPD has not been precisely defined. The PREPOCOL study found a prevalence of 6.7% in people exposed to wood smoke and not to cigarette smoke compared to 7.8% in people exposed to cigarette smoke and not to wood smoke [37]. In rural Puno, Peru, daily use of biomass fuel for cooking among women was associated with COPD (prevalence ratio: 2.22, 95% CI: 1.02–4.81) and the population attributable risk of COPD due to daily exposure to biomass fuel smoke was 55% [31].

Some populational studies, however, found no association between exposure to biomass fuels and COPD [41, 42]. Most of the people evaluated in these studies lived near sea level, where cooking is usually done outdoors or with better ventilation. In contrast, many of the studies which document this association have included areas situated at high or intermediary altitudes, where, due to low temperatures, cooking is done all year round inside poorly ventilated homes as it occurs in winter in regions that have seasons. There is lack of standardization of questionnaires or other tools for evaluating the exposure to biomass smoke derived from cooking or heating. A recent study, from Kyrgyzstan, evaluated the prevalence of COPD associated with indoor contamination at different altitudes and found a higher prevalence of COPD at high altitude versus at low altitude (36.7% vs. 10.4%;  $p < 0.001$ ) associated with exposure to a greater indoor contamination at high altitude [43].

## 2. Differences between WS-COPD and TS-COPD

Although the risk of COPD has been proven for all types of biomass fuels, studies which best characterize COPD due to this type of exposure have focused on COPD caused by inhalation of wood smoke (WS-COPD). Therefore, this chapter also focuses on the differences of WS-COPD and TS-COPD.

**Core differences.** A growing body of evidence supports that WS-COPD, unlike TS-COPD, is predominantly and markedly a disease of the airways with mild or minimum emphysema [44–54]. Although recent publications have focused on the compromise of the small airways in biomass COPD [46, 50, 52], different from TS-COPD, in WS-COPD there is also a notorious compromise of the central airway due to carbon deposition (bronchial anthracofibrosis) with plaque formation and reduction in the caliber of the lobar bronchi [55–57]. In addition to this fundamental pathological difference between WS-COPD and TS-COPD, there are many other differences that we summarize in **Table 1** and review it below.

**Demographic differences.** Women and their children are the most exposed population to indoor air pollution from biomass fuels because of women are usually responsible for cooking meals, particularly in developing countries, spending several hours a day in frequently poor ventilated kitchens, and keeping their children close to them [4, 16, 23, 30, 58, 59].

Most of the studies show that women with WS-COPD are consistently shorter in height and have higher body mass index (BMI) than women with TS-COPD [37, 44, 49, 54, 60–66]. There is not a clear explanation for this difference. In general, women with WS-COPD were born and have lived in rural areas as their ancestry, while women with TS-COPD have lived in urban areas for many years and many of them have urban ancestry. Therefore, it is possible that some ethnic, nutritional and socioeconomics conditions could be part of the explanation of the difference, but there is not consistent information about this.

Moreover, women with WS-COPD are older, suggesting that patients with this type of exposure need more time to develop the disease or are diagnosed later [30, 37, 44, 51, 61, 63–65, 67].

**Clinical differences.** Several studies have shown a high frequency of respiratory symptoms (cough, expectoration, and dyspnea) and chronic bronchitis in subjects exposed to biomass smoke [9, 10, 14, 61]. However, comparative studies between WS-COPD and TS-COPD have found no consistent differences. Some studies show that symptoms, not only cough and phlegm but dyspnea, are more frequent or have more impact in WS-COPD than TS-COPD [48, 61, 68, 69] but other not [63, 65, 70]. Rhonchus and wheezing are more frequent in WS-COPD [68].

| Characteristics  | WS-COPD  | TS-COPD   |
|--|--|---|
| Demographic data   |  |   |
| Sex  | Predominantly women  | Predominantly men                                   |
| Age  | Higher   | Lower   |
| Height   | Lower  | Higher  |
| BMI  | Higher   | Lower   |
| Clinical characteristics   |  |   |
| Cough and expectoration  | Very common  | Common  |
| Chronic bronchitis   | Common   | Common  |
| Rhonchus and wheezing  | Common   | Less common   |
| Lung function tests  |  |   |
| PaCO <sub>2</sub>  | Higher (some studies)  | Less high   |
| PaO <sub>2</sub> and SaO <sub>2</sub>  | Lower  | Less low  |
| Obstruction (FEV <sub>1</sub> /FVC reduced)  | Mild   | More severe   |
| Reduced FEV <sub>1</sub>   | Lower  | Higher  |
| Bronchial hyperresponsiveness  | Higher   | Lower   |
| DLco and DLco/VA   | Normal or mildly reduced   | More reduced  |
| Radiography-tomography   |  |   |
| Emphysema  | Uncommon and mild  | Common and more severe                              |
| Bronchial thickening   | Common   | Less common   |
| Bronchiectasis   | Common   | Uncommon  |
| Atelectasis  | Common   | Uncommon  |
| Histology  |  |   |
| Emphysema  | Mild   | More severe   |
| Anthracosis  | Common   | Less common   |
| Airway fibrosis  | Common   | Less common   |
| Thickening of arteriole intima   | Common   | Less common   |
| Outcomes and clinical phenotypes   |  |   |
| Pulmonary hypertension   | More common  | Less common   |
| Quality of life  | Symptoms and activities more compromised or similar              | Similar or symptoms and activities less compromised |
| Survival   | Similar after adjusting for age.<br>Less after adjusting for age | Similar   |
| Exacerbator phenotype  | Similar  | Similar   |
| Asthma-COPD overlap phenotype  | More common  | Less common   |
| Emphysema phenotype  | Uncommon   | More common   |
| WS: wood smoke; TS: tobacco smoke; BMI: body mass index; DL <sub>CO</sub> : carbon monoxide diffusing capacity; FEV <sub>1</sub> : forced expiratory volume in 1 second; FVC: forced vital capacity; PaCO <sub>2</sub> : carbon dioxide arterial pressure; PaO <sub>2</sub> : oxygen arterial pressure; SaO <sub>2</sub> : oxygen saturation; VA: alveolar volume. |  |   |

**Table 1.**  
*Differences between wood smoke COPD and tobacco smoke COPD.*




The greater bronchial compromise in WS-COPD documented in several publications which use functional and tomographic evaluations supports the studies which show more frequent cough, expectoration, rhonchus and wheezing in WS-COPD.

**Differences in quality of life.** Using the Saint George's Hospital Questionnaire, Camp *et al.* found worse symptoms and more impaired activity indices in women with WS-COPD [48]. González-García *et al.*, in 138 women with COPD, showed that, at the same degree of obstruction, women with WS-COPD had a worse health status (poorer quality of life and worse dyspnea) than those with TS-COPD, without differences in comorbidities (**Figure 1**) [68]. Some studies have not shown differences in quality of life between these two groups of patients [65].

**Differences in lung function.** Airflow obstruction, both overall and adjusted by age, is milder [37, 48, 54, 60–63, 65] and the FEV<sub>1</sub> decline is slower and more homogeneous in WS-COPD [54, 62] than in TS-COPD. Ocakli *et al.* described a more significant compromise of FVC with higher FEV<sub>1</sub>/FVC ratio in people with COPD related to biomass smoke suggesting restrictive ventilatory alteration, but lung volumes were not measured [71]. The chronic airflow limitation in patients with WS-COPD is possibly due not only to the small airways compromise [46, 50, 52, 54], but to anthracofibrosis of the large airways [55–57].

A recent study, aimed to evaluate the lung volumes and the resistance and conductance of the airways using plethysmography, showed that residual volumes (RV),




**Figure 1.**  
*Quality of life. Comparison between WS-COPD and TS-COPD.* WS-COPD: wood-smoke COPD; TS-COPD: tobacco smoke COPD. SGRQ: Saint George Respiratory Questionnaire. S: Symptoms; A: Activity; I: Impact; T: Total. From: González-García M, *et al.* Arch Bronconeumol. 2014; 50 (ALAT congress): 59 (ref. 68).

total lung capacities (TLC) and RV/TLC ratios were significantly increased among both TS-COPD and non-smoking COPD (including biomass smoke COPD) subjects compared to healthy subjects ( $p < 0.0001$ ), with no differences between the two COPD groups [54]. The same study showed that patients with COPD related to biomass smoke had significantly higher airway resistance (sRaw values) than TS-COPD patients ( $p = 0.005$ ) and significantly lower conductance (sGaw values) in biomass COPD than in TS-COPD ( $p = 0.010$ ) [54].


Some studies have showed that carbon dioxide arterial pressure ( $\text{PaCO}_2$ ) is higher (lower ventilation) and oxygen arterial pressure ( $\text{PaO}_2$ ) and oxygen hemoglobin saturation ( $\text{SaO}_2$ ) are lower in WS-COPD than in TS-COPD [48, 61, 62, 65, 69]. Interestingly, Olloquequi *et al.* described that COPD patients who have had mixed exposure to tobacco smoke as well as biomass smoke had lower oxygen saturation than those who were exposed only to cigarette smoke or biomass smoke [72]. The lower oxygenation rates observed in WS-COPD may be explained by the compromise of the small airways and/or by hypoventilation. It remains to be determined if the higher BMI in these patients, most of whom are women over 50 years of age, is involved in this behavior.

One of most consistent differences between WS-COPD and TS-COPD is the significantly lower compromise of the diffusion capacity ( $\text{DL}_{\text{CO}}$ ) in WS-COPD.  $\text{DL}_{\text{CO}}$  and  $\text{DL}_{\text{CO}}$ /alveolar volume ( $\text{DL}_{\text{CO}}/\text{VA}$ ) ratio are normal or mildly altered in WS-COPD patients compared to TS-COPD patients, in who these parameters are significantly reduced [49, 60, 61], and occurs at all levels of COPD severity (**Figure 2A and B**) [61]. This finding correlates well with the lower grade of emphysema found on computed tomography (CT) in patients with WS-COPD in comparison with TS-COPD [48, 49, 54, 73]. The mildly reduced  $\text{DL}_{\text{CO}}$  with normal  $\text{DL}_{\text{CO}}/\text{VA}$  found in women with WS-COPD has been described in patients with significantly compromised small airways with little emphysema (pseudophysiological emphysema) [74]. The correlation between the level of decrease of  $\text{FEV}_1$  and the level of  $\text{DL}_{\text{CO}}$  reduction is significantly better in women with TS-COPD than in those with WS-COPD, highlighting the greater contribution of emphysema to airflow obstruction in TS-COPD (**Figure 3**) [61].

**Differences in bronchial hyperresponsiveness.** Women with WS-COPD have greater bronchial hyperresponsiveness than women with TS-COPD (**Figure 4**) [64].



**Figure 2.**  
**Comparison of diffusion capacity between WS-COPD and TS-COPD according to degree of obstruction.** A.  $\text{DL}_{\text{CO}}$  (%) and B.  $\text{DL}_{\text{CO}}/\text{VA}$  (%). In TS-COPD,  $\text{DL}_{\text{CO}}$  and  $\text{DL}_{\text{CO}}/\text{VA}$  are more significantly compromised than in WS-COPD.  $\text{DL}_{\text{CO}}/\text{VA}$  is normal in WS-COPD at all levels of severity. WS: wood smoke; TS: tobacco smoke;  $\text{DL}_{\text{CO}}$ : carbon monoxide diffusing capacity; VA: alveolar volume. From: González-García M, *et al.* Acta Med Colomb. 2004; 29: 17–25 (Ref. 61).



**Figure 3.**  
**Correlation between  $FEV_1$  (%) and  $DL_{CO}$  by exposure.** Greater correlation is observed between  $FEV_1$  and  $DL_{CO}$  in TS-COPD ( $P < .001$ ,  $r = 0.599$ ) than in WS-COPD ( $P = .014$ ,  $r = 0.320$ ). WS: Wood smoke; TS: Tobacco smoke;  $DL_{CO}$ : carbon monoxide diffusing capacity;  $FEV_1$ : forced expiratory volume in 1 s. From: González-García M, et al. *Acta Med Colomb.* 2004; 29: 17–25 (ref. 61).

This finding could be correlated to the higher frequency of the asthma-COPD overlap phenotype described in biomass-related COPD [63].


**Differences in exercise tolerance.** Some studies which included the 6-minute walking test found no significant differences in distances walked between patients with WS-COPD and TS-COPD [48, 65, 68]. However, the study carried out by Zhao *et al.* found that patients exposed to biomass smoke walked fewer meters in the 6-minute walk than those with TS-COPD [46].

**Tomography and histological differences.** Patients with WS-COPD have consistently less emphysema and more airway changes (bronchial and peribronchial thickening and fibrosis, bronchiectasis, segmental and laminar subsegmental atelectasis, mosaic perfusion pattern, parenchymal bands) than patients with TS-COPD on both chest tomography and histological studies [48–50, 52, 60, 69, 73] (**Figure 5**). Bronchial and lung biopsies from patients with WS-COPD show significant inflammation and thickening of the bronchial wall, mainly of its basal membrane, squamous cell metaplasia with a remarkable anthracotic pigment deposition in the bronchi and pulmonary interstitium [60, 75–77].

**Differences in the distribution of clinical phenotypes.** Golpe *et al.* evaluated the frequency of clinical phenotypes defined by the 2014 Spanish COPD guidelines [78] in patients with COPD caused by biomass or tobacco smoke. They found a greater frequency of emphysema phenotype in TS-COPD and a more common asthma-COPD overlap phenotype in biomass COPD, but the difference disappeared after adjusting for sex [63]. These differences fit perfectly with the findings of the studies reported so far in this chapter. No difference was found in the frequencies of chronic bronchitis or exacerbator phenotypes [63].

**Differences in pulmonary hypertension.** Sertogullarindan *et al.* found that pulmonary hypertension (PH) on echocardiography was more common in WS-COPD than in TS-COPD patients [79]. In previous studies, our group, based






**Figure 4.**  
*Bronchial hyperresponsiveness evaluated by PC<sub>20</sub> by exposure.* PC<sub>20</sub>: methacholine concentration causing ≥ 20% reduction in FEV<sub>1</sub>. White circles: WS-COPD; black circles: TS-COPD. PC<sub>20</sub> geometric mean: WS-COPD versus TS-COPD: 0.39 (0.06–5.13) versus 1.24 (0.34–9.39),  $P = .028$ . WS: wood smoke; TS: tobacco smoke. From: Gonzalez-Garcia M, et al. *Int J Chron Obstruct Pulmon Dis.* 2012; 7: 367–373 (Ref. 64).

on radiographic evaluation, suggested the same higher frequency of PH in patients with severe WS-COPD [61], and Sandoval *et al.* showed a high rate of PH and cor pulmonale among patients with long-term domestic exposure to wood smoke [80]. It has been posed that the PH observed in WS-COPD could be associated not only to airflow obstruction and hypoxemia, but to a direct inflammatory effect of the inhaled smoke. It has been described that mice exposed to biomass exhibited more perivascular inflammation than those exposed to cigarette smoke [81].

**Differences in the frequency of bronchial anthracofibrosis.** The incidence of bronchial anthracofibrosis and its severity in individuals exposed to wood smoke or tobacco smoke has not been evaluated in prospective studies, and no differences are known. However, marked anthracofibrosis is commonly encountered in the airway of subjects exposed to wood smoke [16, 55–57], sometimes accompanied by bronchial stenosis, and it seems more frequent and severe than in tobacco smoke exposed. A significant proportion of these patients have chronic airflow limitation and meet the functional criteria of COPD [55] possibly due to the small airways compromise aggravated by anthracofibrotic central airway stenosis. It is currently impossible to ascertain if bronchial anthracofibrosis is another feature of WS-COPD that appears more commonly and in a more severe form than in TS-COPD, or if it is a specific entity accompanied by obstruction.

**Differences in survival and frequency of exacerbations.** After adjusting for age, sex, and disease severity, no differences have been found in survival between WS-COPD and TS-COPD [65, 82].

Recently, Cho *et al.*, among 1033 patients with COPD, have shown that patients with COPD associated with biomass smoke and those with COPD associated with tobacco smoke had a similar risk of exacerbations [67], confirming previous observations [63].



**Figure 5.** Tomographic differences between smoking and non-smoking COPD. Differences between smoker COPD (S-COPD) ( $n = 38$ ) and non-smoker COPD (NS-COPD) ( $n = 70$ ) on inspiratory-expiratory high resolution computerized tomography (HRCT) imaging (A) Showing HRCT classification of smoker and non-smoker COPD, including airway disease (black bars), emphysema (open bars) and interstitial lung abnormality (ILA) predominance (gray bars); (B) dot plot for emphysema and decreased attenuation of expiratory CT in S-COPD (■) and NS-COPD (Δ) compared to healthy subjects (●), where bars indicate median and interquartile ranges and  $* p < 0.05$ , (C) Representative HRCT images of Smoker-COPD showing extensive centrilobular emphysema in the upper lobes and non-smoker COPD showing generalized decreased attenuation and some bronchial wall thickening. From: Salvi SS, et al. phenotypic comparison between smoking and non-smoking chronic obstructive pulmonary disease. *Respiratory Research*. 2020; 21: 50 [64].

**Differences in inflammatory profile and pathophysiological ways.** It is expected that the greater airway inflammatory involvement and the lower rate of emphysema in biomass COPD, including WS-COPD, compared to TS-COPD have an etiological, pathogenic and physiopathological basis. However, although there is a growing information about the pathogenic mechanisms in COPD due to biomass smoke exposure [83], it is not clear what are the reasons explaining its differences

with TS-COPD. Some studies have focused on looking for differences in the type of inflammation and the proteolytic activity, and recently the gene expression.

The exposure to biomass smoke induces pulmonary macrophages and mononuclear and polynuclear cells to generate numerous inflammatory mediators, including interleukin-6 (IL-6), interleukin-8 (IL-8), monocyte chemoattractant protein 1 (MCP-1), macrophage inflammatory protein 2 (MIP2) and tumoral necrosis factor (TNF) [83, 84]. These can generate a second wave of mediators that include enzymes, such as matrix metalloproteinase 9 (MMP-9) and matrix metalloproteinase 12 (MMP-12) involved in proteolysis and tissue remodeling typical of COPD. A recent study explored differences in chemokine and cytokine concentrations among biomass-COPD versus TS-COPD and exposed controls without COPD. The author identified CCL27 and CXCL13 as putative, plausibly homeostatic/protective biomarkers for biomass COPD [85].


Golpe *et al.* found that serum IL-6, IL-8, IL-5 were significantly higher in TS-COPD patients than in biomass COPD without differences in serum IL-13, periostin, surfactant protein-P, TNF- $\alpha$ , IgE, erythrocyte sedimentation rate, C-reactive protein and fibrinogen [86]. The level of exhaled nitric oxide (FeNO) was higher in biomass COPD ( $39.0 \pm 14.6$  ppb) than in TS-COPD ( $27.6 \pm 16.3$  ppb); although the difference did not reach the statistical significance level, it was borderline ( $p: 0.056$ ) and it could be related to a small sample size [86].

A study by Solleiro-Villavicencio *et al.*, done in women with COPD and healthy controls, found that IL-4 and T<sub>H</sub>2 cells were significantly higher in biomass COPD than in TS-COPD [87]. Frequency of T<sub>H</sub>17 cells in patients with TS-COPD was significantly higher than in patients with biomass COPD. They suggested that a T<sub>H</sub>2 cytokine inflammatory profile could predominate in biomass COPD [87]. Although the majority the authors have not found differences in blood eosinophils counts between biomass COPD and TS-COPD, Fernandes *et al.*, using a cutoff of  $\geq 3\%$ , found more frequent sputum eosinophilia in biomass COPD than in TS-COPD [88]. In the same way of the responses T<sub>H</sub>2, Olloquequi *et al.* found higher levels of total IgE in patients with biomass smoke COPD than in TS-COPD (**Figure 6**) [72].

It seems clear that the development and clinical course of COPD depend on an interaction between genetic and environmental factors. The gene regulation and expression are fundamentally involved in the pathophysiology of COPD and it is known that microRNAs (miRNAs) participate in the control of post-transcriptional regulation in TS-COPD. Recently, Velasco-Torres *et al.* have described the differential role of miR-34a (downregulated) [89] and of the axis miR-22 - histone deacetylase activity (HDAC4) - IL-17 [90]. This axis has been linked to the development of emphysema in rats. Serum miR-22-3p was downregulated in biomass COPD-Bs relative to COPD-TS. In contrast, the concentration of HDAC4 was higher in biomass and exhibited a significant positive correlation with DL<sub>CO</sub>% [90]. This mechanism could be involved in the lower expression of emphysema in WS-COPD.

In summary, inflammation in biomass COPD, including WS-COPD, could be different from that in TS-COPD with a possible predominance of T<sub>H</sub>2 profile, and the lower generation of emphysema could be related to a particular and different response to biomass smoke.

**Differences in therapeutic interventions.** Almost all the studies on pharmacological interventions in COPD supporting clinical guidelines and strategies [91] have been done in developed countries and use the exposure to tobacco smoke as inclusion criteria. So, the derived evidence could not be extrapolated to other causes of COPD. Taking into account the different type of inflammation, higher bronchial hyperresponsiveness and higher frequency of the asthma-COPD overlap phenotype, it is expected a benefit role of inhaled corticosteroids. To our best knowledge, there is only one study in a small sample of patients that showed better results in



**Figure 6.**  
*IgE levels in COPD due to tobacco smoke, wood smoke or both.* α: Different from control subjects ( $p < 0.05$ ). β: Different from TS COPD group ( $p < 0.05$ ). TS: Tobacco smoke; WS: Wood smoke. From: Olloquequi J, et al. *Respiratory Research* (2018) 19:13 [72].

reducing the exacerbation frequency and improving lung function in patients with WS-COPD with the use of ICS [92]. So, there is a lack of information regarding the efficacy and safety of pharmacological interventions in patients with WS-COPD, particularly on the potential benefits of the use of ICS in these patients.

### 3. COPD related to biomass fuels different from wood smoke

Biomass fuels used for cooking include mainly wood, charcoal, agricultural residues, and animal dung. The composition of these types of biomass and the smoke derived from burning it significantly differ [93]. There is growing information about the different animal and human responses to the exposures to different kind of the biomass fuels. As we have presented in this chapter, probably the most studied biomass smoke and the responses to its inhalation is wood [94, 95]. Animal manure contains greater diversity and greater quantities of microorganisms and the inflammatory response could be different [81, 96, 97]. Dung biomass smoke [96] activates inflammatory responses in human epithelial cells from airways. Cow dung exposure, but not wood smoke exposure, mediated a measurable increase in non-tipeable *Haemophilus influenzae* adhesion to airway epithelial cells.

Most of the epidemiologic, systematic reviews and meta-analysis group these types of fuels as the generic term “biomass” studies. The distribution of the types of biomass fuels used for cooking differ depending on the country and region. In certain regions, wood or wood and charcoal are the only or most used fuel but in other ones it could predominate animal dung or agricultural residues. One study in Tanzania shows that 99.5% of participants had exposure to biomass: 92.4% used wood, 14.9% used charcoal, 1.5% used crop residues and 0.6% used animal dung for cooking and heating purposes [98].

Studies that have characterized COPD, this means that have described the clinical, functional, radiographic and histopathological characteristics, have been done in wood smoke exposed, and those that have grouped patients under “biomass”



exposed or “non-smokers” have included mainly people exposed to wood smoke. Therefore, it is possible that the COPD related to dung or crop residues be different and it is not recommendable to generalize the observations done in WS-COPD and described in this chapter to other types of biomass fuels.

#### **4. Respiratory disease due to indoor chronic exposure to wood smoke: a different phenotype of COPD or a separable disease?**

COPD is defined using a functional criterion and an unspecific exposure. So, under this term can be included a very numerous and heterogenous pathologic conditions. Accepting the weakness of the definition of COPD as a disease (it is more a syndrome), the WS-COPD (it is not sure if all type of biomass COPD) could be accepted as a phenotype of COPD. However, if we assume that etiology is different (wood smoke is not cigarette smoke), the inflammatory responses and the pathophysiologic could be different, and the clinical, functional and histopathologic expression are also different, the chronic respiratory disease due to long-term indoor exposure to wood smoke is better understood as a separate nosology entity. Most importantly the actions for prevention significantly differ and the therapeutic interventions could be also different.

#### **Author details**


Carlos A. Torres-Duque<sup>1,2\*</sup>, Felipe Severiche-Bueno<sup>1,2</sup>  
and Mauricio González-García<sup>1,2</sup>

1 Fundación Neumológica Colombiana, Bogotá, Colombia

2 La Sabana University (Universidad de La Sabana), Bogotá, Colombia

\*Address all correspondence to: [ctorres@neumologica.org](mailto:ctorres@neumologica.org)

#### **IntechOpen**

© 2021 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

## References

- [1] Bonjour S, Adair-Rohani H, Wolf J, Bruce NG, Mehta S, Pruss-Ustun A, Lahiff M, Rehfuess EA, Mishra V, Smith KR. Solid fuel use for household cooking: country and regional estimates for 1980-2010. *Environmental Health Perspectives* 2013; 121: 784-790.
- [2] WHO. World Health Organization. Household air pollution and health. 2018. <https://www.who.int/news-room/fact-sheets/detail/household-air-pollution-and-health>. (Accessed on: December 13, 2020). 2018.
- [3] Caballero A, Torres-Duque CA, Jaramillo C, Bolivar F, Sanabria F, Osorio P, Orduz C, Guevara DP, Maldonado D. Prevalence of COPD in five Colombian cities situated at low, medium, and high altitude (PREPOCOL study). *Chest* 2008; 133: 343-349.
- [4] Amegah AK, Jaakkola JJK. Household air pollution and the sustainable development goals. *Bulletin of the World Health Organization* 2016; 94: 215-221.
- [5] Smith KR, Bruce N, Balakrishnan K, Adair-Rohani H, Balmes J, Chafe Z, Dherani M, Hosgood HD, Mehta S, Pope D, Rehfuess E. Millions dead: how do we know and what does it mean? Methods used in the comparative risk assessment of household air pollution. *Annual Review of Public Health* 2014; 35: 185-206.
- [6] Lee KK, Bing R, Kiang J, Bashir S, Spath N, Stelzle D, Mortimer K, Bularga A, Doudehis D, Joshi SS, Strachan F, Gummy S, Adair-Rohani H, Attia EF, Chung MH, Miller MR, Newby DE, Mills NL, McAllister DA, Shah ASV. Adverse health effects associated with household air pollution: a systematic review, meta-analysis, and burden estimation study. *The Lancet Global Health* 2020; 8: e1427-e1434.
- [7] Pathak U, Gupta NC, Suri JC. Risk of COPD due to indoor air pollution from biomass cooking fuel: a systematic review and meta-analysis. *International Journal of Environmental Health Research* 2019; 1-14.
- [8] Hu G, Zhou Y, Tian J, Yao W, Li J, Li B, Ran P. Risk of COPD from exposure to biomass smoke: a metaanalysis. *Chest* 2010; 138: 20-31.
- [9] Po JY, FitzGerald JM, Carlsten C. Respiratory disease associated with solid biomass fuel exposure in rural women and children: systematic review and meta-analysis. *Thorax* 2011; 66: 232-239.
- [10] Kurmi OP, Semple S, Simkhada P, Smith WC, Ayres JG. COPD and chronic bronchitis risk of indoor air pollution from solid fuel: a systematic review and meta-analysis. *Thorax* 2010; 65: 221-228.
- [11] Torres-Duque C, Maldonado D, Pérez-Padilla R, Ezzati M, Viegi G. Biomass fuels and respiratory diseases: a review of the evidence. *Proceedings of the American Thoracic Society* 2008; 5: 577-590.
- [12] Gordon SB, Bruce NG, Grigg J, Hibberd PL, Kurmi OP, Lam KB, Mortimer K, Asante KP, Balakrishnan K, Balmes J, Bar-Zeev N, Bates MN, Breyse PN, Buist S, Chen Z, Havens D, Jack D, Jindal S, Kan H, Mehta S, Moschovis P, Naeher L, Patel A, Perez-Padilla R, Pope D, Rylance J, Semple S, Martin WJ, 2nd. Respiratory risks from household air pollution in low and middle income countries. *The Lancet Respiratory Medicine* 2014; 2: 823-860.
- [13] Capistrano SJ, van Reyk D, Chen H, Oliver BG. Evidence of Biomass Smoke Exposure as a Causative Factor for the Development of COPD. *Toxics* 2017; 5.
- [14] Gonzalez-Garcia M, Caballero A, Jaramillo C, Torres-Duque CA. Chronic

bronchitis: High prevalence in never smokers and underdiagnosis- A population-based study in Colombia. *Chronic Respiratory Disease* 2018; 1479972318769771.

[15] Panigrahi A, Padhi BK. Chronic bronchitis and airflow obstruction is associated with household cooking fuel use among never-smoking women: a community-based cross-sectional study in Odisha, India. *BMC Public Health* 2018; 18: 924.

[16] Siddharthan T, Grigsby MR, Goodman D, Chowdhury M, Rubinstein A, Irazola V, Gutierrez L, Miranda JJ, Bernabe-Ortiz A, Alam D, Kirenga B, Jones R, van Gemert F, Wise RA, Checkley W. Association between Household Air Pollution Exposure and Chronic Obstructive Pulmonary Disease Outcomes in 13 Low- and Middle-Income Country Settings. *American Journal of Respiratory and Critical Care Medicine* 2018; 197: 611-620.

[17] Orozco-Levi M, Garcia-Aymerich J, Villar J, Ramírez-Sarmiento A, Antó JM, Gea J. Wood smoke exposure and risk of chronic obstructive pulmonary disease. *The European Respiratory Journal* 2006; 27: 542-546.

[18] Raju S, Keet CA, Paulin LM, Matsui EC, Peng RD, Hansel NN, McCormack MC. Rural Residence and Poverty Are Independent Risk Factors for Chronic Obstructive Pulmonary Disease in the United States. *American Journal of Respiratory and Critical Care Medicine* 2019; 199: 961-969.

[19] Kalkana T, Moitra S, Jindal SK, Moitra S. Increasing burden of COPD in rural India: an example why India warrants primary healthcare reforms. *ERJ Open Research* 2016; 2: 00032-02016.

[20] McKay AJ, Mahesh PA, Fordham JZ, Majeed A. Prevalence of COPD in

India: a systematic review. *Primary Care Respiratory Journal : Journal of the General Practice Airways Group* 2012; 21: 313-321.

[21] KalagoudaMahishale V, Angadi N, Metgudmath V, Lolly M, Eti A, Khan S. The Prevalence of Chronic Obstructive Pulmonary Disease and the Determinants of Underdiagnosis in Women Exposed to Biomass Fuel in India- a Cross Section Study. *Chonnam Medical Journal* 2016; 52: 117-122.

[22] Liu S, Zhou Y, Wang X, Wang D, Lu J, Zheng J, Zhong N, Ran P. Biomass fuels are the probable risk factor for chronic obstructive pulmonary disease in rural South China. *Thorax* 2007; 62: 889-897.

[23] Sana A, Somda SMA, Meda N, Bouland C. Chronic obstructive pulmonary disease associated with biomass fuel use in women: a systematic review and meta-analysis. *BMJ Open Respiratory Research* 2018; 5.

[24] Smith M, Li L, Augustyn M, Kurmi O, Chen J, Collins R, Guo Y, Han Y, Qin J, Xu G, Wang J, Bian Z, Zhou G, Peto R, Chen Z. Prevalence and correlates of airflow obstruction in ~317 000 never-smokers in China. 2014; 44: 66-77.

[25] Wang C, Xu J, Yang L, Xu Y, Zhang X, Bai C, Kang J, Ran P, Shen H, Wen F, Huang K, Yao W, Sun T, Shan G, Yang T, Lin Y, Wu S, Zhu J, Wang R, Shi Z, Zhao J, Ye X, Song Y, Wang Q, Zhou Y, Ding L, Yang T, Chen Y, Guo Y, Xiao F, Lu Y, Peng X, Zhang B, Xiao D, Chen CS, Wang Z, Zhang H, Bu X, Zhang X, An L, Zhang S, Cao Z, Zhan Q, Yang Y, Cao B, Dai H, Liang L, He J. Prevalence and risk factors of chronic obstructive pulmonary disease in China (the China Pulmonary Health [CPH] study): a national cross-sectional study. *Lancet (London, England)* 2018; 391: 1706-1717.

- [26] Yang Y, Mao J, Ye Z, Li J, Zhao H, Liu Y. Risk factors of chronic obstructive pulmonary disease among adults in Chinese mainland: A systematic review and meta-analysis. *Respiratory Medicine* 2017; 131: 158-165.
- [27] Zhang JJ, Smith KR. Household air pollution from coal and biomass fuels in China: measurements, health impacts, and interventions. *Environmental Health Perspectives* 2007; 115: 848-855.
- [28] Zhu B, Wang Y, Ming J, Chen W, Zhang L. Disease burden of COPD in China: a systematic review. *International Journal of Chronic Obstructive Pulmonary Disease* 2018; 13: 1353-1364.
- [29] The burden of chronic respiratory diseases and their heterogeneity across the states of India: the Global Burden of Disease Study 1990-2016. *The Lancet Global Health* 2018; 6: e1363-e1374.
- [30] Chan KH, Kurmi OP, Bennett DA, Yang L, Chen Y, Tan Y, Pei P, Zhong X, Chen J, Zhang J, Kan H, Peto R, Lam KBH, Chen Z. Solid Fuel Use and Risks of Respiratory Diseases. A Cohort Study of 280,000 Chinese Never-Smokers. *American Journal of Respiratory and Critical Care Medicine* 2019; 199: 352-361.
- [31] Jaganath D, Miranda JJ, Gilman RH, Wise RA, Diette GB, Miele CH, Bernabe-Ortiz A, Checkley W, Group CCS. Prevalence of chronic obstructive pulmonary disease and variation in risk factors across four geographically diverse resource-limited settings in Peru. *Respiratory Research* 2015; 16: 40-40.
- [32] Montes de Oca M, Zabert G, Moreno D, Laucho-Contreras ME, Lopez Varela MV, Surmont F. Smoke, Biomass Exposure, and COPD Risk in the Primary Care Setting: The PUMA Study. *Respiratory Care* 2017; 62: 1058-1066.
- [33] Perez-Padilla R, Regalado J, Vedral S, Pare P, Chapela R, Sansores R, Selman M. Exposure to biomass smoke and chronic airway disease in Mexican women. A case-control study. *American Journal of Respiratory and Critical Care Medicine* 1996; 154: 701-706.
- [34] Regalado J, Perez-Padilla R, Sansores R, Paramo Ramirez JI, Brauer M, Pare P, Vedral S. The effect of biomass burning on respiratory symptoms and lung function in rural Mexican women. *American Journal of Respiratory and Critical Care Medicine* 2006; 174: 901-905.
- [35] Dennis RJ, Maldonado D, Norman S, Baena E, Martinez G. Woodsmoke exposure and risk for obstructive airways disease among women. *Chest* 1996; 109: 115-119.
- [36] Kiraz K, Kart L, Demir R, Oymak S, Gulmez I, Unalacak M, Ozesmi M. Chronic pulmonary disease in rural women exposed to biomass fumes. *Clinical and Investigative Medicine Medecine Clinique et Experimentale* 2003; 26: 243-248.
- [37] Torres-Duque C, Caballero A, González-García M, Jaramillo C, Maldonado D. Chronic obstructive pulmonary disease in people exposed to Wood smoke. PREPOCOL: a population based study. *Am J Respir Crit Care Med* 2013; 187: A364.
- [38] Salvi S, Barnes PJ. Is exposure to biomass smoke the biggest risk factor for COPD globally? *Chest* 2010; 138: 3-6.
- [39] Mortimer K, Gordon SB, Jindal SK, Accinelli RA, Balmes J, Martin WJ, 2nd. Household air pollution is a major avoidable risk factor for cardiorespiratory disease. *Chest* 2012; 142: 1308-1315.
- [40] Bellou V, Belbasis L, Konstantinidis AK, Evangelou E. Elucidating the risk factors



for chronic obstructive pulmonary disease: an umbrella review of meta-analyses. *The International Journal of Tuberculosis and Lung Disease : The Official Journal of the International Union against Tuberculosis and Lung Disease* 2019; 23: 58-66.

[41] Menezes AM, Perez-Padilla R, Jardim JR, Muiño A, Lopez MV, Valdivia G, Montes de Oca M, Talamo C, Hallal PC, Victora CG. Chronic obstructive pulmonary disease in five Latin American cities (the PLATINO study): a prevalence study. *Lancet (London, England)* 2005; 366: 1875-1881.

[42] Lamprecht B, McBurnie MA, Vollmer WM, Gudmundsson G, Welte T, Nizankowska-Mogilnicka E, Studnicka M, Bateman E, Anto JM, Burney P, Mannino DM, Buist SA. COPD in never smokers: results from the population-based burden of obstructive lung disease study. *Chest* 2011; 139: 752-763.

[43] Brakema EA, Tabyshova A, Kasteleyn MJ, Molendijk E, van der Kleij R, van Boven JFM, Emilov B, Akmatolieva M, Mademilov M, Numans ME, Williams S, Sooronbaev T, Chavannes NH. High COPD prevalence at high altitude: does household air pollution play a role? *The European Respiratory Journal* 2019; 53.

[44] Pérez-Padilla R, Ramirez-Venegas A, Sansores-Martinez R. Clinical Characteristics of Patients With Biomass Smoke-Associated COPD and Chronic Bronchitis, 2004-2014. *Chronic Obstructive Pulmonary Diseases: Journal of the COPD Foundation* 2014; 1: 23-32.

[45] Torres-Duque CA, Garcia-Rodriguez MC, Gonzalez-Garcia M. Is Chronic Obstructive Pulmonary Disease Caused by Wood Smoke a Different Phenotype or a Different Entity? *Archivos de Bronconeumologia* 2016; 52: 425-431.

[46] Zhao D, Zhou Y, Jiang C, Zhao Z, He F, Ran P. Small airway disease: A different phenotype of early stage COPD associated with biomass smoke exposure. *Respirology (Carlton, Vic)* 2018; 23: 198-205.

[47] Perret JL, Abramson MJ. Biomass smoke COPD: A phenotype or a different disease? *Respirology (Carlton, Vic)* 2018; 23: 124-125.

[48] Camp PG, Ramirez-Venegas A, Sansores RH, Alva LF, McDougall JE, Sin DD, Pare PD, Muller NL, Silva CI, Rojas CE, Coxson HO. COPD phenotypes in biomass smoke- versus tobacco smoke-exposed Mexican women. *The European Respiratory Journal* 2014; 43: 725-734.

[49] Gonzalez-Garcia M, Maldonado Gomez D, Torres-Duque CA, Barrero M, Jaramillo Villegas C, Perez JM, Varon H. Tomographic and functional findings in severe COPD: comparison between the wood smoke-related and smoking-related disease. *Jornal Brasileiro de Pneumologia : Publicacao Oficial da Sociedade Brasileira de Pneumologia e Tisiologia* 2013; 39: 147-154.

[50] Fernandes L, Gulati N, Fernandes Y, Mesquita AM, Sardesai M, Lammers JJ, Mohamed Hoesein FA, Ten Hacken NHT, van den Berge M, Galban CJ, Siddiqui S. Small airway imaging phenotypes in biomass- and tobacco smoke-exposed patients with COPD. *ERJ Open Res* 2017; 3.

[51] Assad NA, Balmes J, Mehta S, Cheema U, Sood A. Chronic obstructive pulmonary disease secondary to household air pollution. *Seminars in Respiratory and Critical Care Medicine* 2015; 36: 408-421.

[52] Ramirez-Venegas A, Torres-Duque CA, Guzman-Bouilloud NE, Gonzalez-Garcia M, Sansores RH. Small airway disease in COPD associated to biomass exposure. *Revista de*

*Investigacion Clinica; Organo del Hospital de Enfermedades de la Nutricion* 2019; 71: 70-78.

[53] Jindal S, Jindal A. COPD in Biomass exposed nonsmokers: a different phenotype. *Expert Review of Respiratory Medicine* 2020; 1-8.

[54] Salvi SS, Brashier BB, Londhe J, Pyasi K, Vincent V, Kajale SS, Tambe S, Mandani K, Nair A, Mak SM, Madas S, Juvekar S, Donnelly LE, Barnes PJ. Phenotypic comparison between smoking and non-smoking chronic obstructive pulmonary disease. *Respiratory Research* 2020; 21: 50.

[55] Kim YJ, Jung CY, Shin HW, Lee BK. Biomass smoke induced bronchial anthracofibrosis: presenting features and clinical course. *Respiratory Medicine* 2009; 103: 757-765.

[56] Gupta A, Shah A. Bronchial anthracofibrosis: an emerging pulmonary disease due to biomass fuel exposure. *The International Journal of Tuberculosis and Lung Disease : The Official Journal of the International Union against Tuberculosis and Lung Disease* 2011; 15: 602-612.

[57] Kim H, Cha SI, Shin KM, Lim JK, Oh S, Kim MJ, Lee YD, Kim M, Lee J, Kim CH. Clinical relevance of bronchial anthracofibrosis in patients with chronic obstructive pulmonary disease exacerbation. *Tuberculosis and Respiratory Diseases* 2014; 77: 124-131.

[58] Duan JX, Cheng W, Zeng YQ, Chen Y, Cai S, Li X, Zhu YQ, Chen M, Zhou ML, Ma LB, Liu QM, Chen P. Characteristics of Patients with Chronic Obstructive Pulmonary Disease Exposed to Different Environmental Risk Factors: A Large Cross-Sectional Study. *Int J Chron Obstruct Pulmon Dis* 2020; 15: 2857-2867.

[59] Moreira M, Barbosa M, Jardim J, Queiroz M, Inácio L. Doença pulmonar

obstrutiva crônica em mulheres expostas à fumaça de fogão à lenha. . *Revista da Associação Médica Brasileira* 2013; 59: 607-613.

[60] Moran-Mendoza O, Perez-Padilla JR, Salazar-Flores M, Vazquez-Alfaro F. Wood smoke-associated lung disease: a clinical, functional, radiological and pathological description. *The International Journal of Tuberculosis and Lung Disease : The Official Journal of the International Union against Tuberculosis and Lung Disease* 2008; 12: 1092-1098.

[61] González M, Páez S, Jaramillo C, Barrero M, Maldonado D. Enfermedad pulmonar obstructiva crónica (EPOC) por humo de leña en mujeres. *Acta Med Colomb* 2004; 29: 17-25.

[62] Ramirez-Venegas A, Sansores RH, Quintana-Carrillo RH, Velazquez-Uncal M, Hernandez-Zenteno RJ, Sanchez-Romero C, Velazquez-Montero A, Flores-Trujillo F. FEV1 decline in patients with chronic obstructive pulmonary disease associated with biomass exposure. *American Journal of Respiratory and Critical Care Medicine* 2014; 190: 996-1002.

[63] Golpe R, Sanjuan Lopez P, Cano Jimenez E, Castro Anon O, Perez de Llano LA. Distribution of clinical phenotypes in patients with chronic obstructive pulmonary disease caused by biomass and tobacco smoke. *Archivos de Bronconeumologia* 2014; 50: 318-324.

[64] Gonzalez-Garcia M, Torres-Duque CA, Bustos A, Jaramillo C, Maldonado D. Bronchial hyperresponsiveness in women with chronic obstructive pulmonary disease related to wood smoke. *Int J Chron Obstruct Pulmon Dis* 2012; 7: 367-373.

[65] Ramirez-Venegas A, Sansores RH, Perez-Padilla R, Regalado J, Velazquez A, Sanchez C, Mayar ME. Survival of patients with chronic

obstructive pulmonary disease due to biomass smoke and tobacco. *American Journal of Respiratory and Critical Care Medicine* 2006; 173: 393-397.

[66] Pérez-Bautista O, Montaña M, Pérez-Padilla R, Zúñiga-Ramos J, Camacho-Priego M, Barrientos-Gutiérrez T, Buendía-Roldan I, Velasco-Torres Y, Ramos C. Women with COPD by biomass show different serum profile of adipokines, incretins, and peptide hormones than smokers. *RESPIRATORY RESEARCH* 2018; 19: 239.

[67] Cho J, Lee C-H, Hwang S-s, Kim KU, Lee SH, Park HY, Park SJ, Min KH, Oh Y-M, Yoo KH, Jung K-S, on behalf of the K, Investigators K. Risk of acute exacerbations in chronic obstructive pulmonary disease associated with biomass smoke compared with tobacco smoke. *BMC Pulmonary Medicine* 2019; 19: 68.

[68] González-García M GV, Perlaza I, Casas A. Diferencias en el impacto sobre el estado de salud entre la EPOC por cigarrillo y por humo de leña. *Archivos de Bronconeumologia* 2014; 50 (Numero Especial Congreso ALAT): 59.

[69] Meneghini AC, Koenigkam-Santos M, Pereira MC, Tonidandel PR, Terra-Filho J, Cunha FQ, Menezes MBd, Vianna EO. Biomass smoke COPD has less tomographic abnormalities but worse hypoxemia compared with tobacco COPD %J Brazilian Journal of Medical and Biological Research. 2019; 52.

[70] Moreira MAC, Moraes MRd, Silva DGST, Pinheiro TF, Vasconcelos Júnior HM, Maia LFdL, Couto DVD. Estudo comparativo de sintomas respiratórios e função pulmonar em pacientes com doença pulmonar obstrutiva crônica relacionada à exposição à fumaça de lenha e de tabaco %J Jornal Brasileiro de Pneumologia. 2008; 34: 667-674.

[71] Ocakli B, Acarturk E, Aksoy E, Gungor S, Ciyiltepe F, Oztas S, Ozmen I, Agca MC, Salturk C, Adiguzel N, Karakurt Z. The impact of exposure to biomass smoke versus cigarette smoke on inflammatory markers and pulmonary function parameters in patients with chronic respiratory failure. *Int J Chron Obstruct Pulmon Dis* 2018; 13: 1261-1267.

[72] Olloquequi J, Jaime S, Parra V, Cornejo-Córdova E, Valdivia G, Agustí À, Silva O R. Comparative analysis of COPD associated with tobacco smoking, biomass smoke exposure or both. *Respiratory Research* 2018; 19: 13.

[73] Moreira MA, Barbosa MA, Queiroz MC, Teixeira KI, Torres PP, Santana Junior PJ, Montadon Junior ME, Jardim JR. Pulmonary changes on HRCT scans in nonsmoking females with COPD due to wood smoke exposure. *Jornal Brasileiro de Pneumologia : Publicacao Oficial da Sociedade Brasileira de Pneumologia e Tisiologia* 2013; 39: 155-163.

[74] Gelb AF, Zamel N, Hogg JC, Müller NL, Schein MJ. Pseudophysiologic emphysema resulting from severe small-airways disease. *American Journal of Respiratory and Critical Care Medicine* 1998; 158: 815-819.

[75] Rivera RM, Cosio MG, Ghezzi H, Salazar M, Perez-Padilla R. Comparison of lung morphology in COPD secondary to cigarette and biomass smoke. *The International Journal of Tuberculosis and Lung Disease : The Official Journal of the International Union against Tuberculosis and Lung Disease* 2008; 12: 972-977.

[76] Palacios DM, Méndez O. Neumopatía por humo de leña Un estudio en autopsias. *Biomédica* 1998; 18: 153-160.

[77] Restrepo J, Reyes P, de Ochoa P, Patino E. Neumoconiosis por inhalación



del humo de leña. *Acta Med Colomb* 1983; 8: 191-204.

[78] Miravittles M, Soler-Cataluña JJ, Calle M, Molina J, Almagro P, Quintano JA, Riesco JA, Trigueros JA, Piñera P, Simón A, Rodríguez-Hermosa JL, Marco E, López D, Coll R, Coll-Fernández R, Lobo MÁ, Díez J, Soriano JB, Ancochea J. Guía española de la EPOC (GesEPOC). Actualización 2014. *Archivos de Bronconeumología* 2014; 50: 1-16.

[79] Sertogullarindan B, Gumrukcuoglu HA, Sezgi C, Akil MA. Frequency of pulmonary hypertension in patients with COPD due to biomass smoke and tobacco smoke. *International Journal of Medical Sciences* 2012; 9: 406-412.

[80] Sandoval J, Salas J, Martinez-Guerra ML, Gómez A, Martinez C, Portales A, Palomar A, Villegas M, Barrios R. Pulmonary arterial hypertension and cor pulmonale associated with chronic domestic woodsmoke inhalation. *Chest* 1993; 103: 12-20.

[81] Mehra D, Geraghty PM, Hardigan AA, Foronjy R. A comparison of the inflammatory and proteolytic effects of dung biomass and cigarette smoke exposure in the lung. *PloS one* 2012; 7: e52889.

[82] Golpe R, Mengual-Macenlle N, Sanjuan-Lopez P, Cano-Jimenez E, Castro-Anon O, Perez-de-Llano LA. Prognostic Indices and Mortality Prediction in COPD Caused by Biomass Smoke Exposure. *Lung* 2015; 193: 497-503.

[83] Silva R, Oyarzún M, Olloquequi J. Pathogenic mechanisms in chronic obstructive pulmonary disease due to biomass smoke exposure. *Archivos de Bronconeumología* 2015; 51: 285-292.

[84] Dutta A, Roychoudhury S, Chowdhury S, Ray MR. Changes in

sputum cytology, airway inflammation and oxidative stress due to chronic inhalation of biomass smoke during cooking in premenopausal rural Indian women. *International Journal of Hygiene and Environmental Health* 2013; 216: 301-308.

[85] Vishweswaraiah S, Thimraj TA, George L, Krishnarao CS, Lokesh KS, Siddaiah JB, Larsson K, Upadhyay S, Palmberg L, Anand MP, Ganguly K. Putative Systemic Biomarkers of Biomass Smoke-Induced Chronic Obstructive Pulmonary Disease among Women in a Rural South Indian Population. *Disease Markers* 2018; 2018: 4949175.

[86] Golpe R, Martín-Robles I, Sanjuán-López P, Pérez-de-Llano L, González-Juanatey C, López-Campos JL, Arellano-Orden E. Differences in systemic inflammation between cigarette and biomass smoke-induced COPD. *Int J Chron Obstruct Pulmon Dis* 2017; 12: 2639-2646.

[87] Solleiro-Villavicencio H, Quintana-Carrillo R, Falfán-Valencia R, Vargas-Rojas MI. Chronic obstructive pulmonary disease induced by exposure to biomass smoke is associated with a Th2 cytokine production profile. *Clinical Immunology (Orlando, Fla)* 2015; 161: 150-155.

[88] Fernandes L, Rane S, Mandrekar S, Mesquita AM. Eosinophilic Airway Inflammation in Patients with Stable Biomass Smoke- versus Tobacco Smoke-Associated Chronic Obstructive Pulmonary Disease. *Journal of Health & Pollution* 2019; 9: 191209.

[89] Velasco-Torres Y, Ruiz-López V, Pérez-Bautista O, Buendía-Roldan I, Ramírez-Venegas A, Pérez-Ramos J, Falfán-Valencia R, Ramos C, Montaña M. miR-34a in serum is involved in mild-to-moderate COPD in women exposed to biomass smoke. *BMC Pulm Med* 2019; 19: 227.



- [90] Velasco-Torres Y, Ruiz V, Montaña M, Pérez-Padilla R, Falfán-Valencia R, Pérez-Ramos J, Pérez-Bautista O, Ramos C. Participation of the miR-22-HDAC4-DLCO Axis in Patients with COPD by Tobacco and Biomass. *Biomolecules* 2019; 9.
- [91] Vogelmeier CF, Criner GJ, Martinez FJ, Anzueto A, Barnes PJ, BourbeauJ, CelliBR, ChenR, DecramerM, Fabbri LM, Frith P, Halpin DM, Lopez Varela MV, Nishimura M, Roche N, Rodriguez-Roisin R, Sin DD, Singh D, Stockley R, Vestbo J, Wedzicha JA, Agusti A. Global Strategy for the Diagnosis, Management, and Prevention of Chronic Obstructive Lung Disease 2017 Report. GOLD Executive Summary. *American Journal of Respiratory and Critical Care Medicine* 2017; 195: 557-582.
- [92] Deb A, Agarwal M, Reddy V, Kodgule R, Hemalatha V, Awasthi A, al. e. Effects of a fixed dose once-daily triple drug combination in smokers and non-smokers with COPD. . *The European Respiratory Journal* 2016; 48: PA654.
- [93] Chen J, Li C, Ristovski Z, Milic A, Gu Y, Islam MS, Wang S, Hao J, Zhang H, He C, Guo H, Fu H, Miljevic B, Morawska L, Thai P, Lam YF, Pereira G, Ding A, Huang X, Dumka UC. A review of biomass burning: Emissions and impacts on air quality, health and climate in China. *Science of The Total Environment* 2017; 579: 1000-1034.
- [94] Naeher LP, Brauer M, Lipsett M, Zelikoff JT, Simpson CD, Koenig JQ, Smith KR. Woodsmoke health effects: a review. *Inhalation Toxicology* 2007; 19: 67-106.
- [95] Zelikoff JT, Chen LC, Cohen MD, Schlesinger RB. The toxicology of inhaled woodsmoke. *Journal of Toxicology and Environmental Health Part B, Critical Reviews* 2002; 5: 269-282.
- [96] McCarthy CE, Duffney PF, Gelein R, Thatcher TH, Elder A, Phipps RP, Sime PJ. Dung biomass smoke activates inflammatory signaling pathways in human small airway epithelial cells. *American Journal of Physiology Lung Cellular and Molecular Physiology* 2016; 311: L1222-l1233.
- [97] Kc R, Hyland I, Smith J, Shukla S, Hansbro P, Zosky G, Karupiah G, O'Toole R. Cow Dung Biomass Smoke Exposure Increases Adherence of Respiratory Pathogen Nontypeable Haemophilus influenzae to Human Bronchial Epithelial Cells. *Exposure and Health* 2020; 12.
- [98] Magitta NwF, Walker RW, Apte KK, Shimwela MD, Mwaiselage JD, Sanga AA, Namdeo AK, Madas SJ, Salvi SS. Prevalence, risk factors and clinical correlates of COPD in a rural setting in Tanzania. 2018; 51: 1700182.