

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Treatment of Chikungunya Virus (CHIKV) Using Targeted Immunotherapy

*Fleury Augustin Nsole Biteghe,
Chalomie Nyangone Ekome Toung, Jean De La Croix Ndong,
Neelakshi Mungra, Tahir B. Dar and Arnaud John Kombe Kombe*

Abstract

Chikungunya virus (CHIKV) is the most common mosquito-borne *Alphavirus* infecting humans worldwide. Up to date, there are no antiviral treatments or vaccines approved to treat or prevent CHIKV for which treatments remain symptomatic based on clinical manifestations. Hence, designing effective therapies to either prevent or treat CHIKV infection is of paramount importance. Interestingly, monoclonal antibodies (mAbs) are known to be significantly important in mediating protective immunity in CHIV infection. During the last decades, numerous animal studies have reported the protective and prophylactic efficacy of human and mouse anti-CHIKV mAbs isolated from convalescent patients. However, the therapeutic benefits of these anti-CHIKV mAbs can be limited by multiple factors. Thus, it becomes pertinent to better understand the CHIKV infection dynamics, mitigate the undesired mAbs-associated effects and improve therapies. In this review, we critically discuss CHIKV antiviral infectious mechanisms and address how the improved understanding of the latter may pave the way to better targeted immunotherapies.

Keywords: Therapeutics, Antibodies, Vaccines, Prophylactics, Chikungunya virus

1. Introduction

Chikungunya virus (CHIKV) is an arthropod-borne virus firstly discovered during the Tanzanian outbreak in 1952 and isolated a year later (1953) from patient serum and mosquitoes [1]. CHIKV is a worldwide epidemic threat responsible for self-limited fever, maculopapular rashes, and debilitating polyarthralgia in most (90–92%) infected patients [1, 2]. Its name, meaning “that which contorts or bends up” in “Kimakonde”, a Tanzanian and Mozambican vernacular language, stems from the stooped posture exhibited by infected patients [3–6]. In endemic areas, CHIKV can be misdiagnosed as it displays dengue (DENV) or zika virus (ZIKV) like symptoms [1, 7]. Although less lethal, CHIKV-associated mortality rate can be influenced by other factors, including immunocompromised individuals, newborns from high viremic mothers or patients with preexisting arthritis [1]. CHIKV represents a serious economic burden, affecting the physical status of infected

patients, restraining them from working for up to 35 days, as reported during the 2007 Indian outbreak [8]. For that reason, CHIKV was categorized as a biodefense pathogen by the National Institute of Allergy and Infectious Diseases (NIAID) in the USA [9]. Presently, there are no clinically licensed vaccines or therapies to treat CHIKV. Nonetheless, several pre-clinical animal models using antibody-based immunotherapy have shown some promising results in preventing and treating CHIKV infections [1, 7, 9, 10]. Despite several obstacles that have to be overcome to reach clinical fruition of such therapy [11], mAbs therapies offer better therapeutic avenues with respect to emerging disease outbreaks [12]. Hence, this review succinctly describes CHIKV characteristics (Transmission, structure and diagnosis) and highlights the potential therapeutic usage of mAbs based on their protective role in naturally occurring humoral immunity following CHIV infection. Lastly, we briefly discuss some challenges associated with mAbs therapy and propose future alternative therapeutic approaches.

2. Chikungunya's transmission cycle

CHIKV can be transmitted either horizontally from human to human through mosquito bites (*Aedes* species) [1] or vertically from a mother to a child during pregnancy or at birth, thus causing severe neonatal infection [1, 13, 14]. However, the rate of CHIKV transmission can be affected by the geographical location and the vector. It has been historically reported that CHIKV was predominant in Sub-Saharan Africa (SSA), Southeast Asia tropical and sub-tropical regions, where two different transmission cycles occur [15]. In rural areas, CHIKV replicates through a sylvatic transmission cycle (animal to human transmission via the mosquito), involving forest or savanna *Aedes* mosquitoes (*A. furcifer* and *A. africanus*) and animals (domestics and non-domestics), representing the main CHIKV reservoirs within this cycle [16]. The urban transmission cycle is mediated by *A. albopictus* (also known as Asian tiger mosquito) and *A. aegypti*, both maintaining the human-to-human transmission, thus making humans the principal CHIKV reservoirs in the urban epidemic cycle (**Figure 1**). Interestingly, Chikungunya fever outbreaks in Asia have been associated with only the




Figure 1. CHIKV transmission cycle: Two cycles known as sylvatic and urban characterized the CHIKV transmission cycle. While the sylvatic predominantly occurs in Africa and circulates between primates, rodents and other vertebrates, the urban counterparts is commonly observed in Asia, where CHIKV is horizontally transferred from human to human through mosquito bites or blood meal.

urban CHIKV transmission cycle, while the African counterparts are related to both sylvatic and urban transmission cycles [15, 17]. Based on this evidence, it can hypothetically be suggested that the CHIKV transmission cycle before 1952 was restricted to the sylvatic cycle and that its urban transmission resulted from demographic expansion through deforestation or migration of CHIKV infected travelers from rural to the most populated urban cities. While the natural factors sustaining CHIKV infection remain elusive [16], it can be hypothesized that simultaneous presence of *Aedes* specie mosquitoes and CHIKV carriers (even with very low viral load in the blood) in a given area is enough for the resurgence of CHIKV-associated outbreaks.

3. Clinical manifestations of chikungunya fever

Chikungunya virus (CHIKV) is an emerging and re-emerging arbovirus associated with high morbidity, characterized by multiple clinical symptoms, with typical and/or atypical signs. CHIKV infection can be broken down into three phases, including the acute (the first 21st days following the onset of clinical symptoms), post-acute (from the 3rd week till the end of the 3rd month) and chronic phase (3 months following the onset of clinical symptoms) [1]. Therefore, chikungunya can present diverse clinical symptoms, for which 3–25% of infected individuals are asymptomatic (**Figure 2**) [18]. During the symptomatic phase, the majority of infected patients (90–95%) develop long lasting clinical symptoms coinciding with viremia peak (Elevated viral load ranging between 10^5 and 10^{12} viruses per milliliter of blood) and manifested as debilitating polyarthralgia, myalgia and arthralgia which are rarely fatal (1 in 1000) and not observed in dengue fever [18–20]. Atypical cases of CHIKV occur less frequently and mainly affect elderly, immunocompromised individuals and neonates from viremic mothers [1, 18]. The clinical manifestations associated with these atypical cases include severe neurological dysfunctions (Encephalitis), ocular changes, cardiovascular defect, preterm birth, hyperpigmentation, bullous dermatosis and Guillain-Barré syndrome [1, 18].




Figure 2. Acute CHIKV clinical symptoms: During the symptomatic phase, CHIKV reportedly causes self-limited fever, as well as debilitating polyarthralgia in most (85–92%) infected patients [18–20]. Also, CHIKV can manifest atypical dermatological symptoms (42%) including edema, hemorrhagic bullous skin lesions, hyperpigmentation [21] and neurological symptoms, characterized by encephalopathy, encephalitis, Guillain-Barré or encephalomyelora-diculitis [22].

Yet, the mechanisms contributing to the long-lasting clinical symptoms remain elusive. An attempt to elucidate this mechanism was made by Reddy et al., 2017, demonstrating CHIKV's ability to evade host immunity as a result of CHIKV E1 glycoprotein homology to the host human counterpart [23]. Consequently, a more detailed understanding of CHIKV replication and infectious cycle is necessary to mitigate these undesired effects.

4. Chikungunya virus genome structure and infectious cycle

CHIKV is a small enveloped virus with a positive single-strand RNA virus belonging to the *Alphavirus* genus from the *Togaviridae* family [1, 21]. It is also referred to as an arbovirus (for *arthropod-borne virus*) due to arthropods transmission (*Aedes* mosquitoes) [21]. CHIKV has an RNA genome of approximately 11.8 kb, comprising two open reading frames (ORFs), known as 5'capped ORF (7424 nucleotides) and 3'polyadenylated ORF (3732 nucleotides), linked by a junction region [22]. The 5'end is protected by a 7-methylguanosine (7MG) cap, with the ORF accounting for 66% of the genome encoding a precursor protein, which upon processing produces four nonstructural proteins (nsP1–4) responsible for CHIKV replication in host cell cytoplasm upon entry [1, 22]. On the other hand, the 3'ORF ends with a polyadenylation signal (3'poly-A tail) and encodes another precursor protein, generating five structural polyprotein comprising: the capsid C, envelop glycoproteins (E1, E2 and E3) and the 6 K protein [1, 22]. **Figure 3A** give a graphic representation of CHIKV genome. Once matured (after replicating in host cells), the virion particle has about 70 nm diameter and possesses 240 copies of capsid proteins embedded in 80 spikes of envelope trimers (glycoproteins E2/E1), which are inserted within the plasma membrane of the infected cells (to evade host immunity) that they use to bud out of the host cells through a secretory pathway [1, 22] (**Figure 3B**). Therefore, elucidating CHIKV replication cycle in both mammalian and mosquito cells becomes pertinent to develop efficient therapies.

CHIKV primarily replicates in the epithelial cells of mosquito midgut before migrating into the salivary glands where they keep replicating throughout the insect life, prior dissemination into the bloodstream of mammalian host, following mosquito bite/blood meal [1, 25]. Once deposited in the human bloodstream or skin after an infected mosquito bite (or blood meal), CHIKV firstly replicates within fibroblasts and macrophages found at the site of inoculation [16]. Thereafter, CHIKV systemically propagates within the body through the lymphatic system to multiple replication sites, where prominent disease symptoms occur (lymph nodes, spleen, skin, muscles, peripheral joints, brain, liver and tendons) despite innate immunity [16]. Notably, CHIKV replication in peripheral tissues is associated with elevated viral loads ($>10^9$ virus particles/ml), which is conducive for mosquito transmission during episodic bloodmeal or bite [16, 26]. Moreover, CHIKV can be transmitted to humans in various ways, including infected needles, contaminated blood donation, organ graft and from viremic mother to newborn [1, 27]. In this regard, Campos et al., 2017 study primarily highlighted the presence of CHIKV in the breast milk of an infected mother, who tested positive for serum and urine [28]. Fortunately, this study did not reveal a breastfeeding transmission capacity of the infected breast milk, as the 3-month-old newborn CHIKV serology and reverse transcriptase-polymerase chain reaction (RT-PCR) tested negative [1, 28]. In light of the latter results, it is worth mentioning that only the structural proteins (surface glycoproteins E2 and E1) are present on the outer surface of CHIKV envelop and are endowed with antigenic potential, able to elicit activation of host immune defense mechanisms [10].




Figure 3.

Chikungunya virus genome structure and infectious cycle. (A) Chikungunya genome structure, (B) chikungunya (CHIK) internalizes within the target cell through receptor mediated endocytosis. Upon entry, CHIK undergo conformational changes leading to E1 peptide exposure as a result of endosomal microenvironment acidity, which favors virus–host cell membrane fusion and subsequent cytoplasmic release of nucleocapsid core and viral genome. Thereafter, two viral mRNA strands are translated to give to two non-structural proteins (nsPs), which are cleaved to generate nsP1–nsP4. The function of the latter are as followed: 1) nsP1 synthesis the negative viral strand RNA and cap it, 2) nsP2 exhibits RNA triphosphatase, helicase and proteinase functions contributing to host cell transcriptional machinery inhibition, 3) nsP3 forms part of the replicase unit and 4) nsP4 exert viral RNA polymerase function. The coordinated activity of these proteins is to produce the viral replication complex, enabling the synthesis of the full-length negative-strand RNA intermediate serving as the precursor to generate the subgenomic (26S) and genomic (49S) RNAs. Once synthesized, the 26S mRNA is translated into C–pE2–6 K–E1 polypeptide precursors, which are further processed by a serine protease to produce a capsid (C), pE2 and E1 glycoproteins. Then, pE2 and E1 assemble in the Golgi and translocate to the plasma membrane, where pE2 is cleaved into E2 (enabling receptor binding) and E3 (ensuring correct pE2 folding and its subsequent assembly with E1). Completion of this viral replication cycle is achieved through 1) assembly of the viral nucleocapsid with the viral RNA, 2) recruitment of membrane-associated envelope glycoproteins and 3) budding at the virus at the host cell membrane [24].

5. CHIKV induced immune response

5.1 The role of innate immunity in chikungunya infection management

The innate immune system refers to the first non-physical and non-specific defense mechanisms to encounter invading pathogens or foreign elements immediately or hours following the appearance of their antigens within the body [29]. To perform its protective role, the host innate immunity relies on antigen-presenting cells (APC) such as macrophages (Mc) and dendritic cells (DC) to neutralize foreign antigens and subsequently present it in a specific way to specialized lymphocytes B (antibody producing cells) and T-cells of the adaptive immunity in secondary lymphoid organs (spleen and ganglions) [30, 31]. Conventionally, all APCs are endowed with the ability to capture an antigen originating from extracellular or intracellular milieu and present it to CD4+ or CD8+ T cells, using major histocompatibility complex II (MHC-II) or I (MHC I) associated peptides respectively [30–32]. However, DCs are endowed with the unique potency to prime CD4+ and CD8+ T-cells (in secondary lymphoid organs) due to their ability to cross-present captured foreign antigens using MHC-II/I complexes [32–34]. Under normal conditions, DC cells exist in immature and inactivated states, hence acting as the sentinel of the immune system [31, 34, 35]. However, DCs are activated into a mature state once exposed to the antigenic determinant of the foreign organism. This DC protective role was illustrated by Long et al., 2013 reporting increased DCs presence at CHIKV infection site 24 and 36 h post infection [36]. Additionally, the latter report highlighted that mice harboring DCs deficiency for dendritic immunogenic receptor (DCIR) displayed more severe CHIKV related symptoms (increased inflammation, edema, weight loss and damage of inoculated foot and the ankle joint) than wild type control [36]. In this line, Das et al., 2015, demonstrated an increase in CD206+ DCs mobilization at the CHIKV infected astrocytes site [37].

Nevertheless, it is well documented that innate immunity following viral infection can recruit inflammatory cells to infected musculoskeletal tissues, with the potential to cause muscular and articular damages inducing the observed pain and discomfort in muscles, joints, and tendons [16]. This CHIKV dependent polyarthritis-like symptoms were reported by Amdekar et al., 2017, to be linked with host inflammatory response regulated by infiltrating macrophages (in joints), T cells, and viral persistence within the immune inaccessible infected sites [38]. Similarly, numerous studies have demonstrated the dichotomic CHIKV reservoir (during chronic infection phase) and modulatory macrophage functions (keeping

local Th1/Th2 balance) in the affected tissues (muscles, joints, lymphoid tissues and liver) that they co-infiltrated with mononuclear inflammatory cells [39–41]. These aforementioned results corroborated Gardner et al., 2010 study, revealing the equally important role of macrophages in establishing arthritis symptoms and CHIKV clearance [42]. Furthermore, Phuklia et al., 2013 and Schett, 2007 showed that CHIKV-infected synoviocytes (Cells producing the synovial fluid component important for absorption from the joint cavity and for synovial/blood fluid exchange) were able to promote the migration and differentiation of both monocytes and macrophages into joints damaging osteoclast-like cells, capable of producing high levels of arthritis mediators (Interleukin-6 and Tumor Necrosis Factor- α) [43, 44]. In contrast, Haist et al., 2017, underlined the protective role of inflammatory Ly6C^{hi} CCR2⁺ monocytes in controlling CHIKV infectious, as their depletion (using diphtheria toxin on CCR2-DTR⁺ mice) was associated with severe disease symptoms and reduction in Ly6C and NK cells in bloodstream and muscles when compared to wild type mice [45].

6. Chikungunya and adaptive immunity

Humoral and cell-mediated immunity hold the center stage for protection against CHIKV infection through the combined action of T and B-lymphocytes, respectively [46]. Multiple human and animal studies have demonstrated the host immune system's ability to induce the production of neutralizing anti-CHIKV antibodies, to rapidly clear CHIKV viral loads and establish long-term immunity [1, 10, 46]. This adaptive humoral immunity relies on immunoglobulin-M (IgM: representing ~30% of circulating antibodies) to primarily entrap the CHIKV antigen to secondary lymphoid tissues by bridging the innate and adaptive immunity before the onset of the robust and specific life-long protecting IgG response [1, 47]. During the chikungunya fever acute phase, neutralizing IgM are produced as early as 2–6 days post-infection and act to reduce viral loads (**Figure 4**) [48]. This neutralizing IgM capacity perdures up to 10 days following CHIKV infection before synergizing with specific IgG response [1]. Interestingly, Couderc and Lecuit et al., 2015 reported persistent anti-CHIKV IgM in chronic chikungunya fever patients related to repeated CHIKV antigen and RNA exposure [49]. Of late, Tanabe et al., 2018 showed that IgM antibodies preferentially interact with cognate E1-E2 fusion glycoprotein epitopes found on CHIKV outer surface [23, 46] and that long term viral immune protection was achieved through specific IgG production, persisting up to 19 years following the first infectious episode (1991 Thailand CHIKV outbreak) [50]. Among all IgG subtypes, IgG3 was found to be the most predominant, able to serve as a predictive marker of high-risk patients (when minimally produced) or as a prophylactic agent, able to passively transfer from mother to newborn via the placenta and limit viral metastases to distant muscles and joints [1]. This pentameric immunoglobulin structure armed them with the ability to easily mediate CHIKV aggregation, recognition and elimination by cytotoxic CD8⁺ T cells [46]. Unfortunately, CD4⁺ T-cells can be detrimental to CHIKV infected mice, as it was reported by Theo et al., 2017 that they can severely damage the joints of T-cell receptor-deficient mice when adoptively transferred [46, 51]. Mitigation of these undesired CD4⁺ T-cell effects could be performed using fingolimod (phosphorylated sphingosine 1-phosphate receptor 1 agonist: S1PR1) treatment, blocking S1PR1 mediated CD4⁺ T-cells dissemination to the joints of CHIKV infected mice [46, 51]. In a parallel study, Miner et al., 2017 demonstrated the synergy of immunosuppressive cytotoxic T-lymphocyte-antigen-4 (CTLA-4) abatacept (blocking CD4 co-stimulatory activating signals) with anti-CHIKV (reducing viral load) in controlling arthritis and CHIKV infection, respectively [52]. This result was




Figure 4. CHIKV infection and immune response: CHIKV infection begin with mosquito's bites, allowing viral entry into serum and subsequent dissemination into peripheral organs (lymphoid, muscles and joints), leading to the onset of symptoms (acute phase) that can perdure beyond 8 days. During the acute phase (0–4 days), neutralizing IgMs are produced, and their increase inversely correlates with CHIKV viral loads. During the recovery phase or convalescence (beyond 7 days), IgGs are produced and act in concert with IgM to provide long-term immune protection.

particularly relevant, as it may pave the way for novel antibody therapeutic strategies, with the potential to clinically benefit CHIKV infected patients.

7. Chikungunya virus therapeutic treatment and future perspectives

Targeted antibody-based immunotherapy was developed to address the unmet CHIKV therapeutic clinical needs [1]. In this regard, immunotherapy using IgG has widely been developed and has demonstrated the potential to reach clinical fruition by exerting its virucidal (Virus killing activity) and prophylactic activities on infected patients [1]. This IgG virucidal effect was confirmed by Scott et al., 2017 showing its ability to induce complete recovery in encephalitis infected patients, using a therapeutic dose of 400 mg/kg for 5 days [2]. With this in mind, Fernandes et al., 2019 corroborated these latter results, by achieving a 10 day total recovery, following a combination therapy involving antibiotic and IgG treatment (400 mg/kg for 5 days) on a severely infected 56 years old patient, presenting with atypical dermatological form associated with edema and hemorrhagic bullous skin lesions [53]. This IgG immunotherapeutic agent could either be used to immunized healthy or asymptomatic individuals in endemic areas (with the aim to reduce CHIKV propagation) or to treat high-risk patients such as pregnant women, newborns, immunocompromised individuals, elders or patients with preexisting arthritis [1]. For example, Couderc et al. 2009 demonstrated the therapeutic and prophylactic actions of IgGs (purified from convalescent CHIKV infected donor), protecting adult immunocompromised and immunocompetent mice neonates from CHIKV

re-challenge and severe neurological symptoms associated with chronic CHIKV infection [54]. Based on these accumulative reports, a phase I/II clinical trial was initiated to evaluate the therapeutic and prophylactic efficacy of CHIKV IgG therapy to newborns of mothers with acute CHIKV infection, whose infectious rates reached 49% (Clinical trials: NCT02230163) [14]. However, the efficacy of CHIKV IgG immunotherapy can be compromised by immune evasion, arising from monoclonal Abs (mAbs) selective pressure or accumulated genetic mutations able to suppress antigen specificity, thus antigen–antibody interaction [1]. With the advent of recombinant antibody engineering technology, efforts to overcome these unwanted effects were developed and led to the generation of bispecific antibodies that can recognize two or more epitopes; a detailed discussion beyond the scope of this review has been provided by Nyakatura et al., 2017 [55]. Besides immune evasion, other factors like antibody-dependent enhancement (ADE), characterized by sub-neutralizing Abs concentration that is permissive for viral replication, has been identified as a major challenge for CHIKV management [1, 56]. In spite of the single serotype status of CHIKV, ADE cannot be overlooked with respect to CHIKV reemergence, as it has been observed in other viral diseases, including Dengue, Rabies virus, influenza and HIV [1]. Also, antibody-based therapy is generally associated with high production costs, which may restrain therapeutic usage due to affordability. To address these issues, nucleic acid vaccines made up of mRNA or




Figure 5. mRNA vaccine mechanism of action. Conventional mRNA possesses disease specific gene of interest (GOI) encoding the antigen DNA sequence that is respectively capped at the 5' end and polyadenylated (poly-a tail) at the 3' untranslated region (UTR). Upon internalization within the host cell, viral mRNA is released from the endosome to the cytosol, where it is immediately translated by the ribosomes and post-translationally modified by the proteasome. In opposition to the conventional mRNA vaccination, the self-amplifying mRNA method mostly uses positive-sense single-stranded viral RNA encoding the disease specific antigen and viral nonstructural proteins (nsPs) driving intracellular RNA amplification and significant antigen expression. These two vaccination methods need a delivery system for host cell internalization, commonly performed through endocytosis, preceding cytosolic viral mRNA cargo releases from the endosome, driving translation and subsequent viral antigen protein processing for MHC presentation to ensue and activate host immunity [59].

plasmid DNA (pDNA) were developed [57, 58]. These novel vaccines are commonly used to treat mRNA alphavirus and are mainly produced by genetically substituting viral encoding structural protein with target antigen encoding genes while preserving the RNA replication machinery [57]. Hence, when injected into patients, these vaccines (mRNA) mimic the virus by taking advantage of the host cell machinery to efficiently transcribe an antigen encoded RNA into an antigen encoded DNA, which is subsequently translated and post-translationally modified into a form (protein) that resemble the natural antigen that can successively stimulate and activate the innate and adaptive immunity (**Figure 5**) [57, 58]. Recently, Kose et al. (2019) demonstrated CHIKV and arthritis protecting properties of an infused nanoparticle encoding CHIKV antibody mRNA, which was as good as purified IgG mAbs in stimulating protective anti-CHIKV serum concentrations in macaques [60].

So far, there are no clinically approved RNA vaccines to treat human diseases. Yet, in response to the ongoing coronavirus disease 2019 (COVID-19) pandemic, multiple clinical trials such as NCT04283461, have been initiated to test the safety of an mRNA vaccine (encoding SARS-CoV-2 spike protein), which successfully produced SARS-CoV-2 antibodies and had no clinically adverse effects [61, 62]. Currently, it has progressed into phase III clinical trial (NCT04470427) to evaluate its efficacy against COVID-19. Likewise, other clinical trials are presently investigating the safety, tolerability and immunogenicity of zika virus envelop protein encoded mRNA vaccines (NCT04064905; NCT03014089) [61, 63]. Taken altogether, it becomes evident that the rapid advances made in recombinant DNA technology, will revolutionize the way we approach and promptly respond to endemic CHIKV/COVID-19 like diseases.

IntechOpen

IntechOpen

Author details

Fleury Augustin Nsole Biteghe^{1*†}, Chalomie Nyangone Ekome Toug²,
Jean De La Croix Ndong³, Neelakshi Mungra⁴, Tahir B. Dar¹
and Arnaud John Kombe Kombe^{5†}

1 Department of Radiation Oncology and Biomedical Sciences, Cedars-Sinai
Medical, Los Angeles, USA

2 Zhongshan Medical School, Sun Yat-Sen University, Guangzhou, China

3 Federated Departments of Biological Sciences, Rutgers University School of Arts
and Science, USA

4 Faculty of Health Sciences, Medical Biotechnology and Immunotherapy Research
Unit, Institute of Infectious Disease and Molecular Medicine, University of Cape
Town, Cape Town, South Africa

5 Division of Molecular Medicine, Hefei National Laboratory for Physical Sciences
at Microscale, CAS Key Laboratory of Innate Immunity and Chronic Disease,
Division of Life Sciences and Medicine, University of Science and Technology of
China, Hefei, China

*Address all correspondence to: fleury.nsolebiteghe@cshs.org

†Both authors equally contributed

IntechOpen

© 2021 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms
of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium,
provided the original work is properly cited. 

References

- [1] Kumar R, Shrivastava T, Samal S, Ahmed S, Parray HA. Antibody-based therapeutic interventions: possible strategy to counter chikungunya viral infection. *Appl Microbiol Biotechnol*. 2020;104(8):3209-28.
- [2] Scott SS de O, Braga-Neto P, Pereira LP, Nóbrega PR, de Assis Aquino Gondim F, Sobreira-Neto MA, et al. Immunoglobulin-responsive chikungunya encephalitis: two case reports. *J Neurovirol*. 2017;23(4):625-31.
- [3] MC R. An epidemic of virus disease in Southern Province, Tanganyika Territory, in 1952-53. *Trans R Soc Trop Med Hyg*. 1955;49(1):28-32.
- [4] Thiboutot MM, Kannan S, Kawalekar OU, Shedlock DJ, Khan AS, Sarangan G et al. Chikungunya: a potentially emerging epidemic? *PLoS Negl Trop Dis*. 2010;4(4):e63.
- [5] SB H. Reappearance of chikungunya, formerly called dengue, in the Americas. *Emerg Infect Dis*. 2015;21(4):557-61.
- [6] Halstead SB. Reappearance of chikungunya, formerly called Dengue, in the Americas. *Emerg Infect Dis*. 2015;21(4):557-61.
- [7] Clayton AM. Monoclonal antibodies as prophylactic and therapeutic agents against chikungunya virus. *J Infect Dis*. 2016;214(Suppl 5):S506-9.
- [8] Gopalan, S.S.; Das A. Household economic impact of an emerging disease in terms of catastrophic out-of-pocket health care expenditure and loss of productivity: Investigation of an outbreak of chikungunya in orissa, india. *J Vector Borne Dis*. 2009;46:57-64.
- [9] Jin J, Simmons G. Antiviral functions of monoclonal antibodies against chikungunya virus. *Viruses*. 2019;11(4):1-17.
- [10] Powers AM. Vaccine and therapeutic options to control chikungunya virus. *Clin Microbiol Rev*. 2018;31(1):1-29.
- [11] Burt FJ, Chen W, Miner JJ, Lenschow DJ, Merits A, Schnettler E, Kohl A, Rudd PA, Taylor A, Herrero LJ, Zaid A, NgLFP MS. Chikungunya virus: an update on the biology and pathogenesis of this emerging pathogen. *Lancet Infect Dis*. 2017;17(4):e107-17.
- [12] Marston HD, Paules CI FA. Monoclonal antibodies for emerging infectious diseases - borrowing from history. *N Engl J Med* 378:1469-1472. 2018;378:1469-72.
- [13] Gerardin P, Samperiz S, Ramful D, Boumahni B, Bintner M, Alessandri JL et al. Neurocognitive outcome of children exposed to perinatal mother-to-child Chikungunya virus infection: the CHIMERE cohort study on Reunion Island. *PLoS Negl Trop Dis*. 2014;8(7):e2996.
- [14] Cardona-Correa SE, Castaño-Jaramillo LM, Quevedo-Vélez A. Vertical transmission of chikungunya virus infection. Case report. *Rev Chil Pediatr*. 2017;88(2):285-8.
- [15] Burt FJ, Rolph MS, Rulli NE, Mahalingam S, Heise MT. Chikungunya: A re-emerging virus. In: *The Lancet*. Elsevier; 2012. p. 662-71.
- [16] Silva LA, Dermody TS. Chikungunya virus: Epidemiology, replication, disease mechanisms, and prospective intervention strategies. *J Clin Invest*. 2017;127(3):737-49.
- [17] Silva LA DT. Chikungunya virus: epidemiology, replication, disease mechanisms, and prospective intervention strategies. *J Clin Invest*. 2017;127(3):737-49.

- [18] 1. Moizéis RNC, Fernandes TAA de M, Guedes PM da M, Pereira HWB, Lanza DCF, Azevedo JWV de et al. Chikungunya fever: a threat to global public health. *Pathog Glob Heal*. 2018;112(4):182-94.
- [19] Sa PKO, Nunes MM, Leite IR, Campelo M, Leao CFR, Souza JR, Castellano LR FA(. Chikungunya virus infection with severe neurologic manifestations: report offour fatal cases. *Rev Soc Bras Med Trop*. 2017;50: 265-2687.
- [20] Josseran L, Paquet C, Zehgnoun A, Caillere N, Le Tertre A, Solet JL LM. Chikungunya disease outbreak, Reunion Island. *Emerg Infect Dis*. 2006;12: 1994-5.
- [21] Weaver SC, Osorio JE, Livengood JA, Chen R SD. Chikungunya virus and prospects for a vaccine. *Expert Rev Vaccines*. 2012;11(9):1087-101.
- [22] Weaver SC, Osorio JE, Livengood JA, Chen R, Dan T, Weaver SC, et al. Chikungunya virus and prospects for a vaccine Chikungunya virus and prospects for a vaccine. *Expert Rev Vaccines*. 2012;11(9):1087-101.
- [23] Reddy V, Desai A, Krishna SS, Vasanthapuram R. Molecular Mimicry between Chikungunya Virus and Host Components: A Possible Mechanism for the Arthritic Manifestations. *PLoS Negl Trop Dis*. 2017;11(1):1-20.
- [24] Schwartz O, Albert ML. Biology and pathogenesis of chikungunya virus. *Nat Rev Microbiol*. 2010;8(7):491-500.
- [25] Moizéis RNC, Fernandes TAA de M, Guedes PM da M, Pereira HWB, Lanza DCF, Azevedo JWV de, et al. Chikungunya fever: a threat to global public health. *Pathog Glob Health*. 2018;112(4):182-94.
- [26] Das T et al. Chikungunya fever: CNS infection and pathologies of a re-emerging arbovirus. *Prog Neurobiol*. 2010;91(2):121-9.
- [27] Rosso F, Rodriguez S, Cedano JA, Mora BL, Moncada PA VJ(. Chikungunya in solid organ transplant recipients, a case se- ries and literature review. *Transpl Infect Dis*. 2018;20(6):e12978.
- [28] Campos GS, Albuquerque Bandeira AC, Diniz Rocha VF, Dias JP, Carvalho RH SS. First detection of chikungunya virus in breast milk. *Pediatr Infect Dis J*. 2017;36(10):1015-7.
- [29] Dar TB, Henson RM, Shiao SL. Targeting innate immunity to enhance the efficacy of radiation therapy. *Front Immunol*. 2019;9:1-11.
- [30] Woo S-R, Corrales L, Gajewski TF. Innate Immune Recognition of Cancer. *Annu Rev Immunol*. 2015;33(1):445-74.
- [31] Patente TA, Pinho MP, Oliveira AA, Evangelista GCM, Bergami-Santos PC, Barbuto JAM. Human dendritic cells: Their heterogeneity and clinical application potential in cancer immunotherapy. *Front Immunol*. 2019;9(3176):1-18.
- [32] Garg AD, Coulie PG, Van den Eynde BJ, Agostinis P. Integrating Next-Generation Dendritic Cell Vaccines into the Current Cancer Immunotherapy Landscape. Vol. 38, *Trends in Immunology*. Elsevier Ltd; 2017. p. 577-93.
- [33] Cance JC, Crozat K, Dalod M, Mattiuz R. Are conventional type 1 dendritic cells critical for protective antitumor immunity and how? *Front Immunol*. 2019;10(FEB).
- [34] Bol KF, Schreibelt G, Gerritsen WR, De Vries IJM, Figdor CG. Dendritic cell-based immunotherapy: State of the art and beyond. *Clin Cancer Res*. 2016;22(8):1897-906.

- [35] Shang N, Figini M, Shangguan J, Wang B, Sun C, Pan L, et al. Dendritic cells based immunotherapy. Vol. 27, Cell Research. E-Century Publishing Corporation; 2017. p. 74-95.
- [36] Long KM, Whitmore AC, Ferris MT, Sempowski GD, McGee C, Trollinger B, et al. Dendritic Cell Immunoreceptor Regulates Chikungunya Virus Pathogenesis in Mice. *J Virol*. 2013;87(10):5697-706.
- [37] Das T, Hoarau JJ, Bandjee MCJ, Marianne M, Gasque P. Multifaceted innate immune responses engaged by astrocytes, microglia and resident dendritic cells against Chikungunya neuroinfection. *J Gen Virol*. 2015;96(2):294-310.
- [38] Sarika Amdekar, Deepti Parashar and KA. Chikungunya Virus-Induced Arthritis: Role of Host and Viral Factors in the Pathogenesis. *Viral immunology*. 2017;30(10):691-702.
- [39] Labadie, K., Larcher, T., Joubert, C., Mannioui, A., Delache, B., Brochard, P. et al. Chikungunya disease in nonhuman primates involves long-term viral persistence in macrophages. (*J Clin Invest*. 2010;120:894-906.
- [40] Dupuis-Maguiraga, L., Noret, M., Brun, S., Le Grand, R., Gras, G., and Roques P. Chikungunya disease: infection-associated markers from the acute to the chronic phase of arbovirus-induced arthralgia. *PLoS Negl Trop Dis*. 2012;6:e1446.
- [41] Ziegler, S. A., Lu, L., da Rosa, A. P., Xiao, S. Y., and Tesh RB. An animal model for studying the pathogenesis of chikungunya virus infection. *Am J Trop Med*. 2008;79:133-9.
- [42] Gardner J, Anraku I, Le TT, Larcher T, Major L, Roques P, et al. Chikungunya Virus Arthritis in Adult Wild-Type Mice. *J Virol*. 2010;84(16):8021-32.
- [43] Phuklia, W., Kasisith, J., Modhiran, N., Rodpai, E., Thannagith, M., Thongsakulprasert, T. et al. Osteoclastogenesis induced by CHIKV-infected fibroblast-like synoviocytes: a possible interplay between synoviocytes and monocytes/macrophages in CHIKV-induced arthralgia/arthritis. *Virus Res*. 2013;177:179-88.
- [44] Schett G. Cells of the synovium in rheumatoid arthritis. Osteoclasts. *Arthritis Res Ther*. 2007;9(1):1-6.
- [45] Haist KC, Burrack KS, Davenport BJ, Morrison TE. Inflammatory monocytes mediate control of acute alphavirus infection in mice. *PLoS Pathog*. 2017;13(12):1-30.
- [46] Tanabe ISB, Tanabe ELL, Santos EC, Martins W V., Araújo IMTC, Cavalcante MCA, et al. Cellular and Molecular Immune Response to Chikungunya Virus Infection. *Front Cell Infect Microbiol*. 2018;8(October):345.
- [47] Vollmers HP BS. The “early birds”: natural IgM antibodies and immune surveillance. *Histol Histopathol*. 2005;20(3):927-37.
- [48] Petitdemange C, Wauquier N V V. Control of immunopathology during chikungunya virus infection. *J Allergy Clin Immunol*. 2015;135(4):846-55.
- [49] Couderc T LM. Chikungunya virus pathogenesis: from bedside to bench. *Antivir Res*. 2015;121:120-31.
- [50] Nitatpattana N, Kanjanopas K, Yoksan S, Satimai W, Vongba N, Langdatsuan S, Nakgoi K, Ratchakum S, Wauquier N, Souris M, Auewarakul P GJ. Long-term persistence of chikungunya virus neutralizing antibodies in human populations of north eastern Thailand. *Virol J*. 2014;11:183.
- [51] Teo, T. H., Chan, Y. H., Lee, W. W., Lum, F. M., Amrun, S. N., Her, Z. et al.

Fingolimod treatment abrogates chikungunya virus-induced arthralgia. *Sci Transl Med.* 2017;9(375):eaal1333.

[52] Miner JJ, Cook LE, Hong JP, Smith AM, Richner JM, Shimak RM, Young AR, Monte K, Poddar S, Crowe JE Jr, Lenschow DJ DM. Therapy with CTLA4-Ig and an antiviral monoclonal antibody controls chikungunya virus arthritis. *Sci Transl Med.* 2017;9(375):eaah3438.

[53] Fernandes AI V., Souza JR, Silva AR, Cruz SBSC, Castellano LRC. Immunoglobulin therapy in a patient with severe chikungunya fever and vesiculobullous lesions. *Front Immunol.* 2019;10(JUL):1-6.

[54] Couderc T, Khandoudi N, GrandadamM, Visse C, Gangneux N, Bagot S, Prost JF LM. Prophylaxis and therapy for chikungunya virus infection. *J Infect Dis.* 2009;Couderc T, (4):516-23.

[55] Nyakatura EK, Soare AY, Lai JR. Bispecific antibodies for viral immunotherapy. *Hum Vaccines Immunother.* 2017;13(4):836-42.

[56] Lum FM, Couderc T, Chia BS, Ong RY, Her Z, Chow A, Leo YS, Kam YW, Renia L, Lecuit M NL. Antibody-mediated enhancement aggravates chikungunya virus infection and disease severity. *Sci Rep.* 2018;8(1):1860.

[57] Pitman MC, Lau JSY MJ et al. Barriers and strategies to achieve a cure for HIV. *lancet HIV.* 2018;5(317-28).

[58] Dharanidharan Ramamurthy, Trishana Nundalall, Sanele Cingo, Neelakshi Mungra, Maryam Karaan, Krupa Naran1 SB. Recent Advances in Immunotherapies Against Infectious Diseases. *Immunother Adv.* 2020;1-6.

[59] Maruggi G, Zhang C, Li J, Ulmer JB, Yu D. mRNA as a Transformative

Technology for Vaccine Development to Control Infectious Diseases. *Mol Ther.* 2019;27(4):757-72. Available from: <https://doi.org/10.1016/j.ymthe.2019.01.020>

[60] Kose N, Fox JM, Sapparapu G, Bombardi R, Tennekoon RN, de Silva AD, Elbashir SM, Theisen MA, Humphris-Narayanan E, Ciaramella G, Himansu S, Diamond MS CJJ. A lipid-encapsulated mRNA encoding a potentially neutralizing human monoclonal antibody protects against chikungunya infection. *Sci Immunol.* 2019;4(35):eaaw6647.

[61] Kim YC, Dema B, Reyes-Sandoval A. COVID-19 vaccines: breaking record times to first-in-human trials. *npj Vaccines.* 2020;5(1):19-21.

[62] Tung Thanh Le, Zacharias Andreadakis, Arun Kumar, Raúl Gómez Román, Stig Tollefsen, Melanie Saville SM. The COVID-19 vaccine development landscape. *Nat Rev Drug Discov.* 2020;19(5):305-6.

[63] Jackson NAC, Kester KE, Casimiro D, Gurunathan S, DeRosa F. The promise of mRNA vaccines: a biotech and industrial perspective. *npj Vaccines.* 2020;5(1):3-8.