

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Role of Vitamin E in Boosting the Immunity from Neonates to Elderly

Mariyappan Kowsalya, Mohan Prasanna Rajeshkumar, Thangavel Velmurugan, Kattakgounder Govindaraj Sudha and Saheb Ali

Abstract

The vitamin E is a fat-soluble vitamin which occurs as a tocopherol component abundant in humans. The vitamin E supplements in humans and animals have provided numerous health benefits. The vitamin E is rich in antioxidants which slow the aging process and reduce the free radical damage. Vitamin E isoforms play an important role in respiratory health. It is also important in health and well-being of preterm neonates. Vitamin E deficiency in new born includes hemolytic anemia, disease of retina, bronchopulmonary dysplasia. Further, in vitro studies, vitamin E has increased the oxidative resistance and prevents the atherosclerotic plaque. The consumption of vitamin E rich foods reduces coronary heart diseases. This chapter focuses on the treatment of vitamin E deficiency in preterm babies and the role of vitamin E in preventing coronary heart diseases.

Keywords: Vitamin E, neonates, immunity, preterm infants, α -tocopherol

1. Introduction

Vitamins are defined as “organic compounds required in diet in little quantity to execute specific biological functions for normal protection of ideal growth and health of the organism” [1]. In 1922, vitamin E was discovered by Evans and Bishop named it as “X-factor” and got its name after the classification of other vitamins [2]. Vitamin E is generally used as a common word for four tocopherols (α , β , γ and δ) and tocotrienols (α , β , γ and δ) present in food. The main component of the group of compounds in vitamin-E is α -tocopherols. The tocopherols are products of 6-hydroxy chromane (tocol) ring with isoprenoid side chain. Vitamin E with the aid of selenium prevents the non-enzymatic oxidation of cellular components and free radical formation. Vitamin is lipophilic in nature and present on association with derivatives of lipids and cell membranes [1]. The synthetic α -tocopherol is called as *all-rac*- α -tocopherol and naturally available form is RRR- α -tocopherol [3]. The key characteristics of vitamin-E was identified as scavenger of free radical and it is the foremost fat-soluble vitamin responsible for protecting cell membranes against peroxidation [4]. In early stages of life, vitamins are extremely important. Vitamin -E supplies the essential antioxidant protection and stimulates the development of the




Figure 1.
The low level of serum α -tocopherols was observed in certain disease was mentioned which cause high rate of morbidity in neonates. And also, elderly patients have diseases associated with low level of vitamin E and other micronutrients in their diet.

immune system in new-borns [5]. Our human system does not synthesize Vitamin E and it is obtained through dietary intake. Vitamin E rich foods are wheat germ oil, sunflower oils, soybean oil, sunflower seeds, cotton seed, walnut, hazel nuts, peanut butter, corn, palm, spinach, broccoli, kiwi fruit, mango, raspberries, blackberries, blackcurrant, avocado and tomato [6–8]. The vitamin E deficiency was known as the cause of foetal death. Early high dose of vitamin E either through intravenous or intramuscular route decreases the chance of hemorrhage, bronchopulmonary dysplasia, haemolytic anemia, retrolental fibroplasia and retinopathy of prematurity [9]. The low level of vitamin E in serum was observed in certain diseases which was depicted in **Figure 1**.

The antioxidant potential of vitamin E could protect the polyunsaturated fatty acids in the membrane from oxidation, regulating the production of reactive oxygen species, reactive nitrogen species and modulates signal transduction [10]. The effect of antioxidant activity of vitamin E was based on the number of methyl group in its chromane ring. The α -tocopherol have three methyl group whereas the δ -tocopherol has only one methyl group in it [2]. Vitamin E also has anti-cancer potential by stimulating the p⁵³ gene, down regulation of mutant p⁵³ gene and activates the heat shock proteins. Production of PKC and collagenase was inhibited by α -tocopherol. In milieu, γ -tocopherol has effective anti-cancer activity than the α -tocopherol [5]. Intrauterine growth restriction (IUGR) is one of the major causes of neonatal morbidity and mortality. Some studies shows that α -tocopherol aids in the intrauterine development of foetus. 15 million babies are born premature every year and they need special care just to stay alive [11]. During gestational period, the maternal oral intake of vitamin E increases the weight of foetus [12]. The oral administration of vitamin E was given to children to combat malabsorption disorders. Some of the compounds used as vitamin E therapy in newborn and preterm infants are α -tocopherol, tocopherolsol, dl- α -tocopherol, dl- α -tocopheryl acetate [13]. In 2000, the institute of medicine (IOM) chose the hydrogen peroxide induced




Figure 2.
Vitamin E supplementation has many biological activities and it enhances the immune response.

erythrocyte haemolysis test as a marker to determine the vitamin E status. Because, erythrocyte fragility and anemia with higher erythrocyte was observed in children. In adults, vitamin E deficiency occurs due to genetic disorders in α -tocopherol transfer protein (TTP) which causes ataxia. Further, fat malabsorption also leads to vitamin E deficiency due to genetic disorder in TG transfer protein [14]. The people with ataxia are provided with 800-1200 mg of vitamin E as a supplement to prevent the progression of disease [6]. Orally, vitamin E was given in the form of dl- α -tocopherol acetate. Supplementation of α -tocopherol increases insulin sensitivity and decreases the oxidative stress [15]. Oxidative stress was due to the disturbance in balance between generation of free radicals and their exclusion by free radical scavenging activity of the system [16]. Vitamin E supplementation has enhanced the host defense system by heightening the humoral and cell mediated immune system. In elderly people, vitamin E has boosted the resistance against viral diseases and also stimulates immune response to distinctive antigen [5]. Immunostimulatory property of Vitamin E provides enhanced resistance to many pathogens [9]. Earlier studies show that vitamin E possess neuroprotective activity in both foetus and adult rats [17]. The vitamin E supplementation has numerous biological activity which was depicted in the **Figure 2**.

2. Immunological functions of vitamin E in different stages of life

2.1 Preterm infants

Preterm birth was defined as parturition prior to 37 weeks of gestation, the leading cause of mortality and morbidity in neonates. The rate of prematurity complications increases with decreased gestational age and birth weight [10]. New-borns are one of the risk groups of vitamin E deficiency. The limited transfer of α -tocopherol through the placenta which results in low level of vitamin-E in tissues and serum at the time of birth in premature infants than in full term infants [18]. The preterm infants' blood vessels are exposed to high oxygen tension due to the deficient amount of vitamin E [19]. Preterm infants are born with low body weight and have less stored fat-soluble vitamins. The level of α -tocopherol is low at their time of birth was shown by the elevated erythrocyte haemolysis in the presence

of hydrogen peroxide. The enteral dose of vitamin E is given at 50 IU/kg within 4 hours of birth and it increases the serum α -tocopherol level [20]. The preterm infants those less than 1000 g in NICU were given 1.5 ml of vitamin E per day through infuvite pediatric as a portion of parental nutrition. And infants in 1000 g to 3000 g obtain 3.5 ml of vitamin E per day. Each vial (1 ml) of infuvite encompasses 7 mg of α -tocopheryl acetate [3].

The preterm infants administering the oral dose of vitamin E helps in treating the vitamin E deficiency syndromes like anemia, retinopathy, thrombocytosis and congenital malformation [21]. The retrolental fibroplasia occurs often in preterm infants because the last portion of the retinal blood vessel continues to develop till the end of eight month of gestation. The blood vessels of the preterm infants might not be developed well and they need an appropriate environment with arterial oxygen tension, blood supply, nutrient, disposal of wastes and exposure to visible light [18]. The haemolytic anemia was frequently observed in preterm infants, where their intake of fortified iron is higher along with linolenic acids. Nowadays, the formula for preterm babies had increased the vitamin E and reduces their iron content so that this helps to prevent the development of haemolytic anemia [22]. The vitamin E supplements in preterm infants increases the hemoglobin level after 8–10 weeks of birth [23]. Previous studies reported that preterm infants born below 2000 g, when supplemented with 16.5 mg of α -tocopherol acetate per day for about ten days has improved the hemoglobin level and decreases the reticulocyte count. Supplementing 16.5 mg concentration of vitamin E is higher in their study compared to regular dose of 1.5 mg/day but it shows betterment of disease condition, however their consequences have to be studied [24].

The bronchopulmonary dysplasia (BPD) was observed among preterm infants due to free radicals which injures the lung cells. It was generally noticed in preterm infants particularly born before 28 weeks of gestation. Oral supplementation of vitamin E is tolerable in pregnancy and infancy. They are present routinely in the total parenteral nutrition (TPN) which is used in the treatment of BPD. And also, the α -tocopherol plays a key role as anti-inflammatory factor in neonatal lungs [15]. A study with 133 preterm infants were grouped into 68 preterm infants with confirmed case of BPD, 65 preterm infants without BDP. It was observed that as the condition of BPD increases with vitamin E deficiency. The involvement of vitamin E in BDP was confirmed and their supplementation would become a therapy to BPD [25].

2.2 Neonates

The adequate intake of Vitamin E is essential in neonates as they are born with low stores of it. The routine recommended intake of vitamin E is 2.8 IU/kg per day (Maximum 7 IU/kg per day) [8]. The term babies gestational age was between 37 and 41 weeks and their weight at the time of birth was 2500 g -3700 g [26]. It was observed that very low birth weight in infants were deficient in Vitamin E. Hence, vitamin E supplementation is required to enhance the weight gain in infants [27]. The yellow coloration of the skin and sclera of new born babies are referred to as neonatal jaundice which is the result of bilirubin deposition. Vitamin -E is supplemented along with phototherapy of the full-term neonates [28]. In an extrauterine environment, vitamin E acts as defense against oxygen toxicity and their placental transfer to the foetus is limited during gestational period. Hence, maternal milk is important to supply this vitamin to the neonates in their initial period and during lactation, thus, protecting it from haemolytic anemia, bronchopulmonary dysplasia, neurological dysfunction and increased neonatal mortality. The studies show that vitamin E in maternal milk decreases with increase in lactation period i.e., colostrum ($40.5 \pm 15 \mu\text{mol/L}$) provides the highest content of vitamin E to infants.

Whereas it gradually decreases in transitional ($13.9 \pm 5.2 \mu\text{mol/L}$) and mature ($8.0 \pm 3.8 \mu\text{mol/L}$) milk of lactation. The higher concentration of alpha-tocopherol in colostrum acts as complementary mechanism [29]. Hyperbilirubinemia was observed in neonates which was treated with phototherapy and vitamin E supplementation [30]. The world health organization WHO has issued a global public health recommendation that infants should exclusively breastfed for 6 months to attain optimal growth and development as the maternal milk was the infant's best source of nutrients. The average amount of vitamin E in maternal milk is 2.5–2.9 mg it contributes to prevention of free radical propagation in numerous lipid structures within the system [31]. The Dietary Reference Intake (DRI) recommends the intake of 4 mg/day of vitamin E for children between 0 and 6 months. The vitamin E supplementation in children is essential to the development of the immune system, lungs, extracellular matrix of the vascular system and mental development [29].

2.3 Children

Vitamin E enhances the immune responses of the children. Cardiomyopathy was observed as a symptom in children with critical deficiency of vitamin E. The prescribed amount of vitamin E supplementation in children is 1000 mg/d [13]. Low level of maternal plasma α -tocopherol increase the chance of asthma in children within their first ten years [32]. The antioxidant intake of vitamin E during pregnancy and their effect on development of wheezing and eczema in children is needed to be confirmed with the follow up investigation on the children's growth and their dietary intake [33]. In previous report, the association of vitamin E intake during pregnancy and chance of asthma in children was observed in 2000 pregnant women. The α -tocopherol level in plasma was examined during gestation and cord blood after delivery. Their intake of anti-oxidant vitamins are also examined. Then follow up studies was performed up to five-year age of children. The symptoms of asthma, wheezing and dietary intake was collected for 1253 children and 797 children were participated in hospital evaluation. They have concluded that low level of serum vitamin E during pregnancy was observed among the children with phenotypic symptoms of asthma. Though, vitamin E does not directly improve the conditions of asthma patients their deficiency in early stage of life was perceived in children be inflicted with asthma [34].

The children suffering from chronic kidney disease are at high risk of micro-nutrient deficiency. The concentration of vitamin A and vitamin B12 was within the range of reference while vitamin E shows major changes as the kidney disease worsens hence advance studies are essential to determine the required concentration of vitamin supplements in infants and children with this disease [35]. In children, respiratory tract infection is common cause of the morbidity. The vitamin E supplementation helps to treat this disease. A substantial amount of plasma vitamin E improves the hemolytic uremic syndrome in children. Which was due to the defect in production of prostacyclin and thus preventing the mortality and morbidity rate [36]. Supplementation of vitamin E (100 IU/ day) has increased the glutathione level and decreased the lipid peroxidation and concentration of HBA1C in the erythrocytes of type 1 diabetics children [37]. The childhood obesity was growing and it may be due to imbalance in the oxidant and antioxidant level. The obese children were observed with increased level of oxidants like isoprostanes and decreased level of antioxidant like vitamin E. Oxidative stress markers were decreased by vitamin E supplementation in obese children with liver steatosis. Hence, early implementation of vitamin E will reduce the risk of cardiovascular and metabolic changes connected with non-alcoholic fatty liver disease in children and adolescence [38].

2.4 Adults

The reactive oxygen species include (ROS) like hydrogen peroxide, hydroxyl radicals, and superoxide were studied exclusively. The formation of ROS during electron transport chain was presumed due to increased ingestion of oxygen by mitochondria in the cell. Some recent studies reported that exercises increased the ROS production. The studies revealed that vitamin-E supplementation in athlete improved the muscle performance and protects the cell membrane from damage through oxidative stress. This study augment that vitamin E would strengthen the skeletal muscle of the older people [39].

In the previous research, Women with the possibility of pre-eclampsia during pregnancy was provided with 400 IU of vitamin E and 1 g of vitamin C per day. As a result, very low birth weight infants were born to supplemented group compared to placebo group. Hence, the vitamin E concentration should not be exceeded above the required dietary allowances [40]. Furthermore, the greater the concentration of serum vitamin E during pregnancy then there is a chance of macrosomia [41]. Recently, a study with 897 girls in their adolescence found that vitamin E aids in the inflammatory response and oxidative stress during menstrual cycle [42].

Cystic fibrosis was one the genetic disorder which leads to deterioration of pulmonary functions in children and adults, partially due to oxidative stress. The vitamin E supplementation greater than required dietary allowances has enhanced the serum α -tocopherol [43]. In USA, the higher acceptable level of vitamin E per day was 1000 mg/day [44].

2.5 Elderly

Vitamin E group has tocopherol and tocotrienol groups each with identical isomers but with alteration in hydrophobic tridecyl chain saturation point. The tocotrienol was recently discovered to be more effective than tocopherol in the treatment of age-related cardiovascular diseases [45]. The low intake of vitamin E in childhood and adulthood leads possibly to hypertension than those who have taken an adequate amount of vitamin E. The cardiovascular disease risk can be decreased by α -tocopherol which inhibits LDL oxidation and reduces the inflammatory responses [46]. The intake of antioxidants could prevent the lipid peroxidation [47]. The epidemiological study with 27271 men who were smokers and by no records of myocardial infraction in the age group 50–69 was divided into group and supplemented with vitamin E, placebo and beta-carotene. The groups were keenly monitored to detect the possible effect of vitamin E supplementation in cardiovascular disorder. The results revealed that 50 mg/day supplementation of vitamin E has prevented the occurrence of nonfatal myocardial infraction by 4% and fatal coronary heart disease by 8%. The vitamin E (α -tocopherol) supplemented group has lessened the chance of coronary heart disorder in human trails followed up to six years related to the placebo group. The beta-carotene supplementation has no effect on non-fatal myocardial infraction disease [48].

Immunological function especially cell-mediated immune response will decrease with age and hence elders are prone to infectious diseases. The 200-800 mg supplementation of vitamin E enhances the production of antibody to a primary immunization [49]. In the in vivo study, healthy elderly people in the age group greater than 60 years are supplemented with 800 mg of vitamin E per day and the fasting blood sample was collected for the duration of 6 months and this study has evidenced the increase of α -tocopherol level in serum and increase in delayed

type hypersensitivity was observed [50]. In four month of clinical trials, vitamin E supplementation greater than RDA in elderly people had enhanced production of antibodies against various diseases like tetanus, hepatitis B and DTH vaccines [51].

The oxidative stress was one of the reasons for neurodegenerative diseases like Alzheimer's disease, Parkinson's disease and processes in cells related to aging. The brain cells were easily damaged by free radicals due to its consumption of high amount of oxygen along with certainly peroxidable lipid membranes and comparatively lesser enzymes with antioxidant potential. The vitamin E and vitamin C were the antioxidant rich supplements which will scavenge the reactive oxygen species and helps in regeneration of neurons [52]. Vitamin E also protects the nervous system with aging [13]. Neurological abnormalities are observed in children and adults when the serum α -tocopherol level is less than 0.5 mg. Vitamin E deficiency manifests as neuropathic and myopathic disorders. Spinocerebellar syndrome, ataxia, hyporeflexia, vibratory sensation was commonly observed clinically. Moreover, skeletal myopathy and pigmented retinopathy also occurs due to the vitamin E deficiency [6]. Further, Vitamin E supplementation has upgraded the cognitive function and vascular dementia in elderly [53]. The higher level of α -tocopherol in the brain generates an anti-inflammatory milieu to decrease the density of microglial cells. It also protects from the Alzheimer's disease [54]. A study group involved 39,876 women in US, in which 6377 women were greater than 65 years. It was evidenced that vitamin E supplementation in 600 IU alternative days with the follow up for 4 years had enhanced cognitive development. Because the oxidative stress was the major cause for dementia pathogenesis and numerous reports proved that vitamin E has decreased the lipid peroxidation in brain and also prevent the occurrence of Alzheimer's disease [55].

Previous studies reported that intervention of Vitamin E improves the lymphocyte proliferation and delayed type hypersensitivity was enhanced with higher production of IL-2 and lesser production of IL-6 [9]. An intervention of 200 mg of vitamin E has enhanced immune response of elderly people [56]. When the older mice supplemented with vitamin E has enhanced the cell mediated immune response, production of IL-2 and delayed type hyper sensitivity. In elder human subjects also vitamin E supplementation progresses both the innate and adaptive immunity. The phagocytic capability of leucocyte was improved but declined the bactericidal activity which might be due to the antioxidant potential of vitamin and lesser production of hydrogen peroxide. Further, the optimal concentration of vitamin E supplementation is 200 IU per day than the 60 and 800 IU per day of vitamin E supplementation has increased the T-cell proliferation in elder people [57].

In vivo studies of both animals and human have evident that immunity decreases with aging process. It was observed by decreased antibody response, delayed-type hypersensitivity, proliferation of T cell in response to mitogens. Research has proved that antioxidants have helped to improve the immunity in aging process [50]. Aging is a gradual and typical loss of the physiological system along with its immune response. The elderly people are prone to infections and other diseases like cancer due to the age related decrease of immunity. One of the widely consented reason for aging was oxidative stress. The previous study suggested that supplementation of both vitamin C (500 mg/day) and vitamin E (200 mg/day) to the elderly people has improved their immune function by enhancing the humoral and cell-mediated immune response. The α -tocopherol acts as immunomodulator and enhances the cytokine levels in older population of group. Level of INF- γ has increased in tested groups of older people supplemented with vitamin E [58].

S. No	Groups	No. of Participants	Age	Disease Condition	Dosage form of Vitamin E	Quantity	Outcome	Reference
1	Preterm Infants	12	28–30 week gestation	Very low Birth weight	dl-alpha-tocopheryl-acetate	3.5 mg/dl	Reduce the risk of Retinopathy	[62]
2		215	28–30 week gestation	Very low Birth weight	Soybean oil-based lipid emulsion	—	Reduce cholestasis	[63]
3		34	≤ 35 week gestation	Very Low Birth Weight	alpha-tocopheryl acetate	16.5 mg/day	Increases Hemoglobin concentration	[64]
4		168	≤ 30 week gestation	Very Low Birth Weight	Vitamin E	3.3 mg/day	Prevent retrolental fibroplasia	[65]
5		25	< 30 week gestation	Very Low Birth Weight	dl-alpha-tocopheryl acetate	25 mg Ephynal	Enhances serum alpha-tocopherol	[66]
6		36	25.5 week gestation	Very Low Birth Weight	Mixed vitamins in parenteral solution	3.17 mg/day	Increased serum alpha-tocopherol	[67]
7		151	< 35	Very Low Birth Weight	alpha- tocopherol	50 mg/kg	Decreases the bilirubin level	[33]
8	Neonates	77	37–42 week gestation	—	RRR-alpha- tocopheryl acetate; all-rac-alpha-tocopheryl acetate	20 IU; 13.5 IU	Infants discriminates natural and synthetic vitamin E	[15]
9		80	—	Hyperbilirubinemia	Phototherapy & Vitamin E	4 mg/day	Faster Recovery	[23]

Table 1.
Role of vitamin E in improving immune system in various disease condition of preterm infants and neonates.

S. No	Groups	No. of participants	Age	Disease Condition	Dosage form of Vitamin E	Quantity	Outcome	Reference
1	Children	2372	≤ 5 years	Kwashiorkor	Vitamin E, riboflavin, Se	—	No data on morbidity	[68]
2		16	—	Chronic Cholestasis	d- alpha -tocopheryl polyethylene glycol 1000 succinate (TPGS)	20–25 IU	Improves neurological development	[69]
3		61	≤10 years	Acute pyelonephritis	Vitamin E	—	Reduces Renal scarring	[70]
4		141	≤ 14.5 years	Cystic fibrosis	Vitamin E	—	Enhances vitamin E level	[71]
5	Adults	716	Gestation period	HIV infected	alpha -tocopherol acetate	30 mg/day	Improves vitamin status in infants	[72]
6		23	—	Oxidative stress	Vitamin E	400 IU	Prevents exercise induced oxidative stress	[73]
7	Elders	184	—	Cardiovascular disease	alpha -tocopheryl acetate	400 IU	Reduces lipid per oxidation	[74]
8		29133	50–69	Pneumoniae	alpha -tocopherol	50 mg/day	Decrease the risk	[75]

Table 2.
Role of vitamin E in improving immune system in various disease conditions of children, infants and elderly.

3. Immunological functions of vitamin E in infections

3.1 Pneumoniae

The in vivo studies on older mice affected with pneumonia was supplemented with vitamin E at the concentration of 500 mg per kg for four weeks and found that it has help to improve the lung functions. It was observed that migration of neutrophil and production of inflammatory cytokines was reduced with the intake of vitamin E. In human subjects, the older people supplemented with 200 IU of vitamin E per day has subsided the necessary to re-hospitalization of older people with pneumoniae up to 63% and it aids in faster recovery [57].

3.2 Human immunodeficiency virus

The vitamin E possess the anti-inflammatory property and people infected with HIV was found be lack of it and weakened immune system. Vitamin E supplementation in 400 IU per day has reinstate the delayed type hypersensitivity, production of IL-2 and T_H cells. The higher level of α -tocopherol in serum had blocked the advancement of infection. The murine model with HIV infection supplemented with increase in fifteen-fold of dietary vitamin E regularize the distorted immune system caused by infection to normal state [5]. Further, there was greater incidence of non-alcoholic steatohepatitis in HIV infected patients. The previous studies with 27 HIV patients with NASH reported that vitamin E supplementation was a most effective treatment as it has enhanced the ALT level, ck-18 and CAP score. Moreover, it does not cause any adverse effects in participated individuals [59].

3.3 Influenza virus

Influenza virus cause severe damage to lungs and also inflammation leads to greater oxidative stress. The vitamin E acts as an effective antioxidant therapy in influenza diseases. Thus, vitamin E supplementation protects the respiratory system and prevents the occurrence of oxidative damage due to influenza [60].

3.4 Others

The mouse models were infected with coxsackievirus which induces myocarditis. And in vitamin E deficient group the virulence of this virus is greater. In human studies, the coxsackie virus was isolated from infected individual, which was called as keshan an endemic disease commonly observed among children and women. The affected individuals were deficient in vitamin E and Se hence their supplementation in diet can prevent the viral infection [61]. The role of vitamin E in boosting immune system in various diseases in infants, children and elderly are depicted in **Tables 1 and 2.**

4. Discussion

The vitamin E was present in many natural foods which should be taken in adequate amount because the fortified or supplementation of vitamin E does not provide greater health benefits. The natural form of vitamin E rich foods should be incorporated into the diet. The healthy foods should be added to the diet for the efficient action of antioxidants in the human system [47]. The mother's milk

especially colostrum contains higher concentration of vitamin E which prevents the infants from oxidative damage and develops their immune system. Since, very less amount of vitamin E is transferred through the placenta, maternal milk plays a crucial role in enhancing the serum α -tocopherol level in new born [76]. The babies born with lesser gestational age was highly prone to oxidative stress and the maternal milk is the best source of antioxidants than the formulas to protect the new born infants [77]. Still today, α -tocopherol level in the serum was frequently measured through HPLC analysis. A novel method should be developed for their measurements.

The preterm infants were susceptible to oxygen radical disease and it can treat with antioxidant therapy in which the well-known antioxidant nutrient vitamin E can be used to treat this condition in preterm infants [78]. The formula fed preterm infants have the risk of developing high oxidative stress but in the study with 31 healthy preterm infants shows that long chain poly unsaturated fatty acid supplemented group does not affect the solubility of α and γ - tocopherol [79]. Another research also suggest that both the breast fed and formula fed preterm infants possess the ability to tolerate oxidative stress. It was confirmed by the presence of malonaldehyde (MDA) in the urine which was measured by HPLC analysis [80]. The vitamin E supplementation is required in preterm infants to improve certain disease conditions like haemolytic anemia, retrolental fibroplasia, bronchopulmonary dysplasia but their long-term high dosage leads to sepsis, necro colitis and in some cases even death of premature infants. Hence the ideal dosage of vitamin E is prerequisite to treat the disease in preterm infants on the other hand to ensure the safety and longer healthy life to the preterm babies [81].

In children, vitamin E deficiency results in the development of chronic cholestasis. In this case, the children have normal serum α -tocopherol level but decreased ratio to the total lipid content [82]. The vitamin E deficiency could be combated by intaking fortified foods can enhance the vitamin E and it was one of the best methods to reach daily requirements of vitamin E [83]. Vitamin E supplementation along with other micronutrients and trace elements increases our defense barrier system and prevents the development of infection [84].

The oxidative stress in the human body was mainly originated from the free radicals produced by mitochondria and other cellular components. The external factor for oxidative stress includes UV light rays from sun, ozone, pollutants, smoke from cigarette and smog. These factors contribute to the aging process. The antioxidant rich foods would help to balance the oxidant- antioxidant level in human body. The foods enriched with vitamin E are a good choice to delay the aging process and to have healthy and youthful glowing skin in later stages of life. [85].

Further, vitamin E supplementation also improves the cognitive function when their dietary intake is higher at the earlier stage of Alzheimer's disease. And also in the later stages of this disease their admission to centre is greatly reduced but it does not improve any cognitive functions [52]. The vitamin E is an effective antioxidant which shows beneficial report in preventing the progression of Alzheimer's disease, Parkinson's disease and dementia [55]. Though, there was no significant cognitive improvement in healthy individuals supplemented with vitamin E, it helped in the diseased condition We suggest that vitamin E has positive impact in improving the brain disorders associated with oxidative damage of brain cells.

Vitamin E was one of the efficacious nutrients which could modulate the immune system. The concentration of vitamin E is higher in immune cells than other cells in the blood. It has been observed that deficiency of vitamin E worsens the immune system in both animals and humans. Thus, vitamin E supplementation exceeding the required dietary recommendation has contributed to enhance the immune system. It has intensifies the differentiation and proliferation of T cell,

production of IL2, activity of T_H cells, macrophages and phagocytic cells [57]. The vitamin E supplementation in elderly has improved their immune response towards improved antibody response, delayed type hypersensitivity and also T-cell was proliferated was stimulated in response to mitogens. And Vitamin E in combination with vitamin C has also shows an enhanced immune response in elderly group than with the study group administered with vitamin E alone [50]. The higher level of vitamin E (α -tocopherol) in serum the greater the ability to resist viral infection in elder population [5]. Furthermore, the dietary intake of vitamin E enhances the immunity in elders and immunocompromised persons [86].

The vitamin E supplementation has modulated the immune system in both direct and indirect way. Directly it has maintained the cellular integrity and protected the cells from damage caused by oxidative stress. While indirectly it has aid the modulation of inflammatory intermediaries like proinflammatory cytokines and prostaglandin E2. The studies in mice suggests that inflammatory lung disease can be treated with the combination of probiotic strain *Bifidobacterium lactis* and vitamin E, C to lessen the lung inflammation due to air polluting agents [87]. In animal models, the α -tocopherol possess the anti-inflammatory and γ - tocopherol with pro-inflammatory property it could be used in the treatment of asthma [88].

The elderly people in the age group of 65 years and above was investigated for the association of vitamin E supplementation and respiratory tract infection. The study results shows that the vitamin E supplantation does not show any statistical difference among the lower respiratory infection among the supplemented and placebo group. However, the vitamin E supplementation group has a remarkable result in preventing the upper respiratory tract infection and against common cold [89]. Acute and chronic lung injuries was observed in new-borns due to oxidative damage. Usually, surfactant lipids protect the type II alveolar cells of lungs from air-borne pathogens. The vitamin E supplementation enhance surfactants and prevent the development of lung diseases like bronchopulmonary dysplasia [90]. Thus, based on these results vitamin E supplementation in our diet would prevent the upper respiratory tract infection and strengthens the lungs alveolar cells which might help us to combat the Covid-19 infection and help us to build a stronger immunity in current pandemic situation.

5. Conclusion

In past decades, vitamin E deficiency was frequently observed in preterm infants and neonates leading to various diseases like bronchopulmonary dysplasia, retrolental fibroplasia, hyperbilirubinemia, haemolytic anemia. Most often, the very low birth weight infants are at the high risk of developing vitamin E deficiency. The adequate amount of vitamin E supplementation has prevented the development of these disease conditions. The alpha tocopherol and alpha tocopheryl acetate are the most common form of vitamin E supplemented to preterm infants and new born infants. Vitamin E supplementation has prevented the development of asthma in children but not in patients with chronic severe asthma. Hence, the mechanism of vitamin E involved in preventing the disease in juvenile stage in children is need to be investigated. Vitamin E also possess numerous health benefits along with its antioxidant property. The aging process decreases the immunological response and increases the chances of infection in elder population. Thus, vitamin E supplementation has improved the T-cell mediated immune response and prevented the progression of Parkinson's disease and dementia in older people. Further, vitamin E is an important micronutrient which plays a crucial role in the early stage of our life to lead a healthy life. Many reports proved that vitamin E supplementation

has enhanced the immunity and prevented many diseases in infants. However, the optimal dosage amount, isoform and duration of vitamin E supplementation in each disease conditions in preterm infants and newborn is still need to be validated. Indeed, the mechanism behind the vitamin E in curing the infectious diseases and improving the immune response is little-known and future research will bring to light the unknown mechanism of vitamin E in boosting immune response in infants.

Conflict of interest

The authors declare no conflict of interest.

Author details

Mariyappan Kowsalya, Mohan Prasanna Rajeshkumar*, Thangavel Velmurugan, Kattakgounder Govindaraj Sudha and Saheb Ali
K. S. Rangasamy College of Arts and Science (Autonomous),
Tiruchengode, Tamil Nadu, India

*Address all correspondence to: prasanna4d@gmail.com

IntechOpen

© 2021 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

References

- [1] Satyanarayana U, Chakrapani U. Essentials of Biochemistry. 2nd edition. Books and Allied P Ltd., 57-65p. ISBN: 81-87134-82-8.
- [2] Miyazawa T, Burdeos GC, Itaya M, Nakagawa K, Miyazawa T. Vitamin E: Regulatory Redox Interactions. IUBMB Journal. 2019. <https://doi.org/10.1002/iub.2008>.
- [3] Han SN, & Meydani SN. Impact of vitamin E on immune function and its clinical implications. Expert Review of Clinical Immunology. 2006; 2(4), 561-567. doi:10.1586/1744666x.2.4.561.
- [4] Gomez-Pomar E, Hatfield E, Garlitz K, Westgate PM, Bada HS. Vitamin E in preterm infant: A forgotten cause of hemolytic anemia. American Journal of Perinatology. 2017. Doi: <http://10.1055/s-0037-160728>.
- [5] Lima MSR, Dimenstein R, Ribeiro KDS. Vitamin E concentration in human milk and associated factors: a literature review. J. Pediatr (Rio J). 2014; 90 (5), 440-480.
- [6] Rizvi S, Raza ST, Ahmed F, Abbas S, Mahdi F. Role of vitamin E in human health and some diseases. SQU Medical Journal. 2014; vol-14, Issue2, e157-165.
- [7] Morsy TA, and Alanazi AD. A mini-overview of vitamin E. J. Egypt. Soc. Parasitol. 2020; 50(2), 247-257.
- [8] How does vitamin E supports your immune system? [Internet] <https://betteryou.com/blogs/health-hub/vitamin-e-immune-system-support>.
- [9] Andrea SV. Effects of Vitamin E in neonates and young infants. International Journal of pediatrics. 2016; 4 (5), 1745-1757. 10.22038/IJP.2016.6736.
- [10] Lee GY and Han SN. The role of vitamin E in Immunity. Nutrients. 2018; 10(11): 1614. Doi: 10.3390/nu10111614.
- [11] Falsaperla R, Lombardo F, Filasco F, Romano C, Saporito MAN, Puglisi F, Piro E, Ruggieri M, Pavone P. Oxidative Stress in Preterm Infants: Overview of Current Evidence and Future Prospects. Pharmaceuticals. 2020; 13(7), 145. Doi: 10.3390/ph13070145.
- [12] Parraguez VH, Sales F, Peralta OA, Narbona E, Lira R, Reyes MD, Gonzalex-Bulnes A. Supplementation of underfed twin-bearing Ewes with herbal vitamic C and E: Impacts on birth weight, postnatal growth and pre-weaning survival of the lambs. Animals. 2020; 10,652. Doi: 10.3390/ani10040652.
- [13] Westergren T, and Kalikstad B. Dosage and formulation issues: Oral vitamin E therapy in children. 2010; 66, 109-118. Doi: 10.1007/s00228-009-0729-1.
- [14] Traber MG. Vitamin E inadequacy in humans: causes and consequences. American society for nutrition. Adv. Nutr. 2014; 5, 503-514. Doi: 10.3945/an.114006254.
- [15] Du O, Luo Z-C, Nuyt AM, Audibert F, Julein P, Wei S-Q, Zhang D-L, Fraser W, Levy E. Vitamin A and E Nutritional status in relation to leptin, adiponectin, IGF-I and IGF-II in early life – a birth cohort study. Scientific reports. 2018; 8, 100. Doi: 10.1038/s41598-017-18531-3.
- [16] Stone CA, McEvoy CT, Aschner JL, Kirk A, Ross-salazar C, Cook- Mills JM, Moore PE, Walsh WF, Hartert TV. Update on vitamin E and its potential role in preventing or treating bronchopulmonary dysplasia. Neonatology. 2018; 113, 366-378. Doi: 10.1159/000487388.
- [17] Erdemli Z, Erdemli ME, Turkoz Y, Yigitcan B, Aladag MA, Cigremis Y, *et al.* Vitamin E effects on developmental disorders in fetuses and cognitive

dysfunction in adults following acrylamide treatment during pregnancy. *Biotechnic & Histochemistry*, 2020; 1-9. doi:10.1080/10520295.2020.1751880.

[18] Silva AGCL, Reboucas AS, Mendonca BMA, Silva DCN, Dimenstein R, Ribeiro KDS. Relationship between the dietary intake, serum, and breast milk concentrations of vitamin A and vitamin E in a cohort of women over the course of lactation. *Maternal and Child Nutrition*. 2019; <https://doi.org/10.1111/mcn.12772>.

[19] Johnson L, Schaffer D, Boggs TR. The premature infant, vitamin E deficiency and reteolental fibroplasia. *The ammerican journal of clinical nutrition*. 1974; 1158-1173.

[20] Bell EF and Filer LG. Role of vitamin E in the nutrition of premature infants. *The American Journal of clinical nutrition*. 1981; 414-422.

[21] Ribeiro KD, Lima MSR, Medeiros JFP, Reboucas AS, Dantas RCS, Bezerra DS, Osorio MM, Dimenstein R. Association between maternal Vitamin E and alpha tocopherol levels in the new born and colostrum. *Maternal & Child Nutrition*. 2016; 12, 801-807. Doi: 10.1111/mcn.12232.

[22] Bieri JG, Corash L, Hubbard VS. *Medical Uses of Vitamin E*. New England Journal of Medicine. 1983; 308(18), 1063-1071. doi:10.1056/nejm198305053081805.

[23] Abrams BA, Gutteridge MC, Stocks J, Friedman M, Dormandy TL. Vitamin E in neonatal hyperbilirubinaemia. *Archives of Disease in childhood*. 1973; 48, 721.

[24] Jansson L, Holmberg L, Nilsson B, Johansson. Vitamin E requirements for preterm infants. *Acta paediatrica*. 1978; 67 (4), 459-463. <https://doi.org/10.1111/j.1651-2227.1978.tb16354.x>.

[25] Ge H, Liu W, Li H, Zhang M, Liu C, Qiao Y. The association of vitamin D and vitamin E levels at birth with bronchopulmonary dysplasia in preterm infants. *Pediatr Pulmonol*. 2021; 1- 6. <https://doi.org/10.1002/ppul.25414>.

[26] Baydas G, Canatan H, Turkoglu A. Comparative analysis of the protective effects of melatonin and vitamin E on streptozocin-induced diabetes mellitus. *JPR*. 2002; <https://doi.org/10.1034/j.1600-079X.2002.01856.x>.

[27] Kositamongkol S, Suthutvoravut U, Chongviriyaphan N, Feungpean B and Nuntnarumit p. Vitamin A and E status in very low birth weight infants. *Journal of Perinatology*. 2011; 31, 471-476.

[28] Al-Banna SM, Riad AN, Anes SS. Fenofibrate and antioxidant vitamins [D, E, and C] as novel approach in treatment of uncomplicated neonatal hyperbilirubinemia. *MJMR*. 2019; 30 (3), 206-210.

[29] Silva ALC, Ribeiro KDS, Rayanne L, Melo M, Bezerra DF, Queiroz JLC, Lima ASR, Pries JF, Bezerra DS, Osorio MM, Dimenstein. Vitamin E in human milk and its retention to the nutritional requirement of the term newborn. *Rey Paul Pediatr*. 2017; 36(2), 58-164.

[30] El Mashad GM, El Sayed HM, El Refaey NAS. Role of vitamin E supplementation in neonates with hyperbilirubinemia. *Faculty of Medicine, Menouia University*. 2021; IP: 223.182.216.4.

[31] Han J, Kang L, Liang D, Li H, Su Y, Zhang Y, Yang Y. Composition requirements of follow up formula for 6-12 months-old infants: recommendations of a Chinese expert group. *Asia Pac J Clin Nutr*. 2019; 28(2),347-355. <http://dx.doi.org/10.1016/j.jped.2014.04.006>.

[32] Allan KM, Prabhu N, Craig LCA, McNeill G, Kirby B, McLay J, Helms PJ,

Ayres JG, Seaton A, Turner SE and Devereux G. Epidemiology and paediatric asthma. 2015; 45, 1027-1036. Doi: 10.1183/09031936.00102214.

[33] Martindale S, McNeil G, Devereux G, Campbell D, Russell G, and Seaton A. Antioxidant intake in pregnancy in relation to wheeze and eczema in the first two years of life. 2004; 171(2). <https://doi.org/10.1164/rccm.200402-220OC>

[34] Devereux G, Turner SW, Craig LCA, McNeil G, Martindale S, Harbour PJ, Helms PJ and Seaton A. Low maternal vitamin E intake during pregnancy is associated with asthma in 5-year-old children. Am. J. Respir. Crit. Care Med. 2006; 174 (5). <https://doi.org/10.1164/rccm.200512-1946OC>.

[35] Joyce T, Rasmussen P, Melhem N, Clothier J, Booth C, Sinha MD. Vitamin and trace element concentrations in infants and children with chronic kidney disease. Pediatric Nephrology. 2020; 35, 1443-1470. <http://doi.org/10.1007/s00467-020-04536-0>.

[36] Powell HR, McCredie DA, Taylor CM, Burke JR, Walker RG. Vitamin E treatment of haemolytic uraemic syndrome. Archives of disease in Childhood. 1984; 59, 401-404.

[37] Jain SK, McVie R and Smith T. Vitamin E supplementation restores and malondialdehyde to normal concentrations in erythrocytes of type I diabetic children. Diabeted care. 2000; 23 (9), 1389-1394. <https://doi.org/10.2337/diacare.23.9.1389>.

[38] Adamo ED, Marcovecchio ML, Giannini C, de Giorgis T, Chiavaroli V, Chiarelli F & Mohn A. Improved oxidative stress and cardio-metabolic status in obese prepubertal children with liver steatosis treated with lifestyle combined with Vitamin E. Free Radical Research. 2013; 47(3), 146-153. DOI:10.3109/10715762.2012.755262.

[39] Dutra MT, Martins WR, Ribeiro ALA, and Bottaro M. The Effects of Strength Training Combined with Vitamin C and E Supplementation on Skeletal Muscle Mass and Strength: A Systematic Review and Meta-Analysis. Journal of sports Medicine. 2020; Article ID 3505209. <https://doi.org/10.1155/2020/3505209>.

[40] Poston L, Briley AL, Seed PT, Kelly FJ, Shennan H. Vitamin C and vitamin E in pregnant women at risk for pre-eclampsia (VIP trial): randomised placebo-controlled trial. The Lancet. 2006; 367 (9517), 1145-1154. [https://doi.org/10.1016/S0140-6736\(06\)68433-X](https://doi.org/10.1016/S0140-6736(06)68433-X).

[41] Yang W, Jiao M, Xi L, Luo S, Zhou Q, Wang H. The association between maternal fat-soluble vitamin concentrations during pregnancy and infant birth weight in China. The British Journal of Nutrition. 2020. Doi: 10.1017/S0007114520003347.

[42] Bahrami A, Bahrami-Taghanaki H, Khorasanchi Z. *et al.* Menstrual problems in adolescence: relationship to serum vitamins A and E, and systemic inflammation. Arch Gynecol Obstet. 2020; 301, 189-197. <https://doi.org/10.1007/s00404-019-05343-1>.

[43] Woestenenk JW, Broos N, Stellato RK, Arets GM *et al.* Vitamin E intake, α -tocopherol levels and pulmonary functions in children and adolescents with cystic fibrosis. British Journal of Nutrition. 2015; 113, 1096-1101.

[44] Han SN, Adolfsson O, Lee CK, Prolla TA, Ordovas J, Meydani SN. Age and vitamin E-induced changes in gene expression profiles of T cells. J. Immunol. 2006; 177:6052-6061. doi: 10.4049/jimmunol.177.9.6052.

[45] Ramanathan N, Tan E, Loh LJ, Soh BS, Yap WN. Tocotrienol is a cardioprotective agent against ageing-associated cardiovascular disease and its

associated morbidities. *Nutrition & Metabolism*. 2018; 15 (6).

[46] Mishra GD, Malik NS, Paul AA, Wadsworth MEJ, Bolton-Smith C. Childhood and adulthood dietary vitamin E intake and cardiovascular risk factors in mid-life in the 1946 British Birth Cohort. *European Journal of Clinical Nutrition*. 2003; 57, 1418-1425.

[47] Jordao KSLU, Assumpção D, Barros MBA, Filho AAB. Vitamin E intake and food sources in adolescent diet: A cross-sectional population-based study. *Rev. paul. Pediatr*. 2021; 39. <https://doi.org/10.1590/1984-0462/2021/39/2019295>.

[48] Virtamo J, Rapola JM, Ripatti S, Heinonen OP, Taylor PR, Albanes D, Huttunen JK. Effect of Vitamin E and Beta Carotene on the Incidence of Primary Nonfatal Myocardial Infarction and Fatal Coronary Heart Disease. *Arch Intern Med*. 1998;158(6):668-675. doi:10.1001/archinte.158.6.668.

[49] High KP. Micronutrient supplementation and Immune function in the elderly. *Infectious disease society of America*. 1999; 28, 717-22.

[50] Rall L, Meydani SN. Vitamin E, Vitamin C and Immune Response: Recent Advances. Institute of Medicine (US) Committee on Military Nutrition Research. *Military Strategies for Sustainment of Nutrition and Immune Function in the Field*. Washington (DC): National Academies Press (US). 1999; 13. <https://www.ncbi.nlm.nih.gov/books/NBK230984/>

[51] Meydani SN, Meydani M, Blumberg JB, Leka LS, Siber G, Loszewski, Thompson C, Pedrosa MC, Diamond RD, Stollar BD. Vitamin E Supplementation and In Vivo Immune Response in Healthy Elderly Subjects A Randomized Controlled Trial. *JAMA*. 1997; 277(17):1380-1386. doi:10.1001/jama.1997.03540410058031.

[52] Morris MC, Evans DA, Bienias JL, Tangney CC, Wilson RS. Vitamin E and cognitive decline in older persons. *Arch Neurol*. 2002; 59 (7), 1125-1132. doi:10.1001/archneur.59.7.1125.

[53] Masaki KH, Losonczy KG, Izmirlian G, Foley DG, Ross GW, Petrovitch H, Havlik R, White LR. Association of Vitamin E and C supplement use with cognitive function and dementia in elderly men. *Neurology*. 2000; 54(6). Doi: <http://doi.org/10.1212/WNL.54.6.1265>.

[54] de Leeuw FA, Schneider, JA, Agarwal S, Leurgans SE, Morris MC. Brain tocopherol levels are associated with lower activated microglia density in elderly human cortex. *Translational research & Clinical interventions*. 2020. <https://doi.org/10.1002/trc2.12021>.

[55] Kang JH, Cook N, Manson J, Buring JE, Grodstein F. A Randomized Trial of Vitamin E Supplementation and Cognitive Function in Women. *Arch Intern Med*. 2006;166(22):2462-2468. doi:10.1001/archinte.166.22.2462.

[56] Graat JM, Schouten EG, Kok FJ. Effect of vitamin E and multivitamin-mineral supplementation on acute respiratory tract infection in elderly. *JAMA*. 2002; 288(6), 715-721. Doi: 10.1001/jama.288.6.715.

[57] Lewis ED, Meydani SN, Wu D. Regulatory Role of Vitamin E in the Immune System and Inflammation. *International Union of Biochemistry and Molecular Biology*. 2019; 71(4), 487-494. DOI 10.1002/iub.1976.

[58] Funte MD, Sánchez C, Vellejo C, Cerro ED, Arnalich F, Hernanz A. Vitamin C and vitamin C plus E improve the immune function in the elderly. *Experimental Gerontology*, 2020. <https://doi.org/10.1016/j.exger.2020.111118>.

[59] Giada S, Sahar S, Bertrand L, Alexandra P, Jason S, Louis-Patrick H,

- Jean-Pierre R, Philip W, Marc D, Peter G, Marina K. Vitamin E is an effective treatment for nonalcoholic steatohepatitis in HIV mono-infected patients. *AIDS*. 2020; 34(2), 237-244. doi: 10.1097/QAD.0000000000002412.
- [60] Mileva M, Galabov AS. Vitamin E and Influenza virus Infection. *IntechOpen*. 2018. Doi: 10.5772/intechopen.80954.
- [61] Beck MA. Increased Virulence of Coxsackievirus B3 in Mice Due to Vitamin E or Selenium Deficiency. *The Journal of Nutrition*. 1997; 127(5), 966S-970S. <https://doi.org/10.1093/jn/127.5.966S>.
- [62] Brion L, Bell E, Raghuveer T. *et al*. What Is The Appropriate Intravenous Dose Of Vitamin E For Very-Low-Birth-Weight Infants?. *J Perinatol*. 2004; 24, 205-207. <https://doi.org/10.1038/sj.jp.7211078>
- [63] Franco S, Goriacko P, Rosen O, Morgan-Joseph T. The incidence of complications associated with parenteral nutrition in preterm infants < 32 weeks with a mixed oil lipid emulsion versus a soybean oil lipid emulsion in a level IV neonatal intensive care unit. *Journal of Parenteral and Enteral Nutrition*. 2020; doi:10.1002/jpen.2011.
- [64] Maria Pacifici G. Effects of Vitamin E in Neonates and Young Infants. *Int J Pediatr* 2016; 4(5): 1745-57.
- [65] Hittner HM, Speer, ME, Rudolph AJ, Blifield C, Chadda P, Holbein B, Godio LB, Kretzer FL. Retrolental Fibroplasia and Vitamin E in the Preterm Infant—Comparison of Oral Versus Intramuscular:Oral Administration. *Pediatrics*. 1984; 73 (2), 238-249.
- [66] Jansson L, Lindreoth M, Tyoppoen J. Intestinal Absorption of Vitamin E in Low Birth Weight Infants. *Acta Paediatrica*. 1984; 73(3), 329-332. doi:10.1111/j.1651-2227.1994.tb17743.x.
- [67] Porcelli PJ, Greene H, Adcock E. A Modified Vitamin Regimen for Vitamin B2, A, and E Administration in Very-Low-Birth-Weight Infants. *Journal of Pediatric Gastroenterology and Nutrition*. 2004; 38 (4), 392-400.
- [68] Odigwe CC, Smedslund G, Ejemot-Nwadiaro RI, Anyanechi CC, Krawinkel MB. Supplementary vitamin E, selenium, cysteine and riboflavin for preventing kwashiorkor in preschool children in developing countries. *Cochrane Database of Systematic reviews*. 2010; 4. Doi: 10.1002/14651858.CD008147.pub2.
- [69] Sokol R J, Butler-Simon N, Conner C, Heubi JE, Sinatra FR, Suchy FJ, *et al*. Multicenter trial of d- α -tocopheryl polyethylene glycol 1000 succinate for treatment of vitamin E deficiency in children with chronic cholestasis. *Gastroenterology*. 1993; 104(6), 1727-1735. doi: 10.1016/0016-5085(93)90652-s.
- [70] Sobouti B, Hooman N, Movahed M. The effect of vitamin E or vitamin A on the prevention of renal scarring in children with acute pyelonephritis. *Pediatric Nephrology*. 2013; 28, 277-283. <https://doi.org/10.1007/s00467-012-2308-4>.
- [71] Okebukola PO, Kansra S, Barrett. Vitamin E supplementation in people with cystic fibrosis. *Cochrane Database of Systematic Reviews*. 2017; Issue 3. Doi: 10.1002/14651858.CD0094.pub3.
- [72] Baylin A, Villamor E, Rifai N, Msamanga G, Fawzi WW. Effect of vitamin supplementation to HIV-infected pregnant women on the micronutrient status of their infants. *European Journal of Clinical Nutrition*. 2005; 59, 960-968. <https://doi.org/10.1038/sj.ejcn.1602201>.
- [73] Goldfarb AH, McKenzie MJ, Bloomer RJ. Gender comparison of exercise-induced oxidative stress:

influence of antioxidant supplementation. *Applied physiology, Nutrition and Metabolism*. 2007; 32(6): 1124-1131. <https://doi.org/10.1139/H07-078>.

[74] Huang, H, Appel LJ, Croft KD, Miller ER, Moru TA, Puddey IB. Effects of vitamin C and vitamin E on in vivo lipid peroxidation: results of a randomized controlled trial. *The American Journal of Clinical Nutrition*. 2002; 76(3), 549-555. <https://doi.org/10.1093/ajcn/76.3.549>.

[75] Hemila H, Virtamo J, Albanes D, Kaprio J. Vitamin E and Beta-carotene supplementation and hospital treated pneumonia incidence in male smokers. *CHEST*. 2004; 125(2), 557-565.

[76] Debier C. Vitamin E During Pre- and Postnatal Periods. *Vitamins & Hormones*. 2007; 357-373. doi:10.1016/s0083-6729(07)76013-2.

[77] Oveisi MR, Sadeghi N, Jannat B, Hajimahmoodi M, Behfar A, Jannat F, MohktariNasab. Human Breast milk provides better antioxidant capacity than infant formula. *Iran J Res*. 2010; 9 (4) 445-449.PMID: 24381611.

[78] Katti K, Ayasolla KR, Lurcotta T, Potak D, Codipilly C, Weinberger B. Lipid peroxidation products as predictors of oxidant-mediated disease in preterm infants. *The Journal of Maternal-Fetal & Neonatal Medicine*. 2020; <https://doi.org/10.1080/14767058.2020.1869934>.

[79] Kaempf-Rotzoll, Hellstern and Linderamp. Influence of Long-Chain Polyunsaturated Fatty Acid Formula Feeds on Vitamin E Status in Preterm. *International Journal of vitamin and nutrition research*. 2013; 73 (5), 377-387. <https://doi.org/10.1024/0300-9831.73.5.377>.

[80] Korchazhkina O, Jones E, Czauderna M, Spencer SA. Effects of

exclusive formula or breast milk feeding on oxidative stress in healthy preterm infants. *Archives of Disease in Childhood*. 2006; 91, 327-329. <http://dx.doi.org/10.1136/adc.2005.084798>.

[81] Mino M. Use and safety of elevated dosages of vitamin E in infants and children. *International Journal of vitamin and Nutrition research*. 1989; 30, 69-80. PMID: 2507708.

[82] Sokol RJ, Heubi JE, Iannaccone ST, Bove, KE and Balistreri, WF. Vitamin E Deficiency with Normal Serum Vitamin E Concentrations in Children with Chronic Cholestasis. *N Engl J Med*. 1984; 310, 1209-1212. Doi: 10.1056/NEJM1984051031011901.

[83] Utsugi MT, Nakade M, Imai E, Kasaoka NT, Nozue M, et al. Distribution of vitamin E intake among Japanese Dietary supplement and fortified food users: A secondary analysis from the nutritional health and nutrition survey,2003-2009. *J. Nutr.Sci. vitaminol*. 2013; 59, 576-583.

[84] Maggini S, Wintergerst ES, Beveridge S, Horning DH. Selected vitamins and trace elements support immune function by strengthening epithelial barriers and cellular and humoral immune responses. *British Journal of Nutrition*. 2007; 98(S1), s29-s35. <http://doi.org/10.1017/S0007114507832971>.

[85] Meydani M. Effect of functional food ingredients: vitamin E modulation of cardiovascular diseases and immune status in the elderly. *The American Journal of Clinical Nutrition*. 2000; 71(6), 1665S-1668S. <https://doi.org/10.1093/ajcn/71.6.1665S>.

[86] Gay R, Meydani SN. The effect of vitamin, vitamin B₆ and vitamin B₁₂ on immune function. *Nutrition in clinical care*. 2002; 4(4), 188-198. <https://doi.org/10.1046/j.1523-5408.2001.00142.x>.

[87] Panebianco C, Bou Nasser EF, Forlani G, Palmiweri G, Tatangelo L, Villani A, Xu L, Accolla R, Pazienza V. Probiotic *Bifidobacterium lactis*, anti-oxidant vitamin E/C and anti-inflammatory dha attenuate lung inflammation due to pm2.5 exposure in mice. *Beneficial Microbes*. 2019; 10 (1), 69-75(7). <https://doi.org/10.3920/BM2018.0060>.

[88] Cook-Mills JM, Avila PC. Vitamin E and D regulation of allergic asthma immunopathogenesis. *International Immunopharmacology*. 2014; 23(1), 364-372. doi:10.1016/j.intimp.2014.08.007.

[89] Meydani SN, Leka LS, Fine BC, Dallal GE, Keusch GT, Singh MF, Hamer DH. Vitamin E and Respiratory Tract Infections in Elderly Nursing Home Residents: A Randomized Controlled Trial. *JAMA*. 2004; 292(7); 828-836. doi:10.1001/jama.292.7.828

[90] Kolleck I, Sinha P, Rustow B. Vitamin E as an antioxidant of the lung mechanisms of vitamin E delivery to alveolar type II cells. *American Journal of Respiratory and Critical care medicine*. 2002; 166 (1), s62-s66. <https://doi.org/10.1164/rccm.2206019>.