We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists



186,000

200M



Our authors are among the

TOP 1% most cited scientists





WEB OF SCIENCE

Selection of our books indexed in the Book Citation Index in Web of Science™ Core Collection (BKCI)

# Interested in publishing with us? Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected. For more information visit www.intechopen.com



#### Chapter

# Potential of Cyanobacteria in Wound Healing

Laxmi Parwani, Mansi Shrivastava and Jaspreet Singh

# Abstract

The wound care market is rapidly expanding due to the development of innumerable dressings that exhibit specific healing requirements for different wound types. The use of biomaterials as suitable wound dressing material is highly advantageous due to their biocompatibility, biodegradability, and non-toxicity. Cyanobacteria have been widely explored for their potential applications in wound healing, as they are the rich source of bioactive compounds with antibacterial, antitumor, antiviral, antioxidant, and antifungal activities. In recent years this group of organisms has been widely studied due to their immense potential in biomedical applications. Although their different bioactivities can support wound healing in different ways, very few forms have proven utility as a wound-healing agent. This chapter gives an insight into the potential of cyanobacteria in wound healing. Different bioactive compounds present in variable forms of cyanobacteria and their associated activities were reported to support tissue regeneration and wound healing acceleration. As the demand for cost-effective, bioactive wound care products is ever increasing, these organisms have immense potential to be utilized for the development of bioactive wound dressings. Hence, various bioactive compounds of cyanobacteria, their associated activities, and roles in wound healing have been briefly reviewed in this chapter.

**Keywords:** Cyanobacteria, wound healing, bioactive compounds, hemostatic, antioxidant activities

## 1. Introduction

Wound healing involves various interactions between cellular, molecular, biochemical, and physiological activities, making the process very complex, dynamic, and precisely programmed. Wound healing involves four phases: hemostasis, inflammation, proliferation, and remodeling to restore the structural, functional, and physiological integrities of injured tissues [1]. This process results in the regeneration and replacement of injured tissue at the wound site [2]. Several nutritional factors are required for proper cellular differentiation, immune functioning, and collagen formation [3]. Any interruption, aberrance, or prolongation in the healing process would extend the tissue damage and thus prolong the repair process, contributing to chronic wound healing [4]. In recent years there has been accelerating demand for various wound dressings, each with specific characteristics, considering the distinctive healing requirements of various wound types. Various synthetic and natural products have been widely explored for their efficiency and accelerating wound healing abilities to accomplish the need for suitable dressing for a particular wound type. The nonabsorbent and non-biodegradable nature of synthetic products makes these products unsuitable for healing purposes [5]. Medicinal plants are widely being explored for their utility in wound healing, and the ancient knowledge of medicinally essential plants, their increased popularity and utility further raised interest in exploring new natural products useful for the healing process. Various studies confirmed the anti-inflammatory, pro-collagen synthesis, antioxidant and antibacterial activities of natural products from plants and microbes, potentially beneficial for healing purposes. Biocompatibility and the presence of various bioactive phytochemicals in natural products efficiently promote the healing process and make them economically suitable for designing and fabricating dressings [6]. To satisfy the demand for new natural therapeutic agents for wound healing and to decrease the average costs involved in their development, researchers are screening organisms from overlooked microbial sources having the potential to be used as effective wound healing agents such as proteobacteria, bacterioidetes, and cyanobacteria.

Cyanobacteria are ubiquitous, oxygenic photosynthetic bacteria having diverse nature and can be found in various forms like unicellular and filamentous, marine and freshwater, free-living symbiotic, edible, and poisonous [7, 8]. This group of organisms has immense potential to produce many primary and secondary metabolites thus are known to perform potent biological activities. Produced secondary metabolites are low molecular weight, natural organic compounds, essential for average growth and development of these organisms. These metabolites have a wide range of applications in the field of medicines, industries, and biotechnology. These metabolites are a rich source of bioactive compounds, and cyanobacteria are the most promising organisms to produce them. In the last few decades, cyanobacteria have gained lots of attention for their medicinal values and wound healing properties. They are the choice of organisms due to their easy availability, fast regeneration, and huge diversity which have further expanded interest of researchers in their values as medicine and functional foods [9]. Thus, their potential as a good source of new therapeutic lead compounds has been realized during the last two decades. Cyanobacterial secondary metabolites show different medicinally essential activities such as antitumor, antibacterial, antifungal, antiviral, anti-inflammatory, immunomodulatory effects, and protease inhibition [10]. These biological activities are helpful to promote wound healing in different ways. Despite their potent biological activities, very few cyanobacterial forms are known to be useful in wound healing acceleration. Thus, this chapter presents an overview of bioactive compounds of cyanobacteria responsible for their various biological activities that promote the wound healing process by affecting different phases and factors of wound healing.

#### 2. Important wound healing properties of cyanobacteria

#### 2.1 Antibacterial activity

Antibacterial compounds are the molecules that kill or inhibit the growth of bacteria by affecting their physiological processes. Increased temperature due to inflammation caused by immune response and humidity caused by accumulation of fluid at the wound surface, make the wounds more vulnerable for bacterial infections. Once the wounded site came in contact with bacteria, they can penetrate underlying tissues, leading to life-threatening infections. To deal with such pathological conditions, effective wound management practices are essentially required. To improve the healing process and to reduce the wound bacterial colonization as

well as infection at the wound site, the use of systematic antibiotics and the application of antimicrobial-loaded wound dressings are viable options to overcome the issue of infection and associated delay in healing. Considering the fact that usage of antibiotics has led to the increasing emergence of multidrug-resistant (MDR) strains of bacteria which negatively affects the healing process and worsens this issue [11]. Thus, there is a need to develop/discover an effective wound management material which can efficiently cure wound in such conditions. Several cyanobacteria are known to produce intracellular and extracellular bioactive compounds that possess antibacterial activities [12]. These compounds are effective against various bacterial strains like Staphylococcus epidermidis, Bacillus cereus, Escherichia coli, Staphylococcus aureus, Salmonella typhi, Staphylococcus albus, Micrococcus flavus, Bacillus subtilis, Pseudomonas aeruginosa, Serratia marcescens, Mycobacterium smeg*matis, Streptococcus pyogenes*, and some other Gram-positive and Gram-negative bacteria [13–19]. However, the production of antibacterial agents depends upon the composition of the culture medium, incubation period, pH, temperature, and light intensity during the growth of the selected cyanobacteria [20]. Table 1 shows the bioactive compounds produced by different cyanobacterial strains, potentially useful for their antibacterial activity.

## 2.2 Antifungal activity

Devastating chronic wound infection is a significant reason for trauma worldwide, which causes serious public health problems. Majorly bacteria are responsible for infections in wounds, but very few recent studies analyzed both fungal and bacterial communities in the microbiome of chronic wounds, suggesting the role of fungi as underappreciated agents that leads to complications at the wound site [33].

Cyanobacterial Strain	Bioactive Compounds	Reference
Nostoc commune	Noscomin,Comnostins	[13, 21]
<i>Nostoc</i> sp.	Comnostin, Muscoride A, Dodecahydrophenanthrene, 4-methylchrysazin, Norharmane and 4-hydroxy-7- methylindan-1,carbamidocyclophane, Nostocarboline	[13, 16, 22]
Nostoc muscorum	Muscoride A	[23]
Lyngbya majuscula	Tanikolide	[24]
Microcoleus lacustris	Abietane	[25]
Oscillatoria redekei	Coriolic acid	[14]
<i>Lyngbya</i> sp.	Lyngbyazothrin, pahayokolide A	[26, 27]
Scytonema ocellatum, Tolypothrix conglutinate	Tolytoxin	[28]
Lyngbya majuscula	Pitipeptolides, malyngolid	[29, 30]
<i>Scytonema</i> sp.	Scytonemin	[31]
Nostocspongiaeforme	Tenuecyclamides	[31]
Micrococcus lacustris	Norbietane	[32]
<i>Fischerella</i> sp.	hapalindole T, ambiguine-I isonitrile	[14, 18]
Anabaena	Exopolysaccharide	[32]
Tolypothrix tenuis	Exopolysaccharide	[32]

#### Table 1.

Bioactive compounds of cyanobacterial strains possess antibacterial activity.

Hard to heal wounds like diabetic foot ulcers (DFUs) are majorly infected by members of the genus *Candid*a [34, 35]. Prolong infection along with delayed healing makes these wound types chronic and challenging to manage. Over 75% Candida species were isolated by DFUs. In allergic rhinitis, respiratory disease, diabetic foot ulcers, other allergic fungi such as *Cladosporium* spp., *Aspergillus* spp., *Penicillium* spp., Alternaria spp., Pleospora spp., and Fusarium spp. are reported as critical infection-causing agents. Pathogenic and opportunistic fungi such as Candida spp., *Trichosporon asahii*, and *Rhodotorula* spp. are reported to associate with stalled open wounds or wounds resulting in an amputation [33]. Antifungal drugs to cure infected wounds are limited due to their high cost and side effects. So there is an emerging need to discover a new natural fungicidal agent. Many cyanobacterial compounds have been reported to possess inhibitory effects against different pathogenic strains of fungi [36]. They are an excellent source of various bioactive compounds that belong to several different chemical classes like peptides, polyketides, and alkaloids [37]. Bioactive antifungal compounds like nostodione, nostocyclamide, carazostatin, scytophycins, tolytoxin, phytoalexin, fisherellin, hapalindole are isolated from different strains of cyanobacteria [38]. Organic extracts of several cyanobacteria like Oscillatoria latevirens, Chrococcus minor, Phormidium corium, Lyngbya martensiana, Lyngbya aestuarii, Aphanothece bullosacrude and Microcystis aeruginosa have potential antifungal activity against Candida albicans and Aspergillus *flavus* [19]. **Table 2** shows various bioactive compounds of cyanobacteria possessing antifungal activity.

#### 2.3 Antioxidant activity

Antioxidants are substances that prevent oxidation. They balance oxidative stress by eliminating free radicals and allowing the regeneration of tissue by repairing the cells [50]. Respiratory burst generates oxidants during wound healing; production of these oxidants supports acceleration in healing. These produced oxidants act as a messenger and thus promote healing. However, a delicate balance between oxidants and antioxidants required to control the progress of the wound. For normal wounds, low physiology levels of reactive oxygen species and oxidative stress are required at wound sites. Whereas oxidative stress and impaired wound healing led by their overexposure. To improve the level of oxidative stress, increased level of antioxidants is expected, which further help in healing acceleration [51]. Antioxidants also preserve

Cyanobacterial Strain	Bioactive Compounds	Reference
Scytonema hofmanni	Cyanobacterin	[39]
Calothrix fusca	Calophycin	[40]
Hapalosiphon fontinalis	Fontonamide, hapalindole	[38, 41]
Tolypothrix tenuis	Toyocamycin, Tubercidin	[42, 43]
Lyngbya majuscula	Majusculamide C, Hectochlorin, Tanikolide	[24, 44, 45]
Hyella caespitosa	Carazostatin	[46]
Nostoc sp.	Nostocyclamide	[47]
Nostoc commune	Nostofungicidine, Nostodione	[48, 49]
Scytonema sp.	tolytoxin, phytoalexin, scytophycins	[38]
Fischerella muscicola	fisherellin	[38]

#### Table 2.

Antifungal activity of bioactive compounds isolated from different cyanobacterial strains.

and stimulate the function of immune cells against homeostatic disorders. Therefore, their increased levels can improve the immune response and accelerate wound healing [52]. Accumulation of low molecular weight iron at the wound site also increases inflammation and microbial invasion of the wound [53], suggesting the requirement of a chelating agent to achieve proper healing. Production of various, chemically diverse groups of secondary metabolites from cyanobacteria established their industrial significance and made them an excellent source of antioxidants that facilitate the formation of the body's defense mechanism against free radical induced damages to cells. Their antioxidant and metal chelating ability are reportedly due to phytonutrients and pigments present in them [54]. Their cell-free extracts possess free radical scavenging property, metal chelating activity, and deoxyribose protection [55]. The antioxidant properties of the cyanobacterial cell extracts are imparted by the total phenol and total flavonoid content present in the extracts. The free radical scavenging, metal chelating, and antioxidative damage protecting properties of cyanobacterial cell extracts are presumably linked with varied quantities of polyphenolics, gallic, chlorogenic, caffeic, vanillic, and ferulic acids, flavonoids, quercetin, and kaempferol present in them. Exopolysaccharides of cyanobacteria also exhibit good antioxidant and anticoagulant activities. They can also induce oxidants and antioxidant enzymes and are known as immunostimulators [56]. Exopolymers of three strains of Anabaena and Tolypothrix tenuis exhibited antioxidant activities against  $O_2$ ,  $H_2O_2$ ,  $OH_2$ , and  $NO_2$ . These polymers also possess Fe2<sup>+</sup> chelating activity, helps in preventing the invasion of pathogenic organisms at the wound site. A strong correlation of sulfate content against superoxide and nitric oxide radicals scavenging activity of exopolymers was found, whereas  $H_2O_2$  scavenging ability and reducing power were contributed by phenols present in them. The overall reducing power and superoxide control strongly related to their iron chelation ability [57]. The highest phenolic, flavanoid, and phycobiliprotein content in Lyngbya sp. possessed the highest 2,2-diphenyl-1-picrylhydrazyl (DPPH) activity. The antioxidant potential of *Spirulina* sp. was presumable related to the presence of alcoholic and phenolic OH groups in the cellular structure of the organism [54]. Phenolic content in methanolic extracts of Anabaena sp., Nostoc sp., Nostoc commune, Nodularia spumigena, Leptolyngbya protospira, Phormidiochaete sp. and Arthrospira platensis reported higher antioxidant activity than their aqueous extracts [58, 59]. Similarly, ethanolic extracts of Phormidium fragile, Lyngbya limnetica, Scytonema bohnerii and Calothrix fusca possess antioxidant ability [60]. Cyanobacterial pigments also possess antioxidant activities. In Lyngbya and Phormidium sp. peroxyl and hydroxyl radicals, scavenging ability was reported to be linked with the presence of covalently linked tetrapyrrole chromophore with phycocyanobilin [61]. Phycocyanin, a type of phycobiliprotein isolated from Spirulina platensis and Geitlerinema sp., reported to have antioxidant and anti-inflammatory properties and ability to scavenge peroxyl, hydroxyl, and superoxide radicals [62]. Phytonutrients, pigments, and polysaccharides of cyanobacteria exhibit good antioxidant, anticoagulant, and metal chelating ability [54, 56]. Polysaccharidic exopolymers of three strains of Anabaena and Tolypothrix tenuis possessed antiradical and Fe<sup>2+</sup> chelating activity. All these exopolymers exhibited antioxidant activities against  $O_2^{\bullet}$ ,  $H_2O_2$ ,  $OH^{\bullet}$ , and  $NO^{\bullet}$ , thus helps in the rapid healing of the wound [57].

#### 2.4 Immunomodulatory and anti-inflammatory effects

Inflammation is a local, protective and physiological response to microbial invasion or injury. The magnitude of the inflammatory response is crucial: insufficient responses result in immunodeficiency, whereas excessive responses cause morbidity and mortality. Therefore, homeostasis and health are restored when inflammation

is limited by anti-inflammatory responses [63]. Natural compounds have gained lots of attention in treating various types of inflammations to reduce the reaction of the immune system against pathogens, toxic compounds, and damaged cells. The immune system actively participates in homeostasis, re-establishment, following tissue injury via multiple mechanisms, and plays a critical role throughout the wound healing process. Immune system control to promote tissue repair and regeneration is an attractive approach when designing regenerative strategies. Now a day, the multifunctional immunomodulatory properties of cyanobacteria are gaining much attention in the field of medicine. Cyanobacteria produce various metabolites with different chemical structures, including small molecules of peptides and proteins, polysaccharides, fatty acids, and their derivatives, possessing anti-inflammatory activities [64]. Different cyanobacterial components control the release of certain cytokines from human monocytes and macrophages. Depending on the wound microenvironment, they also can reduce or activate the production of reactive oxygen species from neutrophils [65]. Immunomodulatory effects of cyanobacteria highly induce activation of both types of immune cells which could promote wound debridement, accelerate re-epithelization, and wound closure [66]. Edible cyanobacterial forms are known for their immuneboosting abilities, and many studies have proven their immunomodulatory and anti-inflammatory effects. Spirulina is a widely consumed cyanobacterial strain because of its extraordinary neutraceutical and pharmaceutical values. Dietary effects of Spirulina platensis showed increased phagocytic and increased antigen production and increased natural killer cell-mediated antitumor activity in the test animals under study [67]. Daily consumption of Spirulina stimulates and promotes the immune system by increasing the phagocytic activity of macrophages, induces antibodies and cytokines production, increases accumulation of natural killer cells into tissues, and activates T and B cells [68]. Daily consumption of Spirulina for 16 consecutive weeks reported a significant rise in plasma IL- 2 concentration and a significant reduction in IL-6 concentration in humans [69]. Nonadecane and 9-Eicosyne in *Spirulina* were supposed to be responsible for the anti-inflammatory effects [70]. A high-molecular-weight polysaccharide extracted from the cyanobacterium *Spirulina* is known as Immulina<sup>®</sup>, a potent activator of innate immune cells and exerts inhibitory effects in the induced allergic inflammatory response [71, 72]. It also exhibits anti-inflammatory properties and can inhibit the release of histamine from mast cells [47]. Gama and alpha-linolenic acid are the fatty substances extracted from Spirulina and Aphanizomenon flos-aquae, respectively, which inhibit the formation of inflammatory mediators [48]. Similarly, Nostoc is an edible, largely consumed cyanobacterial form after Spirulina. Polysaccharide-rich extracts of Nostoc commune could be potentially used for macrophage activation and consequently inhibit the leukemic cell growth and induce monocytes/macrophages when used to treat human monoblastoid leukemia U937 cells [73]. Different immunomodulating activities of *Microcystis aeruginosa*, Synechocystis aquatilis, Oscillatoria redekei, Anabaena flos-aque, Aphanizomenon flos-aquae, Oscillatoria rubescens, Oscillatoria tenuis have also been reported [74] their different cell extracts variably promote the proliferation of lymphocytes. On the other hand, some species of cyanobacteria that contain cytotoxic metabolites (cyanotoxins) can cause immunotoxicity and immunosuppression. The property of immunotoxicity is well reported in cyanobacterial bloom extracts containing microcystin [75]. Treatment with microcystin showed inhibition of lipopolysaccharide-induced lymphoproliferation and decreased numbers of antibody-forming cells; this results in immunosuppression in mice that were immunized using T-dependent antigen sheep red blood cells. A form of cyanobacterial toxin, cylindrospermopsin, is an important water pollutant having broad biological activity. Cylindrospermopsin promotes

significant production of pro-inflammatory mediator tumor necrosis factor α and reactive oxygen species when introduced in macrophages cells [76]. Effmert in 1991 reported aqueous extracts of different cyanobacterial species Microcystis aeruginosa, Synechocystis aquatilis, Oscillatoria redekei, Anabaena flos-aque, Aphanizomenon flos-aquae, Oscillatoria rubescens, Oscillatoria tenuis possess strong immunomodulating activities [74]. Aqueous extract of the marine cyanobacterium, *Trichodesmium* erythraeum, ethyl acetate extract of non-edible blue-green algae Geitlerinema *splendidum* exhibit anti-inflammatory activity [77, 78]. In chronic wounds where a coordinated wound healing cascade fails to progress beyond the inflammatory phase, the anti-inflammatory activity of cyanobacteria helps in the wound healing by proliferation, matrix deposition, and ultimately, wound resolution [79]. During the last decade, several bioactive compounds have been isolated from cyanobacteria having anti-inflammatory properties that suggest their importance and potential in developing a huge market of wound healing aids [10]. Different classes of cyanobacterial bioactive compounds like phycobilins, phenols, polysaccharides, steroids, and terpenoids possess potential anti-inflammatory effects. Selected anti-inflammatory compounds of different cyanobacterial strains are listed in Table 3.

#### 2.5 Hemostatic activity

For people throughout the world, traumatic injuries have been a challenge. Considering technological advancements made through age's trauma remains a leading cause of human morbidity and mortality [90]. Excess bleeding can cause delayed wound healing, hematoma formation, infection, dehiscence, and necrosis. Patients suffering from trauma and its consequent hemorrhage essentially require the establishment of hemostasis by topical wound dressings. Constriction of blood vessels, the activation of the coagulation cascade, and the formation of blood clots are essential

Chemical group of bioactive compound	Bioactive Compound	Cyanobacterial Strain	References
Amino acids and peptides	Aeruginosin	Nostoc sp.	[80]
	Porphyra	Aphanizomenon flos-aquae	[81]
	Shinorine	Anabaena variabilis	[82]
	Phycocyanin	<i>Spirulina</i> sp	[72]
	Ethyl tumonoate A	Oscillatoria margaritifera	[83]
	Cyanopeptolin	Microcystis spp	[84]
Polysaccharide	Sacran	Aphanothece sacrum	[85]
Lipids	Monogalactosyl diacylglycerol, Digalactosyl diacylglycerol, Sulphoquinovosyl diacylglycerol, Phosphatidyl glycerol	Phormidium sp.	[86]
Pigment	Scytonemin	Scytonema	[87]
Others	Coibacin A	<i>Oscillatoria</i> sp.	[88]
	Honaucins A–C	Leptolyngbya crossbyana	[89]

#### Table 3.

Bioactive compounds of cyanobacterial strains possessing anti-inflammatory properties.

and significant steps of hemostasis. Therefore, any effort made to accelerate any or all phases above can help achieve hemostasis [91]. The role of cyanobacteria and algae in hemostasis is significantly less known and identified. In the majority, their antithrombic activities useful to cure thrombosis-related diseases are reported. The antithrombic activity of Spirulan, a sulfated polysaccharide of Arthrospira platensis has been widely known. Spirulan is helpful to directly decrease the activity of thrombin and factor X activated, procoagulant proteins that can be used to prevent thrombus formation and partial lysis of thrombus [92]. Similarly, *Microcystis aeruginosa* and different cyanobacteria blooms produce probable fVIIa-soluble Tissue Factor (fVIIasTF) inhibitors [93]. Vitamin-K-dependent clotting factors like thrombin and fVIIa are associated with bleeding, related complications, and disorders. These factors can induce excessive bleeding when treated with vitamin-K antagonists. M. aeruginosa produces bioactive compound Aeruginosin, which shows positive fVIIa-sTF inhibitory activity. Daily consumption of aqueous extract of Spirulina platensis, containing a high dose of phycocyanin was found safe for when tried on the human to test its anticoagulant activity and platelet activation ability. The studied extract is also suitable for providing rapid and robust relief in chronic pain. The extract also improved the liver function and metabolism by reducing the activity of aspartate transaminase and alanine transaminase enzymes [94]. C-phycocyanins extracted from Spirulina platensis possess significant anticoagulation, antioxidant, and prevention of DNA damage activities [95]. Exopolysaccharides of A. platensis possess antiatherogenic and anti-thrombogenic activities [96]. Although the significant roles of cyanobacterial and algal polysaccharides as antioxidants, antiviral, antitumoral, and anticoagulant have been well-documented and reviewed [10] the single report was found to date mentioning the role of cyanobacterial exopolymers (EPs) isolated from four desert cyanobacteria (Tolypothrix tenuis and three species of Anabaena) in hemostasis. These exopolymers were potentially beneficial to reduce activated partial thromboplastin time (APTT, a measure of how long blood takes to clot) and prothrombin time (PT, a measure of how long blood plasma takes to clot) by 16–41% and 12–65% respectively. The gravimetric method of thrombogenicity assessment showed that the blood clot formed by the cyanobacterial EPs was heavier vis a vis glass (positive control) and thus was thrombogenic. Similar studies can open the treasure of hidden potential of bioactive compounds of cyanobacteria in hemostasis which needs to be explored more.

#### 2.6 Wound healing effect of cyanobacteria

Specific healing requirements of a wound widen the scope of identifying more natural, economic, and effective wound healing agents, which gave extraordinary rise to the development of many synthetic and natural products useful as suitable wound healing dressing materials. Various natural products obtained by plants are widely known for their medicinal properties, facilitating wound healing. Bioactive secondary metabolites like alkaloids, essential oils, flavonoids, tannins, terpenoids, saponins, and phenolics are present in plant-based natural products. These compounds possess various activities like anti-inflammatory, antioxidant, antibacterial, procollagen synthesis, etc., and efficacy to modulate one or more phases of the wound healing process which further help in accelerated tissue regeneration during healing [97]. Similarly, cyanobacterial bioactive compounds possess important medicinal properties like immunostimulating, antiviral, antioxidant, antibacterial, antifungal, antialgal, and anticancerous activities [38, 98] but very few are known for their wound healing potential. The blue-green microalgae Spirulina has been widely studied for its wound healing potential and possess antioxidant, antimutagenic, antiviral, anticancer, anti-allergic, immune-enhancing, hepatoprotective, blood vessel relaxing, and blood lipid-lowering activities [99, 100]. The C-phycocyanin pigment

of Spirulina promoted proliferation, regeneration, and migration of cells tested on cultured human keratinocytes. A similar set of experiments on Sprague–Dawley male rats showed re-epithelization, neovascularization, presence of inflammatory cells, granulation of tissue amount, and maturation during wound healing [101]. Aqueous extract of *Spirulina platensis* showed proliferation, migration, and enhanced closure rate of wound area within 24 hours when tested on human dermal fibroblast cells (HDF) [102]. Compounds like cinnamic acid, narigenin, kaempferol, temsirolimus, phosphatidylserine, and isomeric derivatives were presumably involved in the accelerated wound healing activity of the studied aqueous extract of Spirulina platensis. Antibacterial and antioxidant activities of Spirulina are useful to improve the immunity of skin cells, and thus burn cream of ethanolic extract of Spirulina along with gold nanoparticles has been formulated. To reduce the pain by numbing the injured area, Lidocaine has been added to the cream as an anesthetic, sodium dodecyl sulfate, and glycerine used for skin moisturization [103]. Polylactic acid (PLA), polyethylene oxide, and PHB (polyhydroxybutyrate) extracted from Arthrospira LEB 18 strain used to prepare biodegradable, biocompatible nanofibrous scaffolds. Properties of prepared scaffolds like increased conductivity, higher mechanical durability with enhanced elasticity, tensile strength, and breaking elongation supported nutrient, growth factors, and metabolism byproducts distribution at the wound site [104]. Scaffolds of poly-D, L-lactic acid (PDLLA) associated with LEB 18 biomass showed better adherence to the wound, increased cell viability, and more moldable when compared with classic PDDLA [105]. Natural wound healing processes can be promoted by such biomatrices produced from Arthrospira, and also they can minimize the risk of infection [104]. PCL nanofiber loaded with Spirulina extract was fabricated as a cutaneous wound dressing material. The designed dressing showed enhanced wound regeneration ability by modulating intra- and extracellular ROS with enhancing antioxidant mechanism and increased fibroblast viability under oxidative stress [106]. Synechococcus elongatusa, a naturally occurring photoautotrophic cyanobacterium promoted angiogenesis and burn wound repair in mice. The promotion of interleukin-6 expression and secretion of extracellular vesicles probably induced pro-angiogenic and wound healing effects in the studied animals [107]. In diverse marine, freshwater, and terrestrial organisms, ultraviolet-absorbing mycosporine-like amino acids (MAAs) are found as secondary metabolites. Nostoc flagel*liforme* and *Nostoc commune*, are widely known for their edible values and reported as natural resources of MAAs. These secondary metabolites possess several beneficial effects and are useful as sun-screening cosmetics, antioxidants, and pharmaceuticals [108]. In wound healing, MAAs modulate skin fibroblast proliferation and activate adhesion kinases, extracellular signal-regulated kinases that promote acceleration in wound healing. Extensive researches are conducting on MAAs to identify their potential as a new wound healing agent to be explored as a novel biomaterial for wound healing therapies [109]. Cyanobacterial glycans are known as hydrogels that imbibe large amounts of biofluids [110]. This property of glycans extremely useful for targeted drug delivery in chronic wound cases [111], where they have been exploited for delivery of low molecular weight drugs and macro-molecular payloads (hormones, peptide, and protein drugs) [112]. Cyanobacteria have been widely explored and studied for their immense potential in various biomedical applications, but very few are reported for their accelerated wound healing abilities. Most of the studies have been conducted on Spirulina which is known for its extraordinary neutraceutical and pharmaceutical values. As these organisms are phototrophic and have minimal growth requirements, they can be easily regenerated, and also their huge diversity makes them suitable candidates for research, having hidden potential for wound healing. Further, they can be explored to develop bioactive and cost-effective wound dressing materials, suitable for all sections of society.

# 3. Conclusions

The ever-increasing demand for effective bioactive dressings suitable for different types of wounds is rapidly expanding the wound care market at diverse levels. Cyanobacteria are known for their numerous biomedical applications; thus, recently, they are a widely explored group of organisms. The cyanobacterial bioactive compounds possess antiviral, antifungal, antibacterial, antitumor, antiinflammatory, and antioxidant activities suitable to accelerate wound healing even in chronic conditions by controlling different phases and factors of the wound. Cyanobacteria are known for the abundant availability of versatile bioactive compounds and their associated properties, but unfortunately, very few forms of cyanobacteria are known for their role in wound healing. Their easy cultivation, colossal diversity, and different biological activities can make them suitable candidates for research. Further, they can be explored more in the field of biomaterials for designing and fabrication of low-cost, biocompatible, and biodegradable wound dressings.

# Acknowledgements

The authors are indebted to the Department of Science and Technology (DST), Govt. of India New Delhi, for providing opportunity and financial support (SP/YO/2019/1502) for interdisciplinary research work. The authors are also grateful to Professor Ashish Bhatnagar and Professor Monica Bhatnagar, Department of Microbiology, M.D.S. University, Ajmer, India for their support in the subject study.

# **Conflict of interest**

The authors declare no conflict of interest.

# **Author details**

Laxmi Parwani<sup>1\*</sup>, Mansi Shrivastava<sup>1</sup> and Jaspreet Singh<sup>2</sup>

1 Department of Bioscience and Biotechnology, Banasthali Vidyapith, P.O. Banasthali, Rajasthan, India

2 School of Life Sciences, SIILAS Campus, Jaipur National University, Jaipur, Rajasthan, India

\*Address all correspondence to: laxmi\_parwani@rediffmail.com

# IntechOpen

© 2021 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

# References

[1] Judith R, Nithya M, Rose C,
Mandal AB. Application of a PDGF containing novel gel for cutaneous wound healing. Life Sciences. 2010;87:
1-8. DOI: 10.1016/j.lfs.2010.05.003

[2] Velnar T, Bailey T, Smrkolj V. The wound healing process: an overview of the cellular and molecular mechanisms. Journal of International Medical Research. 2009;37(5):1528-1542. DOI: 10.1177/147323000903700531

[3] Mackay D, Miller AL. Nutritional support for wound healing. Alternative Medicine Review. 2003;8:359-377. DOI: 10.1002/jor.23656

[4] Kurahashi T, Fujii J. Roles of antioxidative enzymes in wound healing. Journal of Developmental Biology. 2015;3(2):57-70. DOI: https:// doi.org/10.3390/jdb3020057

[5] Mir M, Ali MN, Barakullah A, Gulzar A, Arshad M, Fatima S, Asad M. Synthetic polymeric biomaterials for wound healing: a review. Progress in biomaterials. 2018;7(1):1-21. DOI: 10.1007/s40204-018-0083-4

[6] Thakur R, Jain N, Pathak R, Sandhu SS. Practices in wound healing studies of plants. Evidence-based complementary and alternative medicine. 2011;2011:1-17. DOI: https:// doi.org/10.1155/2011/438056

[7] Taton A, Grubisic S, Balthasart P, Hodgson DA, Laybourn-Parry J, Wilmotte A. Biogeographical distribution and ecological ranges of benthic cyanobacteria in East Antarctic lakes. FEMS microbiology ecology. 2006;57(2):272-289. DOI: https://doi. org/10.1111/j.1574-6941.2006.00110.x

[8] Dorador C, Vila I, Imhoff JF, Witzel KP. Cyanobacterial diversity in Salar de Huasco, a high altitude saline wetland in northern Chile: an example of geographical dispersion? FEMS microbiology ecology. 2008;64(3):419-432. DOI: 10.1111/j.1574-6941.2008. 00483.x

[9] Barrow C, Shahidi F. Marine nutraceuticals and functional foods.1st ed. CRC Press; 2007. 512 p. DOI: https:// doi.org/10.1201/9781420015812

[10] Singh R, Parihar P, Singh M, Bajguz A, Kumar J, Singh S, Singh VP, Prasad SM. Uncovering potential applications of cyanobacteria and algal metabolites in biology, agriculture and medicine: current status and future prospects. Frontiers in microbiology. 2017;8:515. DOI: https://doi.org/10.3389/ fmicb.2017.00515

[11] Li B, Webster TJ. Bacteria antibiotic resistance: New challenges and opportunities for implant-associated orthopedic infections. Journal of Orthopaedic Research®. 2018;36(1):22-32. DOI: 10.1002/jor.23656

[12] Schlegel I, Doan NT, de Chazal N, Smith GD. Antibiotic activity of new cyanobacterial isolates from Australia and Asia against green algae and cyanobacteria. Journal of Applied Phycology. 1998;10(5):471-479. DOI: https://doi.org/10.1023/A:10080426 19686

[13] Ostensvik O, Skulberg OM, Underdal B, Hormazabal V. Antibacterial properties of extracts from selected planktonic freshwater cyanobacteria-A comparative study of bacterial bioassays. Journal of applied microbiology.1998;84(6):1117-1124. DOI: 10.1046/j.1365-2672.1998.00449.x

[14] Mundt S, Kreitlow S, Jansen R. Fatty acids with antibacterial activity from the cyanobacterium *Oscillatoria redekei* HUB 051. Journal of Applied Phycology. 2003;15(2):263-267. DOI: https://doi. org/10.1023/A:1023889813697 [15] Bui HT, Jansen R, Pham HT, Mundt S. Carbamidocyclophanes A-E, chlorinated paracyclophanes with cytotoxic and antibiotic activity from the Vietnamese cyanobacterium *Nostoc* sp. Journal of natural products. 2007;70(4):499-503. DOI: 10.1021/ np060324m

[16] Nowruzi B, Haghighat S, Fahimi H, Mohammadi E. *Nostoc* cyanobacteria species: a new and rich source of novel bioactive compounds with pharmaceutical potential. Journal of Pharmaceutical Health Services Research. 2018;9(1):5-12. DOI: https:// doi.org/10.1111/jphs.12202

[17] El-Sheekh MM, Osman ME, Dyab MA, Amer MS. Production and characterization of antimicrobial active substance from the cyanobacterium *Nostoc muscorum*. Environmental toxicology and pharmacology. 2006;21(1):42-50. DOI: 10.1016/j. etap.2005.06.006

[18] Raveh A, Carmeli S. Antimicrobial ambiguines from the cyanobacterium *Fischerella* sp. collected in Israel. Journal of natural products. 2007;70(2):196-201. DOI: 10.1021/np060495r

[19] Ananya AK, Ahmad IZ. Cyanobacteria"the blue green algae" and its novel applications: A brief review. International Journal of Innovation and Applied Studies. 2014;7(1):251-261

[20] Noaman NH, Fattah A, Khaleafa M, Zaky SH. Factors affecting antimicrobial activity of *Synechococcus leopoliensis*.
Microbiological Research. 2004;159(4): 395-402. DOI: 10.1016/j.micres.2004.
09.001

[21] Jaki B, Orjala J, Heilmann J, Linden A, Vogler B, Sticher O. Novel extracellular diterpenoids with biological activity from the cyanobacterium *Nostoc commune*. Journal of natural products. 2000;63(3):339-343. DOI: 10.1021/ np9903090

[22] Becher PG, Keller S, Jung G,
Süssmuth RD, Jüttner F. Insecticidal activity of 12-epi-hapalindole J isonitrile. Phytochemistry.
2007;68(19):2493-2497. DOI: 10.1016/j. phytochem.2007.06.024

[23] Nagatsu A, Kajitani H, Sakakibara J. Muscoride A: A new oxazole peptide alkaloid from freshwater cyanobacterium *Nostoc muscorum*. Tetrahedron Letters. 1995;36(23):4097-4100. DOI: https://doi.org/10.1016/ 0040-4039(95)00724-Q

[24] Pandey VD. Cyanobacterial natural products as antimicrobial agents.International Journal of Current Microbiology and Applied Sciences.2015;4(1):310-317.

[25] Thajuddin N, Subramanian G. Cyanobacterial biodiversity and potential applications in biotechnology. Current science. 2005;89(1):47-57. DOI: https://www.jstor.org/stable/24110431

[26] Asthana RK, Srivastava A, Singh AP, Singh SP, Nath G, Srivastava R, Srivastava BS. Identification of an antimicrobial entity from the cyanobacterium *Fischerella* sp. isolated from bark of *Azadirachta indica* (Neem) tree. Journal of applied phycology. 2006;18(1):33-39. DOI: https://doi. org/10.1007/s10811-005-9011-9

[27] Zainuddin EN, Jansen R, Nimtz M, Wray V, Preisitsch M, Lalk M, Mundt S. Lyngbyazothrins A– D, Antimicrobial cyclic undecapeptides from the cultured cyanobacterium *Lyngbya* sp. Journal of natural products. 2009;72(8):1373-1378. DOI: 10.1021/np8007792

[28] Moore RE, Patterson GM, Mynderse JS, Barchi J, Norton TR, Furusawa E, Furusawa S. Toxins from cyanophytes belonging to the *Scytonemataceae*. Pure and Applied

Chemistry. 1986;58(2):263-271. DOI: 10.1351/pac198658020263

[29] Berry JP, Gantar M, Gawley RE, Wang M, Rein KS. Pharmacology and toxicology of pahayokolide A, a bioactive metabolite from a freshwater species of *Lyngbya* isolated from the Florida Everglades. Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology. 2004;139(4):231-238. DOI: 10.1016/j. cca.2004.11.005

[30] Helms GL, Moore RE, Niemczura WP, Patterson GM, Tomer KB, Gross ML. Scytonemin A, a novel calcium antagonist from a bluegreen alga. The Journal of Organic Chemistry. 1988;53(6):1298-1307. DOI: https://doi.org/10.1021/jo00241a033

[31] Gutiérrez RM, Flores AM, Solís RV, Jimenez JC. Two new antibacterial norabietane diterpenoids from cyanobacteria, *Microcoleous lacustris*. Journal of natural medicines. 2008;62(3):328-331. DOI: 10.1007/ s11418-008-0238-z

[32] Bhatnagar M, Parwani L, Sharma V, Ganguly J, Bhatnagar A. Exopolymers from *Tolypothrix tenuis* and three *Anabaena* sp. (Cyanobacteriaceae) as novel blood clotting agents for wound management. Carbohydrate polymers. 2014;99:692-699. DOI: 10.1016/j. carbpol.2013.09.005

[33] Kalan L, Grice EA. Fungi in the wound microbiome. Advances in wound care. 2018;7(7):247-255. DOI: 10.1089/ wound.2017.0756

[34] Chellan G, Shivaprakash S, Ramaiyar SK, Varma AK, Varma N, Sukumaran MT, Vasukutty JR, Bal A, Kumar H. Spectrum and prevalence of fungi infecting deep tissues of lowerlimb wounds in patients with type 2 diabetes. Journal of clinical microbiology. 2010;48(6):2097-2102. DOI: 10.1128/JCM.02035-09 [35] Dowd SE, Delton Hanson J, Rees E, Wolcott RD, Zischau AM, Sun Y, White J, Smith DM, Kennedy J, Jones CE. Survey of fungi and yeast in polymicrobial infections in chronic wounds. Journal of wound care. 2011;20(1):40-47. DOI: 10.12968/ jowc.2011.20.1.40

[36] Pawar ST, Puranik PR. Screening of terrestrial and freshwater halotolerant cyanobacteria for antifungal activities. World Journal of Microbiology and Biotechnology. 2008;24(7):1019-1025. DOI: 10.1007/s11274-007-9565-6

[37] Shishido TK, Humisto A, Jokela J, Liu L, Wahlsten M, Tamrakar A, Fewer DP, Permi P, Andreote AP, Fiore MF, Sivonen K. Antifungal compounds from cyanobacteria. Marine drugs. 2015;13(4):2124-2140. DOI: https://doi.org/10.3390/md13042124

[38] Abed RM, Dobretsov S, Sudesh K. Applications of cyanobacteria in biotechnology. Journal of applied microbiology. 2009;106(1):1-2. DOI: 10.1111/j.1365-2672.2008.03918.x

[39] Mason CP, Edwards KR, Carlson RE, Pignatello J, Gleason FK, Wood JM. Isolation of chlorinecontaining antibiotic from the freshwater cyanobacterium Scytonema hofmanni. Science. 1982;215(4531):400-402. DOI: 10.1126/science.6800032

[40] Moon SS, Chen JL, Moore RE, Patterson GM. Calophycin, a fungicidal cyclic decapeptide from the terrestrial blue-green alga *Calothrix fusca*. The Journal of Organic Chemistry. 1992;57(4):1097-1103. DOI: https://doi. org/10.1021/jo00030a013

[41] Moore RE, Yang XQ, Patterson GM. Fontonamide and anhydrohapaloxindole A, two new alkaloids from the bluegreen alga *Hapalosiphon fontinalis*. The Journal of Organic Chemistry. 1987;52(17):3773-3777. DOI: https://doi. org/10.1021/jo00226a009 [42] Moore RE. Toxins, anticancer agents, and tumor promoters from marine prokaryotes. Pure and applied chemistry. 1982;54(10):1919-1934. DOI: https://doi.org/10.1351/pac198254 101919

[43] Banker R, Carmeli S. Tenuecyclamides A– D, Cyclic Hexapeptides from the Cyanobacterium *Nostoc s pongiaeforme* var. t enue. Journal of natural products. 1998;61(10):1248-1251. DOI: 10.1021/np980138j

[44] Swain SS, Paidesetty SK, Padhy RN. Antibacterial, antifungal and antimycobacterial compounds from cyanobacteria. Biomedicine & Pharmacotherapy. 2017;90:760-776. DOI: 10.1016/j.biopha.2017.04.030

[45] Singh IP, Milligan KE, Gerwick WH. Tanikolide, a toxic and antifungal lactone from the marine cyanobacterium *Lyngbya m ajuscula*. Journal of natural products. 1999;62(9):1333-1335.DOI: https://doi. org/10.1021/np990162c

[46] Burja AM, Banaigs B, Abou-Mansour E, Burgessd JG, Wrighta PC. Marine cyanobacteria - A prolific source of natural products. Tetrahedron. 2001;57:9347-9377. DOI: https://doi.org/10.1016/ S0040-4020(01)00931-0

[47] Moore RE, Patterson GM, Carmichael WW. New pharmaceuticals from cultured blue-green algae. Biomedical importance of marine organisms. 1988;13(1988):143-150. DOI: https://doi.org/10.1007/ BF00003544

[48] Kajiyama SI, Kanzaki H, Kawazu K, Kobayashi A. Nostofungicidine, an antifungal lipopeptide from the fieldgrown terrestrial blue-green alga *Nostoc commune*. Tetrahedron letters. 1998;39(22):3737-3740. DOI: https:// doi.org/10.1016/S0040-4039(98) 00573-5 [49] Bhadury P, Wright PC. Exploitation of marine algae: biogenic compounds for potential antifouling applications. Planta. 2004;219(4):561-578. DOI: 10.1007/s00425-004-1307-5

[50] Houghton PJ, Hylands PJ, Mensah AY, Hensel A, Deters AM. In vitro tests and ethnopharmacological investigations: wound healing as an example. Journal of Ethnopharmaco logy. 2005;100(1-2):100-107. DOI: 10.1016/j.jep.2005.07.001

[51] Campa-Córdova AI, Hernández-Saavedra NY, Ascencio F. Superoxide dismutase as modulator of immune function in American white shrimp (*Litopenaeus vannamei*). Comparative biochemistry and physiology part C: Toxicology & pharmacology. 2002;133(4):557-565. DOI: 10.1016/s1532-0456(02)00125-4

[52] De la Fuente M, Victor VM. Antioxidants as modulators of immune function. Immunology and cell biology. 2000;78(1):49-54. DOI: 10.1046/j.1440-1711.2000.00884.x

[53] Weinberg ED. Iron withholding: a defense against infection and neoplasia. Physiological reviews. 1984;64(1):65-102. DOI: 10.1152/physrev.1984.64.1.65

[54] Hossain MF, Ratnayake RR, Meerajini K, Wasantha Kumara KL. Antioxidant properties in some selected cyanobacteria isolated from fresh water bodies of Sri Lanka. Food science & nutrition. 2016;4(5):753-758. DOI: 10.1002/fsn3.340

[55] Singh DP, Prabha R, Verma S, Meena KK, Yandigeri M. Antioxidant properties and polyphenolic content in terrestrial cyanobacteria. 3 Biotech. 2017;7(2):1-4.

[56] Costa LS, Fidelis GP, Cordeiro SL, Oliveira RM, Sabry DD, Câmara RB, Nobre LT, Costa MS, Almeida-Lima J, Farias EH, Leite EL. Biological activities

of sulfated polysaccharides from tropical seaweeds. Biomedicine & Pharmacotherapy. 2010;64(1):21-28. DOI: 10.1016/j.biopha.2009.03.005

[57] Parwani L, Bhatnagar M, Bhatnagar A, Sharma V. Antioxidant and iron-chelating activities of cyanobacterial exopolymers with potential for wound healing. Journal of applied phycology. 2014;26(3):1473-1482. DOI: 10.1007/s10811-013-0180-7

[58] Pant G, Kumar G, Karthik L, Prasuna RG, Rao KB. Antioxidant activity of methanolic extract of blue green algae *Anabaena* sp. (Nostocaceae). European Journal of Experimental Biology. 2011;1(1):156-162.

[59] Jerez-Martel I, García-Poza S, Rodríguez-Martel G, Rico M, Afonso-Olivares C, Gómez-Pinchetti JL. Phenolic profile and antioxidant activity of crude extracts from microalgae and cyanobacteria strains. Journal of Food Quality.2017;2017:1-8. DOI: https://doi. org/10.1155/2017/2924508

[60] Sharathchandra K, Rajashekhar M. Antioxidant activity in the four species of cyanobacteria isolated from a sulfur spring in the Western Ghats of Karnataka. International Journal of Pharma and Bio Sciences. 2013;4(1):275-285.

[61] Gerwick WH, Coates RC, Engene N, Gerwick L, Grindberg RV, Jones AC, Sorrels CM. Giant marine cyanobacteria produce exciting potential pharmaceuticals. Microbe-American Society for Microbiology. 2008;3(6):277-284. DOI: 10.1128/ microbe.3.277.1

[62] Bermejo P, Piñero E, Villar ÁM. Iron-chelating ability and antioxidant properties of phycocyanin isolated from a protean extract of *Spirulina platensis*. Food chemistry. 2008;110(2):436-445. DOI: 10.1016/j.foodchem.2008.02.021 [63] Tracey KJ. The inflammatory reflex. Nature. 2002;420(6917):853-859. DOI: 10.1038/nature01321

[64] Tabarzad M, Atabaki V, Hosseinabadi T. Anti-inflammatory Activity of bioactive compounds from microalgae and cyanobacteria by focusing on the mechanisms of action. Molecular Biology Reports. 2020;47(8):6193-6205. DOI: 10.1007/ s11033-020-05562-9

[65] Stewart I, Schluter PJ, Shaw GR. Cyanobacterial lipopolysaccharides and human health–a review. Environmental Health. 2006;5(1):1-23. DOI: 10.1186/ 1476-069X-5-7

[66] Larouche J, Sheoran S, Maruyama K, Martino MM. Immune regulation of skin wound healing: mechanisms and novel therapeutic targets. Advances in Wound Care. 2018;7(7):209-231. DOI: 10.1089/ wound.2017.0761

[67] Hayashi O, Katoh T, Okuwaki Y. Enhancement of antibody production in mice by dietary *Spirulina platensis*. Journal of Nutritional Science and Vitaminology. 1994;40(5):431-441. DOI: 10.3177/jnsv.40.431

[68] Khan Z, Bhadouria P, Bisen PS. Nutritional and therapeutic potential of *Spirulina*. Current pharmaceutical biotechnology. 2005;6(5):373-379. DOI: 10.2174/138920105774370607

[69] Park HJ, Lee YJ, Ryu HK, Kim MH, Chung HW, Kim WY. A randomized double-blind, placebo-controlled study to establish the effects of *Spirulina* in elderly Koreans. Annals of Nutrition and Metabolism. 2008;52(4):322-328. DOI: 10.1159/000151486

[70] Punitha P, Bharathi V. In vitro anti-inflammatory activity in ethanolic extract of *Spirulina plantensis* and phytochemical analysis underneath GC-MS/MS. World Journal of Pharmaceutical and Medical Research. 2017;3(11):217-225.

[71] Appel K, Munoz E, Navarrete C, Cruz-Teno C, Biller A, Thiemann E. Immunomodulatory and inhibitory effect of Immulina®, and Immunloges® in the Ig-E mediated activation of RBL-2H3 cells. A new role in allergic inflammatory responses. Plants. 2018;7(1):13. DOI: 10.3390/ plants7010013

[72] Singh S, Kate BN, Banerjee UC. Bioactive compounds from cyanobacteria and microalgae: an overview. Critical reviews in biotechnology. 2005;25(3):73-95. DOI: 10.1080/07388550500248498

[73] Liao HF, Wu TJ, Tai JL, Chi MC, Lin LL. Immunomodulatory potential of the polysaccharide-rich extract from edible cyanobacterium Nostoc commune. Medical Sciences. 2015;3(4):112-123. Doi: 10.3390/ medsci3040112

[74] Effmert U, Mundt S, Teuscher E. Investigations of the immunomodulatory effect of cyanobacterial extracts. Allergie und Immunologie. 1991;37(2):97-102.

[75] Shen PP, Zhao SW, Zheng WJ, Hua ZC, Shi Q, Liu ZT. Effects of cyanobacteria bloom extract on some parameters of immune function in mice. Toxicology letters. 2003;143(1):27-36. DOI: https://doi.org/10.1016/ S0378-4274(03)00110-3

[76] Moosova Z, Pekarova M, Sindlerova LS, Vasicek O, Kubala L, Blaha L, Adamovsky O. Immunomodulatory effects of cyanobacterial toxin cylindro spermopsin on innate immune cells. Chemosphere. 2019;226:439-446. DOI: 10.1016/j.chemosphere.2019.03.143

[77] Silambarasan G, Ramanathan T, Nabeel MA, Kalaichelvan VK, Kathiresan K, Balasubramanian T. Anti-inflammatory activity of the marine cyanobacterium *Trichodesmium erythraeum* against carrageenan-induced paw oedema in wistar albino rats. European Journal of Inflammation. 2011;9(1):53-56. DOI: https://doi.org/10. 1177/1721727X1100900108

[78] Krishnaveni J, Alarmal MS, Jency M. In vitro antiinflammatory activity of blue-green algae-*Geitlerinema splendidum* collected from Western Ghats, South India Asian Journal of Pharmaceutical and Clinical Research. 2018;11(5),407-409. DOI: https://doi.org/10.22159/ajpcr.2018. v11i5.25301

[79] Eming SA, Koch M, Krieger A, Brachvogel B, Kreft S, Bruckner-Tuderman L, Krieg T, Shannon JD, Fox JW. Differential proteomic analysis distinguishes tissue repair biomarker signatures in wound exudates obtained from normal healing and chronic wounds. Journal of proteome research. 2010;9(9):4758-4766. DOI: https://doi.org/10.1021/ pr100456d

[80] Kapuścik A, Hrouzek P, Kuzma M, Bártová S, Novák P, Jokela J, Pflüger M, Eger A, Hundsberger H, Kopecký J. Novel Aeruginosin-865 from *Nostoc* sp. as a Potent Anti-inflammatory Agent. ChemBioChem. 2013;14(17):2329-2337. DOI: 10.1002/cbic.201300246

[81] Torres A, Enk CD, Hochberg M, Srebnik M. Porphyra-334, a potential natural source for UVA protective sunscreens. Photochemical & Photobiological Sciences.
2006;5(4):432-435. DOI: https://doi. org/10.1039/B517330M

[82] Singh SP, Klisch M, Sinha RP, Häder DP. Effects of abiotic stressors on synthesis of the mycosporine-like amino acid shinorine in the cyanobacterium *Anabaena variabilis* PCC 7937. Photochemistry and Photobiology.

2008;84(6):1500-1505. DOI: https://doi. org/10.1111/j.1751-1097.2008.00376.x

[83] Engene N, Choi H, Esquenazi E, Byrum T, Villa FA, Cao Z, Murray TF, Dorrestein PC, Gerwick L, Gerwick WH. Phylogeny-guided isolation of ethyl tumonoate A from the marine cyanobacterium cf. *Oscillatoria margaritifera*. Journal of natural products. 2011;74(8):1737-1743. DOI: https://doi.org/10.1021/np200236c

[84] Faltermann S, Hutter S, Christen V, Hettich T, Fent K. Anti-inflammatory activity of cyanobacterial serine protease inhibitors aeruginosin 828A and cyanopeptolin 1020 in human hepatoma cell line Huh7 and effects in Zebrafish (*Danio rerio*). Toxins. 2016;8(219):1-14.DOI: https://doi. org/10.3390/toxins8070219

[85] Motoyama K, Tanida Y, Hata K, Hayashi T, Hashim II, Higashi T, Ishitsuka Y, Kondo Y, Irie T, Kaneko S, Arima H. Anti-inflammatory effects of novel polysaccharide sacran extracted from cyanobacterium *Aphanothece sacrum* in various inflammatory animal models. Biological and Pharmaceutical Bulletin. 2016;39(7):1172-1178. DOI: 10.1248/bpb.b16-00208

[86] Bruno A, Rossi C, Marcolongo G, Di Lena A, Venzo A, Berrie CP, Corda D. Selective *in vivo* anti-inflammatory action of the galactolipid monogalactosyl diacylglycerol. European Journal of Pharmacology. 2005;524(1-3):159-168. DOI: 10.1016/j. ejphar.2005.09.023

[87] Stevenson CS, Capper EA, Roshak AK, Marquez B, Grace K, Gerwick WH, Jacobs RS, Marshall LA. Scytonemin-a marine natural product inhibitor of kinases key in hyperproliferative inflammatory diseases. Inflammation Research. 2002;51(2):112-114. DOI: 10.1007/ BF02684014 [88] Balunas MJ, Grosso MF, Villa FA, Engene N, McPhail KL, Tidgewell K, Pineda LM, Gerwick L, Spadafora C, Kyle DE, Gerwick WH. Coibacins A–D, antileishmanial marine cyanobacterial polyketides with intriguing biosynthetic origins. Organic letters. 2012;14(15): 3878-3881. DOI: https://doi.org/10.1021/ ol301607q

[89] Choi H, Mascuch SJ, Villa FA, Byrum T, Teasdale ME, Smith JE, Preskitt LB, Rowley DC, Gerwick L, Gerwick WH. Honaucins A– C, potent inhibitors of inflammation and bacterial quorum sensing: synthetic derivatives and structure-activity relationships. Chemistry & Biology. 2012;19(5):589-598. Doi: 10.1016/j.chembiol.2012.03.014

[90] Morrison CA. The prehospital treatment of the bleeding patient—Dare to dream. Journal of Surgical Research. 2013;180(2):246-247. DOI: 10.1016/j. jss.2011.12.022

[91] Dai C, Liu C, Wei J, Hong H, Zhao Q. Molecular imprinted macroporous chitosan coated mesoporous silica xerogels for hemorrhage control. Biomaterials. 2010;31(30):7620-7630. DOI: 10.1016/j. biomaterials.2010.06.049

[92] Choi JH, Kim S, Kim SJ. Spirulan from blue-green algae inhibits fibrin and blood clots: Its potent antithrombotic effects. Journal of Biochemical and Molecular Toxicology.
2015;29(5):240-248. DOI: https://doi. org/10.1002/jbt.21690

[93] Anas AR, Nakajima A, Naruse C, Tone M, Asukabe H, Harada KI. Determination of FVIIa-sTF Inhibitors in Toxic Microcystis Cyanobacteria by LC-MS Technique. Marine drugs. 2016;14(1):7.DOI: https://doi. org/10.3390/md14010007

[94] Jensen GS, Drapeau C, Lenninger M, Benson KF. Clinical safety of a high dose of Phycocyanin-enriched aqueous extract from *Arthrospira* (*Spirulina*) *platensis*: Results from a randomized, double-blind, placebocontrolled study with a focus on anticoagulant activity and platelet activation. Journal of Medicinal Food. 2016;19(7):645-653. DOI: 10.1089/ jmf.2015.0143

[95] Kamble SP, Gaikar RB, Padalia RB, Shinde KD. Extraction and purification of C-phycocyanin from dry *Spirulina* powder and evaluating its antioxidant, anticoagulation and prevention of DNA damage activity. Journal of Applied Pharmaceutical Science. 2013;3(8):149. DOI: 10.7324/JAPS.2013.3826

[96] Kaji T, Okabe M, Shimada S, Yamamoto C, Fujiwara Y, Lee JB, Hayashi T. Sodium spirulan as a potent inhibitor of arterial smooth muscle cell proliferation *in vitro*. Life Sciences. 2004 ;74(19):2431-2439. DOI: 10.1016/j. lfs.2003.09.061

[97] Harborne JB. Phytochemical methods; A guide to modern Techniques of plant analysis. Chapman and Hall. New York. 1998:1-98.

[98] Dahms HU, Ying X, Pfeiffer C. Antifouling potential of cyanobacteria: a mini-review. Biofouling. 2006;22(5):317-327. DOI: https://doi. org/10.1080/08927010600967261

[99] Kim HM, Lee EH, Cho HH, Moon YH. Inhibitory effect of mastcellmediated immediate-type allergic reactions in rats by *Spirulina*, Biochemical Pharmacology 1998;55:1071-1076 DOI: 10.1016/ s0006-2952(97)00678-3

[100] Subhashini J, Mahipal SV, Reddy MC, Reddy MM, Rachamallu A, Reddanna P. Molecular mechanisms in C-Phycocyanin induced apoptosis in human chronic myeloid leukemia cell line-K562. Biochemical pharmacology. 2004;68(3):453-462. DOI: 10.1016/j. bcp.2004.02.025 [101] Gur CS, Erdogan DK, Onbasılar I, Atilla P, Cakar N, Gurhan ID. In vitro and in vivo investigations of the wound healing effect of crude *Spirulina* extract and C-phycocyanin. Journal of Medicinal Plants Research. 2013;7(8):425-433. DOI: https://doi. org/10.5897/JMPR12.705

[102] Syarina PN, Karthivashan G, Abas F, Arulselvan P, Fakurazi S. Wound healing potential of *Spirulina platensis* extracts on human dermal fibroblast cells. EXCLI journal. 2015;14:385-393. DOI: 10.17179/excli2014-697

[103] Han SG, Song JM, Lee J. Formulating a therapeutic cream for burn injury using cyanobacterial extracts and gold nanoparticles. International Journal of Medical Research and Pharmaceutical Sciences. 2017;4(3):45-53. DOI: 10.5281/ zenodo.376416

[104] Morais MG, Vaz BD, Morais EG, Costa JA. Biological effects of *Spirulina (Arthrospira)* biopolymers and biomass in the development of nanostructured scaffolds. BioMed Research International. 2014;2014. DOI: https:// doi.org/10.1155/2014/762705

[105] Steffens D, Leonardi D, da Luz Soster PR, Lersch M, Rosa A, Crestani T, Scher C, de Morais MG, Costa JA, Pranke P. Development of a new nanofiber scaffold for use with stem cells in a third degree burn animal model. Burns. 2014;40(8):1650-1660. DOI: 10.1016/j.burns.2014.03.008

[106] Jung SM, Min SK, Lee HC, Kwon YS, Jung MH, Shin HS. *Spirulina*-PCL nanofiber wound dressing to improve cutaneous wound healing by enhancing antioxidative mechanism. Journal of Nanomaterials. 2016;2016 DOI: https://doi.org/10.1155/2016/ 6135727

[107] Yin H, Chen CY, Liu YW, Tan YJ, Deng ZL, Yang F, Huang FY, Wen C,

Rao SS, Luo MJ, Hu XK. *Synechococcus elongatus* PCC7942 secretes extracellular vesicles to accelerate cutaneous wound healing by promoting angiogenesis. Theranostics. 2019;9(9):2678-2693 DOI: 10.7150/thno.31884

[108] Liu W, Xu H, Gao X. Mild Acid Hydrolysis-related Release of Watersoluble Sunscreen Pigments from the Exopolysaccharide Matrix of Edible Terrestrial Cyanobacteria. Journal of Applied Biology & Biotechnology. 2017;5(5):11-15. DOI: 10.7324/ JABB.2017.50502

[109] Choi YH, Yang DJ, Kulkarni A, Moh SH, Kim KW. Mycosporine-like amino acids promote wound healing through focal adhesion kinase (FAK) and mitogen-activated protein kinases (MAP kinases) signaling pathway in keratinocytes. Marine drugs. 2015;13(12):7055-7066. DOI: 10.3390/ md13127056

[110] Peppas NA, Bures P, Leobandung WS, Ichikawa H. Hydrogels in pharmaceutical formulations. European Journal of Pharmaceutics and Biopharmaceutics. 2000;50(1):27-46. DOI: 10.1016/s0939-6411(00)00090-4

[111] Agnihotri SA, Mallikarjuna NN, Aminabhavi TM. Recent advances on chitosan-based micro-and nanoparticles in drug delivery. Journal of Controlled Release. 2004;100(1):5-28. DOI: 10.1016/j.jconrel.2004.08.010

[112] Oh JK, Lee DI, Park JM.
Biopolymer-based microgels/nanogels for drug delivery applications. Progress in polymer science. 2009;34(12):1261-1282. DOI: 10.1016/j.progpolymsci. 2009.08.001

