

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Introductory Chapter: Metabolomics

Xianquan Zhan, Jingru Yang, Shu Zheng, Nannan Li
and Na Li

1. Introduction

The center of multiomics is being moved from genomics to phenomics (Figure 1) [1]. Proteome and metabolome are two main components of phenome, and are equally important. The concept and development of proteoforms significantly enrich the content of a proteome. A book entitled “Proteoforms: Concept and Applications in Medical Sciences” has been published focusing on proteomics at the proteoform level [3]. It is driving the editor to edit another book focusing on metabolomics to discuss (i) the methodology of metabolomics, including sample preparation, targeted metabolomics, and untargeted metabolomics based on nuclear magnetic resonance (NMR) or mass spectrometry (MS), and (ii) applications of metabolomics in the research and practice of life science and medical science.

Metabolomics is an important aspect of phenomics, which is the theory and methodology to study metabolome, including identification of biochemical and molecular characteristics of metabolome, characterization of interactions among




Figure 1.
The imbalance contribution of multiomics to clinical practice. RNAome includes messenger RNAs (mRNAs) and non-coding RNAs (ncRNA). Multiple modifications extensively occur at three different levels of DNAs, RNAs, and proteins to systematically regulate physiological and pathological processes. The center of multiomics is being moved from genomics to phenomics, especially proteomics and metabolomics. PTMs = post-translational modifications. PPPM = predictive, preventive and personalized medicine (3P medicine). Modified and upgraded from Zhan et al. [1], with permission from Elsevier publisher, copyright 2018; and reproduced from Li et al. [2], with permission from Wiley publisher, copyright 2021.

different metabolites or between metabolites and genetic/environmental factors, and evaluation of biochemical mechanisms related to a given condition such as different pathophysiological processes [1]. Metabolome contains all metabolites derived from nucleic acids, proteins, lipids, and sugars in a given cell, tissue, biological system, or body-fluid. The metabolites in a metabolome interact mutually in enzymatic reaction systems to form metabolic network systems. The metabolomic variation is associated with multiple factors, including genetic, environmental, internal, external, drug, or dietary factors [1]. Currently the studies on metabolomic variations are much insufficient in the width and depth of metabolomics. It is necessary to develop high-sensitivity, high-throughput, and high-reproducibility methodology for maximizing the coverage of metabolomic variations. The studies on metabolomic variations directly result in the discovery of effective biomarkers to clarify molecular mechanisms of a disease, determine reliable therapeutic targets, and discover reliable biomarkers for precise prediction, diagnosis, and prognostic assessment in the context of predictive, preventive and personalized medicine (3P medicine, PPPM).

2. Importance of metabolomic variations in medical science

Metabolome contains all metabolites derived from nucleic acids, proteins, lipids, and sugars in a given cell, tissue, biological system, or body-fluid [4–6]. The metabolites in a metabolome interact mutually in enzymatic reaction systems to form metabolic network systems [5]. The change of metabolites is associated with multiple factors, including internal, external, genetic, environmental, drug, or dietary factors. Metabolomics is the theory and methodology to study metabolome, including identifying biochemical and molecular characteristics of metabolome, characterizing interactions among different metabolites or between metabolites and genetic/environmental factors, and evaluating biochemical mechanisms related to a given condition such as different pathophysiological processes [7]. Metabolomic variations can reflect the status of physiological and pathological processes, monitor the progression of a disease, and predict and assess the drug effects compared to the baseline of metabolic profiles, which benefits for disease stratification, and personalized/precise medicine in the context of PPPM [8].

3. Samples used to measure metabolomic variations

The biological samples are very intricate that are used to measure metabolomic variations, including extracts from different cells, tissues, and body-fluids (**Table 1**). Urine and serum/plasma [6, 17, 18] are the most commonly used body-fluids to analyze metabolome for different diseases because these samples are very easily available and are easy to be prepared, without any injury. In addition, tears [19] are the good samples for analyzing metabolome of an eye disease, exhaled air [20, 21] for pulmonary and airway diseases or other diseases, saliva [22] for oral diseases, synovial fluid [23] for arthritis, and cerebrospinal fluid (CSF) [24] for neurological systems disease. Generally speaking, there are many biological samples that are suitable for metabolomics analysis of a disease. The metabolomics studies based on these different samples can directly or indirectly reflect the status of a disease, which may use to understand the molecular mechanism of a disease, and discover therapeutic targets and reliable biomarkers to predict, diagnose, and prognostically evaluate a disease.

Biological sample	Methods	Main results	References
HeLa cells	Gas cluster ion beam-secondary ion MS (GCIB-SIMS)	Purinosomes comprise nine enzymes that act synergistically, channeling the pathway intermediates to synthesize purine nucleotides, increasing the pathway flux, and influencing the adenosine monophosphate/guanosine monophosphate ratio.	[9]
Carcinoma and adjacent normal tissues	UHPLC-Orbitrap MS	This method enables targeted profiling of over 400 biologically important metabolites covering 92 metabolic pathways	[10]
Sweat	GC-MS and LC-MS/MS	As most of the identified metabolites are involved in key biochemical pathways, this study opens interesting possibilities to the use of dry sweat as a source of metabolite markers for specific disorders.	[11]
Urine and plasma	HPLC-ESI-qTOF-MS	A total of 31 and 38 metabolites in plasma and urine, respectively, showed significant differences between healthy volunteers and Sjögren's Syndrome patients and were proposed for their identification.	[12]
Cerebrospinal fluid (CSF)	GC-MS and LC-MS/MS	A total of 274 CSF-derived metabolites were common to the discovery and replication cohorts in cancer-related fatigue.	[13]
Saliva	UHPLC-qTOF-MS	The study identified and classified a total of 211 endogenous and exogenous salivary metabolites. The results reveal a distinct metabolite profile of dog and human saliva as 25 lipid compounds were identified only in canine saliva and eight dipeptides only in human saliva.	[14]
Sputum	LC-MS/MS	The KEGG analysis revealed that the glycerophospholipid metabolism pathway was downregulated in severe COPD. Due to the critical role of glycerophospholipid metabolism in oxidative stress, significant negative correlations were discovered between glycerophospholipid metabolites and three oxidative stress products (SOD, MPO, and 8-iso-PGF2 α). The diagnostic values of SOD, MPO, and 8-iso-PGF2 α in induced sputum were found to exhibit high sensitivities and specificities in the prediction of COPD severity.	[15]
Blood	1D 1H NMR spectroscopy methods	This has led to the absolute quantitation of nearly 70 metabolites in serum and plasma and nearly 80 in whole blood.	[16]

Table 1.
Examples of different types of biological samples used for metabolomics analysis.

4. Methods used to measure metabolomic variations

The appropriate analytical methods for metabolomics are important to detect, identify, and quantify metabolomic variations in a given condition; for example, a disease status versus control, which are mainly classified into targeted metabolomics [25] and untargeted metabolomics [26]. (i) The targeted metabolomics [25] is to mainly quantify hypothesis-driven known metabolite variations in a metabolome (such as

metabolites derived from one or more unknown metabolism pathways) between or among research groups, and then use multivariate statistical analysis to establish mathematical models [27]. This mathematical model then is used to discriminate Diseases from healthy controls, treatment from untreatment, or different stages of diseases. The often used methods for targeted metabolomics are the selected/multiple reaction monitoring (SRM/MRM) analysis with an optimized sample extraction and liquid chromatography-mass spectrometry (LC-MS) conditions using the triple quadrupole mass spectrometry (QqQ-MS) [28]. (ii) The untargeted metabolomics [26] is an none hypothesis-driven approach to globally detect, identify, and quantify metabolite variations in a metabolome in a biological system without any bias, which will benefit the understanding molecular mechanism of a disease, discover new therapeutic targets/drugs and metabolite biomarkers for effective prediction, diagnosis, and prognosis. The often used methods for untargeted metabolomics are the mass spectrometry (MS)-based methods [6, 29], and nuclear magnetic resonance (NMR)-based methods [30, 31] (**Figure 2**). (a) MS-based methods have ion mobility coupled with MS (IM-MS) that can measure time, mass-to-charge (m/z) and intensity variables [1], capillary electrophoresis coupled with MS (CE-MS) that can measure time, m/z and intensity variables [29, 32, 33], gas chromatography coupled with MS (GC-MS) that can measure time, m/z and intensity variables [29, 34], liquid chromatography coupled with MS (LC-MS) that can measure retention time (RT), m/z and intensity variables [26, 29, 35], and direct injection coupled with MS (DI-MS) that can measure m/z and intensity variables [1]. IM-MS is to use a buffer gas and a uniform or periodic electric field for separation of ions based on size and shape of the ions, followed by MS analysis. This is a very high throughput and high selectivity method, which can easily separate isomeric and isobaric compounds. CE-MS is to use electro kinetics for




Figure 2.
The main metabolomic strategies for identification of metabolite profiling and discovery of biomarkers.
Reproduced from Zhan et al. [1], with permission from Elsevier Publisher, copyright 2018.

separation of polar molecules, followed by MS analysis. This is a very good method to analyze polar molecules in aqueous samples for measurement of inorganic and organic anions, with low running costs and relatively low throughput. GC-MS is to use gas chromatography for separation of molecules, followed by MS analysis. This method is suitable for a polar and volatiles compounds, whose advantages are availability of universal database for identification, high sensitivity, and high reproducibility; and whose disadvantages are only detection of a polar and volatile compounds, requirement of derivatization of polar compounds, low ionization discrimination, and requirement of higher amount of samples. LC-MS is to use liquid chromatography for separation of molecules, followed by MS analysis. This method is suitable for polar to hydrophobic compounds, whose advantages are requirement of minimal amount of samples, high sensitivity, high throughput, and flexibility in column chemistry widening the range of detectable compounds; and whose disadvantages are requirement of high ionization discrimination, lack of large metabolite databases, and requirement of specific chromatographic conditions for very polar molecules. DI-MS is to use the nanospray source directly coupled with MS, which does not require chromatography separation, whose advantages are low sample volume requirement, high sensitivity, high-throughput, and low cost; and whose disadvantages are requirement of high ionization discrimination, significant ion suppression phenomenon, and inability to separate isomers and isobaric species. (b) NMR-based methods have one-dimensional, two-dimensional, and three-dimensional NMR methods (1D-NMR, 2D-NMR, and 3D-NMR) [31], which is to use the interaction of spin active nuclei (^{13}C , ^1H , ^{31}P , ^{19}F) in the electromagnetic fields for obtaining structural, chemical, and molecular environment information [30, 31], whose advantages are non-destruction of sample, minimal sample preparation, high reproducibility, relative high throughput, availability of molecular dynamic and compartmental information with diffusional methods, and availability of databases; and whose disadvantages are low sensitivity, overlapping of metabolites, and high instrumentation cost [36]. MS-based methods and NMR-based methods are complimentary for metabolomics analysis, and both will produce very complex data. The processing, analysis, and annotation of data are very important and crucial steps to discover the potential and important metabolic biomarkers [37, 38]. However, compared to the NMR-based metabolomics, MS-based metabolomics has a relatively low cost, high sensitivity and resolution, and very good analytical performance to measure the metabolomic variations for PPPM or PM practice [39].

5. Applications of metabolomics in life science and medical science

Metabolome is the important content of phenome. Metabolomics conducts qualitative and quantitative analysis of all small molecule metabolites in organisms, and searches for the relative relationship between metabolites and physiological and pathological changes. The subjects are mostly small molecules with molecular weights of less than 1,000. With the development of high throughput technology, the study of living organisms has developed from single small molecule to multi-omics; such as genomics, transcriptomics, proteomics, metabolomics. Multiomics reflects molecular changes in a disease or biological process, and molecules that can be identified can be used as valuable biomarkers. Metabolites are substances produced or consumed through the metabolic process. Metabolites are the final expression products subject to genetic control and environmental influence. Imprints with genomic, transcriptomic, epigenetic and environmental effects are called “associations between genotypes and phenotypes” [40]. Metabolomics has been extensively applied in fields of medical science and life science (Table 2). It has important applications in medicine and life sciences, agriculture, food safety and so on. Metabolites,

Metabolomics methods	Biological samples	Main discoveries	References
1D-NMR	Plasma samples from SARS-CoV-2 rRT-PCR-positive patients (n = 15, with multiple sampling timepoints) and age-matched healthy controls (n = 34, confirmed rRT-PCR negative), together with patients with COVID-19/ influenza-like clinical symptoms who tested SARS-CoV-2 negative (n = 35).	The study observed four plasma cytokine clusters that expressed complex differential statistical relationships with multiple lipoproteins and metabolites. These included the following: cluster 1, comprising MIP-1 β , SDF-1 α , IL-22, and IL-1 α , which correlated with multiple increased LDL and VLDL subfractions; cluster 2, including IL-10 and IL-17A, which was only weakly linked to the lipoprotein profile; cluster 3, which included IL-8 and MCP-1 and were inversely correlated with multiple lipoproteins. IL-18, IL-6, and IFN- γ together with IP-10 and RANTES exhibited strong positive correlations with LDL1–4 subfractions and negative correlations with multiple HDL subfractions.	[41]
2D-NMR	Different aging regimes (crust from dry-aged beef, inner edible flesh of dry-aged beef, and wet-aged beef striploin)	NMR-based multivariable analyses could be used to distinguish the method, degree, and doneness of beef aging.	[42]
3D-NMR	<i>E. coli</i> cell lysate	For 19 of the 25 model metabolites, “Structure of unknown metabolomic mixture components by MS/NMR” yielded complete structures that matched those in the mixture independent of database information.	[43]
DI-MS	A parasite–host cell system	The study applied a metabolic fingerprinting approach to evaluate metabolic changes induced by six different (candidate) drugs in a parasite–host cell system.	[44]
LC–MS	<i>P. aeruginosa</i> (35 clinical strains)	Those clinical strains that differed in their virulence and biofilm phenotype also had pronounced divergence in their metabolomes, as underlined by 332 features that were significantly differentially abundant with fold changes greater than 1.5 in both directions.	[45]
GC–MS	Embryonic zebrafish	A total of 87 important endogenous metabolites such as citric acid and hypoxanthine were identified by universal databases or standards among 270 extracted metabolites, which consisted of sugars, amines, amino acids, nucleotides, fatty acids, and sterols.	[46]
CE-MS	Plasma samples of acute corneal seizure mouse model.	Both electrically induced seizures showed decreased values of methionine, lysine, glycine, phenylalanine, citrulline, 3-methyladenine and histidine in mice plasma. However, a second provoked seizure, 13 days later, showed a less pronounced decrease of the mean concentrations of these plasma metabolites, demonstrated by higher fold change ratios.	[47]

Metabolomics methods	Biological samples	Main discoveries	References
IM-MS	Breast cancer plasma samples	Analysis of the resulting data showed that phosphatidylcholines, triglycerides and diglycerides exhibited lower expression and phosphatidylserine showed increased expression in the breast cancer samples compared to those of healthy subjects. The coefficients of variation, determined by reference to the QC data, for all of the features identified as potential markers of disease, were 6% or less.	[48]

Table 2.
Examples of different metabolomics applied in life science and medical science.

as the end products of gene expression, have been implicated in many diseases. For example, metabolomics has great potential for diabetes research, metabolic markers hold the potential to detect diabetes-related complications already under subclinical conditions in the general population [49]. Metabolomics is used to identify key disease-related metabolic changes and disease-progression-related changes, and defining metabolic changes during AD disease trajectory and its relationship to clinical phenotypes provided a powerful roadmap for drug and biomarker discovery [50]. Carmen Peña-Bautista's work shows that the untargeted analysis carried out in human plasma samples from early Alzheimer's disease patients and healthy individuals, and the use of sophisticated statistical tools, identified some metabolic pathways and plasma biomarkers [51]. Nina P Paynter's work shows metabolomics also has important applications in cancer. The processes of life accompany metabolism, such as glycolysis, protein synthesis and metabolism. These fundamental features of cellular metabolism are reprogrammed in cancer cells to support their pathological levels of growth and proliferation. Metabolic reprogramming in malignant cells is likely the result of the multifactorial effects of genomic alterations (i.e. mutations of oncogenes and tumor suppressors), the tumor microenvironment (which imposes metabolic stress caused by compromised nutrients and oxygen availability), and other influences [52]. These changes may be the result of changes in the genome or environmental impacts and a variety of other factors. We need to understand the complete breadth of metabolic abnormalities in cancer because some metabolic changes provide opportunities to develop novel therapeutic targets and predictive biomarkers [52]. As mentioned in Yousra Ahmed-Salim's study, generally, combinations of more than one significant metabolite as a panel, in different studies, achieved a higher sensitivity and specificity for diagnosis than a single metabolite [53]. Metabolomics has become the most powerful platform for studying tissue samples. A common application of metabolomics is the discovery of biomarkers for diagnosis or prediction of treatment sensitivity and prognosis. For example, Yousra Ahmed-Salim et al. conducted a systematic review of the application of metabolomics in the treatment of ovarian cancer. The most frequently described metabolite difference between the biological fluids and tissues of patients with ovarian cancer and those of healthy controls have been in phospholipids [53]. Su et al. interrogated metabolomics and gene-expression from the NCI-60 cell lines to study relationships between metabolite and transcripts [54]. They observed that the metabolome can distinguish cancer subtypes and that metabolite levels correlate well with gene expression under strong correlation models [54]. In conclusion, metabolomics can more accurately determine pathophysiological changes of diseases and identify effective biomarkers through

the high-throughput study of metabolites in organisms with abundant sources of samples, so as to further understand the molecular mechanism of diseases. Thus, it is beneficial to the prevention, diagnosis and treatment of diseases.

6. Conclusion

Metabolomics as the important aspect of phenomics is emerging as the frontier field in life science and medical science. Many biological samples have been used to measure metabolomic variations, including extracts from different cells, tissues, organisms, and body-fluids (for example, urine, serum/plasma, tear, exhaled air, saliva, synovial fluid, CSF, and sputum). Metabolomics is classified into targeted metabolomics and untargeted metabolomics. Targeted metabolomics is used to analyze the known metabolite profiling with SRM/MRM methods. Untargeted metabolomics is used to globally analyze the unknown metabolite profiling with NMR-based methods (1D-NMR, 2D-NMR, and 3D-NMR) and MS-based methods (DI-MS, LC-MS, GC-MS, CE-MS, and IM-MS). Metabolomics has been extensively applied in the research and practice of life science and medical science. However, currently the studies on metabolomic variations are much insufficient in the width and depth. The development of high-sensitivity, high-throughput, and high-reproducibility methodology is needed to maximize the coverage of metabolomic variations for clarification of molecular mechanism of a disease, determination of effective therapeutic targets, and discovery of reliable biomarkers for prediction, diagnosis, and prognostic assessment in the context of PPPM practice.

Acknowledgements

The authors acknowledge the financial supports from the Shandong First Medical University Talent Introduction Funds (to X.Z.), the Hunan Provincial Hundred Talent Plan (to X.Z.), and the Hunan Provincial High-Level Health Talents “225” Plan – Medical Academic Leader Funds (to X.Z.).

Author’s contributions

X.Z. conceived the concept, designed the manuscript, wrote and critically revised the manuscript, coordinated and was responsible for the correspondence work and financial support. J.Y., S.Z., N.L., and N.L. participated in the literature analysis, and wrote partial manuscript.

Conflict of interest

We declare that we have no financial and personal relationships with other people or organizations.

Acronyms and abbreviations

CE-MS	Capillary electrophoresis coupled with mass spectrometry
CSF	Cerebrospinal fluid
DI-MS	Direct injection coupled with mass spectrometry

GC-MS	Gas chromatography coupled with mass spectrometry
IM-MS	Ion mobility coupled with mass spectrometry
LC-MS	liquid chromatography coupled with mass spectrometry
LC	Liquid chromatography
MS	Mass spectrometry
m/z	Mass-to-charge
NMR	Nuclear magnetic resonance
QqQ-MS	Triple quadrupole mass spectrometry
PPPM	Predictive, preventive and personalized medicine
PTM	Posttranslational modification
SRM/MRM	Selected/multiple reaction monitoring
1D-NMR	One-dimensional NMR
2D-NMR	Two-dimensional NMR
3D-NMR	Three-dimensional NMR
3P medicine/PPPM	Predictive, preventive and personalized medicine

Author details

Xianquan Zhan^{1,2*}, Jingru Yang², Shu Zheng², Nannan Li^{1,2} and Na Li²

1 Shandong Key Laboratory of Radiation Oncology, Cancer Hospital of Shandong First Medical University, Jinan, Shandong, P.R. China

2 Medical Science and Technology Innovation Center, Shandong First Medical University, Jinan, Shandong, P.R. China

*Address all correspondence to: yjzhan2011@gmail.com

IntechOpen

© 2021 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

References

- [1] Zhan X, Long Y, Lu M. Exploration of variations in proteome and metabolome for predictive diagnostics and personalized treatment algorithms: Innovative approach and examples for potential clinical application. *J Proteomics*, 2018, 188: 30-40. DOI: 10.1016/j.jprot.2017.08.020.
- [2] Li N, Desiderio DM, Zhan X. The use of mass spectrometry in a proteome-centered multiomics study of human pituitary adenomas. *Mass Spectrom Rev*, 2021, 40: 1-50. DOI:10.1002/mas.21710.
- [3] Zhan X (ed.). *Proteoforms: Concept and Applications in Medical Sciences*. InTech - Open science publisher, London, United Kingdom. Published: July 15th 2020. ISBN: 978-1-83880-034-5. Copyright year: 2020. DOI: 10.5772/intechopen.83687.
- [4] Nicholson JK, Lindon JC, Holmes E. "Metabonomics": understanding the metabolic responses of living systems to pathophysiological stimuli via multivariate statistical analysis of biological NMR spectroscopic data. *Xenobiotica*, 1999, 29: 1181-1189. DOI: 10.1080/004982599238047.
- [5] Dunn WB, Bradhurst DI, Atherton HJ, Goodacre R, Griffin JL. Systems level studies of mammalian metabolomes: the roles of mass spectrometry and nuclear magnetic resonance spectroscopy. *Chem Soc Rev*, 2011, 40: 387-426. DOI: 10.1039/b906712b
- [6] Khamis MM, Adamko DJ, El-Aneed A. Mass spectrometric based approaches in urine metabolomics and biomarker discovery. *Mass Spectrom Rev*, 2017, 36: 115-134. DOI: 10.1002/mas.21455
- [7] Tebani A, Abily-Donval L, Afonso C, Marret S, Bekri S. Clinical metabolomics: the new metabolic window for inborn errors of metabolism investigations in the post-genomic era. *Int J Mol Sci*, 2016, 17: 1167. DOI: 10.3390/ijms17071167
- [8] Everett JR. Pharmacometabonomics in humans: a new tool for personalized medicine. *Pharmacogenomics*, 2015, 16: 737-754. DOI: 10.2217/pgs.15.20
- [9] Pareek V, Tian H, Winograd N, Benkovic SJ. Metabolomics and mass spectrometry imaging reveal channeled de novo purine synthesis in cells. *Science*, 2020, 368(6488): 283-290. doi: 10.1126/science.aaz6465.
- [10] Cao G, Song Z, Hong Y, Yang Z, Song Y, Chen Z, Chen Z, Cai Z. Large-scale targeted metabolomics method for metabolite profiling of human samples. *Anal Chim Acta*, 2020, 1125: 144-151. doi: 10.1016/j.aca.2020.05.053.
- [11] Delgado-Povedano MM, Castillo-Peinado LS, Calderón-Santiago M, Luque de Castro MD, Priego-Capote F. Dry sweat as sample for metabolomics analysis. *Talanta*, 2020, 208: 120428. doi: 10.1016/j.talanta.2019.120428.
- [12] Fernández-Ochoa Á, Borrás-Linares I, Quirantes-Piné R, Alarcón-Riquelme ME, Beretta L, Segura-Carretero A; Precisesads Clinical Consortium. Discovering new metabolite alterations in primary sjögren's syndrome in urinary and plasma samples using an HPLC-ESI-QTOF-MS methodology. *J Pharm Biomed Anal*, 2020, 179: 112999. doi: 10.1016/j.jpba.2019.112999.
- [13] Brown AL, Sok P, Taylor O, Woodhouse JP, Bernhardt MB, Raghubar KP, Kahalley LS, Lupo PJ, Hockenberry MJ, Scheurer ME. Cerebrospinal Fluid Metabolomic Profiles Associated With Fatigue During Treatment for Pediatric Acute

Lymphoblastic Leukemia. *J Pain Symptom Manage*, 2021, 61(3): 464-473. doi: 10.1016/j.jpainsymman.2020.08.030.

[14] Turunen S, Puurunen J, Auriola S, Kullaa AM, Kärkkäinen O, Lohi H, Hanhineva K. Metabolome of canine and human saliva: a non-targeted metabolomics study. *Metabolomics*, 2020, 16(9): 90. doi: 10.1007/s11306-020-01711-0.

[15] Zhu T, Li S, Wang J, Liu C, Gao L, Zeng Y, Mao R, Cui B, Ji H, Chen Z. Induced sputum metabolomic profiles and oxidative stress are associated with chronic obstructive pulmonary disease (COPD) severity: potential use for predictive, preventive, and personalized medicine. *EPMA J*, 2020, 11(4): 645-659. doi: 10.1007/s13167-020-00227-w.

[16] Nagana Gowda GA, Raftery D. Analysis of Plasma, Serum, and Whole Blood Metabolites Using ¹H NMR Spectroscopy. *Methods Mol Biol*, 2019, 2037: 17-34. doi: 10.1007/978-1-4939-9690-2_2.

[17] Want EJ, Wilson ID, Gika H, Theodoridis G, Plumb RS, Shockcor J, Holmes E, Nicholson JK. Global metabolic profiling procedures for urine using UPLC-MS. *Nat Protoc*, 2010, 5: 1005-1018. DOI: 10.1038/nprot.2010.50

[18] Dunn WB, Broadhurst D, Begley P, Zelena E, Francis-McIntyre S, Anderson N, Brown M, Knowles JD, Halsall A, Haselden JN, Nicholls AW, Wilson ID, Kell DB, Goodacre R. Human Serum Metabolome (HUSERMET) Consortium: procedures for large-scale metabolic profiling of serum and plasma using gas chromatography and liquid chromatography coupled to mass spectrometry. *Nat Protoc*, 2011, 6: 1060-1083. DOI: 10.1038/nprot.2011.335

[19] Ahamad SR, Raish M, Yaqoob SH, Khan A, Shakeel F. Metabolomics and

trace element analysis of camel tear by GC-MS and ICP-MS. *Biol Trace Elem Res*, 2017, 177(2): 251-257. DOI: 10.1007/s12011-016-0889-7

[20] Beale DJ, Jones OA, Karpe AV, Dayalan S, Oh DY, Kouremenos KA, Ahmed W, Palombo EA. A review of analytical techniques and their application in disease diagnosis in breathomics and salivaomics research. *Int J Mol Sci*, 2016, 18(1): 24. DOI: 10.3390/ijms18010024

[21] Boots AW, Bos LD, van der Schee MP, van Schooten FJ, Sterk PJ. Exhaled molecular fingerprinting in diagnosis and monitoring: validating volatile promises. *Trends Mol Med*, 2015, 21: 633-644. DOI: 10.1016/j.molmed.2015.08.001

[22] Mikkonen JJ, Singh SP, Herrala M, Lappalainen R, Myllymaa S, Kullaa AM. Salivary metabolomics in the diagnosis of oral cancer and periodontal diseases. *J Periodontal Res*, 2016, 51: 431-437. DOI: 10.1111/jre.12327

[23] Mickiewicz B, Kelly JJ, Ludwig TE, Weljie AM, Wiley JP, Schmidt TA, Vogel HJ. Metabolic analysis of knee synovial fluid as a potential diagnostic approach for osteoarthritis. *J Orthop Res*, 2015, 33: 1631-1638. DOI: 10.1002/jor.22949

[24] Graham SF, Chevallier OP, Roberts D, Hölscher C, Elliott CT, Green BD. Investigation of the human brain metabolome to identify potential markers for early diagnosis and therapeutic targets of Alzheimer's disease. *Anal Chem*, 2013, 85: 1803-1811. DOI: 10.1021/ac303163f

[25] Siskos AP, Jain P, Römisch-Margl W, Bennett M, Achaintre D, Asad Y, Marney L, Richardson L, Koulman A, Griffin JL, Raynaud F, Scalbert A, Adamski J, Prehn C, Keun HC. Interlaboratory reproducibility of a targeted metabolomics platform for

analysis of human serum and plasma. *Anal Chem*, 2017, 89: 656-665. DOI: 10.1021/acs.analchem.6b02930

[26] Mizuno H, Ueda K, Kobayashi Y, Tsuyama N, Todoroki K, Min JZ, Toyo'oka T. The great importance of normalization of LC-MS data for highly-accurate nontargeted metabolomics. *Biomed Chromatogr*, 2017, 31: 1. DOI: 10.1002/bmc.3864. DOI: 10.1002/bmc.3864

[27] Kitteringham NR, Jenkins RE, Lane CS, Elliott VL, Park BK. Multiple reaction monitoring for quantitative biomarker analysis in proteomics and metabolomics. *J Chromatogr B Anal Technol Biomed Life Sci*, 2009, 877: 1229-1239. DOI: 10.1016/j.jchromb.2008.11.013

[28] Zhou J, Yin Y. Strategies for large-scale targeted metabolomics quantification by liquid chromatography-mass spectrometry. *Analyst*, 2016, 141: 6362-6373. DOI: 10.1039/c6an01753c

[29] Naz S, Moreira dos Santos DC, García A, Barbas C. Analytical protocols based on LC-MS, GC-MS and CE-MS for nontargeted metabolomics of biological tissues. *Bioanalysis*, 2014, 6: 1657-1677. DOI: 10.4155/bio.14.119

[30] Marchand J, Martineau E, Guitton Y, Dervilly-Pinel G, Giraudeau P. Multidimensional NMR approaches towards highly resolved, sensitive and highthroughput quantitative metabolomics. *Curr Opin Biotechnol*, 2016, 43: 49-55. DOI: 10.1002/bmc.3864

[31] Kruk J, Doskocz M, Jodłowska E, Zacharzewska A, Łakomiec J, Czaja K, Kujawski J. NMR techniques in metabolomic studies: a quick overview on examples of utilization. *Appl Magn Reson*, 2017, 48: 1-21. DOI: 10.1007/s00723-016-0846-9

[32] Týčová A, Ledvina V, Klepárník K. Recent advances in CE-MS coupling: instrumentation, methodology, and applications. *Electrophoresis*, 2017, 38: 115-134. DOI: 10.1002/elps.201600366

[33] Ramautar R, Somsen GW, de Jong GJ. CE-MS for metabolomics: developments and applications in the period 2014-2016. *Electrophoresis*, 2017, 38: 190-202. DOI: 10.1002/elps.201600370

[34] Mastrangelo A, Ferrarini A, Rey-Stolle F, García A, Barbas C. From sample treatment to biomarker discovery: a tutorial for untargeted metabolomics based on GC-(EI)-Q-MS. *Anal Chim Acta*, 2015, 900: 21-35. DOI: 10.1016/j.aca.2015.10.001

[35] Kohler I, Giera M. Recent advances in liquid-phase separations for clinical metabolomics. *J Sep Sci*, 2017, 40: 93-108. DOI: 10.1002/jssc.201600981

[36] Markley JL, Brüschweiler R, Edison AS, Eghbalnia HR, Powers R, Raftery D, Wishart DS. The future of NMR-based metabolomics. *Curr Opin Biotechnol*, 2016, 43: 34-40. DOI: 10.1016/j.copbio.2016.08.001

[37] Aretz I, Meierhofer D. Advantages and pitfalls of mass spectrometry based metabolome profiling in systems biology. *Int J Mol Sci*, 2016, 17(5): 632. DOI: 10.3390/ijms17050632

[38] Uppal K, Walker DI, Liu K, Li S, Go YM, Jones DP. Computational metabolomics: a framework for the million metabolome. *Chem Res Toxicol*, 2016, 29: 1956-1975. DOI: 10.1021/acs.chemrestox.6b00179

[39] Zampieri M, Sekar K, Zamboni N, Sauer U. Frontiers of high-throughput metabolomics. *Curr Opin Chem Biol*, 2017, 36: 15-23. DOI: 10.1016/j.cbpa.2016.12.006

- [40] Fiehn O. Metabolomics – the link between genotypes and phenotypes. *Plant Mol Biol*, 2002, 48: 155-171.
- [41] Lodge S, Nitschke P, Kimhofer T, Coudert JD, Begum S, Bong SH, Richards T, Edgar D, Raby E, Spraul M, Schaefer H, Lindon JC, Loo RL, Holmes E, Nicholson JK. NMR Spectroscopic Windows on the Systemic Effects of SARS-CoV-2 Infection on Plasma Lipoproteins and Metabolites in Relation to Circulating Cytokines. *J Proteome Res*, 2021, 20(2):1382-1396. doi: 10.1021/acs.jproteome.0c00876.
- [42] Kim HC, Baek KH, Ko YJ, Lee HJ, Yim DG, Jo C. Characteristic Metabolic Changes of the Crust from Dry-Aged Beef Using 2D NMR Spectroscopy. *Molecules*, 2020, 25(13): 3087. doi: 10.3390/molecules25133087.
- [43] Wang C, He L, Li DW, Bruschweiler-Li L, Marshall AG, Bruschweiler R. Accurate Identification of Unknown and Known Metabolic Mixture Components by Combining 3D NMR with Fourier Transform Ion Cyclotron Resonance Tandem Mass Spectrometry. *J Proteome Res*, 2017, 16(10): 3774-3786. doi: 10.1021/acs.jproteome.7b00457.
- [44] Hennig K, Abi-Ghanem J, Bunescu A, Meniche X, Biliaut E, Ouattara AD, Lewis MD, Kelly JM, Braillard S, Courtemanche G, Chatelain E, Béquet F. Metabolomics, lipidomics and proteomics profiling of myoblasts infected with *Trypanosoma cruzi* after treatment with different drugs against Chagas disease. *Metabolomics*, 2019, 15(9): 117. doi: 10.1007/s11306-019-1583-5.
- [45] Depke T, Thöming JG, Kordes A, Häussler S, Brönstrup M. Untargeted LC-MS Metabolomics Differentiates Between Virulent and Avirulent Clinical Strains of *Pseudomonas aeruginosa*. *Biomolecules*. 2020, 10(7): 1041. doi: 10.3390/biom10071041.
- [46] Yan SC, Chen ZF, Zhang H, Chen Y, Qi Z, Liu G, Cai Z. Evaluation and optimization of sample pretreatment for GC/MS-based metabolomics in embryonic zebrafish. *Talanta*. 2020 Jan 15;207:120260. doi: 10.1016/j.talanta.2019.120260. Epub 2019 Aug 14. PMID: 31594598.
- [47] Segers K, Zhang W, Aourz N, Bongaerts J, Declerck S, Mangelings D, Hankemeier T, De Bundel D, Vander Heyden Y, Smolders I, Ramautar R, Van Eeckhaut A. CE-MS metabolic profiling of volume-restricted plasma samples from an acute mouse model for epileptic seizures to discover potentially involved metabolomic features. *Talanta*, 2020, 217: 121107. doi: 10.1016/j.talanta.2020.121107.
- [48] King AM, Trengove RD, Mullin LG, Rainville PD, Isaac G, Plumb RS, Gethings LA, Wilson ID. Rapid profiling method for the analysis of lipids in human plasma using ion mobility enabled-reversed phase-ultra high performance liquid chromatography/mass spectrometry. *J Chromatogr A*, 2020, 1611: 460597. doi: 10.1016/j.chroma.2019.460597.
- [49] Suhre K, Meisinger C, Döring A, Altmaier E, Belcredi P, Gieger C, Illig T. Metabolic footprint of diabetes: A multiplatform metabolomics study in an epidemiological setting. *PloS One*, 2010, 5(11): e13953. doi:10.1371/journal.pone.0013953.
- [50] Toledo JB, Arnold M, Kastenmüller G, Chang R, Baillie RA, Han X, Kaddurah-Daouk R. Metabolic network failures in Alzheimer's disease: A biochemical road map. *Alzheimer's & Dementia*, 2017, 13(9): 965-984. doi:10.1016/j.jalz.2017.01.020.
- [51] Peña-Bautista C, Roca M, Hervás D, Cuevas A, López-Cuevas R, Vento M, Baquero M, García-Blanco A, Cháfer-Pericás C. Plasma metabolomics

in early Alzheimer's disease patients diagnosed with amyloid biomarker. *J Proteomics*, 2019, 200: 144-152. doi: 10.1016/j.jpro.2019.04.008.

[52] Kaushik AK, DeBerardinis RJ. Applications of metabolomics to study cancer metabolism. *Biochim Biophys Acta Rev Cancer*, 2018, 1870(1): 2-14. doi: 10.1016/j.bbcan.2018.04.009.

[53] Ahmed-Salim Y, Galazis N, Bracewell-Milnes T, Phelps DL, Jones BP, Chan M, Munoz-Gonzales MD, Matsuzono T, Smith JR, Yazbek J, Krell J, Ghaem-Maghami S, Saso S. The application of metabolomics in ovarian cancer management: a systematic review. *Int J Gynecol Cancer*, 2021, 31(5): 754-774. doi: 10.1136/ijgc-2020-001862.

[54] Su G, Burant CF, Beecher CW, Athey BD, Meng F. Integrated metabolome and transcriptome analysis of the NCI60 dataset. *BMC Bioinformatics*, 2011, 12 Suppl 1 (Suppl 1): S36. doi: 10.1186/1471-2105-12-S1-S36.