

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



A Review on the Ethnobotanical Uses, Phytochemistry and Pharmacological Effect of *Luffa cylindrica*

Kazeeem Akinyinka Akinwumi, Oluwole Olusoji Eleyowo
and Omolara Omowunmi Oladipo

Abstract

Luffa cylindrica, popularly known as sponge gourd is a tropic and sub-tropical fibrous plant with fruits containing black seeds. The fruit is consumed by humans as a vegetable in many parts of Asia, while different parts of the plant are used for cosmetics and as medicine in many parts of the globe. The plant has been used in the treatment of many ailments including nose cancer, snake venom, wound healing, edema, enterobiasis, filaria, whooping cough, stomach upset, stomach pain and malaria. Many health-promoting compounds such as flavonoids (apigenin-7-glucuronide luteolin-7-O- β -D-glucuronide methyl ester, -O-feruloyl- β -D-glucose, luteolin-7-O- β -D-glucuronide methyl ester), phenolics acids (p-Coumaric, gallic, caffeic, chlorogenic), triterpenoids (oleanolic acid and echinocystic acid), saponins (Lucyoside A-M), tannins (catechin), ribosome-inactivating proteins (α -luffin), carotenoids (9-cis neoxanthin, all-trans-lutein, all-trans- β -carotene), chlorophylls (chlorophyll a and b, pheophytin), cucurbitacin B and gypsogenin have been detected or isolated from different parts of the plants. Extracts of the plant and isolated compounds have wide spectrum pharmacological activities and have been shown to possess antiemetic, antidiabetic, antiviral, wound healing, anticancer, antipyretic, anti-inflammatory, antifungal, anti-bacteria, anthelmintic, hypoglycemic and antihyperglycemic, anti-inflammatory, antioxidant activity, and hepatoprotective effects in animal models. However, further information is needed on its safety and mechanisms of action. The present article is an updated review of the ethnobotanical uses, pharmacological actions, phytochemistry, safety, and future application of *Luffa cylindrica* in translational medicine.

Keywords: *Luffa cylindrica*, medicinal plants, phytochemicals, antioxidant

1. Introduction

Luffa cylindrica is an important edible and medicinal plant that belong to the Cucurbitaceae family. It has many common names including smooth luffa, sponge luffa, vegetable sponge gourd, climbing okra, dishcloth gourd, and Chinese okra [1]. Locally in Nigeria, it is commonly referred to as kankan or kankan oyibo in Yoruba, while the Hausas call it soosoo. The Igbos named it Asisa. The plant is a




Figure 1.
Different parts of *Luffa cylindrica* (A) leaves with flower (B) matured fruits (C) dried fruits (D) *Luffa* sponge (E) seeds.

sub-tropical crop which when planted in northern latitudes warrants hot and humid climates as well as prolonged chilly planting conditions. Also, a vast abseiling shrub with such a greenish, scrumptious vine that is slender but very stiff, extending to a length of 30 feet is *Luffa cylindrica*. It has a fleshy egg-shaped dehiscent fruit with a green papillary dermis, transversely characterized with black crevasses which generally range from 10 to 15 in total. A taut piny filament is discovered under each one of these protrusions [2]. Its leaves are 7–20 cm across and have three lobes. Flowers are bright yellow in color. The fruits which grow to about 60 cm in length are oval-shaped, smooth and, are made up of many seeds. The fruit is brown when matured and dries on the vine to resemble an inedible sponge-like structure. The picture of different parts of the plant is shown in **Figure 1**.

2. Distribution

Luffa cylindrica is commonly found in the humid tropics and Asia. Although the plant is of medieval or primitive origin, it is difficult to decide if Africa or Asia is the ancestral home. In West Africa, the plant grows naturally, but this is also presumed to be a consequence of evasion from planting since the crop is known in many cultures in the area as ‘white people’s sponge.’ Proof of Asian ancestry is however scarce. It is also not clear how well the vine has dispersed over the vast coastal area. Many dispute that rising temperatures are a detonation factor, but the likely cause of the extensive distribution of *Luffa cylindrica* is most often human dispersal. It is documented to have evolved from India but it extensively grows as a weed in Nigeria and many other African countries.

3. Traditional and medicinal uses

Luffa cylindrica has diverse ethnomedicinal uses especially in Africa and Asia. The thawed fiber is used in Ghana for the filtration of water and palm wine [3]. Leaf

formulations are used topically for oedemas and for treating malaria in Togo. The fruit is used on tumor and inflammation in Guinea, while the pulp of the fruit is used as a desiccant in Guinea and Nigeria. There are culinary conifers cultivated in Guinea and Côte d’Ivoire. Zulu people in South Africa take a leaf decoction to treat stomach pain. The root formulation and leaf syrup were documented to be ingested in Tanzania to decrease the probability of pregnancy termination. The leaves are used for the stimulation of wound healing and abscess cognitive development. The leaves are grounded with water and the juice is used for stomach upset medication in Rwanda. Leaf decoctions are used to make childbearing smoother in Uganda. Pulverized leaves are anally inserted for enterobiasis therapy in the Central African Republic. Decoction of the leaves is fully viable against filaria and a colloidal solution of fresh leaves is used to combat whooping cough in Congo-Brazaville. Root formulation is used in Gabon as medicine for nose cancer. A root and leaf aqueous extract is documented to be consumed and used as an aborticide in an enema in the Democratic Republic of the Congo. The seed is used in Egypt for managing diabetes. The pulp of the whole crop is often used as a remedy for acid reflux in African

Country	Local name	Medicinal value	Plant part used	Preparation/application
Togo	Bassarii-Bindumpo, Gudscha, Tem	Oedema and malaria treatment ^a	Leaf	Formulation/ oral
Guinea	Manding-Mandinka	Treatment of tumor, inflammation and as an emollient ^a	Fruit Fruit pulp	
Nigeria	Kan-kan oyibo (Yoruba), Ihion-osa (Edo), sooso (Hausa), asisa (Igbo)	Use as an emollient. ^a	Fruit pulp	
South Africa		Treatment of Stomach pain ^a	Leaf	Decoction/oral
Uganda		Aiding child birth ^a	Leaf	Decoction/oral
Rwanda		Treatment of stomach upset and wounds ^a	Leaf	Decoction/oral Decoction/topical
Tanzania		Anti-abortion ^a	Leaves/root	Formulation/oral
Central African Republic		Enteriobiasis therapy ^a	Leaves	Pulverization/Rectal insertion
Congo-Brazaville		Filaria and whooping cough treatment ^a	Leaves	Decoction/oral
Gabon		Treatment of nose cancer ^a	Root	Formulation/oral
DR Congo		Aborticide ^a	Leaf	Formulation/oral
Egypt		Managing diabetes	Seed	

^a Available at <https://uses.plantnet-project.org>.

Table 1.
Local names and medicinal uses of L.cylindrica in Africa.

indigenous medication. The traditional uses of *Luffa cylindrica* in different parts of Africa are summarized in **Table 1**.

Production of edible forms has taken place in India and the Philippines where the crop is mainly bred. A brand of curry which is produced from the fruit is stripped, chopped and, fried in China and India. The fruit is consumed fresh or diced and processed in Japan for later consumption. The fruit is also employed as a therapy for the treatment of cynocytosis and flu in Asia. Traditional medicine practitioners in China use the seed and sponge of the old fruits of the plant as stomachic, antipyretic and anthelmintic medicine. In addition, dried fruit is used as therapy for abdominal, chest, muscle, and joint pains [4]. Moreover, the fruits are employed in the treatment of rheumatism, dyspnea, cough and skin inflammation in Chinese folk medicine [5]. The fruit reduces breast swellings and it is combined with other Chinese herbs as a remedy against cancer. The fibrovascular bundle of *Luffa cylindrica* dried fruit is officially listed as a treatment for paralytic diseases in Chinese pharmacopeia. In Korea, *Luffa cylindrica* fruit pulp is used to treat fever, induce hemostasis, stimulate menstrual flow, strengthen the network vessels, invigorate blood and clear phlegm [6]. In Japan, the water extract of the vascular bundle of the plant, 'Hechimasui' is used as diuretic, antitussive and skin lotion [7]. In Java-Indonesia, the leaf juice is used for amenorrhea, while it is used for treating snake bites and dysentery in India [8]. The Santals people of Indian, use the plant in treating cramps, convulsion, tetanus, leprosy and syphilis [9]. Oil extracted from the seed is used for treating skin infection, while the fruit or its tincture is used as a therapy against intestinal and biliary colitis, jaundice, hepatomegaly, splenomegaly, dropsy, nephritis, bronchitis, tuberculosis and ascites in Indian herbal medicine practice [9]. The Filipinos used the leaves for orchitis and skin diseases [8].

4. Other ethnomedicinal uses

Raw or prepared as a vegetable, the new fruit can be consumed, but it must be selected before entrenching the spongy cotyledons and before generating the extinguishing substances. The leaves are also consumed as a vegetable, while the charred seeds contain edible oil that is safe for consumption. The gritty and hazardous seed cake is not ideal for livestock feed but could be used as a compost since the plant is rich in nitrogen and phosphorus [3]. The plant is used for treating bowel and bladder hemorrhage, hemorrhoids, toothache, scarlet fever and smallpox [8]. *Luffa cylindrica* seeds are used for treating fever and respiratory disorders including sinusitis, bronchitis and asthma [10, 11]. The seed oil is used as a lubricant and atopically applied to the skin in the treatment of leprosy, shingles, boils and other skin diseases. The oil also found application in several cosmetic products including sunscreens, anti-aging creams, moisturizers, sunless tanners, facial cleansers and sunscreens. The oil is used in sunscreens because of its toxicity to skin cancer cells.

Goats feed on the fruits and leaves [3], while bees prey on their flowers. The root formulations are also used to relieve of stomach pain and as a muscle relaxant. The leaves are used for stimulation of wound repair and echogenic cognitive development. The fruit sag is consumed as a powerful prophylactic, while the seeds are consumed for their anti-parasitic and relaxing properties. The fruit is also used in the treatment of piles and hematuria [12]. Additionally, the fresh fruit is demulcent, cooling and beneficial to the intestine, stomach and genital organs [8]. The flower of *Luffa cylindrica* is used as a therapy against migraine [13].

5. Pharmacological activities

5.1 Antioxidant activity

The methanol and chloroform *Luffa cylindrica* leaves extract exhibited anti-oxidant property via enhanced scavenging of DPPH and superoxide radicals in a dose-dependent fashion [14]. Similarly, its methanol extract displayed free radical scavenging ability against hydrogen peroxide, hydroxyl and nitric oxide radicals. Ethanol extract of the fruit of *Luffa cylindrica* was earlier reported to possess strong antioxidant activity against DPPH radical [15]. Methanol extract of *Luffa cylindrica* vegetable thermally processed by different methods was recently found to show varying degrees of antioxidant properties as measured by thiobarbituric acid, DPPH, ferric thiocyanate and ferric reducing antioxidant power radicals scavenging assays [16]. Similarly, Bulbul *et al.* [17] using DPPH scavaging assay obtained IC₅₀ values of 50.32, 56.27 and 61.24 µg/ml for ethyl acetate, n-hexane and chloroform extracts of the leaves of *Luffa cylindrica* respectively as compared to an IC₅₀ value of 43.22 µg/ml obtained for ascorbic acid, which was used as a standard. *In vivo*, anti-oxidant capacity of the fruit extract of *L. cylindrica* was recently demonstrated in a rat model of cataract. The extract delayed the initiation and inhibit the progression of H₂O₂- induced cataract by inhibiting lipid peroxidation and modulating cellular antioxidants and antioxidant enzyme activity [18].

5.2 Anti-inflammatory activity

Anti-inflammatory activity was exhibited by chloroform extract of *Luffa cylindrica* whole plant through marked reduction of carrageenan-induced rat paw edema in experimental animals that received 50 mg/kg body weight of the extract [19]. Ethyl acetate and ethanol extracts of *Luffa cylindrica* peel and pulp displayed anti-inflammatory action against LPS-induced inflammation in RAW 264.7 cells by modulating NO, IL-6, PGE₂, iNOS, pIκβ and p-ERK expression [20]. Moreover, two fractions from the petroleum ether and benzene extracts of the seed exhibited anti-inflammatory activity in the same experimental animal model [9]. Lucyoside B, a triterpenoid saponin extracted from the fruit of *Luffa cylindrica* also exhibited anti-inflammatory effects through subdual of proinflammatory mediators such as iNOS, IL-6 and MCP-1 at the transcriptional and translational levels coupled with the production of NO [21].

5.3 Anticancer activity

The aqueous-ethanol extract of *Luffa cylindrica* leaves displayed anticancer effects against MCF-7, BT-474, and MDA-MB-231 cell lines which epitomize three sub-types of breast cancer: luminal A, luminal B, and triple-negative [22]. The observed effect was attributed to the presence of phytochemicals such as apigenin and luteolin. The hot water extract of *Luffa cylindrica* whole plant also exhibited anticancer activity against circulating tumor cells of hepatocellular carcinoma especially the cells subpopulation CD133+ /CD44+ with little effect among CD133+ /CD44- subpopulation [23]. Aqueous-ethanol extract of *Luffa cylindrica* leaves showed anticancer activity on three different subtypes of breast cancer including luminal A, luminal B and Her2/neu enriched through reduction of total cell viability, CD44+/24- and total CD24+ cell sub-populations percentages after treatment with the extract [24]. More recently, the anti-cancer activity of hydro-ethanol extract of *Luffa cylindrica* against CD34+/CD38+ and CD34+/CD38+ leukemic stem cells obtained from patients with acute lymphoblastic leukemia was investigated

by Yehia *et al.* [25]. The extract effectively induced cell cycle arrest and apoptosis in both populations of cells as well as exert inhibitory effects against proliferation and colonogenicity of leukemic cells. Aqueous extract of the whole plant displayed cytotoxicity against blood-derived cancer stem cells [23, 24]. The cytotoxic activity of the whole plant ethanol extract to the HT-29 and HCT-15 cell lines has also been documented [26]. The anti-tumor activity of *L. cylindrica* seeds was linked to its luffin content [27].

5.4 Anti-viral effects

The *L. cylindrica* vine demonstrated 66.7–80% protection against Japanese B encephalitis virus when given pre-treatment to mice before viral infection, while the protection diminished when given 210 minutes post-infection to the virus [28]. Luffin P1, a ribosome-inactivating peptide isolated from *Luffa cylindrica* seeds displayed anti-HIV-1 activity in infected C8166 T-cell lines by binding HIV reverse response element and possibly via charge complementation with cellular or viral proteins [29, 30]. Recently, *in silico* analysis revealed that four saponins namely lucyoside H, lucyoside F, 3-O- β -D-glucopyranosyl-oleanolic acid and 3-O- β -D-glucopyranosyl-spinasterol from air-dried fruits of *L. cylindrica* showed strong affinity for the substrate-binding pocket of SARS-CoV-2 Mpro with docking energy scores of -7.54, 7.47, -7.29 and -7.13 kcal/mol, respectively as compared with the binding ability equivalent of N3 protease inhibitor (-7.51 kcal/mol), which is an established inhibitor [31]. Therefore, suggesting that *L. cylindrica* and these aforementioned compounds could find application in the prevention and treatment of SARS-CoV-2.

5.5 Antifungal activity

The ethyl acetate extract of *Luffa cylindrica* leaves displayed antifungal activity against *Candida albicans*, *Candida tropicalis*, *Trichophyton rubrum* together with four clinical isolates of *C. albicans*, *C. tropicalis*, *Microsporum canis* and *Epidermophyton floccosum* [32]. Some compounds isolated from the benzene and petroleum ether of *Luffa cylindrica* seeds also displayed anti-fungal properties against *Candida albicans* [9]. The petroleum ether crude extract of *Luffa cylindrica* fruits exhibited anti-fungal property against *Candida albicans* and *Aspergillus niger* [33]. The butanol extract displayed profound antifungal action against *Trichophyton longifusus* and *Fusarium solani*, while the ethyl acetate fraction of the crude methanol extract markedly inhibited the growth of *Microsporum canis* [34]. *In vivo* anti-fungal activity was also exhibited by crude ethyl acetate extract of *Luffa cylindrica* leaves in laboratory animals by promoting plodding healing of the infected skin of experimental animals [32].

5.6 Antibacterial activity

The petroleum ether extract obtained from *Luffa cylindrica* fruit showed potent antibacterial activity against bacteria *Bacillus cereus*, *Bacillus megaterium*, *Bacillus subtilis*, *Staphylococcus aureus*, *Sarcina lutea*, *Escherichia coli*, *Salmonella typhi*, *Pseudomonas aeruginosa*, *Salmonella paratyphi*, *Shigella dysenteriae*, *Vibrio mimicus*, and *Vibrio parahemolyticus* [33]. Crude methanol and n-hexane fraction of *Luffa cylindrica* also exhibited antibacterial activity against *Bacillus subtilis*, while the butanol fraction exhibited relative activity against *S. flexenari* [34]. The chloroform and n-hexane extract of *Luffa cylindrica* leaves showed potent antibacterial activity against gram-positive and gram-negative bacteria [17].

5.7 Anthelmintic activity

Both crude ethanol and methanol extracts of *Luffa cylindrica* leaves displayed anthelmintic activity against *Pheretima posthuma* [1, 14]. In fact, the anthelmintic activity of the ethanol extract was comparable to the standard drug mebendazole [14].

5.8 Anti-pyretic activity

Methanol extract of *Luffa cylindrica* leaves displayed antipyretic activity by decreasing the rectal temperature of experimental animals at the studied doses and also impeding the compression of abdominal walls in experimental animals, which were induced with agony sensation depending on the dose [35].

5.9 Hypoglycemic and anti-diabetic activity

Methanol extract of *Luffa cylindrica* fruit exhibited excellent hypoglycemic properties in alloxan-induced rat models by decreasing blood glucose level [36]. A significant reduction in blood sugar of glucose-loaded mice after administration of methanol extract of *Luffa cylindrica* fruit further demonstrated the antihyperglycemic activity of *Luffa cylindrica* fruit [37]. Moreover, hydro and ethanol extracts of the fruits exhibited comparable β -cells regeneration with glibenclamide in the alloxan model of diabetes in rats [38]. El-Fiky *et al.* [39] also investigated the effect of oral administration of ethanol seed extract of *Luffa cylindrica* on a streptozotocin rat model of diabetes. The results showed the extract drastically reduced blood glucose level in diabetic rats within three hours of treatment and the efficacy of the extract in reducing blood glucose was similar to a standard anti-diabetic drug, metformin. Tryptic and alcalase protein hydrolysates from the seed have strong inhibitory action against angiotensin-converting enzymes, α -amylase and α -glucosidase [40]. The authors therefore opined that plant holds strong potential in the treatment of hypertension and diabetes.

5.10 Hepatoprotective activity

Methanol extract of *L. cylindrica* leaves displayed hepatoprotective effects through the reduction of serum liver enzymes in a paracetamol model of hepatic injury [41]. The hydroalcoholic extract of *Luffa cylindrica* leaves also exhibited a similar hepatoprotective property in the erythromycin estolate-induced model of liver damage [42]. Increased serum liver enzyme levels in paracetamol-induced rats reduced drastically on treatment with alcohol and aqueous extracts of *Luffa cylindrica* fruits coupled with conservation of the structural integrity of liver membrane [43].

5.11 Sedative and anti-epileptic effects

The sedative, anti-epileptic and anti-convulsant activities of alcohol extract of *L. cylindrica* fruits were investigated in rats by Sunil *et al.* [44]. The results showed that the extract at 400 mg/kg body weight lessened the sleep induction time and prolonged the sleeping time in rats exposed to diazepam. At the same dose, the extract lengthened the latency time, but reduced the time of seizure in the pentyl-enetetrazole-induced model of convulsion, while it decreased total seizure time as well as clonic tonic time in the maximal electroshock model of convulsion. These effects were however lower than that of standard drugs, diazepam and phenytoin.

5.12 Skin protection

Umehara *et al.* [7] investigated the effect of *L.cylindrica* fruit extract on UVB-irradiation-induced mice model of dry skin and demonstrated that the extract and isolated phenylpropanoids inhibited trans-dermal water loss in hairless mice. The extract and p-coumaric acid isolated from the extract stimulated dome formation by MDCK I cells. Additionally, p-coumaric acid increased mRNA expression of water permeability and reabsorption protein, AQP3. The authors, therefore, concluded that p-coumaric was responsible for *L. cylindrica* related water permeability and that *L. cylindrica* could contribute to the treatment of disease relating to the inability to retain moisture including dry syndrome. Furthermore, an extract obtained from the fruit pulp prevented the development of atopic dermatitis-like skin lesions in mice exposed to *Dermatophagoides farinae* [6].

5.13 Anti- emetic activity

The ethanol extract of the fruit peel of *L.cylindrica* showed significant anti-emetic activity in young chicks at a dose of 150 mg/kg body weight [45]. The ethanol and hexane extracts of leaves and male flowers of *L. cylindrica* exhibited anti-emetic effect against chick emesis facsimilia. The anti-emetic effects of hexane extract of male *L. cylindrica* flowers and leaves were at 71.5% and 43.5% inhibition of reteches respectively, whereas the ethanol extract of leaves and male flowers of *L. cylindrica* was at 68.66% and 68.46% inhibition of reteches respectively [46].

5.14 Wound healing activity

Different parts of *L. cylindrica* have been reported to possess wound healing capacity. Chloroform extract of the whole plant showed wound healing activity in a rat model by reducing the wound area and time of epithelization [19]. Diethyl ether, n-hexane, chloroform, ethyl acetate, butanol and methanol seed extract also promoted wound healing in rats [47]. Diethyl ether extract showed the highest wound healing activity, while the weakest activity was displayed by chloroform extract [47].

5.15 Effects on hematological parameters

Raw and thermally processed *Luffa cylindrica* seed meal fed to albino rats had no adverse effects on the hematological indices of the experimental animals [48].

However, administration of methanol extract of *L.cylindrica* leaves to rats produced an elevation in the hematological parameters in experimental animals [49].

5.16 Oxytocic activity

Aqueous extract of *L.cylindrica* leaves increased uterine motility in an isolated rat uterus [50]. Thus suggesting that *L.cylindrica* is oxytocic and giving credence for its use in Uganda to facilitate labour and treat postpartum issues.

6. Chemical composition

L.cylindrica is rich in nutrients and phytochemicals. Several recent studies have revealed that *L. cylindrica* leaves, fruits, and seeds are a rich source of

carbohydrates, protein, fiber, fats, amino acids and minerals [48, 51]. The mineral found in the seeds of *Luffa cylindrica* plant include sodium, iron, phosphorus, calcium, zinc, potassium, manganese, copper, chromium, magnesium [51]. Moderate amounts of K (13.86 mg/100 g) and Na (8.18 mg/100 g) were found in the seed, but the concentration of Cr (0.25 mg/100 g) was low [50]. *L.cylindrica* fruit is an excellent source of Vitamins A, B5, B6, C, and dietary fiber.

According to the preliminary phytochemical screening test carried out on different extracts of *L.cylindrica* leaf and seed extracts, all the extracts were found to contain saponin, alkaloids and cardiac glycosides [52]. Only the seed extracts contained the steroidal rings, while anthraquinones, tannins and phlobatinnins were not detected in any of the extract [52]. Reducing carbohydrates, flavonoids, tannins, saponins and glycosides were also detected in the plant extract [2]. The phytochemical screening carried out on the methanol and ethyl acetate extract of the leaves of *L.cylindrica* indicated that the extracts of leaves of *L.cylindrica* contain carbohydrates, sterols, saponins, flavonoids, alkaloid and phenols, while resins, tannins, terpenes, balsams and anthraquinones were not found [32]. Another preliminary phytochemical screening of aqueous methanol extract of *L.cylindrica* leaves revealed the presence of sugar molecules including glucose, fructose and galactose as well as the presence of amino acids such as phenylalanine, glycine and tyrosine [53]. Moreover, phytate and oxalate were found in the methanol extract of flowers and leaves of *L. cylindrica* [54]. The *L. cylindrica* seeds contain high amount of saponins, alkaloids and phlobotannins. The butanol extract of *Luffa cylindrica* seeds contains alkaloids and deoxy sugars, while the diethyl ether extract contains deoxy sugars, cardiac glycosides, alkaloids and carbohydrates [47]. The chloroform extract contains deoxy sugars and cardiac glycosides [47]. Quantitative analysis of the sponge revealed that it contains 1.2, 0.5, 17.94 and 20.74 mg/g of ascorbic acid, total anthocyanins, flavonoids and phenolics respectively [55]. The total flavonoid and phenol in the aqueous and ethanol extracts of *Luffa* pulp and peels were reported to range from 0.94–14.02 mg/g GAE and 0.33–18.09 mg/g QE respectively, while olenolic acid, carotenoids, and chlorophylls were in the range 0.01–25.79, 0.01–14.87 and 0.04–37.29 mg/g extract respectively [20].

A penotacyclic triterpenoid saponin, lucyoside O was isolated from the leaves of *L.cylindrica*. Lucyin A, lucyosides G, N, O, Q, P, R, ginsenosides Re and Rg1, 21 β - hydroxyoleanic acid and 3-O- β -D-glucopyranosyl- maslinic acid, were also identified in the leaves of *Luffa cylindrica* [22]. In addition, Lucyoside K was isolated from the hydro-ethanol extract of the leaf extract, while Lucyoside A-M were identified in the fruit extract of the plant [22]. A peptide with luf-facyclin that possessed antifungal activity was also isolated from *L. cylindrica* seeds. Sapogenins (I & II) were isolated from ethanol extract of the seeds and were both found to exhibit immunomodulatory effects [56]. Some triterpenoids and fibrinolytic saponins were isolated from seeds and fruits of the plant [47]. Moreover 22, 23-dihydroxy spinasterol and 3-hydroxy-1-methylene-2,3,4,4-tetrahydroxynaphthalene-2-carbaldehyde were separated from the petroleum ether extract of the fruit [57]. Very recently [31] isolated lucyoside F, lucyoside H, 3-O- β -D-glucopyranosyl-spinasterol and 3-O- β -D-glucopyranosyl-oleanolic acid from the dried fruits.

L. cylindrica is very rich in polyphenols. Aqueous ethanol extract of *L. cylindrica* leaves contains phenolics such as apigenin 7 glucuronide, eriodictyol -7 glucoside, kaempferide, luteolin-O-diglucoside, neodiosmin, diosmin and kaempferol 3 - [2'',3'',4''-triacytl - α - L -arabinopyranosyl -(1-6) -glucoside] or its isomer kaempferol 3 -[2'',3'',5''- triacytl - α - L -arabinofuranosyl -(1-6) -glucoside [22].

Similarly, Sunnil *et al.* [44] recently used LC-ESI-MS/MS to identified several flavonoids and polyphenol including hyperoside, kaempferol-3,7-O-bis- α -L-rhamnoside, quercetrin, tiliroside, acacetin, datiscin, fortunellin, linarin, luteolin, bobin, vitexin, vitexin-2''-O-rhamnoside, saponarin from the alcohol extract of the whole fruit of *L. cylindrica*. In addition, Yadav *et al* [16] also identified the following phenolics and flavonoids: gallic acid (0–26.8 $\mu\text{g/ml}$), caffeic acid (0.23–18.4 $\mu\text{g/ml}$), cinnamic acid (1.52–8.6 $\mu\text{g/ml}$), ferulic acid (9.31–49.6 $\mu\text{g/ml}$), ellagic acid (0–78.8 $\mu\text{g/ml}$), rutin (0–79.3 $\mu\text{g/ml}$), quercetin (45.18–55.42), myrecetin (20.95–35.79 $\mu\text{g/ml}$), catechin (66.24–77.87 $\mu\text{g/ml}$) from methanol extract of *L. cylindrica* fruits thermally processed by different methods. Furthermore, Hlel *et al* (2017) using HPLC/TOF-MS identified chlorogenic acid, gentisic acid, gallic acid, vanillic acid, salicylic acid, ferulic acid, 4-hydroxy benzoic acid, p-coumaric acid, naringenin, catechol and rutin in *L. cylindrica* fruits at different stages of maturation. The amount of quercetin, luteolin and myricetin in the sprout extract of *L. cylindrica* was quantified using UPLC-MS/MS as 32.5, 12.5 and 32.4 $\mu\text{g/g}$ respectively. Meanwhile, five derivatives of cinnamic acid including 1-O-p-coumaroyl--D-glucose, 1-O-feruloyl-, -D-glucose, 1-O-caffeoyl--D-glucose and p-coumaric acid well as three flavonoids glycosides namely: apigenin-7-O-, -D-glucuronide methyl ester, diosmetin-7-O--D-glucuronide methyl ester, and luteolin 7-O--D-glucuronide methyl ester) were earlier identified in *Luffa cylindrica* [15]. A phenylpropanoid glucoside, 4-O-feruloyl-glucose was isolated from a natural source for the first time in *L. cylindrica* fruit [7].

Other phenylpropanoid glucosides that were isolated from the edible part of *Luffa cylindrica* are 4-O-caffeoyl-glucose, 1-O-caffeoyl- β -glucose, 6-O-caffeoyl-glucose, 4-O-p-coumaroyl-glucose, 1-O-p-coumaroyl- β -glucose, 6-O-p-coumaroyl-glucose, 4-O-feruloyl-glucose, 1-O-feruloyl- β -glucose, 6-O-feruloyl-glucose [58].

Other compounds there were recently isolated from the dried fruits of *L. cylindrica* includes: 3,5-dihydroxy- δ -valerolactone, phenanthrene, 1,2-naphthoquinone, cinnamic acid, (S)-dehydrovomifoliol, 2,6-dimethyl-1,4-benzenediol, litchiol B, pinoresinol phthalic acid, 4-(hydroxymethyl)benzene-1,2-diol, tridecan-7-one, apigenin and henicosan-11-one [31]. Similarly, fifty-three volatile compounds including aromatics (10.1%), acids (15.1%), ketones (38.2%), alcohols (51.6%) and aldehydes/furans (66.2%) were recently identified by [59] in young and matured fruits of *Luffa cylindrica* using headspace SPME-GS-MS and UPLC-MS. Hydrocarbons including noctacosane, n-heptacosane, n-hexacosane, n-tetracosane, n-tricosane, tetraeicosane-6-ol, nanodecane-6-ol, dieicosane-6-ol and eicosane-6-ol have earlier been identified in the fruit of *L. cylindrica* [12]. The diverse types of phytochemicals found in *L. cylindrica* have various biological effects (Table 2) and could account for its wide pharmacological activities.

6.1 Acute toxicity studies

Etim *et al* [49] administered methanol leave extract of *L. cylindrica* to Swiss albino mice at doses up to 4000 mg/kg body weight. Treated animals did not die nor display signs of toxicity. In addition, there was no mortality in animals exposed to *L. cylindrica* fruit extracts at doses between 100 and 2,000 mg/kg body weight [145]. Similarly, Oyeyemi *et al.* [146] reported that administration of 5000 mg/kg of both hydro-methanol and aqueous extracts of *L. cylindrica* leaves did not induce acute toxicity in mice. However, the extracts given at doses between 200 and 1600 mg/kg increased bone marrow micronucleated polychromatic erythrocytes formation, but of a lower degree to the positive control, methyl methanesulfonate (Oyeyemi *et al*, 2015). An LD₅₀ of 450 mg/kg body weight was reported for crude petroleum ether extract of the fruit [123].

Phyto chemicals	Compounds	Pharmacological action	Ref.
Flavonoids	Apigenin 7 glucuronide,	Anti-oxidant, anti-complement, anti-inflammatory, and aldose reductase inhibitory activities	[60]
	Kaempferol 3 - [2'',3'',4''- triacetyl - α - L -arabinopyranosyl -(1-6) -glucoside],	NRPA	
	Kaempferide,	Anticancer, cardioprotective and Osteo-protective, anti-oxidant, anti-inflammatory, anti-bacterial anti-viral	[61, 62].
	Diosmin,	Anti-ulcer, anti-inflammation, anti-oxidation, anti-diabetes, anti-cancer, anti-microbial, hepato- protective, neuroprotective cardio-protective, nephroprotective, and retinal protection.	[63, 64]
	Neodiosmin, Eriodictyol -7 glucoside,	NRPA Antioxidant, anti-inflammatory, anti-cancer	[65]
	Quercetin	Antioxidant, anti-inflammatory, anticancer, cardio-protective, neuroprotective, pneumo-proective, hepatoprotective	[66, 67]
	Myrecetin	Antioxidant, antitumor, anti-inflammatory, neuroprotective, immunomodulatory, antimicrobial, antiviral, hepatoprotective, anti-obesity, cardiovascular protection	[68, 69]
	Rutin	Antioxidant, anti-inflammatory neuroprotective, antitumor, sedative, anti-convulsant, anti-Alzheimer, anti-cholesteremic, anti-asthmatic, antiosteoporotic, anticataract, immunostimulatory, antimicrobial, antiviral, antihypertensive	[70]
	Catechin	Anticardiovascular, antioxidant, neuroprotection, hepatoprotection, anti-infectious, anti-diabetic	[71]
	Luteolin	Antioxidant, anti-inflammatory, antitumor, antiapoptotic, anti-allergy,	[72]
	Hyperoside	Anticancer, anti-inflammatory, Anti-oxidant, antiparasite, anti-cholesterolemic, cardioprotection, antidepressant, anti-aging.	[73-75]
	Kaempferitrin	antioxidant, anti-inflammatory, antitumor, anti-angiogenic	[76]
	Quercetrin	Antioxidant, antitumor,	[77]
	Tiliroside	Antioxidant, antidiabetic, anti-obesity, anti-inflammatory, hepatoprotective, anti-allergy, anti-thrombotic, neuroprotective osteogenic, antiobesity, antimicrobial, antiviral, antiprotozoal, antihypertensive antiaging	[78, 79]
	Acacetin	antioxidant, anti-inflammatory, anti-depressant, antinociceptive antitumor, neuroprotective	[80-82]
	Datiscin Fortunellin	NRPA Antioxidant, anti-inflammatory, antidiabetic, cardioprotective	[83]
	Linarin	Antioxidant, anti-inflammatory, osteogenic, anti-cholinesterase neuroprotection, analgesic cardioprotective,	[84, 85]
	Robinin	NRPA	
	Vitexin	Antioxidant, neuroprotective anti-inflammatory,antidiabetic, anti-tumor, hepatoprotection, cardio-protective, amtiviral, antibacteria	[86, 87]
	Vitexin-2''-O-rhamnoside	Antioxidant	[88]
	Saponarin	Hepatoprotective, antioxidant anti-lipid peroxidation	[82, 89]

Phyto chemicals	Compounds	Pharmacological action	Ref.
Phenols	p- coumaric acid,	Antioxidant, antiviral, anti-inflammatory, anti-cancer, anti-lipidemic, anti-gout, antimicrobial, immunomodulatory, antiplatelet aggregation, anti-diabetic, anxiolytic, anti-arthritis, antipyretic, analgesic.	[90, 91]
	1-O-feruloyl- β -D-glucose,	Anti-adipogenic	[92]
	1-O-(4- hydroxyl benzoyl) glucose.	NRPA	
	Gallic acid,	Antioxidant, antiobesity, antihyperglycaemic, antidiabetic, anti-lipid peroxidative, wound healing, anti-inflammatory, 10.1016/j.phrs.2018.08.002 neuroprotective, cardioprotective, antimicrobial, gastroprotective	[93]
	Gentisic acid	Antioxidant, anti-inflammatory, neuroprotective, antigenotoxic, hepatoprotective, antimicrobial, anticancer, analgesic, skin-lightening, muscle relaxation, cardioprotective	[94]
	Chlorogenic acid	Antioxidant, cardioprotective, neuroprotective, renoprotective, antidiabetic, antitumour, Gastro-intestinal protection, antitumour	[95]
	4 -hydroxy benzoic acid	NRPA	[96–98]
	Vanillic acid,	Anti-obesity, anti-inflammatory, antioxidant, neuroprotective cardioprotective	
	Salicylic acid,	Anti-inflammatory, analgesic	[99]
	Naringenin,	Antioxidant, antibacterial, antitumor, anti-inflammatory, cardioprotective, antiadipogenic immunomodulatory, antiviral	[100]
	Catechol	Antioxidant, anticancer	[101]
	Caffeic acid	Antioxidant, anticancer, antiviral, antimicrobial, anti-inflammatory, Antidiabetic, cardioprotective, immunostimulatory,	[102]
	Ferulic acid	Antioxidant anti-inflammatory, anticancer, antidiabetic, antimicrobial, antithrombotic, anti-arrhythmic, antidiabetic, immunostimulatory, anti-aging, neuroprotective, photoprotective	[103]
Triterpenoids	Cinnamic acid	Antioxidant, antidiabetic, antimicrobial, anti-melanogenesis, UV-protective	[104, 105]
	Ellagic acid	Antioxidant, hepatoprotective antitumor, antiangiogenic, antimetastatic, anti-inflammatory ,neuroprotective, antidiabetic, anti-atherogenic	[106, 107]
	Oleanolic acid	Antioxidant immunomodulatory, antiviral, antimicrobial, hepatoprotective, cardioprotective anti-inflammatory, analgesic antihypertensive, anticancer, immunostimulatory	[108, 109]
	Echinocystic acid	Antioxidant, anti-inflammatory, antibacterial, antiapoptotic, antiviral, antitumor antioxidant, Immunostimulatory	[110, 111]
	Gypsogenin 3-O- β -D-glucopyranosyl-maslinic acid,	Antitumour Immunostimulatory	[112]
	Dehydrovomifoliol	Anticancer, anticholinesterase	[113, 114]

Phyto chemicals	Compounds	Pharmacological action	Ref.
Saponin	Lucyoside K, Lucyoside O,	NRPA	
	Lucyoside B	Anti-inflammatory	[21, 115];
	Lucyosides N &P Lucyosides A,C-M,R Ginsenosides Re Anti-arrhythmic, modulation of insulin resistance [116–118]	Fibrinolytic activity NRPA	[119]
	Ginsenosides Rg1	Neuroprotective, anti-depressant, anti-inflammatory, anti-sepsis	[27, 120–122]
	21 β - hydroxyoleanoic acid	NRPA	
Cucurbitacins	Cucurbitacin B	Anticancer	[123]
	Cucurbitacin E	Immunomodulatory, anti-inflammatory, neuroprotective anti-tumorigenic,	[124].
Peptides	Luffacyclin	Antifungal	[125]
	Luffin –a Luffin -b	Abotificient, Antitumour, Abotificient, Antitumour	[27, 125] [27, 125]
	Luffins P1,	Anti-HIV, antitumour, antifungal	[29, 30, 125]
	Luffin B	Antitumor, Antiviral	[8]
	Luffins S,	Antitumor, Antiviral	[29, 30, 125]
	Luffin – α	Antitumor	[27]
	Bryonolic acid	Inhibits passive cutaneous anaphylaxis and delayed hypersensitivity	[126]
Phenyl propanoid glucosides	1-O-feruloyl- β -D-glucose, 1-O-caffeoyl- β -D- glucose 4-O-caffeoyl- glucose.	Antioxidant NRPA	[15, 127]
	6-O-caffeoyl- glucose 4-O-p-coumaroyl- glucose 1-O-p-coumaroyl- β -glucose	Antiradical and antioxidant NRPA NRPA	[128]
	6-O-p-coumaroyl- glucose 6-O-feruloyl- glucose 1-O-feruloyl- β - glucose 4-O-feruloyl- glucose	NRPA Antioxidant NRPA	[129]

Phyto chemicals	Compounds	Pharmacological action	Ref.
Volatiles and other compounds	3-hydroxy1-methylene-tetrahydroxy-napthalene-2-carbaldehyde	Antimicrobial	[57]
	22, 23-dihydroxy spinasterol	Antimicrobial	[57]
	Phenanthrene	Antimicrobial, spasmolytic, anti-inflammatory, antiplatelet aggregation, antiallergic activities and phytotoxic effects	[130, 131]
	Litchiol B	Antioxidant,anti-bacteria	[132]
	Pinoresinol	Anti-proliferative, antioxidant	[133]
	Pentanoic acid	NRPA	
	Nonanoic acid	Bioherbicides	[134]
	1-Octen-3-ol	Pesticides	[135]
	β-linalool	Antioxidant, antibacterial	[136]
	α-terpineol Nonanal	Antioxidant, insecticidal, anticancer, anti-nociceptive, anticonvulsant, antihypertensive, antiulcer Antifungal, antidiarrheal	[137, 138]
	Nonadienal	Anti-bacteria	[59]
	Decanal	Antimicrobial	[139]
	Eugenol	Mosquito repellent, antifungal, antibacterial, antiinflammatory antioxidant	[140]
	Limonene	Antitumour, anti-inflammatory, antioxidant, antiviral,antibacterial	[141]
	Trihydroxy-octadecadienoic acid	Antifungal and bacteria	[141, 142]
	Trihydroxy-octadecenoic acid	Antiviral	[143]
	Octadecadienoic acid	Antidiabetic	[144]

NRPA = no reported pharmacological action

Table 2.
Some phytochemicals in Luffa cylindrica and their pharmacological action.

7. Conclusion and future perspective

The review provides an up-to-date and comprehensive summary of the traditional uses, pharmacology and phytochemical composition of *Luffa cylindrica*. *Luffa cylindrica* has been eaten as food and used in folk medicine for several years especially in Africa and Asia for the treatment of many diseases including malaria, stomach disorders, whooping cough, oedemas, wounds, tumor, filarial, rheumatism, dyspnea, inflammation, leprosy, syphilis, bronchitis, tuberculosis, dysentery and amenorrhea. In the last few decades, the plant has attracted attention due to its potential pharmacological actions including anti-inflammatory, anticancer antioxidant, anti-viral, antimicrobial, anti-diabetic, hepatoprotective, sedative, anthelmintic, anti-pyretic, anti-epileptic, hypoglycemic, skin protection. Anti-emetic and wound healing. The repertoire of beneficial and health-promoting phytochemicals that are present in *Luffa cylindrica* could be responsible for diverse

ethnomedicinal uses and pharmacological activity recorded for the plant. However, like many other medicinal plants, efforts should be made to standardize its usage in different disease models through activity-guided bioassays and isolation of active principle(s). Formulated standardization of *Luffa cylindrica* extract is needed to have reproducible results that can be integrated into translation medicine. In addition, more mechanistic and comprehensive safety studies on *Luffa cylindrica* are needed to enhance its pharmaceutical potentials and know the long-term effects of consumption of *L. cylindrica* as medicine since most of the studies found in literature only addressed its acute toxicity. Moreover, more clinical studies are warranted to confirm the pharmacological activities of *Luffa cylindrica* extracts and its constituents in order to translate results obtained from animal studies into human.

Taken together, *Luffa cylindrica* holds great potential as a repository of beneficial phytochemicals that can be leveraged on for the betterment of human health. Research efforts should therefore be directed at optimizing the bioactive extracts and/or phytochemicals for health promotion and improving the quality of life.

Author details


Kazeem Akinyinka Akinwumi^{1*}, Oluwale Olusoji Eleyowo^{1,2}
and Omolara Omowunmi Oladipo¹

¹ Department of Chemical and Food Sciences, Bells University of Technology Ota, Nigeria

² Department of Science Laboratory Technology, Lagos State Polytechnic, Lagos, Nigeria

*Address all correspondence to: qaakinwumi@yahoo.co.uk

IntechOpen

© 2021 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

References

- [1] Partap S, Kumar A, Sharma NK, Jha KK. *Luffa Cylindrica*: An important medicinal plant. *J. Nat. Prod. Plant Resour*, 2012; 2(1): 127-134.
- [2] Gandhamalla P, Shiva GB, Pravalika R, Ramya DM, Boggula Narender. Plant preliminary phytochemical analysis and thrombolytic screening of *Luffa cylindrica* Linn. Fruits an *in-vivo* study. *IJIPSR* 2018, 6 (01):61-742018.
- [3] Achigan-Dako EG, N'danikou, S., Vodouhê, RS, 2011. *in vivo* (L.) M. Roem. Record from PROTA4U. Brink, M. & Achigan-Dako, E.G. (editors). PROTA (plant resources of tropical Africa / Ressources végétales de l'Afrique tropicale), Wageningen, Netherlands. 2011 Available at : <http://www.prota4u.org/search.asp>. Accessed 10 February 2021.
- [4] Duke, J. A., 2002. CRC handbook of medicinal herbs, 2nd ed. 30-35.
- [5] Sangh P, Amit K, Neeraj KS., Jha KK. *Luffa cylindrica* : An important medicinal plant. *J. Nat. Prod. Plant Resour.*, 2012, 2, 127-134. Abdel-salam *et al*, 2019
- [6] Ha H, Lim H.S, Lee M.Y, Shin, IS, Jeon, WY, Kim, JH., Shin, H.K. *Luffa cylindrica* suppresses development of *Dermatophagoides farinae* induced atopic dermatitis like skin lesions in Nc/Nga mice. *Pharmaceutical Biology* 2015, 53(4), 555-562.
- [7] Umehara M, Yamamoto T, Ito R, Nonaka S, Yanae K, Sai M. Effects of phenolic constituents of *Luffa cylindrica* on UVB-damaged mouse skin and on dome formation by MDCK I cells, *Journal of Functional Foods* 2018, 40: 477-483,
- [8] Lim T.K. *Luffa cylindrica*. In: Edible medicinal and non-medicinal plants. Springer 2012, Dordrecht. https://doi.org/10.1007/978-94-007-1764-0_46
- [9] Muthumani P, Meera R, Subin Mary Jeenamathew, P Devi, B Kameswari, B Eswara Priya. Phytochemical screening anti-inflammatory, bronchodilator and antimicrobial activities of the seeds of *Luffa cylindrica*. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*, 2010; 1(4): 11-22.
- [10] Simeon, K. A., Imeh, J., Gbola, O. (2017). Plants in respiratory disorders I anti asthmatics, a review. *British Journal of Pharmaceutical Research*, 16(2), 1-22.
- [11] Stephens, J.M., 2003. Ground Luffa. *J. hort. Sci.*, 3, 19-21.
- [12] Nirmal SA, Kothawade PC. Datir SB, Pal SC, Mandal SC Pattan S.R. Nonpolar compounds from *Luffa cylindrica*. *FACTA UNIVERSITATIS Series: Physics, Chemistry and Technology* 2009, 7(1): 69 - 72
- [13] Sutharshana V. (2013). Protective role of *Luffa cylindrica*. *Journal of Pharmaceutical Sciences and Research*, 5(9), 184 – 186
- [14] Tripathi A, Tandon M, Chandekar A, Soni N Upmanyu N. *In vitro* antioxidant and anthelmintic activity on *Luffa cylindrica* leaf extract, *Journal of Herbs, Spices & Medicinal Plants*, 22:4, 348-355 2016
- [15] Du Q, Xu Y, Li L, Zhao Y, Jerz G, Winterhalter P. Antioxidant constituents in the fruits of *Luffa cylindrica* (L.) Roem. *J Agric Food Chem* 2006, 54(12): 4186 – 4190 .
- [16] Yadav, R., Yadav, B. S., & Yadav, R. B. (2016). Phenolic profile and antioxidant activity of thermally processed sponge gourd (*Luffa cylindrica*) as studied by using high performance thin layer chromatography

(HPTLC). International Journal of Food Properties 20(9):2096-2112

[17] Bulbul, J., Zulfiker, A., Hamid, K., Khatun, M., and Begum, Y..Comparative study of in vitro antioxidant, antibacterial and cytotoxic activity of two bangladeshi medicinal plants-*Luffa cylindrica* L. and *Luffa acutangula*. Pharmacogn. J 2011;3:59-66.

[18] Dubey S, Saha S, Kaithwas G and Saraf SA. Effect of standardized fruit extract of *Luffa cylindrica* on oxidative stress markers in hydrogen peroxide induced cataract. Indian J Pharmacol 2015;47:644-648.

[19] Abirami MS, Indhumathy R, Devi GS, Kumar DS, Sudarvoli M and Nandini R. Evaluation of the wound healing and anti-inflammatory activity of whole plant of *Luffa cylindrica* (Linn) in Rats. Pharmacologyonline, 2011; 3: 281-285

[20] Kao TH, Huang CW and Chen BH, Functional components in *Luffa cylindrica* and their effects on anti-inflammation of macrophage cells. Food Chem 2012;135:386– 395

[21] Han Y, Zhang X, Qi R, Li X, Gao Y, Zou Z, Cai R, Qi Y. Lucyoside B, a triterpenoid saponin from *Luffa cylindrica*, inhibits the production of inflammatory mediators via both nuclear factor- κ B and activator protein-1 pathways in activated macrophages. Journal of Functional Foods 2020;69 doi.org/10.1016/j.jff.2020.103941

[22] Abdel-Salam IM, Ashmawy AM, Hilal AM, Eldahshan OA and Ashour M. Chemical composition of aqueous ethanol extract of *Luffa cylindrica* leaves and its effect on representation of caspase-8, caspase-3, and the proliferation marker Ki67 in intrinsic molecular subtypes of breast cancer in vitro. Chem Biodivers 2018; 15(8):e1800045. doi: 10.1002/cbdv.201800045

[23] Abdel-Salam IM^A, Awadein NE, Ashour M. Cytotoxicity of *Luffa cylindrica* (L.) M. Roem. Extract against circulating cancer stem cells in hepatocellular carcinoma. J Ethnopharmacol 2019a; 229:89-96.

[24] Abdel-Salam IMB, Abou-Bakr AA, Ashour M. Cytotoxic effect of aqueous ethanol extract of *Luffa cylindrica* leaves on cancer stem cells CD44+/24- in breast cancer patients with various molecular sub-types using tissue samples *in vitro*. J. Ethnopharmacol. 2019;238:111877. doi: 10.1016/j.jep.2019.111877.

[25] Yehia S, Abdel-Salam IM., El-agamy B and Aldesouki HM. Cytotoxic and apoptotic effects of *Luffa cylindrica* leaves extract against acute lymphoblastic leukemic stem cells. Asian Pacific Journal of Cancer Prevention 21(12):3661-3668 2020

[26] Sharma D, Rawat I and Goel HC. Anticancer and anti-inflammatory activities of some dietary cucurbits. Indian J Exp Biol 2015;53(4):216-221

[27] Liu L., Wang R., He W., He F and Huang G. Cloning and soluble expression of mature a-luffin from *Luffa cylindrica* and its antitumor activities *in vitro* Acta Biochim Biophys Sin 2010, 42 :585-592

[28] Xu, Z.X., L.Q. Li, Z.Q. Zhou, F.Z. Qu and L.L. Tong,. Antiviral effect of an extract of *Luffa cylindrica* (L 043) on Japanese B encephalitis virus infection in vivo. Wei Sheng Wu Xue Bao 1985; 25: 66-68.

[29] Ng YM, Yang Y, Sze KH, Zhang X, Zheng YT, Shaw PC. Structural characterization and anti-HIV-1 activities of arginine/glutamaterich polypeptide Luffin P1 from the seeds of sponge gourd (*Luffa cylindrica*). Journal of Structural Biology 2011;174:164-172

[30] Ng YM, Yang Y, Sze KH, Zhang X, Zheng YT, Shaw PC. Structural

characterization and anti-HIV-1 activities of arginine/glutamaterich polypeptide Luffin P1 from the seeds of sponge gourd (*Luffa cylindrica*). Journal of Structural Biology 2011;174, 164-172

[31] Cao TQ, Kim JA., Woo MH., Min BS. SARS-CoV-2 main protease inhibition by compounds isolated from *Luffa cylindrica* using molecular docking, Bioorganic & Medicinal Chemistry Letters 2021, doi: <https://doi.org/10.1016/j.bmcl.2021.127972>

[32] Aboh IM, Fidelis S, Oladosu OP, Adeshina GO, Olayinka BO, Olonitola SO. Antifungal potentials of *Luffa cylindrica* (Roem) ethyl acetate leaf extract. Journal of Phytopharmacology 2020; 9(3): 178-18

[33] Hossain et al. Phytochemical and antimicrobial investigation of *Luffa cylindrica*. Boletín Latinoamericano y del Caribe de Plantas Medicinales y Aromáticas 2010;9 (5): 328

[34] Ahmad B, Khan AA. Antibacterial, antifungal and phytotoxic activities of *Luffa cylindrica* and *Momordica charantia*. Journal of Medicinal Plants Research 2013;7(22):1593-1599

[35] Saliu OA, Akanji MA., Saliu NB, Idowu OA. Pharmacological evidence favouring the ethnomedicinal use of *Luffa cylindrica* (L.) Roem leaf in the relief of pain and fever. Journal of Health and Pharmacology 2019; 7 (4): 38-42.

[36] Hazra M, Sen SK, Bhattacharya S, Halder PK. Evaluation of hypoglycemic and antihyperglycemic effects of *Luffa cylindrica* fruit extract in rats. Journal of Advanced Pharmacy Education Research 2011; 2: 138-146

[37] Akther F, Ashif R, Jajiratul JP, Zahirul K, Prashanta K Paul, Mohammed R. Methanolic extract of *Luffa cylindrica* fruits show antihyperglycemic potential in Swiss

albino mice Adv. in Nat. Appl. Sci., 2014; 8(8): 62-65

[38] Balakrishnan N and Sharma A. Preliminary phytochemical and pharmacological activities of *Luffa cylindrica* fruit. Asian J Pharm Clin Res 2013; 6(2):113-116

[39] El-Fiky FK, Abou-Karam MA, Afify EA. Effect of *Luffa aegyptiaca* (seeds) and *Carissa edulis* (leaves) extracts on blood glucose level of normal and streptozotocin diabetic rats. J Ethnopharmacol. 1996;50(1):43-47.

[40] Arise, R. O. et al. In vitro Angiotensin-1-converting enzyme, α -amylase and α -glucosidase inhibitory and antioxidant activities of *Luffa cylindrica* (L.) M. Roem seed protein hydrolysate. Heliyon 5: e01634 (2019)

[41] Balakrishnan N, Huria T. Protective effect of *Luffa cylindrica* L. fruit in paracetamol induced hepatotoxicity in rats. Int J Pharm Biol Arch 2011;2(6): 1761-1764

[42] Pawashe PM, Shete RV, Kore KJ, Otari KV. Protective role of *Luffa cylindrica* Linn against erythromycin estolate-induced hepatotoxicity. Curr Pharma Res 2011;1:315-319

[43] Pal RK, Manoj J. Hepatoprotective activity of aqueous and alcoholic extracts of *L. cylindrica* in rats. Ann. Bio. Res. 2011; 2:132-141.

[44] Sunil KM, Yadav B, Upadhyay P, Kumar P, Singh C, Dixit J and Tiwari KN LC-ESI MS/MS profiling, antioxidant and anti-epileptic activity of *Luffa cylindrica* (L.) Roem extract. Journal of Pharmacology and Toxicology 2018; 13: 1-18

[45] Kanwal W, Syed AW, Salman A, Mohtashee HM. Anti-emetic and anti-inflammatory activity of fruit peel of *Luffa cylindrica* (L.) Roem. Asian J Nat Appl Sci. 2013;2(2):175-180.

- [46] Khan KW, Ahmed SW, Ahmed S, Hasan MM. Antiemetic and anti-inflammatory activity of leaves and flower extracts of *Luffa cylindrica* (L.) Roem. J Ethnobiol Trad Med Photon 2013; 118:258-263
- [47] Antia BS, Essien EE, Okokon JE., Alalade IG. Wound healing, phytochemical and antimicrobial properties of *Luffa cylindrica* (Linn.) seed extracts. Int.J. Pharm. Sci Drug Res. 2015; 7(4): 340-344
- [48] Onigemo MA., Dairo FAS and Oso YA.A.. Amino acids profile of loofah gourd, *Luffa cylindrica* (M J Roem) seeds subjected to different heat processing methods. Nig. J. Anim. Prod. 2020; 47(2): 280-288.
- [49] Etim EA, Adebayo YA, and Ifeanyi OE. Effect of *Luffa cylindrica* leaf extract on hematological parameters of Swiss albino mice. Medicinal and Aromatic Plants, 2018; 7(5): 1-5.
- [50] Kamatenesi-Mugisha M, Makawiti DW, Oryem-Origa H, Nganga J. The oxytocic properties of *Luffa cylindrica* (L.) M. Roem. and *Bidens pilosa* L., traditionally used medicinal plants from western Uganda. African J Ecology 2007; 45(3): 88-93.
- [51] Ogunyemi, T. C., Ekuma, C. M., Egwu, J. E., & Abbey, D. M. (2020). Proximate and mineral composition of sponge gourd (*Luffa cylindrica*) seed grown in South-Western Nigeria. Journal of Scientific Research and Reports 26(4): 61-67.
- [52] Oyetayo FL, Oyetayo VO and Ajewole V. Phytochemical profile and antibacterial properties of the seeds and leaf of the Luffa plant (L. *cylindrica*). J Pharmacol Toxicol 2007; 2: 586-589
- [53] Howlader, AH Iqbal, Shamim SM, Sirajul I and Quader MA. Phytochemical constituents of some vegetables Dhaka Univ. J . Sci. 2013; 61(2): 147-151
- [54] Aladejimokun AO. Adesina IA, Falusi VO, Edagbo DE. Comparative study of antimicrobial potency and phytochemical analysis of methanolic extracts of the leaf and flower of *Luffa cylindrica*. Journal of Natural Sciences Research 2014; 4(8):7-10
- [55] Reddy BP, Reddy AR, Reddy BS, Mohan SV and Sarma PN. Apoptosis inducing activity of *Luffa acutangula* fruit in leukemia cells [HL-60]. International Journal of Pharmaceutical Research and Development 2010; 2:109-122.
- [56] Khajuria A, Gupta A, Garai S and Wakhloo BP. Immunomodulatory effects of two saponins 1 and 2 isolated from *Luffa cylindrica* in Balb/C mice. Bioorg Med Chem Lett 2007;17(6):1608-1612.
- [57] Ismail, Hussain MM, Dastagir MG, Billah M and Quader A. Phytochemical and antimicrobial investigation of *Luffa cylindrica*. Boletín Latinoamericano y del Caribe de Plantas Medicinales y Aromáticas 2010; 9 (5): 327 – 332.
- [58] Hleh, TB, Belhadj F. et al . Variations in the bioactive compounds composition and biological activities of loofah (*Luffa cylindrica*) fruits in relation to maturation stages. Chemistry & Biodiversity 2019;14: e1700178
- [59] Maamoun AA., El-akkad RH, Farag MA. Mapping metabolome changes in *Luffa aegyptiaca* mill fruits at different maturation stages via MS-based metabolomics and chemometrics, Journal of Advanced Research 2021; 29: 179-189,
- [60] Hu W., Wang X., Wu L., Shen T., Ji L., Zhao X., Si C. L., Jiang Y. and Wang G. Apigenin-7-O-D-glucuronide inhibits LPS-induced inflammation

- through the inactivation of AP-1 and MAPK signaling pathways in RAW264.7 macrophages and protects mice against endotoxin shock, *Food Funct.* 2016;7: 1002-1013.
- [61] Nath LR, Gorantla JN, Joseph SM, Antony J, Thankachan S, Menon DB, Sankar S, Lankalapalli RS, Anto RJ. Kaempferide, the most active among the four flavonoids isolated and characterized from *Chromolaena odorata*, induces apoptosis in cervical cancer cells while being pharmacologically safe. *RSC Adv*, 2015;5:100912-100922
- [62] Jiao, Z., Xu, W., Zheng, J. *et al.* Kaempferide prevents titanium particle induced osteolysis by suppressing JNK activation during osteoclast formation. *Sci Rep* 2017, 7: 16665 <https://doi.org/10.1038/s41598-017-16853-w>
- [63] Arab HH, Salama SA, Omar HA, Arafa E-SA, Maghrabi IA. Diosmin protects against ethanol-induced gastric injury in rats: Novel anti-ulcer actions. *PLoS ONE* 2015;10(3): e0122417. <https://doi.org/10.1371/journal.pone.0122417>.
- [64] Zheng Y, Zhang R, Shi W, Li L, Liu H, Chen Z, Wu L. Metabolism and pharmacological activities of the natural health-benefiting compound diosmin. *Food Funct.* 2020;11, 8472-8492. doi. [org/10.1039/D0FO01598A](https://doi.org/10.1039/D0FO01598A)
- [65] Lee SE, Yang H., Son GW, Park HR, Park CS, Jin YH, Park YS. Eriodictyol protects endothelial cells against oxidative stress-induced cell death through modulating ERK/Nrf2/ARE-dependent heme oxygenase-1 expression. *Int.J Mol.Sci.*, 2015; 16: 14526-14539 .
- [66] Xu, D, Hu MJ, Wang, YQ, Cui YL. Antioxidant activities of quercetin and its complexes for medicinal application. *Molecules* 2019;24: 1123.<https://doi.org/10.3390/molecules24061123>
- [67] Rauf A, Imran M, Khan IA, Rehman M, Gilani SA, Mehmood Z, Mubarak MS. Anticancer potential of quercetin :comprehensive review. *Phyto Ther Res* 2018;32 (11):2109-2130
- [68] Song X, Tan L, Wang M, Ren C, Guo C, Yang B, Ren Y, Cao Z, Li Y, Pei J. Myricetin: a review of the most recent research *Biomed Pharmacother.* 2021;134: 111017. doi.org/10.1016/j.biopha.2020.111017. *Asia* 2019; 45:532-537
- [69] Li, M.; Chen, J.; Yu, X.; Xu, S.; Li, D.; Zheng, Q.; Yin, Y. Myricetin suppresses the propagation of hepatocellular carcinoma via Down-regulating expression of YAP. *Cells* 2019, 8, 358. <https://doi.org/10.3390/cells8040358>
- [70] Ganeshpurkar A, Saluja AK.. The pharmacological potential of rutin. *Saudi Pharmaceutical Journal* 2017; 25(2): 149-164.
- [71] Isemura M Catechin in human health and disease. *Molecules* 2019, 24(3):528
- [72] Imran M, Rauf A, Abu-Izneid T, Nadeem M, Shariati MA, Khan IA, Imran A, Orhan E , Rizwan M, Atif Md, Gondal TA, Mubarak MS. Luteolin, a flavonoid, as an anticancer agent: a review, *Biomedicine & Pharmacotherapy* 2019;112,108612, <https://doi.org/10.1016/j.biopha.2019.108612>
- [73] Qiu, J.; Zhang, T.; Zhu, X.; Yang, C.; Wang, Y.; Zhou, N.; Ju, B.; Zhou, T.; Deng, G.; Qiu, C. Hyperoside induces breast Cancer cells apoptosis via ROS-mediated NF-κB signaling pathway. *Int. J. Mol. Sci.* 2020, 21, 131. <https://doi.org/10.3390/ijms21010131>
- [74] Wang X, Liu Y, Xiao L, Li L, Zhao X, Yang L, Chen N, Gao L, Zhang J: Hyperoside protects against pressure overload-induced cardiac remodeling

via the AKT signaling pathway. *Cell Physiol Biochem* 2018;51:827-841. doi: 10.1159/000495368

[75] Boukes GJ, van de Venter M. The apoptotic and autophagic properties of two natural occurring prodrugs, hyperoside and hypoxoside, against pancreatic cancer cell lines. *Biomed. and Pharmacotherapy* 2016;83:617-626. <http://dx.doi.org/10.1016/j.biopha.2016.07.029>.

[76] Govindarasu M, Thangaraj K, Murugesan V, Vaiyapuri M. Kaempferitrin cause cell cycle arrest at G2/M phase and reactive oxygen species mediated apoptosis in human colon cancer HT-29 cells. *IJRTE* 2019; 8(3S2) : 2277-3878..

[77] Zhang Y, Guo Y, Wang M, Dong H, Zhang J and Zhang L: Quercetrin from *Toona sinensis* leaves induces cell cycle arrest and apoptosis via enhancement of oxidative stress in human colorectal cancer SW620 cells. *Oncol Rep* 38: 3319-3326, 2017

[78] Grochowski DM, Locatelli M, Granica S, Cacciagrano F, Tomczyk M. A review on the dietary flavonoid tiliroside. *Comprehensive Reviews in Food Science and Food Safety* 2018; 17 (5): 1395-1421

[79] Velagapudi, R., El-Bakoush, A. & Olajide, O.A. Activation of Nrf2 pathway contributes to neuroprotection by the dietary flavonoid Tiliroside. *Mol Neurobiol* 55, 8103-8123 (2018). <https://doi.org/10.1007/s12035-018-0975-2>

[80] Kwon EB, Kang MJ, Ryu HW, Lee S, Lee JW, Lee MK, Lee HS, Lee SU, Oh SR, Kim MO. Acacetin enhances glucose uptake through insulin-independent GLUT4 translocation in L6 myotubes, *Phytomedicine*, 68, 2020, 153178 :doi.org/10.1016/j.phymed.2020.153178

[81] Xiao WZ, Zhou WH, Ma Q, Cui WG, Mei QY, Zhao X.

Serotonergically dependent antidepressant-like activity on behavior and stress axis responsivity of acacetin. *Pharmacological Research*,2019; 146:104310. doi.org/10.1016/j.phrs.2019.104310

[82] Kim SM, Park YJ, Shin MS, Kim HR, Kim MJ, Lee SH, Yun S, Kwon SH. Acacetin inhibits neuronal cell death induced by 6-hydroxydopamine in cellular Parkinson's disease model. *Bioorganic & Medicinal Chemistry Letters* 2017;27 (23): 5207-5212

[83] Zhao C, Zhang Y, Liu H, Li, P, Zhang H, Cheng G,. Fortunellin protects against high fructose-induced diabetic heart injury in mice by suppressing inflammation and oxidative stress via AMPK/Nrf-2 pathway regulation. *Biochemical and Biophysical Research Communications* 2017; 490: (2):552-559

[84] Wang J, Fu B, Lu F, Hu X, Tang J, Huang L. Inhibitory activity of linarin on osteoclastogenesis through receptor activator of nuclear factor κ B ligand-induced NF- κ B pathway, *Biochem Biophys. Res. Comm.* 2018; 495(3): 2133-2138

[85] Yu Q, Li X, Cao X. Linarin could protect myocardial tissue from the injury of ischemia reperfusion through activating Nrf-2, *Biomed. Pharmacother.* 2018 90 (2017a):1e7.

[86] Ni M, Hu X, Gong D, Zhang G. Inhibitory mechanism of vitexin on α -glucosidase and its synergy with acarbose, *Food Hydrocolloids* 2020;105:105824

[87] Peng, Y., Gan, R., Li, H., Yang, M., McClements, D. J., Gao, R., & Sun, Q. Absorption, metabolism, and bioactivity of vitexin: Recent advances in understanding the efficacy of an important nutraceutical. *Critical Reviews in Food Science and Nutrition*, 2021;61:1-16

- [88] Wang Y, Liu T, Li MF, Yang YS, Li R, Tan J, Tang SH, Jiang ZT. Composition, cytotoxicity and antioxidant activities of polyphenols in the leaves of star anise (*Illicium verum* Hook. f.). *ScienceAsia* 2019;45:532-537
- [89] Simeonova R, Vitcheva V, Kondeva-Burdina M, Krasteva I, Manov V, Mitcheva M., Hepatoprotective and antioxidant effects of saponarin, isolated from *Gypsophila trichotoma* Wend. on paracetamol-induced liver damage in rats. *BioMed research international* 2013(5):757126
- [90] Pei K, Ou J, Huang J, Ou S. p-Coumaric acid and its conjugates: Dietary sources, pharmacokinetic properties and biological activities. *J Sci Food Agric*. 2016;96(9):2952-2962.
- [91] Shen Y, Song X, Li L, Jian S, Jaiswal Y, Huang J, Liu C, Yang W, Williams L, Zhang H, Guan Y. Protective effects of p-coumaric acid against oxidant and hyperlipidemia-an in vitro and in vivo evaluation. *Biomedicine & Pharmacotherapy* 2019;111:579-587
- [92] Kwak SH, Kim YH. Anti-adipogenic effect of 1-O-feruloyl- β -D-glucose on 3T3L1preadipocytes. *Korean J. Food Preserv*. 2018; 25(6): 689-695
- [93] Kahkeshani N, Farzaei F, Fotouhi M, Alavi SSH, Bahramsoltani R, Naseri R, Momtaz S, Abbasabadi Z, Rahimi R, Farzaei MH, Bishayee A. Pharmacological effects of gallic acid in health and diseases: a mechanistic review. *Iran J Basic Med Sci* 2019; 22:225-237.
- [94] Abedi F, Razavi BM. Hosseinzadeh H. A review on gentisic acid as a plant derived phenolic acid and metabolite of aspirin: Comprehensive pharmacology, toxicology, and some pharmaceutical aspects. *Phyto. Ther. Res*. 2020; 34 (4): 729-741
- [95] Lu H, Tian Z , Cui Y, Liu Z, Ma X. Chlorogenic acid: A comprehensive review of the dietary sources, processing effects, bioavailability, beneficial properties, mechanisms of action, and future directions. *Comp. Rev. Food Sci Food Saf* 2020;9(6): 3130-3158
- [96] Baniahmad B, Safaeian L, Vaseghi G, Rabbani M, Mohammadi B. Cardioprotective effect of vanillic acid against doxorubicin-induced cardiotoxicity in rat. *Res Pharm Sci*. 2020; 15(1):87-96.
- [97] Han X, Guo J, You Y, Yin M, Liang J, Ren C, Zhan J and Huang W. Vanillic acid activates thermogenesis in brown and white adipose tissue. **Food Funct**. 2018; 9: 4366-4375
- [98] Amin, F., Shah, S. & Kim, M. Vanillic acid attenuates A β ₁₋₄₂-induced oxidative stress and cognitive impairment in mice. *Sci Rep* 2017; 7: 40753 <https://doi.org/10.1038/srep40753>
- [99] Fadeyi OO, Obafemi CA, Adewunmi CO, Iwalewa EO Antipyretic, analgesic, anti-inflammatory and cytotoxic effects of four derivatives of salicylic acid and anthranilic acid in mice and rats. *African Journal of Biotechnology* 2004; 3 (8):426-431
- [100] Zeng W, Jin L, Zhang F, Zhang C, Liang W. Naringenin as a potential immunomodulator in therapeutics, *Pharmacological Research* 2018; 13: 122-126
- [101] Moon JY, Ediriweera MK, Ryu JY, Kim HY and Cho SK: Catechol enhances chemo- and radio-sensitivity by targeting AMPK/hippo signaling in pancreatic cancer cells. *Oncol Rep* 2021; 45: 1133-1141.
- [102] Espíndola KMM, Ferreira RG, Narvaez LEM, Silva Rosario ACR, da Silva AHM, Silva AGB, Vieira APO and

Monteiro MC. Chemical and pharmacological aspects of caffeic acid and its activity in hepatocarcinoma. *Front.Oncol.* 2019; 9:541. doi: 10.3389/fonc.2019.00541

[103] Zduńska K, Dana A, Kolodziejczak A, Rotsztein H: Antioxidant properties of ferulic acid and its possible application. *Skin Pharmacol Physiol* 2018;31:332-336.

[104] Gunia-Krzyżak A., Słoczyńska K, Popiół J, Koczurkiewicz P, Marona H, Pękala E. Cinnamic acid derivatives in cosmetics: Current use and future prospects. *Int. J. Cos.* 2018 Sci 40 (4): 356-366.

[105] Adisakwattana S. Cinnamic acid and its derivatives: mechanisms for prevention and management of diabetes and its complications. *Nutrients.* 2017; 9(2):163. <https://doi.org/10.3390/nu9020163>

[106] Ceci C, Lacal PM, Tentori L, De Martino MG, Miano R, Graziani G. Experimental evidence of the antitumor, antimetastatic and antiangiogenic activity of ellagic acid. *Nutrients.* 2018; 10(11):1756. <https://doi.org/10.3390/nu10111756>

[107] Ríos JL, Giner RM, Marín M, Recio MC. A pharmacological update of ellagic acid. *Planta Med* 2018; 84(15): 1068-1093

[108] Ayeleso TB, Matumba MG, Mukwevho E. Oleanolic acid and its derivatives: Biological activities and therapeutic potential in chronic diseases. *Molecules.* 2017; 22(11):1915. <https://doi.org/10.3390/molecules22111915>

[109] Khwaza V, Oyedele OO, Aderibigbe BA. Antiviral activities of oleanolic acid and its analogues. *Molecules.* 2018; 23(9):2300. <https://doi.org/10.3390/molecules23092300>

[110] Yu Q, Li X, Cao X. Linarin could protect myocardial tissue from the injury of ischemia reperfusion through

activating Nrf-2, *Biomed. Pharmacother.* 2017; 90: 1e7.

[111] Garai, S.G.R., Bandopadhyay, P.P., Mondal, N.C., Chattopadhyay, A., Anti-microbial and anti-cancer properties of echinocystic acid extracted from *Luffa cylindrica*. *J. Food Process.* 2018; 9:1-4

[112] Öztürk S. E., Karayıldırım T., Çapcı-Karagöz A., Alanku O., Özmen A., and Poyrazoğlu-Çoban E. Synthesis, antimicrobial and cytotoxic activities, and structure-activity relationships of gypsogenin derivatives against human cancer cells. *European Journal of Medicinal Chemistry* 2014; 1:1-47

[113] Ren, Y.; Shen, L.; Zhang, D.; Dai, S. Two new Sesquiterpenoids from *Solanum lyratum* with cytotoxic activities. *Chem. Pharm. Bull.* 2009; 57 (4):408-410.

[114] Fang, Z.; Jeong, S. Y.; Jung, H. A.; Choi, J. S.; Min, B. S.; Woo, M. H. Anticholinesterase and antioxidant constituents from *Gloiopeltis furcata*. *Chem. Pharm. Bull.* 2010;58 (9): 1236-1239.

[115] Kulkarni, SS, Bhalke D, Pande VV, Kendre PN. Herbal plants in photo protection and sun screening action: An overview. *Indo American Journal of Pharmaceutical Research* 2014; 4(2):1104-1113

[116] Peng L, Sun S, Xie LH, Wicks SM, Xie JT. Ginsenoside Re: Pharmacological effects on cardiovascular system. *Cardiovasc.Ther.* 2012; 30: e183-e188

[117] Gao Y, Yang MF, Su YP, Jiang HM, You XJ, Yang YJ, Zhang HL. Ginsenoside Re reduces insulin resistance through activation of PPAR- α pathway and inhibition of TNF- α production. *J. Ethnopharmacol.* 2013; 147: 509-516.

[118] Liu Z, Qi Y, Cheng Z, Zhu X, Fan C, Yu SY. The effects of ginsenoside Rg1 on

chronic stress induced depression-like behaviors, BDNF expression and the phosphorylation of PKA and CREB in rats. *Neuroscience* 2016; 322: 358-369

[119] Yoshikawa K, Arihara S, Wang JD, Narui T and Okuyama T. Structures of two new fibrinolytic saponins from the seed of *Luffa cylindrica* Roem. *Chem Pharm Bull (Tokyo)* 1991;39(5): 1185-1188.

[120] Zhou T, Zu G, Zhang X, Wang X, Li S, Gong X, Liang Z, Zhao J. Neuroprotective effects of ginsenoside Rg1 through the Wnt/b-catenin signaling pathway in both in vivo and in vitro models of Parkinson's disease. *Neuropharmacology* 2016;101: 480-489

[121] Li Y, Wang F, Luo Y. Ginsenoside Rg1 protects against sepsis-associated encephalopathy through beclin 1-independent autophagy in mice. *J. Surg. Res.* 2017; 207: 181-189

[122] Xin Y, Wei J, Chunhua M, Danhong Y, Jianguo Z, Zongqi C, Jian-an B. Protective effects of ginsenoside Rg1 against carbon tetrachloride-induced liver injury in mice through suppression of inflammation. *Phytomedicine* 2016;23: 583-588

[123] El-Gengaihi S, Abd El-Hamid SR and Kamel AM. Anti-inflammatory effect of some cucurbitaceous plants. *Herba Polonica* 2009, 55(4): 119-126.

[124] Attard E, Martinoli MG. Cucurbitacin E, An experimental Lead triterpenoid with anticancer, immunomodulatory and novel effects against degenerative diseases. A Mini-Review. *Curr Top Med Chem.* 2015;15(17):1708-1713.

[125] Ng TB, Wong RNS, Yeung HW. Two proteins with ribosome-inactivating, cytotoxic and abortifacient activities from seeds of *Luffa cylindrica*

Roem (Cucurbitaceae). *Biochemistry International* 1992;27:197-207

[126] Tanaka S, Uno C, Akimoto M, et al. Anti-allergic effect of bryonolic acid from *Luffa cylindrica* cell suspension cultures. *Planta Med.* 1991, 57:527-530

[127] Du Q and Wang K. Preparative separation of phenolic constituents in the fruits of *Luffa cylindrica* (L.) Roem using slow rotary countercurrent chromatography. *Journal of Liquid Chromatography and Related Technologies* 2007, 30 (13):1915-1922.

[128] Gao JJ, Igalashi K, Nukina M () Radical Scavenging Activity of Phenylpropanoid Glycosides in *Caryopteris incana*, Bioscience, Biotechnology, and Biochemistry 1999, 63:6: 983-988

[129] Kylli P, Nousiainen P, Biely P, Sipila J, Tenkanen M, Heinonen M (2008) Antioxidant potential of hydroxycinnamic acid glycoside esters. *J Agric Food Chem* 56:4797-4805

[130] Tóth B, Hohmann J, Vasas A. Phenanthrenes: a promising group of plant secondary metabolites. *J. Nat. Prod.* 2018, 81, 3, 661-678

[131] Kovács A., Vasas A, Hohmann J. Natural phenanthrenes and their biological activity. *Phytochemistry* 2008;69:1084-1110

[132] Wang L, Lou G, Ma Z, Liu X. Chemical constituents with antioxidant activities from litchi (*Litchi chinensis* Sonn.) seeds. *Food Chemistry* 2011; 126(3):1081-1087

[133] López-Biedma A, Sánchez-Quesada C, Beltrán G, Delgado-Rodríguez M, Gaforio JJ. Phytoestrogen (+)-pinoresinol exerts antitumor activity in breast cancer cells with different oestrogen receptor statuses. *BMC Complement Altern Med.* 2016;16(1):350. doi: 10.1186/s12906-016-1233-7.

- [134] Travlos, I.; Rapti, E.; Gazoulis, I.; Kanatas, P.; Tataridas, A.; Kakabouki, I.; Papastilianou, P. The herbicidal potential of different Pelargonic acid products and essential oils against several important weed species. *Agronomy* 2020; 10:1687. <https://doi.org/10.3390/agronomy10111687>
- [135] Kaidi Cui, Song Yang, Nan Zou, Leiming He, Tao Zhang, Feng Liu, Wei Mu. Residual behavior of the potential grain fumigant 1-octen-3-ol in wheat during fumigation and ventilation processes. <https://doi.org/10.1002/ps.6329>
- [136] Wang CY, Chen YW, Hou CY. Antioxidant and antibacterial activity of seven predominant terpenoids, *International Journal of Food Properties* 2019; 22: 230-238
- [137] Khaleel C, Tabanca N, Buchbauer G. α -Terpineol, a natural monoterpene: A review of its biological properties. *Chemistry Open* 2018b; 16 (1):): 122-135
- [138] Zavala-Sanchez MA, Pérez-Gutiérrez S, Prez-González C, Sánchez-Saldivar D, Arias-García L. Antidiarrhoeal activity of nonanal, an aldehyde isolated from *Artemisia ludoviciana*. *Pharmaceutical Biology* 2012; 40 (4): 263-268
- [139] Mahboubi M., Feizabadi MM Antimicrobial activity of *Ducrosia anethifolia*. Essential oil and main component, decanal against methicillin-resistant and methicillin-susceptible *Staphylococcus aureus*, *Journal of Essential Oil Bearing Plants* 2009, 12:5, 574-579 Brian *et al*, 2018,
- [140] Barboza JN, Filho CSMB, Silva RO, Medeiros JVR. de Sousa, DP. An Overview on the anti-inflammatory potential and antioxidant profile of eugenol. *Oxidative Medicine and Cellular Longevity* 2018; Article ID 3957262, 9 pages, 2018. <https://doi.org/10.1155/2018/3957262>
- [141] Mukhtar YM, Adu-Frimpong M, Xu X, Yu J. Biochemical significance of limonene and its metabolites: future prospects for designing and developing highly potent anticancer drugs. *Biosci Rep.* 2018;38(6):BSR20181253. Published 2018 Nov 13. doi:10.1042/BSR20181253
- [142] Prost, I. Evaluation of the antimicrobial activities of plant oxylipins supports their involvement in defense against pathogens. *Plant Physiol.* 2005; 139: 1902-1913
- [143] Nagai, T.; Kiyohara, H.; Munakata, K.; Shirahata, T.; Sunazuka, T.; Harigaya, Y.; Yamada, H. Pinellic acid from the tuber of *Pinellia ternata* Breitenbach as an effective oral adjuvant for nasal influenza vaccine. *Int. Immunopharmacol.* 2002; 2: 1183-1193
- [144] Yoshida J, Uesugi S, Kawamura T, Kimura K, Hu D, Xia S, Toyooka N, Ohnishi M Kawashima H. (4Z,15Z)-octadecadienoic acid inhibits glycogen synthase kinase-3 β and glucose production in H4IIE cells *Lipids* 2017;52:. 295-230
- [145] Thayyil AH, Surulivel MKM, Ahmed MF, et al. Hypolipidemic activity of *Luffa aegyptiaca* fruits in cholesterol fed hypercholesterolemic rabbits. *Int J Pharm Appl.* 2011;2(1): 81-88.
- [146] Oyeyemi IT, Yekeen OM, Odusina PO, Ologun TM, Ogbaide OM, Olaleye OI, Bakare AA. Genotoxicity and antigenotoxicity study of aqueous and hydro-methanol extracts of *Spondias mombin* L., *Nymphaea lotus* L. and *Luffa cylindrical* L. using animal bioassays. *Interdiscip Toxicol.* 2015; 8(4):184-92.