

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

185,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



# The Role of Cryptosporidiosis in Sheep Welfare

*María Uxúa Alonso Fresán and Alberto Barbabosa Pliego*

## Abstract

Welfare in animal production has been defined as the optimal mental and physiological state of the animals. It has been recently redefined according to animals' freedoms. As systems, individual sheep and herds are dynamic with constant interaction with each other and the environment. In this interaction, diseases play a fundamental role in welfare. Parasitism is common in sheep, and several management practices have been established to maintain the herds healthy. *Cryptosporidium* represents a special case, because it is a highly resistant environmental parasite, that can easily infect lambs, producing weakening diarrheas and even death. In this chapter, the role of cryptosporidiosis in sheep welfare and economic loss will be analyzed, as means of providing information on how to minimize and deal with the infection.

**Keywords:** *Cryptosporidium*, cryptosporidiosis, sheep, lamb, welfare, diarrhea, economic loss, prevention, control, treatment

## 1. Introduction

Animal welfare and health status in herds go hand in hand. When pathogens are present, disease may develop and will be reflected on decreased productivity. Parasites are pathogens which may be found in the environment as well as in the hosts, some of them as opportunists that can cause disease. Such is the case of *Cryptosporidium spp.*, a ubiquitous parasite with worldwide distribution, which causes diarrhea in newborn lambs that can lead to death or self-limiting diarrheas in immunocompetent hosts. In this chapter, welfare and cryptosporidiosis is discussed in an attempt to provide information on how to prevent and control the disease.

## 2. Welfare generalities

According to the World Organization for Animal Health (OIE), animal health and welfare is defined as “the physical and mental state of an animal in relation to the conditions in which it lives and dies”, and covers “the five freedoms”: “1) freedom from hunger, malnutrition and thirst; 2) freedom from fear and anxiety; 3) freedom from heat stress or physical discomfort; 4) freedom from pain; and 5) freedom to express normal patterns of behaviour”, [1]. These were developed by the UK Animal Welfare Council (FAWC) in 1979, after researching on farmed animals in intensive systems, [2].

Derived from these freedoms and based on [3], Codes of Best Practices for Welfare Establishments have been developed, such as the one from the Welsh Government [4], in which a series of issues are covered, in order to keep the animals healthy and productive. As an example, this one indicates not only how the animals should be managed, but also how staff and volunteers should be managed too. Regarding animal management, a conscientious record of animal admissions, behavior and assessment, housing and environment, cleaning and hygiene among others are defined. Animal health and disease are also included, as well as rehabilitation, rehoming or release, and transportation, all based on the corresponding legislation.

There is a close relationship between animal health and welfare. If they are optimal, they will promote high productivity and the reduction in the use of antimicrobials, as well as reduced risk for foodborne diseases to humans [5].


### 3. Welfare and disease

In this sense, the AWIN welfare assessment protocol for sheep [6], defines welfare indicators according to principles and criteria, in which good health is measured by the absence of disease, and according to this protocol and the objective of this chapter, it is indicated by fecal soiling. The European Food Safety Authority (EFSA) has identified gastroenteric disorders among the main welfare consequences in lambs [1]. Regarding gastroenteric disorders, diarrhea is defined as a “complex, multifactorial disease involving the animal, the environment, nutrition, and infectious agents” [7], in which *Cryptosporidium spp.* is among the most common. It is an apicomplexan protozoa that causes profuse diarrhea in neonatal lambs (4–9 days old), with low mortality, increasing concurrent infection and deficiencies in nutrition and husbandry. In well-fed animals, it often persists for 5–7 days, lessens and lambs recover. Diarrhea leads to dehydration, inappetence, abdominal pain and lethargy. Diarrhea is liquid and yellowish, varying from mild and self-limited to severe (when mixed with other pathogens) but relapse is quite common [8]. Cryptosporidiosis produces severe villous atrophy, caused by loss of enterocytes, with crypt hyperplasia by the replacement of epithelial cell loss. This occurs due to the cytotoxic effect of the parasite and apoptosis it causes, resulting in a malabsorptive diarrhea, with prostaglandin-mediated anion secretion [9].

Other pathogens might be present in diarrheas caused by *Cryptosporidium spp.*, such as rotaviruses, *E. coli*, *Salmonella spp.* or coronaviruses. In this case, prognosis degenerates and clinical signs and treatment become complicated [10].

### 4. Species and life cycle

There are more than 38 species of *Cryptosporidium*, but only three main species are reported in small ruminants: *C. parvum*, *C. xiaoi* and *C. ubiquitum* [11]. *Cryptosporidium spp.* oocysts size ranges from 4 to 4.5  $\mu\text{m}$ , with spherical to ovoid shape, and four sporozoites per oocyst. They can be identified through Ziehl-Neelsen stain, where oocysts are shown as bright pink round bodies. The transmission of the infection happens by direct ingestion of oocysts present in food and water. Once in the host, oocysts excyst in the gut and sporozoites are released, entering the intestinal epithelial cells through the brush border [12]. After excystation, sporozoites are found in the extracytoplasmatic



**Figure 1.**  
*Cryptosporidium spp.* life cycle [14].

parasitophorus vacuole, where the parasite reproduces either asexually or by schizogony, leading to 8 merozoites within a type I meront. Merozoites are capable of invading neighboring epithelial cells and other sites of the intestine, where they either reproduce asexually forming thin-wall oocysts that cause autoinfection or a sexually forming type II meronts differentiating into microgametocytes and macrogametocytes, which will unite to form the zygote. This zygote will form 4 sporozoites in thick or thin-walled oocysts, by a process called sporogony. The ones with the thick wall are released to the environment through feces [13] (**Figure 1**). Up to  $1 \times 10^6$ - $2 \times 10^6$  oocysts/g of feces can be excreted by the host (which are immediately infectious) becoming a source of infection to susceptible lambs [15]. Oocysts can survive in water and in the environment several months, as soon as cool temperatures and suitable moisture exists, and can be easily transported through air and water [16].

## 5. Prevalence

*Cryptosporidium* was incidentally discovered by Tyzzer in 1907, but until 1978 oocyst identification in feces samples was confirmed through microscopy, making it the diagnostic method. Several stains have been proven, but the most common is modified Ziehl-Neelsen. Next came immunological tests which are more sensitive and specific for the detection of antigens, giving rise to immunofluorescent tests. These are specially used in water samples. Finally, genetic tools such as PCR were developed to identify the species [17, 18]. Nevertheless, microscopy using stains still remains as the simplest, cheapest and the most common way for detection.

Ever since it was first detected and confirmed as pathogen, there have been many studies regarding prevalence in sheep from different geographical regions, in which its presence may be attributed to either infectivity or the contamination of the environment [19]. In an attempt trying to explain the wide variation in prevalence, it is shown in tables according to Koeppen's climate classification (**Tables 1-5**).

Author	Year	Country	Detection test	Population	Prevalence (%)	Climate
[20]	1990	Guaiba District, Brasil	Z-N	10 lambs	20	Aw
[21]	1993	Trinidad and Tobago	Z-N	51 diarrheic lambs 31 healthy lambs	24.5 10	Am
[22]	1995	Western Malaysia	Z-N	25 lambs	36.0	Af
[23]	1996	Morogoro Region, Tanzania	Z-N	121 lambs	97.5	Aw
[24]	2001	Ibadan, Nigeria	Z-N		43.3	Aw
[25]	2003	Aba, Nigeria	Z-N	29 sheep	2.1	Am
[26]	2014	Papua New Guinea	PCR	276 adult sheep	2.2	Aw
[27]	2015	India	IF	20 (5 sheep/pool)	45	Aw
[28]	2017	Sinaloa, Mexico	Z-N	1144 lambs (1–90 days old)	41.58	As
[29]	2018	Northern Veracruz, Mexico	Kinyoun	210 healthy lambs	19.5	Aw

Af: Tropical wet; Am: Tropical monsoon; As: Tropical dry savanna; Aw: Tropical savanna.

**Table 1.**  
*Cryptosporidium spp. prevalences reported in climate a (tropical).*

Author	Year	Country	Detection test	Population	Prevalence (%)	Climate
[30]	1994	Ismalia, Egypt	—	100 lambs	24	BWh
[31]	1998	Gharbia province, Egypt	—	52 s lamb	7.69	BWh
[32]	2002	Tehran, Iran		87 sheep	Lambs (less than a month old): 14.28 Ewes (3 years or more): 17.69	BSk
[33]	2002	Basrah, Iraq	Z-N	45 sheep	13.3	BWh
[34]	2002	Dakahlia Governorate, Egypt	Z-N	200 lambs	13.3	BWh
[35]	2002	Zaragoza, Spain	Z-N	344 lambs	59	BSk
[36]	2004	Sharkia and Dakahlyia Provinces, Egypt		470 lambs and kids (310 diarrheic y 160 healthy)	16.8 in diarrheic animals	BWh
[37]	2007	Aguascalientes, Mexico	Kinyoun	40 sheep	100	Bsh
[38]	2014	Kafr El Sheikh, Egypt	Z-N	45 lambs 75 sheep	4.4 1.3	BWh
[39]	2015	Lahore, Pakistan	Z-N	150 diarrheic and healthy lambs: 0–3 months 4–6 months 7–12 months 1 year	40 22.5 12.5 6.67	Bsh
[40]	2019	Yazd Province, Iran	PCR	192 slaughtered adult sheep	5.7	Bwk

BWh: Tropical and subtropical desert; Bsh: Mid-latitude steppe and desert; BSk: Tropical and subtropical steppe.

**Table 2.**  
*Cryptosporidium spp. prevalences reported in climate B (dry).*

Author	Year	Country	Detection test	Population	Prevalence (%)	Climate
[41]	1990	V, VI and Metropolitan regions, Chile	Z-N	57 diarrheic lambs 114 healthy lambs	3.5 (diarrheic) 7.9 (healthy)	Csa/Csb
[42]	1990	Elazig, Turkey	—	267 lambs	12	Csa
[43]	1991	Lorestan, Iran	Z-N	215 healthy lambs	17.2	Csa
[44]	1991	Valdivia, Chile	Z-N	196 dead lambs (0–28 days of age)	7.7	Cfb
[45]	1991	Galicia, Spain	Safranin methylene blue and Köster stain	69 lambs (6 days–6 months of age)	1.45	Csb
[46]	1994	Larissa, Greece	Z-N	65 diarrheic lambs	4.61	Csa
[47]	1996	Castille and Leon, Spain	Z-N	183 diarrheic lambs	45	Csb
[48]	1999	China	—	3250 lambs	23.08	Cfa Cwa
[49]	2000	Izmir, Turkey	—	150 diarrheic lambs 50 healthy lambs	23.3 2	Csa
[50]	2000	Poznan, Poland	Z-N	205 lambs	10.1	Cfb
[51]	2001	Poland	Coproantigen	17 lambs	11.76	Cfb
[52]	2004	São Paolo, Brasil	Z-N	184 lambs in rainy season 179 lambs in dry season	55.4 17.3	Cfa
[53]	2004	Aydin Province, Turkey	Heine	67 diarrheic lambs 77 healthy lambs	79.1 18.2	Csa
[54]	2004	Guangdong, China	Fenol, auramine and Z-N	Sheep	21.7	Cwa
[55]	2005	São Paolo, Brasil	Auramine O and Z-N	20 sheep 20 lambs	26.7 31.9	Cfa



Author	Year	Country	Detection test	Population	Prevalence (%)	Climate
[56]	2005	Central Mexico	Z-N	1200 (559 lambs and 641 ewes)	General: 34.3 Lambs: 32.5 Ewes: 35.9	Cwb
[57]	2005	Konya, Turkey	Z-N	471 lambs (1–60 days of age)	2.97	Csa
[58]	2006	Hirta, St. Kilda, Scotland	Z-N	Lambs and sheep	2001–2028.6 2002–9.0 2003–2011.9	Csa
[59]	2006	Serbia	Z-N and Kinyoun	126 lambs (1–90 days of age)	42.1	Cfa
[60]	2007	Maryland, USA	IF	32 ewes 31 lambs	9.4 32.5	Cfa
[61]	2007	Galicia, Spain	IF	446 adult sheep	5.3	Csb
[62]	2007	Tunisia	PCR	89 sheep (healthy lambs and adults)	11.2	Csa
[63]	2007	Eastern England	FAT	80 asymptomatic lambs	9.7	Cfb
[64]	2008	Central Macedonia, Greece	Z-N	207 diarrheic lambs 79 healthy lambs 237 healthy ewes	55.07 15.18 10.97	Csa
[65]	2008	East Flanders, Belgium	IF and PCR	137 lambs (1–10 weeks old)	13.1	Cfb
[66]	2009	Australia	PCR	477 pre-weaned sheep	24.5	Csb
[67]	2014	Crete, Greece	IF	425 lambs	5.1	Csa
[68]	2015	Jammu District, India	Z-N	55 diarrheic lambs	45	Cfa
[69]	2016	Veracruz, Mexico	Z-N	80 sheep	70	Cfb

Author	Year	Country	Detection test	Population	Prevalence (%)	Climate
[70]	2016	Lorestan,Iran	Z-N	345 sheep (lambs and adults, diarrheic and healthy)	5.8	Csa
[71]	2017	Poland	PCR	234 asymptomatic lambs	19.2	Cfb
[72]	2018	Algeria	PCR	62 lambs	14.5	Csa
[73]	2020	French Basque Country, France	Heine	Asymptomatic sheep: 79 lambs 72 ewes	1.3–77.8 1.4–50	Cfb
[74]	2020	Sardinia Provinces, Italy	Z-N	915 lambs (diarrheic and healthy)	10.1	Csa
[75]	2021	Kenya	Z-N	388 sheep	19.6	Csb

*Cfa: Humid subtropical; Cfb: Marine West Coast; Csa: Warm summer Mediterranean; Csb: Hot summer Mediterranean; Cwa: Monsoon-influenced humid subtropical.*

**Table 3.**  
*Cryptosporidium spp. prevalences reported in climate C (temperate).*



Author	Year	Country	Detection test	Population	Prevalence (%)	Climate
[76]	1993	Ohio, U.S.A.	IF	9 newborn lambs (5–10 days of age) 23 lambs (2–3 weeks of age) 23 healthy ewes	100 78.3 17.4	Dfa/Dfb
[77]	1995	Hungary	Z-N	53 lambs	22.64	Dfb
[78]	2009	Kars, Turkey	Z-N	400 diarrheic lambs (up to one month old)	38.8	Dfb
[79]	2013	Iran	Z-N	231 diarrheic and healthy lambs	2.5	Dsa
[80]	2016	Qinghai, China	PCR	350 sheep	12.3	Dwc
[81]	2019	Ladakh, India	Z-N	37 sheep (0–5 years old, diarrheic and asymptomatic animals)	54.0	Dwc
[16]	2020	Azerbaijan	Z-N	1,823 sheep	34.17	Dfb

*Dfa/Dfb Humid continental; Dfb Warm-summer humid continental; Dsa Hot summer continental; Dwc Monsoon-influenced subartic.*

**Table 4.**  
*Cryptosporidium spp. prevalences reported in climate D (continental).*

Author	Year	Country	Detection test	Population	Prevalence (%)	Climate
[82]	1997	Different regions in Canada	IF	89 sheep	24	Dfb Dfc Dsb Cfb
[83]	2013	Uttar Pradesh, Uttarakhand, Bihar, Karnataka and Kerala States, India	Z-N	55 lambs	1.8	Csa Cwa Aw Am
[84]	2014	Iran	Z-N	1749 asymptomatic sheep	11.3	BWh Bwk Csa BSk
[85]	2014	Australia	PCR	1182 lambs (12–29 weeks old)	16.9	Cfb Bsh Csa Csb Cfa BWh desert
[86]	2021	Iran	—	3901 sheep Metanalysis	9.9	BWh Bwk Csa BSk

*Am: Tropical monsoon; Aw: Tropical wet savanna; Bsh: Mid-latitude steppe and desert; BSk: Cold semiarid; BWh: Tropical and subtropical desert; Bwk: Tropical and subtropical desert; Dfb: Warm-summer humid continental; Cfa: Humid subtropical; Cfb: Marine West Coast; Csa: Warm summer Mediterranean; Csb: Hot summer Mediterranean; Cwa: Monsoon-influenced humid subtropical; Dfc: Subartic; Dsb: Mediterranean-influenced warm-summer humid continental.*

**Table 5.**  
*Cryptosporidium spp. prevalences reported in studies from countries in which several regions were sampled.*

Results show that it can be found in all different climates. One important factor is the high resistance of the oocysts when shed to the environment. Whenever there are conditions in which the parasite can survive (protection from desiccation or avoidance of direct sunlight) it will be present, even in zones with low humidity [87]. As cryptosporidiosis is multifactorial, it is also very important to take into account the immune status of the host. Immunocompetency varies with age, therefore making lambs the most susceptible. When sheep are immunocompetent, they develop the disease with weakening diarrhea and excrete high loads of the parasite, but it is self-limiting. They remain as healthy carriers, still excreting infective oocysts to the environment. Therefore, it is also important to analyze the prevalence results according to the health status of the host. In some countries, such as Iran, where studies with different climates have been undertaken (Table 5), prevalence has maintained with almost no shifts through time (2014: 11.3%; 2021: 9.9%). Nevertheless, in this same country, in studies where samples were taken in specific zones (dry climate), results show ample variation (2002: 14.28 and 17.69%; 2019: 5.7%), in samples taken from sheep with different ages. In other cases such as Mexico, prevalence was reported in 2005 in a temperate region (34.3%), 2007 in a dry region (100%) and in 2017 (41.58%) and 2018 (19.5%) in a tropical region, where variation in prevalence is wide. India is another example of wide variation in results. In 2013 a prevalence of 1.8% was reported in several climate regions, in 2015, 45% (tropical and temperate climate) and in 2019, 54.05% in continental climate. The studies in Nigeria with only a difference of two years show a prevalence of 43.3% in 2001 and 2.1% in 2003 in tropical zones. Turkey reports in 2004 a prevalence of 79.1% and in 2005 of 2.97% in lambs with wide variation too in the temperate climate zone. With all these examples, it is evident that generalizations cannot be made, neither by country nor continent. These results only prove the parasite's worldwide distribution throughout different climate zones, ages and health status of sheep.

## 6. Economic loss

Economic loss is reflected not only in mortality, but in retarded growth rates, decreased feed conversion rate, poor carcass quality, veterinary assistance and increased costs due to extra care. Moreover, healthy sheep can shed oocysts, specially periparturient ewes, with the possibility of maintaining the infection within the flock [88] and contaminate the environment [73].

*Cryptosporidium* causes diarrheas in developing countries, and it has been found that in developed countries it has become a serious problem, due to the limited number of parasitic protozoa in their surveillance programmes, as well as being a neglected disease caused by high confidence in hygiene, municipal sanitization services and good agriculture and livestock practices [18].

The model GloWPa-Crypto was used in [89] and differences were found in oocyst load in extensive and intensive systems, in which intensive ones provided a higher load directly on land.

## 7. Prevention, control and treatment

Transmission of *Cryptosporidium spp.* is influenced by reservoirs and environmental characteristics. Oocysts are sensible to high temperature and desiccation. Direct sunlight for several hours entirely inactivates oocysts [90]. To prevent and protect lambs from cryptosporidiosis, colostrum is fundamental. Hygiene measures are very important to destroy the parasite and prevent its transmission in between

animals and from the environment. Cleaning and disinfecting pens and buildings, the use of clean straw beds, avoiding high number of ewes in the parturition area and separating healthy from ill animals during diarrhea outbreaks, as well as an appropriate and in time administration of colostrum to lambs, aid in the prevention of cryptosporidial outbreaks and decreases the morbidity and mortality in sheep herds [91].


There are three main ways for treating cryptosporidiosis in farms: using drugs (most of them have been proven at laboratory level), fluid therapy and reducing the quantity of oocysts excreted to the environment [92].

There is no specific drug targeted against cryptosporidiosis, although more than 140 active principles have been tested *in vivo* and *in vitro* but none of them have been useful for eliminating cryptosporidiosis from the infected animal. This can be explained because of its particular intracellular but extracytoplasmatic location in the intestine, making it resistant to antimicrobials as well as a difficult drug target. The first attempts to cure cryptosporidiosis were based on using the same drugs as the ones for genetically related pathogens (Apicomplexa) such as *Toxoplasma* and *Plasmodium*, but are not suitable against *Cryptosporidium* because of differences in its cell biology and biochemistry: it has no apicoplast organelle, neither citric acid cycle nor cytochrome respiratory chain. Due to the fact that cryptosporidiosis has had a higher impact in underdeveloped countries, the strategy for treatment has been the use of existing medications [14].

Significant results have been found when using halofuginone lactate [93], which is the only one licensed treatment for cryptosporidiosis in calves, thought to interfere with the merozoite and sporozoite stages of *Cryptosporidium* and proven effective in the control of oocyst shedding [94]. It is an antiprotozoal drug for *Eimeria* and *Theileria*, derivative of quinazolinone [95], used both for prevention and treatment [96]. Other drugs that have also reduced the excreted number of oocysts [40] are: paramomycin, (an aminoglycoside broad-spectrum antibiotic, poorly absorbed in the gut, remaining active in the lumen [97]), metronidazole (a nitroimidazole used against giardiasis [95]), benzoxaborole (a synthetic boron-heterocyclic compound that inhibits essential enzymes, used as an antimalarial) and occidiofungin (a broad-spectrum glycolipopeptide with antifungal activity) [98, 99]. Nitazoxanide (nitratiazole benzamide derivative, licensed in humans as treatment for cryptosporidiosis [95]), beta-cyclodextrin (a cyclic oligosaccharide made up of glucose residues [100], used as pharmaceutical excipient [101]) as well as colostrum preparations, probiotics and decoquinate (coccidiostat quinolone derivative [102]) have also been used [103]. Therefore, the following sanitary and breeding recommendations are essential: sectioning of age groups, assurance of sufficient colostrum supply and intake, isolation of sick animals from the healthy ones, avoiding over-population, cleanliness and dryness of the environment, disinfection of premises and daily cleaning of equipment [93].

As alternative treatments, experiments in mice using watery and alcoholic extracts from *Curcuma longa* and *Coriandrum sativum* have been effective in reducing the number of excysted oocysts [104]. *Peganum harmala*, *Artemisia herb-alba* and *Olea europea* have shown similar results [105].

The most important measure to lessen the clinical signs of cryptosporidiosis is oral or intravenous fluid therapy. To decrease the spread of the disease, it is imperative to avoid oocyst shedding to the environment. For decontamination of surfaces, over 35 disinfectants have been tested, but only 5 are effective: 50% ammonia, 3% hydrogen peroxide, 10% formalin, Exspor and Oo-cide. Formaldehyde or ammonia gas used for steam heat sterilization and fumigation are also recommended [103]. UV radiation and ethylene oxide have been reported to be effective [106].



**Figure 2.**  
*Cryptosporidium spp.* in fecal smear (Ziehl-Neelsen stain, oil immersion, 100X). Arrows show deep pink oocysts. (photograph by Alonso-Fresán).

Not only lambs may be infected through the fecal-oral route, but they can also get infected by drinking contaminated water. This may happen due to the high number of oocysts released to the environment and their resistance to chlorine and other conventional disinfectants used in water treatment processes. Oocyst concentrations of more than 150/L of surface water have been found in agricultural run-off water [15]. Therefore, another effective way of preventing *Cryptosporidium* spread into the environment and watersheds is using vegetation buffers, which aid in the retention of oocysts (**Figure 2**) [107].

Proper composting of manure (>60°C) inactivates oocysts and reduces the risk of their viability. Slurry storage produces ammonia and low pH, and anaerobic digestion using mesophilic and thermophilic bacteria also helps reducing oocyst viability. Fencing sheep away from water sources prevents water contamination with feces [108].

Even though many antigenic target candidates have been characterized, there is still no vaccine with proven effectiveness or ideal cost–benefit ratio [109].

## 8. Conclusion

*Cryptosporidium spp.* is a ubiquitous parasite that directly affects animal welfare by producing self-limiting diarrheas in immunocompetent hosts. In lambs, it is capable of producing a weakening diarrhea which may lead to death. The disease may be complicated by the interaction of other pathogens such as *E. coli* or *Salmonella spp.* Asymptomatic sheep may excrete oocysts, becoming a source of infection and contaminating the environment. Prevalence worldwide in sheep herds may be up to 100%. To prevent the disease, hygiene measures should be taken to avoid environmental contamination. Pregnant ewes should be separated from the herd during and after parturition. If lambs get infected, they should be treated in the same way as with any other diarrheic disease, with fluid therapy.

Many active principles have been tested, as well as alternative natural remedies such as plant extracts in an attempt to cure cryptosporidiosis which have only shown to reduce the number of oocysts excreted, mildening the disease in lambs and reducing environmental contamination, but none of them cures the disease. Vaccines are still in development as means of prevention. Even though there is neither a specific treatment for curing cryptosporidiosis nor a vaccine, manure should be properly composted and to avoid water contamination, vegetation buffers may be used.

### **Author details**

María Uxúa Alonso Fresán<sup>1\*</sup> and Alberto Barbabosa Pliego<sup>2</sup>

1 Faculty of Veterinary Medicine and Zootechnique, Autonomous University of the State of Mexico, Toluca, Mexico

2 Research Investigation Center on Animal Health, Faculty of Veterinary Medicine and Zootechnique, Autonomous University of the State of Mexico, Toluca, Mexico

\*Address all correspondence to: [muaf@uaemex.mx](mailto:muaf@uaemex.mx)

### **IntechOpen**

© 2021 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 



## References

- [1] Åkerfeldt MP, Gunnarsson S, Bernes G, Blanco-Penedo I. Health and welfare in organic livestock production systems—a systematic mapping of current knowledge. *Org agric* [Internet]. 2020; Available from: <http://dx.doi.org/10.1007/s13165-020-00334-y>
- [2] McCausland C. The five freedoms of animal welfare are rights. *J Agric Environ Ethics*. 2014;27(4):649-662.
- [3] Oie. *Terrestrial Animal Health Code* 2019. 28th ed. Office International Des Epizooties; 2019.
- [4] Animal Welfare Network Wales. Code of best practice for animal welfare establishments [Internet]. Welsh Government; [cited 2021 Jun 19]. Available from: <https://gov.wales/sites/default/files/publications/2020-09/animal-welfare-establishments-code-of-best-practice.pdf>
- [5] Magnusson U. Animal Health and Welfare for a Sustainable Livestock sector [Internet]. K-state.edu. [cited 2021 Jun 19]. Available from: <https://www.ksre.k-state.edu/news/stories/2019/09/Animal-Health-and-Welfare.pdf>
- [6] AWIN Welfare assessment protocol for sheep [Internet]. Uni-sz.bg. [cited 2021 Jun 19]. Available from: [http://uni-sz.bg/truni11/wp-content/uploads/biblioteka/file/TUNI10015667\(1\).pdf](http://uni-sz.bg/truni11/wp-content/uploads/biblioteka/file/TUNI10015667(1).pdf)
- [7] Schoenian S. Diarrhea (scours) in small ruminants [Internet]. Osu.edu. [cited 2021 Jun 19]. Available from: <https://u.osu.edu/sheep/2019/06/11/diarrhea-scours-in-small-ruminants/>
- [8] Kerie Y. Cryptosporidiosis: Its importance and diagnostic techniques in farm animals. *Int j vet health sci res*. 2019;236-241.
- [9] Foster DM, Smith GW. Pathophysiology of diarrhea in calves. *Vet Clin North Am Food Anim Pract*. 2009;25(1):13-36, xi.
- [10] Raja Abdul Rahman RNRI, Md Isa ML, Mat Yusof A. A review of *Cryptosporidium spp.* Infection in livestock. *J Teknol* [Internet]. 2017;79(6). Available from: <http://dx.doi.org/10.11113/jt.v79.10330>
- [11] Guy RA, Yanta CA, Muchaal PK, Rankin MA, Thivierge K, Lau R, et al. Molecular characterization of *Cryptosporidium* isolates from humans in Ontario, Canada. *Parasit Vectors*. 2021;14(1):69.
- [12] Shanmathi S, Yashica KA, Gunasekaran P. Epidemiology and diagnosis of cryptosporidiosis: A review [Internet]. *Entomoljournal.com*. [cited 2021 Jun 19]. Available from: <https://www.entomoljournal.com/archives/2020/vol8issue4/PartH/8-3-183-107.pdf>
- [13] Gerace E, Lo Presti VDM, Biondo C. *Cryptosporidium* infection: Epidemiology, pathogenesis, and differential diagnosis. *Eur J Microbiol Immunol (Bp)*. 2019;9(4):119-123.
- [14] Bones AJ, Jossé L, More C, Miller CN, Michaelis M, Tsaousis AD. Past and future trends of *Cryptosporidium in vitro* research. *Experimental Parasitology*. 2019; 196:28-37.
- [15] Hassan EM, Örmeci B, DeRosa MC, Dixon BR, Sattar SA, Iqbal A. A review of *Cryptosporidium spp.* and their detection in water. *Water Sci Technol*. 2021;83(1):1-25.
- [16] Mamedova S, Karanis P. *Cryptosporidium spp.* Infections in Livestock and Wild Animals in Azerbaijan Territory. *Environmental Sciences Proceedings*. 2020;2(1):44.

- [17] Tzipori S, Widmer G. A hundred-year retrospective on cryptosporidiosis. *Trends in Parasitology*. 2008;24(4):184-189.
- [18] Ahmed SA, Karanis P. *Cryptosporidium* and cryptosporidiosis: the perspective from the Gulf countries. *International Journal of Environmental Research and Public Health*. 2020;17(18):6824.
- [19] Raja Abdul Rahman, RNRI, Md Isa ML, Mat Yusaf A. A review of *Cryptosporidium spp* infection in livestock. 2017;79(6):99-109.
- [20] Silva NRS, Falci VSN, Azevedo JSC, Chaplin EL, Araujo FAP. *Cryptosporidium parvum* em ovinos no município de Guaíba, RS [Internet]. 1990. Available from: <http://www.ufrgs.br/actavet/1-29/1990.pdf>
- [21] Kaminjolo JS, Adesiyun AA, Loregnard R, Kitson-Piggott W. Prevalence of *Cryptosporidium* oocysts in livestock in Trinidad and Tobago. *Vet Parasitol*. 1993;45(3-4):209-213.
- [22] Fatimah CTNI, Lee CC, Azri A, Rafie D, Fazlina B, Salim NB, et al. The occurrence and epidemiology of enteropathogens and diarrhea in neonatal lambs. *Jurnal Veterinar Malaysia*. 1995;7(1):27-29.
- [23] Kambarage DM, Kimera SI, Kusiluka LJM, Mtambo MMA. Prevalence of *Eimeria* and *Cryptosporidium* oocysts in cattle, sheep and goats in Morogoro region, Tanzania. *Journal of Applied Animal Research*. 1996;9(1):73-78.
- [24] Agunyole CA, Cadmus SIB, Sansi JAA. A survey of *Cryptosporidium parvum* infection in calves, goat, kids and lambs in Ibadan, Nigeria. *Bulletin of Animal Health and Production in Africa*. 2001;49:264-267.
- [25] Ohaeri CC, Iwu CJ. Prevalence of *Cryptosporidium* species in some domestic animals. *Journal of Sustainable Agriculture and the Environment*. 2003;5(2):309-312.
- [26] Koinari M, Lymbery AJ, Ryan UM. *Cryptosporidium* species in sheep and goats from Papua New Guinea. *Exp Parasitol*. 2014;141:134-137.
- [27] Daniels ME, Shrivastava A, Smith WA, Sahu P, Odagiri M, Misra PR, et al. *Cryptosporidium* and *Giardia* in humans, domestic animals, and village water sources in rural India. *Am J Trop Med Hyg*. 2015;93(3):596-600.
- [28] Castro-del Campo N, Castro-del Campo N, Enriquez-Verdugo, I, Portillo-Loera, JJ, y Gaxiola-Camacho SM.. Prevalencia y factores de riesgo asociados a *Cryptosporidium spp.* en corderos de pelo del municipio de Culiacán, Sinaloa, México. *Rev Cient-Fac Cien V*. 2017;XXVII(4):148-156.
- [29] Vitela-Mendoza I, Guillen Lorenzo V, Cruz Vazquez CR, Medina Esparza LE, Ramos Parra M. Frecuencia de casos e identificación molecular de *Cryptosporidium spp.* en corderos lactantes mantenidos en pastoreo en el estado de Veracruz, México. *Rev Mex Cienc Pecu*. 2018;9(3):420-434.
- [30] Abou-Eisha AM. Cryptosporidial infection in man and farm animals in Ismailia Governorate. *Veterinary Medical Journal Giza*. 1994;42(2):107-111.
- [31] El-Hohary AH, Abdel-Latif AM. Zoonotic importance of cryptosporidiosis among some animals at Gharbia Province in Egypt. *Indian Journal of Animal Sciences*. 1998;67(4):305-307.
- [32] Dezfouli MRM, Mesghi B. Epidemiological study of cryptosporidial infestation of man and animals. *Journal of the Faculty of Veterinary Medicine, University of Tehran*. 2002;57(1):87-92.



- [33] Mahdi NK, Ali NH. Cryptosporidiosis among animal handlers and their livestock in Basrah, Iraq. *East Afr Med J*. 2002;79(10):550-553.
- [34] Nassif MN, Amer SA, A OS. Some studies on ovine and caprine cryptosporidiosis concerning prevalence and electrophoretic pattern of blood serum protein. *Assiut Veterinary Medical Journal*. 2002;47(94):249-262.
- [35] Causapé AC, Quílez J, Sánchez-Acedo C, del Cacho E, López-Bernad F. Prevalence and analysis of potential risk factors for *Cryptosporidium parvum* infection in lambs in Zaragoza (northeastern Spain). *Vet Parasitol*. 2002;104(4):287-298.
- [36] Eisa MI, Mohamed AA. Role of enteric pathogens in enteritis in lambs, goat kids and children and their zoonotic importance. *Veterinary Medical Journal Giza*. 2004;52(1):41-59.
- [37] Martínez GA, Vitela MIV, Cruz-Vázquez CR, Ramos PM. Presencia de oocistos de *Cryptosporidium spp.*, durante el parto y posparto, en borregos pelibuey y corderos latantes de un rebaño en el Estado de Aguascalientes. In: *Memorias de la XLIII Reunión Nacional de Investigación Pecuaria Sinaloa*. 2007.
- [38] Mahfouz ME, Mira N, Amer S. Prevalence and genotyping of *Cryptosporidium spp.* in farm animals in Egypt. *J Vet Med Sci*. 2014;76(12):1569-1575.
- [39] Shafiq MAB, Maqbool A, Khan UJ, Lateef M, Ijaz M. Prevalence, water borne transmission and chemotherapy of cryptosporidiosis in small ruminants. *Pakistan J Zool*. 2015;47(6):1715-1721.
- [40] Firoozi Z, Sazmand A, Zahedi A, Astani A, Fattahi-Bafghi A, Kiani-Salmi N, et al. Prevalence and genotyping identification of *Cryptosporidium* in adult ruminants in central Iran. *Parasit Vectors*. 2019;12(1):510.
- [41] Gorman T, Alcaíno H, Mandry P. Cryptosporidiosis en ovinos y caprinos de la zona central de Chile. *Archivos de Medicina Veterinaria*. 1990;XXII(2):155-158.
- [42] Özer E, Erdoğan SZ, Köroğlu E. Studies on the incidence of *Cryptosporidium* in calves and lambs in the Eilaziğ area. *Doğa, Türk Veterinerlik ve Hayvancılık Dergisi*. 1990;14(3):439-445.
- [43] Nouri M, Karami M. Asymptomatic cryptosporidiosis in nomadic shepherds and their sheep. *J Infect*. 1991;23(3):331-3.
- [44] Valenzuela G, Sci MV, Grandón W, Quintana I, Tadich TM. Prevalencia de *Cryptosporidium spp.* en corderos muertos en la provincia de Valdivia, Chile. *Archivos de Medicina Veterinaria*. 1991;XXIII(1):81- 83.
- [45] Villacorta I, Ares-Mazas E, Lorenzo MJ. *Cryptosporidium parvum* in cattle, sheep and pigs in Galicia (N.W. Spain). *Vet Parasitol*. 1991;38(2-3):249-252.
- [46] Minas A, Koutsoukou-Hartona W, Papisavvas M. Epidemiological survey of diarrhoeic faeces of lambs and kids for cryptosporidial oocysts in the Larissa region. *Bulletin of the Hellenic Veterinary Medical Society*. 1994;44(2).
- [47] Muñoz M, Alvarez M, Lanza I, Cármenes P. Role of enteric pathogens in the aetiology of neonatal diarrhoea in lambs and goat kids in Spain. *Epidemiol Infect*. 1996;117(1):203-211.
- [48] JianMin Y, JiaMing B, TingHe C, Xiao S, GuoHuo L, GuoXin C, et al. Investigation of cryptosporidiosis in domestic animals and poultry and experimental infection test. *Chinese*

Journal of Veterinary Medicine.  
1999;25(12):5-7.

[49] Erman H, Beyazit A, Öz I. The prevalence of cryptosporidiosis in lambs and goat kids in Izmir. Bornova Veteriner Kontrol ve Arestirma Enstitüsü Dergisi. 2000;25(39):33-38.

[50] Majewska AC, Werner A, Sulima P, Luty T. Prevalence of *Cryptosporidium* in sheep and goats bred on five farms in west-central region of Poland. Veterinary Parasitology. 2000;89(4).

[51] Pilarczyk B, Balicka-Ramisz A. Prevalence of *Cryptosporidium spp.* in animals in Western Pomerania. Folia Universitatis Agriculturae Stetinensis, Zootechnica. 2001;42:141-144.

[52] Green RE, Amarante AFT, Mascarini LM. The seasonal distribution of *Cryptosporidium* oocysts in sheep raised in the State of São Paulo. Revista Brasileira de Parasitologia Veterinária. 2004;13(3):125-127.

[53] Ulutaş B, Voyvoda H. Cryptosporidiosis in diarrhoeic lambs on a sheep farm. Türkiye Parazitoloji Dergisi. 2004;28(1):15-17.

[54] Zhuang JC, Xie HY, Liu WL, Deng YX, Huo RS, Pan XP, et al. Investigation on six kinds of animals infected with *Cryptosporidium* in the poultry and domestic animals wholesale market of Guangzhou City. Journal of Tropical Medicine (Guangzhou). 2004;4(6):710-711.

[55] Green RE, Amarante AFT, Mascarini LM. Patterns of shedding of cryptosporidial oocysts by ewes and lambs kept indoors. Brazilian Journal of Veterinary Research and Animal Science. 2005;42(3):204-209.

[56] Alonso-Fresán MU, García-Alvarez A, Salazar-García F, Vázquez-Chagoyán JC, Pescador-Salas N, Saltijeral-Oaxaca J.

Prevalence of *Cryptosporidium spp.* in asymptomatic sheep in family flocks from Mexico State. J Vet Med B Infect Dis Vet Public Health. 2005;52(10):482-483.

[57] Sevinç F, Uslu U, Derinbay Ö. The prevalence of *Cryptosporidium parvum* in lambs around Konya. Türk Veterinerlik ve Hayvancılık Dergisi. 2005;29(5):1191-1194.

[58] Craig BH, Pilkington JG, Kruuk LEB, Pemberton JM. Epidemiology of parasitic protozoan infections in Soay sheep (*Ovis aries* L.) on St Kilda. Parasitology. 2007;134(Pt 1):9-21.

[59] Zorana M B, Sofija K-R P, Zoran K. *Cryptosporidium* infection in lambs and goat kids in Serbia. Acta Vet (Beogr). 2006;56(1):49-54.

[60] Santín M, Trout JM, Fayer R. Prevalence and molecular characterization of *Cryptosporidium* and *Giardia* species and genotypes in sheep in Maryland. Vet Parasitol. 2007;146(1-2):17-24.

[61] Castro-Hermida JA, Almeida A, González-Warleta M, Correia da Costa JM, Rumbo-Lorenzo C, Mezo M. Occurrence of *Cryptosporidium parvum* and *Giardia duodenalis* in healthy adult domestic ruminants. Parasitol Res. 2007;101(5):1443-1448.

[62] Soltane R, Guyot K, Dei-Cas E, Ayadi A. Prevalence of *Cryptosporidium spp.* (Eucoccidiorida: Cryptosporiidae) in seven species of farm animals in Tunisia. Parasite. 2007;14(4):335-338.

[63] Pritchard GC, Marshall JA, Giles M, Chalmers RM, Marshall RN. *Cryptosporidium parvum* infection in orphan lambs on a farm open to the public. Vet Rec. 2007;161(1):11-14.

[64] Panousis N, Diakou A, Giadinis N, Papadopoulos E, Karatzias H,

Haralampidis S. Prevalence of *Cryptosporidium* infection in sheep flocks with a history of lambs' diarrhoea [Internet]. Revmedvet.com. [cited 2021 Jun 21]. Available from: [https://www.revmedvet.com/2008/RMV159\\_528\\_531.pdf](https://www.revmedvet.com/2008/RMV159_528_531.pdf)

[65] Geurden T, Thomas P, Casaert S, Vercruysse J, Claerebout E. Prevalence and molecular characterisation of *Cryptosporidium* and *Giardia* in lambs and goat kids in Belgium. *Vet Parasitol.* 2008;155(1-2):142-5.

[66] Yang R, Jacobson C, Gordon C, Ryan U. Prevalence and molecular characterisation of *Cryptosporidium* and *Giardia* species in pre-weaned sheep in Australia. *Vet Parasitol.* 2009;161(1-2):19-24.

[67] Tzanidakis N, Sotiraki S, Claerebout E, Ehsan A, Voutzourakis N, Kostopoulou D, et al. Occurrence and molecular characterization of *Giardia duodenalis* and *Cryptosporidium spp.* in sheep and goats reared under dairy husbandry systems in Greece. *Parasite.* 2014;21:45.

[68] Ahamed I, Yadav A, Katoch R, Godara R, Saleem T, Nisar NA. Prevalence and analysis of associated risk factors for *Cryptosporidium* infection in lambs in Jammu district. *J Parasit Dis.* 2015;39(3):414-417.

[69] Romero-Salas D, Alvarado-Esquivel C, Cruz-Romero A, Aguilar-Domínguez M, Ibarra-Priego N, Merino-Charrez JO, et al. Prevalence of *Cryptosporidium* in small ruminants from Veracruz, Mexico. *BMC Vet Res.* 2016;12(1):14.

[70] Shafieyan H, Alborzi A, Hamidinejat H, Tabandeh MR, Hajikolaei MRH. Prevalence of *Cryptosporidium spp.* in ruminants of Lorestan province, Iran. *J Parasit Dis.* 2016;40(4):1165-1169.

[71] Kaupke A, Michalski MM, Rzeżutka A. Diversity of *Cryptosporidium* species occurring in sheep and goat breeds reared in Poland. *Parasitol Res.* 2017;116(3):871-879.

[72] Baroudi D, Hakem A, Adamu H, Amer S, Khelef D, Adjou K, et al. Zoonotic *Cryptosporidium* species and subtypes in lambs and goat kids in Algeria. *Parasit Vectors.* 2018;11(1):582.

[73] Bordes L, Houert P, Costa D, Favennec L, Vial-Novella C, Fidelle F, et al. Asymptomatic *Cryptosporidium* infections in ewes and lambs are a source of environmental contamination with zoonotic genotypes of *Cryptosporidium parvum*. *Parasite.* 2020;27:57.

[74] Dessì G, Tamponi C, Varcasia A, Sanna G, Pipia AP, Carta S, et al. *Cryptosporidium* infections in sheep farms from Italy. *Parasitol Res.* 2020;119(12):4211-4218.

[75] Essendi MW, Muleke C, Otachi E, Miheso M, Kyule D. Prevalence of zoonotic *Cryptosporidium spp.* Isolates in Njoro sub-county, Nakuru county, Kenya. *Afr J Infect Dis.* 2021;15(2):3-9.

[76] Xiao L, Herd RP, Rings DM. Diagnosis of *Cryptosporidium* on a sheep farm with neonatal diarrhea by immunofluorescence assays. *Vet Parasitol.* 1993;47(1-2):17-77.

[77] Nagy B. Epidemiological information on *Cryptosporidium parvum* infection of domestic mammals in Hungary. *Magyar Állatorvosok Lapja.* 1995;50(3):139-144.

[78] Sari B, Arslan MO, Gicik Y, Kara M, Taşçi GT. The prevalence of *Cryptosporidium* species in diarrhoeic lambs in Kars province and potential risk factors. *Trop Anim Health Prod.* 2009;41(5):819-826.



- [79] Jafari R, Maghsood AH, Fallah M. Prevalence of *Cryptosporidium* infection among livestock and humans in contact with livestock in Hamadan district, Iran, 2012. J Res Health Sci. 2013;13(1):86-89.
- [80] Li P, Cai J, Cai M, Wu W, Li C, Lei M, et al. Distribution of *Cryptosporidium* species in Tibetan sheep and yaks in Qinghai, China. Vet Parasitol. 2016;215:58-62.
- [81] Bhat AM, Malik HU, Wani NM, Paul S, Gupta S, Dolma T, et al. First report of *Cryptosporidium spp.* infection in sheep population of Ladakh, India. J Parasit Dis. 2019;43(3):513-516.
- [82] Olson ME, Thorlakson CL, Deselliers L, Morck DW, McAllister TA. *Giardia* and *Cryptosporidium* in Canadian farm animals. Vet Parasitol. 1997;68(4):375-381.
- [83] Maurya PS, Rakesh RL, Pradeep B, Kumar S, Kundu K, Garg R, et al. Prevalence and risk factors associated with *Cryptosporidium spp.* infection in young domestic livestock in India. Trop Anim Health Prod. 2013;45(4):941-946.
- [84] Gharekhani J, Heidari H, Youssefi M. Prevalence of *Cryptosporidium* infection in sheep in Iran. Turkiye Parazitolo Derg. 2014;38(1):22-25.
- [85] Yang R, Jacobson C, Gardner G, Carmichael I, Campbell AJD, Ng-Hublin J, et al. Longitudinal prevalence, oocyst shedding and molecular characterisation of *Cryptosporidium* species in sheep across four states in Australia. Vet Parasitol. 2014;200(1-2):50-58.
- [86] Bokaie S, Jokar M, Rabiee M, Rahmanian V, Dehesh P, Hasannejad H, et al. Prevalence of cryptosporidiosis in animals in Iran: A systematic review and metaanalysis. Asian Pac J Trop Med. 2021;14(3):99.
- [87] King B, Monis PT. Critical processes affecting *Cryptosporidium* oocyst survival in the environment. Parasitology. 2007;134:309-323.
- [88] Santín M. Clinical and subclinical infections with *Cryptosporidium* in animals. N Z Vet J. 2013;61(1):1-10.
- [89] Vermeulen LC, Benders J, Medema G, Hofstra N. Global *Cryptosporidium* loads from livestock manure. Environ Sci Technol. 2017;51(15):8663-8671.
- [90] Samie A, Al-Qahtani A, El Bakri A, Ehdaie B. Challenges and innovative strategies to interrupt *Cryptosporidium* transmission in resource-limited settings. Curr Trop Med Rep. 2015;2(3):161-170.
- [91] Kerie, Doctor of Veterinary Medicine (DVM) and Master of Microbiology (MSc), Bahir dar Animal Health Diagnostic and Investigation Laboratory (BAHDIL), Ethiopia. Cryptosporidiosis: Its importance and diagnostic techniques in farm animals. Int J Vet Health Sci Res. 2019;236-41.
- [92] Dinler C; Ulutas B. Cryptosporidiosis in ruminants: update and current therapeutic approach. American Journal of Animal and Veterinary Sciences. 2017;12(3):96-103.
- [93] Hoste H, Ehrhardt N, Paraud C, Rieux A, Mercier P, Valas S, et al. Recherche en pathologie caprine : applications et perspectives. INRA Prod Anim. 2012;25(3):245-258.
- [94] Aydogdu U, Isik N, Ekici OD, Yildiz R, Sen I, Coskun A. Comparison of the effectiveness of halofuginone lactate and paromomycin in the treatment of calves naturally infected

with *Cryptosporidium parvum*. Acta Sci Vet. 2018;46(1):9.

[95] Akinci IY, Albay MK. Etiological treatment of protozoer diarrhea in neonatal ruminants. MAE Vet Fak Derg. 2020;5(1):28-33.

[96] Thomson S, Hamilton CA, Hope JC, Katzer F, Mabbott NA, Morrison LJ, Innes EA. Bovine cryptosporidiosis: impact, host-parasite interaction and control strategies. Vet Res. 2017;48:42.

[97] Hameed OA, Abu-Zeid TESA, Rasool G, Makki A, Taha MK, Duquesne B. The prevalence of *Cryptosporidium* infection in neonatal diarrhoeic sheep lambs with observation of resolution using Paromomycin treatment. Journal of Animal Sciences and Livestock Production. 2019;3(2).

[98] Lunde CS, Stebbins EE, Jumani RS, Hasan MM, Miller P, Barlow J, et al. Identification of a potent benzoxaborole drug candidate for treating cryptosporidiosis. Nat Commun. 2019;10(1):2816.

[99] Ma J, Guo F, Jin Z, Geng M, Ju M, Ravichandran A, et al. Novel antiparasitic activity of the antifungal lead occidiofungin. Antimicrob Agents Chemother [Internet]. 2020;64(8). Available from: <http://dx.doi.org/10.1128/AAC.00244-20>

[100] Martini A, Torricelli C, Muggetti, De Ponti R. Use of dehydrated beta-cyclodextrin as pharmaceutical excipient. Drug development and industrial pharmacy. 1994;20(15):2381-2393.

[101] Castro-Hermida JA, González-Losada Y, Freire-Santos F, Mezo-Menéndez M, Ares-Mazás E. Evaluation of  $\beta$ -cyclodextrin against natural infections of cryptosporidiosis in calves. Veterinary Parasitology. 2001;101(2):85-89.

[102] Taylor MA, Bartram DJ. The history of decoquinate in the control of coccidial infections. Journal of Veterinary Pharmacology and Therapeutics. 2012;35(5):417-427.

[103] Ghazy AA, Shafy SA-, Shaapan RM. Cryptosporidiosis in animals and man: 3. Prevention and control. Asian J Epidemiol. 2016;9(1-3):1-9.

[104] Obiad HM, Al-Alousi TI, Al-Jboori AH. The in vivo effect of some medicinal plant extracts on *Cryptosporidium* parasite. J. of University of Anbar for Pure Science. 2012;6(3).

[105] Al-Hamairy AK, Mugheer AH, Al-Dulaimi FHA. Prevalence of *Cryptosporidium spp.* and treatment by using some plant extracts in Al-Hilla City, Babylon Province. Egypt. J. Exp. Biol. (Zool.). 2012;8(2):287-293.

[106] Spickler AR. Cryptosporidiosis [Internet]. Iastate.edu. 2018 [cited 2021 Jun 19]. Available from: <http://www.cfsph.iastate.edu/DiseaseInfo/factsheets.php>.

[107] Atwill ER, Tate KW, Pereira MDGC, Bartolome J, Nader G. Efficacy of natural grassland buffers for removal of *Cryptosporidium parvum* in rangeland runoff. J Food Prot. 2006;69(1):177-184.

[108] Innes EA, Chalmers RM, Wells B, Pawlowic MC. A One Health approach to tackle cryptosporidiosis. Trends Parasitol. 2020;36(3):290-303.

[109] Silva DRR da, Oliveira TCB de, Marta BBF, Baptista CB, Bottaro MCZ, Bresciani KDS. Vaccine development for cryptosporidiosis: Systematic review. Res Soc Dev. 2021;10(6):e18910615540.