

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



# A Study of Nonlinear Boundary Value Problem

*Noureddine Bouteraa and Habib Djourdem*

## Abstract

In this chapter, firstly we apply the iterative method to establish the existence of the positive solution for a type of nonlinear singular higher-order fractional differential equation with fractional multi-point boundary conditions. Explicit iterative sequences are given to approximate the solutions and the error estimations are also given. Secondly, we cover the multi-valued case of our problem. We investigate it for nonconvex compact valued multifunctions via a fixed point theorem for multivalued maps due to Covitz and Nadler. Two illustrative examples are presented at the end to illustrate the validity of our results.

**Keywords:** Positive solution, Uniqueness, Iterative sequence, Green's function, Fractional differential equation and inclusion, Existence, Nonlocal boundary value problem, Fixed point theorem

## 1. Introduction

In this chapter, we are interested in the existence of solutions for the nonlinear fractional boundary value problem (BVP)

$$\begin{cases} D_{0+}^{\alpha}u(t) + f(t, u(t)) = 0, & t \in (0, 1), \\ u^{(i)}(0) = 0, i \in \{0, 1, 2, \dots, n-2\}, D_{0+}^{\beta}u(1) = \sum_{j=1}^p a_j D_{0+}^{\beta}u(\eta_j). \end{cases} \quad (1)$$

We also cover the multi-valued case of problem

$$\begin{cases} -D_{0+}^{\alpha}u(t) \in F(t, u(t)), & t \in (0, 1), \\ u^{(i)}(0) = 0, i \in \{0, 1, 2, \dots, n-2\}, D_{0+}^{\beta}u(1) = \sum_{j=1}^p a_j D_{0+}^{\beta}u(\eta_j), \end{cases} \quad (2)$$

where  $D_{0+}^{\alpha}$ ,  $D_{0+}^{\beta}$  are the standard Riemann-Liouville fractional derivative of order

$$\alpha \in (n-1, n], \beta \in [1, n-2] \text{ for } n \in \mathbb{N}^* \text{ and } n \geq 3,$$

where  $D_{0+}^{\alpha}$ ,  $D_{0+}^{\beta}$  are the standard Riemann-Liouville fractional derivative of order  $\alpha \in (n-1, n], \beta \in [1, n-2]$  for  $n \geq 3$ , the function  $f \in C((0, 1) \times \mathbb{R}, \mathbb{R})$ , the

multifunction  $F : [0, 1] \times \mathbb{R} \rightarrow 2^{\mathbb{R}}$  are allowed to be singular at  $t = 0$  and/or  $t = 1$  and  $a_j \in \mathbb{R}^+$ ,  $j = 1, 2, \dots, p$ ,  $0 < \eta_1 < \eta_2 < \dots < \eta_p < 1$ , for  $p \in \mathbb{N}^*$ .

The first definition of fractional derivative was introduced at the end of the nineteenth century by Liouville and Riemann, but the concept of non-integer derivative and integral, as a generalization of the traditional integer order differential and integral calculus, was mentioned already in 1695 by Leibniz [1] and L'Hospital [2]. In fact, fractional derivatives provide an excellent tool for the description of memory and hereditary properties of various materials and processes. The mathematical modeling of systems and processes in the fields of physics, chemistry, aerodynamics, electrodynamics of complex medium, polymer rheology, Bode's analysis of feedback amplifiers, capacitor theory, electrical circuits, electro-analytical chemistry, biology, control theory, fitting of experimental data, involves derivatives of fractional order. In consequence, the subject of fractional differential equations is gaining much importance and attention. For more details we refer the reader to [1–6] and the references cited therein.

Boundary value problems for nonlinear differential equations arise in a variety of areas of applied mathematics, physics and variational problems of control theory. A point of central importance in the study of nonlinear boundary value problems is to understand how the properties of nonlinearity in a problem influence the nature of the solutions to the boundary value problems. The multi-point boundary conditions are important in various physical problems of applied science when the controllers at the end points of the interval (under consideration) dissipate or add energy according to the sensors located, at intermediate points, see [7, 8] and the references therein. We quote also that realistic problems arising from economics, optimal control, stochastic analysis can be modeled as differential inclusion. The study of fractional differential inclusions was initiated by El-Sayed and Ibrahim [9]. Also, recently, several qualitative results for fractional differential inclusion were obtained in [10–13] and the references therein.

The techniques of nonlinear analysis, as the main method to deal with the problems of nonlinear differential equations (DEs), nonlinear fractional differential equations (FDEs), nonlinear partial differential equations (PDEs), nonlinear fractional partial differential equations (FPDEs), nonlinear stochastic fractional partial differential equations (SFPDEs), plays an essential role in the research of this field, such as establishing the existence, uniqueness and multiplicity of solutions (or positive solutions) and mild solutions for nonlinear of different kinds of FPDEs, FPDEs, SFPDEs, inclusion differential equations and inclusion fractional differential equations with various boundary conditions, by using different techniques (approaches). For more details, see [14–37] and the references therein. For example, iterative method is an important tool for solving linear and nonlinear Boundary Value Problems. It has been used in the research areas of mathematics and several branches of science and other fields. However, Many authors showed the existence of positive solutions for a class of boundary value problem at resonance case. Some recent development for resonant case can be found in [38, 39]. Let us cited few papers. In [40], the authors studied the boundary value problems of the fractional order differential equation:

$$\begin{cases} D_{0+}^{\alpha} u(t) = f(t, u(t)) = 0, & t \in (0, 1), \\ u(0) = 0, \quad D_{0+}^{\beta} u(1) = a D_{0+}^{\beta} u(\eta), \end{cases}$$

where  $1 < \alpha \leq 2$ ,  $0 < \eta < 1$ ,  $0 < a, \beta < 1$ ,  $f \in C([0, 1] \times \mathbb{R}^2, \mathbb{R})$  and  $D_{0+}^{\alpha}$ ,  $D_{0+}^{\beta}$  are the standard Riemann-Liouville fractional derivative of order  $\alpha$ . They obtained the

multiple positive solutions by the Leray-Schauder nonlinear alternative and the fixed point theorem on cones.

In 2020 Li et al. [41] consider the existence of a positive solution for the following BVP of nonlinear fractional differential equation with integral boundary conditions:

$$\begin{cases} ({}^C D_{0+}^q u)(t) + f(t, u(t)) = 0, & t \in [0, 1], \\ u''(0) = 0, \\ \alpha u(0) - \beta u'(0) = \int_0^1 h_1(s)u(s)ds, \\ \gamma u(1) + \delta ({}^C D_{0+}^\sigma u)(1) = \int_0^1 h_2(s)ds, \end{cases}$$

where  $2 < q \leq 3$ ,  $0 < \sigma \leq 1$ ,  $\alpha, \gamma, \delta \geq 0$ , and  $\beta > 0$  satisfying  $0 < \rho(\alpha + \beta)\gamma + \frac{\alpha\delta}{\Gamma(2-\sigma)} < \beta \left[ \gamma + \frac{\delta\Gamma(q)}{\Gamma(q-\sigma)} \right]$ ,  $f : [0, 1] \times [0, +\infty) \rightarrow [0, +\infty)$  and  $h_i (i = 1, 2) : [0, 1] \rightarrow [0, +\infty)$  are continuous. To obtain the existence results, the authors used the well-known GuoKrasnoselskiis fixed point theorem.

In 2017, Rezapour et al. [42] investigated a Caputo fractional inclusion with integral boundary condition for the following problem

$$\begin{cases} {}^c D^\alpha u(t) \in F(t, u(t), {}^c D^\beta u(t), u'(t)), \\ u(0) + u'(0) + {}^c D^\beta u(0) = \int_0^\eta u(s)ds, \\ u(1) + u'(1) + {}^c D^\beta u(1) = \int_0^\nu u(s)ds, \end{cases}$$

where  $1 < \alpha \leq 2$ ,  $\eta, \nu, \beta \in (0, 1)$ ,  $F : [0, 1] \times \mathbb{R} \times \mathbb{R} \times \mathbb{R} \rightarrow 2^{\mathbb{R}}$  is a compact valued multifunction and  ${}^c D^\alpha$  denotes the Caputo fractional derivative of order  $\alpha$ .

In 2018, Bouteraa and Benaicha [10] studied the existence of solutions for the Caputo fractional differential inclusion

$${}^c D^\alpha u(t) \in F(t, u(t), u'(t)), \quad t \in J = [0, 1],$$

subject to three-point boundary conditions

$$\begin{cases} \beta u(0) + \gamma u(1) = u(\eta), \\ u(0) = \int_0^\eta u(s)ds, \\ \beta {}^c D^p u(0) + \gamma {}^c D^p u(1) = {}^c D^p u(\eta), \end{cases}$$

where  $2 < \alpha \leq 3$ ,  $1 < p \leq 2$ ,  $0 < \eta < 1$ ,  $\beta, \gamma \in \mathbb{R}^+$ ,  $F : [0, 1] \times \mathbb{R} \times \mathbb{R} \rightarrow 2^{\mathbb{R}}$  is a compact valued multifunction and  ${}^c D^\alpha$  denotes the Caputo fractional derivative of order  $\alpha$ .

In 2019, Ahmad et al. [43] investigated the existence of solutions for the boundary value problem of coupled Caputo (Liouville-Caputo) type fractional differential inclusions:

$$\begin{cases} {}^C D^\alpha x(t) \in F(t, x(t), y(t)), & t \in [0, T], \quad 1 < \alpha \leq 2, \\ {}^C D^\beta y(t) \in F(t, x(t), y(t)), & t \in [0, T], \quad 1 < \beta \leq 2, \end{cases}$$

subject to the coupled boundary conditions:

$$\begin{aligned} x(0) &= \nu_1 y(T), & x'(0) &= \nu_2 y'(T), \\ y(0) &= \mu_1 x(T), & y'(0) &= \mu_2 x'(T), \end{aligned}$$

where  ${}^C D^\alpha$ ,  ${}^C D^\beta$  denote the Caputo fractional derivatives of order  $\alpha$  and  $\beta$  respectively,  $F, G : [0, T] \times \mathbb{R} \times \mathbb{R}$  are given multivalued maps,  $P(\mathbb{R})$  is the family of all nonempty subsets of  $\mathbb{R}$ , and  $\nu_i, \mu_i, i = 1, 2$  are real constants with  $\nu_i \mu_i \neq 1, i = 1, 2$ .

Inspired and motivated by the works mentioned above, we focus on the uniqueness of positive solutions for the nonlocal boundary value problem (1) with the iterative method and properties of  $f(t, u)$ , explicit iterative sequences are given to approximate the solutions and the error estimations are also given. We also cover the multi-valued case of problem (2) when the right-hand side is nonconvex compact valued multifunctions via a fixed point theorem for multivalued maps due to Covitz and Nadler.

The chapter is organized as follows. In Section 2, we present some notations and lemmas that will be used to prove our main results of problem (1) and we discuss the uniqueness of problem (1). Finally, we give an example to illustrate our result. In Section 3, we introduce some definitions and preliminary results about essential properties of multifunction that will be used in the remainder of the chapter and we present existence results for the problem (2) when the right-hand side is a non-convex compact multifunction. We shall use the fixed point theorem for contraction multivalued maps due to Covitz and Nadler [44] to prove the uniqueness of solution of problem (2). Finally, we give an example to ascertain the main result.

## 2. Existence and uniqueness results for problem (2)

### 2.1 Preliminaries

In this section, we recall some definitions and facts which will be used in the later analysis. These details can be found in the recent literature; see [2, 4, 6, 45–47] and the references therein.

Let  $AC^i([0, 1], \mathbb{R})$  denote the space of  $i$  – times differentiable functions  $u : [0, 1] \rightarrow \mathbb{R}$  whose  $i$  – th derivative  $u^{(i)}$  is absolutely continuous and  $[\alpha]$  denotes the integer part of number  $\alpha$ .

Definition 2.1. Let  $\alpha > 0$ ,  $n - 1 < \alpha < n$ ,  $n = [\alpha] + 1$  and  $u \in AC^n([0, \infty), \mathbb{R})$ .

The Caputo derivative of fractional order  $\alpha$  for the function  $u : [0, +\infty) \rightarrow \mathbb{R}$  is defined by

$${}^C D^\alpha u(t) = \frac{1}{\Gamma(n - \alpha)} \int_0^t (t - s)^{n - \alpha - 1} u^{(n)}(s) ds.$$

The Riemann-Liouville fractional derivative order  $\alpha$  for the function  $u : [0, +\infty) \rightarrow \mathbb{R}$  is defined by

$$D_{0+}^\alpha u(t) = \frac{1}{\Gamma(n - \alpha)} \frac{d^n}{dt^n} \int_0^t (t - s)^{n - \alpha - 1} u(s) ds, \quad t > 0,$$

provided that the right hand side is pointwise defined in  $(0, \infty)$  and the function  $\Gamma : (0, \infty) \rightarrow \mathbb{R}$ , defined by

$$\Gamma(u) = \int_0^\infty t^{u-1} e^{-t} dt,$$

is called Euler's gamma function.

Definition 2.2. The Riemann-Liouville fractional integral of order  $\alpha > 0$  of a function  $u : (0, \infty) \rightarrow \mathbb{R}$  is given by

$$I^\alpha u(t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} u(s) ds, \quad t > 0,$$

provided that the right hand side is pointwise defined in  $(0, \infty)$ .

We recall in the following lemma some properties involving Riemann-Liouville fractional integral and Riemann-Liouville fractional derivative or Caputo fractional derivative which are need in Lemma 2.4.

Lemma 2.1. (([45], Prop.4.3), [46]) Let  $\alpha, \beta \geq 0$  and  $u \in L^1(0, 1)$ . Then the next formulas hold.

- i.  $(D^\beta I^\alpha u)(t) = I^{\alpha-\beta} u(t)$ ,
- ii.  $(D^\alpha I^\alpha u)(t) = u(t)$ ,
- iii.  $I_{0+}^\alpha I_{0+}^\beta u(t) = I_{0+}^{\alpha+\beta} u(t)$ .
- iv. If  $\beta > \alpha > 0$ , then  $D^\alpha t^{\beta-1} = \frac{\Gamma(\beta)t^{\beta-\alpha-1}}{\Gamma(\beta-\alpha)}$ . where  $D^\alpha$  and  $D^\beta$  represents Riemann-Liouville's or Caputo's fractional derivative of order  $\alpha$  and  $\beta$  respectively.

Lemma 2.2 [47]. Let  $\alpha > 0$  and  $y \in L^1(0, 1)$ . Then, the general solution of the fractional differential equation  $D_{0+}^\alpha u(t) + y(t) = 0$ ,  $0 < t < 1$  is given by

$$u(t) = -\frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} y(s) ds + c_1 t^{\alpha-1} + c_2 t^{\alpha-2} + \dots + c_n t^{\alpha-n}, \quad 0 < t < 1,$$

where  $c_0, c_1, \dots, c_{n-1}$  are real constants and  $n = [\alpha] + 1$ .

Based on the previous Lemma 2.2, we will define the integral solution of our problem (1).

Lemma 2.3. Let  $\sum_{j=1}^p a_j \eta_j^{\alpha-\beta-1} \in [0, 1)$ ,  $\alpha \in (n-1, n]$ ,  $\beta \in [1, n-2]$ ,  $n \geq 3$  and  $y(\cdot) \in C[0, 1]$ . Then the solution of the fractional boundary value problem

$$\begin{cases} D_{0+}^\alpha u(t) + y(t) = 0, \\ u^{(i)}(0) = 0, \quad i \in \{0, 1, 2, \dots, n-2\}, \\ D_{0+}^\beta u(1) = \sum_{j=1}^p a_j D_{0+}^\beta u(\eta_j), \end{cases} \quad (3)$$

is given by

$$u(t) = \int_0^1 G(t,s) y(s) ds, \quad (4)$$

where

$$G(t, s) = g(t, s) + \frac{t^{\alpha-1}}{d} \sum_{j=1}^p a_j h(\eta_j, s), \quad (5)$$

$$g(t, s) = \frac{1}{\Gamma(\alpha)} \begin{cases} t^{\alpha-1}(1-s)^{\alpha-\beta-1} - (t-s)^{\alpha-1}, & 0 \leq s \leq t \leq 1, \\ t^{\alpha-1}(1-s)^{\alpha-\beta-1}, & 0 \leq t \leq s \leq 1, \end{cases} \quad (6)$$

$$h(t, s) = \frac{1}{\Gamma(\alpha)} \begin{cases} t^{\alpha-\beta-1}(1-s)^{\alpha-\beta-1} - (t-s)^{\alpha-\beta-1}, & 0 \leq s \leq t \leq 1, \\ t^{\alpha-\beta-1}(1-s)^{\alpha-\beta-1}, & 0 \leq t \leq s \leq 1, \end{cases} \quad (7)$$

where  $d = 1 - \sum_{j=1}^p a_j \eta_j^{\alpha-\beta-1}$ .

Proof. By using Lemma 2.2, the solution of the equation  $D_{0+}^\alpha u(t) + y(t) = 0$  is

$$u(t) = -\frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} y(s) ds + c_1 t^{\alpha-1} + c_2 t^{\alpha-2} + \dots + c_n t^{\alpha-n},$$

where  $c_1, c_2, \dots, c_n$  are arbitrary real constants.

From the boundary condition in (1), one can  $c_2 = c_3 = \dots = c_{n-2} = c_{n-1} = c_n = 0$ .

Hence

$$u(t) = -\frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} y(s) ds + c_1 t^{\alpha-1}.$$

By the last above equation and Lemma 2.1(i), we get

$$D_{0+}^\beta u(t) = \frac{1}{\Gamma(\alpha-\beta)} \left[ c_1 \Gamma(\alpha) t^{\alpha-\beta-1} - \int_0^t (t-s)^{\alpha-\beta-1} y(s) ds \right],$$

this and by  $D_{0+}^\beta u(1) = \sum_{j=1}^p a_j D_{0+}^\beta u(\eta_j)$ , we have

$$c_1 = \frac{1}{d \Gamma(\alpha)} \left[ \int_0^1 (1-s)^{\alpha-\beta-1} y(s) ds - \sum_{j=1}^p a_j \int_0^{\eta_j} (\eta_j - s)^{\alpha-\beta-1} y(s) ds \right].$$

Then, the unique solution of the problem (1) is given by

$$\begin{aligned}
 u(t) &= \frac{t^{\alpha-1}}{d\Gamma(\alpha)} \left[ \int_0^1 (1-s)^{\alpha-\beta-1} y(s) ds - \sum_{j=1}^p a_j \int_0^{\eta_j} (\eta_j - s)^{\alpha-\beta-1} y(s) ds \right] - \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} y(s) ds, \\
 &= \frac{1}{\Gamma(\alpha)} \left[ \int_0^t [t^{\alpha-1}(1-s)^{\alpha-\beta-1} - (t-s)^{\alpha-1}] y(s) ds + \int_t^1 t^{\alpha-1}(1-s)^{\alpha-\beta-1} y(s) ds \right. \\
 &\quad \left. + \frac{1-d}{d} \int_0^1 t^{\alpha-1}(1-s)^{\alpha-\beta-1} y(s) ds - \frac{t^{\alpha-1}}{d} \sum_{j=1}^p a_j \int_0^{\eta_j} (\eta_j - s)^{\alpha-\beta-1} y(s) ds \right] \\
 &= \int_0^1 g(t,s) y(s) ds + \frac{t^{\alpha-1}}{d} \sum_{j=1}^p a_j \left[ \int_{\eta_j}^1 \eta_j^{\alpha-\beta-1} (1-s)^{\alpha-\beta-1} y(s) ds \right. \\
 &\quad \left. + \int_0^{\eta_j} [\eta_j^{\alpha-\beta-1} (1-s)^{\alpha-\beta-1} - (\eta_j - s)^{\alpha-\beta-1}] y(s) ds \right] \\
 &= \int_0^1 g(t,s) y(s) ds + \frac{t^{\alpha-1}}{d} \sum_{j=1}^p a_j h(\eta_j, s) y(s) ds \\
 &= \int_0^1 G(t,s) y(s) ds.
 \end{aligned}$$

The proof is completed. □

**Lemma 2.4.** Let  $\sum_{j=1}^p a_j \eta_j^{\alpha-\beta-1} \in [0, 1]$ ,  $\alpha \in (n-1, n]$ ,  $\beta \in [1, n-2]$ ,  $n \geq 3$ . Then, the functions  $g(t, s)$  and  $h(t, s)$  defined by (6) and (7) have the following properties:

- i. The functions  $g(t, s)$  and  $h(t, s)$  are continuous on  $[0, 1] \times [0, 1]$  and for all  $t, s \in (0, 1)$

$$g(t, s) > 0, \quad h(t, s) > 0.$$

- ii.  $g(t, s) \leq \frac{t^{\alpha-1}}{\Gamma(\alpha)}$  for all  $t, s \in [0, 1]$ .

- iii.  $g(t, s) \geq t^{\alpha-1} g(1, s)$  for all  $t, s \in [0, 1]$ , where

$$g(1, s) = \frac{1}{\Gamma(\alpha)} \left[ (1-s)^{\alpha-\beta-1} - (1-s)^{\alpha-1} \right].$$

From the above properties, we deduce the following properties:

- iv. The function  $G(t, s) \geq 0$  is continuous on  $[0, 1] \times [0, 1]$  and  $G(t, s) > 0$  for all  $t, s \in (0, 1)$ .

- v.  $\max_{t \in [0, 1]} G(t, s) = G(1, s)$ , for all  $s \in [0, 1]$ , where

$$G(1, s) = g(1, s) + \frac{1}{d} \sum_{j=1}^p a_j h(\eta_j, s) \leq \frac{(1-s)^{\alpha-\beta-1}}{d\Gamma(\alpha)}.$$

Proof. It is easy to check that (i), (v), (vi) holds. So we prove that (ii) is true. Note that (6) and  $0 \leq (1-s)^{\alpha-\beta-1} \leq 1$ . It follows that  $g(t, s) \leq \frac{t^{\alpha-1}}{\Gamma(\alpha)}$  for all  $t, s \in [0, 1]$ . It remains to prove (iii). We divide the proof into two cases and by (1), we have.

Case1. When  $0 \leq s \leq t \leq 1$ , we have

$$g(t, s) = \frac{1}{\Gamma(\alpha)} t^{\alpha-1} \left[ (1-s)^{\alpha-\beta-1} - \left(1 - \frac{s}{t}\right)^{\alpha-1} \right] \geq t^{\alpha-1} g(1, s).$$

Case2. When  $0 \leq t \leq s \leq 1$ , we have

$$g(t, s) = \frac{1}{\Gamma(\alpha)} t^{\alpha-1} (1-s)^{\alpha-\beta-1} \geq t^{\alpha-1} g(1, s).$$

Hence  $g(t, s) \geq t^{\alpha-1} g(1, s)$  for all  $t, s \in [0, 1]$ . □

## 2.2 Existence results

First, for the uniqueness results of problem (1), we need the following assumptions.

(A<sub>1</sub>)  $f(t, u_1) \leq f(t, u_2)$  for any  $0 < t < 1$ ,  $0 \leq u_1 \leq u_2$ .

(A<sub>2</sub>) For any  $r \in (0, 1)$ , there exists a constant  $q \in (0, 1)$  such that

$$f(t, ru) \geq r^q f(t, u), \quad (t, u) \in (0, 1) \times [0, \infty). \quad (8)$$

(A<sub>3</sub>)  $0 < \int_0^1 f(s, s^{\alpha-1}) ds < \infty$ .

We shall consider the Banach space  $E = C[0, 1]$  equipped with the norm  $\|u\| = \max_{0 \leq t \leq 1} |u(t)|$  and let

$$D = \{u \in C^+[0, 1] : \exists M_u \geq m_u \geq 0, m_u t^{\alpha-1} \leq u(t) \leq M_u t^{\alpha-1}, \text{ for } t \in [0, 1]\}, \quad (9)$$

where

$$C^+[0, 1] = \{u \in E : u(t) \geq 0, t \in [0, 1]\}.$$

In view of Lemma 2.3, we define an operator  $T$  as

$$(Tu)(t) = \int_0^1 G(t, s) y(s) ds, \quad (10)$$

where  $G(t, s)$  is given by (5).

By (A<sub>1</sub>) it is easy to see that the operator  $T : D \rightarrow C^+[0, 1]$  is increasing. Observe that the BVP (1) has a solution if and only if the operator  $T$  has a fixed point.

Obviously, from (A<sub>1</sub>) we obtain

$$f(t, ru) \leq r^q f(t, u), \quad \forall r > 1, q \in (0, 1), \quad (t, u) \in (0, 1) \times [0, \infty).$$

In what follows, we first prove  $T : D \rightarrow D$ . In fact, for any  $u \in D$ , there exist a positive constants  $0 < m_u < 1 < M_u$  such that

$$m_u s^{\alpha-1} \leq u(s) \leq M_u s^{\alpha-1}, \quad s \in [0, 1].$$

Then, from  $(A_1)$ ,  $f(t, u)$  non-decreasing respect to  $u$  and  $(A_2)$ , we can imply that for  $s \in (0, 1)$ ,  $q \in (0, 1)$

$$(m_u)^q f(s, s^{\alpha-1}) \leq f(s, u(s)) \leq (M_u)^q f(s, s^{\alpha-1}), \quad s \in (0, 1). \quad (11)$$

From (11) and Lemma 2.4, we obtain

$$\begin{aligned} Tu(t) &= \int_0^1 g(t, s) f(s, u(s)) ds + \frac{t^{\alpha-1}}{d} \sum_{j=1}^p a_j \int_0^1 h(\eta_j, s) f(s, u(s)) ds, \\ &\leq t^{\alpha-1} \left[ \frac{1}{\Gamma(\alpha)} \int_0^1 f(s, u(s)) ds + \frac{1}{d} \sum_{j=1}^p a_j \int_0^1 h(\eta_j, s) f(s, u(s)) ds \right], \\ &\leq t^{\alpha-1} \left[ \frac{(M_u)^q}{\Gamma(\alpha)} \int_0^1 f(s, s^{\alpha-1}) ds + \frac{(M_u)^q}{d} \sum_{j=1}^p a_j \int_0^1 h(\eta_j, s) f(s, s^{\alpha-1}) ds \right], \quad t \in [0, 1], \end{aligned} \quad (12)$$

and

$$\begin{aligned} Tu(t) &= \int_0^1 g(t, s) f(s, u(s)) ds + \frac{t^{\alpha-1}}{d} \sum_{j=1}^p a_j \int_0^1 h(\eta_j, s) f(s, u(s)) ds, \\ &\geq t^{\alpha-1} \left[ \frac{1}{\Gamma(\alpha)} \int_0^1 g(1, s) f(s, u(s)) ds + \frac{1}{d} \sum_{j=1}^p a_j \int_0^1 h(\eta_j, s) f(s, u(s)) ds \right], \\ &\geq t^{\alpha-1} \left[ \frac{(m_u)^q}{\Gamma(\alpha)} \int_0^1 g(1, s) f(s, s^{\alpha-1}) ds + \frac{(m_u)^q}{d} \sum_{j=1}^p a_j \int_0^1 h(\eta_j, s) f(s, s^{\alpha-1}) ds \right], \quad t \in [0, 1]. \end{aligned} \quad (13)$$

Eqs. (12) and (13) and assumption  $(A_3)$  imply that  $T : D \rightarrow D$ .

Now, we are in the position to give the first main result of this chapter.

**Theorem 1.1** Suppose  $(A_1) - (A_3)$  hold. Then problem (1) has a unique, nondecreasing solution  $u^* \in D$ , moreover, constructing successively the sequence of functions

$$h_n(t) = \int_0^1 G(t, s) f(s, h_{n-1}(s)) ds, \quad t \in [0, 1], \quad n = 1, 2, \dots, \quad (14)$$

for any initial function  $h_0(t) \in D$ , then  $\{h_n(t)\}$  must converge to  $u^*(t)$  uniformly on  $[0, 1]$  and the rate of convergence is

$$\max_{t \in [0, 1]} |h_n(t) - u^*(t)| = O(1 - \theta^n), \quad (15)$$

where  $0 < \theta < 1$ , which depends on the initial function  $h_0(t)$ .

Proof. For any  $h_0 \in D$ , we let

$$l_{h_0} = \sup\{l > 0 : lh_0(t) \leq (Th_0)(t), t \in [0, 1]\}, \quad (16)$$

$$L_{h_0} = \inf\{L > 0 : Lh_0(t) \geq (Th_0)(t), t \in [0, 1]\}, \quad (17)$$

$$m = \min\left\{1, (l_{h_0})^{\frac{1}{1-q}}\right\}, \quad M = \max\left\{1, (L_{h_0})^{\frac{1}{1-q}}\right\}, \quad (18)$$

and

$$u_0(t) = mh_0(t), \quad v_0(t) = Mh_0(t), \quad (19)$$

$$u_n(t) = Tu_{n-1}(t), \quad v_n(t) = Tv_{n-1}(t), \quad n = 0, 1, \dots, \quad (20)$$

Since the operator  $T$  is increasing,  $(A_1)$ ,  $(A_2)$  and (16)–(20) imply that there exist iterative sequences  $\{u_n\}$ ,  $\{v_n\}$  satisfying

$$u_0(t) \leq u_1(t) \leq \dots \leq u_n(t) \leq \dots \leq v_n(t) \leq \dots \leq v_1(t) \leq v_0(t), \quad t \in [0, 1]. \quad (21)$$

In fact, from (19) and (20), we have

$$u_0(t) \leq v_0(t), \quad (22)$$

$$\begin{aligned} u_1(t) &= Tu_0(t) = \int_0^1 G_1(t, s) f(s, mh_0(s)) ds + \frac{t^{\alpha-1}}{d} \sum_{i=1}^n a_j \int_0^1 G_2(\eta_j, s) f(s, mh_0(s)) ds, \\ &\geq m^q \left[ \int_0^1 G_1(t, s) f(s, h_0(s)) ds + \frac{t^{\alpha-1}}{d} \sum_{i=1}^n a_j \int_0^1 G_2(\eta_j, s) f(s, h_0(s)) ds \right], \\ &\geq m^q Th_0(t) \geq mh_0(t) = u_0(t), \end{aligned} \quad (23)$$

and

$$\begin{aligned} v_1(t) &= Tv_0(t) = \int_0^1 G_1(t, s) f(s, Mh_0(s)) ds + \frac{t^{\alpha-1}}{d} \sum_{i=1}^n a_j \int_0^1 G_2(\eta_j, s) f(s, Mh_0(s)) ds, \\ &\leq M^q \left[ \int_0^1 G_1(t, s) f(s, h_0(s)) ds + \frac{t^{\alpha-1}}{d} \sum_{i=1}^n a_j \int_0^1 G_2(\eta_j, s) f(s, h_0(s)) ds \right] \\ &\leq M^q Th_0(t) \leq Mh_0(t) = v_0(t). \end{aligned} \quad (24)$$

Then, by (22)–(24) and induction, the iterative sequences  $\{u_n\}$ ,  $\{v_n\}$  satisfy

$$u_0(t) \leq u_1(t) \leq \dots \leq u_n(t) \leq \dots \leq v_n(t) \leq \dots \leq v_1(t) \leq v_0(t), \quad \forall t \in [0, 1].$$

Note that  $u_0(t) = \frac{m}{M} v_0(t)$ , from  $(A_1)$ , (10), (19) and (20), it can be obtained by induction that

$$u_n(t) \geq \theta^{qn} v_n(t), \quad t \in [0, 1], \quad n = 0, 1, 2, \dots, \quad (25)$$

where  $\theta = \frac{m}{M}$ .

From (21) and (25) we know that

$$0 \leq u_{n+p}(t) - u_n(t) \leq v_n(t) - u_n(t) \leq (1 - \theta^{qn})Mh_0(t), \quad \forall n, p \in \mathbb{N}, \quad (26)$$

and since  $(1 - \theta^{qn})Mh_0(t) \rightarrow 0$ , as  $n \rightarrow \infty$ , this yields that there exists  $u^* \in D$  such that

$$u_n(t) \rightarrow u^*(t), \quad (\text{uniformly on } [0, 1]).$$

Moreover, from (26) and

$$\begin{aligned} 0 \leq v_n(t) - u^*(t) &= v_n(t) - u_n(t) + u_n(t) - u^*(t), \\ &\leq (1 - \theta^{qn})Mh_0(t) \rightarrow 0, \quad \text{as } n \rightarrow \infty, \end{aligned}$$

we have

$$v_n(t) \rightarrow u^*(t), \quad (\text{uniformly on } [0, 1]),$$

so,

$$u_n(t) \rightarrow u^*(t), \quad v_n(t) \rightarrow u^*(t), \quad (\text{uniformly on } [0, 1]). \quad (27)$$

Therefore

$$u_n(t) \leq u^*(t) \leq v_n(t), \quad t \in [0, 1], n = 0, 1, 2, \dots, \quad (28)$$

From  $(A_1)$ , (19) and (20), we have

$$u_{n+1}(t) = Tu_n(t) \leq Tu^*(t) \leq Tv_n(t) = v_{n+1}(t), n = 0, 1, 2, \dots, .$$

This together with (27) and uniqueness of limit imply that  $u^*$  satisfy  $u^* = Tu^*$ , that is  $u^* \in D$  is a solution of BVP (1) and (2).

From (19)–(21) and  $(A_1)$ , we obtain

$$u_n(t) \leq h_n(t) \leq v_n(t), n = 0, 1, 2, \dots, . \quad (29)$$

It follows from (26)–(29) that

$$\begin{aligned} |h_n(t) - u^*(t)| &\leq |h_n(t) - u_n(t)| + |u_n(t) - u^*(t)|, \\ &\leq |h_n(t) - u_n(t)| + |u^*(t) - u_n(t)|, \\ &\leq 2|v_n(t) - u_n(t)|, \\ &\leq 2M(1 - \theta^{qn})|h_0(t)|. \end{aligned}$$

Therefore

$$\max_{t \in [0, 1]} |h_n(t) - u^*(t)| \leq 2M(1 - \theta^{qn}) \max_{t \in [0, 1]} |h_0(t)|.$$

Hence, (15) holds. Since  $h_0(t)$  is arbitrary in  $D$  we know that  $u^*(t)$  is the unique solution of the boundary value problem (1) in  $D$ .  $\square$

We construct an example to illustrate the applicability of the result presented.

Example 2.1. Consider the following boundary value problem

$$\begin{cases} D_{0^+}^{\frac{5}{2}} u(t) + \frac{(u)^{\frac{2}{3} - \frac{1}{6} \cos(t)}}{\sqrt{t}} = 0, & t \in (0, 1), \\ u(0) = u'(0) = 0, \quad u'(1) = \frac{\sqrt{2}}{2} u'\left(\frac{1}{2}\right), \end{cases} \quad (30)$$

where  $\alpha = \frac{5}{2}, \beta = 1, a_1 = \frac{\sqrt{2}}{2}, \eta_1 = \frac{1}{2}$  and  $f(t, u) = \frac{(u)^{\frac{2}{3} - \frac{1}{6} \cos(t)}}{\sqrt{t}}$  is increasing function with respect to  $u$  for all  $t \in (0, 1)$ , so, assumption  $(A_1)$  satisfied.

By simple calculation we have  $d = 1 - \frac{\sqrt{2}}{2} \left( \sqrt{\frac{1}{2}} \right) = \frac{1}{2}$ .

For any  $r \in (0, 1)$ , there exists  $q = \frac{1}{2} \in (0, 1)$  such that

$$f(t, ru) = \frac{(ru)^{\frac{2}{3} - \frac{1}{6} \cos(t)}}{\sqrt{t}} \geq r^{\frac{1}{2}} \frac{(u)^{\frac{2}{3} - \frac{1}{6} \cos(t)}}{\sqrt{t}} = r^{\frac{1}{2}} f(t, u),$$

thus,  $f(t, u)$  satisfies  $(A_2)$  and is singular at  $t = 0$ .

On the other hand,

$$\int_0^1 f(t, t^{2,5-1}) dt \leq \int_0^1 t^{\frac{1}{4}} dt = \frac{4}{5} < \infty,$$

so, assumption  $(A_3)$  is satisfied.

Hence, all the assumptions of Theorem 1.1 are satisfied. Which implies that the boundary value (30) has an unique, nondecreasing solution  $u^* \in D$ .

### 3. Existence result for inclusion problem (2)

We provide another result about the existence of solutions for the problem (2) by using the assumption of nonconvex compact values for multifunction. Our strategy to deal with this problem is based on the Covitz-Nadler theorem for the contraction multivalued maps [44] for lower semi-continuous maps with decomposable values.

First, we will present notations, definitions and preliminary facts from multivalued analysis which are used throughout this chapter. For more details on the multivalued maps, see the book of Aubin and Cellina [48], Demling [49], Gorniewicz [50] and Hu and Papageorgiou [51], see also [44, 48, 49, 52–54].

Here  $(C[0, 1], \mathbb{R})$  denotes the Banach space of all continuous functions from  $[0, 1]$  into  $\mathbb{R}$  with the norm  $\|u\| = \sup\{|u(t)| : \text{for all } t \in [0, 1]\}$ ,  $L^1([0, 1], \mathbb{R})$ , the Banach space of measurable functions  $u : [0, 1] \rightarrow \mathbb{R}$  which are Lebesgue integrable, normed by  $\|u\|_{L^1} = \int_0^1 |u(t)| dt$ .

Let  $(X, d)$  be a metric space induced from the normed space  $(X, \|\cdot\|)$ . We denote

$$\begin{aligned} P_0(X) &= \{A \in P(X) : A \neq \emptyset\}, \\ P_b(X) &= \{A \in P_0(X) : A \text{ is bounded}\}, \\ P_{cl}(X) &= \{A \in P_0(X) : A \text{ is closed}\}, \\ P_{cp}(X) &= \{A \in P_0(X) : A \text{ is compact}\}, \\ P_{b,cl}(X) &= \{A \in P_0(X) : A \text{ is closed and bounded}\}, \end{aligned}$$

where  $P(X)$  is the family of all subsets of  $X$ .

Definition 3.1. A multivalued map  $G : X \rightarrow P(X)$ .

1.  $G(u)$  is convex (closed) valued if  $G(u)$  is convex (closed) for all  $u \in X$ ,

2. is bounded on bounded sets if  $G(B) = \bigcup_{u \in B} G(u)$  is bounded in  $X$  for all  $B \in P_b(X)$  i.e.,  $\sup_{u \in B} \{\sup\{|v|, v \in G(u)\}\} < \infty$ ,

3. has a fixed point if there is  $u \in X$  such that  $u \in G(u)$ . The fixed point set of the multivalued operator  $G$  will be denote by  $\text{Fix } G$ .

Definition 3.2. A multivalued map  $G : [0, 1] \rightarrow P_{cl}(\mathbb{R})$  is said to be measurable if for every  $y \in \mathbb{R}$  the function

$$t \mapsto d(y, G(t)) = \inf \{ \|y - z\| : z \in G(t) \},$$

is measurable.

Definition 3.3. Let  $Y$  be a nonempty closed subset of a Banach space  $E$  and  $G : Y \rightarrow P_{cl}(E)$  be a multivalued operator with nonempty closed values.

- i.  $G$  is said to be lower semi-continuous (l.s.c) if the set  $\{x \in X : G(x) \cap U \neq \phi\}$  is open for any open set  $U$  in  $E$ .
- ii.  $G$  has a fixed point if there is  $x \in Y$  such that  $x \in G(x)$ .

For each  $u \in (C[0, 1], \mathbb{R})$ , define the set of selection of  $F$  by

$$S_{F,u} = \{v \in AC([0, 1], \mathbb{R}) : v \in F(t, u(t)), \text{ for almost all } t \in [0, 1]\}.$$

For  $P(X) = 2^X$ , consider the Pompeiu-Hausdorff metric (see [55]).

$H_d : 2^X \times 2^X \rightarrow [0, \infty)$  given by

$$H_d(A, B) = \max \left\{ \sup_{a \in A} d(a, B), \sup_{b \in B} d(b, A) \right\},$$

where  $d(a, B) = \inf_{b \in B} d(a, b)$  and  $d(b, A) = \inf_{a \in A} d(a, b)$ . Then  $(P_{b,cl}(X), H_d)$  is a metric space and  $(P_{cl}(X), H_d)$  is a generalized metric space see [8].

Definition 3.4. Let  $A$  be a subset of  $[0, 1] \times \mathbb{R}$ .  $A$  is  $L \otimes B$  measurable if  $A$  belongs to the  $\sigma$ -algebra generated by all sets of the  $J \times D$ , where  $J$  is Lebesgue measurable in  $[0, 1]$  and  $D$  is Borel measurable in  $\mathbb{R}$ .

Definition 3.5. A subset  $A$  of  $L^1([0, 1], \mathbb{R})$  is decomposable if all  $u, v \in A$  and measurable  $J \subset [0, 1] = j$ , the function  $u\chi_J + v\chi_{j^c} \in A$ , where  $\chi_J$  stands for the characteristic function of  $J$ .

Definition 3.6. Let  $Y$  be a separable metric space and  $N : Y \rightarrow P(L^1([0, 1], \mathbb{R}))$  be a multivalued operator. We say  $N$  has property (BC) if  $N$  is lower semi-continuous (l.s.c) and has nonempty closed and decomposable values.

Let  $F : [0, 1] \times \mathbb{R} \rightarrow P(\mathbb{R})$  be a multivalued map with nonempty compact values. Define a multivalued operator

$$\Phi : C([0, 1], \mathbb{R}) \rightarrow P(L^1([0, 1], \mathbb{R})),$$

by letting

$$\Phi(u) = \{w \in L^1([0, 1], \mathbb{R}) : w(t) \in F(t, u(t)) \text{ for a.e. } t \in [0, 1]\}.$$

Definition 3.7. The operator  $\Phi$  is called the Niemytzki operator associated with  $F$ . We say  $F$  is of the lower semi-continuous type (l.s.c type) if its associated Niemytzki operator  $\Phi$  has (BC) property.

Definition 3.8. A multivalued operator  $N : X \rightarrow P_{cl}(X)$  is called.

i.  $\rho$ -Lipschitz if and only if there exists  $\rho > 0$  such that  $H_d(N(u), N(v)) \leq \rho d(u, v)$  for each  $u, v \in X$ ,

ii. a contraction if and only if it is  $\rho$ -Lipschitz with  $\rho < 1$ .

Lemma 3.1. ([44] Covitz-Nadler). Let  $(X, d)$  be a complete metric space. If  $N : X \rightarrow P_{cl}(X)$  is a contraction, then  $\text{Fix}N \neq \emptyset$ , where  $\text{Fix}N$  is the fixed point of the operator  $N$ .

Definition 3.9. A measurable multivalued function  $F : [0, 1] \rightarrow P(X)$  is said to be integrably bounded if there exists a function  $g \in L^1([0, 1], X)$  such that, for all  $v \in F(t)$ ,  $\|v\| \leq g(t)$  for a.e.  $t \in [0, 1]$ .

Let us introduce the following hypotheses.

(A<sub>4</sub>)  $F : [0, 1] \times \mathbb{R} \rightarrow P_{cp}(\mathbb{R})$  be a multivalued map verifying.

i.  $(t, u) \mapsto F(t, u)$  is  $L \otimes B$  measurable.

ii.  $u \mapsto F(t, u)$  is lower semi-continuous for a.e.  $t \in [0, 1]$ .

(A<sub>5</sub>)  $F$  is integrably bounded, that is, there exists a function  $m \in L^1([0, 1], \mathbb{R}^+)$  such that  $\|F(t, u)\| = \sup\{|v| : v \in F(t, u)\} \leq m(t)$  for almost all  $t \in [0, 1]$ .

Lemma 3.2. [56] Let  $F : [0, 1] \times \mathbb{R} \rightarrow P_{cp}(\mathbb{R})$  be a multivalued map. Assume (A<sub>4</sub>) and (A<sub>5</sub>) hold. Then  $F$  is of the l.s.c. type.

Definition 3.10. A function  $u \in AC^2([0, 1], \mathbb{R})$  is called a solution to the boundary value problem (2) if  $u$  satisfies the differential inclusion in (2) a.e. on  $[0, 1]$  and the conditions in (2).

Finally, we state and prove the second main result of this Chapter. We prove the existence of solutions for the inclusion problem (2) with a nonconvex valued right hand side by applying a fixed point theorem for multivalued maps due to Covitz and Nadler. For investigation of the problem (2) we shall provide an application of the Lemma 3.4 and the following Lemma.

Lemma 3.3. ([13]) A multifunction  $F : X \rightarrow C(X)$  is called a contraction whenever there exists  $\gamma \in (0, 1)$  such that  $H_d(N(u), N(v)) \leq \gamma d(u, v)$  for all  $u, v \in X$ .

Now, we present second main result of this section.

Theorem 1.2 Assume that the following hypotheses hold.

(H<sub>1</sub>)  $F : J \times \mathbb{R} \rightarrow P_{cp}(\mathbb{R})$  is an integrable bounded multifunction such that the map  $t \mapsto F(t, u)$  is measurable,

(H<sub>2</sub>)  $H_d(F(t, u_1), F(t, u_2)) \leq m(t)|u_1 - u_2|$  for almost all  $t \in J$  and  $u_1, u_2 \in \mathbb{R}$  with  $m \in L^1(J, \mathbb{R})$  and  $d(0, F(t, 0)) \leq m(t)$  for almost all  $t \in J$ . Then the problem (2) has a solution provided that

$$l = \int_0^1 G(1, s)m(s)ds < 1.$$

Proof. We transform problem (2) into a fixed point problem. Consider the operator  $N : C[0, 1] \rightarrow P(C[0, 1], \mathbb{R})$  defined by

$$N(u) = \left\{ h \in X, \exists y \in S_{F,u} \setminus h(t) = \int_0^1 G(t,s)y(s)ds, t \in J \right\}, \quad (31)$$

where  $G(t,s)$  defined by (5). It is clear that fixed points of  $N$  are solution of (2).

We shall prove that  $N$  fulfills the assumptions of Covitz-Nadler contraction principle.

Note that, the multivalued map  $t \mapsto F(t, u(t))$  is measurable and closed for all  $u \in AC^1([0, \infty))$  (e.g., [52] Theorem III.6). Hence, it has a measurable selection and so the set  $S_{F,u}$  is nonempty, so,  $N(u)$  is nonempty for any  $u \in C([0, \infty))$ .

First, we show that  $N(u)$  is a closed subset of  $X$  for all  $u \in AC^1([0, \infty), \mathbb{R})$ . Let  $u \in X$  and  $\{u_n\}_{n \geq 1}$  be a sequence in  $N(u)$  with  $u_n \rightarrow u$ , as  $n \rightarrow \infty$  in  $u \in C([0, \infty))$ . For each  $n$ , choose  $y_n \in S_{F,u}$  such that

$$u_n(t) = \int_0^1 G(t,s)y_n(s)ds.$$

Since  $F$  has compact values, we may pass onto a subsequence (if necessary) to obtain that  $y_n$  converges to  $y \in L^1([0, 1], \mathbb{R})$  in  $L^1([0, 1], \mathbb{R})$ . In particular,  $y \in S_{F,u}$  and for any  $t \in [0, 1]$ , we have

$$u_n(t) \rightarrow u(t) = \int_0^1 G(t,s)y(s)ds,$$

i.e.,  $u \in N(u)$  and  $N(u)$  is closed.

Next, we show that  $N$  is a contractive multifunction with constant  $l < 1$ . Let  $u, v \in C([0, 1], \mathbb{R})$  and  $h_1 \in N(u)$ . Then there exist  $y_1 \in S_{F,u}$  such that

$$h_1(t) = \int_0^1 G(t,s)y_1(s)ds, \quad t \in J.$$

By  $(H_2)$ , we have

$$H_d(F(t, u(t)), F(t, v(t))) \leq m(t)(|u(t) - v(t)|),$$

for almost all  $t \in J$ .

So, there exists  $w \in S_{F,v}$  such that

$$|y_1(t) - w| \leq m(t)(|u(t) - v(t)|),$$

for almost all  $t \in J$ .

Define the multifunction  $U : J \rightarrow P(\mathbb{R})$  by

$$U(t) = \{w \in \mathbb{R} : |y_1(t) - w| \leq m(t)(|u(t) - v(t)|) \text{ for almost all } t \in J\}.$$

It is easy to check that the multifunction  $V(\cdot) = U(\cdot) \cap F(\cdot, v(\cdot))$  is measurable (e.g., [52] Theorem III.4).

Thus, there exists a function  $y_2(t)$  which is measurable selection for  $V$ . So,  $y_2 \in S_{F,v}$  and for each  $t \in J$ , we have

$$|y_1(t) - y_2(t)| \leq m(t)(|u(t) - v(t)|).$$

Now, consider  $h_2 \in N(u)$  which is defined by

$$h_2(t) = \int_0^1 G(t,s)y_2(s)ds, \quad t \in J,$$

and one can obtain

$$\begin{aligned} |h_1(t) - h_2(t)| &\leq \int_0^1 G(t,s)|y_1(s) - y_2(s)|ds \\ &\leq \int_0^1 G(1,s)m(s)|u(s) - v(s)|ds. \end{aligned}$$

Hence

$$\|h_1(t) - h_2(t)\| \leq \|p\|_\infty \|u - v\| \left[ \int_0^1 G(1,s)m(s)ds \right].$$

Analogously, interchanging the roles of  $u$  and  $v$ , we obtain

$$H_d(N(u), N(v)) \leq \|u - v\| \left[ \int_0^1 G(1,s)m(s)ds \right].$$

Since  $N$  is a contraction, it follows by Lemma 3.1 (by using the result of Covitz and Nadler) that  $N$  has a fixed point which is a solution to problem (2).  $\square$

We construct an example to illustrate the applicability of the result presented.

**Example 3.1.** Consider the problem

$$-D^\alpha u(t) \in F(t, u(t)), \quad t \in [0, 1], \quad (32)$$

subject to the three-point boundary conditions

$$u^{(i)}(0) = 0, i \in \{0, 1\}, \quad D_{0+}^\beta u(1) = \sum_{j=1}^2 a_j D_{0+}^\beta u(\eta_j), \quad (33)$$

where  $\alpha = \frac{5}{2}, \beta = 1, a_1 = \frac{1}{2}, a_2 = \frac{3}{2}, \eta_1 = \frac{1}{16}, \eta_2 = \frac{5}{16}$  and  $F(t, u(t)) : [0, 1] \times \mathbb{R} \rightarrow 2^{\mathbb{R}}$  multivalued map given by

$$u \mapsto F(t, u) = \left( 0, \frac{t|u|}{2(1+|u|)} \right), u \in \mathbb{R},$$

verifying  $(H_1)$ .

Obviously,

$$\sup\{|f| : f \in F(t, u)\} \leq \frac{t+1}{2},$$

we have

$$H_d(F(t, u), F(t, v)) \leq \left(\frac{t+1}{2}\right) |u - v|, u, v \in \mathbb{R}, t \in [0, 1],$$

which shows that  $(H_2)$  holds

So, if  $m(t) = \frac{t+1}{2}$  for all  $t \in [0, 1]$ , then

$$H_d(F(t, u), F(t, v)) \leq m(t) |u - v|.$$

It can be easily found that  $d = 1 - \frac{1}{2} \left(\frac{1}{16}\right)^{\frac{5}{2}} - \frac{3}{2} \left(\frac{5}{16}\right)^{\frac{5}{2}} = 0,9176244637$ .

Finally,

$$l = \int_0^1 G(1, s) m(s) ds = 0,4636273746 < 1.$$

Hence, all assumptions and conditions of Theorem 1.2 are satisfied. So, Theorem 1.2 implies that the inclusion problem (32) and (33) has at least one solution.

## 4. Conclusions

This chapter concerns the boundary value problem of a class of fractional differential equations involving the Riemann-Liouville fractional derivative with nonlocal boundary conditions. By using the properties of the Green's function and the monotone iteration technique, one shows the existence of positive solutions and constructs two successively iterative sequences to approximate the solutions. In the multi-valued case, an existence result is proved by using fixed point theorem for contraction multivalued maps due to Covitz and Nadler. The results of the present chapter are significantly contribute to the existing literature on the topic.

## Acknowledgements

The authors want to thank the anonymous referee for the thorough reading of the manuscript and several suggestions that help us improve the presentation of the chapter.

## Conflict of interest

The authors declare no conflict of interest.

IntechOpen

## Author details

Noureddine Bouteraa<sup>1,2\*†</sup> and Habib Djourdem<sup>1,3\*†</sup>

1 Laboratory of Fundamental and Applied Mathematics of Oran (LMFAO),  
University of Oran1, Ahmed Benbella, Algeria

2 Oran Graduate School of Economics. Algeria

3 University of Ahmed Zabbana, Relizane, Algeria

\*Address all correspondence to: [bouteraa-27@hotmail.fr](mailto:bouteraa-27@hotmail.fr);  
[djourdem.habib7@gmail.com](mailto:djourdem.habib7@gmail.com)

† These authors contributed equally.

## IntechOpen

---

© 2021 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

## References

- [1] R. P. AGARWAL, D. BALEANU, V. HEDAYATI and S. H. REZAPOUR, *Two fractional derivative inclusion problems via integral boundary condition*, Appl. Math. Comput. **vol.257** (2015), 205-212.
- [2] V. KAC and P. CHEUNG, *Quantum Calculus*, Springer, New-York, 2002.
- [3] V. Lakshmikantham and A. S. Vatsala, General uniqueness and monotone iterative technique for fractional differential equations. Appl. Math. Lett. 21 (8)(2008), 828-834.
- [4] S. Miller and B. Ross, An introduction to the fractional calculus and fractional differential equations, John Wiley and Sons, Inc. New-York, (1993).
- [5] W. Rudin, Functional Analysis, 2nd edn. International Series in Pure and Applied Mathematics. Mc Graw-Hill, New York (1991).
- [6] S. G. Samko, A. A. Kilbas and O. I. Marichev, Fractional Integrals and Derivatives: Theory and Applications. Gordon & Breach, Yverdon (1993).
- [7] F. Jarad, T. Abdeljaw and D. Baleanu, On the generalized fractional derivatiives and their Caputo modification. J. Nonl. Sci. Appl. 10 (5) (2017), 2607-2619.
- [8] Y. Tian, Positive solutions to m-point boundary value problem of fractional differential equation. Acta Math. Appl. Sinica (Engl. Ser.) 29 (2013), 661-672.
- [9] A.M.A. El-Sayed and A.G. Ibrahim, Multivalued Fractional differential equations of arbitrary orders. Springer-Verlag, Appl. Math. Comput. 68 (1995), 15-25.
- [10] N. Bouteraa and S. Benaicha, Existence of solutions for nonlocal boundary value problem for Caputo nonlinear fractional differential inclusion, Journal of Mathematical Sciences and Modelling, 1 (1) (2018), 45-55.
- [11] N. Bouteraa and S. Benaicha, Existence results for fractional differential inclusion with nonlocal boundary conditions, Ri. Mat. Parma, Vol.11(2020),181-206.
- [12] A. CERNIA, *Existence of solutions for a certain boundary value problem associated to a fourth order differential inclusion*, Inter. Jour. Anal. Appl. **vol.14** (2017), 27-33.
- [13] S. K. Ntouyas, S. Etemad, J. Tariboon and W. Sutsutad, Boundary value problems for Riemann-Liouville nonlinear fractional differential inclusions with nonlocal Hadamard fractional integral conditions, Meditter. J. Math. 2015 (2015), 16 pages.
- [14] N. Bouteraa and S. Benaicha, Triple positive solutions of higher-order nonlinear boundary value problems, Journal of Computer Science and Computational Mathematics, Volume 7, Issue 2, June 2017, 25-31.
- [15] N. Bouteraa and S. Benaicha, Existence of solutions for three-point boundary value problem for nonlinear fractional equations, Analele Universitatii Oradia Fasc. Mathematica, Tom XXIV (2017), Issue No. 2, 109-119.
- [16] S. Benaicha and N. Bouteraa, Existence of solutions for three-point boundary value problem for nonlinear fractional differential equations, Bulltin of the Transilvania University of Brasov, Serie III: Mathematics, Informtics, Physics. Volume 10 (59), No. 2-2017.
- [17] N. Bouteraa and S. Benaicha, Existence of solutions for third-order three-point boundary value problem, Mathematica. 60 (83), N 0 1, 2018, pp. 21-31.

- [18] N. Bouteraa and S. Benaicha, The uniqueness of positive solution for higher-order nonlinear fractional differential equation with nonlocal boundary conditions, *Advances in the Theory of Nonlinear and its Application*, 2(2018) No 2, 74-84.
- [19] N. Bouteraa, S. Benaicha and H. Djourdem, Positive solutions for nonlinear fractional differential equation with nonlocal boundary conditions, *Universal Journal of Mathematics and Applications*, 1 (1) (2018), 39-45.
- [20] N. Bouteraa and S. Benaicha, The uniqueness of positive solution for nonlinear fractional differential equation with nonlocal boundary conditions, *Analele universitatii Oradia. Fasc. Matematica. Tom XXV (2018), Issue No. 2*, 53-65.
- [21] N. Bouteraa, S. Benaicha, H. Djourdem and N. Benatia, Positive solutions of nonlinear fourth-order two-point boundary value problem with a parameter, *Romanian Journal of Mathematics and Computer Science*, 2018, Volume 8, Issue 1, p.17-30.
- [22] N. Bouteraa and S. Benaicha, Positive periodic solutions for a class of fourth-order nonlinear differential equations, *Siberian Journal of Numerical Mathematics*, Volume 22, No. 1 (2019), 1-14.
- [23] N. Bouteraa, Existence of solutions for some nonlinear boundary value problems, [thesis]. University of Oran1, Ahmed Benbella, Algeria; 2018.
- [24] N. Bouteraa, S. Benaicha and H. Djourdem, ON the existence and multiplicity of positive radial solutions for nonlinear elliptic equation on bounded annular domains via fixed point index, *Maltepe Journal of Mathematics*, Volume I, Issue 1, (2019), 30-47.
- [25] N. Bouteraa, S. Benaicha and H. Djourdem, Positive solutions for systems of fourth-order two-point boundary value problems with parameter, *Journal of Mathematical Sciences and Modeling*, 2(1) (2019), 30-38.
- [26] N. Bouteraa and S. Benaicha, Existence and multiplicity of positive radial solutions to the Dirichlet problem for the nonlinear elliptic equations on annular domains, *Stud. Univ. Babeş-Bolyai Math*, 65(2020), No. 1, 109-125.
- [27] S. Benaicha, N. Bouteraa and H. Djourdem, Triple positive solutions for a class of boundary value problems with integral boundary conditions, *Bulletin of Transilvania University of Brasov, Series III : Mathematics, Informatics, Physics*, Vol. 13 (62), No. 1 (2020), 51-68.
- [28] N. Bouteraa, H. Djourdem and S. Benaicha, Existence of solution for a system of coupled fractional boundary value problem, *Proceedings of International Mathematical Sciences*, Vol. II, Issue 1 (2020), 48-59.
- [29] N. Bouteraa and S. Benaicha, Existence results for second-order nonlinear differential inclusion with nonlocal boundary conditions, *Numerical Analysis and Applications*, 2021, Vol. 14, No. 1, pp. 3039.
- [30] N. Bouteraa M. In, M. A. Akinlar, B. Almohsen, Mild solutions of fractional PDE with noise, *Math. Meth. Appl. Sci.* 2021;115.
- [31] N. Bouteraa, S. Benaicha, Existence Results for Second-Order Nonlinear Differential Inclusion with Nonlocal Boundary Conditions, *Numerical Analysis and Applications*, 2021, Vol. 14, No. 1, pp. 3039.
- [32] N. Bouteraa, S. Benaicha, A study of existence and multiplicity of positive solutions for nonlinear fractional

differential equations with nonlocal boundary conditions, *Stud. Univ. Babeş-Bolyai Math.* 66(2021), No. 2, 361-380.

[33] H. Djourdem, S. Benaicha and N. Bouteraa, Existence and iteration of monotone positive solution for a fourth-order nonlinear boundary value problem, *Fundamental Journal of Mathematics and Applications*, 1 (2) (2018), 205-211.

[34] H. Djourdem, S. Benaicha, and N. Bouteraa, Two Positive Solutions for a Fourth-Order Three-Point BVP with Sign-Changing Greens Function, *Communications in Advanced Mathematical Sciences*. Vol. II, No. 1 (2019), 60-68.

[35] H. Djourdem and N. Bouteraa, Mild solution for a stochastic partial differential equation with noise, *WSEAS Transactions on Systems*, Vol. 19, (2020), 246-256.

[36] R. GHORBANIAN, V. HEDAYATI M. POSTOLACHE, and S. H. REZAPOUR, *On a fractional differential inclusion via a new integral boundary condition*, *J. Inequal. Appl.* (2014), 20 pages.

[37] M. Inc, N. Bouteraa, M. A. Akinlar, Y. M. Chu, G. W. Weber and B. Almohsen, New positive solutions of nonlinear elliptic PDEs, *Applied Sciences*, 2020, 10, 4863; doi :10.3390/app10144863, 17 pages.

[38] N. Bouteraa and S. Benaicha, Nonlinear boundary value problems for higher-order ordinary differential equation at resonance, *Romanian Journal of Mathematic and Computer Science*. 2018. Vol 8, Issue 2 (2018), p. 83-91.

[39] N. Bouteraa, S. Benaicha, A class of third-order boundary value problem with integral condition at resonance, *Maltepe Journal of Mathematics*, Volume II, Issue 2,(2020), 43-54.

[40] X. Lin, Z. Zhao and Y. Guan, Iterative Technology in a Singular Fractional Boundary Value Problem With q-Difference, *Appl. Math.* 7 (2016), 91-97.

[41] M. Li, JP. Sun and YH. Zhao, Existence of positive solution for BVP of nonlinear fractional differential equation with integral boundary conditions. *Adv Differ Equ* 2020, 177 (2020). <https://doi.org/10.1186/s13662-020-02618-9>

[42] S. H. Rezapour and V. Hedayati, On a Caputo fractional differential inclusion with integral boundary condition for convex-compact and nonconvex-compact valued multifunctions, *Kragujevac Journal of Mathematics*. **vol.41** (2017), 143-158.

[43] B. AHMAD, SK. NTOUYAS and A. ALSAEDI, Coupled systems of fractional differential inclusions with coupled boundary conditions, *Electronic Journal of Differential Equations*, Vol. 2019 (2019), No. 69, pp. 121.

[44] H. COVITZ and S. B. JR. NADLER, *Multivalued contraction mappings in generalized metric spaces*, *Israel J. Math.* **vol.8** (1970), 5-11.

[45] M. H. ANNABY, and Z. S. Mansour, *q-Fractional calculus and equations*, *Lecture Notes in Mathematics*. **vol.2056**, Springer-Verlag, Berlin 2012.

[46] O. AGRAWAL, *Some generalized fractional calculus operators and their applications in integral equations*, *Fract. Cal. Appl. Anal.* **Vol.15** (2012), 700-711.

[47] A. A. KILBAS, H. M. SRIVASTAVA and J. J. TRIJULL, *Theory and applications of fractional differential equations*, Elsevier Science B. V, Amsterdam, 2006.

[48] J. P. AUBIN and A. CELLINA, *Differential Inclusions*, Springer-Verlag, 2012.

[49] K. DEMLING, *Multivalued Differential equations*, Walter De Gryter, Berlin-New-York 1982.

[50] L. GORNIEWICZ, *Topological Fixed Point Theory of Multivalued Mappings, Mathematics and Its Applications*, Vol. 495, Kluwer Academic Publishers, Dordrecht 1999.

[51] S. HU and N. PAPAGEORGIOU, *Handbook of Multivalued Analysis, vol.I, Theory*, Kluwer Academic. J. Diff. Equ. No.147, (2013), 1-11.

[52] C. CASTAING and M. VALADIER, *Convex analysis and measurable multifunctions, Lecture Notes in Mathematics*, Springer-Verlage, Berlin-Heidelberg, New-York, 580, 1977.

[53] A. LASOTA and Z. OPIAL, *An application of the Kakutani-Ky Fan theorem in the theory of ordinary differential equations*, Bull. Acad. Pol. Sci. Ser. Sci. Math. Astronom. Phy. **vol.13** (1965), 781-786.

[54] J. MUSIELAK, *Introduction to functional analysis*, PWN, Warsaw, 1976, (in Polish).

[55] V. BERINDE and M. PACURAR, *The role of the Pompeiu-Hausdorff metric in fixed point theory*, Creat. Math. Inform. **vol.22** (2013), 35-42.

[56] M. FRIGON and A. GRANAS, *Theoremes d'existence pour des inclusions differentielles sans convexite*, C. R. Acad. Sci. Paris, SerI. **vol.310** (1990), 819-822.