We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists



186,000

200M



Our authors are among the

TOP 1% most cited scientists





WEB OF SCIENCE

Selection of our books indexed in the Book Citation Index in Web of Science™ Core Collection (BKCI)

### Interested in publishing with us? Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected. For more information visit www.intechopen.com



#### Chapter

## Relaxation Dynamics of Point Vortices

### Ken Sawada and Takashi Suzuki

# Abstract

We study a model describing relaxation dynamics of point vortices, from quasistationary state to the stationary state. It takes the form of a mean field equation of Brownian point vortices derived from Chavanis, and is formulated by our previous work as a limit equation of the patch model studied by Robert-Someria. This model is subject to the micro-canonical statistic laws; conservation of energy, that of mass, and increasing of the entropy. We study the existence and nonexistence of the global-in-time solution. It is known that this profile is controlled by a bound of the negative inverse temperature. Here we prove a rigorous result for radially symmetric case. Hence  $E/M^2$  large and small imply the global-in-time and blowup in finite time of the solution, respectively. Where E and M denote the total energy and the total mass, respectively.

Keywords: point vortex, quasi-equilibrium, relaxation dynamics

#### 1. Introduction

Our purpose is to study the system

$$\omega_{t} + \nabla \cdot \omega \nabla^{\perp} \psi = \nabla \cdot (\nabla \omega + \beta \omega \nabla \psi) \quad \text{in } \Omega \times (0, T),$$

$$\frac{\partial \omega}{\partial \nu} + \beta \omega \frac{\partial \psi}{\partial \nu}\Big|_{\partial \Omega} = 0, \quad \omega|_{t=0} = \omega_{0}(x)$$
(1)
with
$$-\Delta \psi = \omega \quad \text{in } \Omega, \quad \psi|_{\partial \Omega} = 0, \quad \beta = -\frac{\int_{\Omega} \nabla \omega \cdot \nabla \psi}{\int_{\Omega} \omega |\nabla \psi|^{2}},$$
(2)

where  $\Omega \subset \mathbf{R}^2$  is a bounded domain with smooth boundary  $\partial \Omega$ ,  $\nu$  is the outer unit normal vector on  $\partial \Omega$ , and

$$\nabla = \begin{pmatrix} \frac{\partial}{\partial x_1} \\ \frac{\partial}{\partial x_2} \end{pmatrix}, \quad \nabla^{\perp} = \begin{pmatrix} \frac{\partial}{\partial x_2} \\ -\frac{\partial}{\partial x_1} \end{pmatrix}, \quad x = (x_1, x_2).$$
(3)

The unknown  $\omega = \omega(x, t) \in \mathbf{R}$  stands for a mean field limit of many point vortices,

#### IntechOpen

$$\omega(x,t)dx = \sum_{i=1}^{N} \alpha_i \delta_{x_i(t)}(dx).$$
(4)

It was derived, first, for Brownian point vortices by [1, 2], with  $\beta = \beta(t)$  standing for the inverse temperature. Then, [3, 4] reached it by the Lynden-Bell theory [5] of relaxation dynamics, that is, as a model describing the movement of the mean field of many point vortices, from quasi-stationary state to the stationary state. This model is consistent to the Onsager theory [6–12] on stationary states and also the patch model proposed by [13, 14], that is,

$$\omega(x,t) = \sum_{i=1}^{N_p} \sigma_i \mathbf{1}_{\Omega_i(t)}(x), \tag{5}$$

where  $N_p$ ,  $\sigma_i$ , and  $\Omega_i(t)$  denote the number of patches, the vorticity of the *i*-th patch, and the domain of the *i*-th patch, respectively [15–17].

This chapter is concerned on the one-sided case of

$$\omega_0 = \omega_0(x) > 0. \tag{6}$$

If this initial value is smooth, there is a unique classical solution to (1)–(4) local in time, denoted by  $\omega = \omega(x, t)$ , with the maximal existence time  $T = T_{\text{max}} \in (0, +\infty]$ . More precisely, the strong maximum principle to (1) guarantees

$$\omega = \omega(x,t) > 0 \quad \text{on } \overline{\Omega} \times [0,T). \tag{7}$$

Then, the Hopf lemma to the Poisson equation in (2) ensures

$$\left. \frac{\partial \psi}{\partial \nu} \right|_{\partial \Omega} < 0, \tag{8}$$

and hence the well-definedness of

$$-\beta = \frac{\int_{\Omega} \nabla \omega \cdot \nabla \psi}{\int_{\Omega} \omega |\nabla \psi|^2}.$$
(9)

We confirm that system (1)-(3) satisfies the requirements of isolated system of thermodynamics. First, the mass conservation is derived from (1) as

$$\frac{d}{dt}\int_{\Omega}\omega=0, \qquad (10)$$

because

$$u \cdot 
abla^{\perp} \psi \Big|_{\partial\Omega} = 0$$
(11)

holds by (2). Second, the energy conservation follows as

$$\frac{1}{2}\frac{d}{dt} \|\nabla\psi\|_{2}^{2} = (\nabla\psi, \nabla\psi_{t}) = (\omega_{t}, \psi)$$

$$= (\omega\nabla^{\perp}\psi, \nabla\psi) - (\nabla\omega + \beta\omega\nabla\psi, \nabla\psi)$$

$$= -(\nabla\omega, \nabla\psi) - \beta \int_{\Omega} \omega |\nabla\psi|^{2} = 0$$
(12)

by (1) and (2), because

$$\nabla^{\perp}\psi\cdot\nabla\psi=0, \tag{13}$$

where ( , ) denotes the  $L^2$  inner product. Third, the entropy increasing is achieved, writing (1) as

$$\omega_{t} = \nabla \cdot \omega \left( -\nabla^{\perp} \psi + \nabla (\log \omega + \beta \psi) \right), \quad \frac{\partial}{\partial \nu} (\log \omega + \beta \psi) \bigg|_{\partial \Omega} = 0.$$
(14)  
In fact, it then follows that  
$$\int_{\Omega} \omega_{t} (\log \omega + \beta \psi) = \int_{\Omega} \omega \nabla^{\perp} \psi \cdot \nabla (\log \psi + \beta \psi) - \omega |\nabla (\log \omega + \beta \psi)|^{2} dx$$
(15)

with

$$\int_{\Omega} \omega \nabla^{\perp} \psi \cdot \nabla (\log \omega + \beta \psi) = \int_{\Omega} \nabla \omega \cdot \nabla^{\perp} \psi$$

$$= \int_{\partial \Omega} \omega \nu \cdot \nabla^{\perp} \psi - \int_{\Omega} \omega \nabla \cdot (\nabla^{\perp} \psi) = 0$$
(16)

from (11) and

$$\nabla^{\perp} \cdot \nabla = \nabla \cdot \nabla^{\perp} = 0. \tag{17}$$

Since

$$\int_{\Omega} \omega_t \log \omega = \frac{d}{dt} \int_{\Omega} \omega(\log \omega - 1), \quad \int_{\Omega} \omega_t \psi = \frac{1}{2} \frac{d}{dt} \|\nabla \psi\|_2^2 = 0,$$
(18)

We thus end up with the mass conservation



and the entropy increasing

$$\frac{d}{dt} \int_{\Omega} \omega (\log \omega - 1) = -\int_{\Omega} \omega |\nabla (\log \omega + \beta \psi)|^2 \le 0.$$
(21)

Henceforth, C > 0 stands for a generic constant. In the previous work [4] we studied radially symmetric solutions and obtained a criterion for the existence of the solution global in time. Here, we refine the result as follows, where B(0, 1) denotes the unit ball.

Theorem 1 Let

$$\Omega = B(0,1), \quad \omega_0 = \omega_0(r), \quad \omega_{0r} < 0, \quad 0 < r = |x| \le 1.$$
(22)

Then there is  $C_0 > 0$  such that

$$C_0 \|\omega_0\|_2^3 \le E\underline{\omega} \Rightarrow T = +\infty, \ \|\omega(\cdot, t)\|_{\infty} \le C, \ t \ge 0,$$
(23)

where

$$\underline{\omega} = \min_{\overline{\Omega}} \omega_0 > 0.$$
 (24)

**Theorem 2** Under the assumption of (22) there is  $\delta_0 > 0$  such that

$$\frac{E}{M^2} < \delta_0 \Rightarrow T < +\infty.$$
(25)

**Remark 1** Since

$$\|\omega_0\|_2^3 = \left(\int_{\Omega} \omega_0^2\right)^{3/2} \ge \left(\underline{\omega}^{2/3} \int_{\Omega} \omega_0^{4/3}\right)^{3/2}$$

$$= \underline{\omega} \left(\int_{\Omega} \omega_0^{4/3}\right)^{3/2} \ge \underline{\omega} |\Omega|^{-1/2} \left(\int_{\Omega} \omega_0\right)^2 = \underline{\omega} |\Omega|^{-1/2} M^2$$
(26)

the assumption (23) implies

$$\frac{E}{M^2} \ge C_0 |\Omega|^{-1/2}.$$
(27)

Therefore, roughly, the conditions  $E/M^2 \gg 1$  and  $E/M^2 \ll 1$  imply  $T = +\infty$  and  $T < +\infty$ , respectively.

Remark 2 The assumption (22) implies

$$\beta = \beta(t) < 0, \quad 0 \le t < T, \tag{28}$$

and then we obtain Theorem 1. In other words, the conclusion of this theorem arises from (28), without (22).

**Remark 3** Since

$$\frac{E}{M^2} = \frac{\int_{\Omega} |\nabla \psi|^2}{\left(\int_{\Omega} \omega\right)^2}$$
(29)

$$\frac{E}{M^2} = \|\nabla c\|_2^2, \quad c = \frac{(-\Delta)^{-1}\omega_0}{\int_\Omega \omega_0} = \frac{\psi_0}{\int_{\partial\Omega} - \frac{\partial\psi_0}{\partial\nu}},\tag{30}$$

where  $\psi_0 = (-\Delta)^{-1} \omega_0$ .

The system (1)-(4) thus obays a profile of the micro-canonical ensemble. In a system associated with the canonical ensemble, the inverse temperature  $\beta$  is a constant in (1) independent of *t*, with the third equality in (2) elimiated:

$$\omega_{t} + \nabla \cdot \omega \nabla^{\perp} \psi = \nabla \cdot (\nabla \omega + \beta \omega \nabla \psi), \quad \frac{\partial \omega}{\partial \nu} + \beta \omega \frac{\partial \psi}{\partial \nu}\Big|_{\partial \Omega} = 0, \quad \omega|_{t=0} = \omega_{0}(x) > 0$$
$$-\Delta \psi = \omega, \quad \psi|_{\partial \Omega} = 0. \tag{31}$$

4

Then there arise the mass conservation

$$\frac{d}{dt}\int_{\Omega}\omega=0,$$
(32)

and the free energy decreasing

$$\frac{d}{dt} \int_{\Omega} \omega (\log \omega - 1) + \frac{\beta}{2} |\nabla \psi|^2 \ dx = -\int_{\Omega} \omega |\nabla (\log \omega + \beta \psi)|^2 \le 0.$$
(33)

The system (31) without vortex term,

$$\omega_{t} = \nabla \cdot (\nabla \omega + \beta \omega \nabla \psi), \quad \frac{\partial \omega}{\partial \nu} + \beta \omega \frac{\partial \psi}{\partial \nu}\Big|_{\partial \Omega} = 0, \quad \omega|_{t=0} = \omega_{0}(x) > 0$$

$$-\Delta \psi = \omega, \quad \psi|_{\partial \Omega} = 0.$$
(34)

is called the Smoluchowski-Poisson equation. This model is concerned on the thermodynamics of self-gravitating Brownian particles [18] and has been studied in the context of chemotaxis [19–23]. We have a blowup threshold to (34) as a consequence of the quantized blowup mechanism [19, 23]. The results on the existence of the bounded global-in-time solution [24–26] and blowup of the solution in finite time [27] are valid even to the case that  $\beta$  is a function of t as in  $\beta = \beta(t)$ . provided with the vortex term  $\nabla \cdot \omega \nabla^{\perp} \psi$  on the right-hand side. We thus obtain the following theorems.

Theorem 3 It holds that

$$-\beta(t) \le \delta, \quad \|\omega_0\|_1 < 8\pi\delta^{-1} \Rightarrow T = +\infty, \quad \|\omega(\cdot, t)\|_\infty \le C \tag{35}$$

in (31), where  $\delta > 0$  is arbitrary. **Theorem 4** It holds that

 $-\beta(t) \ge \delta$ ,  $\|\omega_0\|_1 > 8\pi\delta^{-1} \Rightarrow \exists \omega_0 > 0$ ,  $\|\omega_0\|_1 > 8\pi\delta^{-1}$  such that  $T < +\infty$  (36)

in (31), where  $\delta > 0$  is arbitrary.

**Remark 4** In the context of chemotaxis in biology, the boundary condition of  $\psi$  is required to be the form of Neumann zero. The Poisson equation in (34) is thus replaced by

$$-\Delta \psi = \omega - \frac{1}{|\Omega|} \int_{\Omega} \omega, \quad \frac{\partial \psi}{\partial \nu} \Big|_{\partial \Omega} = 0$$
(37)

or

$$-\Delta \psi + \psi = \omega, \quad \frac{\partial \psi}{\partial \nu}\Big|_{\partial \Omega} = 0$$
 (38)

by [28] and [29], respectively. In this case there arises the boundary blowup, which reduces the value  $8\pi$  in Theorems 3–4 to  $4\pi$ . The value  $8\pi$  in Theorems 3–4, therefore, is a consequence of the exclusion of the boundary blowup [30]. This property is valid even for (37) or (38) of the Poisson part, if (22) is assumed.

**Remark 5** The requirement to  $\omega_0$  in Theorem 4 is the concentration at an interior point, which is not necessary in the case of (22). Hence Theorems 3 and 4 are refined as

$$-\beta(t) \le \delta, \quad \|\omega_0\|_1 < 8\pi\delta^{-1} \Rightarrow T = +\infty, \quad \|\omega(\cdot, t)\|_{\infty} \le C$$
(39)

and

$$-\beta(t) \ge \delta, \quad \|\omega_0\|_1 > 8\pi\delta^{-1} \Rightarrow T < +\infty, \tag{40}$$

if (22) holds in (35). The main task for the proof of Theorems 1 and 2, therefore, is a control of  $\beta = \beta(t)$  in (1).

This paper is composed of four sections and an appendix. Section 2 is devoted to the study on the stationary solutions, and Theorems 1 and 2 are proven in Sections 3 and 4, respectively. Then Theorem 4 is confirmed in Appendix.

#### 2. Stationary states

First, we take the canonical system (31) with  $\beta$  independent of *t*. By (32) and (33), its stationary state is defined by

$$\log \omega + \beta \psi = \text{constant}, \ \omega = \omega(x) > 0, \quad \int_{\Omega} \omega = M.$$
 (41)

Then it holds that

$$\omega = \frac{Me^{-\beta\psi}}{\int_{\Omega} e^{-\beta\psi}} \tag{42}$$

and hence

$$-\Delta \psi = \frac{M e^{-\beta \psi}}{\int_{\Omega} e^{-\beta \psi}}, \quad \psi|_{\partial \Omega} = 0.$$
(43)

There arises an oredered structure arises in  $\beta < 0$ , as observed by [11], as a consequence of a quantized blowup mechanism [19, 20, 31]. In the microcanonical system (1) and (2), the value  $\beta$  in (43) has to be determined by *E* besides *M*.

Equality (21), however, still ensures (41) and hence (42) in the stationary state even for (1)-(3). Writing



we obtain

$$-\Delta v = \mu e^{v} \text{ in } \Omega, \quad v|_{\partial\Omega} = 0, \quad \frac{E}{M^{2}} = \frac{\|\nabla v\|_{2}^{2}}{\left(\int_{\Omega} - \frac{\partial v}{\partial v}\right)^{2}}$$
(45)

by (30) and (43).

This system is the stationary state of (1) and (2) introduced by [4]. The first two equalities

$$-\Delta v = \mu e^{v}, \quad v|_{\partial\Omega} = 0 \tag{46}$$

comprise a nonlinear elliptic eigenvalue problem and the unknown eigenvalue  $\mu$  is determined by the third equality,

$$\frac{E}{M^2} = \frac{\|\nabla v\|_2^2}{\left(\int_{\Omega} - \frac{\partial v}{\partial \nu}\right)^2}.$$
(47)

The elliptic theory ensures rather deailed features of the set of solutions to (46). Here we note the following facts [31].

- 1. There is  $\overline{\mu} = \mu(\Omega) > 0$  such that the problem (46) does not admit a solution for  $\mu > \overline{\mu}$ .
- 2. Each  $\mu \leq 0$  admits a unique solution.
- 3. Each  $0 < \delta < \overline{\mu}$  admits a constant  $C = C(\delta) > 0$  such that  $||v||_{\infty} \le C$  for any solution v = v(x).

4. There is a family of solutions  $\{(\mu, v)\}$  such that  $\mu \downarrow 0$  and  $\|v\|_{\infty} \to +\infty$ .

We show the following theorem, consistent to Theorem 2. **Theorem 5** If  $\Omega = B(0, 1) \subset \mathbb{R}^2$ , there is  $\delta > 0$  such that any solution  $(v, \mu)$  to (45) admits

$$\frac{E}{M^2} \ge \delta. \tag{48}$$

*Proof:* If  $\mu = 0$ , it holds that v = 0. We have  $\pm v > 0$  exclusively in  $\Omega$ , provided that  $\pm \mu > 0$ , respectively. By the elliptic theory [32], therefore, any solution v to (46) is radially symmetric as in v = v(r), r = |x|. We have, furthermore,  $\pm v_r < 0$  in  $0 < r \le 1$ , if  $\pm \mu > 0$ , respectively.

Then it holds that  $\psi = \psi(r)$ , and hence

$$-\frac{1}{r}(r\psi_r)_r = \omega \text{ in } 0 < r \le 1, \quad \psi|_{r=1} = 0$$
(49)

by (42) and (43), which implies

$$-r\psi_r(r) = \int_0^r s\omega(s)ds > 0, \quad 0 < r \le 1.$$
(50)

We thus obtain  $\mu \neq 0$ , in particular.

If  $\mu < 0$  we have  $\beta > 0$  by (44), and therefore,  $\psi_r > 0$  in  $0 < r \le 1$  by  $v_r > 0$  there. It is a contradiction, and hence  $\mu > 0$ . In this case, the solution v = v(r) to (46) is explicit [31]. The numbers of the solution is 0, 1, and 2, according to  $\mu > 2$ ,  $\mu = 2$ , and  $0 < \mu < 2$ , respectively, and if  $0 < \mu \le 2$  the solutions  $v = v_{\pm}$  are given as

$$v_{\pm}(r) = \log \frac{8\gamma_{\pm}r}{\left(1 + \gamma_{\pm}r^{2}\right)^{2}}, \quad \gamma_{\pm} = \frac{4}{\mu} \left\{ 1 - \frac{\mu}{4} \pm \left(1 - \frac{\mu}{2}\right)^{1/2} \right\}.$$
 (51)

In fact, we have  $\gamma_+ = \gamma_-$  for  $\mu = 2$ . This solution is parametrized by

$$\sigma = \int_{\Omega} \mu e^{\nu} \in (0, 8\pi).$$
(52)

Hence each  $0 < \sigma < 8\pi$  admits a unique solution  $(v, \mu)$  to (46), and  $v = v_+$  and  $v = v_-$  according as  $\sigma \ge 4\pi$  and  $\sigma \le 4\pi$ , respectively. It holds also that  $\mu \downarrow 0$  if either

 $\sigma \uparrow 8\pi$  or  $\sigma \downarrow 0$ . Thus we have only to confirm that  $E/M^2$  is bounded, both as  $\sigma \uparrow 8\pi$  and  $\sigma \downarrow 0$ .

As  $\sigma \uparrow 8\pi$ , we have

$$v = v_+(x) \rightarrow 4 \log \frac{1}{|x|}$$
 locally uniformly on  $\overline{\Omega} \setminus \{0\}$  (53)

and hence

which implies  

$$\|\nabla v\|_{2}^{2} \to +\infty, \quad \int_{\partial\Omega} -\frac{\partial v}{\partial\nu} \to 8\pi, \quad (54)$$

$$\lim_{\sigma \uparrow 8\pi} \frac{E}{M^{2}} = +\infty. \quad (55)$$

As  $\sigma \downarrow 0$ , on the other hand, we have

$$v = v_{-}(x) \rightarrow 0$$
 uniformly in  $\overline{\Omega}$ . (56)

Since  $\mu \downarrow 0$ , furthermore, there arises that

$$\gamma = \gamma_{-} = \frac{4}{\mu} \left\{ 1 - \frac{\mu}{4} - \left( 1 - \frac{\mu}{2} \right)^{1/2} \right\} = \mu (1 + o(1)).$$
(57)

It holds also that

$$v(r) = \log \frac{8\gamma}{\mu} - 2\log\left(1 + \mu r^2\right)$$
(58)

and hence

$$v_r(r) = -\frac{4\mu r}{(1+\mu r^2)^2} = -4\mu r(1+o(1))$$
 uniformly on  $\overline{\Omega}$ . (59)

Then, (59) implies  
$$\|\nabla v\|_{2}^{2} = 2\pi \int_{0}^{1} v_{r}^{2} r \, dr = 2\pi \cdot 16\mu^{2} \cdot \int_{0}^{1} r^{3} \, dr \cdot (1+o(1))$$
(60)
$$= 8\pi \mu^{2} (1+o(1))$$

as well as

$$\left(\int_{\partial\Omega} -\frac{\partial v}{\partial\nu}\right)^2 = 16\mu^2 \cdot 2\pi (1+o(1)).$$
(61)

It thus follows that

$$\lim_{\sigma \downarrow 0} \frac{E}{M^2} = \frac{1}{4} \tag{62}$$

and hence the conclusion.

#### 3. Proof of Theorem 1

The first observation is the following lemma. **Lemma 1** *Under the assumption of* (22), *it holds that* 

$$\beta = \beta(t) < 0, \quad \omega_r(r,t) < 0, \quad 0 < r \le 1, \quad 0 \le t < T.$$
(63)

*Proof:* We have (7) and hence

$$\psi_{r}(r,t) < 0, \quad 0 < r \le 1, \quad 0 \le t < T$$
(64)  
by (49), which implies, in particular,  
$$\beta = -\frac{(\nabla \omega, \nabla \psi)}{\int_{\Omega} \omega |\nabla \psi|^{2}} < 0$$
(65)

at t = 0 by (22). Since  $\omega = \omega(r, t)$  and  $\psi = \psi(r, t)$ , we obtain  $\nabla^{\perp} \psi = 0$ , and hence

$$\omega_t = \omega_{rr} + \frac{1}{r}\omega_r + \beta\psi_r\omega_r - \beta\omega^2$$
(66)

by (1). Then  $z = \omega_r$  satisfies

$$z_{t} = z_{rr} - \frac{1}{r^{2}}z + \frac{1}{r}z_{r} + \beta \psi_{rr}z + \beta \psi_{r}z_{r} - 2\beta \omega z, \quad 0 < r \le 1, \quad 0 \le t < T$$

$$z|_{r=0} = 0, \qquad z|_{t=0} = \omega_{0r}(r) < 0, \quad 0 < r \le 1$$
(67)

and

$$z = -\beta \omega \psi_r, \quad r = 1, \quad 0 \le t < T.$$
(68)

Putting

$$m(t) = \min_{\partial \Omega} z(\cdot, t) = \omega_r(\cdot, t)|_{r=1},$$
(69)

we obtain 
$$m(0) < 0$$
 from the assumption. If there is  $0 < t_0 <$  such that  
 $m(t) < 0, \quad 0 \le t < t_0 < T, \quad m(t_0) = 0,$ 
(70)

we obtain z(r,t) > 0 for  $0 \le t < t_0$ ,  $0 < r \le 1$ , and  $t = t_0$ , 0 < r < 1 by the strong maximum principle. By (64), we have (65) for  $0 \le t \le t_0$ , that is,

$$\beta = -\frac{\int_0^1 \psi_r zr \, dr}{\int_0^1 \omega \psi_r^2 r \, dr} < 0, \quad 0 \le t \le t_0,$$
(71)

and hence

$$z = -\beta \omega \psi_r < 0 \quad r = 1, \ t = t_0, \tag{72}$$

a contradiction. It holds that  $z = \omega_r < 0$  for  $0 \le t < T$ , r = 1, and hence

$$\beta = -\frac{\int_0^1 \psi_r \omega_r r dr}{\int_0^1 \omega \psi_r^2 r \ dr} < 0, \quad 0 \le t < T. \qquad \Box$$
(73)

The proof of Theorem 3 relies on the fact

$$\beta \ge -C, \quad \int_{\Omega} \omega(\log \omega - 1) \le C \Rightarrow T = +\infty, \quad \|\omega(\cdot, t)\|_{\infty} \le C.$$
 (74)

This property is known for the Smoluchoski-Poisson equation (34), but the proof is valid even to (31) with vortex term. Having (21), therefore, we have to provide the inequality  $\beta \ge -C$ .

The inequality  $\beta < 0$ , on the other hand, is sufficient for the following arguments.

Lemma 2 If 
$$\beta \le 0$$
,  $0 \le t < T$ , it holds that  
 $\omega \ge \underline{\omega} \equiv \min_{\overline{\Omega}} \omega_0 > 0$  on  $\overline{\Omega} \times [0, T]$ . (75)

*Proof:* Since (17) we obtain

$$\omega_{t} + \nabla^{\perp}\psi \cdot \nabla\omega = \Delta\omega + \beta\nabla\psi \cdot \nabla\omega + \beta\Delta\psi$$
$$= \Delta\omega + \beta\nabla\psi \cdot \nabla\omega - \beta\omega^{2}$$
$$\geq \Delta\omega + \beta\nabla\psi \cdot \nabla\omega \quad \text{in } \Omega \times (0, T)$$
(76)

with

$$-\frac{\partial\omega}{\partial\nu} = \beta\omega\frac{\partial\psi}{\partial\nu} > 0 \quad \text{on } \partial\Omega \times [0,T)$$
(77)

**Lemma 3** Under the assumption of the previous lemma, there is  $C_0 = C_0(\Omega) > 0$  such that

$$C_0 \|\omega_0\|_2^3 \le E\underline{\omega} \Rightarrow \|\omega(\cdot, t)\|_2 \le \|\omega_0\|_2, \quad -\beta(t) \le \alpha \equiv \frac{\|\omega_0\|_2^2}{\underline{E}\underline{\omega}}, \quad 0 \le t < T.$$
(78)

*Proof:* Using (11) and (17), we obtain

$$\int_{\Omega} \left[ \nabla \cdot \left( \omega \nabla^{\perp} \psi \right) \right] \omega = \int_{\Omega} \omega \nabla \omega \cdot \nabla^{\perp} \psi = \frac{1}{2} \int_{\Omega} \nabla \omega^{2} \cdot \nabla^{\perp} \psi$$

$$= -\frac{1}{2} \int_{\Omega} \omega^{2} \nabla \cdot \nabla^{\perp} \psi = 0.$$
(79)

Hence (1) with (2) implies

$$\frac{1}{2}\frac{d}{dt}\|\omega\|_{2}^{2} + \|\nabla\omega\|_{2}^{2} = -\beta \int_{\Omega} \omega \nabla\psi \cdot \nabla\omega = -\frac{\beta}{2} \left(\nabla\psi, \nabla\omega^{2}\right)$$
$$= -\frac{\beta}{2} \int_{\partial\Omega} \omega^{2} \frac{\partial\psi}{\partial\nu} + \frac{\beta}{2} \left(\Delta\psi, \omega^{2}\right) \leq -\frac{\beta}{2} \|\omega\|_{3}^{3}$$
(80)

by  $\beta \leq 0$  and (88). Since

$$\int_{\Omega} \nabla \omega \cdot \nabla \psi = \int_{\partial \Omega} \omega \frac{\partial \psi}{\partial \nu} + \int_{\Omega} \omega (-\Delta \psi) \le \int_{\Omega} \omega^2$$
(81)

follows from (8), furthermore, it holds that

$$-\beta = \frac{\int_{\Omega} \nabla \omega \cdot \nabla \psi}{\int_{\Omega} \omega |\nabla \psi|^2} \le \underline{\omega}^{-1} \cdot \frac{\|\omega\|_2^2}{\|\nabla \psi\|_2^2} = \frac{1}{E\underline{\omega}} \|\omega\|_2^2.$$
(82)

Then ineqality (80) induces

$$\frac{1}{2}\frac{d}{dt}\|\omega\|_{2}^{2} + \|\nabla\omega\|_{2}^{2} \le \frac{1}{2E\underline{\omega}}\|\omega\|_{2}^{2} \cdot \|\omega\|_{3}^{3}.$$
(83)

Here we use the Gagliardo-Nirenberg inequality (see (4.16) of [19]) in the form of

$$\|\omega\|_{3}^{3} \leq C \|\omega\|_{H^{1}} \cdot \|\omega\|_{2}^{2} = C \|\omega\|_{2}^{2} (\|\nabla\omega\|_{2} + \|\omega\|_{2}),$$
(84)

to obtain

$$\frac{1}{2}\frac{d}{dt}\|\omega\|_{2}^{2} + \|\nabla\omega\|_{2}^{2} \leq \frac{C}{E\underline{\omega}}\|\omega\|_{2}^{4}(\|\nabla\omega\|_{2} + \|\omega\|_{2}) \\
\leq \frac{1}{2}\|\nabla\omega\|_{2}^{2} + \frac{C^{2}}{8(E\underline{\omega})^{2}}\|\omega\|_{2}^{8} + \frac{C}{2E\underline{\omega}}\|\omega\|_{2}^{5}$$
(85)

and hence

$$\frac{d}{dt} \|\omega\|_2^2 + \|\nabla\omega\|_2^2 \le \frac{C}{E\underline{\omega}} \|\omega\|_2^5 \left(\frac{C}{E\underline{\omega}} \|\omega\|_2^3 + 1\right).$$
(86)

Then, Poincaré-Wirtinger's inequality ensures

$$\frac{d}{dt} \|\omega\|_2^2 + \mu \|\omega\|_2^2 \le \frac{C}{E\underline{\omega}} \left(\frac{C}{E\underline{\omega}} \|\omega\|_2^6 + \|\omega\|_2^3\right) \|\omega\|_2^2, \tag{87}$$

where  $\mu = \mu(\Omega) > 0$  is a constant. Writing

$$y(t) = \frac{C}{E\underline{\omega}} \|\omega\|_2^3,$$
(88)
we obtain

$$\frac{d}{dt} \|\omega\|_2^2 + \mu \|\omega\|_2^2 \le (y^2 + y) \|\omega\|_2^2,$$
(89)

and therefore, if

$$y^2 + y < \mu/2 \tag{90}$$

holds at t = 0, it keeps to hold that

$$\frac{d}{dt}\|\omega\|_2^2 \le 0 \tag{91}$$

and (90) for  $0 \le t < T$ . Then, we obtain

$$\|\omega(\cdot, t)\|_{2} \le \|\omega_{0}\|_{2}, \quad 0 \le t < T,$$
(92)

and hence

$$-\beta(t) \le \frac{\|\omega_0\|_2^2}{E\underline{\omega}} = \alpha, \quad 0 \le t < T$$
(93)

by (82).

The condition  $y(0) < \frac{\mu}{2}$  means

 $C_0 \|\omega_0\|_2 \le E\underline{\omega}$  (94) for  $C_0 > 0$  sufficiently large, and hence we obtain the conclusion. *Proof of Theorem 1:* By the parabolic regularity, it suffices to show that

$$\|\omega(\cdot,t)\|_{\infty} \le C, \quad 0 \le t < T \tag{95}$$

under the assumption. We have readily shown

$$\|\omega(\cdot,t)\|_2 \le C, \ \ 0 \le -\beta(t) \le C, \ \ 0 \le t < T$$
 (96)

by Lemma 3. Then, the conclusion (95) is obtained similarly to (34). See [26] for more details.

In fact, we have

$$\int_{\Omega} \left[ \nabla \cdot \left( \omega \nabla^{\perp} \psi \right) \right] \omega^{p} = -\int_{\Omega} \omega \nabla^{\perp} \psi \cdot \nabla \omega^{p} = -p \int_{\Omega} \omega^{p} \nabla^{\perp} \psi \cdot \nabla \omega$$
$$= -\frac{p}{p+1} \int_{\Omega} \nabla^{\perp} \psi \cdot \nabla \omega^{p+1} = \frac{p}{p+1} \int_{\Omega} \omega^{p+1} \nabla \cdot \left( \nabla^{\perp} \psi \right) = 0$$
(97)

for p > 0 by (11) and (34). Then it follows that

$$\frac{1}{p+1}\frac{d}{dt}\int_{\Omega}\omega^{p+1} + \frac{4p}{(p+1)^2} \|\nabla\omega^{\frac{p+1}{2}}\|_2^2 = -\beta \int_{\Omega}\omega\nabla\psi\cdot\nabla\omega^p$$

$$= -\beta \cdot \frac{p}{p+1}\int_{\Omega}\nabla\psi\cdot\nabla\omega^{p+1} \le -\beta \frac{p}{p+1}\int_{\Omega}\omega^{p+1}(-\Delta\psi)$$

$$= -\beta \frac{p}{p+1}\int_{\Omega}\omega^{p+1} \le C \int_{\Omega}\omega^{p+2}$$
(98)

by  $\beta < 0$  and (8). Then, Moser's iteration scheme ensures (95) as in [33].

#### 4. Proof of Theorem 2

We begin with the following lemma. **Lemma 4** *Under the assumption of* (22), *it holds that* 

$$-\beta(t) \ge \delta, \quad 0 \le t < T, \quad M = \|\omega_0\|_1 > \frac{8\pi}{\delta} \Rightarrow T < +\infty$$
(99)

in (31), where  $\delta > 0$  is a constant.

*Proof:* We have  $\omega = \omega(r, t)$  and  $\psi = \psi(r, t)$  for r = |x| under the assumption, which implies  $\nabla^{\perp}\psi = 0$ . Then we obtain

$$\nabla \cdot \omega \nabla^{\perp} \psi = \nabla \omega \cdot \nabla^{\perp} \psi = 0 \tag{100}$$

by (17). It holds also that

$$\nabla \cdot (\omega \nabla \psi) = \nabla \cdot \left( \omega \psi_r \frac{x}{r} \right) = \left( \nabla \cdot \frac{x}{r} \right) \omega \psi_r + \frac{x}{r} \cdot \nabla (\omega \psi_r)$$

$$= \frac{1}{r} \omega \psi_r + (\omega \psi_r)_r = \frac{1}{r} (r \omega \psi_r)_r,$$
and therefore, there arises that
$$(101)$$

$$\omega_t = \frac{1}{r} (r\omega_r + \beta r \omega \psi_r)_r, \quad \omega_r + \beta \omega \psi_r|_{r=1} = 0.$$
 (102)

from (31). Then (102) implies

$$\frac{d}{dt} \int_{0}^{1} \omega r^{3} dr = \int_{0}^{1} \omega_{t} r^{3} dr = \int_{0}^{1} (r\omega_{r} + \beta r \omega \psi_{r})_{r} r^{2} dr$$

$$= -\int_{0}^{1} 2r^{2} (\omega_{r} + \beta \omega \psi_{r}) dr$$

$$= -2r^{2} \omega \Big|_{r=0}^{r=1} + \int_{0}^{1} 4r\omega - 2\beta \omega \psi_{r} r^{2} dr.$$
(103)

Here we use (50) derived from the Poisson part of (31), that is,

$$-r\psi_r(r,t) = A(r,t) \equiv \int_0^r s\omega(s,t)ds.$$
 (104)

Putting

$$\lambda = \int_{0}^{1} \omega r \, dr = \frac{M}{2\pi}, \qquad (105)$$
we obtain
$$\frac{d}{dt} \int_{0}^{1} \omega r^{3} \, dr = -2\omega|_{r=1} + 4\lambda + 2\beta \int_{0}^{1} AA_{r} \, dr$$

$$= -2\omega|_{r=1} + 4\lambda + \beta A^{2}|_{r=0}^{r=1}$$

$$= -2\omega|_{r=1} + 4\lambda + \beta \lambda^{2}$$

$$< 4\lambda \left(\beta + \frac{M}{8\pi}\right) \le 4\lambda \left(-\delta + \frac{M}{8\pi}\right).$$

Since  $-\delta + \frac{M}{8\pi} < 0$ , therefore,  $T = +\infty$  is impossible, and we obtain  $T < +\infty$ . **Lemma 5** Under the assumption (22), there is  $\delta > 0$  such that

$$\frac{E}{M^2} < \delta, \quad \beta(t) \le 0, \quad 0 \le t < T \Rightarrow \beta(t) \le -\frac{1}{CE^{1/2}} \quad 0 \le t < T.$$
(107)

Proof: First, Lemma 1 implies

$$\omega \ge \omega_* \equiv \omega|_{r=1}.\tag{108}$$

Second, we have

$$\int_{\Omega} \nabla \psi \cdot \nabla \omega = \int_{\partial \Omega} \frac{\partial \psi}{\partial \nu} \omega + \int_{\Omega} (-\Delta \psi) \omega = \omega_* \int_{\partial \Omega} \frac{\partial \psi}{\partial \nu} + \|\omega\|_2^2$$
(109)  

$$= \omega_* \int_{\Omega} \Delta \psi + \|\omega\|_2^2 = \|\omega\|_2^2 - \omega_* M,$$

$$-\beta = \frac{\int_{\Omega} \nabla \psi \cdot \nabla \omega}{2} = \frac{\|\omega\|_2^2 - \omega_* M}{2}.$$
(110)

 $-p = \frac{1}{\int_{\Omega} \omega |\nabla \psi|^2} = \frac{1}{\int_{\Omega} \omega |\nabla \psi|^2}.$ 

Here, we use the Gagliardo-Nirenberg inequality in the form of

$$\|w\|_{4}^{2} \le C \|w\|_{2} \|w\|_{H^{1}}, \tag{111}$$

which implies

$$\int_{\Omega} \omega |\nabla \psi|^{2} \leq ||\omega||_{2} ||\nabla \psi||_{4}^{2} \leq C ||\omega||_{2} ||\nabla \psi||_{2} ||\nabla \psi||_{H^{1}}$$

$$\leq C E^{1/2} ||\omega||_{2}^{2}$$
(112)

by the elliptic estimate of the Poisson equation in (2),

$$\|\psi\|_{H^2} \le C \|\omega\|_2. \tag{113}$$

We have, on the other hand,

$$\omega_* M \le \frac{M}{E} \int_{\Omega} \omega |\nabla \psi|^2$$
(114)  
by (110), and therefore,  
$$-\beta \ge \frac{1}{CE^{1/2}} - \frac{E}{M} \ge \frac{1}{2CE^{1/2}},$$
(115)

provided that

$$\frac{E}{M^2} < \left(\frac{1}{2C}\right)^2. \tag{116}$$

Then the conclusion follows.

*Proof of Theorem 2:* By Lemma 5, there is  $\delta_0 >$  such that

$$\frac{E}{M^2} < \delta \Rightarrow -\beta \ge \frac{1}{CE^{1/2}} \equiv \delta_1, \tag{117}$$

and then, Lemma 4 ensures

$$M > \frac{8\pi}{\delta_1} \Rightarrow T < +\infty.$$
(118)

The assumption in (118) means



#### **Appendix Proof of Theorem 4**

This theorem is valid to the general case of  $\Omega$  and  $\omega_0$  without (22). We assume  $\delta = 1$  without loss of generation, so that

$$\beta \le -1. \tag{120}$$

We follow the argument [27] concerning (34) with the Poisson part replaced by (42) or (43). Thus we have to take case of the vortex term  $\nabla \cdot \omega \nabla^{\perp} \psi$ , time varying  $\beta = \beta(t)$ , and the Dirichlet boundary condition in (31).

We recall the cut-off function used in [34] (see also Chapter 5 of [19]). Hence each  $x_0 \in \overline{\Omega}$  and  $0 < R \le 1$  admit  $\varphi = \varphi_{x_0,R} \in C^2(\overline{\Omega})$  with

$$\frac{\partial \varphi}{\partial \nu}\Big|_{\partial \Omega} = 0, \quad 0 \le \varphi \le 1, \quad \varphi = 1 \text{ in } \Omega \cap B(x_0, R/2), \quad \varphi = 0 \text{ in } \Omega \setminus B(x_0, R),$$
(121)

and

$$|\nabla \varphi| \le CR^{-1} \varphi^{1/2}, \quad |\nabla^2 \varphi| \le CR^{-2} \varphi^{1/2}.$$
 (122)

In more details, we take a cut-off function, denoted by  $\psi$ , satisfying (121), using a local conformal mapping, and then put  $\varphi = \psi^4$ . Let

$$\varphi \in C^2(\overline{\Omega}), \quad \frac{\partial \varphi}{\partial \nu}\Big|_{\partial \Omega} = 0.$$
 (123)

be given. First, we have

$$\frac{d}{dt} \int_{\Omega} \omega \varphi = \int_{\Omega} \omega \nabla^{\perp} \psi \cdot \nabla \varphi - (\nabla \omega + \beta \omega \nabla \psi) \cdot \nabla \varphi \, dx$$

$$= \int_{\Omega} \omega \nabla^{\perp} \psi \cdot \nabla \varphi + \omega \Delta \varphi - \beta \omega \nabla \psi \cdot \nabla \varphi \, dx$$
(124)

by (11). It holds that

$$\begin{split} \int_{\Omega} &\omega \nabla \psi \cdot \nabla \varphi = \iint_{\Omega \times \Omega} \omega(x,t) [\nabla_x G(x,x') \cdot \nabla \varphi(x)] \omega(x',t) \ dx dx' \\ &= \iint_{\Omega \Omega} \omega(x,t) \varphi_{x_0,2R}(x') [\nabla_x G(x,x') \cdot \nabla \varphi(x)] \omega(x',t) \ dx dx' \\ &+ \iint_{\Omega \times \Omega} \omega(x,t) \left(1 - \varphi_{x_0,2R}(x')\right) [\nabla_x G(x,x') \cdot \nabla \varphi(x)] \omega(x',t) \ dx dx' \\ &= I + II. \end{split}$$
(125)  
Let, furthermore,  $x_0 \in \Omega$  and  $0 < R \ll 1$  in the above equality. Then,  
 $\varphi = |x - x_0|^2 \varphi_{x_0,R}$ (126)

satisfies the requirement (123). It holds that

$$\nabla \varphi = 2(x - x_0)\varphi_{x_0,R} + |x - x_0|^2 \nabla \varphi_{x_0,R}$$
(127)

and hence

$$|\nabla \varphi| \le C|x - x_0| \left( \varphi_{x_0,R} + |x - x_0| R^{-1} \varphi_{x_0,R}^{1/2} \right) \le C|x - x_0| \varphi_{x_0,R}^{1/2}.$$
(128)

We obtain, furthermore,

$$|x' - x_0| \ge 2R, \ |x - x_0| \le R \Rightarrow |x - x'| \ge R,$$
 (129)

and hence

$$|\nabla_x G(x, x')| \le CR^{-1} \tag{130}$$

in this case. Then it follows that

$$|II| \le CR^{-1}M \int_{\Omega} |x - x_0| \varphi_{x_0,R}^{1/2} \omega(x,t) \, dx \le CR^{-1}M^{3/2}A^{1/2}, \tag{131}$$
  
where  
$$A = \int_{\Omega} |x - x_0|^2 \varphi_{x_0,R} \omega. \tag{132}$$

We have, on the other hand,

$$I = \iint_{\Omega \times \Omega} \omega(x,t) \varphi_{x_0,2R}(x') [\nabla_x G(x,x') \cdot \nabla \varphi(x)] \omega(x',t) \ dxdx'$$
  
$$= \frac{1}{2} \iint_{\Omega\Omega} \left[ \varphi_{x_0,2R}(x') \nabla \varphi(x) \cdot \nabla_x G(x,x') + \varphi_{x_0,2R}(x) \nabla \varphi(x') \cdot \nabla_{x'} G(x,x') \right] \omega \otimes \omega,$$
  
(133)

where  $G=G(x,x^\prime)$  is the Green's function to

$$-\Delta \psi = \omega, \quad \omega|_{\partial\Omega} = 0 \tag{134}$$

and

$$\omega \otimes \omega = \omega(x,t)\omega(x',t) \ dxdx'. \tag{135}$$

Here we use the local property of the Green's function

$$G(x,x') = \Gamma(x-x') + K(x,x'), \quad K \in C^2(\overline{\Omega} \times \Omega) \cap C^2(\Omega \times \overline{\Omega}),$$
(136)

where

$$\Gamma(x) = \frac{1}{2\pi} \log \frac{1}{|x|}$$
(137)  
stands for the fundamental solution to  $-\Delta$ .  
Let  
 $\rho_{x_0,R}^2(x,x() = \varphi_{x_0,2R}(x')\nabla\varphi(x) \cdot \nabla_x K(x,x') + \varphi_{x_0,2R}\nabla\varphi(x') \cdot \nabla_{x'} K(x,x').$ (138)

Since (128) implies

$$\begin{aligned} |\varphi_{x_0,2R}(x')\nabla\varphi(x)| &\leq C\varphi_{x_0,2R}(x')|x - x_0|\varphi_{x_0,R}^{1/2}(x) \\ &\leq C|x - x_0|\varphi_{x_0,R}^{1/2}(x), \end{aligned}$$
(139)

it holds that

$$|\rho_{x_0,R}^1(x,x')| \le C \Big( |x-x_0| \varphi_{x_0,R}^{1/2}(x) + |x'-x_0| \varphi_{x_0,R}^{1/2}(x') \Big).$$
(140)

Then, we obtain

$$I = \frac{1}{2} \iint_{\Omega \times \Omega} \rho_{x_0, R}^0(x, x') \omega \otimes \omega + III$$
(141)

with

$$|III| \le CM^{3/2} A^{1/2} \le CR^{-1} M^{3/2} A^{1/2}, \tag{142}$$

where  

$$\rho_{x_0,R}^0(x,x') = \nabla \Gamma(x-x') \cdot \left(\varphi_{x_0,2R}(x') \nabla \varphi(x) - \varphi_{x_0,2R}(x) \nabla \varphi(x')\right). \quad (143)$$
Here, we have  

$$\nabla \Gamma(x) = -\frac{x}{2\pi |x|^2}, \quad (144)$$

and therefore,

$$\rho_{x_0,R}^0(x,x') = \rho_{x_0,R}^2(x,x') + \rho_{x_0,R}^3(x,x')$$
(145)

fo

$$\rho_{x_0,R}^2(x,x') = -\frac{1}{2\pi} \frac{x-x'}{|x-x'|^2} \varphi_{x_0,2R}(x') \cdot (\nabla \varphi(x) - \nabla \varphi(x'))$$
(146)

$$\rho_{x_0,R}^3(x,x') = -\frac{1}{2\pi} \frac{x-x'}{|x-x'|^2} \left( \varphi_{x_0,2R}(x') - \varphi_{x_0,2R}(x) \right) \cdot \nabla \varphi(x).$$
(147)

Since (128) implies

$$|\rho_{x_0,R}^3(x,x')| \le CR^{-1} |\nabla \varphi(x)| \le CR^{-1} |x - x_0| \varphi_{x_0,R}^{1/2}(x),$$
(148)

there arises that

with

$$I = \frac{1}{2} \iint_{\Omega \times \Omega} \rho_{x_0,R}^2(x, x') \quad \omega \otimes \omega + IV,$$
(149)

 $|IV| \le CR^{-1}M^{3/2}A^{1/2},$  (150) we have, furthermore,

$$\nabla\varphi(x) - \nabla\varphi(x') = 2(x - x')\varphi_{x_0,R}(x) + 2(x' - x_0)\left(\varphi_{x_0,R}(x) - \varphi_{x_0,R}(x')\right) \\ + |x' - x_0|^2 \left(\nabla\varphi_{x_0,R}(x) - \nabla\varphi_{x_0,R}(x')\right) + \left(|x - x_0|^2 - |x' - x_0|^2\right)\nabla\varphi_{x_0,R}(x),$$
(151)

and hence

$$\rho_{x_0,R}^2(x,x') = -\frac{1}{\pi} \varphi_{x_0,2R}(x') \varphi_{x_0,R}(x) + \rho_{x_0,R}^4(x,x') + \rho_{x_0,R}^5(x,x') + \rho_{x_0,R}^6(x,x')$$
(152)

with

$$\begin{aligned} |\rho_{x_0,R}^4(x,x')| &\leq C|x-x'|^{-1}\varphi_{x_0,2R}(x')|x'-x_0||\varphi_{x_0,R}(x)-\varphi_{x_0,R}(x')| \\ &\leq CR^{-1}|x'-x_0|\varphi_{x_0,2R}(x'), \end{aligned}$$
(153)

$$\begin{aligned} |\rho_{x_0,R}^5(x,x')| &\leq C|x-x'|^{-1}\varphi_{x_0,2R}(x')|x'-x_0|^2|\nabla\varphi_{x_0,R}(x) - \varphi_{x_0,R}(x')| \\ &\leq CR^{-2}|x'-x_0|^2\varphi_{x_0,2R}(x') \\ &\leq CR^{-1}|x'-x_0|\varphi_{x_0,2R}(x'), \end{aligned}$$
(154)

and  

$$\begin{aligned} |\rho_{x_{0},R}^{6}(x,x')| &\leq C|x-x'|\varphi_{x_{0},2R}(x')||x-x_{0}|^{2} - |x'-x_{0}|^{2}| \cdot |\nabla\varphi_{x_{0},R}(x)| \\ &\leq CR^{-1}(|x-x_{0}| + |x'-x_{0}|)\varphi_{x_{0},R}(x)\varphi_{x_{0},2R}(x') \\ &\leq C\left(R^{-1}|x-x_{0}|\varphi_{x_{0},R}(x) + R^{-1}|x'-x_{0}|\varphi_{x_{0},2R}(x')\right) \end{aligned}$$
(155)

by

$$||x - x_0|^2 - |x' - x_0|^2| = |(x - x', x + x' - 2x_0)| \le |x - x'|(|x - x_0| + |x' - x_0|).$$
(156)

The residual terms are thus treated similarly, and it follows that

$$\left| I + \frac{1}{2\pi} \int_{\Omega} \omega \varphi_{x_0,R} \cdot \int_{\Omega} \omega \varphi_{x_0,2R} \right| \le C R^{-1} M^{3/2} A^{1/2}, \tag{157}$$

which results in

$$\left| \int_{\Omega} \omega \nabla \psi \cdot \nabla \varphi + \frac{1}{2\pi} \int_{\Omega} \omega \varphi_{x_0,R} \cdot \int_{\Omega} \omega \varphi_{x_0,2R} \right| \le CR^{-1} M^{3/2} A^{1/2}.$$
(158)

We can argue similarly to the vortex term in (124). This time, from

$$\nabla^{\perp} \Gamma(x) \cdot x = 0 \tag{159}$$

it follows that

$$\left| \int_{\Omega} \omega \nabla^{\perp} \psi \cdot \nabla \varphi \right| \le CR^{-1} M^{3/2} A^{1/2}.$$
(160)  
Concerning the principal term of (124), we use

$$\Delta \varphi = 4\varphi_{x_0,R} + 4(x - x_0) \cdot \nabla \varphi_{x_0,R} + |x - x_0|^2 \Delta \varphi_{x_0,R}.$$
 (161)

From

$$|(x - x_0) \cdot \nabla \varphi_{x_0,R}| \le CR^{-1} |x - x_0| \varphi_{x_0,R}^{1/2}$$
(162)

and

$$|-x_{0}|^{2} \Delta \varphi_{x_{0},R}| \leq CR^{-2} |x - x_{0}|^{2} \varphi_{x_{0},R}^{1/2}$$

$$\leq CR^{-1} |x - x_{0}| \varphi_{x_{0},R}^{1/2},$$
(163)

it follows that

$$\left| \int_{\Omega} \omega \Delta \varphi - 4 \int_{\Omega} \omega \varphi_{x_0,R} \right| \leq C \int_{\Omega} R^{-1} |x - x_0| \varphi_{x_0,R} \omega$$

$$\leq C R^{-1} M^{1/2} A^{1/2}.$$
(164)

Let  $M_1 = M_{x_0,R}$  and  $M_2 = M_{x_0,2R}$  for

$$M_{x_0,R} = \int_{\Omega} \omega \varphi_{x_0,R}.$$
 (165)

Then, using (120), we end up with

$$\frac{dA}{dt} \le 4M_1 - \frac{M_1^2}{2\pi} + CR^{-1} \left( M^{3/2} + M^{1/2} \right) A^{1/2} + C(M_2 - M_1).$$
(166)

Inequalily (166) implies  $T < +\infty$  if  $A(0) \ll 1$ , as is observed by [27] (see also Chapter 5 of [19]). Here we describe the proof for completeness.

The first observation is the monotoniity formula

$$\left|\frac{d}{dt}\int_{\Omega}\omega\varphi\right| \le C\left(M+M^2\right) \|\nabla\varphi\|_{C^1},\tag{167}$$

derived from (124) and the symmetry of the Green's function: G(x, x') = G(x', x). The proof is the same as in (34) and is omitted.

Second, we put  $I_1 = I_{x_0,R}$  and  $I_2 = I_{x_0,2R}$  for

$$I_{x_0,R} = \int_{\Omega} |x - x_0|^2 \omega \varphi_{x_0,R}.$$
 (168)

Then it holds that

$$M_{2} - M_{1} \leq \int_{R < |x - x_{0}| < 2R} \varphi_{x_{0}, 2R} \omega$$

$$\leq 2R^{-1} \int_{\Omega} |x - x_{0}| \varphi_{x_{0}, 2R} \omega \leq 2M^{1/2} R^{-1} I_{2}^{1/2}$$
(169)
and

$$A_{2} = A_{1} + \int_{\Omega} |x - x_{0}|^{2} (\varphi_{x_{0},2R} - \varphi_{x_{0},R}) \omega$$

$$\leq A_{1} + 4R^{2} \int_{\Omega} (\varphi_{x_{0},2R} - \varphi_{x_{0},R}) \omega,$$
(170)

which implies

$$\frac{dA_{1}}{dt} \leq 4M_{1} - \frac{M_{1}^{2}}{2\pi} + CR^{-1} \left(M^{3/2} + M^{1/2}\right) A_{1}^{1/2} + C\left(M^{3/2} + M^{1/2}\right) \left\{ \int_{\Omega} \left(\varphi_{x_{0},2R} - \varphi_{x_{0},R}\right) \omega \right\}^{1/2}.$$
(171)

Here, we use (167) to ensure

$$\left|\frac{d}{dt}\left(4M_{1} - \frac{M^{2}}{2\pi}\right| \le C(M + M^{2})R^{-2}$$
(172)

and

$$\left|\frac{d}{dt}\int_{\Omega} (\varphi_{x_0,2R} - \varphi_{x_0,R})\omega\right| \le C(M + M^2)R^{-2}.$$
(173)  
Then, it follows that

$$4M_1 - \frac{M_1^2}{2\pi} \le 4M_1(0) - \frac{M_1(0)^2}{2\pi} + CBa\left(R^{-1}t^{1/2}\right)$$
(174)

and

$$\int_{\Omega} (\varphi_{x_{0},2R} - \varphi_{x_{0},R}) \omega \leq \int_{\Omega} (\varphi_{x_{0},2R} - \varphi_{x_{0},R}) \omega_{0} + CBa \left( R^{-1} t^{1/2} \right) \\
\leq 2R^{-2} A_{2}(0) + CBa \left( R^{-1} t^{1/2} \right)$$
(175)

for

$$B = M^{3/2} + M^{1/2}, \quad a(s) = s^2 + s.$$
 (176)

Thus we obtain

$$\frac{dA_1}{dt} \le 4M_1(0) - \frac{M_1(0)^2}{2\pi} + CR^{-1}BA_1^{1/2} + CBA_2(0)^{1/2} + CBa\left(R^{-1}t^{1/2}\right)$$

$$= J(0) + CBa\left(R^{-1}t^{1/2}\right) + CBR^{-1}A_1^{1/2}$$
(177)

for

$$J = 4M_1 - \frac{M_1^2}{4\pi} + CBR^{-1}A_2^{1/2}.$$
(178)  
Assume  $M_1(0) > 8\pi$ , and put

$$-4\delta = 4M_1(0) - \frac{M_1(0)^2}{2\pi} < 0.$$
(179)

Let, furthemore,

$$\frac{1}{R^2} \int_{\Omega} |x - x_0|^2 \varphi_{x_0, 2R} \omega_0 \le \eta.$$
(180)

Now we define  $s_0$  by

$$CBa(s_0) = \delta \tag{181}$$

in (177), and take  $0 < \eta \!\ll\! 1$  such that

$$\eta \le \delta s_0^2. \tag{182}$$

Then, if R and  ${T}_0$  satisfy  $R^{-2}{T}_0=\eta\delta^{-1},$  it holds that

$$A_1(0) \le R^2 \eta < 2\delta T_0. \tag{183}$$

Making  $0 < \eta \ll 1$ , furthermore, we may assume

$$J(0) + CBR^{-1}A_{1}(0)^{1/2} \leq -4\delta + CBR^{-1}A_{2}(0)^{1/2}$$
  
$$\leq -4\delta + CB\eta^{1/2} \leq -3\delta,$$
 (184)

which results in

$$\frac{dA_1}{dt} \le J(0) + CBa\left(R^{-1}T_0^{1/2}\right) + BR^{-1}A_1(t)^{1/2}$$
$$= J(0) + \delta + CBR^{-1}A_1^{1/2}, \quad 0 \le t < T_0,$$
(185)

provided that  $T \ge T_0$ .

A continuation argument to (184)–(185) guarantees

$$\frac{dA_1}{dt} \le -2\delta, \quad 0 \le t < T_0, \tag{186}$$

and then we obtain

$$A_1(T_0) \le A_1(0) - 2\delta T_0 < 0 \tag{187}$$

by (183), a contradiction.

Acknowledgements

This work was supported by JSPS Grand-in-Aid for Scientific Research 19H01799.



1 Meteorological College, Asahi-cho, Kashiwashi, Japan

2 Center for Mathematical Modeling and Data Science, Osaka University, Toyonakashi, Japan

\*Address all correspondence to: suzuki@sigmath.es.osaka-u.ac.jp

#### **IntechOpen**

© 2021 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

#### References

[1] P.H. Chavanis, Generalized thermodynamics and Fokker-Planck equations: Applications to stellar dynamics and two-dimensional turbulence, Phys. Rev. E68, (2003) 036108.

[2] P.-H. Chavanis, *Two-dimensional Brownian vortices*, Physica A 387 (2008) 6917-6942.

[3] P.H. Chavanis, J. Sommeria, R. Robert, *Statistical mechanics of twodimensional vortices and collisionless stellar systems*, Astrophys. J. 471 (1996) 385-399.

[4] K. Sawada, T, Suzuki, *Relaxation theory for point vortices*, In; Vortex
Structures in Fluid Dynamic Problems (H. Perez-de-Tejada ed.), INTECH
2017, Chapter 11, 205-224.

[5] D. Lynden-Bell, *Statistical mechanics of violent relaxation in stellar systems*, Monthly Notices of Royal Astronmical Society 136 (1967) 101-121.

[6] E. Caglioti, P.L. Lions, C. Marchioro, M. Pulvirenti, *A special class of stationary flows for two-dimensional Euler equations: a statistical mechanics description* Comm. Math. Phys. 143 (1992) 501.

[7] G.L. Eyink, H. Spohn, Negativetemperature states and large scale, longvisited vortices in two-dimensional turbulence, Statistical Physics 70 (1993) 833.

[8] G. Joyce, D. Montgomery, *Negative temperature states for two-dimensional guiding-centre plasma*, J. Plasma Phys. 10 (1973) 107.

[9] M.K.H. Kiessling, *Statistical mechanics of classical particles with logarithmic interaction*, Comm. Pure Appl. Math. 46 (1993) 27-56.

[10] K. Nagasaki, T. Suzuki, *Asymptotic* analysis for two-dimensional elliptic

*eigenvalue problems with exponentially dominated nonlinearities*, Asymptotic Analysis 3 (1990) 173-188.

[11] L. Onsager, *Statistical hydrodynamics*, Suppl. Nuovo Cimento 6 (1949) 279-287.

[12] Y.B. Pointin, T.S. Lundgren, *Statistical mechanics of two-dimensional vortices in a bounded container*, Phys. Fluids 19 (1976) 1459-1470.

[13] R. Robert, J. Sommeria, *Statistical equilibrium states for two-dimensional flows*, J. Fluid Mech. 229 (1991) 291-310.

[14] R. Robert, J. Sommeria, *Relaxation towards a statistical equilibrium state in two-dimensional perfect fluid dynamics*, Phys. Rev. Lett. 69 (1992) 2776-2779.

[15] R. Robert, *A maximum-entropy* principle for two-dimensional perfect fluid dynamics J. Stat. Phys. **65** (1991) 531-553.

[16] R. Robert, C. Rosier, *The modeling of small scales in two-dimensional turbulent flows: A statistical Mechanics Approach*, J. Stat. Phys. 86 (1997) 481-515.

[17] K. Sawada, T. Suzuki, Derivation of the equilibrium mean field equations of point vortex system and vortex filament system, Theor. Appl. Mech. Japan 56 (2008) 285-290.

[18] C. Sire, P.-H. Chavanis, *Thermodynamics and collapse of selfgravitating Brownian particles in D dimensions*, Phys. Rev. E 66 (2002) 046133.

[19] T. Suzuki, *Free Energy and Self-Interacting Particles*, Birkhäuser, Boston, 2005.

[20] T. Suzuki, *Mean Field Theories and Dual Variation - Mathematical Structures of Mesoscopic Model*, 2nd edition, Atlantis Press, Paris, 2015. [21] T. Suzuki, *Chemotaxis, Reaction, Network - Mathematics for Self-Organization*, World Scientific, Singapore, 2018.

[22] T. Suzuki, Liouville's Theory in Linear and Nonlinear Partial Differential Equations - Interaction of Analysis, Geometry, Physics, Springer, Berlin, (to appear).

[23] T. Suzuki, *Applied Analysis -Mathematics for Science, Engineering, Technology*, 3rd edition, Imperial College Press, London, (to appear).

[24] P. Biler, Local and global solvability of some parabolic systems modelling chemotaxis, Adv. Math. Sci. Appl. 8 (1998) 715-743.

[25] H. Gajewski, K. Zacharias, *Global* behaviour of a reaction-diffusion system modelling chemotaxis, Math. Nachr. 195 (1998) 77-114.

[26] T. Nagai, T. Senba, and K. Yoshida, *Application of the Trudinger-Moser inequality to a parabolic system of chemotaxis*, Funkcial. Ekvac. 40 (1997) 411-433.

[27] T. Senba and T. Suzuki, *Parabolic system of chemotaxis; blowup in a finite and in the infinite time*, Meth. Appl. Anal. 8 (2001) 349-368.

[28] W. Jäger, S. Luckhaus, On explosions of solutions to a system of partial differential equations modelling chemotaxis, Trans. Amer. Math. Soc. 329 (1992) 819-824.

[29] T. Nagai, *Blow-up of radially* symmetric solutions to a chemotaxis system, Adv. Math. Sci. Appl. 5 (1995) 581-601.

[30] T. Suzuki, Exclusion of boundary blowup for 2D chemotaxis system provided with Dirichlet condition for the Poisson part, J. Math. Pure Appl. 100 (2013) 347-367. [31] T. Suzuki, *Semilinear Elliptic Equations - Classical and Modern Theories*, De Gruyter, Berlin, 2020.

[32] B. Gidas, W.M. Ni, L. Nirenberg, Symmetry and related properties via the maximum principle, Comm. Math. Phys.
68 (1979) 209-243.

[33] N.D. Alikakos,  $L^p$  bounds of solutions of reaction-diffusion equations, Comm. Partial Differential Equations 4 (1979) 827-868.

[34] T. Senba and T. Suzuki, *Chemotactic collapse in a parabolic-elliptic system of mathematical biology*, Adv. Differential Equations 6 (2001) 21-50.

