

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Chapter

Use of Cell Biology to Identify Cellular Targets in Drug Development Process against *Leishmania* Sp.

*Gabrielle dos Santos da Silva e Miranda,
Joseane Lima Prado Godinho, Sara Teixeira de Macedo-Silva,
Brunno Renato Farias Verçoza, Alisson Amaral da Rocha,
Milena Barenco Pires de Abreu Sodré,
Victor Feliciano dos Santos Ramos
and Juliany Cola Fernandes Rodrigues*

Abstract

Leishmaniasis is one of the most important neglected tropical diseases. The chemotherapy for its treatment uses very toxic compounds with a low efficacy rate. Thus, there is an urgent need to develop new chemotherapeutic agents to help countries control this devastating disease. In drug development, different approaches can be used to identify potential cellular targets that allow us to understand better the cell biology of eukaryotic cells. Several groups are dedicated to studying new molecules, searching for promising candidates against *Leishmania*. Different techniques have been used to characterize the cell biology, biochemistry, and molecular biology alterations induced by the treatments, trying to understand the mechanisms of action. The main goal of this chapter is to describe an overview of the literature exploring the several studies published about the chemotherapy of anti-*Leishmania* concerning the mechanisms of action of different classes of molecules or therapeutic alternatives.

Keywords: chemotherapy, drug development, cell biology, ergosterol, histone deacetylases, organometallic compounds, therapeutic combination, nanotechnology

1. Introduction

Leishmaniasis is a neglected tropical disease that comprises a large and complex group of infections caused by the *Leishmania* genus protozoan parasites. *Leishmania* parasites have intrinsic biological features that make chemotherapy challenging, presenting high adaptability and plasticity. Its life cycle has two different developmental stages: intracellular amastigotes that live in the mammalian host cells and promastigotes that develop in the insect vectors (**Figure 1**) [1].




Figure 1.
Illustration of the amastigote (left) and promastigote (right) forms of *Leishmania* sp. showing the main organelles and structures.

The ultrastructure of *Leishmania* parasites (**Figure 2**) presents some conserved features and a classical internal organization of eukaryotic cells, with an individualized nuclear envelope, a single and ramified mitochondrion, endoplasmic reticulum, and Golgi complex, which can lead to difficulties in the development of *Leishmania*-specific drugs with low toxic side effects to mammalian hosts [1]. In addition, however, these protozoans have essential and exclusive organelles and structures such as acidocalcisomes, glycosomes, megasomes, and subpellicular microtubules (**Figure 1**) that can be exploited as drug targets [2].

The *Leishmania* plasma membrane comprises a lipid bilayer associated with proteins and a glycocalyx consisting of a myriad of glycoconjugates. The lipid bilayer has a trilaminar aspect with about 9 nm of thickness. The lipidic composition of the Trypanosomatidae family members is dependent on genus and species. In general, *Leishmania* has 24-methylated sterols, such as episterol, 5-dehydroepisterol, and traces of ergosterol as the major endogenous sterols' constituents, which are absent in mammalian host cells, where cholesterol is the main source of membrane sterols [3]. This divergence in sterol profiles has been exploited to develop drugs that affect the sterols biosynthesis pathway, including azoles,




Figure 2.
Ultrathin sections of *L. amazonensis* amastigotes (A) and promastigote (B). F, flagellum; HC, host cell; K, kinetoplast; LB, lipid body; M, mitochondrion; N, nucleus.

azasterols, and others [2]. In addition, many attempts to develop drugs targeting *Leishmania* glycolyx have also been performed [4].

Leishmania parasites present a single and ramified mitochondrion frequently associated with the plasma membrane, subpellicular microtubules, and endoplasmic reticulum. As well as in other eukaryotes, *Leishmania* mitochondrion operates in energetic metabolism, compartmentalizing the Krebs cycle and performing cellular respiration. The amphotericin B and pentamidine, two of the currently used drugs for the treatment of leishmaniasis, target mitochondria, resulting in a decrease of the mitochondrial membrane potential. Moreover, other drugs targeting mitochondria, such as hydroxynaphthoquinones, have been evaluated against *Leishmania* [5].

To maintain its morphological structure, *Leishmania* has a cytoskeleton composed mainly of subpellicular microtubules, which are filaments finely associated to plasma membrane inner leaflet, regularly spaced, and longitudinally oriented throughout the parasite's cellular body. Despite the phylogenetic conservation of α and β -tubulins, structural divergences in specific tubulin drug binding sites have suggested these proteins as a potential target, as described for podophyllotoxin derivatives and others [6, 7].

In the Medicinal Chemistry field, several approaches have been attempted, trying to find potential cellular targets for developing anti-*Leishmania* drugs. First, nanotechnology-based drug delivery systems have been applied, improving efficacy and enhancing pharmacokinetics properties of currently available drugs [8]. Second, molecular hybridization techniques can combine two drugs or chemical groups with previously known biological activity, producing a single and novel molecule with increased activity [9]. Finally, another alternative is the transposition of a drug already used to treat another disease; it is the case of miltefosine, the last treatment included in the arsenal of chemotherapeutic agents against leishmaniasis. Miltefosine was initially designed as an anticancer medicine, and in 2002 it was registered as first-line treatment, mainly for visceral leishmaniasis (VL), in Asia, Africa, and some regions of Europe.

2. Challenging the target: phospholipid and ergosterol biosynthesis

The first metabolically stable analogs derived from lysophosphatidylcholine were synthesized in the late 1960s. Two decades later, Eibl and Unger synthesized the first alkyl phospholipids (APLs), also called miltefosine, first administrated by an intravenous route to treat systemic tumors [10]. However, the treatment failed, and miltefosine was evaluated against topical cutaneous metastases from breast cancer [11]. In the late 1990s, miltefosine was assessed *in vitro* against the *Leishmania* genus [12, 13] and in murine models infected with *L. donovani* and *L. infantum* [14]. More recently, miltefosine was also evaluated against murine models of cutaneous leishmaniasis, alone or in combination with paromomycin [15–17].

After the evidence of the excellent anti-*Leishmania* activity of miltefosine demonstrated *in vitro* and *in vivo*, clinical tests began to be carried out immediately. Thus, miltefosine was the first oral drug approved for the treatment of visceral leishmaniasis, and for several years it was used as the first choice for the treatment of visceral leishmaniasis (VL), mainly in India [13, 18].

Miltefosine was also evaluated in patients infected with cutaneous leishmaniasis in Colombia, Guatemala, and Brazil [19, 20]. The efficacy in this clinical manifestation was variable, depending on the species. For patients infected with *L. panamensis* in Colombia, the cure rate was 91%, while in Guatemala, the cure rate was 53% for infections with *L. braziliensis* and *L. mexicana* [19]. Furthermore, in Brazil, in patients infected with *L. braziliensis*, the cure rate was 75%, compared to 53% cure achieved with pentavalent antimonials [20].

Several studies have demonstrated that the primary target of miltefosine is the cell membranes, affecting cellular processes such as signal transduction, lipid metabolism, and calcium homeostasis [21]. The selectivity for the plasma membrane is related to its chemical structure formed by a polar choline head bound to a long non-polar hydrocarbon chain, which easily inserts into the lipid bilayer, presenting detergent properties that lead to cell lysis in high concentrations [10]. In *L. donovani* and *T. cruzi*, miltefosine inhibits the phosphatidylcholine biosynthesis pathway (Greenberg pathway), being more selective for the protozoan parasites than mammalian cells, where the main route for phospholipid synthesis is the Kennedy pathway [22, 23]. Miltefosine also interfered with the ergosterol biosynthesis and promoted a disturbance in GPI synthesis [21, 24–26], leading to membrane permeability and fluidity changes. In addition, miltefosine interfered with the host immune response by inducing the production of interferon γ cytokine, leading to a biased immune response towards Th1, which would be a beneficial outcome for immunosuppressed patients [26, 27]. About the ultrastructure of *Leishmania* treated with miltefosine (**Figure 3A**), some studies revealed that it induced several alterations, mainly observed in the mitochondrion, in the plasma membrane, an increase of autophagic structures and phenotypes related to cell cycle arrest and apoptosis-like cell death [26, 28, 29].

With the success of miltefosine, several groups worldwide began to study new chemical routes to synthesize ether phospholipid derivatives in searching for novel molecules more active and selective against *Leishmania* [30]. Our group studied a novel hybrid derivative called alkyl phosphocholine-dinitroaniline, which presented a potent effect against *L. amazonensis* at least around 15-times better than miltefosine [31].

Another essential metabolic route for *Leishmania* and other protozoan parasites is the biosynthesis of ergosterol (or 24-methyl sterols) [32]. *Leishmania* has in these cell membranes three major sterol components that are absent in mammalian cells, 5-dehydroepisterol, episterol, and ergosterol. In mammals, cholesterol is the principal sterol present in cell membranes. Thus, the differences between some steps and enzymes in the biosynthetic route of the protozoan parasites and mammalian host cells have been exploited as targets to develop novel drugs as candidates to chemotherapeutic agents [32].

At least 20 metabolic steps are necessary to synthesize ergosterol, and several enzymes participate in these reaction sequences [32–34]. Furthermore, several works have shown that multiple classes of compounds targeting 24-methyl sterol biosynthesis exhibit suitable anti-trypanosomatid activities *in vitro* (**Figure 3B, C**) and *in vivo* [33–39].




Figure 3. Ultrathin sections of *L. amazonensis* promastigotes treated with miltefosine and two ergosterol biosynthesis inhibitors. (A) 30 μ M miltefosine for 72 h; (B) 3 μ M ravuconazole for 48 h; (C) 1 μ M posaconazole for 48 h. Mitochondrion was the organelle more affected in all treatments, presenting alterations in the cristae and kDNA structure (asterisk, thin arrow, and arrowheads) and mitochondrial swelling and disorganization of its ultrastructure. In panel C, several lipid bodies appeared after treatment. K, kinetoplast; LB, lipid body; M, mitochondrion; N, nucleus.

More than 30 drugs have been studied in the last 30 years. These drugs are included in large classes of inhibitors, such as 1) Statins, which inhibit the enzyme 3-hydroxy-3-methyl-glutaryl-CoA reductase (HMG-CoA reductase), also evaluated as cholesterol-lowering drugs; 2) Bisphosphonates that act in the enzyme farnesyl pyrophosphate synthase and inhibit the isoprenoid pathway, mainly used to treat hypercalcemia; 3) Quinuclidines and zaragozic acid, developed to inhibit the squalene synthase, the enzyme that catalyzes the first committed step in the sterol biosynthesis pathway. This class of drugs was developed as an alternative to statin use because they do not inhibit the synthesis of the isoprenoids. 4) Allylamines, which include the known antifungal terbinafine that inhibits the squalene epoxidase; 5) Azoles, which are essential medicines to treat many fungal diseases and inhibit the C14 α -demethylase. Several azoles were developed, always trying to find new tolerate and efficacy drugs, also searching to novel molecules to solve the problems with antifungal resistance; finally, 6) Azasterols, which inhibit the enzyme $\Delta^{24,25}$ sterol methyltransferase absent in mammalian cells, one enzyme that catalyzes the methylation of steroid nucleus of sterols, producing 24-methyl sterols, essential for *Leishmania*, *T. cruzi* and fungi (Reviewed in [32]).

In summary, several works have pointed to the importance of looking for the biochemical properties of each enzyme involved in the pathway and its relevance as an essential target for the parasite viability; this feature characterizes the enzyme as a promising target for the development of potential chemotherapeutic candidates for the treatment of leishmaniasis.

3. Challenging the target: histone deacetylases

Histone deacetylase (HDACs) inhibitors are a relatively new class of potential agents in treating neurodegenerative diseases, various types of cancer, and parasitic infections. HDACs have broad importance in the cellular environment. They regulate histone and non-histone proteins affecting the cell cycle, energy metabolism, and inducing cell death. Some HDAC inhibitors were already approved by the FDA (Food and Drug Administration) to treat lymphoma and myeloma, such as vorinostat, romidepsin, belinostat, and panobinostat, in combination with bortezomib and dexamethasone [40]. Given the results obtained *in vitro* and *in vivo* in several disease models, the advancement of clinical trials in tumors, and the transposition of drugs as an old ally in the treatment of leishmaniasis, HDAC inhibitors are a promising approach in the understanding of cell biology of the parasite, especially concerning its chemotherapy.

There are 18 histone deacetylases in humans, which can be grouped according to cell location and the molecule used as a cofactor for its enzymatic action. These HDACs are divided into 1) zinc-dependent HDACs, also called “classical” histone deacetylases; and 2) nicotinamide and adenine dinucleotide [NAD⁺]-dependent HDACs. The first one comprises class I (HDACs 1–3, 8), IIa (HDACs 4, 5, 7, and 9), IIb (HDACs 6 and 10), and IV (HDAC 11). While the second one, the NAD⁺-HDAC, belongs to class III and is also known as sirtuins (SIRT 1–7). HDACs are still poorly understood and characterized in *Leishmania*. So far, four Zn²⁺-dependent histones deacetylases and three NAD⁺-dependent histones deacetylases were discovered in the parasite [41, 42]. Although four homologs of classical HDACs were identified, none was functionally characterized. Among the information in the literature, a *L. major* HDAC (gene LmjF21.0680) was shown to be expressed during the differentiation of promastigotes to amastigotes, with a possible role in chromatin structure and impacts on gene transcription [43]. Furthermore, Prasanna et al. [44] managed to isolate, express, and purify an *L. donovani* histone deacetylase

(LD_HDAC), with less than 40% identity with class I human HDACs. Information about classical HDACs in the *Leishmania* genus is still very early and deserves further development.

Unlike classical HDACs, there are several studies about NAD⁺-dependent HDACs in the *Leishmania* genus. Three sirtuins were already found in *Leishmania*, SIR2RP1, SIR2RP2, and SIR2RP3 [45]. These sirtuins were not found in the nucleus, as described for *Saccharomyces cerevisiae* [45]. SIR2RP1 is expressed from a single copy gene in *L. amazonensis* (LaSIR2RP1), resulting in a monomeric protein with NAD⁺-dependent deacetylase action immunodetected in its glycosylated form [46]. LaSIR2RP1 has dispersed localization in the cytosol or cytoplasmic granules [47] and is secreted in lesions derived from intracellular amastigotes [48]. In addition, the *L. donovani* SIR2RP1 was 46% similar to the human SIRT2 [49]. In *L. infantum* and *L. major*, the SIR2RP1 proteins present two functional sites for NAD⁺-dependent deacetylation activity and ADP-ribosyl transferase activity [49].

In *L. major* SIR2RP1, removing the acetyl group of lysine 40 from α -tubulin was demonstrated *in vitro* and *in vivo* [50]. However, in these parasites, the ribosylation function of α -tubulin ADP was also shown, resulting in its depolymerization or even inhibiting its assembly [51]. Thus, sirtuins in *Leishmania* have demonstrated an essential role in cytoskeleton dynamics and may have significant implications for remodeling the parasite's morphology and its interaction with the host cell. LmSIR2RP1 also showed a close relationship with the HSP83 protein, an orthologous chaperone of the human HSP90 chaperone, although the intracellular levels of LmSIR2RP1 do not influence the acetylation status of HSP83 [52]. The use of geldanamycin, an HSP90 inhibitor, induced alterations in the cytodifferentiation of promastigotes to intracellular amastigotes [53]; the same was observed for protozoa overexpressing or knockout for the LmSIR2RP1 gene [52]. This study confirms that the SIR2RP1/HSP83 interaction may play an essential role in the differentiation of the parasite.

The work in [54] demonstrated that *L. major* sirtuins may be related to the success of infection through interaction with macrophage surface proteins and, therefore, play a regulatory role in immune responses [55]. This work also showed the capacity of *L. major* sirtuins to trigger the effector response of B cells, promoting a robust humoral response with the secretion of specific antibodies, such as IgG1 and IgG2a [55]. Meanwhile, the overexpression of *L. major* sirtuin, also observed for *L. infantum* sirtuin [48], revealed that it would be involved in the proliferation rate, besides participating in the regulation of death factors, thus preventing death by apoptosis.

The SIR2RP2 present in *L. infantum* mitochondrion is related to the growth rates of promastigote forms with a direct relationship with NAD⁺ homeostasis [56]. In 2017, a study revealed that the *L. donovani* SIR2RP2 also has a mitochondrial location [49]. The deletion of this gene led to a reduction in the proliferation rate, further resulting in the interruption of the cell cycle in the G2/M phase. Furthermore, the deletion of LdSIR2RP2 resulted in greater susceptibility of the parasite to commercially available sirtuin inhibitors and a reduction in the mitochondrial membrane potential, resulting in a low concentration of ATP in the mitochondrion. Interestingly, in this SIR2RP2 knockout *Leishmania*, there was an intensification of glycolysis that could be an attempt to compensate for the disbalance of the mitochondrial metabolism. LdSIR2RP2 showed NAD⁺-dependent ADP-ribosyl transferase activity with 39% similarity to SIRT4 from humans [49].

The third sirtuin, SIR2RP3, still has few descriptions in the literature. The *L. donovani* SIR2RP3 was 37% similar to the human SIRT5 [49], so that molecular coupling assays, using known inhibitors and SIR2RP3, revealed a strong analogy with SIRT5, which refers to the form of interaction with inhibitors. However, the

interaction of these inhibitors with SIR2RP3 also demonstrated significant molecular differences compared to SIRT5 in humans, which could act as selective targets for the treatment of leishmaniasis [57].

A recent study with a histone deacetylase inhibitor in *L. amazonensis* revealed the sirtuins' potential to develop novel molecules with anti-*Leishmania* activity [58]. Furthermore, this study from our group demonstrated the potent inhibition of parasite proliferation that is probably related to essential functions for HDACs in *Leishmania*, which include the control of the cell cycle and the induction of cell death [58]. Other effects already observed with HDAC inhibitors are different levels of chromatin compaction, increased number of lipid bodies randomly distributed throughout the cytosol, increased production of reactive oxygen species, changes in *Leishmania* morphology, and increased expression of acetylated α -tubulin (**Figure 4**) [58].

However, the parasite's ability to modulate the histone deacetylases of the mammalian host to establish the infection has already been observed. For example, the upregulation of the macrophage HDAC1 was observed during infection with *L. amazonensis*, resulting in deacetylation of the histone tail of the gene's promoter region responsible for producing nitric oxide. This deacetylation prevents the access of transcription factors, which culminates in the repression of nitric oxide production, which allows the establishment of intracellular amastigote forms in the parasitophorous vacuole [59].

Besides, HDAC inhibitors have also been used in combination therapy to treat antimony-resistant *L. donovani* infection, where the upregulation of the multiresistant protein depends on IL-10 production. Thus, the imipramine antidepressant positively regulates HDAC11, inhibiting the acetylation of the IL-10 promoter, which leads to a decrease in its production [60]. This imipramine-mediated lowering of the IL-10 level reduces MDR-1 expression and aids in the elimination of the parasite.

Thus, histone deacetylase inhibitors belong to a class of compounds with potential application to develop novel molecules with anti-*Leishmania* activity and for the treatment of leishmaniasis, alone or in combination with other medications already established. Furthermore, the World Health Organization (WHO) has already recommended the therapeutic combination to reduce the doses and toxicity and develop a more effective and safe treatment. Finally, HDAC inhibitors seem to be an exciting tool for a better understanding of the Cell Biology of *Leishmania* and enabling the knowledge of new routes for the development of novel drug candidates.




Figure 4.
 Ultrathin sections of *L. amazonensis* promastigotes treated with HDAC inhibitors for 48 h. (A) 1.5 μ M NIH119; (B) 1 μ M NIH119; (C) 15 μ M tubastatina. A. NIH119 and tubastatin induced different ultrastructural alterations such as 1) increase in the number of lipid bodies, some of them presenting different morphologies (A, B); 2) presence of membrane protrusions (A, B – Arrow); 3) alterations in the kinetoplast (A); 4) decondensation of chromatin (B, C); and, 5) changes in mitochondrial ultrastructure. FP, flagellar pocket; K, kinetoplast; LB, lipid body; M, mitochondrion; N, nucleus.

4. Challenging the target: organometallic compounds

Although metals have been used in Medicine for centuries, most compounds produced by the pharmaceutical industry are still based on organic molecules. Nevertheless, new perspectives about metal-based drugs and their therapeutic potential against cancer, bacteria, virus, and even trypanosomatid infections have emerged in the last few years. In this context, *Leishmania* infections have been treated with pentavalent antimonials since the 1940s [1], which indicates that new compounds containing other kinds of metals may present anti-*Leishmania* activity, thus opening the possibility to reduce the toxicity based on the metal-drug synergism.

Platinum-derived metals are well known to have antitumor effects due to their ability to bind to DNA molecules. So, since tumor cells and kinetoplastid parasites present similar metabolic pathways [61], the coordination of these metals to organic compounds might be efficient in treating *Leishmania* infections. For example, the (2,2':6'2"-terpyridine)platinum(II) complexes can inhibit 100% of the growth of *L. donovani* amastigotes at a concentration of 1 μM through the intercalation of terpyridine and platinum(II) into the DNA, probably binding to guanine bases or some enzyme active sites [62]. More recently, cisplatin-derived complexes (*cis*-diamminedichloroplatinum(II)), an anticancer drug, were tested on *L. infantum* promastigotes and amastigotes, revealing a remarkable anti-*Leishmania* activity with IC_{50} values of 1.03 μM and 0.10 μM , respectively. Furthermore, the treatment with *cis*-DDP induced loss of mitochondrial transmembrane potential and DNA fragmentation, thus leading to apoptosis-like death [63].

In addition to platinum and its derivatives, other transition metals have been widely studied in terms of antiprotozoal activity. For example, organometallic complexes containing ruthenium(II) and anti-inflammatories were evaluated active against *L. amazonensis* and *L. infantum* promastigotes, presenting IC_{50} values comparable to the meglumine antimoniate, one of the first-line drugs for *Leishmania* infections [64]. Moreover, the coordination of gold compounds with organic ligands was efficient against *L. amazonensis* and *L. braziliensis* promastigotes while presenting low toxicity to host cells [65]. Thus, the therapeutic mechanism of these organometallic molecules may be related to induction of oxidative damage and alterations in the membrane permeabilization by the inhibition of specific membrane protein channels and zinc-binding proteins. These alterations can lead to parasite cell death by apoptosis-like and necrosis.

Among transition metals, the essential ones, such as zinc and copper, are present in cell structures and involved in many cellular processes, becoming indispensable for host-parasite interactions. Zinc regulates gene transcription processes and cell signaling, while copper is also a critical enzymatic cofactor for organ functioning and multiple metabolic processes [66]. Therefore, the coordination of organic molecules to essential metals regarding new antiprotozoal treatments might increase the drug uptake and contribute to the parasite's elimination. The zinc(II)-dipicolylamine (ZnDPA) coordination complexes were active against *L. major* promastigotes *in vitro* with IC_{50} values between 12.7 μM and 0.3 μM and minimal mammalian cell toxicity. The compounds also showed *in vivo* activity, with a high affinity for intracellular amastigotes and low toxicity to mice [67]. The effects of a copper dimethoxy bipyridine (CuDMOBP) complex were also investigated against *L. major*; the complex presented significant *in vitro* activity with high selectivity index [68]. Results from quantitative real-time PCR also indicate a significant reduction in cellular expression of IL-10 and TNF- α in macrophages treated with CuDMOBP, probably due to a reduction of the parasite population.

Metals have also been combined with ergosterol biosynthesis inhibitors, including the azoles family of drugs, which are used to treat fungal infections and present activity against protozoan parasites. For example, a recent study from our group revealed the potent effect of the combination of itraconazole (ITZ) with zinc (Zn) against *L. amazonensis* promastigotes (**Figure 5B**) and intracellular amastigotes [69]. The biological effects were significantly increased when the itraconazole was coupled with zinc, resulting in IC₅₀ values in nanomolar ranges and cell death of parasites in low concentrations [69]. Likewise, the coordination of clotrimazole, ketoconazole (**Figure 5A**), and miconazole to manganese (Mn) provide novel molecules with better activity against *L. major* when compared to the original organic antifungals [70].

Despite Medicine's advances in the past decades, information about organometallic drugs is still lacking. More profound studies must be done to understand the role of metals in host–parasite interactions, thereby better comprising the mechanisms of organometallics drugs against the parasites and mammalian host cells. Nevertheless, the literature available indicates that organometallics are a promising class of drugs for treating leishmaniasis.




Figure 5.
 Ultrathin sections of *L. amazonensis* promastigotes treated with organometallic compounds. (A) Treatment with 300 nM ketoconazole-ruthenium induced alterations in the mitochondria, such as swelling and loss of the mitochondrial matrix and the formation of large autophagic vacuoles. (B) Treatment with 0.5 μM itraconazole-zinc resulted in mitochondrial alterations and an increase in the secretion of vesicles (arrowhead) at the flagellar pocket. In both panels, the nuclear chromatin appeared altered. F, flagellum; M, mitochondrion; N, nucleus.

5. Therapeutic combination: what do we know?

There are many strategies to treat leishmaniasis; however, several studies have shown the numerous advantages of therapeutic combination, like observing for other diseases. For example, combining drugs from different chemical classes could reduce the total drug doses or treatment duration. These aspects are important to minimize toxic side effects, submission at treatment, less load on the public health system and reduce cases of drug resistance. Also, the therapeutic combination could improve treatment efficacy for refractory or complicated cases, such as in patients coinfecting with HIV. A successful study conducted by the Drugs for Neglected Diseases initiative (DNDi), in partnership with Médecins Sans Frontières (MSF) and other institutions, pointed to evidence of the high efficacy of the combination therapy to treat patients with visceral leishmaniasis (VL) in coinfection with HIV [71]. Although the current WHO guidelines recommend the treatment of HIV/VL coinfection with liposomal amphotericin B (AmBisome®), this work strongly supports a change in the treatment recommendations, from AmBisome monotherapy to

combination therapy as the first-line treatment. Moreover, they suggest the combination with miltefosine once this combination therapy has a good safety profile and is highly efficacious [71].

Nowadays, combination therapy is an efficient tool to treat many microbial infections such as AIDS, tuberculosis, malaria, and several other diseases. Recent works have shown that combination therapy for leishmaniasis has progressively been recommended to increase treatment tolerance and efficacy, reduce cost and treatment duration, and limit the growth of drug resistance [72–75]. For the treatment of leishmaniasis, WHO has recommended combination therapy based on many studies showing the efficacy of this therapeutic tool; the combinations include novel synthesized drugs, nanoparticles developed for drug delivery, repositioned drugs, old medicines, and immunomodulatory agents [76–79]. Indeed, several studies have reported the superior efficacies of combination therapies against leishmaniasis. Some of them demonstrated the synergic effects of combinations between amphotericin B with other available medicines, such as meglumine antimoniate, miltefosine, paromomycin, or azithromycin [80–82].

Analysis of drug interactions aimed to show if the interaction between them is classified as synergistic, antagonistic, or indifferent. *In vitro* data are based on an extended ratio and concentration range. However, *in vivo* combinations are more complex and less defined, with the number of doses limited. The synergy between two (or more drugs) occurs when their combined activities are improved over the sum of their separate individual effects. Synergistic drug combinations provide lower concentrations of both compounds, enhancing therapy outcomes by increasing efficacy and reducing side effects. Moreover, synergistic combinations could reintroduce those that have lost activity against drug-resistant strains. The FICI (fractional inhibitory concentration index) value is considered the standard reference parameter to quantify interactions between pairs of drugs. Odds in 2003 [83] established more restrictive criteria to analyze experiments and defined “synergy” as a ≥ 4 -fold reduction in the MICs of both compounds in combination when compared to their MICs alone, where the FICI value must be ≤ 0.5 . MIC means the minimum inhibitory concentration, i.e., the lowest drug concentration with no visible cell growth.

In vitro studies against *L. amazonensis* suggest that combinations between compounds that act in different biosynthetic pathways of the parasite, such as sterol biosynthesis, are promising [32]. Interestingly, a recent study showed that sterol biosynthesis inhibitors and alkylphosphocholine analogs, combined with medicines available to treat other diseases, are efficient against trypanosomatids [32, 36, 78, 84]. Besides, another study showed the *in vivo* efficacy of the combination therapy between miltefosine, an alkylphosphocholine, and amphotericin B or paromomycin [85], a therapeutic alternative to treat antimony-resistant VL cases in India. Furthermore, topical treatment against cutaneous leishmaniasis might be effective when amphotericin B and miltefosine are co-loaded at second-generation ultra-deformable liposomes since *in vivo* studies displayed a significant reduction in the parasitic burden [86]. However, a Brazilian survey against *L. infantum* revealed a decrease of the miltefosine concentration when combined with lopinavir (anti-HIV drug); yet, the synergistic effect was not evidenced [87]. Unexpectedly, the combination of nelfinavir with miltefosine presented better results, with FICI ≤ 0.5 . Thus, this study also concluded that the combination might be helpful to treat patients with visceral leishmaniasis who also are infected with HIV [88].

Combination therapy is a promising strategy to treat several diseases. Therefore, it is urgent to investigate synergistic and other drug combinations to increase novel probabilities of therapeutic protocols to treat leishmaniasis. The discovery and the analysis of drug combinations can be facilitated by the collective use of different

approaches and methods. Drug combinations have proved to be a successful strategy to shorten the course of therapy and reduce toxicity through lower dosage administration; these strategies should reduce the appearance of new resistant parasites. Thus, recent proposals of combinations have been suggested as state-of-the-art for the treatment of leishmaniasis. In the short run, combination therapy is an interesting way to improve the treatment for leishmaniasis.

6. Where are we going? Nanotechnology

Recent advances in Nanotechnology have had a profound impact on health sciences, especially Medicine, because of the development of different nanomaterials designed as intracellular carriers to deliver drugs and genes. The development and use of nanocarriers have also been established in the field of Pharmaceutical Sciences by enabling the encapsulation of drugs creating stable and controlled environments, and improving the biocompatibility of these drugs in various biological systems. These nanocarriers were developed to improve the solubility of poorly soluble drugs, control or maintain their release, and protect them from degradation. These characteristics increase drug bioavailability, reduce systemic side effects, and increase drug specificity for biological targets. For these reasons, Nanotechnology is a new field that allows the construction of versatile diagnostic and therapeutic platforms using nanocarriers as molecular machinery for different clinical applications.

The development of nanocarriers began in the 1960s to always seek to improve biocompatibility and reduce the toxicity of nanomaterials. The second generation of nanocarriers was developed around the 1980s and sought to improve the surface of these materials by increasing their stability, stealth, and targeting ability. The third generation of nanocarrier introduced the idea of intelligent nanomedication to enhance the targeting mechanisms and theranostic capabilities of these nanomaterials [89]. The word *nanoparticle* has been widely applied to describe numerous pharmaceutical carriers or imaging systems based on nanoscale materials. Nanoparticles are particulate materials in their solid or dispersed state present in a size scale between 10 nm and 100 nm (ISO/TS 80.004–1:2015). Due to the great diversity of these nanomaterials, the scientific community elaborates a classification based on their characteristics and properties.

Initially, the nanomaterials are divided into two classes: inorganic and organic. However, they are divided into three subgroups, according to some of their characteristics. The first would concentrate single-chain polymer-drug conjugates, polymeric colloids prepared by techniques such as emulsion polymerization, cross-linked nanogel matrices, dendrimers, and carbon nanotubes, where the nanocarrier is a single synthetic molecule with covalent bonds and a relatively large molar mass. The second subgroup of nanocarriers would comprise self-assembly of smaller molecules such as 1) liposomes and polyplexes, the most studied members of this group of nanoparticles; 2) polymersomes and other sets of block copolymers; 3) colloidosomal aggregates of latex particles and sets of proteins or peptides. In this case, the dynamic nature of these types of systems depends on intermolecular forces and biological conditions. Finally, the last subgroup of nanocarriers would include complexes based on fullerenes, silica, colloidal gold, gold nanoshells, quantum dots, and superparamagnetic particles.

Another critical point in developing nanocarriers is the synthesis, which can be rationally divided into two fundamental stages: nucleation and growth. Understanding and manipulating these two steps have created new possibilities allowing researchers to easily control the synthesis of nanoparticles in terms of

size, morphology, and monodispersity. The choice of the synthesis route provides a characteristic set of advantages and disadvantages in nanomaterial production.

In *Leishmania* sp., the use of nanocarriers has been studied since the late 1970s with the development of liposomal amphotericin B [90, 91]. Since its development, liposomal amphotericin B has become more efficient and bioavailable, less toxic, and better tolerated by patients [92, 93]. This formulation also has a broad and rapid biodistribution reaching steady-state plasma concentrations faster with higher total plasma concentration when compared to its deoxycholate form. Furthermore, liposomal amphotericin B is probably inactive because it is fixed to the liposome; thus, the biologically active drug is released only after direct contact with protozoa or fungal cell walls [94]. In 2010, the WHO proposed the administration of liposomal amphotericin B.

The success achieved by the liposomal formulation of amphotericin B is related to its properties as a nanocarrier system, which has numerous advantages. However, despite these advantages, this system has some disadvantages, including its high cost, limiting its use [94]. Thus, the development of new, cheaper, and more efficient nanoformulations is necessary. Furthermore, different nanocarriers have been developed in recent decades, searching for new therapeutic alternatives to treat leishmaniasis, including nanoparticles based on liposomal, lipid, polymeric, and metallic nanomaterials [8, 95–98]. Therefore, choosing the correct nanocarriers is crucial to define properties and characteristics for this proposed new therapeutic approach. Thereby, this enormous diversity of available nanoparticles makes the development of nanocarriers for the treatment of leishmaniasis very promising since each of these carrier systems has advantages and limitations over each other [8].

Some nanoparticles have been generating significant repercussions for presenting theranostic properties, thus allowing them to be used simultaneously for diagnosis and therapy. Superparamagnetic iron oxide nanoparticles (SPIONs) are one example of this type of nanomaterials. SPIONs have excellent biocompatibility, degradability in moderate acid conditions, magnetic properties, and the ability to generate heat when subjected to an alternating current magnetic field [99, 100]. In addition, this type of nanomaterial still has a large surface area presenting great chemical diversity, which can increase the efficacy of the treatment. Finally, these nanomaterials can also be conjugated to specific molecules to facilitate selective and efficient drug delivery to a diseased tissue or organ [101]. The application of this type of nanomaterial for the treatment of leishmaniasis has been studied by different groups and has shown promising results over the past few years (Figure 6) [102–105].




Figure 6. Ultrathin sections of *L. amazonensis* promastigotes (A) and amastigotes (B) after treatment with 100 µg/mL SPIONs for 24 h. In both developmental stages, SPIONs were found inside membrane-bound structures (arrowheads). Some alterations were also observed in the treated parasites, such as increased number of lipid bodies (arrows), presence of autophagosome (thin arrows), and secretion of extracellular vesicles (large arrow).

In summary, Nanotechnology and the use of nanoparticles inaugurated a new field in health science called Nanomedicine, one of the most promising branches of contemporary Medicine. Thus, the development of new nanomaterials to treat leishmaniasis significantly increases the possibility of finding novel therapeutic alternatives, mainly considering the great diversity of clinical manifestations. An excellent example of this is the use of nanoparticles to develop a topical treatment that can revolutionize the treatment of cutaneous leishmaniasis.

7. Conclusions and perspectives

The treatment of infectious diseases depends on a better understanding of Cell Biology, mainly for parasites that are intracellular obligate eukaryotes, such as *Leishmania*. The knowledge about these parasites allows the identification of essential metabolic pathways and the mechanisms of action involved in drug inhibition, enabling the application of new treatments and conventional treatments combined with new therapies, also using Nanotechnology. Thus, the improvement of tested molecules associated with drug delivery techniques is the path to success for novel leishmaniasis treatments.

Acknowledgements

We acknowledge the Brazilian funding agencies Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), Conselho Nacional de Desenvolvimento e Pesquisa (CNPq), and Fundação Carlos Chagas Filho de Amparo à Pesquisa do Estado do Rio de Janeiro (FAPERJ) for financial support.

Author details

Gabrielle dos Santos da Silva e Miranda¹, Joseane Lima Prado Godinho^{1,2}, Sara Teixeira de Macedo-Silva³, Brunno Renato Farias Verçoza¹, Alisson Amaral da Rocha¹, Milena Barenco Pires de Abreu Sodré¹, Victor Feliciano dos Santos Ramos¹ and Juliany Cola Fernandes Rodrigues^{1*}

¹ Federal University of Rio de Janeiro, Campus UFRJ Duque de Caxias Prof. Geraldo Cidade, Duque de Caxias, Brazil

² Graduate Program in Public Health, Federal University of Maranhão, Maranhão, Brazil

³ CENABIO - Advanced Microscopy Unit, Federal University of Rio de Janeiro, Rio de Janeiro, Brazil

*Address all correspondence to: juliany.rodriques@xerem.ufrj.br

IntechOpen

© 2021 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

References

- [1] Rodrigues JCF, Godinho JLP, de Souza W. Biology of human pathogenic trypanosomatids: Epidemiology, lifecycle and ultrastructure. *Subcellular Biochemistry*. 2014;**74**:1-42
- [2] Chawla B, Madhubala R. Drug targets in *Leishmania*. *Journal of Parasitic Diseases*. 2010;**34**(1):1-13
- [3] de Souza W, Rodrigues JCF. Sterol biosynthesis pathway as target for anti-trypanosomatid drugs. *Interdisciplinary Perspectives on Infectious Diseases*. 2009;**2009**:1-19
- [4] Cabezas Y, Legentil L, Robert-Gangneux F, Daligault F, Belaz S, Nugier-Chauvin C, et al. *Leishmania* cell wall as a potent target for antiparasitic drugs. A focus on the glycoconjugates. *Organic and Biomolecular Chemistry*. 2015;**13**(31): 8393-8404
- [5] Fidalgo LM, Gille L. Mitochondria and trypanosomatids: Targets and drugs. *Pharmaceutical Research*. 2011;**28**(11):2758-2770
- [6] Escudero-Martínez JM, Pérez-Pertejo Y, Reguera RM, Castro MÁ, Rojo MV, Santiago C, et al. Antileishmanial activity and tubulin polymerization inhibition of podophyllotoxin derivatives on *Leishmania infantum*. *International Journal for Parasitology: Drugs and Drug Resistance*. 2017;**7**(3):272-285
- [7] Chatterji BP, Jindal B, Srivastava S, Panda D. Microtubules as antifungal and antiparasitic drug targets. *Expert Opinion on Therapeutic Patents*. 2011;**21**(2):167-186
- [8] Saleem K, Khursheed Z, Hano C, Anjum I, Anjum S. Applications of nanomaterials in leishmaniasis: A focus on recent advances and challenges. *Nanomaterials*. 2019;**9**(12):1-18
- [9] Walsh J, Bell A. Hybrid drugs for malaria. *Current Pharmaceutical Design*. 2009;**15**(25):2970-2985
- [10] van Blitterswijk W, Verheij M. Anticancer alkylphospholipids: mechanisms of action, cellular sensitivity and resistance, and clinical prospects. *Current Pharmaceutical Design*. 2008;**14**(21):2061-2074
- [11] Leonard R, Hardy J, van Tienhoven G, Houston S, Simmonds P, David M, et al. Randomized double-blind, placebo-controlled, multicenter trial of 6% miltefosine solution, a topical chemotherapy in cutaneous metastases from breast cancer. *Journal of Clinical Oncology*. 2001;**19**(21): 4150-4159
- [12] Achterberg V, Gercken G. Cytotoxicity of ester and ether lysophospholipids on *Leishmania donovani* promastigotes. *Molecular and Biochemical Parasitology*. 1987;**23**(2): 117-122
- [13] Croft SL, Neal RA, Pendergast W, Chan JH. The activity of alkyl phosphorylcholines and related derivatives against *Leishmania donovani*. *Biochemical Pharmacology*. 1987;**36**(16):2633-2636
- [14] Kuhlencord A, Maniera T, Eibl H, Unger C. Allgemeine Hygiene I. Hexadecylphosphocholine: Oral treatment of visceral leishmaniasis in mice. *Antimicrobial Agents and Chemotherapy*. 1992;**36**(8):1630-1634
- [15] Aguiar MG, Pereira AMM, Fernandes AP, Ferreira LAM. Reductions in skin and systemic parasite burdens as a combined effect of topical paromomycin and oral miltefosine treatment of mice experimentally infected with *Leishmania (Leishmania) amazonensis*. *Antimicrobial Agents and Chemotherapy*. 2010;**54**(11):4699-4704

- [16] Aguiar MG, Silva DL, Nunan FA, Nunan EA, Fernandes AP, Ferreira LAM. Combined topical paromomycin and oral miltefosine treatment of mice experimentally infected with *Leishmania* (*Leishmania*) *major* leads to reduction in both lesion size and systemic parasite burdens. *Journal of Antimicrobial Chemotherapy*. 2009;**64**(6):1234-1240
- [17] Godinho JLP, Simas-Rodrigues C, Silva R, Ürmenyi TP, de Souza W, Rodrigues JCF. Efficacy of miltefosine treatment in *Leishmania amazonensis*-infected BALB/c mice. *International Journal of Antimicrobial Agents*. 2012;**39**(4):326-331
- [18] Sundar S, Jha TK, Thakur CP, Engel J, Sindermann H, Fischer C, et al. Oral miltefosine for indian visceral leishmaniasis. *New England Journal of Medicine*. 2002;**347**(22):1739-1746
- [19] Soto J, Arana BA, Tolado J, Rizzo N, Vega JC, Diaz A, et al. Miltefosine for new world cutaneous leishmaniasis. *Clinical Infectious Diseases*. 2004;**38**(9):1266-1272
- [20] Machado PR, Ampuero J, Guimarães LH, Villasboas L, Rocha AT, Schriefer A, et al. Miltefosine in the treatment of cutaneous leishmaniasis caused by *Leishmania braziliensis* in Brazil: A randomized and controlled trial. *PLoS Neglected Tropical Diseases*. 2010;**4**(12):1-6
- [21] Rakotomanga M, Blanc S, Gaudin K, Chaminade P, Loiseau PM. Miltefosine affects lipid metabolism in *Leishmania donovani* promastigotes. *Antimicrobial Agents and Chemotherapy*. 2007;**51**(4):1425-1430
- [22] Lira R, Contreras LM, Santa Rita RM, Urbina JA. Mechanism of action of anti-proliferative lysophospholipid analogues against the protozoan parasite *Trypanosoma cruzi*: Potentiation of *in vitro* activity by the sterol biosynthesis inhibitor ketoconazole. *Journal of Antimicrobial Chemotherapy*. 2001;**47**(5):537-546
- [23] Saraiva VB, Gibaldi D, Previato JO, Mendonça-Previato L, Bozza MT, Freire-de-Lima CG, et al. Proinflammatory and cytotoxic effects of hexadecylphosphocholine (miltefosine) against drug-resistant strains of *Trypanosoma cruzi*. *Antimicrobial Agents and Chemotherapy*. 2002;**46**(11):3472-3477
- [24] Seifert K. Structures, targets and recent approaches in anti-leishmanial drug discovery and development. *The Open Medicinal Chemistry Journal*. 2011;**5**:31-39
- [25] Ouellette M, Drummelsmith J, Papadopolou B. Leishmaniasis: Drugs in the clinic, resistance and new developments. *Drug Resistance Updates*. 2004;**7**(4-5):257-266
- [26] Dorlo TPC, Balasegaram M, Beijnen JH, de vries PJ. Miltefosine: A review of its pharmacology and therapeutic efficacy in the treatment of leishmaniasis. *Journal of Antimicrobial Chemotherapy*. 2012;**67**(11):2576-2597
- [27] Maltezou HC. Drug resistance in visceral leishmaniasis. *Journal of Biomedicine and Biotechnology*. 2010;**2010**:1-8
- [28] Santa-Rita RM, Henriques-Pons A, Barbosa HS, de Castro SL. Effect of the lysophospholipid analogues edelfosine, ilmofosine and miltefosine against *Leishmania amazonensis*. *Journal of Antimicrobial Chemotherapy*. 2004;**54**(4):704-710
- [29] Verma NK, Singh G, Dey CS. Miltefosine induces apoptosis in arsenite-resistant *Leishmania donovani* promastigotes through mitochondrial dysfunction. *Experimental Parasitology*. 2007;**116**(1):1-13

- [30] Calogeropoulou T, Angelou P, Detsi A, Fragiadaki I, Scoulica E. Design and synthesis of potent antileishmanial cycloalkylidene-substituted ether phospholipid derivatives. *Journal of Medicinal Chemistry*. 2008;**51**(4): 897-908
- [31] Godinho JLP, Georgikopoulou K, Calogeropoulou T, de Souza W, Rodrigues JCF. A novel alkyl phosphocholine-dinitroaniline hybrid molecule exhibits biological activity *in vitro* against *Leishmania amazonensis*. *Experimental Parasitology*. 2013;**135**(1): 153-165
- [32] de Macedo-Silva S, Souza W, Rodrigues J. Sterol biosynthesis pathway as an alternative for the anti-protozoan parasite chemotherapy. *Current Medicinal Chemistry*. 2015;**22**(18): 2186-2198
- [33] Rodrigues JCF, Attias M, Rodriguez C, Urbina JA, de Souza W. Ultrastructural and biochemical alterations induced by promastigote and amastigote forms of *Leishmania amazonensis*. *Antimicrobial Agents and Chemotherapy*. 2002;**46**(2):487-499
- [34] Emami S, Tavangar P, Keighobadi M. An overview of azoles targeting sterol 14 α -demethylase for antileishmanial therapy. *European Journal of Medicinal Chemistry*. 2017;**135**:241-259
- [35] Urbina JA, Docampo R. Specific chemotherapy of Chagas disease: Controversies and advances. *Trends in Parasitology*. 2003;**19**(11):495-501
- [36] Benaim G, Sanders JM, Garcia-Marchán Y, Colina C, Lira R, Caldera AR, et al. Amiodarone has intrinsic anti-*Trypanosoma cruzi* activity and acts synergistically with posaconazole. *Journal of Medicinal Chemistry*. 2006;**49**(3):892-899
- [37] de Macedo-Silva ST, Urbina JA, de Souza W, Rodrigues JCF. *In vitro* activity of the antifungal azoles itraconazole and posaconazole against *Leishmania amazonensis*. *PLoS One*. 2013;**8**(12): e83247
- [38] Mukherjee S, Basu S, Zhang K. Farnesyl pyrophosphate synthase is essential for the promastigote and amastigote stages in *Leishmania major*. *Molecular and Biochemical Parasitology*. 2019;**230**:8-15
- [39] Gadelha APR, Brigagao CM, da Silva MB, Rodrigues ABM, Guimarães ACR, Paiva F, et al. Insights about the structure of farnesyl diphosphate synthase (FPPS) and the activity of bisphosphonates on the proliferation and ultrastructure of *Leishmania* and *Giardia*. *Parasites and Vectors*. 2020;**13**(1):1-18
- [40] Shah RR. Safety and tolerability of histone deacetylase (HDAC) inhibitors in oncology. *Drug Safety*. 2019;**42**(2):235-245
- [41] Andrews KT, Haque A, Jones MK. HDAC inhibitors in parasitic diseases. *Immunology and Cell Biology*. 2012; **90**(1):66-77
- [42] Kumar D, Rajanala K, Minocha N, Saha S. Histone H4 lysine 14 acetylation in *Leishmania donovani* is mediated by the MYST-family protein HAT4. *Microbiology*. 2012;**158**(2):328-337
- [43] Saxena A, Lahav T, Holland N, Aggarwal G, Anupama A, Huang Y, et al. Analysis of the *Leishmania donovani* transcriptome reveals an ordered progression of transient and permanent changes in gene expression during differentiation. *Molecular and Biochemical Parasitology*. 2007;**152**(1): 53-65
- [44] Prasanna P, Kumar R, Singh VK, Upadhyay A. Cloning, purification, and homology modeling of histone deacetylase in *Leishmania donovani*. *Infection, Genetics and Evolution*. 2021;**89**:1-6

- [45] Yahiaoui B, Taibi A, Ouaisi A. A *Leishmania major* protein with extensive homology to silent information regulator 2 of *Saccharomyces cerevisiae*. *Gene*. 1996;**169**(1):115-118
- [46] Ritagliati C, Alonso VL, Manarin R, Cribb P, Serra EC. Overexpression of cytoplasmic TcSIR2RP1 and mitochondrial TcSIR2RP3 impacts on *Trypanosoma cruzi* growth and cell invasion. *PLoS Neglected Tropical Diseases*. 2015;**9**(4):1-22
- [47] Vergnes B, Sereno D, Madjidian-Sereno N, Lemesre JL, Ouaisi A. Cytoplasmic SIR2 homologue overexpression promotes survival of *Leishmania* parasites by preventing programmed cell death. *Gene*. 2002;**296**(1-2):139-150
- [48] Vergnes B, Sereno D, Tavares J, Cordeiro-Da-Silva A, Vanhille L, Madjidian-Sereno N, et al. Targeted disruption of cytosolic SIR2 deacetylase discloses its essential role in *Leishmania* survival and proliferation. *Gene*. 2005;**363**(1-2):85-96
- [49] Mittal N, Muthuswami R, Madhubala R. The mitochondrial SIR2 related protein 2 (SIR2RP2) impacts *Leishmania donovani* growth and infectivity. *PLoS Neglected Tropical Diseases*. 2017;**11**(5):e0005590
- [50] North BJ, Marshall BL, Borra MT, Denu JM, Verdin E. The human Sir2 ortholog, SIRT2, is an NAD⁺-dependent tubulin deacetylase. *Molecular Cell*. 2003;**11**(2):437-444
- [51] Tavares J, Ouaisi A, Santarém N, Sereno D, Vergnes B, Sampaio P, et al. The *Leishmania infantum* cytosolic SIR2-related protein 1 (LiSIR2RP1) is an NAD⁺-dependent deacetylase and ADP-ribosyltransferase. *Biochemical Journal*. 2008;**415**(3):377-386
- [52] Adriano MA, Vergnes B, Poncet J, Mathieu-Daude F, da Silva AC, Ouaisi A, et al. Proof of interaction between *Leishmania* SIR2RP1 deacetylase and chaperone HSP83. *Parasitology Research*. 2007;**100**(4):811-818
- [53] Wiesgigl M, Clos J. Heat shock protein 90 homeostasis controls stage differentiation in *Leishmania donovani*. *Molecular Biology of the Cell*. 2001;**12**:3307-3316
- [54] Zemzoumi K, Sereno D, François C, Guilvard E, Lemesre JL, Ouaisi A. *Leishmania major*: Cell type dependent distribution of a 43 kDa antigen related to silent information regulatory-2 protein family. *Biology of the Cell*. 1998;**90**(3):239-245
- [55] Silvestre R, Cordeiro-Da-Silva A, Tavares J, Sereno D, Ouaisi A. *Leishmania* cytosolic silent information regulatory protein 2 deacetylase induces murine B-cell differentiation and *in vivo* production of specific antibodies. *Immunology*. 2006;**119**(4):529-540
- [56] Vergnes B, Gazanion E, Grentzinger T. Functional divergence of SIR2 orthologs between trypanosomatid parasites. *Molecular and Biochemical Parasitology*. 2016;**207**(2):96-101
- [57] Sacconnay L, Smirlis D, Queiroz EF, Wolfender JL, Soares MBP, Carrupt PA, et al. Structural insights of SIR2rp3 proteins as promising biotargets to fight against Chagas disease and leishmaniasis. *Molecular BioSystems*. 2013;**9**(9):2223-2230
- [58] Verçoza BRF, Godinho JLP, de Macedo-Silva ST, Huber K, Bracher F, de Souza W, et al. KH-TFMDI, a novel sirtuin inhibitor, alters the cytoskeleton and mitochondrial metabolism promoting cell death in *Leishmania amazonensis*. *Apoptosis*. 2017;**22**(9):1169-1188
- [59] Calegari-Silva TC, Vivarini ÁC, Pereira R de MS, Dias-Teixeira KL,

- Rath CT, Pacheco ASS, et al. *Leishmania amazonensis* downregulates macrophage iNOS expression via Histone Deacetylase 1 (HDAC1): A novel parasite evasion mechanism. *European Journal of Immunology*. 2018;**48**(7):1188-1198
- [60] Mukherjee S, Mukherjee B, Mukhopadhyay R, Naskar K, Sundar S, Dujardin J-C, et al. Imipramine exploits histone deacetylase 11 to increase the IL-12/IL-10 ratio in macrophages infected with antimony-resistant *Leishmania donovani* and clears organ parasites in experimental infection. *The Journal of Immunology*. 2014;**193**(8):4083-4094
- [61] Perez J, Fuertes M, Nguewa P, Castilla J, Alonso C. Anticancer compounds as leishmanicidal drugs: Challenges in chemotherapy and future perspectives. *Current Medicinal Chemistry*. 2008;**15**(5):433-439
- [62] Lowe G, Droz AS, Vilaivan T, Weaver GW, Tweedale L, Pratt JM, et al. Cytotoxicity of (2,2',6',2''-Terpyridine) platinum(II) Complexes to *Leishmania donovani*, *Trypanosoma cruzi*, and *Trypanosoma brucei*. *Journal of Medicinal Chemistry*. 1999;**42**(6):999-1006
- [63] Tavares J, Ouaisi M, Ouaisi A, Cordeiro-da-Silva A. Characterization of the anti-*Leishmania* effect induced by cisplatin, an anticancer drug. *Acta Tropica*. 2007;**103**(2):133-141
- [64] Miranda VM, Costa MS, Guilardi S, Machado AEH, Ellena JA, Tudini KAG, et al. *In vitro* leishmanicidal activity and theoretical insights into biological action of ruthenium(II) organometallic complexes containing anti-inflammatories. *Biometals*. 2018;**31**(6):1003-1017
- [65] Minori K, Rosa LB, Bonsignore R, Casini A, Miguel DC. Comparing the antileishmanial activity of gold(I) and gold(III) compounds in *L. amazonensis* and *L. braziliensis* *in vitro*. *ChemMedChem*. 2020;**15**(22):2146-2150
- [66] Morey JR, McDevitt CA, Kehl-Fie TE. Host-imposed manganese starvation of invading pathogens: Two routes to the same destination. *Biometals*. 2015;**28**(3):509-519
- [67] Rice DR, Vacchina P, Norris-Mullins B, Morales MA, Smith BD. Zinc(II)-dipicolylamine coordination complexes as targeting and chemotherapeutic agents for *Leishmania major*. *Antimicrobial Agents and Chemotherapy*. 2016;**60**(5):2932-2940
- [68] Mirzaei M, Nadushan AS, Nooshadokht M, Abiri A, Anjomshoa M, Sharifi I, et al. *In silico* and *in vitro* inhibitory potential of an organometallic Cu (II) complex on *Leishmania major* stages. *Annals of Parasitology*. 2021;**67**(1):45-54
- [69] de Azevedo-França JA, Granado R, de Macedo Silva ST, dos Santos-Silva G, Scapin S, Borba-Santos LP, et al. Synthesis and biological activity of novel zinc-itraconazole complexes in protozoan parasites and *Sporothrix* spp. *Antimicrobial Agents and Chemotherapy*. 2020;**64**(5):e01980-e01919
- [70] Ravera M, Moreno-Viguri E, Paucar R, Pérez-Silanes S, Gabano E. Organometallic compounds in the discovery of new agents against kinetoplastid-caused diseases. *European Journal of Medicinal Chemistry*. 2018;**155**:459-482
- [71] Diro E, Blesson S, Edwards T, Ritmeijer K, Fikre H, Admassu H, et al. A randomized trial of AmBisome monotherapy and AmBisome and miltefosine combination to treat visceral leishmaniasis in HIV co-infected patients in Ethiopia. *PLoS Neglected Tropical Diseases*. 2019;**13**(1):1-19
- [72] Miret JA, Moreno J, Nieto J, Carter KC, Mullen AB, Ambros L, et al.

Antileishmanial efficacy and tolerability of combined treatment with non-ionic surfactant vesicle formulations of sodium stibogluconate and paromomycin in dogs. *Experimental Parasitology*. 2021;**220**:108033

[73] Rashid HU, Ullah I, Adeeb H, Zeb M, Mohammad A, Rehman N. Synergistic effect of oral allopurinol and intralesional sodium stibogluconate in the treatment of cutaneous leishmaniasis. *Journal of Ayub Medical College, Abbottabad : JAMC*. 2020;**32**(4):558-561

[74] Brustolin AÁ, Ramos-Milaré ÁCFH, de Castro KR, Mota CA, Pelloso SM, Silveira TGV. Successful combined therapy with Glucantime™ and pentoxifylline for the nasal mucosal lesion recently developed in a leishmaniasis patient having untreated cutaneous lesion for seven decades. *Parasitology International*. 2021;**85**:102422

[75] van Griensven J, Balasegaram M, Meheus F, Alvar J, Lynen L, Boelaert M. Combination therapy for visceral leishmaniasis. *The Lancet Infectious Diseases*. 2010;**10**(3):184-194

[76] Kar A, Jayaraman A, Charan Raja MR, Srinivasan S, Debnath J, Mahapatra SK. Synergic effect of eugenol oleate with amphotericin B augments anti-leishmanial immune response in experimental visceral leishmaniasis *in vitro* and *in vivo*. *International Immunopharmacology*. 2021;**91**:107291

[77] Albalawi AE, Abdel-Shafy S, Khalaf AK, Alanazi AD, Baharvand P, Ebrahimi K, et al. Therapeutic potential of green synthesized copper nanoparticles alone or combined with meglumine antimoniate (Glucantime®) in cutaneous leishmaniasis. *Nanomaterials*. 2021;**11**(4):891

[78] Bahrami S, Oryan A, Bemani E. Efficacy of amiodarone and

voriconazole combination therapy in cutaneous leishmaniasis in the mice experimentally infected with *Leishmania major*. *Journal of Infection and Chemotherapy*. 2021;**27**(7):984-990

[79] Caridha D, Sciotti RJ, Sousa J, Vesely B, Teshome T, Bonkougou G, et al. Combination of subtherapeutic doses of tretazicar and liposomal amphotericin B suppresses and cures *Leishmania major*-induced cutaneous lesions in murine models. *ACS Infectious Diseases*. 2021;**7**(2):506-517

[80] de Moraes-Teixeira E, Gallupo MK, Rodrigues LF, Romanha á lvaro J, Rabello A. *In vitro* interaction between paromomycin sulphate and four drugs with leishmanicidal activity against three New World *Leishmania* species. *Journal of Antimicrobial Chemotherapy*. 2014;**69**(1):150-154

[81] Sundar S, Rai M, Chakravarty J, Agarwal D, Agrawal N, Vaillant M, et al. New treatment approach in Indian visceral leishmaniasis: Single-dose liposomal amphotericin b followed by short-course oral miltefosine. *Clinical Infectious Diseases*. 2008;**47**(8):1000-1006

[82] Singh N, Kumar M, Singh RK. Leishmaniasis: Current status of available drugs and new potential drug targets. *Asian Pacific Journal of Tropical Medicine*. 2012;**5**(6):485-497

[83] Odds FC. Synergy, antagonism, and what the chequerboard puts between them. *Journal of Antimicrobial Chemotherapy*. 2003;**52**(1):1

[84] Ahmed H, Curtis CR, Tur-Gracia S, Olatunji TO, Carter KC, Williams RAM. Drug combinations as effective anti-leishmanials against drug resistant: *Leishmania mexicana*. *RSC Medicinal Chemistry*. 2020;**11**(8):905-912

[85] Seifert K, Croft SL. *In vitro* and *in vivo* interactions between miltefosine

and other antileishmanial drugs. Antimicrobial Agents and Chemotherapy. 2006;**50**(1):73-79

[86] Dar MJ, Khalid S, McElroy CA, Satoskar AR, Khan GM. Topical treatment of cutaneous leishmaniasis with novel amphotericin B-miltefosine co-incorporated second generation ultra-deformable liposomes. International Journal of Pharmaceutics. 2020;**573**:118900

[87] Rebello KM, Andrade-Neto VV, Gomes CRB, de Souza MVN, Branquinho MH, Santos ALS, et al. Miltefosine-lopinavir combination therapy against *leishmania infantum* infection: *In vitro* and *in vivo* approaches. Frontiers in Cellular and Infection Microbiology. 2019;**9**:1-8

[88] Valdivieso E, Mejías F, Carrillo E, Sánchez C, Moreno J. Potentiation of the leishmanicidal activity of nelfinavir in combination with miltefosine or amphotericin B. International Journal of Antimicrobial Agents. 2018;**52**(5): 682-687

[89] Banik BL, Fattahi P, Brown JL. Polymeric nanoparticles: The future of nanomedicine. Wiley Interdisciplinary Reviews: Nanomedicine and Nanobiotechnology. 2016;**8**(2):271-299

[90] Black CDV, Watson GJ, Ward RJ. The use of pentostam liposomes in the chemotherapy of experimental leishmaniasis. Transactions of the Royal Society of Tropical Medicine and Hygiene. 1977;**71**(6):550-552

[91] Alving CR, Steck EA, Chapman WL, Waits VB, Hendricks LD, Swartz GM, et al. Therapy of leishmaniasis: Superior efficacies of liposome encapsulated drugs. Proceedings of the National Academy of Sciences of the United States of America. 1978;**75**(6):2959-2963

[92] Müller RH, Jacobs C, Kayser O. Nanosuspensions as particulate drug

formulations in therapy: Rationale for development and what we can expect for the future. Advanced Drug Delivery Reviews. 2001;**47**(1):3-19

[93] Brown M, Noursadeghi M, Boyle J, Davidson RN. Successful liposomal amphotericin B treatment of *Leishmania braziliensis* cutaneous leishmaniasis. British Journal of Dermatology. 2005;**153**(1):203-205

[94] Shirzadi MR. Liposomal amphotericin B: A review of its properties, function, and use for treatment of cutaneous leishmaniasis. Research and Reports in Tropical Medicine. 2019;**10**:11-18

[95] Gutiérrez V, Seabra AB, Reguera RM, Khandare J, Calderón M. New approaches from nanomedicine for treating leishmaniasis. Chemical Society Reviews. 2016;**45**(1):152-168

[96] Akbari M, Oryan A, Hatam G. Application of nanotechnology in treatment of leishmaniasis: A Review. Acta Tropica. 2017;**172**:86-90

[97] de Souza A, Marins DSS, Mathias SL, Monteiro LM, Yukuyama MN, Scarim CB, et al. Promising nanotherapy in treating leishmaniasis. International Journal of Pharmaceutics. 2018;**547**(1-2):421-431

[98] Nafari A, Cheraghipour K, Sepahvand M, Shahrokhi G, Gabal E, Mahmoudvand H. Nanoparticles: New agents toward treatment of leishmaniasis. Parasite Epidemiology and Control. 2020;**10**:e00156

[99] Barick KC, Aslam M, Lin YP, Bahadur D, Prasad VP, Dravid VP. Novel and efficient MR active aqueous colloidal Fe₃O₄ nanoassemblies. Journal of Materials Chemistry. 2009;**19**(38): 7023-7029

[100] Lee JH, Jang JT, Choi JS, Moon SH, Noh SH, Kim JW, et al.

Exchange-coupled magnetic nanoparticles for efficient heat induction. *Nature Nanotechnology*. 2011;**6**(7):418-422

[101] Barick KC, Singh S, Jadhav NV, Bahadur D, Pandey BN, Hassan PA. PH-responsive peptide mimic shell cross-linked magnetic nanocarriers for combination therapy. *Advanced Functional Materials*. 2012;**22**(23): 4975-4984

[102] Kumar R, Pandey K, Sahoo GC, Das S, Das VNR, Topno RK, et al. Development of high efficacy peptide coated iron oxide nanoparticles encapsulated amphotericin B drug delivery system against visceral leishmaniasis. *Materials Science and Engineering C*. 2017;**75**:1465-1471

[103] Abazari R, Mahjoub AR, Molaie S, Ghaffarifar F, Ghasemi E, Slawin AMZ, et al. The effect of different parameters under ultrasound irradiation for synthesis of new nanostructured Fe_3O_4 @bio-MOF as an efficient anti-leishmanial *in vitro* and *in vivo* conditions. *Ultrasonics Sonochemistry*. 2018;**43**:248-261

[104] Zomorodian K, Veisi H, Mousavi SM, Ataabadi MS, Yazdanpanah S, Bagheri J, et al. Modified magnetic nanoparticles by PEG-400-immobilized ag nanoparticles (Fe_3O_4 @PEG-Ag) as a core/shell nanocomposite and evaluation of its antimicrobial activity. *International Journal of Nanomedicine*. 2018; **13**:3965-3973

[105] Berry SL, Walker K, Hoskins C, Telling ND, Price HP. Nanoparticle-mediated magnetic hyperthermia is an effective method for killing the human-infective protozoan parasite *Leishmania mexicana* *in vitro*. *Scientific Reports*. 2019;**9**(1):1-9