

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

185,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Fibre Bundle Models and 3D Object Recognition

Fangxing Li¹, Linyu Peng² and Huafei Sun³

^{1,3}Beijing Institute of Technology

²University of Surrey

^{1,3}China P. R.

²UK

1. Introduction

Surface modeling is a fundamental issue in 3D images, shape synthesis and recognition. Recently, a surface model called the fibre bundle model is proposed (Chao et al., 2000; Chao & Kim, 2004; Chao et al., 2004; Chao & Li, 2005; Suzuki & Chao, 2002). The fibre bundle model becomes powerful in its full generality however still needs much information, the information of these two curves at all points, which are roughly equivalent to the information of pointwise representation itself. On the other hand, to apply such a surface model efficiently in shape generation and representation, one needs to know the geometrical quantities of the model.

Object recognition techniques using 3D image data are expected to play an important role in recognition-synthesis image coding of 3D moving pictures or animations in virtual environments and image communications. Currently used 3D free object representations seem insufficient in the sense that, for models such as generalized cylinders or super-quadratics, it is usually difficult to find the invariant features, especially to find the complete set of invariants, which is defined as the smallest number of invariants in order to uniquely determine and reproduce the shapes (Chao & Ishii, 1999; Chao & Suzuki, 2002; Chao et al., 1999; Kawano et al., 2002; Sano et al., 2001).

From the view of differential geometry and Riemannian geometry, the authors obtained the geometric structures about fibre bundle models (Li et al., 2008, 1; 2). Meanwhile, an algorithm of 3D object recognition using the linear Lie algebra models is presented, including a convenient recognition method for the objects which are symmetrical about some axis (Li et al., 2009).

2. Fibre bundle model and its geometry

2.1 Fibre bundle model of surfaces

A surface $F = \{F(u, v)\}$ is called a fibre bundle on a given base curve $b = \{b(v), v \in \mathbb{R}\}$, if locally (i.e. at a neighborhood of any point) F is a direct product of b and another curve called a fibre curve. More specifically, as shown in Fig. 1, there is a projection map $\pi : F \rightarrow b$, such that for a point $x \in b$, there is a curve $f_x = \{f_x(u), u \in \mathbb{R}\}$ on F

$$f_x := \pi^{-1}(x) \subset F, \quad (1)$$

which is called a fibre at the base point x . For any $x \in b$, there is a neighborhood of x in $b : U_x \in b$ such that

$$\pi^{-1}(U_x) \cong U_x \times f_x. \quad (2)$$




Fig. 1. Fibre bundle model of surfaces.

The fibre bundle model can represent any surface F , e.g. generalized cylinders and ruled surfaces are special cases of this model.

On the other hand, the fibre bundle model is a local direction product, which means the fibre curves could be very different at each base point. Such a nontrivial fibre bundle thus can represent arbitrary complicated surface.

2.2 Fibre bundle model of 1-parameter lie groups of linear lie algebra and its geometry

Let the fibre curve of the fibre bundle model be a 1-parameter Lie group

$$g_v = \{g_v(u) = e^{Au}b(v), u \in \mathbb{R}\}, \quad (3)$$

where A is a 3×3 matrix called the representation matrix of the fibre curve. Therefore, the surface is defined as

$$F = \{x(u, v) = e^{Au}b(v), u, v \in \mathbb{R}\}. \quad (4)$$

The points $b(v)$ on the base curve b are initial points of integral flows of 1-parameter Lie groups g_v .

The Lie algebra of this fibre bundle (roughly can be regarded as its tangent vector field) is a linear Lie algebra

$$L := \frac{\partial x}{\partial u} = x_u = Ae^{Au}b(v) = Ax. \quad (5)$$

A major advantage of this model is that the Lie algebras of fibre curves are uniquely determined by a complete set of invariants I under Euclidean transformation

$$I = \{\sigma_1, \sigma_2, \sigma_3, \phi_1, \phi_2, \phi_3\}, \quad (6)$$

where $\{\sigma_i\}$ are the singular values of A , assuming the singular value decomposition of $A^R = R^T A R, R \in SO_3(\mathbb{R})$ is $A^R = U^T \Lambda V, \Lambda = \text{diag}\{\sigma_i\}$. $\{\phi_i\}$ are the Euler angles of $VU^T \in SO_3(\mathbb{R})$. Here, a complete set of invariants is the minimal set of invariants which can uniquely determine the curve. Thus, information to describe the fibre-bundle surface model is the base curve and the six invariants of the linear Lie algebra, i.e. of the representation matrix A .




Fig. 2. Fibre bundle of 1-parameter groups of linear Lie algebra.

Lemma 2.1 For any $n \times n$ matrix $A = (a_{ij})$ where $a_{ij} \in \mathbb{R}$ and any $u \in \mathbb{R}$, one has

- 1. $\frac{d}{du}(e^{Au}) = Ae^{Au} = e^{Au}A$;
- 2. $\det(e^{Au}) = e^{\text{tr}(Au)} = e^{\text{tr}(A)u}$, where $\text{tr}(A)$ is the trace of A ;
- 3. when $AB = BA, e^{A+B} = e^Ae^B = e^Be^A$;
- 4. $(e^A)^T = e^{A^T}$.

Lemma 2.2 $A \in so_3(\mathbb{R}) = \{B|B^T = -B\}$, then $e^{Au} \in SO_3(\mathbb{R}) = \{B|B^TB = I, |B| = 1\}$, where $u \in \mathbb{R}$.

Proof. If $A \in so_3(\mathbb{R})$, we can get that $A^T = -A$ and $\text{tr}(A) = 0$. Then from Lemma 2.1, we get $(e^{Au})^Te^{Au} = e^{A^Tu}e^{Au} = e^{(A^T+A)u} = I$ and $\det(e^{Au}) = e^{\text{tr}(A)u} = 1$. Therefore, $e^{Au} \in SO_3(\mathbb{R})$.

Theorem 2.1 For the fibre bundle surface defined by

$$F := \{x(u, v) = e^{Au}b(v), u, v \in \mathbb{R}\}, \tag{7}$$

where $A \in so_3(\mathbb{R})$, and $b(v)$ is a two order differentiable vector, the Gaussian curvature K and the mean curvature H of the fibre bundle surface are given by

$$K = \frac{\det(Ab, b', A^2b) \det(Ab, b', b'') - (\det(Ab, b', Ab'))^2}{(|Ab|^2|b'|^2 - ((Ab) \cdot b')^2)^2} \tag{8}$$

and

$$H = \frac{1}{2} \frac{|b'|^2 \det(Ab, b', A^2b) - 2((Ab) \cdot b') \det(Ab, b', Ab') + |Ab|^2 \det(Ab, b', b'')}{(|Ab|^2 |b'|^2 - ((Ab) \cdot b')^2)^{\frac{3}{2}}}, \quad (9)$$

respectively, where (\cdot) denotes the inner product of vectors and \det the determinant of an $n \times n$ matrix, respectively.

Proof. Firstly, from the Lemmas we can get

$$x_u = Ae^{Au}b, x_v = e^{Av}b'$$

and

$$x_{uu} = e^{Au}A^2b, x_{uv} = x_{vu} = e^{Au}Ab', x_{vv} = e^{Av}b''.$$

Then we can get

$$E = (x_u \cdot x_u) = |Ab|^2, F = (x_u \cdot x_v) = ((Ab) \cdot b'), G = (x_v \cdot x_v) = |b'|^2,$$

$$|x_u \times x_v|^2 = (x_u \cdot x_u)(x_v \cdot x_v) - (x_u \cdot x_v)^2 = |Ab|^2 |b'|^2 - ((Ab) \cdot b')^2$$

and

$$\begin{aligned} L &= \frac{(x_u, x_v, x_{uu})}{|x_u \times x_v|} = \frac{\det(Ab, b', A^2b)}{\sqrt{|Ab|^2 |b'|^2 - ((Ab) \cdot b')^2}}, \\ M &= \frac{(x_u, x_v, x_{uv})}{|x_u \times x_v|} = \frac{\det(Ab, b', Ab')}{\sqrt{|Ab|^2 |b'|^2 - ((Ab) \cdot b')^2}}, \\ N &= \frac{(x_u, x_v, x_{vv})}{|x_u \times x_v|} = \frac{\det(Ab, b', b'')}{\sqrt{|Ab|^2 |b'|^2 - ((Ab) \cdot b')^2}}. \end{aligned}$$

Therefore the Gaussian curvature is given by

$$\begin{aligned} K &= \frac{LN - M^2}{EG - F^2} \\ &= \frac{\det(Ab, b', A^2b) \det(Ab, b', b'') - (\det(Ab, b', Ab'))^2}{(|Ab|^2 |b'|^2 - ((Ab) \cdot b')^2)^2} \end{aligned}$$

and the mean curvature is given by

$$\begin{aligned} H &= \frac{1}{2} \frac{LG - 2MF + NE}{EG - F^2} \\ &= \frac{1}{2} \frac{|b'|^2 \det(Ab, b', A^2b) - 2((Ab) \cdot b') \det(Ab, b', Ab') + |Ab|^2 \det(Ab, b', b'')}{(|Ab|^2 |b'|^2 - ((Ab) \cdot b')^2)^{\frac{3}{2}}}. \end{aligned}$$

Example 2.1 Taking $b(v) = (0, \sin(v), \cos(v))^T$, $A = \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$, from Lemma 2.2, we know

that $e^{Au} \in SO_3(\mathbb{R})$, then we can get $x(u, v) = e^{Au}b(v)$. Thus, we can get

$$\begin{aligned} b' &= (0, \cos(v), -\sin(v))^T, b'' = (0, -\sin(v), -\cos(v))^T, \\ Ab &= (-\sin(v), 0, 0)^T, Ab' = (-\cos(v), 0, 0)^T, A^2b = (0, -\sin(v), 0)^T. \end{aligned}$$

Then the Gaussian curvature is given by

$$K = 1$$

and the mean curvature is

$$H = \begin{cases} 1, & \sin(v) \geq 0 \\ -1, & \sin(v) < 0 \end{cases}.$$




Fig. 3. The figure of $x(u, v)$ in Example 2.1.

Example 2.2 Taking $b(v) = (1, 1, v)^T$, $A = \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$, we get $x(u, v) = e^{Au}b(v)$. Therefore we can get

$$b' = (0, 0, 1)^T, b'' = (0, 0, 0)^T, \\ Ab = (-1, 1, 0)^T, Ab' = (0, 0, 0)^T, A^2b = (-1, -1, 0)^T.$$

Then the Gaussian curvature is given by

$$K = 0$$

and the mean curvature is

$$H = -\frac{\sqrt{2}}{4}.$$




Fig. 4. The figure of $x(u, v)$ in Example 2.2.

Example 2.3 Taking $b(v) = (\cos^3(v), \sin^3(v), v)^T$, $A = \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$, we get $x(u, v) = e^{Au}b(v)$.

Hence

$$b' = (-3 \cos^2(v) \sin(v), 3 \sin^2(v) \cos(v), 1)^T,$$

$$b'' = (-3 \cos^3(v) + 6 \sin^2(v) \cos(v), -3 \sin^3(v) + 6 \sin(v) \cos^2(v), 0)^T,$$

$$Ab = (-\sin^3(v), \cos^3(v), 0)^T, Ab' = (-3 \sin^2(v) \cos(v), -3 \cos^2(v) \sin(v), 0)^T,$$

$$A^2b = (-\cos^3(v), -\sin^3(v), 0)^T.$$

Then the Gaussian curvature is given by

$$K = \frac{48 - 96 \sin^2(2v) + 36 \sin^4(2v)}{(4 + 6 \sin^2(2v) - 9 \sin^4(2v))^2}$$

and the mean curvature is given by

$$H = \frac{18 \sin^2(2v) - 16}{(4 + 6 \sin^2(2v) - 9 \sin^4(2v))^{3/2}}.$$

We can see that

$$K = \begin{cases} -12, & \sin(2v) = 1 \\ 3, & \sin(2v) = 0 \\ 0, & \sin(2v) = \frac{\sqrt{6}}{3} \end{cases}.$$




Fig. 5. The figure of $x(u,v)$ in Example 2.3.




Fig. 6. Curvatures of $x(u,v)$ in Example 2.3.

2.3 Fibre bundle model of 1-parameter lie groups of Hamiltonian lie algebra and its geometry

Consider a spatial curve on a surface M as

$$x(t) := \begin{pmatrix} x_1(t) \\ x_2(t) \\ x_3(t) \end{pmatrix} \in M \subset \mathbb{R}^3 \tag{10}$$

and a state vector as $y(t) = (y_1(t), y_2(t), \dots, y_6(t))^T \in \mathbb{R}^6$

$$y(t) := \begin{pmatrix} \dot{x}(t) \\ x(t) \end{pmatrix} \in M \otimes T_x M, \tag{11}$$

where $T_x M$ is the tangent space of M at point x . A Hamiltonian Lie algebra of tangent vector fields is defined by the infinitesimal generator

$$\begin{pmatrix} \ddot{x}(t) \\ \dot{x}(t) \end{pmatrix} = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \begin{pmatrix} \dot{x}(t) \\ x(t) \end{pmatrix} \quad (12)$$

or $\dot{y} = Hy$, where $H := \begin{pmatrix} A & B \\ C & D \end{pmatrix}$ is called a representation matrix.

For a special fibre bundle model of 1-parameter Lie groups of Hamiltonian Lie algebra defined by

$$F := \{y(u, v) = e^{Hu}b(v), u, v \in \mathbb{R}\},$$

which is an embedded surface of \mathbb{R}^6 , where $e^{Hu} \in SO_6(\mathbb{R})$ and $b(v)$ is a two order differentiable vector, one can get the following

Theorem 2.2 The Gaussian curvature $K(u, v)$ of the model is given by

$$\begin{aligned} K(u, v) = & \frac{1}{\det(g_{ij})} ((Hb \cdot Hb'') + |Hb'|^2) \\ & + \frac{(Hb \cdot Hb')}{\det^2(g_{ij})} ((Hb \cdot b')(Hb \cdot b'') + (Hb \cdot b')(Hb' \cdot b') \\ & - (Hb \cdot Hb')|b'|^2 - (b' \cdot b'')|Hb|^2), \end{aligned} \quad (13)$$

where (\cdot) denotes the inner product of vectors and \det the determinant of an $n \times n$ matrix, respectively.

Proof. Using Lemma 2.1, we can get

$$x_u = He^{Hu}b, \quad x_v = e^{Hu}b'$$

and

$$x_{uu} = H^2e^{Hu}b, \quad x_{uv} = x_{vu} = He^{Hu}b', \quad x_{vv} = e^{Hu}b''.$$

Then from $g_{ij} = (x_i \cdot x_j)$, we can get the Riemannian metric as

$$(g_{ij}) = \begin{pmatrix} |Hb|^2 & (Hb \cdot b') \\ (Hb \cdot b') & |b'|^2 \end{pmatrix}.$$

The determinant and inverse of (g_{ij}) are respectively given by

$$\det(g_{ij}) = |Hb|^2|b'|^2 - (Hb \cdot b')^2$$

and

$$(g^{ij}) = \frac{1}{\det(g_{ij})} \begin{pmatrix} |b'|^2 & -(Hb \cdot b') \\ -(Hb \cdot b') & |Hb|^2 \end{pmatrix}.$$

From

$$\Gamma_{ij}^k = \frac{1}{2}g^{kl} (\partial_j g_{li} + \partial_i g_{lj} - \partial_l g_{ij}),$$

one can get

$$\begin{aligned}\Gamma_{11}^1 &= \frac{1}{\det(g_{ij})}(Hb \cdot b')(Hb \cdot Hb'), \quad \Gamma_{11}^2 = -\frac{1}{\det(g_{ij})}(Hb \cdot Hb')|Hb|^2, \\ \Gamma_{12}^1 &= \Gamma_{21}^1 = \frac{1}{\det(g_{ij})}(Hb \cdot Hb')|b'|^2, \quad \Gamma_{21}^2 = \Gamma_{12}^2 = \frac{1}{\det(g_{ij})}(Hb \cdot b')(Hb \cdot Hb'), \\ \Gamma_{22}^1 &= \frac{1}{\det(g_{ij})}(((Hb' \cdot b') + (Hb \cdot b''))|b'|^2 - (Hb \cdot b')(b' \cdot b'')), \\ \Gamma_{22}^2 &= -\frac{1}{\det(g_{ij})}(((Hb' \cdot b') + (Hb \cdot b''))(Hb \cdot b') - (b' \cdot b'')|Hb|^2).\end{aligned}$$

Using

$$R_{ijkl} = R_{ikl}^h g_{hj},$$

where

$$R_{ikl}^h = \partial_k \Gamma_{il}^h - \partial_l \Gamma_{ik}^h + \Gamma_{il}^j \Gamma_{jk}^h - \Gamma_{ik}^j \Gamma_{jl}^h,$$

we can get the nonzero component of the curvature tensors

$$\begin{aligned}R(u, v, u, v) &= (Hb \cdot Hb'') + |Hb'|^2 \\ &\quad + \frac{(Hb \cdot Hb')}{\det(g_{ij})}((Hb \cdot b')(Hb \cdot b'') + (Hb \cdot b')(Hb' \cdot b')) \\ &\quad - (Hb \cdot Hb')|b'|^2 - (b' \cdot b'')|Hb|^2).\end{aligned}$$

Then from the definition of the Gaussian curvature

$$K(u, v) = \frac{R(u, v, u, v)}{\det(g_{ij})},$$

one can the the conclusion directly.

In all of the following examples, $A = \begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$, $B = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}$, $C = -B$ and $D = -A^T$,

then $H = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$.

Example 2.4 Taking $b(v) = (v, 1, 1, 2, 3, 4)^T$, therefore, $b'(v) = (1, 0, 0, 0, 0, 0)^T$, $b''(v) = (0, 0, 0, 0, 0, 0)^T$. We can get

$$K = \frac{109}{(2v^2 - 6v + 59)^2} > 0.$$

Example 2.5 Taking $b(v) = (\cos v, \sin v, 0, 1, 1, 0)^T$, then $b'(v) = (-\sin v, \cos v, 0, 0, 0, 0)^T$, $b''(v) = (-\cos v, -\sin v, 0, 0, 0, 0)^T$. We can get

$$K = -\frac{2\sqrt{2} + 1}{144} < 0.$$




Fig. 7. Gaussian curvature of the surface in Example 2.4.

Example 2.6 Taking $b(v) = (v,0,a,-a,0,a)^T$, since $b'(v) = (1,0,0,0,0,0)^T, b''(v) = (0,0,0,0,0,0)^T$. We can get

$$K = 0.$$

3. 3D object recognition based on fibre bundle models

Recall the fibre bundle model in (4) as

$$F = \{x(u,v) = e^{Au}b(v), u,v \in \mathbb{R}\}, \tag{14}$$

where A is the representation matrix of the fibre curve. The base curve $b(v)$ can also be described by a voluntariness initial point $x_0 \in \mathbb{R}$ and its representation matrix B given by

$$b(v) = e^{Bv}x_0. \tag{15}$$

And its tangent vector field is also a tangent vector field shown as

$$\frac{\partial x}{\partial v} := Be^{Bv}x_0 = Bb(v). \tag{16}$$

Therefore, the information to describe the fibre bundle model is the base curve and the invariants of the linear Lie algebra, i.e., the representation matrix A or representation matrix A, B and the initial point x_0 .

3.1 Simulations with known representation matrixes and the initial points

Next simulation results of shape synthesis using the proposed models are shown together with the invariants of representation matrix A or representation matrixes A, B and the initial point x_0 .




Fig. 8. Simulations with representation matrixes A and base curves $b(v)$. The representation matrixes A are $\begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$ and $\begin{pmatrix} 0 & 0 & -1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}$ in (a) and (b), respectively.




Fig. 9. Simulations with representation matrixes A, B and initial points x_0 . In (a), the representation matrixes are $A = \begin{pmatrix} 2 & 1.5 & 2 \\ -2 & -1.5 & -2 \\ -3 & -1.5 & -1 \end{pmatrix}$, and $B = \begin{pmatrix} 0 & 0 & -2 \\ -2 & 0 & 2 \\ 2 & 0 & 0 \end{pmatrix}$. And in (b), $A = \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$, $B = \begin{pmatrix} 0 & -1 & -1 \\ 2 & 1 & -2 \\ 2 & 2 & -1 \end{pmatrix}$.

3.2 Algorithm of 3D object recognition

Here we introduce an algorithm to obtain the invariants of a linear Lie algebra model from local image data of 3D objects as follows.

1. Take more than $N > 9$ position vectors $\{x_i\}_{i=1}^N$ on the object. Calculate the normal vectors at all positions $\{n_i\}_{i=1}^N$ from geometric relation, and normalize them.
2. From the position and normal vectors to build the following equations in entries of the representation matrix A under a rotation, where A stems from normalization

$$L := \frac{\partial x}{\partial u} = x_u = Ae^{Au}b(v) = Ax.$$

According to that the tangent vector field perpendicular to the normal vectors, we can get that n_i perpendiculars to $\frac{\partial x}{\partial u}|_{x_i} = Ax|_{x_i}$, then we can get

$$n_i^T Ax_i,$$

namely,

$$n_i^T \begin{pmatrix} a & b & c \\ d & e & f \\ g & h & i \end{pmatrix} x_i.$$

3. From linear system, we get the components of fibre representation matrix A .
4. Just liking the fibre representation matrix extraction, we can take more than nine position vectors y_i and the corresponding normal vectors to build this equation to solve the coefficient of the base curve representation matrix B .
5. Given a discretional initialization point x_0 and using the obtained representation matrix A and B to restore the primary 3D objects.

When we give the fibre representation matrix $A = \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$, then we can use Taylor series to calculate its linear Lie algebra

$$\exp(Au) = \begin{pmatrix} \cos u & -\sin u & 0 \\ \sin u & \cos u & 0 \\ 0 & 0 & 1 \end{pmatrix}. \quad (17)$$

So if we know the initialization point x_0 and the base curve representation matrix B , we can use this representation matrix A to express a form symmetry about an axis without solve the representation matrix A . That can help us easy to express a large number of shapes of 3D objects. (Fig. 8 and Fig. 9)

3.3 Practicality recognition

Next we consider the recognition of two objects, the sphere and vase. Our target sphere is shown in Fig. 10.

Firstly we give a base curve data (easy to confirm) and take more than nine position vectors from the sphere to recognize the object. Here we take data number as 30. From the algorithm, we can get the representation matrix as

$$A = \begin{pmatrix} -0.01 & 1 & -0.0043 \\ -0.9825 & -0.0032 & 0.0204 \\ 0.0202 & 0.0013 & -0.0001 \end{pmatrix}. \quad (18)$$

For the target vase shown in Fig. 12, we have to consider it as three parts, capsule, middle part and bottom, and do the simulations, respectively.

The target vase is a form symmetry about an axis, as presented we can use matrix $A = \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$ as the fibre representation matrix, then we only need to extract the base curve representation matrix B .



Fig. 10. Recognize target sphere.




Fig. 11. Recognition for a sphere.

For all the simulations in Fig. 13, the initial points are the same as $x_0 = (1, -1, 0)^T$. Meanwhile, from the algorithm, we obtain the base curve representation matrixes B in (a), (b) and (c) as $\begin{pmatrix} -6.0859 & 1 & -0.0689 \\ 0.7695 & -0.0865 & 0.0066 \\ -1.8276 & 0.3112 & -0.0212 \end{pmatrix}$, $\begin{pmatrix} -6.5247 & 1 & 0.2672 \\ -3.9099 & 0.6090 & 0.1608 \\ -0.6605 & 0.1122 & 0.0277 \end{pmatrix}$ and $\begin{pmatrix} -0.3293 & 1 & -0.0442 \\ -3.8194 & 3.0828 & -0.3170 \\ 13.1096 & -5.3890 & 0.8316 \end{pmatrix}$, respectively. By the way, all the recognition results are exactly calculated and restored under strong noisy environments.



Fig. 12. Recognize target vase.

4. Conclusion

As we know, differential geometry and Riemannian geometry are powerful in applications in kinds of fields, and many methods and subjects are proposed, e.g. geometric mechanics, human face recognition using the method of manifold and geometry in statistics. In this chapter, we show the beauty of the geometry of the fibre bundle models of 1-parameter Lie groups of linear Lie algebra and Hamiltonian Lie algebra.

Fibre bundle model is effective in its representation for objects. Any object can represent as the form of a fibre bundle model, theoretically. Nevertheless, in practice, it's a challenging task for one to get the parameters of a special object. When the representation matrixes and the initial points are given, one can obtain beautiful photos for 3D objects. However, once one have the object, how to recognize it, namely, how to realize it in one's computer is the special challenging but the most important work. We propose an algorithm for 3D object recognition mainly based on the geometric relationship of the positions and normal vectors. The recognition results of sphere and vase demonstrate the algorithm perfectly.

5. Acknowledgements

This work is partially supported by the National Natural Science Foundations of China, No. 10871218, No. 10932002.





Fig. 13. Recognition for a vase.

6. References

- Chao, J. & Ishii, S. (1999). Invariant Recognition and Segmentation of 3D Object Using Lie Algebra Models, *Proceedings of International Conference on Image Processing*, Vol.1: 550-554.
- Chao, J., Karasudani, A. & Minowa, K. (2000). Invariant Representation and Recognition of Objects Using Linear Lie Algebra of Tangent or Normal Vector Fields, *IEICE Japan Trans.(D-II)*, J83-D-II(9): 1870-1878.
- Chao, J. & Kim, J. (2004). A Fibre Bundle Model of Surfaces and its Generalization, *Proceedings of the 17th International Conference on Pattern Recognition*, Vol.1: 560-563.
- Chao, J., Kim, J. & Nagakura, A. (2004). A New Surface Model Based on a Fibre-bundle of 1-Parameter Groups, *IEEE International Conference on Multimedia and Expo*, Vol.1: 129-132.
- Chao, J. & Li, F. (2005). A Surface Model Based on a Fibre Bundle of 1-Parameter Groups of Hamiltonian Lie Algebra, *IEEE International Conference on Image Processing*, Vol.1: 1021-1024.
- Chao, J. & Suzuki, M. (2002). Invariant Extraction and Segmentation of 3D Objects Using Linear Lie Algebra Model, *Proceedings of International Conference on Image Processing*, Vol.1: 205-208.

- Chao, J., Ura, K. & Honma, G. (1999). Generation of 3D Objects Using Lie Algebra Models Based on Curvature Analysis and Comparison with B-spline Fitting, *Proceedings of International Conference on Image Processing*, Vol.4: 366-370.
- Chern, S.S., Chen, W. & Lam, K.S. (1999). *Lectures on Differential Geometry*, Series on University Mathematics, Vol.1, World Scientific Publishing Company.
- Hall, B.C. (2003). *Lie Groups, Lie Algebras, and Representations: An Elementary Introduction*, Springer-Verlag, New York.
- Kawano, S., Makino, M., Ishii, S. & Chao, J. (2002). Meshing Technology with Quality Assurance for Curved Surfaces Defined by Linear Lie Algebra, *Systems and Computers in Japan*, Vol.33(No.10): 64-73.
- Li, F., Peng, L. & Sun, H. (2008). On Fibre Bundle Surfaces and Their Curvatures, *Journal of Beijing Institute of Technology*, Vol.17(No.4): 451-454.
- Li, F., Wu, P., Peng, L. & Sun, H. (2008). Surface Model Based on a Fibre Bundle of Hamiltonian Lie Algebra and Gaussian Curvature, *Science & Technology Review*, Vol.26(No.8): 37-39.
- Li, F., Wu, P., Sun, H. & Peng, L. (2009). 3D Object Recognition Based on Linear Lie Algebra Model, *Journal of Beijing Institute of Technology*, Vol.18(No.1): 46-50.
- Olver, P.J. (1986). *Applications of Lie Groups to Differential Equations*, Springer-Verlag.
- Petersen, P. (2006). *Riemannian Geometry (Second Edition)*, Springer.
- Sano, H., Makino, M. & Chao, J. (2001). An Adaptive High Quality Mesh Generation for Surface Defined by Linear Lie Algebra, *Japan Society for Simulation Technology*, Vol.20: 251-258.
- Suzuki, M. & Chao, J. (2002). Invariant Extraction and Segmentation of 3D Objects with Linear Lie Algebra Model, *IEICE Japan Trans. on Information Systems(D)*, E85-D(8): 1306-1313.

IntechOpen



Object Recognition

Edited by Dr. Tam Phuong Cao

ISBN 978-953-307-222-7

Hard cover, 350 pages

Publisher InTech

Published online 01, April, 2011

Published in print edition April, 2011

Vision-based object recognition tasks are very familiar in our everyday activities, such as driving our car in the correct lane. We do these tasks effortlessly in real-time. In the last decades, with the advancement of computer technology, researchers and application developers are trying to mimic the human's capability of visually recognising. Such capability will allow machine to free human from boring or dangerous jobs.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Fangxing Li, Linyu Peng and Huafei Sun (2011). Fibre Bundle Models and 3D Object Recognition, Object Recognition, Dr. Tam Phuong Cao (Ed.), ISBN: 978-953-307-222-7, InTech, Available from:
<http://www.intechopen.com/books/object-recognition/fibre-bundle-models-and-3d-object-recognition>

INTeCH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike-3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/), which permits use, distribution and reproduction for non-commercial purposes, provided the original is properly cited and derivative works building on this content are distributed under the same license.

IntechOpen

IntechOpen