

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Effects of Domestic Waste Water on Water Quality of Three Reservoirs Supplying Drinking Water in Kaduna State - Northern Nigeria

Yahuza Tanimu, Sunday Paul Bako and John Ameh Adakole

*Department of Biological Sciences,
Ahmadu Bello University,
Nigeria*

1. Introduction

Waste water management in Nigeria does not receive the attention it deserves. Domestic waste water is discharged into streams and reservoirs that supply drinking water without any treatment (Tiseer *et al.*, 2008). Chemical substances from agricultural activities (fertilizers, pesticides and herbicides) in the catchment of reservoirs may introduce nutrients and heavy metals at concentrations higher than that which the environment can handle (WHO, 2006). Nigeria has a number of environmental regulatory laws which include: the National Environmental Standards and Regulations Enforcement Agency (Establishment) Act of 2007 (The NESREA Act), Nigerian Radioactive Waste Management Regulations 2006, Environmental Impact Assessment Act of 1992 (EIA Act), Harmful Wastes (Special Criminal Provisions etc.) Act of 1988 (Harmful Wastes Act), the National Oil Spill Detection and Response Agency (Establishment) Act 2006 (the NOSDRA Act) and Nigerian Radioactive Waste Management Regulations 2006. However, the enforcement of these regulations has not been effective (Onaruwa and Fakayode, 2002 and Adegroye, 2008) and thus pollution of both rural and urban water sources commonly occurs. In rural areas, natural sources of drinking water, such as streams, wells and other reservoirs are usually polluted by organic substances from upstream users who use water for Agricultural activities and other domestic purposes. In urban areas, population pressure, industrial activities and agricultural activities place pollution stress on reservoirs of water (Adakole *et al.*, 2002, Fakayode, 2005 Kimura, 2005, Tiseer *et al.*, 2008). The water in these reservoirs is sometimes taken directly without any form of treatment.

Contamination of sources of water by waste alters water quality (the physical, chemical and biological characteristics). When the physical and chemical conditions of ecosystems are changed beyond their normal ranges, changes may be expected to occur in individual organisms, populations and communities of the ecosystem (Lenat *et al.*, 1980, Akin-Oriola, 2003, Kadiri, 2006). Assemblages of cyanobacteria are good indicators of eutrophic water bodies (Reynolds, 1998). Some species of cyanobacteria could contain cyanotoxins in their cells but do not release these into the water, and as such are harmful only when consumed while others release toxins directly into the water (Chorus and Batram, 1999 and WHO, 2006). They can also alter taste and odor problems, cause water discoloration, or form large

mats that can interfere with boating, swimming, and fishing (Borgh, 2004). Cyanobacteria present a range of characteristics that give them clear competitive growth advantage over planktonic algae under certain environmental conditions. Such include; a requirement of low light intensity and little energy to maintain cell structure and function (Mur *et al.*, 1999); possession of gas vacuoles within their cells as a buoyancy regulation mechanism to avoid light damage in high-light environments, such as in tropic lakes or to access light in turbid or low-clarity water (Haider *et al.*, 2003). Cyanobacteria can also store phosphorus (luxury uptake), as a useful adaptation that allows continued growth under conditions of fluctuating nutrient concentrations. They are also not grazed by zooplankton, since they are not the preferred food for these aquatic organisms (Chorus and Batram, 1999). Data on levels of aquatic pollution and its implication to human health is generally lacking for most aquatic ecosystems in Nigeria. This study was therefore designed to evaluate the impact of waste water on three reservoirs receiving varying degrees of waste water.

2. Materials and methods

2.1 Study area

The three reservoirs studied were Gimbawa reservoir in Ikara Local Govt. (Long.10°6'N and Lat.8°35'E), Saminaka reservoir in Lere Local Govt. (10°70'N and 8°75'E) and Zaria reservoir, Zaria Local Government (7°38'N and 11°11'E) of Kaduna State. Kaduna State is located in the northern guinea savannah vegetative zone of Nigeria and has a tropical continental climate, with distinct wet and dry seasons. Three sampling stations were studied in each reservoir based on the different activities in the catchment from May 2008 to April 2009.

2.2 Phytoplankton collection:

Phytoplankton was collected using a conical shape plankton net of 20 cm diameter with a 50 ml collection vial attached to it (Perry, 2003). Samples were collected at three sampling points in each reservoir to reflect the various activities in the catchment. Phytoplankton was identified by consulting texts by Presscott (1977) and Perry (2003).

2.3 Physico-chemical parameters

Physico-chemical parameters of water were analyzed once a month from May 2008 to April 2009. Surface water temperature was measured *in situ* using a mercury thermometer. pH and Electrical Conductivity were measured using HANNA instrument (pH/Electrical Conductivity/Temperature meter model 210). Total Hardness, Dissolved oxygen (DO), Biological Oxygen Demand (BOD), Nitrate-Nitrogen (NO₃-N) and Phosphate-phosphorus (PO₄-P) were determined by methods described by APHA (1998).

2.4 Metal analysis

Metal concentration in the water samples was determined by Atomic Absorption Spectrophotometry (AAS). Water samples were digested by Nitric acid (HNO₃) digestion (as described by APHA, 1998).

3. Statistical analysis

Analysis Of Variance (ANOVA) was used to compare the means of physicochemical parameters; heavy metals concentration and abundance of phytoplankton from the different

reservoirs. Pearson’s correlation coefficient was used to determine the relationship between physicochemical parameters; physicochemical parameters and phytoplankton. Shannon-Wiener diversity index was used to determine phytoplankton diversity while Simpson’s Index was used to determine evenness of phytoplankton distribution.

4. Results

Mean monthly Air Temperature varied from 27.67 to 34.17°C with mean ± standard error of 31.76±0.62°C (Table1), for Gimbawa reservoir, whereas in Saminaka reservoir it ranged between 25°C and 36.67°C with mean ± SE of 30.96±0.97°C. In Zaria reservoir, air temperature ranged from 26 to 35.33°C mean ± SE of 29.67±0.68°C(Table 1). This observed difference was however not statistically significant.

The three reservoirs had mean ± SE of Surface water temperature was 26.16±1.00°C (Gimbawa), 26.19±1.07°C (Saminaka) and 26.08±0.63°C (Zaria) (Table 1). The differences were however, not statistically significant between months, seasons and reservoirs (P > 0.05).

	Gimbawa			Saminaka			Zaria		
	Min	Max	Mean ± SE	Min	Max	Mean ± SE	Min	Max	Mean ± SE
Air Temperature (°C)	27.67	34.67	31.76 ± 0.62	25	36.67	30.96 ± 0.97	26	35.33	29.67 ± 0.68
Water Temperature (°C)	20.33	31.67	26.16 ± 1.00	20	31	26.19 ± 1.07	20.67	28	26.08 ± 0.63
Secchi disc Transparency (cm)	13.67	69.67	17.67 ± 6.06	8.17	19.33	7.29 ± 2.19	13.67	47	21.48 ± 4.46
pH	6.87	8.76	7.54 ± 0.15	6.46	8.21	7.34 ± 0.15	6.42	7.9	7.31 ± 0.14
Electrical Conductivity (µS/cm)	45.1	573.33	120.50 ± 41.95	12.33	496	128.07 ±40.00	31.67	518	97.20 ±38.59
Dissolved Oxygen (Mg/L)	6.87	8.76	6.71 ± 0.39	3.52	9.1	6.16 ± 0.53	3.73	10.22	6.44 ± 0.58
BOD (Mg/L)	0.16	4.37	2.17 ± 0.41	0.37	5.57	2.60 ± 0.5	0.06	3.54	1.68 ± 0.38
Alkalinity (Mg/L)	2.87	6.7	5.05 ± 0.32	2.43	14.8	6.77 ± 1.16	2.5	5.8	4.29 ± 0.31
Hardness (Mg/L)	0.5	3.93	1.26 ± 0.26	0.43	4.53	1.46 ± 0.30	0.6	5.1	
NO ₃ -N (Mg/L)	0.03	0.19	0.12 ± 0.01	0.02	0.16	0.09 ± 0.01	0.01	0.55	0.13 ± 0.05
P ₀₄ -P (Mg/L)	0.06	0.62	0.29 ± 0.06	0.06	0.76	0.39 ±0.08	0.03	0.8	0.39 ±0.08

SE = Standard Error, BOD = Biochemical Oxygen Demand, NO₃-N = Nitrate-Nitrogen, P₀₄-P = Phosphate-phosphorus

Table 1. Physico-chemical characteristics of Gimbawa, Saminaka and Zaria reservoirs

Secchi Disc Transparency values in Gimbawa reservoir had the highest value of 69.67cm and lowest of 13.67cm. In Saminaka reservoir, the values ranged from 4.36 to 19.33cm, while in the Zaria reservoir it ranged from 13.67 to 47cm. The mean $\pm$ Standard Error of the reservoirs are Gimbawa: 17.67 ± 6.06 cm, Saminaka: 7.29 ± 2.19 cm and Zaria: 21.48 ± 4.46 cm (Table 1). This observed difference was statistically significant between reservoirs ($P < 0.05$) and between seasons ($P < 0.05$).

pH values in Gimbawa reservoir varied from 6.87 to 8.76. In Saminaka reservoir, the highest pH value was 8.21 and lowest was 6.46. While in Zaria reservoir, the highest pH value was 7.9 and lowest of 6.42. The mean $\pm$ SE observed in the reservoirs were: Gimbawa, 7.54 ± 0.15 ; Saminaka, 7.44 ± 0.15 and Zaria, 7.31 ± 0.14 (Table 1). The observed differences were not significant between reservoirs ($P > 0.05$) but significant between months ($P < 0.05$) and seasons ($P < 0.01$).

The mean $\pm$ SE Electrical of Conductivity (EC) for Gimbawa, Saminaka and Zaria reservoirs observed were $120.50 \pm 41.95 \mu\text{S/cm}$, $128.07 \pm 40.00 \mu\text{S/cm}$ and $97.20 \pm 38.59 \mu\text{S/cm}$ respectively (Table 1). The variation of EC was significant only between months ($P < 0.05$).

Dissolved Oxygen (DO) varied between 8.58mg/L and 3.9 mg/L in Gimbawa reservoir. Saminaka reservoir had values ranging between 9.1mg/L to 3.52mg/L while in Zaria reservoir had range of values for DO from 3.73 mg/L to 10.22 mg/L. The mean $\pm$ SE of Gimbawa, Saminaka and Zaria reservoirs observed were 6.71 ± 0.39 mg/L, 6.16 ± 0.53 mg/L and 6.44 ± 0.58 respectively (Table 1). The variation of DO was significant between months and seasons ($P < 0.05$).

Biochemical Oxygen Demand (BOD) values in Gimbawa reservoir ranged from 4.37mg/L to 0.16mg/L, In Saminaka reservoir the values range from 0.37 to 5.57mg/L whereas in Zaria reservoir the values ranged from 0.06mg/L to 3.54mg/L. The mean $\pm$ SE of Gimbawa, Saminaka and Zaria reservoirs observed were 2.17 ± 0.41 mg/L, 2.60 ± 0.50 mg/L and 1.68 ± 0.38 mg/L respectively (Table 1). The variation of BOD was significant between months and seasons ($P < 0.01$).

The mean $\pm$ SE of Alkalinity for Gimbawa, Saminaka and Zaria reservoirs observed were 5.05 ± 0.32 mg/L, 4.29 ± 0.31 mg/L and 6.77 ± 1.16 mg/L respectively (Table 1). The variation of Alkalinity was significant between months, reservoirs ($P < 0.05$) and between seasons ($P < 0.01$).

The mean $\pm$ SE of Hardness for Gimbawa, Saminaka and Zaria reservoirs observed were 1.26 ± 0.26 mg/L, 1.46 ± 0.30 mg/L and 1.49 ± 0.36 mg/L respectively (Table 1). These variations however, were only significant between months ($P < 0.05$) and not between months and seasons ($P > 0.05$).

Nitrate-nitrogen concentration for Gimbawa reservoir had a highest value of 0.19 mg/L and lowest of 0.03mg/L. Saminaka reservoir had a highest value of 0.16 mg/L and lowest of 0.02mg/L. Zaria reservoir had its highest value of 0.55 mg/L and lowest of 0.01 mg/L. The mean $\pm$ SE Nitrate-nitrogen concentration for Gimbawa, Saminaka and Zaria reservoirs observed were of 0.01 mg/L, 0.09 ± 0.05 mg/L and 0.13 ± 0.05 mg/L respectively (Table 1). These variations however, were not statistically significant between reservoirs, months and seasons ($P > 0.05$).

For phosphate-phosphorus concentration, Gimbawa had its highest value of 0.62mg/L and lowest of 0.18mg/L. Saminaka reservoir had the highest concentration of 0.76mg/L and lowest of 0.04mg/L. Zaria reservoir had its highest value of 0.8mg/L and lowest of 0.04mg/L. The mean $\pm$ SE of Gimbawa, Saminaka and Zaria reservoirs observed were 0.29 ± 0.06 mg/L, 0.39 ± 0.08 mg/L and 0.39 ± 0.08 mg/L respectively (Table 1). These variations however, were only significant between months ($P < 0.01$) but not between reservoirs and seasons ($P > 0.05$).

4.1 Metal ions

The lowest concentrations of Cu, Zn, Mn, Fe and Cr were below detectable limits in the three reservoirs. The highest concentration of Cu, Zn and Cr was recorded in Zaria reservoir (0.39, 0.50 and 1.10 mg/L respectively). Gimbawa reservoir had the highest concentration of Mn (1.01mg/L) and Fe (1.14mg/L). The mean $\pm$ SE of these metals in Gimbawa, Saminaka and Zaria respectively are Cu: 0.03 ± 0.03 mg/L, 0.03 ± 0.02 mg/L and 0.04 ± 0.03 mg/L; Zn: 0.03 ± 0.03 mg/L, 0.02 ± 0.01 mg/L and 0.04 ± 0.04 mg/L; Mn : 0.08 ± 0.08 , 0.09 ± 0.06 mg/L and 0.06 ± 0.06 mg/L mg/L; Fe: 0.28 ± 0.1 mg/L, 0.89 ± 0.43 mg/L and 0.51 ± 0.28 mg/L and Cr: 0.43 ± 0.07 mg/L, 0.36 ± 0.06 mg/L and 0.34 ± 0.08 .

Concentrations of Nickel in the three reservoirs showed the highest concentrations of 1.06, 1.0 and 1.17 mg/L; and lowest concentrations of 0.17, 0.26 and 0.17 mg/L for Gimbawa, Saminaka and Zaria reservoirs respectively (Table 2). The mean $\pm$ Standard Error for the reservoirs were 0.64 ± 0.08 mg/L, 0.62 ± 0.06 mg/L and 0.69 ± 0.10 mg/L for Gimbawa, Saminaka and Zaria reservoirs respectively (Table 2). These differences were however not significant between reservoirs, months and seasons ($P > 0.05$).

	Gimbawa			Saminaka			Zaria			MPL
	Min	Max	Mean $\pm$ SE	Min	Max	Mean $\pm$ SE	Min	Max	Mean $\pm$ SE	
Copper (mg/L)	ND	0.34	0.03 $\pm$ 0.03	$\pm$ ND	0.25	0.03 $\pm$ 0.02	$\pm$ ND	0.39	0.04 $\pm$ 0.03	$\pm$ 2mg/L*
Zinc (mg/L)	ND	0.3	0.03 $\pm$ 0.03	$\pm$ ND	0.17	0.02 $\pm$ 0.01	ND	0.5	0.04 $\pm$ 0.04	$\pm$ 3mg/L*
Manganese (mg/L)	ND	1.01	0.08 $\pm$ 0.08	ND	0.58	0.09 $\pm$ 0.06	$\pm$ ND	0.72	0.06 $\pm$ 0.06	$\pm$ 0.5mg/L*
Cadmium (mg/L)	0.06	0.22	0.14 $\pm$ 0.01	0.06	1.87	0.16 $\pm$ 0.02	$\pm$ 0.06	0.25	0.11 $\pm$ 0.02	$\pm$ 0.003mg/L*
Iron (mg/L)	ND	1.14	0.28 $\pm$ 0.1	$\pm$ ND	5.5	0.89 $\pm$ 0.43	$\pm$ ND	3.55	0.51 $\pm$ 0.28	$\pm$ 0.3mg/L*
Nickel (mg/L)	0.17	1.06	0.64 $\pm$ 0.08	$\pm$ 0.26	1	0.62 $\pm$ 0.06	$\pm$ 0.17	1.17	0.69 $\pm$ 0.10	$\pm$ 0.02mg/L*
Chromium (mg/L)	ND	0.96	0.43 $\pm$ 0.07	$\pm$ ND	0.67	0.36 $\pm$ 0.06	ND	1.1	0.34 $\pm$ 0.08	$\pm$ 0.05mg/L*
Calcium (mg/L)	2.33	41.67	7.70 $\pm$ 3.10	$\pm$ 1	20	6.4 $\pm$ 1.93	$\pm$ 1	40	5.6 $\pm$ 3.14	$\pm$ 200mg/L*
Magnesium (mg/L)	1.6	4.7	3.01 $\pm$ 0.24	$\pm$ 0.9	8.3	3.19 $\pm$ 0.74	0.8	5.1	2.59 $\pm$ 0.31	$\pm$ 0.02mg/L**
Potassium (mg/L)	2.6	8.5	4.80 $\pm$ 0.56	$\pm$ 2.4	9.4	4.8 $\pm$ 0.55	$\pm$ 2.8	6	4.2 $\pm$ 0.26	$\pm$ 200mg/L*
Sodium (mg/L)	8.9	14.5	12.19 $\pm$ 0.53	$\pm$ 6.4	27.3	11.49 $\pm$ 1.75	$\pm$ 6.8	15.9	9.84 $\pm$ 0.74	$\pm$ 200mg/L*

ND= not detectable, Min= minimum, Max= maximum, SE= Standard Error *WHO, 2006 ** Standard Organisation of Nigeria, 2007, MPL = maximum permissible limit

Table 2. Mean Values of Metal ions Observed in Gimbawa, Saminaka and Zaria reservoirs

The highest concentrations of 1.01, 0.58 and 0.5 mg/L of Manganese were observed in Gimbawa, Saminaka and Zaria reservoirs, the lowest concentrations of Manganese were below detectable limits in the three reservoirs. The mean $\pm$ SE concentration of Manganese was 0.08 ± 0.08 mg/L, 0.09 ± 0.06 mg/L and 0.11 ± 0.06 mg/L in Gimbawa, Saminaka and Zaria reservoirs respectively (Table 2).

The concentrations of Cadmium in the three reservoirs showed highest values of 0.22, 0.25 and 0.19 mg/L in Gimbawa, Saminaka and Zaria reservoirs respectively. The three reservoirs had lowest concentrations of 0.06 mg/L during the study period. The mean $\pm$ Standard Error for the reservoirs were 0.14 ± 0.01 mg/L, 0.16 ± 0.02 mg/L and 0.11 ± 0.02 mg/L for Gimbawa, Saminaka and Zaria reservoirs respectively (Table 2). These differences were however not significant between reservoirs, months and seasons ($P > 0.05$).

Magnesium concentration in the three reservoirs showed a highest concentration of 4.7 mg/L, 8.3 mg/L and 5.1 mg/L and lowest of 1.6, 0.9 and 0.8 for Gimbawa, Saminaka and Zaria reservoirs respectively. The mean $\pm$ SE for the reservoirs were 7.70 ± 3.10 mg/L, 6.4 ± 1.93 mg/L and 5.6 ± 3.14 mg/L for Gimbawa, Saminaka and Zaria reservoirs respectively (Table 2). These differences were however not significant between reservoirs and months ($P > 0.05$) but significant between seasons ($P < 0.05$).

The highest Sodium concentrations observed were 14.5, 27.3 and 15.9 mg/L and lowest of 8.9, 6.4, and 6.8 mg/L in Gimbawa, Saminaka and Zaria reservoirs respectively. The mean $\pm$ SE for the reservoirs were 12.19 ± 0.53 mg/L, 11.49 ± 1.75 mg/L and 9.84 ± 0.74 mg/L for Gimbawa, Saminaka and Zaria reservoirs respectively (Table 2). These differences were however not significant between reservoirs and months ($P > 0.05$) but significant between seasons ($P < 0.05$), with significant interaction between reservoirs and seasons ($P < 0.01$).

Gimbawa, Saminaka and Zaria reservoirs had the highest concentration of Potassium of 8.5, 9.4 and 6 mg/L and lowest of 2.6, 2.4 and 2.8 mg/L respectively. The mean $\pm$ Standard Error for the reservoirs were 4.80 ± 0.56 mg/L, 4.80 ± 0.53 mg/L and 4.2 ± 0.26 mg/L for Gimbawa, Saminaka and Zaria reservoirs respectively (Table 2). These differences were however not significant between reservoirs, months and seasons ($P > 0.05$).

The three reservoirs had the highest Iron concentrations of 1.14 mg/L (Gimbawa), 5.4 mg/L (Saminaka) and 3.55 mg/L (Zaria). The lowest concentrations of Iron were below detectable limits in the three reservoirs. The mean $\pm$ Standard Error for the reservoirs were 0.28 ± 0.1 mg/L, 0.89 ± 0.43 mg/L and 0.51 ± 0.28 mg/L for Gimbawa, Saminaka and Zaria reservoirs respectively (Table 2). These differences were however not significant between reservoirs, months and seasons ($P > 0.05$).

4.2 Cyanobacteria

Gimbawa reservoir had its highest number of cyanobacteria cells/L in the month of December (112) and lowest in the month June and August (0 cells/L). Saminaka reservoir had its highest number in the month of March (292 cells/L) and lowest in the months of June and January

(4 cells/L). Zaria reservoir had its highest abundance in October (88 cells/L) and lowest in the month of May (32 cells/L) (Table 3).

Number of taxa (8), number of individuals (308), Shannon Index (1.59) and Simpson index (0.76) was observed in Gimbawa reservoir during the dry season was higher than that observed in the wet season (4, 152, 1.11 and 0.62 respectively). Dominance was higher in the wet season (0.38) than dry season (0.24).

Reservoir	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr	Total
Gimbawa	16	0	28	0	68	12	60	112	12	40	40	44	432
Saminaka	24	12	40	32	60	40	48	52	4	68	292	220	892
Zaria	32	68	36	36	88	188	120	76	68	76	48	4	840

Table 3. Monthly Abundance (cells/L) of Cyanobacteria in Gimbawa, Saminaka and Zaria reservoirs

In Saminaka reservoir the trend was similar, number of individuals (464), Shannon Index (0.97) and Simpson index (0.56) observed during the dry season was higher than that observed in the wet season (236, 0.81 and 0.41 respectively), dominance was higher in the wet season (0.38) than dry season (0.24). The only exception was that the number of taxa observed in both seasons was equal (5).

Diversity Index	Gimbawa		Saminaka		Zaria	
	Wet	Dry	Wet	Dry	Wet	Dry
Taxa_S	4	8	5	5	4	6
Individuals	152	308	236	464	448	392
Dominance_D	0.38	0.24	0.59	0.44	0.53	0.31
Shannon_H	1.11	1.59	0.81	0.97	0.88	1.38
Simpson_1-D	0.62	0.76	0.41	0.56	0.47	0.69

Table 4. Seasonal Diversity Indices of Cyanobacteria in Gimbawa, Saminaka and Zaria reservoirs

In Zaria reservoir, the dry season a higher number of taxa (6), Shannon index (1.4) and Simpson index (0.69) were observed than the wet season (4, 0.88 and 0.47 respectively). While dominance (0.53) and number of individuals (448) observed in the wet season were higher than that observed in the dry season (0.31 and 392 respectively) (Table 4).

4.3 Relationship between physico-chemical characteristics and phytoplankton

In Gimbawa reservoir significant positive correlation was observed between Mg and *Sacconema* sp (r = 0.43) and *Trichodesmium* sp (r = 0.43) and between Fe and *Arthrospira* sp (0.43) and *Borzia* sp (0.43) (Table 5). pH and Electrical Conductivity showed significant positive correlation with *Arthrospira* sp (0.75 and 0.98 respectively); *Borzia* sp (0.75 and 0.98

respectively) and *Merismopedia* sp (0.51 and 0.64 respectively); BOD with *Merismopedia* sp (0.55) (Table 6).
In Saminaka reservoir, significant positive correlation was observed between Chromium with *Oscillatoria* sp ($r = 0.40$); Nickel with *Gleocystis* sp ($r = 0.63$), *Microcystis* sp (0.67) and *Trichodesmium* sp (0.45); Cadmium with *Gleocystis* sp ($r = 0.82$), *Microcystis* sp (0.88) and Iron with *Microcystis* sp (0.66). Significant negative correlation was observed between Potassium and *Spirulina* sp (-0.45); Sodium with *Oscillatoria* sp (-0.48) and *Sacconema* sp (-0.64); Chromium with *Merismopedia* sp (-0.49) (Table 5). *Microcystis* sp was observed to show significant positive correlation with DO (0.42), BOD (0.49), Alkalinity (0.64), $\text{NO}_3\text{-N}$ (0.45). It showed significant negative correlation with Transparency (-0.40) and $\text{PO}_4\text{-P}$ (-0.54). *Oscillatoria* sp showed significant positive correlation with Air Temperature (0.53), DO (0.50) and Alkalinity (0.50). *Spirulina* sp showed significant positive correlation with BOD (0.52) (Table 6).

	K	Na	Mg	Cr	Ni	Cd	Fe
Gimbawa							
<i>Arthrospira</i> sp	0.13	-0.12	0.04	-0.13	0.07	0.36	0.43*
<i>Borzia</i> sp	0.13	-0.12	0.04	-0.13	0.07	0.36	0.43*
<i>Merismopedia</i> sp	-0.09	0.11	-0.09	0.15	-0.06	0.1	0.24
<i>Oscillatoria</i> sp	-0.35	0.23	-0.15	0.18	-0.3	-0.22	0.04
<i>Sacconema</i> sp	0.01	-0.12	0.43*	0.1	-0.08	0.12	-0.11
<i>Spirulina</i> sp	0.34	-0.18	-0.06	-0.39	0.14	0.26	0.17
<i>Spondylosium</i> sp	-0.36	-0.3	-0.04	-0.3	0.29	0.25	-0.23
<i>Trichodesmium</i> sp	0.01	-0.12	0.43*	0.1	-0.08	0.12	-0.11
Saminaka							
<i>Gleocystis</i> sp	-0.19	0.02	0.08	0.24	0.63**	0.82**	0.76
<i>Merismopedia</i> sp	0.11	0.55*	0.33	-0.49*	0.07	0.04	0.12
<i>Microcystis</i> sp	-0.24	-0.17	0.10	0.13	0.67**	0.88**	0.66**
<i>Nostoc</i> sp	0.12	0.26	0.02	-0.18	0.05	-0.05	-0.02
<i>Oscillatoria</i> sp	-0.11	-0.48*	-0.31	0.40*	0.19	-0.22	0.29
<i>Rivularia</i> sp	-0.04	0.20	-0.18	-0.31	-0.27	-0.25	0.02
<i>Sacconema</i> sp	0.35	-0.64*	0.34	0.01	-0.32	-0.19	-0.21
<i>Spirulina</i> sp	-0.45*	-0.33	-0.08	-0.12	-0.31	-0.10	-0.26
<i>Spondilosium</i> sp	0.13	0.28	0.06	-0.01	0.28	0.30	-0.16
<i>Trichodesmium</i> sp	0.27	0.09	-0.04	-0.01	0.45*	0.08	-0.01
Zaria							
<i>Merismopedia</i> sp	-0.38	-0.02	-0.14	-0.20	-0.18	-0.31	-0.16
<i>Nostoc</i> sp	0.12	0.26	0.02	-0.18	0.05	-0.05	-0.02
<i>Oscillatoria</i> sp	-0.11	-0.48**	-0.31	0.40*	0.19	-0.22	0.29
<i>Sacconema</i> sp	-0.49*	0.03	-0.15	-0.18	0.05	-0.01	-0.21
<i>Spirulina</i> sp	-0.45*	-0.33	-0.08	-0.12	-0.31	-0.10	-0.26
<i>Spondilosium</i> sp	0.13	0.28	0.06	-0.01	0.28	0.30	-0.16
<i>Trichodesmium</i> sp	0.27	0.09	-0.04	-0.01	0.45	0.08	-0.01

*Significant $P < 0.05$, ** Significant $P < 0.05$

Table 5. Correlation Coefficient (r) between Cyanobacteria and Metal Ion Concentration in Gimbawa, Saminaka and Zaria reservoirs

In Zaria reservoir, Significant negative correlation was observed between Potassium and *Sacconema* sp ($r = -0.49$), *Spirulina* sp ($r = -0.45$); Sodium with *Oscillatoria* sp ($r = -0.48$). Chromium showed significant positive correlation with *Oscillatoria* sp (0.40) (Table 5).

Species	Air Temp	Water Temp	Transp	pH	EC	DO	BOD	Alkalinity	Hardness	NO ₃ -N	PO ₄ -P
Gimbawa											
<i>Arthrospira</i> sp	0.08	0.32	-0.26	0.75*	0.98**	0.25	0.25	0.27	-0.11	-0.04	0.22
<i>Borzia</i> sp	0.08	0.32	-0.26	0.75*	0.98**	0.25	0.25	0.27	-0.11	-0.04	0.22
<i>Merismopedia</i> sp	0.31	0.33	0.02	0.51*	0.64**	0.32	0.55**	0.04	0.00	-0.07	0.23
Saminaka											
<i>Microcystis</i> sp	0.13	0.08	-0.40*	0.37	0.34	0.42*	0.49*	0.64**	0.03	0.45*	-0.54**
<i>Oscillatoria</i> sp	0.53**	0.18	0.35	0.29	0.12	0.50**	0.27	0.50**	-0.15	0.14	0.15
<i>Spirulina</i> sp	0.14	0.39	-0.06	0.22	0.05	0.08	0.52**	0.28	-0.12	0.26	-0.01
Zaria											
<i>Nostoc</i> sp	-0.50**	-0.80**	0.62**	0.36	-0.15	0.11	0.02	0.14	0.00	-0.01	-0.39
<i>Spirulina</i> sp	0.14	0.24	-0.22	-0.29	-0.20	0.00	0.01	-0.15	-0.28	-0.09	0.48*
<i>Spondilosium</i> sp	0.76**	0.23	-0.02	0.20	-0.06	0.59**	0.44*	0.45*	-0.22	0.02	0.19

*Significant $P < 0.05$, **Significant $P < 0.05$

Table 6. Correlation Coefficient between Cyanobacteria and Physico-chemical Parameters in Gimbawa, Saminaka and Zaria reservoirs

5. Discussion

The statistically significant monthly variation of mean Air Temperature in the three reservoirs could be attributed to the low temperatures experienced between the months of November and February as a result of the harmattan wind (Ezra and Nwankwo, 2001). The significantly higher Transparency of the water in Gimbawa reservoir may be attributed to the low human pressure in its catchment as it is location in the outskirts of a major human settlement. Thus, receiving low amount of silt and nutrients that stimulate algal and cyanobacterial growth. Silt and plankton abundance have been implicated in Transparency fluctuations (Davies *et al*, 2009). The circum-neutral mean pH of water in the reservoirs may be attributed to the relatively high alkalinity values of the reservoirs, which is effective as a buffer to fluctuations of pH that might be caused by introduction of waste water, photosynthesis and other metabolic processes. The wide fluctuations of EC (SE in the range of 38.59 to 41.95) an significantly monthly variations in the three

reservoirs may be attributed to concentration of Electrical Conducting ions due to evaporation during the extensive (six) months of dry season. Similar results were obtained by Chia and Bako (2008). DO concentration was found within the limit of 5-9 mg/l, drinking water (UNESCO/WHO/UNEP, 1996). The mean BOD values of Gimbawa and Saminaka were slightly above the 2 mg/l. BOD above 2 mg/l is associated with waste water contamination (UNESCO/WHO/UNEP, 1996). The mean hardness values (< 1.5 mg/l) may be due to the uptake of the ions (calcium and magnesium) responsible for harness of water by aquatic organisms. Calcium and Magnesium are essential for aquatic organisms such as algae, cyanobacteria, aquatic macrophytes as well as other reptiles. The Mean NO₃-N (1.2 and 1.3 mg/l in Gimbawa and Zaria reservoirs) and PO₄-P (0.29 in Gimbawa reservoir and 0.39 in Saminaka and Zaria reservoirs) were found to be above expected concentration range of natural unpolluted waters of 0.1mg/l and 0.001mg/l respectively, and are capable of stimulating cyanobacterial bloom (UNESCO/WHO/UNEP, 1996).

Metal ions such as Manganese, Iron, Cadmium, Nickel, Chromium, Magnesium showed concentrations higher than the maximum permissible limit for WHO (2006) and SON (2007), other metals like Copper, Zinc, Sodium and Potassium were found to be below the maximum permissible limit. The implication of high concentrations of metal ions in drinking water include: Manganese causes neurological disorder and at concentrations exceeding 0.1mg/L it causes undesirable taste in beverages, stains laundry and may lead to the accumulation of deposit in water distribution system. Iron at levels above 0.3mg/L stains laundry and plumbing fixtures (WHO, 2006). Cadmium is toxic to the kidney, Chromium is carcinogenic, and Magnesium affects consumer acceptability of drinking water (SON, 2007). Zinc imparts an undesirable astringent taste to water at threshold concentration of 4mg/L, water containing Zinc at excess of 3-5mg/L may appear apalacent and develop greasy film on boiling (WHO, 2006). With the exception of Magnesium, all the others are heavy metals capable of accumulating in increasing concentration as they move up the food chain (Chindah *et al*, 2004).

The dynamics of the concentration of these metals may be attributed to inflow of waste water from residential areas (as they are components of many household products such as pesticides, fungicides, paints, batteries and plumbing facilities), remobilization from sediment due to fluctuations of pH, inflow of agro-chemicals such as fertilizers, pesticides and herbicides from farms in the catchment of the reservoirs, chemicals from washing of automobiles. Changes in pH affects the solubility of metal ions, lowering of pH may remobilize insoluble metal complexes adsorbed on clay and silica in the sediments into the water column, for example at pH 6.7, Zinc is available to form complexes with organic matter while at pH > 7 Zinc begin to hydrolyze and form stable Zn(OH)₂ at pH 8. It is important to note that there is a difference between the presence of a metal and its bioavailability. A metal may be present in a form that is not available for utilization by algae and other organisms (Kalis, 2006).

The variation in abundance of Cyanobacteria (Saminaka > Zaria > Gimbawa) in the reservoirs during the study period may be attributed to the variation in N: P ratio (Gimbawa, 0.41; Saminaka, 0.23 and Zaria, 0.33) of the water bodies. Lower N: P ratio promotes cyanobacteria abundance (Tisser *et al.*, 2008). They take advantage of their ability

to fix nitrogen into the aquatic environment, thus enabling them to out-compete other divisions (Chorus and Batram, 1999). High phosphate concentration may result from detergents from sewage, washing of cars, clothes and from fertilization of farms in the catchment of the reservoirs (Reynolds, 1998). High abundance of Cyanobacteria in drinking water may be a serious problem in drinking water as they produce toxins which are harmful to fish, livestock, other aquatic organisms and ultimately man (WHO, 2006). The presence of a bloom of species *Microcystis*, a toxin producing genus in the Saminaka reservoir is worrisome.

The differences in number of taxa and number of individuals between seasons may be due differences in temperatures and pH as different species obtain nutrition at different pH and temperatures. Wilm and Dorris (1966) have suggested a relationship between species diversity and pollution status of aquatic system and classified as follows; > 1 = Clean water, 1-3 = moderately-polluted < 1 = Heavily- polluted. Based on this classification, Gimbawa reservoir was moderately polluted in both seasons, Saminaka reservoir heavily polluted in both seasons while the Zaria reservoir was heavily polluted in the wet season and moderately polluted in the dry season. A similar classification was also used by Shehata *et al.* (2009). Simpson index gives the evenness of species distribution; the higher evenness in species distribution in the dry season may be an indication that the water quality was better to support the growth of most of the species observed.

Significant positive correlation between cyanobacteria and metal ions may be an indication of the possibility of using as indicators of the levels of these metals in a water body. These metals are either essential (Fe, Cu, Zn, Na, Ca, Mn, Co and K) or beneficial (Ni and As) in phytoplankton physiological processes (Paerles-Vela, *et al.*, 2006). Significant negative correlation between metal ions and cyanobacteria may be an indication of toxicity of these metals at high concentrations level exceeding the requirement for nutrition or increased utilization of metals in periods of high abundance. Some of the metals that show significant negative correlation with cyanobacteria abundance are either essential (Fe, Cu, Zn, Na, Ca, Mn, Co and K) or beneficial (Ni and As) in cyanobacteria physiological processes (Daffus, 2002). Significant negative correlation between metal ions and cyanobacteria may be an indication of toxicity of these metals at high concentrations level exceeding the requirement for nutrition or increased utilization of metals in periods of high abundance ((Paerles-Vela, *et al.*, 2006).

Significant positive correlation between cyanobacteria with pH and alkalinity may be due to the fact that some essential elements are bioavailable at certain required pH, on the other hand, the bioavailability of certain elements at toxic concentrations as affected by pH may cause a significant negative correlation between pH and alkalinity with cyanobacterial abundance. A significant positive correlation between nutrient (N and P) load and cyanobacteria abundance may be due to the fact that increased nutrient concentrations leads to a resultant increase in cyanobacteria abundance and a significant negative correlation may due to the reason that increased cyanobacteria abundance may lead to increased utilization of such nutrients by cyanobacteria (Rabalais, 2002). The significant relationship(s) between cyanobacteria abundance and DO, BOD, EC, Hardness, Temperature and Transparency is an indication of the inter-dependance between these important water quality characteristics and the Biota (Shehata *et al.*, 2009).

6. Conclusion

The introduction of waste water into these reservoirs greatly impairs the water quality of these reservoirs. The consequence is seen as the elevated concentration of heavy metals such as Cadmium, Iron, Nickel and Chromium above WHO permissible limit in drinking water. Waste water is also implicated in the increased nutrient (N and P) levels. These nutrients were found to be below concentrations to cause any harm directly to consumers but indirectly by their ability to stimulate cyanobacterial growth. Shannon-Weiner diversity index showed that the water quality of the three reservoirs follows this order Gimbawa > Zaria > Saminaka.

7. References


- Adakole J.A, Balogun J.K. and Lawal, F.A. (2002). Water Quality Impacts Assessment Associated with an Urban Stream in Zaria, Nigeria. *NISEB Journal* 2(3): 195-203.
- Adegoroye, A. (2008). The Challenges of Environmental Enforcement in Africa: The Nigerian Experience. *Proceedings of the Third International Conference of Environmental Enforcement*, Pp 43 – 52.
- Akin-Oriola, G.A. (2003). On the Phytoplankton of Awba reservoir, Ibadan, Nigeria. *Revista de Biol.* 51: 1-15.
- APHA (1998). Standard Methods for the Analysis of Water and Wastewater. American Public Health Association, New York. 1287pp.
- Borgh, M.V. (2004).
[http:// h2o.enr.state.nc.us/esb/EU.html](http://h2o.enr.state.nc.us/esb/EU.html)
- Chia, A.M and Bako, S.P. (2008). Seasonal Variation of Cyanobacteria in Relation to Physicochemical Parameters of some Fresh Water Ecosystems in the Nigerian Guinea Savanna. *Proceedings of Taal 2007: 12th World Lake Conference, Published by Ministry of Environment and Forests, India and International Lake Environment Committee Foundation (ILEC).* Pp.1383-1387.
- Chindah, A.C., Braide, A.S. and Sibendu, O.C. (2004). Distribution of Hydrocarbons and Heavy Metals in Sediment and Cinstaceen (Shrimps-Pannens Notialis) from the Bonny/New Calabar River Eastuary. *A JEAM-RAGEE*, 9:1-17.
- Chorus, I. and Bartram, J. (1999). *Toxic Cyanobacteria in Water: a Guide to Public Health Consequence, Monitoring and Management*. World Health Organization, London, pp.15-40.
- Daffus, J.H. (2002). "Heavy Metals" – A Meaningless Term? *Pure Appl. Chem.*, 74(5):793-807.
- Davies, O.A., Abowel, J.F.N. and Tawari, C.C. (2009). Phytoplankton Community of Elechi Creek, Niger Delta Nigeria – A Nutrient-Polluted Tropical Creek. *American Journal of Applied Science* 6(6):1143-1152.
- Ezra, A.G. and Nwankwo (2001). Composition of Phytoplankton Algae in Gubi Reservoir, Bauchi, Nigeria. *Journal of Aquatic Science* 16(2)115-118.
- Fakayode, S.O. (2005). Impact Assessment of Industrial effluent on Water Quality of the Receiving Alaro River in Ibadan, Nigeria, *AJEAM-RAGEE*: 10:1-13.
- Haider, S., Naithani V., Viswanathan, P.N. and Kakkar P. (2003). Cyanobacterial Toxins: A Growing Environmental Concern. *Chemosphere*, 52:1-21.

- Kadiri, M.O. (2006). Phytoplankton Flora and Physicochemical attributes of some waters in the Eastern Niger-Delta area of Nigeria. *Nigerian Journal of Botany*, 19(2): 188-200.
- Kalis, E.J.J. (2006). Chemical speciation and bioavailability of heavy metals in soil and surface water. Ph.D. thesis Wageningen University, Wageningen, the Netherlands. Pp 142.
- Kimura, H. (2005). The Japanese New Law for Preservation of Lake Water Quality. In *Order of Proceeding World Lake Conference*, Nairobi, Kenya. Pp13.
- Lenat, O.E., Smock, L.A. and Penrose, O.L.(1980). Use of Benthic macroinvertebrates as indicators of environmental effect. Douglas L.W. (Ed) Pub. Lexington Books, Toronto, 99.
- Mur, L.R., Skurlberg, O.M. and Utkilen, H. (1999). Cyanobacteria in the Environment. In: Sperling, E.V. and Gomes, L.N.L. (2008). Cyanotoxin Generation in Tropical Water Supply reservoir. *Proceedings of Taal 2007: 12th World Lake Conference*, Published by Ministry of Environment and Forests, India and International Lake Environment Committee Foundation (ILEC. pp) 451-455.
- Onrauwa, P.C and Fakayode, S.O. (2002). Heavy Metals Contamination of Soil and Bioaccumulation in Annex Evacs (*Panicum maxima*) Around Ikaja Industrial Estate, Lagos, Nigeria. *Environmental Geology*, 43:45-150.
- Perales-Vela, H.V., Pena-Castro, J.M. and Canizares-Villanueva (2006). Heavy Metal detoxification in eukaryotic microalgae. *Chemosphere* 64: 1-10.
- Perry, R. (2003). A Guide to the marine plankton of southern California. <http://www.msc.ucla.edu/oceanglobe>
- Prescott, G.W. (1977). *The Fresh water Algae*. WMC Brown Company Publishers Dubugue, IOWA. P 12.
- Rabalais, N.N. (2002). Nitrogen in Aquatic Ecosystems. *BioOne*, 31: 102-112.
- Reynolds C.S (1998). What Factors Influence the Species Composition in Lakes of Different Trophic Status? *Hydrobiology*, (369/370):11-26.
- Shehata, S.A., Badr, S.A., Ali, G.H., Ghazy, M.M., Moawad, A.K., and Wahba, S.Z. (2009). Assessment of Nile Water Quality via Phytoplankton changes and Toxicity Bioassay Test. *Journal of Applied Sciences*, 5(12): 2083-2095.
- SON (2007). Nigerian Standard for Drinking Water Quality. *Published by the Standard Organisation of Nigeria, Abuja, Nigeria*. Pp 30.
- Tiseer, F. A., Tanimu, Y. and Chia, A. M. (2008). Seasonal occurrence of algae and physico-chemical parameters of Samaru stream, Zaria, Nigeria. *Asian journal of Earth Science* 1(1):31 – 37.
- UNESCO, WHO, UNEP (1996). *Water quality assessment- A guide to the use of biota, sediments and water in environmental monitoring*. E and FN Spon. Cambridge, Great Britain, 609 pp.
- Wilm, J.L. and Dorris, T.C.(1966). Species Diversity of Benthic Macroinvertebrates. In: A Stream Receiving Domestic and Oil Refinery Effluents. In: Islam, S.M. Phytoplankton diversity index with reference to mucalinda serovar Bodh-Gaya. *Order of proceedings of Taal 2007: 12th World Lack Conference*. Published by Ministry of Environment and Forests, India and International Lake Environment Committee Foundation (ILEC). pp 462-463.

World Health Organisation(2006). *Guidelines for drinking water Quality*. (2ed) (addendum to vol.1). Recomendations. WHO Press, Geneva Switzerland. Pp 595.

IntechOpen

IntechOpen



Waste Water - Evaluation and Management

Edited by Prof. Fernando Sebasti  n Garc  a Einschlag

ISBN 978-953-307-233-3

Hard cover, 470 pages

Publisher InTech

Published online 01, April, 2011

Published in print edition April, 2011

Fresh water resources are under serious stress throughout the globe. Water supply and water quality degradation are global concerns. Many natural water bodies receive a varied range of waste water from point and/or non point sources. Hence, there is an increasing need for better tools to assess the effects of pollution sources and prevent the contamination of aquatic ecosystems. The book covers a wide spectrum of issues related to waste water monitoring, the evaluation of waste water effect on different natural environments and the management of water resources.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Yahuza Tanimu, Sunday Paul Bako and John Ameh Adakole (2011). Effects of Domestic Waste Water on Water Quality of Three Reservoirs Supplying Drinking Water in Kaduna State - Northern Nigeria, Waste Water - Evaluation and Management, Prof. Fernando Sebasti  n Garc  a Einschlag (Ed.), ISBN: 978-953-307-233-3, InTech, Available from: <http://www.intechopen.com/books/waste-water-evaluation-and-management/effects-of-domestic-waste-water-on-water-quality-of-three-reservoirs-supplying-drinking-water-in-kad>

INTech
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike-3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/), which permits use, distribution and reproduction for non-commercial purposes, provided the original is properly cited and derivative works building on this content are distributed under the same license.

IntechOpen

IntechOpen