

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

185,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Short-Term Exposure to Outdoor Pollutants and Loss of Pulmonary Function in the Elderly

Yun-Chul Hong and Jin Hee Kim

*Department of Preventive Medicine,
Seoul National University College of Medicine,
Institute of Environmental Medicine,
Seoul National University Medical Research Center,
Korea*

1. Introduction

The reduction of pulmonary function has come to be considered one of the most important health effects of air pollution because it not only affects individual respiratory health status but also predicts cardiovascular morbidity and mortality in the general population (1-3). Although numerous studies have provided an understanding of air pollution's general effects on pulmonary function, the length of the lag period for the effects of short-term exposure to air pollution has not been well elucidated.

In addition, most of the information about the pulmonary effects of air pollution has come from studies of children and adults, whereas pulmonary effects in the elderly are less well understood. Because of the considerable increase in the prevalence of respiratory diseases in the elderly population and this population's known susceptibility to toxic chemicals in the environment, the effect of air pollution on pulmonary function in the elderly has emerged as a serious concern in both public health and clinical medicine.

Although the diminution of pulmonary function because of exposure to air pollutants has been reported in previous studies (4-6), most of these reports were time-series or cross-sectional analyses that did not capture within-subject changes. To account for within-subject changes in pulmonary function, a longitudinal study, such as a cohort or panel study, is required. A cohort study would be more appropriate for evaluating the long-term health effects of air pollution, while a panel study is better suited for evaluating the short-term effects (7). Because each subject can be used as his or her own control in a panel study with repeated measurements, only rapidly changing covariates for each given subject have to be considered in a statistical analysis.

Therefore, we conducted a longitudinal panel study using repeated measurements to evaluate the effects of air pollutants on pulmonary function in the elderly, who are known to be especially susceptible to air pollution (8-10).

2. Methods

2.1 Subject population

The study population consisted of 560 elderly people (all 60 years old or older) living in Seoul, Korea. Up to 5 spirometric tests were performed on different days for each subject.

During each subject's first visit, demographic data and information, such as medical history, dietary habits, smoking and alcohol consumption were gathered using a questionnaire. Written informed consent was obtained from every subject; this study was approved by the institutional review board of the Seoul National University College of Medicine.

2.2 Air pollution and meteorological data

Air pollution data were collected from the Research Institute of Public Health and Environment; values from the monitoring site nearest to each subject's residence were used as the dataset for that subject's exposure to ambient pollutants. Daily average values for concentrations of particulate matter with a diameter of 10 μm or less (PM_{10}), sulfur dioxide (SO_2), nitrogen dioxide (NO_2), carbon monoxide (CO) and ozone (O_3) were computed for each site. Data on the daily average temperature ($^{\circ}\text{C}$) and relative humidity (%) for each day of the study period were obtained from the Korean Meteorological Administration.

2.3 Spirometric measurements

Spirometric testing was conducted at each health evaluation visit according to the 2005 European Respiratory Society/American Thoracic Society recommendations. All tests were performed at the same hour of the morning by one trained technician, using a Microlab® (Sensormedic, USA) spirometer. The spirometric measurements used in our analyses were forced vital capacity (FVC), forced expiratory volume in one second (FEV_1), FEV_1 as a percentage of FVC (FEV_1/FVC), and forced expiratory flow between 25 and 75% of the FVC (FEF_{25-75}).

2.4 Statistical analysis

Linear mixed-effect models were used to estimate the effects of the air pollutants on the pulmonary function test parameters, which controlled for temperature, relative humidity, smoking, dietary habits, alcohol consumption, and individual characteristics. We treated age, sex, height, weight, dietary habits, alcohol consumption, date, temperature, relative humidity, smoking and pollutant levels as fixed effects. Participant identity was treated as a random effect in the models. We examined the ongoing effects of air pollution on lung-function parameters over a lag time of 7 days after exposure. We counted the day of the air-pollutant exposure as lag day 0; lag day 7 was therefore defined as an exposure the 7 days prior. We also evaluated the cumulative lag effects of air pollution using a lag-distribution model.

3. Results

The mean age of the participants was 70.7 years old; 26.1% were male and 73.9% were female. Subjects underwent a mean of 3.3 health examination visits out of 5 possible visits. The male participants were more likely to be smokers than the female Participants (Table 1). Table 2 presents the levels of air pollutants and the meteorological data for the time periods preceding each health examination visit. The mean levels of PM_{10} , SO_2 , O_3 , NO_2 and CO were 42.6 $\mu\text{g}/\text{m}^3$, 3.9 ppb, 19.2 ppb, 35.6 ppb and 0.6 ppm, respectively.

To determine the length of the effective time period for pulmonary impact, we assessed the effects of air pollutants on lung function from the day of exposure (lag day 0) through lag day 7 (Figure 1). To estimate the effects of the air pollution, we calculated the estimated values of lung function parameters assuming average values for age, sex, height, weight, smoking, dietary habits, alcohol consumption, outdoor temperature and humidity. We

found that the effective time period for an air pollution-induced reduction of pulmonary function was from the day of exposure through the following 2 days. Compared to other air pollutants, O₃ did not show a significant association in reducing pulmonary function. We specifically calculated changes in pulmonary function due to exposure to air pollution within the past one day, because the calculated reduction in pulmonary function was most prominent for most air pollutants at lag day 1. Table 3 shows that the levels of PM₁₀, SO₂, NO₂ and CO were negatively associated with FEV₁/FVC and FEF₂₅₋₇₅ but that O₃ levels were not significantly associated with changes in pulmonary function parameters.

	Male	Female	Total
No. of participants (%)	146 (26.1)	414 (73.9)	560 (100)
No. of visits [mean ± SE]	3.1 ± 0.1	3.4 ± 0.1	3.3 ± 0.1
Age [mean (range)]	71.4 (62-84)	70.5 (60-87)	70.7 (60-87)
Height [mean ± SE (cm)]	164.3 ± 0.4	151.3 ± 0.2	154.7 ± 0.3
Weight [mean ± SE (Kg)]	65.7 ± 0.8	57.1 ± 0.4	59.3 ± 0.4
No. of smokers (%)	64 (43.4)	5 (1.2)	67 (12.2)
Lung-function indicators:			
FVC [mean ± SE (L)]	2.86 ± 0.05	20.4 ± 0.02	2.27 ± 0.03
FEV ₁ [mean ± SE (L)]	2.31 ± 0.04	1.74 ± 0.02	0.92 ± 0.02
FEV ₁ /FVC [mean ± SE (%)]	81.37 ± 0.64	86.70 ± 0.35	85.21 ± 0.33
FEF ₂₅₋₇₅ [mean ± SE (L/sec)]	2.31 ± 0.07	2.10 ± 0.04	2.16 ± 0.03

FVC: forced vital capacity; FEV₁: forced expiratory volume in one second; FEF₂₅₋₇₅: forced expiratory flow between 25% and 75% of forced vital capacity.

Table 1. Subject characteristics at first health evaluation visit

Pollutant	Number of measurements	Mean (SD)	Selected percentiles				
			10th	25th	50th	75th	90th
PM ₁₀ (µg/m ³)	1850	42.6 (17.6)	21.9	29.3	39.7	53.1	63.0
SO ₂ (ppb)	1850	3.9 (1.5)	2.4	2.7	3.4	4.5	6.0
O ₃ (ppb)	1850	19.2 (8.4)	8.1	10.9	19.3	26.9	30.2
NO ₂ (ppb)	1850	35.6 (9.1)	24.0	29.3	35.8	39.7	45.3
CO (ppm)	1850	0.6 (0.2)	0.4	0.5	0.6	0.7	0.9
Temperature (°C)	1915	17.3 (9.0)	4.0	10.1	19.1	25.1	27.0
Humidity (%)	1915	63.2 (12.8)	47.9	53.0	63.6	72.5	80.4

Levels of air pollutants for the health-examination day and the seven preceding days; meteorological data for the health examination day.

Table 2. Air-pollutant levels and meteorological data




Fig. 1. The effect of PM₁₀, SO₂, O₃, NO₂, and CO on FEV₁/FVC and FEF₂₅₋₇₅ by single lag days. Changes in FEV₁/FVC (%) and FEF₂₅₋₇₅ (L/sec) were calculated per 10 µg/m³ of change in PM₁₀ levels, per 1 ppb of change in SO₂, per 10 ppb of change in O₃, per 10 ppb of change in NO₂, and per 0.1 ppm of change in CO, adjusted for age, sex, body mass index (BMI), smoking status, alcohol consumption, dietary habits, temperature and relative humidity.

Pollutant	Outcome	No.	Estimated Change	SE	p-Value	Lower 95% CI	Upper 95% CI
PM ₁₀	FEV ₁ (L)	1146	0.003	0.002	0.1541	-0.001	0.008
	FEV ₁ /FVC (%)	1146	-0.287	0.063	<0.0001	-0.410	-0.164
	FEF ₂₅₋₇₅ (L/sec)	1146	-0.014	0.005	0.0034	-0.023	-0.005
SO ₂	FEV ₁ (L)	1146	0.003	0.026	0.8972	-0.047	0.054
	FEV ₁ /FVC (%)	1146	-3.447	0.718	<0.0001	-4.856	-2.037
	FEF ₂₅₋₇₅ (L/sec)	1146	-0.157	0.055	0.0047	-0.266	-0.048
O ₃	FEV ₁ (L)	1146	0.004	0.004	0.3123	-0.004	0.013
	FEV ₁ /FVC (%)	1146	0.228	0.124	0.0671	-0.016	0.472
	FEF ₂₅₋₇₅ (L/sec)	1146	0.016	0.009	0.0968	-0.003	0.034
NO ₂	FEV ₁ (L)	1146	0.002	0.004	0.6592	-0.006	0.009
	FEV ₁ /FVC (%)	1146	-0.542	0.102	<0.0001	-0.743	-0.341
	FEF ₂₅₋₇₅ (L/sec)	1146	-0.027	0.008	0.0005	-0.043	-0.012
CO	FEV ₁ (L)	1143	0.001	0.002	0.4677	-0.002	0.005
	FEV ₁ /FVC (%)	1143	-0.300	0.053	<0.0001	-0.403	-0.197
	FEF ₂₅₋₇₅ (L/sec)	1143	-0.014	0.004	0.0005	-0.022	-0.006

Change in lung-function indicators per 10 µg/m³ of change in PM₁₀ levels; per 10 ppb of change in SO₂; per 10 ppb of change in O₃; per 10 ppb of change in NO₂; per 0.1 ppm of change in CO. The effect of each air pollutant on lung function was adjusted for age, sex, body mass index (BMI), smoking status, alcohol consumption, dietary habits, temperature and relative humidity.

Table 3. The relationship between air pollutants and lung-function indicators on lag day one.

To evaluate the cumulative effect of air pollutants on pulmonary function, we performed a lag-distribution model analysis and found that a significant reduction in pulmonary function occurred when subjects had been exposed to air pollutants during the previous 2 days, as in the single-day lag model analysis (Figure 2).

3.1 Discussion

Our study demonstrates that exposure to outdoor air pollutants exerts harmful effects on lung function in the elderly and that the lag period for this effect is from the day of exposure to 2 days later.

All of these effects were observed at pollution levels within the current air-quality standard in Korea, which sets the maximum 24-hour average at 100 µg/m³ for PM₁₀, 50 ppb for SO₂, and 60 ppb for NO₂ and the maximum 8-hour average at 60 ppb for O₃ and 9 ppm for CO.

Spirometry provides a simple, convenient and reproducible measure of pulmonary function. We examined the association between exposure to ambient air pollutants and lung function test results and found statistically significant correlations between air pollutant levels of PM₁₀, NO₂, SO₂ and CO and results of FEV₁/FVC and FEF₂₅₋₇₅ measurements. In contrast, FVC and FEV₁ results did not show any association with air pollutant levels. Though declines in FVC and FEV₁ have been found to be associated with occupational exposure to air pollutants, these two parameters do not change quickly during the early stages of disease (11). This stasis suggests that the primary site affected by air pollutants might be the airways, especially the small airways, rather than the pulmonary parenchyma.




Fig. 2. The effect of PM₁₀, SO₂, O₃, NO₂, and CO on FEV₁/FVC and FEF₂₅₋₇₅ by distributed lag days. Changes in FEV₁/FVC (%) and FEF₂₅₋₇₅ (L/sec) were calculated per 10 μg/m³ of change in PM₁₀ levels, per 1 ppb of change in SO₂, per 10 ppb of change in O₃, per 10 ppb of change in NO₂, and per 0.1 ppm of change in CO, adjusted for age, sex, body mass index (BMI), smoking status, alcohol consumption, dietary habits, temperature and relative humidity.

The estimated changes for the FEV₁/FVC and FEF₂₅₋₇₅ results were -0.287% and -14 ml/sec with a change in PM₁₀ levels of 10 µg/m³. An increase of 10 ppb for SO₂ reduced FEV₁/FVC results by -3.447% and FEF₂₅₋₇₅ results by -157 ml/sec. An increase of 10 ppb in NO₂ reduced FEV₁/FVC by -0.542% and FEF₂₅₋₇₅ by -27 ml/sec, and an increase of 100 ppb in CO reduced FEV₁/FVC by -0.300% and FEF₂₅₋₇₅ by -14 ml/sec. Although these levels of reduction do not seem to be clinically significant for young adults, the impact of such air pollution on the already declining lung function of the elderly could be considerable.

The effect of air-pollution exposure on lung function is not necessarily contemporaneous (7). When we examined the pattern of lag in the relationship between air pollutants and lung function using single-lag-day and distributed-lag-day models, we found that the adverse effects of exposure to air pollutants could be detected immediately on the day of the exposure and that these effects had a lag period of 2 days except in the case of O₃, which did not show any effect during the test period. Lag structures in relationships between air pollution and respiratory health outcomes have been discussed in several studies. Goldberg et al. reported that the effect on the rate of respiratory-related deaths in persons older than 65 years was greatest following the same-day exposure (12). Braga et al. reported that respiratory deaths were affected more by exposure in the previous 1 to 2 days (13). Peters et al. reported stronger cumulative effects than the same-day effects of air pollution for asthmatic symptoms and peak expiratory flow rates (14). Our results are consistent with the short-term lag effects found in previous studies, showing that the cumulative effects of air pollution exposure on lung function were greater than the single-day lag effects, although the effect was no longer significant after lag day 2 for most of the air pollutants.

In our study, O₃ levels were not associated with changes in pulmonary function, although other pollutants were significantly associated. Estimating the effects of O₃ on lung function is challenging because of the strong correlation between O₃ levels and outdoor temperatures. However, in our results, the estimated risk level resulting from O₃ was constant, whether or not temperature was analyzed as a confounding variable.

The strengths of the present study merit further discussion. First, the panel study design, with repeated measurements on spirometric tests for each participant, provided a good opportunity to evaluate the short-term effects of the changes in air pollutant levels over time. In this longitudinal study, the subjects served as their own controls because personal characteristics could be assumed to remain constant over the study period. We used mixed-effects models to assess the effects of visit-to-visit variations in ambient air pollution on lung function to account for inter- and intra-subject variability. Individual exposure to other pollutants or other confounding factors could have biased the results if they had increased concomitantly with the measured air-pollution levels; however, this scenario is not likely.

There were also some limitations to this study. We could not completely control for those environmental factors that could have changed concomitantly with the levels of air pollutants, which could have biased the results for air-pollution effects. For example, possible environmental factors that could have confounded the results were meteorological variables, such as temperature and humidity. However, the relationships between air pollutants and pulmonary function can be significantly maintained, even after controlling for these meteorological variables. In addition, we did not measure precisely each individual's exposure level to ambient air pollutants, but we used monitoring data for the nearest site to their home. However, the misclassification of exposure levels according to the monitoring data may have contributed to a null association. Although we adjusted for other

potential factors affecting lung function, such as smoking, in the statistical analyses, concomitant exposure to other pollutants, such as volatile organic chemicals or heavy metals, could also have had some impact on lung function.

In conclusion, our findings suggest that short-term exposure to air pollutants reduces lung function in the elderly for a period lasting from the day of exposure to 2 days after exposure. These results contribute to our understanding of how short-term exposure to air pollution affects lung function.

4. References

- [1] Hole DJ, Watt GC, Davey-Smith G, *et al.* Impaired lung function and mortality risk in men and women: findings from the Renfrew and Paisley prospective population study. *BMJ* 1996; 313: 711-715.
- [2] Mannino DM, Buist AS, Petty TL, *et al.* Lung function and mortality in the United States: data from the First National Health and Nutrition Examination Survey follow up study. *Thorax* 2003; 58: 388-393.
- [3] Sircar K, Hnizdo E, Petsonk E, *et al.* Decline in lung function and mortality: implications for medical monitoring. *Occup Environ Med* 2007; 64: 461-466.
- [4] Downs SH, Schindler C, Liu LJS, *et al.* Reduced Exposure to PM10 and Attenuated Age-Related Decline in Lung Function. *N Engl J Med* 2007; 357: 2338-2347.
- [5] Abbey DE, Burchette RJ, Knutsen SF, *et al.* Long-term particulate and other air pollutants and lung function in nonsmokers. *Am J Respir Crit Care Med* 1998; 158: 289-298.
- [6] Ackermann-Liebrich U, Leuenberger P, Schwartz J, *et al.* Lung function and long term exposure to air pollutants in Switzerland. Study on Air Pollution and Lung Diseases in Adults (SAPALDIA) Team. *Am J Respir Crit Care Med* 1997; 155: 122-129.
- [7] Segala C, Fauroux B, Just J, Pascual L, Grimfeld A, Neukirch F. Short-term effect of winter air pollution on respiratory health of asthmatic children in Paris. *Eur Respir J.* 1998 Mar;11(3):677-685.
- [8] Cakmak S, Dales RE, Vidal CB. Air pollution and mortality in Chile: susceptibility among the elderly. *Environ Health Perspect* 2007; 115: 524-527.
- [9] Fischer P, Hoek G, Brunekreef B, *et al.* Air pollution and mortality in The Netherlands: are the elderly more at risk? *Eur Respir J Suppl.* 2003; 40: 34s-38s.
- [10] Medina-Ramon M, Schwartz J. Who is more vulnerable to die from ozone air pollution? *Epidemiology* 2008; 19: 672-679.
- [11] Haynes GB, Christiani DC (1993) Measures of small airways disease as predictors of chronic obstructive pulmonary disease. *Occup Med* 1993; 8: 375-395
- [12] Goldberg MS, Burnett RT, Bailar JC III, *et al.* The association between daily mortality and ambient air particle pollution in Montreal, Quebec. 2. Cause-specific mortality. *Environ Res [A]* 2001; 86: 26-36
- [13] Braga A, Zanobetti A, Schwartz J (2001) Structure between particulate air pollution and respiratory and cardiovascular deaths in 10 US cities. *J Occup Environ Med* 2001; 43: 927-933
- [14] Peters A, Goldstein IF, Beyer U, *et al.* Acute health effects of exposure to high levels of air pollution in eastern Europe. *Am J Epidemiol* 1996; 144: 570-581



Advanced Topics in Environmental Health and Air Pollution Case Studies

Edited by Prof. Anca Moldoveanu

ISBN 978-953-307-525-9

Hard cover, 470 pages

Publisher InTech

Published online 29, August, 2011

Published in print edition August, 2011

The book describes the effects of air pollutants, from the indoor and outdoor spaces, on the human physiology. Air pollutants can influence inflammation biomarkers, can influence the pathogenesis of chronic cough, can influence reactive oxygen species (ROS) and can induce autonomic nervous system interactions that modulate cardiac oxidative stress and cardiac electrophysiological changes, can participate in the onset and exacerbation of upper respiratory and cardio-vascular diseases, can lead to the exacerbation of asthma and allergic diseases. The book also presents how the urban environment can influence and modify the impact of various pollutants on human health.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Yun-Chul Hong and Jin Hee Kim (2011). Short-Term Exposure to Outdoor Pollutants and Loss of Pulmonary Function in the Elderly, Advanced Topics in Environmental Health and Air Pollution Case Studies, Prof. Anca Moldoveanu (Ed.), ISBN: 978-953-307-525-9, InTech, Available from:

<http://www.intechopen.com/books/advanced-topics-in-environmental-health-and-air-pollution-case-studies/short-term-exposure-to-outdoor-pollutants-and-loss-of-pulmonary-function-in-the-elderly>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike-3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/), which permits use, distribution and reproduction for non-commercial purposes, provided the original is properly cited and derivative works building on this content are distributed under the same license.

IntechOpen

IntechOpen