

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Interaction of Urban Vegetation Cover to Sequester Air Pollutants from Ambient Air Environment

Sharda Dhadse, D. G. Gajghate, P.R. Chaudhari,
D. R. Satapathy and S. R. Wate
*National Environmental Engineering research Institute, Nagpur
India*

1. Introduction

Acute air pollution problem is being faced in urban agglomeration due to economic expansion, increase in population, increased industrial activities and exponential growth in automobiles. The air pollution from these sources is imposing threat to urban human health. The morbidity and mortality caused by air pollution result in long term reduction of productivity and ultimately in overall deterioration of economic condition (Dockery & Pope, 1994; Anderson et al., 1992; Schwartz et al., 1996). In India the particulate matter problem is very significant due to the huge number of vehicles plying on the road, number of power plants, combustion processes, dust storms and domestic emissions (Gurjara et al., 2004). In the recent studies, exceeding levels of PM10 are observed (TERI, 1997; Chelani et al., 2001).

The trees in urban environment are continuously exposed to air pollutants, which play an important role in maintaining ecological balance by actively participating in the nutrients cycle. Many trees are effective for trapping and absorbing air pollutants and acts as sink to several air pollutants (Allan & Krupa, 1986; Bell & Treshow, 2002; Farmer, 1993; Barker & Tingey, 1992; De Kok & Whalley 1984; De Kok & Stulen, 1998; Treshaw & Anderson, 1989; Nowak et al. 1997; Shyam et al., 2006). Hence it is more beneficial to see the impact of pollution on vegetation especially on roadside trees. (Gajghate & Hasan, 1999; Kotoh et al. 1989; Kozhauharov et.al. 1985; Ninave et.al., 2001; Mellios et al., 2006; Mutena, 2004; Tommervik et al., 1995).

Nagpur City is the best place to study the interactions of atmospheric pollutants such as SO₂, NO₂ and suspended particulate matter (SPM) on vegetation, as it is high traffic zone with industrial area on the outskirts as well as have good vegetation cover in the city. Nagpur city is very well known as second Green City in India. Plantations are actively carried out every year in the city with the active participation of local administration and non-governmental organizations (NGOs). The city is also richly dotted by well maintained parks, plantations, forest patches and agricultural fields. This has given lush green aesthetic appearance to the Nagpur city.

Present research is carried out, to study the status of urban pollution in relation to biodiversity in the Nagpur city using ambient air quality monitoring, remote sensing for landuse cover, ground truth and anatomical and biochemical responses of the trees to air pollution.

2. Materials and methods

2.1 Sampling locations

The Nagpur city is situated in between 20° 30' and 21° 30' N latitude and 78° 30' and 79° 30' E longitude. The strategic situation of the Nagpur City in the central part of India has lead a rapid expansion of city and ever increasing environmental problems with reference to pollution of air, water and soil. Ambient air quality monitoring, vegetation survey and remote sensing study have been carried out in three different areas of the city having different activities namely industrial (MIDC), commercial (Itwari) and residential (NEERI Campus).

2.2 Remote sensing study

In order to strengthen the baseline information on existing land use pattern, the remote sensing data has been collected for Nagpur area lying between (21° 03' – 21° 13') N longitude (79° 00' – 79° 10') E latitude. The satellite data was acquired from the IRS P6 (RESOURCESAT -1) LISS III Scene (Path 99, Row 57 dated 04 Nov-2004; CD format) and the collateral data were used from Reference map, Toposheets 55 o/4 and 55 k/16.

2.3 Biological survey

Biological survey was carried out at these three sampling sites. The species of plants, their abundance and diversity were recorded and correlated with land use pattern of vegetation obtained through remote sensing imagery.

2.4 Air pollution tolerance index (APTI)

The plants namely *Bougainvella spectabilis*, *Azadirachta indica*, *Pongamia pinnata* and *Polyalthia longifolia* growing in these area were selected for studying their biochemical responses to the impact of air pollution. The leaf samples were analyzed for pH (P) of leaf extract (Singh and Rao, 1993), Chlorophyll (T) (Arnon, 1949), ascorbic acid (A) (Singh, 1977) and Relative water content (R). Air pollution Tolerance Index (APTI) which gives an empirical value representing tolerance level of a plant to air pollution was used to interpret the impact of pollution on the plants. The APTI is calculated by the formula as $A(T+P)+R/10$.

2.5 Air monitoring

SPM in ambient air was measured by using standard High volume (Hi-vol) sampling technique. Gaseous samples were collected in absorbing solutions by tapping air and collected samples were analyzed using standard wet chemical method (Katz, 1977).

3. Results and discussions

3.1 Vegetation cover in study area through remote sensing

The land use / land cover status in Nagpur urban area as per IRS P6 LISS-III is shown in figure. 1. Eight different classes are identified in the Nagpur namely settlement, water bodies, fallow land, scrubland, bare soil/sand, vegetation-1, vegetation-2 and vegetation-3. Normalized Differentiation Vegetation Index (NDVI) values were computed for Nagpur City. The entire area has been classified into five categories namely non-vegetation, vegetation-1, vegetation-2, vegetation-3, and vegetation-4. The percentage compositions of

these classes in Nagpur City were non-vegetation (58.79%), vegetation-1 (17.40%), vegetation-2 (14.86%), vegetation-3 (3.77%), and vegetation-4 (5.19%) respectively. The percentage compositions of different classes of vegetation (Table 1, Figure 1) in residential area are non-vegetation (71.23%), vegetation-1 (8.87%), vegetation-2 (8.82%), vegetation-3 (4.89%) and vegetation-4 (6.19%). The percentage compositions of different classes in industrial area are non-vegetation (74.08%), vegetation-1 (10.18%), vegetation-2 (7.31%), vegetation-3 (4.51%), and vegetation-4 (3.92%). Similarly, the percentage compositions of different classes in commercial area are non-vegetation (89.38%), vegetation-1 (6.18%), vegetation-2 (3.34%), vegetation-3 (0.49%) and vegetation-4 (0.61%). Table 1 show that the vegetation cover is highest at residential area i.e. 28.77%, lowest in commercial area i.e. 10.62% and moderate in industrial area i.e. 25.92%. Similarly the density, abundance and diversity of plants are highest at residential area while it is lowest in commercial area. The density and diversity of plants is medium at industrial area.

Classification Category	Residential Area		Industrial Area		Commercial Area	
	(Area in Km ²)	Area in %	(Area in Km ²)	Area in %	(Area in Km ²)	Area in %
Non-vegetation	19.547	71.23	20.330	74.08	24.529	89.38
Vegetation – 1	2.434	8.87	2.793	10.18	1.695	6.18
Vegetation – 2	2.420	8.82	2.007	7.31	0.917	3.34
Vegetation – 3	1.342	4.89	1.238	4.51	0.133	0.49
Vegetation – 4	1.700	6.19	1.075	3.92	0.618	0.61
Total (Vegetation)	7.896	28.77	7.113	25.92	3.363	10.62
Total Area	27.443	100	27.443	100	27.443	100

Table 1. Inventory of vegetation cover at different air quality monitoring locations

3.2 Floral diversity and abundance

The field survey of plants in Nagpur City showed a total of 103 plant species consisting of 45 tree species, 13 shrub species, 30 herb species, 8 species of climbers, 1 species of bamboo and 5 species of grasses. The density and diversity of plants were not equally distributed throughout the Nagpur City. The numbers of plants are more in well-planned residential area with adequate space for maintenance of green cover. Density and diversity of the plants is more in residential areas. Around 100 plants species were observed in residential areas with higher abundance of most of the species (around 800 trees/ha). The dominant plants were *Azadirachta indica*, *Albizzia procera*, *Annona squamosa*, *Leucaena leucocephala*, *Ipomoea fistulosa*, *Lantana camara*, *Alternanthera sessilis*, *Ageratum conizoides*, *Calotropis gigantia*, *Parthenium hysterophorus*, *Cyanodon dactylon* and *Dactyloctenium*. Among trees *Azadirachta indica*, *Leucaena leucocephala* are most abundant species. The shrub *Calotropis procera* and *Lantana camara* are abundant species. (Table 2).




Fig. 1. Vegetation cover at Nagpur city

The industrial area has moderate amount of vegetation cover, which consist of avenue plantations, greenbelts and vegetation around industrial complexes. A total of 59 species of

plants were recorded from this area. The density of trees is around 200 trees./ha. The dominant plants were *Azadirachta indica*, *Cassia fistula* and *Cassia siamea*. While among herbs *Parthenium hysterophorus*, *Ageratum conizoides* and *Alternanthera sessilis* were the most abundant species.

Sr. No.	Abundance Class	Abundance Scale (Species cover)	Species		
			Tree	Shrub	Herb
1	5	76-100%	<i>Azadirachta indica</i>	<i>Calotropis procera</i>	<i>Parthenium hysterophorus</i>
2	4	51-75%	<i>Leucaena leucocephala</i>	<i>Lantana camara</i>	<i>Ageratum conyzoides</i> , <i>Alternanthera sessile</i>
3	3	26-50%	<i>Albizzia procera</i>	<i>Ricinus communis</i>	<i>Malvastrum tricuspidatum</i>
4	2	6-25%	<i>Cassia siamea</i>	<i>Vitex negundo</i>	<i>Tridax procumbens</i> , <i>Ocimum sanctum</i>
5	1	up to 5%	<i>Psidium guajava</i>	<i>Indigofera tinctoria</i>	<i>Ricinus communis</i> , <i>Tinospora cordifolia</i>

Table 2. Abundance of major plant species in Nagpur urban area

The commercial area is the old part of the city with narrow lanes, closely set houses, shops and hectic commercial activity. Avenue plantation is negligible. Gardens and other plantation are almost absent. Thus, less number of plant species (32 species) were observed, the important species were *Azadirachta indica* and *Cassia siamea*.

3.3 Air quality status

Annual ambient air quality data is presented in Table 3. The annual average values of SPM varied ranged from 124, 134 and 195 g/m³ at residential, industrial and commercial sites respectively. Highest concentrations of SPM were recorded in commercial site. The reason for high SPM at commercial site is due to vehicular emissions, re-suspension of dust, commercial and domestic use of fuel etc. (NEERI Report, 2001).
The annual average values of NO₂ concentration were ranged from 18, 15 and 21 g/m³ at residential, industrial and commercial sites and those of SO₂ in air ranged from 6, 9 and 7 at residential, industrial and commercial sites respectively.
The residential area showed lowest levels of air pollutants as compared to those in industrial and commercial places. Though the values of SO₂ and NO₂ in residential, industrial and commercial areas are well below the standards, SPM values exceed the standards for commercial area i.e. 140 g/m³. Thus the ambient air quality is satisfactory in Nagpur city. However the rapid expansion of city, increasing number of automobiles and

proposed big industrial area indicate that Nagpur would also be classified as polluted city if preventive environmental management measures are not undertaken.

Air Pollutant	Residential Site			Industrial Site			Commercial Site			Co-relation Coefficient
	N	AM	SD	N	AM	SD	N	AM	SD	
SPM	70	124	77	54	134	101	51	195	107	0.00278
SO ₂	92	6	2	83	9	9	82	7	4	0.65
NO ₂	92	18	18	83	15	14	82	21	21	0.709

N: Sampling Days
AM: Arithmetic Mean
SD: Standard Deviation

Table 3. Annual Arithmetic Mean of SPM, SO₂ & NO₂

3.4 Sequestration of air pollutants by vegetation cover

The concentration of air pollutants was observed to be highest in industrial area followed by commercial area and residential area in decreasing order. It appears that ambient air quality in these microhabitats of Nagpur city is governed by sources of pollution as well as vegetation cover in that area. Table 3 shows that the biodiversity is highest in residential area followed by industrial area and commercial area in decreasing order.

Considering the air pollutant concentration in industrial area as 100%, the air pollutants SPM and SO₂ showed decrease in their concentration by 31.28% and 36.39% for SPM and 22.22% and 33.33% for SO₂ in commercial and residential area respectively. Thus filtering capacity of the plants is well known and these are utilized in greenbelt and in avenue plantation for filtration and reduction of dust concentration in air (Rao *et al*, 1993; Olszyk, 1984). The trees are also well known for acting as sink for SO₂ (Butterbach *et al.*, 1997; Shen *et al.*, 1995). The densities of trees (No/hector) were observed to be co-relating with reduction in ambient SPM and SO₂ concentration (Fig. 2 and 3).

Similarly the biodiversity of plants is also observed to be positively co-related with reduction in ambient concentrations of SO₂ and SPM. It is thus evident that increase in number of trees and increase in biodiversity helps in effective sequestration of air pollutants especially SO₂ and SPM. The ambient NO₂ concentration was observed to vary from 15 to 21 µg/m³ (Fig 4). Each distribution pattern did not co-relate with number of trees or biodiversity of the area.

It has been reported that plant has high preference for SO₂ and extremely law preference for NO₂ to be absorbed and metabolized in the plant tissue. The preference for the plants for air pollutants has been investigated by Hill (1971) and Bennet and Hill (1973, 1975) that appear to follow the following order.

$$HF > SO_2 > Cl > NO_2 > O_3 > PAN > NO > CO$$

Smith (1981) assess pollutant removal efficiency of a model forest hector developed by USEPA which consisted of 95% year old plants will observed that, this forest patch annually removes 748 tonnes of SO₂ while only 0.38 tonnes of NO₂ is removed from atmosphere.




Fig. 2. Relationship between number of trees (per ha) and ambient SPM concentration (Slope: 0.04 and Intercept: 166.17)




Fig. 3. Relationship between number of trees (per ha) and ambient SO₂ concentration (Slope: 0.00238; Intercept: 8.17)




Fig. 4. Relationship between number of trees (per ha) and ambient NO₂ concentration (Slope: 0.00238; Intercept: 8.17)

3.5 Air pollution tolerance index (APTI)

The forgoing discussion shows that present level of pollutants are lower than the prescribed limits except SPM. However these chronic levels are prevailing for a long time i.e. more than 20 years. It is the vegetation cover that is acting as sinks to the air pollutants and playing important role in keeping the levels of pollution well below the standards. However the vegetation cover is relatively less in the commercial area and shows impact of air pollution. The vegetation of Nagpur city is exposed to dust pollution, which may affect the biochemical make up and tolerance capability of plants to the air pollution. The Air Pollution Tolerance Index for the plants from residential, commercial and industrial area is shown in figure III respectively. The APTI of the plants from residential area ranged from 13.09 to 61.39 and from 16.69 to 31.44 in commercial area whereas from 22.02 to 83.15 for plants in industrial area. APTI values of the four common trees in polluted area were found to vary from 16.69 to 83.15 and the plants are listed according to their tolerance or sensitivity to pollution. The four species selected for APTI studies are tolerant to air pollutants in the geographical area of Nagpur city. Next to *Azadiracta sp.* with APTI value (61.36) *Baugainvella sp.* was found to have APTI index values of 16.87 at residential area, 27.44 at commercial area and 27.83 in industrial area. *Polyalthia sp.* which ranks next showed APTI values of 18.40 at residential area, 21.64 and 22.12 at commercial and industrial area respectively (Fig. 5). Though these species showed different sensitivity, they all appear to be tolerant to air pollution as their APTI values showed to be increased in polluted environment. This may probably be the reason that the plants develop tolerance and detoxify the absorbed pollutants and as a result other biochemical constituents are not affected in such plants. This

can be taken as an indication of the development of detoxification mechanism in the plant necessary for the tolerance.

This work has indicated the suitability of *Azadirachta indica* as the most tolerant species, suitable as sink for air pollution. It can be utilized for urban plantation and greenbelt development in industrial area to reduce the level of air pollution. However more research is necessary on a wide variety of trees, shrubs and herbs to prepare a biological sensitivity map of flora. The vast database would be useful in identifying tolerant plants, sink plants and management program.




Fig. 5. APTI values of four tree species in different sampling sites

4. Conclusion

The consequences of complex effects of combinations of several atmospheric pollution and climate change in particular may threaten vegetation in ecosystem of urban atmosphere. The levels of pollutants SPM, NO₂ and SO₂ were highest at commercial site followed by industrial and then residential site. Status of air quality with respect to NO₂ and SO₂ showed the values within the range while SPM level is higher than the standard value except at commercial place. In response to these changing conditions plants adopt to their changed environment by showing different air pollution tolerance index. Out of four species namely *Azadirachta*, *Polyalthia*, *Baugainvillia* and *Pongamia*, only the *Azadirachta* is having the best air pollution tolerance index. This study can be helpful to identify the plant species for greenbelt development as one of the control measures for reduction of ambient concentration of air pollution. This tolerant species may be used for avenue plantation and beautification of the city. The vegetation is denser at residential site, while it is moderate at industrial site. Though the industrial area shows moderate air pollution but the small green

belts in and around the industries do not allow the air pollutants to raise upto higher limits. Similarly at commercial location the lack of vegetation causes increase in the levels of NO₂, SO₂ and SPM. This proves that the plants are the best sinks for all types of air pollutants. More work in this area is needed for more precise information on expected changes in vegetation exposed to air pollutants. Similar studies can be extended to identify air pollution tolerant species at different environmental pollution level in different climatic conditions, which can be used to maintain the air quality level within the permissible level.

5. Acknowledgment

Authors are grateful to the Director, NEERI Nagpur and Council of Scientific and Industrial Research (CSIR) for providing the infrastructure and financial support.

6. References


- Allan, H., L & Krupa, S. V. (1986). Air pollutants and their effects on the terrestrial Ecosystem. *Series in advances in environmental Sci & Tech*.
- Anderson, K. R., Avol, E. L., Edward, S.A., Shamoo, D.A., Peng, R.C., Linn, W.S. & Hackny, J.S. (1992). Controlled exposures of volunteers to respirable carbon and sulphuric acid aerosols. *J Air Waste Manag Assoc*, 42, 770-776.
- Arnon, D.F. (1983). Copper enzymes in isolated chloroplasts polyphenol oxidase in *Beta vulgaris*. in: *Proc Symp on Air Pollution Control held at IIT, Delhi*, 17-18.
- Barker, J. R. & Tingey D. T. (1992). *Air pollution effects on biodiversity*. pub. Van Nostrand Reinhold, New York.
- Bell J.N.B. & Treshow, M. (2002). *Air pollution and plant life*. pub. John Wiley & Sons Ltd.
- Benett, J.H. and Hill, A.C. (1973). Absorption of gaseous air pollutants by a standardized plan canopy. *J. Air. Pollut. Contr. Ass.*, 28, 203-206.
- Bennett, J.H. and Hill, A.C. (1975). Interaction of air pollutants with canopies of vegetation. J.S. Mudd and T.T. Koslowski (eds). *Response of plants to Air Pollution*, Academic Press, New York, 273-306.
- Butterbach -Bahl K. R., Gasche, L. Breuer & H. Papen (1997). Fluxes of NO and N₂O from temperate forest soils: impact of forest type, N deposition and of liming on the NO and N₂O emissions. *Impacts of air pollutants on vegetation in developing countries*, 48, 1-2, 79-90.
- Chelani, A.B., Gajghate, D.G. & Hasan, M.Z.. (2001). Airborne toxic metals in air of Mumbai city, India. *Bull Environ Contam Toxicol*, 66, 196-205.
- De KoK Luit, J. & Stulen, B. I. (1998). *Response of plant metabolism, Air Pollution and Global change*. Publishers Leiden.
- De KoK, M.L. & Whalley, F.R. (1984). *Gaseous air pollutants and plants metabolism*, Butterworth London: 480.
- Dockery, D.W. & Pope, C.A. (1994). Acute respiratory effects of particulate air pollution. *Ann Rev Public health*, 15, 107-132.
- Farmer, A.M. (1993). The Effect of dust on vegetation – A review. *Environment Pollution*, 79: 63-75.
- Gajghate, D.G. & Hasan, M.Z. (1999). Ambient lead level in urban Area. *Environmental contamination and Toxicology*, 62, 403-408.

- Gurjara, B.R., Aardennea, J.A., Lelieveld, J. & Mohanb, M. (2004). Emission estimates and trends (1990-2000) for mega city Delhi and implications. *Atmos Environ*, 38, 5663-5681.
- Hill, A.C. (1971). Vegetation: A sink for atmospheric pollutants. *J. Air Pollution Control Assoc.*, 21: 341-346.
- Katz, M. (1977). *Standard methods for air sampling and analysis*. 2nd edition, APHA Press Inc., Spring Field VA.
- Kotoh, T., Kasuya, M., Kagamimork, S., Kazuka, S. & Kawano, S. (1989). Effects For air pollution on Tannin biosynthesis and predation damage in *Cryptomeria Japonica*. *Phytochem*, 28, 439-445.
- Kozhauharov, S.I., Petrova, A.V. & Veroukova, L. (1985). Some aspects of using plants as indicators of environmental pollution. *In symposium on biomonitoring state of environment, Indian National Science Academy, New Delhi-110002*, 98-105.
- Mellios, G., Aalst, R. V & Samaras Z. (2006). Validation of road urban emission inventories by means of concentration data measured at air quality monitoring station in Europe. *Atmospheric Environment*, 40, 7362-7377.
- Mutena, F. (2004). Measuring air quality over large urban area: development and application of an air pollution index at the urban area of Naples. *Atmospheric Environment*, 38, 6195-6202.
- NEERI Report, (2001). Ambient air quality status for ten cities of India. No.11. submitted to Central Pollution Control Board, Government of India, New Delhi.
- Ninave, S.Y., Chaudhari, P.R., Gajghate, D.G. & Tarar, J.L. (2001). Foliar Biochemical features of plants as indicators of air Pollution. *Bull. Environ. Contam. Toxicolol.*, 67, 133-140.
- Nowak, D.L., Michale, P.J., Ibarra, M., Crane, D., Stevens, J. C. and Luley C.J. (1997). The effect of urban vegetation on air pollution in 22nd Nato/CCMS International Technical Meeting on air Pollution modeling and its application, 276-282 Nato/Csms, Brussels.
- Olszyk, D.M. and Tingey, D.T. (1984). Phytotoxicity of air pollutants evidence for Phyto Detoxification of SO₂ but not O₃. *Plant Physiol*, 74, 999-1005.
- Rao, M. V., Khujneri, S, Dubey, P. S. & Kumawa, D. M. (1993). Response of eight tropical plants to enhanced ammonia deposition under field conditions prevalent with SO₂ and NH₃. *J. Water, Air & Soil Pollution*, 71, 331-345.
- Schwartz, J., Dockery, D.W. & Neas, L.M. (1996). Is daily mortality associated specifically with fine particles?. *J Air Waste Manag Assoc.* 46,927-939.
- Shen, J., Zhao, Q., Tang, H., Zhang, F., Feng, Z., Okita, T., Ogura, N. & Totsuka, T. (1995). Concentrations and depositions of SO₂, SO₄²⁻ etc. in a Changing suburban forested area. *Water, Air & Soil Pollution*, 85, 3, 1299-1304.
- Shyamshilpa, H. N., Verma, S. & Bhangana. (2006). Air pollution & its impact on plant Growth. New India Publishing Agency.
- Singh, A. (1977). Practical Plant Physiology. Kalyani Publishers, New Delhi, p. 266.
- Singh, S.K. & Rao, D.N. (1993). Evaluation of plants for their tolerance to air pollution . *In: Proc. Symp on Air Pollution control held at IIT Delhi, Nov, 1983*, 218-224.
- Smith, W.H. (1981). Air pollution and forests interactions between Air Contaminants and Forest Ecosystems Springer – Verlag, New York.

- TERI. (1997). Environmental aspects of energy use in large Indian metropolitan cities. Tata Energy Research Institute (TERI), New Delhi.
- Tommervik, H., Johansen, B.E. & Pedersen, J.P. (1995). Monitoring the effects of air pollution on terrestrial ecosystem in Varanger (Norway) and Nikel-Pechenga (Russia) using remote sensing. *The Science of Total Environment*. 160,161, 753-767.
- Treshaw, M. & Anderson, F. K. (1989). *Plant stress from air Pollution*. John Wiley and Sons.

IntechOpen

IntechOpen



Air Pollution - New Developments

Edited by Prof. Anca Moldoveanu

ISBN 978-953-307-527-3

Hard cover, 324 pages

Publisher InTech

Published online 06, September, 2011

Published in print edition September, 2011

Today, an important issue is environmental pollution, especially air pollution. Due to pollutants present in air, human health as well as animal health and vegetation may suffer. The book can be divided in two parts. The first half presents how the environmental modifications induced by air pollution can have an impact on human health by inducing modifications in different organs and systems and leading to human pathology. This part also presents how environmental modifications induced by air pollution can influence human health during pregnancy. The second half of the book presents the influence of environmental pollution on animal health and vegetation and how this impact can be assessed (the use of the micronucleus tests on *TRADESCANTIA* to evaluate the genotoxic effects of air pollution, the use of transplanted lichen *PSEUDEVERNIA FURFURACEA* for biomonitoring the presence of heavy metals, the monitoring of epiphytic lichen biodiversity to detect environmental quality and air pollution, etc). The book is recommended to professionals interested in health and environmental issues.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Sharda Dhadse, D. G. Gajghate, P.R. Chaudhari, D. R. Satapathy and S. R. Wate (2011). Interaction of Urban Vegetation Cover to Sequester Air Pollutants from Ambient Air Environment, *Air Pollution - New Developments*, Prof. Anca Moldoveanu (Ed.), ISBN: 978-953-307-527-3, InTech, Available from:
<http://www.intechopen.com/books/air-pollution-new-developments/interaction-of-urban-vegetation-cover-to-sequester-air-pollutants-from-ambient-air-environment>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike-3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/), which permits use, distribution and reproduction for non-commercial purposes, provided the original is properly cited and derivative works building on this content are distributed under the same license.

IntechOpen

IntechOpen