

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

185,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



# Epileptic Channelopathies and Dysfunctional Excitability - From Gene Mutations to Novel Treatments

Sigrid Marie Blom<sup>1</sup> and Henrik Sindal Jensen<sup>2</sup>

<sup>1</sup>*Department of Physiology, University of Bern*

<sup>2</sup>*Neuroscience Drug Discovery, H. Lundbeck A/S*

<sup>1</sup>*Switzerland*

<sup>2</sup>*Denmark*

## 1. Introduction

Epilepsy is not a single disorder, but a collection of disorders that all are characterized by episodic abnormal synchronous electrical activity in the brain. This abnormal activity represents a disturbance of the balance between excitatory and inhibitory neurotransmission. The majority (50%) of epilepsies are cryptogenic, meaning there is a presumptive but no identifiable underlying etiology. Approximately 20% of epilepsies have an identifiable cause (i.e. they are symptomatic) and are usually a result of trauma to the head, stroke, brain tumours, or infections. The remaining 30% are idiopathic, meaning there is no apparent underlying cause (Berg et al., 1999). However, as they are usually associated with a family history of similar seizures, they are mostly considered to be genetic. Mutations in over 70 genes have been found to cause epilepsy (Noebels, 2003). Given the dependence of seizures on synaptic transmission and neuronal excitability, it is not surprising that many of these mutations affect the function of ion channels. Since the identification of the first epilepsy-causing ion channel mutation, scientists have come a long way in the understanding of the pathogenesis of the disease. This chapter deals with some of the main questions that have been asked, and looks at some of the proposed answers to the questions. How do mutations in certain ion channels lead to hyperexcitability and seizures? Why do mutations in one ion channel cause a particular epilepsy syndrome? Why are the seizures often initiated during specific physiological events? And why do most of the childhood epilepsies remit with age? Furthermore, ion channels as targets for antiepileptic drugs will be discussed.

## 2. Idiopathic epilepsies

In most cases genetic epilepsy syndromes have a complex rather than a simple inheritance pattern. Although the epilepsies described here are thought to be monogenic, not even those considered inherited in a dominant fashion have a penetrance of 100%. Mutations within the same gene can result in clinically distinct phenotypes. Variable expressivity is also a common feature of inherited epilepsy demonstrated by family members with the same mutation that exhibit differences in the clinical severity of the disease (Hayman et al., 1997).

On the other hand, some of the disorders display locus heterogeneity where mutations in distinct genes result in the same syndrome. This indicates that other factors beside the primary mutation influence the clinical manifestation of the epilepsy, e.g. environmental factors, developmental events, or differences in inheritance of genetic susceptibility alleles. The latter is supported by mouse models where differences between the genetic backgrounds of two mouse strains influence the severity of a disease caused by the same sodium channel mutation (Bergren et al., 2005).

Unfortunately, discovery of the responsible gene for an epilepsy syndrome have not led to a prompt understanding of the pathogenesis of the disease. Many of the mutated channels have been characterized in expression systems, but only in some cases have this led to a better understanding of the disease. In other cases this have led to more confusion, as some mutations in a particular channel are found to enhance channel function while others appear to cause a loss of function, even though the clinical manifestation are similar. There are also large discrepancies between results depending on the expression system used to characterize the channels. The mutated channel can e.g. show enhanced function when expressed in *Xenopus laevis* oocytes, while the opposite is shown when expressed in mammalian cells (Meadows et al., 2002). To make it even more difficult, it has been demonstrated that depending on the type of neuron in which a mutated channel is expressed, it can have strikingly different effects on the excitability of the cell (Waxman, 2007). While a mutation can make one type of neuron hyperexcitable, the same mutation can make another neuron hypoexcitable. So changes in neuronal function are not necessarily predictable solely from the change in the behaviour of the mutated channel itself, but have to be considered in the cell background in which the mutated channel is expressed. Further, depending on whether a mutated channel mainly is expressed in excitatory or inhibitory neurons, it can have completely opposite effects on the excitability status of the neuronal network (Yu et al., 2006).

The ion channel mutations are bound to cause relative subtle changes in neuronal function. Mutations that cause dramatic changes would likely result in a more severe phenotypes or lethality. The mutations apparently allow normal behaviour under most circumstances, but disturb the equilibrium between excitatory and inhibitory neuronal networks, so that small external perturbations such as fever are sufficient to break the homeostasis and induce seizures.

### 3. Mutations in sodium channel subunit genes

#### 3.1 Voltage-gated sodium channels

Voltage-gated sodium channels play an essential role in the initiation and propagation of action potentials. These channels open as the membrane depolarizes and inactivate within a few milliseconds of opening. As the membrane polarizes again, the inactivation is removed and a second depolarizing stimulus is able to reopen the channel.

Sodium channels are large, multimeric complexes composed of an  $\alpha$  subunit and one or more auxiliary  $\beta$  subunits. The  $\alpha$  subunit has four homologous domains, each consisting of six transmembrane helices. The  $\beta$  subunit has one transmembrane segment and an extracellular domain with an immunoglobulin-like fold and belongs to the Ig superfamily of cell adhesion molecules (CAMs) (Catterall, 2000). The association with  $\beta$  subunits modulate cell surface expression and localization, voltage-dependence and kinetics of activation and inactivation, as well as cell adhesion and association with signalling and cytoskeletal

molecules (Patino and Isom, 2010). Nine  $\alpha$  subunits (Nav1.1 – Nav1.9 encoded by SCN1A-SCN11A) and four  $\beta$  subunits (encoded by SCN1B-SCN4B) have been characterized so far. In addition, the enigmatic NaX channel, which appears not to be gated by voltage but rather by sodium, is encoded by the SCN7A gene (previously assigned as SCN6A) (Hiyama et al., 2002). Nav1.1, Nav1.2, Nav1.3 and Nav1.6 are the sodium channel  $\alpha$  subunits most abundantly expressed in the brain (Yu and Catterall, 2003).

### 3.2 GEFS+ and SMEI

Febrile seizures, i. e. seizures induced by elevated body temperature, affect approximately 3% of children under 6 years of age and are by far the most common seizure disorder. Generalized Epilepsy with Febrile Seizures Plus (GEFS+) is an autosomal dominant epileptic syndrome where the febrile seizures may persist beyond 6 years of age and which may be associated with afebrile generalized seizures (Scheffer and Berkovic, 1997). The disease has a penetrance of approximately 60%. In 1998, GEFS+ was linked to mutation in SCN1B, the voltage-gated sodium channel  $\beta$ 1 subunit gene (Wallace et al., 1998). GEFS+ can also result from mutations in the sodium channel  $\alpha$  subunit genes SCN1A (Escayg et al., 2000) and SCN2A (Sugawara et al., 2001), and from mutations in the GABRG2 gene which encodes the  $\gamma$ 2 subunit of the GABA<sub>A</sub> receptor (Baulac et al., 2001). Heterozygous mutations in SCN1A can also result in Severe Myoclonic Epilepsy of Infancy (SMEI), also known as Dravet syndrome (Claes et al., 2001). This rare form of epilepsy is characterized by generalized tonic, clonic, and tonic-clonic seizures that are initially induced by fever, light, sound, or physical activity and typically begin around 6-9 months of age. Later, SMEI patients also manifest other seizure types including absence, myoclonic, and simple and complex partial seizures. Psychomotor development stagnates around the second year of life and the patients often respond poorly to antiepileptic drugs. The disorder usually occurs in isolated patients as a result of *de novo* mutations (Claes et al., 2003; Ohmori et al., 2002).

### 3.3 How mutations in sodium channels can cause seizures

As sodium channels are responsible for the upstroke of the action potential one might expect that epilepsy-causing mutations in sodium channel genes increase the activity of the channel, thereby allowing increased influx of sodium ions and consequently neuronal hyperexcitability. Indeed, biophysical analyses of the mutant channels have shown that several of the mutations are gain-of-function mutations that increase sodium currents, e. g. by impairing inactivation or by causing a hyperpolarizing shift in the voltage-dependence of the channel (Lossin et al., 2002; Spanpanato et al., 2003; Spanpanato et al., 2004). The first identified GEFS+ mutation, a C121W missense mutation that disrupts a conserved disulphide bridge in the extracellular Ig domain of the  $\beta$ 1 subunit, causes subtle changes in modulation of sodium channel function and alter the ability of  $\beta$ 1 to mediate protein-protein interactions that are critical for channel localization (Meadows et al., 2002; Wallace et al., 1998). Electrophysiological and biochemical studies on the mutant C121W  $\beta$ 1 subunit co-expressed with Nav1.2 or Nav1.3 have shown that the C121W mutation causes a reduction in current rundown during high-frequency channel activation and increases the fraction of sodium channels that are available to open at subthreshold membrane potentials (Meadows et al., 2002). The mutation is therefore thought to enhance sodium channel function, thereby increasing neuronal excitability and predisposing to seizures.

On the other hand, many of the characterized sodium channel mutations are found to cause attenuation of sodium current (Barela et al., 2006; Lossin et al., 2003; Sugawara et al., 2001).

While it seems like the mild phenotype of GEFS+ mostly is associated with missense mutations that alter the biophysical properties of the channels, the more severe SMEI phenotype is usually caused by nonsense or frameshift mutations that prevent production of functional channels (Claes et al., 2003; Claes et al., 2001; Nabbout et al., 2003; Ohmori et al., 2002). But how can loss-of-function mutations in a sodium channel cause epilepsy when reduced sodium current should lead to hypoexcitability rather than hyperexcitability? The answer seems to be related to the expression pattern of the channels. Nav1.1 is predominantly found in inhibitory interneurons and is thought to conduct most of the sodium current in these cells, whereas excitatory pyramidal neurons express only negligible levels of Nav1.1 (Ogiwara et al., 2007). Catterall and co-workers showed that haploinsufficiency of Nav1.1 channels in heterozygous knock-out mice led to a phenotype resembling that of SMEI (Oakley et al., 2009; Yu et al., 2006). In these mice, sodium currents in GABAergic interneurons in the hippocampus were substantially reduced, whilst the effect in pyramidal cells was much less severe. Loss of one SCN1A copy led to a reduction in action potential number, frequency and amplitude in the interneurons (Yu et al., 2006). Similarly, studies in several animal models carrying nonsense or missense mutations in SCN1A show impaired interneuron function (Martin et al., 2010; Mashimo et al., 2010; Ogiwara et al., 2007; Tang et al., 2009). These studies indicate that functional loss of one copy of SCN1A reduces the inhibitory function of GABAergic interneurons and enhances the excitability of downstream synaptic targets, thereby predisposing to epileptic seizures.

But if this is true, how does the predicted changed Nav1.1 function in many of the patients lead to hyperexcitability when the consequence should be increased GABA action? One possibility is that enhanced sodium current in the interneurons causes too much inhibition, and that this leads to synchronization of the downstream synaptic targets, as has been suggested in the pathogenesis of autosomal dominant nocturnal frontal lobe epilepsy (ADNFL) (Klaassen et al., 2006) (discussed later). Another possibility is that the functional consequences of the mutations in vivo are different from that predicted after in vitro characterization of the mutant channels, and that all of the mutations actually cause a reduction of sodium current in inhibitory neurons. This is supported by studies on knock-out mice lacking the  $\beta 1$  subunit (Chen et al., 2004). These mice show downregulated Nav1.1 expression, indicating that  $\beta 1$  function might be necessary for normal expression of Nav1.1. As the inhibitory interneurons seem to be most affected by a reduction in Nav1.1, the consequences of the  $\beta 1$  mutations might be reduced sodium current in interneurons rather than, or in addition to, increased Nav1.2 and Nav1.3 function.

As mutations in SCN1A most often are associated with febrile seizures the mutations seem not to be sufficient to cause spontaneous seizure themselves. Why are the seizures triggered by fever? Why are the seizures most prevalent in young children? And what is the reason for the age-specific onset of SMEI? It is known that an increase in body temperature leads to an increase in the rate of respiration, especially in young children (Gadomski et al., 1994). This increased respiration can cause respiratory alkalosis in the immature brain, and alkalosis of brain tissue can lead to enhanced neuronal activity and to epileptiform activity (Lee et al., 1996). Studies on rat pups showed that seizure activity induced by hyperthermia had a well-defined pH threshold and that a rise in brain pH to the threshold level by injection of bicarbonate could provoke seizures (Schuchmann et al., 2006). By suppressing the alkalosis with a moderate elevation of ambient CO<sub>2</sub> to 5%, seizures could be abolished within 20 seconds without affecting body temperature. Bicarbonate-induced pH changes and seizures could also be blocked by elevation of ambient CO<sub>2</sub>. In older rats, hyperthermia



only led to a moderate increase in the respiration rate and did not cause respiratory alkalosis and seizures (Schuchmann et al., 2006). Fever and the accompanying elevated pH and enhanced neuronal activity seem therefore to be the drop that makes the barrel overflow and induce the seizures. As several ion channels are sensitive to changes in pH (Jensen et al., 2005; Prole et al., 2003), it will be interesting to see whether some mutations in sodium channel genes render the channels pH-sensitive, which could make the affected individuals specifically susceptible to febrile seizures.

SMEI patients are normal until their first seizure that typically occurs around 6-9 months of age. This age-specificity may be related to the time-specific expression of sodium channels. Nav1.1 is undetectable during prenatal and early postnatal development, a stage where Nav1.3 is preferentially expressed. Nav1.3 expression declines at the expression of Nav1.1 increases. An animal model of SMEI has shown that loss of inhibition and seizure onset correlates in time with an increase in Nav1.1 levels and decline in Nav1.3 levels (Oakley et al., 2009).

## 4. Mutations in GABA<sub>A</sub> receptor subunit genes

### 4.1 GABA receptors

GABA is the major inhibitory neurotransmitter in the central nervous system. There are three types of GABA receptors: GABA<sub>A</sub>, GABA<sub>B</sub>, and GABA<sub>C</sub>. GABA<sub>A</sub> and GABA<sub>C</sub> receptors are ionotropic while GABA<sub>B</sub> receptors are G-protein coupled and often act by activating potassium channels. Most of the cortical inhibitory effects of GABA are mediated by GABA<sub>A</sub> receptors (Chebib and Johnston, 1999).

The GABA<sub>A</sub> receptors are pentameric chloride channels formed by various combinations of different types of  $\alpha$  ( $\alpha 1$  to  $\alpha 6$ ),  $\beta$  ( $\beta 1$  to  $\beta 3$ ),  $\gamma$  ( $\gamma 1$  to  $\gamma 3$ ),  $\delta$ ,  $\epsilon$ ,  $\pi$ ,  $\theta$ , and  $\rho$  ( $\rho 1$  to  $\rho 3$ ) subunits, that each have four transmembrane segments, M1 to M4 (Benarroch, 2007). The most prevalent subunit combination consists of  $\alpha 1\beta 2\gamma 2$  (McKernan and Whiting, 1996). The subunit composition determines the functional and pharmacological characteristics of the receptors (Meldrum and Rogawski, 2007; Sieghart and Sperk, 2002). Binding of GABA to the receptor triggers opening of the chloride channel, allowing rapid influx of chloride that hyperpolarizes the neuron and thereby decreases the probability of generation of an action potential.

### 4.2 GEFS+ and ADJME

As mentioned, GEFS+ can also result from mutation in the GABRG2 gene encoding the  $\gamma 2$  subunit of the GABA<sub>A</sub> receptor (Baulac et al., 2001). Mutations in the  $\alpha 1$  subunit gene (GABRA1) have been linked to Autosomal Dominant Juvenile Myoclonic Epilepsy (ADJME) (Cossette et al., 2002), an idiopathic epilepsy that is not associated with febrile seizures. This disorder typically manifests itself between the ages of 12 and 18 with myoclonic seizures occurring early in the morning and with additional tonic-clonic and absence seizures in some patients.

### 4.3 How mutant GABA<sub>A</sub> receptor subunits can cause seizures

It has been shown that mutations in the  $\gamma 2$  subunit of the GABA<sub>A</sub> receptor cause retention of the receptor in the endoplasmatic reticulum (ER) (Harkin et al., 2002; Kang and Macdonald, 2004). Similarly, the A322D mutation in the  $\alpha 1$  subunit causes rapid ER associated degradation of the subunit through the ubiquitin-proteasome system (Gallagher et al., 2007).

This reduced cell surface expression would result in decreased inhibitory GABA<sub>A</sub> receptor current, and consequently an increase in neuronal excitability and seizure susceptibility. But why are  $\gamma 2$  mutations associated with febrile seizures? And why are mutations in  $\alpha 1$  not?

Variations in temperature have effects on most cellular events. For example, synaptic vesicle recycling has been shown to be temperature dependent with increased temperature speeding both endo- and exocytosis, and there is evidence that inhibitory synaptic strength can be modulated within 10 min through recruitment of more functional GABA<sub>A</sub> receptors to the postsynaptic plasma membrane (Wan et al., 1997). Studies on cultured hippocampal neurons showed that while trafficking of wild-type  $\alpha 1\beta 2\gamma 2$  receptors is slightly temperature dependent with a small decrease in surface expression after incubation at 40°C for 2h, trafficking of receptors with mutations in the  $\gamma 2$  subunit is highly temperature dependent (Kang et al., 2006). Increases in temperature from 37°C to 40°C impaired trafficking and/or accelerated endocytosis of the mutant receptors within 10 min, suggesting that the febrile seizures may be a result of a temperature-induced reduction in GABA-mediated inhibition. The study also showed that the A322D mutation in the  $\alpha 1$  subunit did not cause a temperature-dependent reduction in surface expression, consistent with a resulting epilepsy syndrome not associated with febrile seizures (Kang et al., 2006).

## 5. Mutations in potassium channel genes

### 5.1 Kv7 channels and the M-current

The Kv7 family of voltage-gated potassium channels consists of five members, Kv7.1 -5 (also termed KCNQ1-5). All five members share the general structure of voltage-gated potassium channels with four subunits that assemble to form functional tetramers. Each subunit consists of six transmembrane helices, S1-S6, and has a pore forming domain, which is formed by a P-loop between the fifth and the sixth helix. The P-loop contains the GYG (glycine-tyrosine-glycine) sequence, which is highly conserved among potassium channels and confers K<sup>+</sup> selectivity. The fourth helix forms the voltage sensor; it contains several arginine residues and is therefore strongly positive. A S4-S5 linker in one subunit couples the voltage sensor to the intracellular activation gate in S6 of the adjacent subunit (Laine et al., 2003). Although heavily debated, it is believed that when the membrane potential depolarize, the voltage sensor is pushed out leading to bending of the S6 so potassium can enter the channel pore (Long et al., 2005).

All Kv7 channels are strongly inhibited upon activation of muscarinic receptors and are hence called M-channels (Schroeder et al., 2000; Selyanko et al., 2000). The current conducted by these channels, the M-current, was first described in bullfrog sympathetic ganglia as a slowly activating, slowly deactivating, sub-threshold voltage-dependent K<sup>+</sup> current that showed no inactivation (Brown and Adams, 1980). The Kv7 channels have slow activation and deactivation kinetics, and in line with other voltage-gated potassium channels they open upon membrane depolarization. However, the threshold for activation is low compared to most other channels, approximately -60 mV. Since the channels open at voltages that are around or below the threshold for generation of an action potential they allow potassium flow that opposes the depolarization required to generate action potentials, and hence make the neuron less excitable. If the M-channels remain open during excitation of the nerve, the spike frequency is dampened, while inhibition of the M-current by activation of muscarinic acetylcholine receptors enables repetitive firing (Hille, 2001).

Kv7 channels are primarily localized at the axon initial segment, the site where synaptic inputs are integrated and action potentials are generated (Pan et al., 2006; Rasmussen et al., 2007). Additionally, immunohistochemical studies have demonstrated a widespread pre-synaptic distribution of some Kv7 channel subunits (Cooper et al., 2000), where they may play a role in depolarization-induced neurotransmitter release (Martire et al., 2004; Martire et al., 2007). Activation of pre-synaptic M-current may hyperpolarize the nerve endings, thus reducing  $\text{Ca}^{2+}$  influx through voltage-gated  $\text{Ca}^{2+}$  channels and limiting the amount of neurotransmitter released.

### 5.2 Benign neonatal familial convulsions

Benign Neonatal Familial Convulsions (BNFC) is a rare autosomal dominant idiopathic form of epilepsy. It is characterized by tonic-clonic seizures that typically begin around three days after birth and remit after 3-4 months. Yet ~16% of patients also experience seizures later in life (Ronen et al., 1993). BNFC is caused by mutations in the genes encoding Kv7.2 (Biervert et al., 1998; Singh et al., 1998) or Kv7.3 (Charlier et al., 1998).

### 5.3 How mutations in Kv7.2 and Kv7.3 can cause BNFC

All characterized mutations in Kv7.2 and Kv7.3 cause a reduction in M-current, either by changing the channel kinetics (Dedek et al., 2001), by altering the trafficking of the channel to the cell membrane (Schwake et al., 2000), or by decreasing the subunit stability (Soldovieri et al., 2006). The mutations usually cause a reduction in M-current of about 25% (Schroeder et al., 1998). Considering the role of Kv7 channels in controlling neuronal excitability, it is not surprising that a reduction in M-current can cause hyperexcitability and predispose to seizures. Several transgenic strategies have been employed to examine how Kv7.2 and Kv7.3 malfunction lead to BNFC. The traditional knock-out approach resulted in mice that died within a few hours after birth due to pulmonary atelectasis (collapse of the lung) (Watanabe et al., 2000). Even though BNFC often results from Kv7.2 haploinsufficiency, heterozygous *KCNQ2*<sup>+/-</sup> mice did not experience neonatal seizures and appeared normal. They did, however, have increased susceptibility to chemically induced seizures (Watanabe et al., 2000). To overcome the postnatal lethality Dirk Isbrandt and colleagues developed mice conditionally expressing dominant-negative Kv7.2 subunits (Kv7.2-G279S) where expression of the transgene could be turned on after birth (Peters et al., 2005). These mice showed spontaneous seizures, exhibited behavioural hyperactivity and had morphological changes in the hippocampus. Mark Leppert and co-workers developed orthologous mouse models carrying disease-causing mutations in the *KCNQ2* (A306T) or *KCNQ3* (G311V) gene (Singh et al., 2008). Mice heterozygous or homozygous for either mutation had reduced seizure threshold, but only mice homozygous for the mutations exhibited spontaneous seizures. The epileptic phenotype was dependent on the specific mutation, the genetic background, sex, and seizure model (Otto et al., 2009). Hence, none of these transgenic strategies have fully recapitulated the human condition but nevertheless provide us with important clues regarding the pathophysiology of BNFC.

It can be questioned why the disease usually only is clinically manifested in neonates when mutations in the channels cause general hyperexcitability. One possible explanation for the age-dependent remission of seizures is related to the expression of Kv7 channels. There is evidence for a developmental upregulation of Kv7 channels (Weber et al., 2006), suggesting that a reduction in M-current of 25% might have a more prominent effect in the fetal brain



and that this reduction is not sufficient to cause seizures later in life when expression levels of Kv7 channels are higher.

Another possible mechanism is related to developmental changes in GABA function. During the first weeks of life, GABA, the main inhibitory neurotransmitter in the adult brain, provides the main excitatory drive to immature hippocampal neurons (Ben-Ari, 2002). Due to delayed expression of a chloride exporter there is a high intracellular concentration of chloride that leads to a negative shift in the reversal potential for chloride ions, so opening of ionotropic GABA receptors leads to an efflux of negative chloride ions and therefore depolarization. When the chloride-extruding system becomes operative, chloride is efficiently transported out of the cell, and GABA begins to exert its conventional inhibitory action (Ben-Ari, 2002). Because of this, the inhibition in neonatal circuits appears mainly to be mediated through presynaptic control of neurotransmitter release. As Kv7 channels are involved in the release of neurotransmitters (Martire et al., 2004; Martire et al., 2007), it was proposed that these channels serve as the main inhibitor in neonates, and that attenuation of M-current due to mutations in Kv7.2 and Kv7.3 causes reduced inhibition that is sufficient to cause epilepsy (Peters et al., 2005). If Kv7 channels are expressed in GABAergic neurons, the reduced M-current would possibly also cause increased GABA release which further would increase excitability at this point of development. It is also important to note that the neonatal brain is particularly prone to seizures (Holmes and Ben-Ari, 1998). As development continues and overall excitability decreases, reduced M-current apparently becomes less problematic and the seizures abate.

Yet, if this is the fact, why do ~16% of the patients experience seizures later in life after GABA has gained its inhibitory function? There is evidence to indicate that suppression of M-current within the first postnatal week can cause developmental defects, and that the resulting morphological changes in the brain, rather than reduced M-current causes the seizures in adulthood (Peters et al., 2005). In other words, it appears to become a symptomatic epilepsy with an idiopathic aetiology.

## 6. Mutations in nAChR subunit genes

### 6.1 Nicotinic acetylcholine receptors

Nicotinic acetylcholine receptors (nAChRs) consist of five subunits that assemble to form functional pentamers. Each subunit consists of a long extracellular N-terminal domain, four transmembrane helices (TM1-TM4), and a short extracellular C-terminal end. The second transmembrane segment (TM2) from each subunit lines the channel pore. The amino acids that compose the TM2 are arranged in such a way that three rings of negatively charged amino acids are oriented toward the central pore of the channel. These provide a selectivity filter that ensures that only cations can pass through the pore. Brief exposure to high concentrations of Ach causes opening of the water-filled pore and permits an influx of  $\text{Na}^+$  and  $\text{Ca}^{2+}$  and an efflux of  $\text{K}^+$  (Waxham, 2003). After a few milliseconds, the receptor closes to a nonconducting state. Prolonged exposure to agonist causes desensitization of the channel, which stabilizes the receptor in an unresponsive, closed state (Dani and Bertrand, 2007).

The neuronal nAChRs can be either homomeric, consisting of five  $\alpha$  subunits, or heteromeric with two  $\alpha$  subunits and three  $\beta$  subunits. So far, 12 nAChR subunits expressed

in the brain have been identified ( $\alpha 2$ - $\alpha 10$  and  $\beta 2$ - $\beta 4$ ). The most widely distributed nAChRs in the human brain are the homomeric  $\alpha 7$  and the heteromeric  $\alpha 4\beta 2$ . The biochemical, pharmacological and biophysical characteristic of the channels are dependent on the subunit composition (Dani and Bertrand, 2007).

## 6.2 Autosomal Dominant Nocturnal Frontal Lobe Epilepsy

Autosomal Dominant Nocturnal Frontal Lobe Epilepsy (ADNFLE) is a focal epilepsy characterized by clusters of brief nocturnal motor seizures with hyperkinetic or tonic manifestations. Seizures typically occur very soon after falling asleep or during the early morning hours, initiated during stage 2 non-rapid-eye-movement (NREM) sleep. Onset usually occurs around age 10 and the seizures often persist through adult life. However, in most patients the seizures tend to peter out in adulthood.

In 1995, a missense mutation (S248F) in the nAChR  $\alpha 4$  subunit gene (CHRNA4) was found to underlie ADNFLE (Steinlein et al., 1995). Additional disease causing mutations in this gene (Hirose et al., 1999; Steinlein et al., 1997), and mutations in the  $\beta 2$  subunit gene (CHRNA2) (Phillips et al., 2001) have later been identified. Because  $\alpha 4$  and  $\beta 2$  subunits combine, it is not strange that mutations in either subunit produce comparable epileptic symptoms. Almost every known mutation is found in the TM2 region of the two subunits. A common finding among the mutations is that the sensitivity to ACh is increased (Bertrand et al., 2002).

## 6.3 How mutations in nAChRs can cause ADNFLE

Stage 2 NREM sleep is characterized by the appearance of sleep spindles and slow waves, transient physiological rhythmic oscillations that are produced by synchronized synaptic potentials in cortical neurons. It appears that the seizure often arise from a sleep spindle that transforms into epileptic discharges (Picard et al., 2007). Thalamocortical circuits are thought to play a role in the generation of these sleep spindles.

Thalamic relay neurons are reciprocally connected to cortical neurons by excitatory synapses. Both thalamic and cortical neurons also excite GABAergic interneurons of the nucleus reticularis, which in turn inhibit thalamic relay neurons thus forming a feedback-loop (Kandel et al., 2000). The thalamic relay neurons have two different modes of signalling activity: a transmission mode during wakefulness and rapid eye movement (REM) sleep and a burst mode during NREM sleep. In the transmission mode, the resting membrane potential of the thalamic relay neurons is near the firing threshold and incoming excitatory synaptic potentials drive the neuron to fire in a pattern that reflects the sensory stimuli. During the burst mode the thalamic relay neurons are hyperpolarized and respond to brief depolarization with a burst of action potentials, which indicates that the thalamus is unable to relay sensory information to the cortex (Kandel et al., 2000). The thalamic relay neurons can be in different modes because they possess special  $\text{Ca}^{2+}$  channels named T-type channels. These channels are inactivated at the resting membrane potential but become available for activation when the cell is hyperpolarized, and can then be transiently opened by depolarization (Contreras, 2006). The interneurons of the nucleus reticularis that form synapses at the relay neurons hyperpolarize the relay neurons upon activation of GABA<sub>A</sub> receptors, thus removing the inactivation of the T-type  $\text{Ca}^{2+}$  channels. Incoming excitatory synaptic potentials can then trigger transient opening of the T-type  $\text{Ca}^{2+}$  channels and the

resulting influx of  $\text{Ca}^{2+}$  brings the neuron's membrane potential above threshold. The cell now fires a burst of action potentials that produces the synchronized postsynaptic potentials in cortical neurons that cause the spindle waves seen on the electroencephalogram. When sufficient  $\text{Ca}^{2+}$  has entered the cell a  $\text{Ca}^{2+}$ -activated  $\text{K}^{+}$  current is triggered that hyperpolarizes the relay neurons and terminate the spindle (Contreras, 2006). Because of the feedback loops from the relay neurons and cortical neurons that innervate the interneurons of the nucleus reticularis, these are again activated, which allows for another round of burst firing.

There is a high expression of the  $\alpha 4\beta 2$  nAChR subtype in the thalamus and the subtype can be found diffusely distributed onto pyramidal cells and GABAergic interneurons in the cortex. The thalamic relay neurons and the nucleus reticularis receive input from two groups of cholinergic neurons in the upper brain stem: the pedunculo pontine and laterodorsal tegmental nuclei (LDT). These are critical for keeping the thalamic relay neurons in transmission mode during wakefulness and REM sleep. The LDT releases Ach during NREM sleep at the time of an arousal that interrupts the sleep spindle oscillations through depolarization of thalamic relay neurons (Lee and McCormick, 1997). The cortex receives cholinergic input from neurons in the nucleus basalis of Meynert that enhance cortical response to incoming sensory stimuli (Kandel et al., 2000).

After studying transgenic ADNFLE mice with heterozygous expression of an ADNFLE mutation ( $\text{Chrna4}^{\text{S252F}}$  or  $\text{Chrna4}^{\text{+264}}$ ), Boulter and colleagues suggested that an Ach-dependent sudden increase in the response of GABAergic cortical interneurons contributes to the epileptogenesis in ADNFLE patients (Klaassen et al., 2006). They showed that asynchronously firing cortical pyramidal cells got synchronized when relieved from a large GABAergic inhibition triggered by cholinergic over-activation of mutant nAChRs in the cortical interneurons.

Dani and Bertrand suggested that this, together with an increased positive response of cortical pyramidal cells through cholinergic stimulation of their hypersensitive  $\alpha 4\beta 2$  nAChRs, contributes to induction of the seizures (Dani and Bertrand, 2007). As the cortical pyramidal cells project back to the thalamus, excitatory stimuli will be boosted on to the interneurons of the nucleus reticularis and the relay neurons and induce synchronous activity and seizures (Dani and Bertrand, 2007).

In addition, PET studies using a high affinity  $\alpha 4\beta 2$  agonist have showed a clear difference in the pattern of the nAChR density in the brains of ADNFLE patients compared to control subjects (Picard et al., 2006). The studies showed that patients had increased  $\alpha 4\beta 2$  density in the epithalamus, the interpeduncular nucleus (IPN) of the ventral mesencephalon and in the cerebellum, while the density in the right dorsolateral prefrontal region was decreased. As the IPN projects to the LDT, the authors proposed that prolonged depolarizations in IPN neurons, because of both increased density of  $\alpha 4\beta 2$  and hypersensitive receptors, could result in over-activation of the LDT and consequently the thalamic relay neurons (Picard et al., 2006). At the time of arousals, ACh acting on sensitized thalamic relay neurons could prevent the normal arousal-induced interruption of the sleep spindle oscillations and transform them into pathological Thalamocortical oscillations, triggering epileptic seizures (Picard et al., 2006).

Neither of these mechanisms excludes the others. In fact, they might work together yielding even stronger possibilities for induction of seizures.

## 7. Ion channel modulators as antiepileptic drugs

Many of the antiepileptic drugs on the market today exert strong effects on ionic currents. Sodium channel blockers were used in the treatment of epilepsy as early as in 1940, and since then several other sodium channel blockers have been developed. Carbamazepine, first introduced in 1968, stabilizes the inactive conformation of sodium channels and is widely used in the treatment of partial and generalized tonic-clonic seizures. Lamotrigine blocks sodium channels in a voltage- and use-dependent manner and is efficient for partial, absence, myoclonic and tonic-clonic seizures (Kwan et al., 2001). Drugs that potentiate GABA receptor function, such as benzodiazepines, have also been used as anticonvulsants for several years. These act by binding to the interface between the  $\alpha$  and  $\gamma$  subunits of the GABA<sub>A</sub> receptor, resulting in allosteric activation of the receptor (Kwan et al., 2001). Several established and novel antiepileptic drugs have been reported to act on various K<sup>+</sup> currents, but none of them exert their main effect on potassium channels (Meldrum and Rogawski, 2007). Carbamazepine also exerts an effect on nAChRs by blocking the open conformation of the channel and is very effective in treatment of patients with ADNFLE. This effect might be related to the fact that several of the disease causing mutations in CHRNA4 and CHRNA2 increase the sensitivity to the drug (Ortells and Barrantes, 2002).

The knowledge of how mutations in ion channels cause neuronal hyperexcitability and epilepsy has led to new therapeutic strategies for prevention of seizures. The mutated proteins serve as mechanistic proof of concept that pharmacological antagonization of the epilepsy causing mechanisms could have desirable effects on neuronal excitability. These mechanisms are not necessarily targeted for treatment of the respective epilepsy syndrome, but as they are showed to cause hyperexcitability in the affected patients, antagonizing drugs may reduce excitability in patients with other types of epilepsy. An excellent example of this is Retigabine (other names are Trobalt, Potiga, or Ezogabine), which was approved by both the European Medicines Agency and the U.S. Food and Drug Administration in 2011 as adjunctive treatment for partial onset seizures. It was originally synthesized as a GABA modulator, but it was later shown that its main molecular target is the neuronal Kv7 channels (Main et al., 2000; Rundfeldt and Netzer, 2000; Wickenden et al., 2000). Retigabine induces a large hyperpolarizing shift in the voltage-dependence of activation of Kv7 channels, accelerates the activation kinetics and slows deactivation kinetics. Enhancement of M-current would clearly be effective in prevention of seizures in BNFC patients, but mutations in the Kv7 channels might render them insensitive to such drugs, which hampers the use of Kv7 channel openers in the treatment of BNFC. The hypothesis of reduced Nav1.1 current in the pathogenesis of GEFS+ and SMEI, together with its primary expression in inhibitory interneurons, indicates that this sodium channel subtype plays a role in reducing neuronal excitability (Ogiwara et al., 2007; Yu et al., 2006). This opens the intriguing possibility that selective Nav1.1 openers can function as anticonvulsants, even though sodium channel openers generally are known as epileptogenic substances.

An understanding of the etiology of the epilepsy syndromes and the causal factors in individual patients are also critical for selection of the right medication. Sodium channel blockers, which normally would exhibit anticonvulsant properties, would clearly not be the right choice to treat patients with SMEI, where a lack of sodium current seems to be the underlying cause. Similarly, as too much inhibition seems to play a role in the pathogenesis



of ADNFLE, GABA receptor agonist might have a negative effect in these patients. In fact, Boulter and co-workers showed that the GABA<sub>A</sub> receptor antagonist picrotoxin, which normally exhibits convulsive properties, was efficient in preventing seizures in the ADNFLE mice (Klaassen et al., 2006).

## 8. Conclusion

As about 30% of all patients with epilepsy respond poorly to antiepileptic drugs, there is clearly a need for development of new therapies. The growing understanding of the epileptic channelopathies and the structural and functional characterization of the mutated channels provide several opportunities for creation of novel and improved drugs.

## 9. References

- Barela, A.J., Waddy, S.P., Lickfett, J.G., Hunter, J., Anido, A., Helmers, S.L., Goldin, A.L., & Escayg, A. (2006). An Epilepsy Mutation in the Sodium Channel SCN1A That Decreases Channel Excitability. *Journal of Neuroscience* Vol. 26, No. 10, pp. 2714-2723, ISSN 0270-6474
- Baulac, S., Huberfeld, G., Gourfinkel-An, I., Mitropoulou, G., Beranger, A., Prud'homme, J.F., Baulac, M., Brice, A., Bruzzone, R., & LeGuern, E. (2001). First genetic evidence of GABA<sub>A</sub> receptor dysfunction in epilepsy: a mutation in the [gamma]2-subunit gene. *Nature Genetics* Vol. 28, No. 1, pp. 46-48, ISSN 1061-4036
- Ben-Ari, Y. (2002). Excitatory actions of gaba during development: the nature of the nurture. *Nature Reviews Neuroscience* Vol. 3, No. 9, pp. 728-739, ISSN 1471-003X
- Benarroch, E.E. (2007). GABA<sub>A</sub> receptor heterogeneity, function, and implications for epilepsy. *Neurology* Vol. 68, No. 8, pp. 612-614, ISSN 0028-3878
- Berg, A.T., Shinnar, S., Levy, S.R., & Testa, F.M. (1999). Newly Diagnosed Epilepsy in Children: Presentation at Diagnosis. *Epilepsia* Vol. 40, No. 4, pp. 445-452, ISSN 0013-9580
- Bergren, S., Chen, S., Galecki, A., & Kearney, J. (2005). Genetic modifiers affecting severity of epilepsy caused by mutation of sodium channelScn2a. *Mammalian Genome* Vol. 16, No. 9, pp. 683-690, ISSN 0938-8990
- Bertrand, D., Picard, F., Le Hellard, S., Weiland, S., Favre, I., Phillips, H., Bertrand, S., Berkovic, S.F., Malafosse, A., & Mulley, J. (2002). How Mutations in the nAChRs Can Cause ADNFLE Epilepsy. *Epilepsia* Vol. 43, No. s5, pp. 112-122, ISSN 0013-9580
- Biervert, C., Schroeder, B.C., Kubisch, C., Berkovic, S.F., Propping, P., Jentsch, T.J., & Steinlein, O.K. (1998). A Potassium Channel Mutation in Neonatal Human Epilepsy. *Science* Vol. 279, No. 5349, pp. 403-406, ISSN 0036-8075
- Brown, D.A., & Adams, P.R. (1980). Muscarinic suppression of a novel voltage-sensitive K<sup>+</sup> current in a vertebrate neurone. *Nature* Vol. 283, No. 5748, pp. 673-676, ISSN 0028-0836
- Catterall, W.A. (2000). From Ionic Currents to Molecular Mechanisms: The Structure and Function of Voltage-Gated Sodium Channels. *Neuron* Vol. 26, No. 1, pp. 13-25, ISSN 0896-6273



- Charlier, C., Singh, N.A., Ryan, S.G., Lewis, T.B., Reus, B.E., Robin, J., & Leppert, M. (1998). A pore mutation in a novel KQT-like potassium channel gene in an idiopathic epilepsy family. *Nature Genetics* Vol. 18, No. 1, pp. 53-55, ISSN 1061-4036
- Chebib, M., & Johnston, G.A.R. (1999). THE 'ABC' OF GABA RECEPTORS: A BRIEF REVIEW. *Clinical and Experimental Pharmacology and Physiology* Vol. 26, No. 11, pp. 937-940, ISSN 0305-1870
- Chen, C., Westenbroek, R.E., Xu, X., Edwards, C.A., Sorenson, D.R., Chen, Y., McEwen, D.P., O'Malley, H.A., Bharucha, V., Meadows, L.S., Knudsen, G.A., Vilaythong, A., Noebels, J.L., Saunders, T.L., Scheuer, T., Shrager, P., Catterall, W.A., & Isom, L.L. (2004). Mice Lacking Sodium Channel  $\beta$ 1 Subunits Display Defects in Neuronal Excitability, Sodium Channel Expression, and Nodal Architecture. *Journal of Neuroscience* Vol. 24, No. 16, pp. 4030-4042, ISSN 0270-6474
- Claes, L., Ceulemans, B., Audenaert, D., Smets, K., Lofgren, A., Del-Favero, J., Ala-Mello, S., Basel-Vanagaite, L., Plecko, B., Raskin, S., Thiry, P., Wolf, N.I., Van Broeckhoven, C., & De Jonghe, P. (2003). De Novo SCN1A mutations are a major cause of severe myoclonic epilepsy of infancy. *Human Mutation* Vol. 21, No. 6, pp. 615-621, ISSN 1059-7794
- Claes, L., Del-Favero, J., Ceulemans, B., Lagae, L., Van Broeckhoven, C., & De Jonghe, P. (2001). De novo mutations in the sodium-channel gene SCN1A cause severe myoclonic epilepsy of infancy. *American Journal of Human Genetics* Vol. 68, No. 6, pp. 1327-1332, ISSN 0002-9297
- Contreras, D. (2006). The Role of T-Channels in the Generation of Thalamocortical Rhythms. *CNS & Neurological Disorders - Drug Targets* Vol. 5, pp. 571-585, ISSN 1871-5273
- Cooper, E.C., Aldape, K.D., Abosch, A., Barbaro, N.M., Berger, M.S., Peacock, W.S., Jan, Y.N., & Jan, L.Y. (2000). Colocalization and coassembly of two human brain M-type potassium channel subunits that are mutated in epilepsy. *Proceedings of the National Academy of Sciences of the USA* Vol. 97, No. 9, pp. 4914-4919, ISSN 0027-8424
- Cossette, P., Liu, L., Brisebois, K., Dong, H., Lortie, A., Vanasse, M., Saint-Hilaire, J.M., Carmant, L., Verner, A., Lu, W.Y., Tian Wang, Y., & Rouleau, G.A. (2002). Mutation of GABRA1 in an autosomal dominant form of juvenile myoclonic epilepsy. *Nature Genetics* Vol. 31, No. 2, pp. 184-189, ISSN 1061-4036
- Dani, J.A., & Bertrand, D. (2007). Nicotinic Acetylcholine Receptors and Nicotinic Cholinergic Mechanisms of the Central Nervous System. *Annual Review of Pharmacology and Toxicology* Vol. 47, No. 1, pp. 699-729, ISSN 0362-1642
- Dedek, K., Kunath, B., Kananura, C., Reuner, U., Jentsch, T.J., & Steinlein, O.K. (2001). Myokymia and neonatal epilepsy caused by a mutation in the voltage sensor of the KCNQ2 K<sup>+</sup> channel. *Proceedings of the National Academy of Sciences of the USA* Vol. 98, No. 21, pp. 12272-12277, ISSN 0027-8424
- Escayg, A., MacDonald, B.T., Meisler, M.H., Baulac, S., Huberfeld, G., An-Gourfinkel, I., Brice, A., LeGuern, E., Moulard, B., Chaigne, D., Buresi, C., & Malafosse, A. (2000). Mutations of SCN1A, encoding a neuronal sodium channel, in two

- families with GEFS+2. *Nature Genetics* Vol. 24, No. 4, pp. 343-345, ISSN 1061-4036
- Gadomski, A.M., Permutt, T., & Stanton, B. (1994). Correcting respiratory rate for the presence of fever. *Journal of Clinical Epidemiology* Vol. 47, No. 9, pp. 1043-1049, ISSN 0895-4356
- Gallagher, M.J., Ding, L., Maheshwari, A., & Macdonald, R.L. (2007). The GABAA receptor {alpha}1 subunit epilepsy mutation A322D inhibits transmembrane helix formation and causes proteasomal degradation. *Proceedings of the National Academy of Sciences of the USA* Vol. 104, No. 32, pp. 12999-13004, ISSN 0027-8424
- Harkin, L.A., David, N.B., Leanne, M.D., Rita, S., Fiona, P., Robyn, H.W., Michaela, C.R., David, A.W., John, C.M., Samuel, F.B., Ingrid, E.S., & Steven, P. (2002). Truncation of the GABAA-Receptor  $\gamma 2$  Subunit in a Family with Generalized Epilepsy with Febrile Seizures Plus. *American Journal of Human Genetics* Vol. 70, pp. 530-536, ISSN 0002-9297
- Hayman, M., Scheffer, I.E., Chinvarun, Y., Berlangieri, S.U., & Berkovic, S.F. (1997). Autosomal dominant nocturnal frontal lobe epilepsy: Demonstration of focal frontal onset and intrafamilial variation. *Neurology* Vol. 49, No. 4, pp. 969-975, ISSN 0028-3878
- Hille, B. (2001). *Ion channels of excitable membranes*, (3. edn.) Sinauer Associates Inc., ISBN 0-87893-321-2, Sunderland, USA.
- Hirose, S., Iwata, H., Akiyoshi, H., Kobayashi, K., Ito, M., Wada, K., Kaneko, S., & Mitsudome, A. (1999). A novel mutation of CHRNA4 responsible for autosomal dominant nocturnal frontal lobe epilepsy. *Neurology* Vol. 53, No. 8, pp. 1749, ISSN 0028-3878
- Hiyama, T.Y., Watanabe, E., Ono, K., Inenaga, K., Tamkun, M.M., Yoshida, S., & Noda, M. (2002). Na<sub>v</sub> channel involved in CNS sodium-level sensing. *Nature Neuroscience* Vol. 5, No. 6, pp. 511-512, ISSN 1097-6256
- Holmes, G.L., & Ben-Ari, Y. (1998). Seizures in the Developing Brain: Perhaps Not So Benign after All. *Neuron* Vol. 21, No. 6, pp. 1231-1234, ISSN 0896-6273
- Jensen, H.S., Callo, K., Jespersen, T., Jensen, B.S., & Olesen, S.P. (2005). The KCNQ5 potassium channel from mouse: A broadly expressed M-current like potassium channel modulated by zinc, pH, and volume changes. *Molecular Brain Research* Vol. 139, No. 1, pp. 52-62, ISSN 0169-328X
- Kandel, E.R., Schwartz, J.H., & Jessell, T.M. (eds.) (2000). *Principles of Neural Science*, (4. edn.) McGraw-Hill, ISBN 0-8385-7701-6, New York, USA.
- Kang, J., & Macdonald, R.L. (2004). The GABAA Receptor  $\gamma 2$  Subunit R43Q Mutation Linked to Childhood Absence Epilepsy and Febrile Seizures Causes Retention of  $\alpha 1\beta 2\gamma 2S$  Receptors in the Endoplasmic Reticulum. *Journal of Neuroscience* Vol. 24, No. 40, pp. 8672-8677, ISSN 0270-6474
- Kang, J.Q., Shen, W., & Macdonald, R.L. (2006). Why Does Fever Trigger Febrile Seizures? GABAA Receptor  $\gamma 2$  Subunit Mutations Associated with Idiopathic Generalized Epilepsies Have Temperature-Dependent Trafficking Deficiencies. *Journal of Neuroscience* Vol. 26, No. 9, pp. 2590-2597, ISSN 0270-6474

- Klaassen, A., Glykys, J., Maguire, J., Labarca, C., Mody, I., & Boulter, J. (2006). Seizures and enhanced cortical GABAergic inhibition in two mouse models of human autosomal dominant nocturnal frontal lobe epilepsy. *Proceedings of the National Academy of Sciences of the USA* Vol. 103, No. 50, pp. 19152-19157, ISSN 0027-8424
- Kwan, P., Sills, G.J., & Brodie, M.J. (2001). The mechanisms of action of commonly used antiepileptic drugs. *Pharmacology & Therapeutics* Vol. 90, No. 1, pp. 21-34, ISSN 0163-7258
- Laine, M., Lin, M.c., Bannister, J.P.A., Silverman, W.R., Mock, A.F., Roux, B., & Papazian, D.M. (2003). Atomic Proximity between S4 Segment and Pore Domain in Shaker Potassium Channels. *Neuron* Vol. 39, No. 3, pp. 467-481, ISSN 0896-6273
- Lee, J., Taira, T., Pihlaja, P., Ransom, B.R., & Kaila, K. (1996). Effects of CO<sub>2</sub> on excitatory transmission apparently caused by changes in intracellular pH in the rat hippocampal slice. *Brain Research* Vol. 706, No. 2, pp. 210-216, ISSN 0006-8993
- Lee, K.H., & McCormick, D.A. (1997). Modulation of spindle oscillations by acetylcholine, cholecystokinin and 1S,3R-ACPD in the ferret lateral geniculate and perigeniculate nuclei in vitro. *Neuroscience* Vol. 77, No. 2, pp. 335-350, ISSN 0306-4522
- Long, S.B., Campbell, E.B., & MacKinnon, R. (2005). Voltage Sensor of Kv1.2: Structural Basis of Electromechanical Coupling. *Science* Vol. 309, No. 5736, pp. 903-908, ISSN 0036-8075
- Lossin, C., Rhodes, T.H., Desai, R.R., Vanoye, C.G., Wang, D., Carniciu, S., Devinsky, O., & George, A.L., Jr. (2003). Epilepsy-Associated Dysfunction in the Voltage-Gated Neuronal Sodium Channel SCN1A. *Journal of Neuroscience* Vol. 23, No. 36, pp. 11289-11295, ISSN 0270-6474
- Lossin, C., Wang, D.W., Rhodes, T.H., Vanoye, C.G., & George, A.L. (2002). Molecular basis of an inherited epilepsy. *Neuron* Vol. 34, No. 6, pp. 877-884, ISSN 0896-6273
- Main, M.J., Cryan, J.E., Dupere, J.R.B., Cox, B., Clare, J.J., & Burbidge, S.A. (2000). Modulation of KCNQ2/3 Potassium Channels by the Novel Anticonvulsant Retigabine. *Molecular Pharmacology* Vol. 58, No. 2, pp. 253-262, ISSN 0026-895X
- Martin, M.S., Dutt, K., Papale, L.A., Dube, C.M., Dutton, S.B., de Haan, G., Shankar, A., Tufik, S., Meisler, M.H., Baram, T.Z., Goldin, A.L., & Escayg, A. (2010). Altered Function of the SCN1A Voltage-gated Sodium Channel Leads to gamma-Aminobutyric Acid-ergic (GABAergic) Interneuron Abnormalities. *Journal of Biological Chemistry* Vol. 285, No. 13, pp. 9823-9834, ISSN 0021-9258
- Martire, M., Castaldo, P., D'Amico, M., Preziosi, P., Annunziato, L., & Tagliatela, M. (2004). M Channels Containing KCNQ2 Subunits Modulate Norepinephrine, Aspartate, and GABA Release from Hippocampal Nerve Terminals. *Journal of Neuroscience* Vol. 24, No. 3, pp. 592-597, ISSN 0270-6474
- Martire, M., D'Amico, M., Panza, E., Miceli, F., Viggiano, D., Lavergata, F., Iannotti, F.A., Barrese, V., Preziosi, P., Annunziato, L., & Tagliatela, M. (2007). Involvement of KCNQ2 subunits in [3H]dopamine release triggered by depolarization and pre-synaptic muscarinic receptor activation from rat striatal synaptosomes. *Journal of Neurochemistry* Vol. 102, No. 1, pp. 179-193, ISSN 0022-3042
- Mashimo, T., Ohmori, I., Ouchida, M., Ohno, Y., Tsurumi, T., Miki, T., Wakamori, M., Ishihara, S., Yoshida, T., Takizawa, A., Kato, M., Hirabayashi, M., Sasa, M., Mori,

- Y., & Serikawa, T. (2010). A Missense Mutation of the Gene Encoding Voltage-Dependent Sodium Channel (Na(v)1.1) Confers Susceptibility to Febrile Seizures in Rats. *Journal of Neuroscience* Vol. 30, No. 16, pp. 5744-5753, ISSN 0270-6474
- McKernan, R.M., & Whiting, P.J. (1996). Which GABAA-receptor subtypes really occur in the brain? *Trends in Neurosciences* Vol. 19, No. 4, pp. 139-143, ISSN 0166-2236
- Meadows, L.S., Malhotra, J., Loukas, A., Thyagarajan, V., Kazen-Gillespie, K.A., Koopman, M.C., Kriegler, S., Isom, L.L., & Ragsdale, D.S. (2002). Functional and Biochemical Analysis of a Sodium Channel beta 1 Subunit Mutation Responsible for Generalized Epilepsy with Febrile Seizures Plus Type 1. *Journal of Neuroscience* Vol. 22, No. 24, pp. 10699-10709, ISSN 0270-6474
- Meldrum, B.S., & Rogawski, M.A. (2007). Molecular Targets for Antiepileptic Drug Development. *Neurotherapeutics* Vol. 4, No. 1, pp. 18-61, ISSN 1933-7213
- Nabbout, R., Gennaro, E., Dalla Bernardina, B., Dulac, O., Madia, F., Bertini, E., Capovilla, G., Chiron, C., Cristofori, G., Elia, M., Fontana, E., Gaggero, R., Granata, T., Guerrini, R., Loi, M., La Selva, L., Lispi, M.L., Matricardi, A., Romeo, A., Tzolas, V., Valseriati, D., Veggiotti, P., Vigeveno, F., Vallee, L., Bricarelli, F.D., Bianchi, A., & Zara, F. (2003). Spectrum of SCN1A mutations in severe myoclonic epilepsy of infancy. *Neurology* Vol. 60, No. 12, pp. 1961-1967, ISSN 0028-3878
- Noebels, J.L. (2003). THE BIOLOGY OF EPILEPSY GENES. *Annual Review of Neuroscience* Vol. 26, No. 1, pp. 599-625, ISSN 0147-006X
- Oakley, J.C., Kalume, F., Yu, F.H., Scheuer, T., & Catterall, W.A. (2009). Temperature- and age-dependent seizures in a mouse model of severe myoclonic epilepsy in infancy. *Proceedings of the National Academy of Sciences of the USA* Vol. 106, No. 10, pp. 3994-3999, ISSN 0027-8424
- Ogiwara, I., Miyamoto, H., Morita, N., Atapour, N., Mazaki, E., Inoue, I., Takeuchi, T., Itohara, S., Yanagawa, Y., Obata, K., Furuichi, T., Hensch, T.K., & Yamakawa, K. (2007). Nav1.1 Localizes to Axons of Parvalbumin-Positive Inhibitory Interneurons: A Circuit Basis for Epileptic Seizures in Mice Carrying an Scn1a Gene Mutation. *Journal of Neuroscience* Vol. 27, No. 22, pp. 5903-5914, ISSN 0270-6474
- Ohmori, I., Ouchida, M., Ohtsuka, Y., Oka, E., & Shimizu, K. (2002). Significant correlation of the SCN1A mutations and severe myoclonic epilepsy in infancy. *Biochemical and Biophysical Research Communications* Vol. 295, No. 1, pp. 17-23, ISSN 0006-291X
- Ortells, M.O., & Barrantes, G.E. (2002). Molecular modelling of the interactions of carbamazepine and a nicotinic receptor involved in the autosomal dominant nocturnal frontal lobe epilepsy. *British Journal of Pharmacology* Vol. 136, No. 6, pp. 883-895, ISSN 0007-1188
- Otto, J.F., Singh, N.A., Dahle, E.J., Leppert, M.F., Pappas, C.M., Pruess, T.H., Wilcox, K.S., & White, H.S. (2009). Electroconvulsive seizure thresholds and kindling acquisition rates are altered in mouse models of human Kcnq2 and Kcnq3 mutations for benign familial neonatal convulsions. *Epilepsia* Vol. 50, No. 7, pp. 1752-1759, ISSN 0013-9580



- Pan, Z., Kao, T., Horvath, Z., Lemos, J., Sul, J.Y., Cranstoun, S.D., Bennett, V., Scherer, S.S., & Cooper, E.C. (2006). A Common Ankyrin-G-Based Mechanism Retains KCNQ and NaV Channels at Electrically Active Domains of the Axon. *Journal of Neuroscience* Vol. 26, No. 10, pp. 2599-2613, ISSN 0270-6474
- Patino, G.A., & Isom, L.L. (2010). Electrophysiology and beyond: Multiple roles of Na<sup>+</sup> channel beta subunits in development and disease. *Neuroscience Letters* Vol. 486, No. 2, pp. 53-59, ISSN 0304-3940
- Peters, H.C., Hu, H., Pongs, O., Storm, J.F., & Isbrandt, D. (2005). Conditional transgenic suppression of M channels in mouse brain reveals functions in neuronal excitability, resonance and behavior. *Nature Neuroscience* Vol. 8, No. 1, pp. 51-60, ISSN 1097-6256
- Phillips, H.A., Favre, I., & Kirkpatrick, M. (2001). CHRNA2 is the second acetylcholine receptor subunit associated with autosomal dominant nocturnal frontal lobe epilepsy. *American Journal of Human Genetics* Vol. 68, pp. 225-231, ISSN 0002-9297
- Picard, F., Bruel, D., Servent, D., Saba, W., Fruchart-Gaillard, C., Schollhorn-Peyronneau, M.A., Roumenov, D., Brodtkorb, E., Zuberi, S., Gambardella, A., Steinborn, B., Hufnagel, A., Valette, H., & Bottlaender, M. (2006). Alteration of the in vivo nicotinic receptor density in ADNFLE patients: a PET study. *Brain* Vol. 129, No. 8, pp. 2047-2060, ISSN 0006-8950
- Picard, F., Megevand, P., Minotti, L., Kahane, P., Ryvlin, P., Seeck, M., Michel, C.M., & Lantz, G. (2007). Intracerebral recordings of nocturnal hyperkinetic seizures: Demonstration of a longer duration of the pre-seizure sleep spindle. *Clinical Neurophysiology* Vol. 118, No. 4, pp. 928-939, ISSN 1388-2457
- Prole, D.L., Lima, P.A., & Marrion, N.V. (2003). Mechanisms Underlying Modulation of Neuronal KCNQ2/KCNQ3 Potassium Channels by Extracellular Protons. *The Journal of General Physiology* Vol. 122, No. 6, pp. 775-793, ISSN 0022-1295
- Rasmussen, H.B., Frokjaer-Jensen, C., Jensen, C.S., Jensen, H.S., Jorgensen, N.K., Misonou, H., Trimmer, J.S., Olesen, S.P., & Schmitt, N. (2007). Requirement of subunit co-assembly and ankyrin-G for M-channel localization at the axon initial segment. *Journal of Cell Science* Vol. 120, No. 6, pp. 953-963, ISSN 0021-9533
- Ronen, G.M., Rosales, T.O., Connolly, M., Anderson, V.E., & Leppert, M. (1993). Seizure characteristics in chromosome 20 benign familial neonatal convulsions. *Neurology* Vol. 43, No. 7, pp. 1355-1360, ISSN 0028-3878
- Rundfeldt, C., & Netzer, R. (2000). The novel anticonvulsant retigabine activates M-currents in Chinese hamster ovary-cells transfected with human KCNQ2/3 subunits. *Neuroscience Letters* Vol. 282, No. 1-2, pp. 73-76, ISSN 0304-3940
- Scheffer, I.E., & Berkovic, S.F. (1997). Generalized epilepsy with febrile seizures plus. A genetic disorder with heterogeneous clinical phenotypes. *Brain* Vol. 120, No. 3, pp. 479-490, ISSN 0006-8950
- Schroeder, B.C., Hechenberger, M., Weinreich, F., Kubisch, C., & Jentsch, T.J. (2000). KCNQ5, a Novel Potassium Channel Broadly Expressed in Brain, Mediates M-type Currents. *Journal of Biological Chemistry* Vol. 275, No. 31, pp. 24089-24095, ISSN 0021-9258




- Schroeder, B.C., Kubisch, C., Stein, V., & Jentsch, T.J. (1998). Moderate loss of function of cyclic-AMP-modulated KCNQ2/KCNQ3 K<sup>+</sup> channels causes epilepsy. *Nature* Vol. 396, No. 6712, pp. 687-690, ISSN 0028-0836
- Schuchmann, S., Schmitz, D., Rivera, C., Vanhatalo, S., Salmen, B., Mackie, K., Sipila, S.T., Voipio, J., & Kaila, K. (2006). Experimental febrile seizures are precipitated by a hyperthermia-induced respiratory alkalosis. *Nature Medicine* Vol. 12, No. 7, pp. 817-823, ISSN 1078-8956
- Schwake, M., Pusch, M., Kharkovets, T., & Jentsch, T.J. (2000). Surface Expression and Single Channel Properties of KCNQ2/KCNQ3, M-type K<sup>+</sup> Channels Involved in Epilepsy. *Journal of Biological Chemistry* Vol. 275, No. 18, pp. 13343-13348, ISSN 0021-9258
- Selyanko, A.A., Hadley, J.K., Wood, I.C., Abogadie, F.C., Jentsch, T.J., & Brown, D.A. (2000). Inhibition of KCNQ1-4 potassium channels expressed in mammalian cells via M1 muscarinic acetylcholine receptors. *The Journal of Physiology* Vol. 522, No. 3, pp. 349-355, ISSN 0022-3751
- Sieghart, W., & Sperk, G. (2002). Subunit Composition, Distribution and Function of GABA-A Receptor Subtypes. *Current Topics in Medicinal Chemistry* Vol. 2, No. 8, pp. 795-816, ISSN 1568-0266
- Singh, N.A., Charlier, C., Stauffer, D., DuPont, B.R., Leach, R.J., Melis, R., Ronen, G.M., Bjerre, I., Quattlebaum, T., Murphy, J.V., McHarg, M.L., Gagnon, D., Rosales, T.O., Peiffer, A., Anderson, V.E., & Leppert, M. (1998). A novel potassium channel gene, KCNQ2, is mutated in an inherited epilepsy of newborns. *Nature Genetics* Vol. 18, No. 1, pp. 25-29, ISSN 1061-4036
- Singh, N.A., Otto, J.F., Dahle, E.J., Pappas, C., Leslie, J.D., Vilaythong, A., Noebels, J.L., White, H.S., Wilcox, K.S., & Leppert, M.F. (2008). Mouse models of human KCNQ2 and KCNQ3 mutations for benign familial neonatal convulsions show seizures and neuronal plasticity without synaptic reorganization. *The Journal of Physiology* Vol. 586, No. 14, pp. 3405-3423, ISSN 0022-3751
- Soldovieri, M.V., Castaldo, P., Iodice, L., Miceli, F., Barrese, V., Bellini, G., del Giudice, E.M., Pascotto, A., Bonatti, S., Annunziato, L., & Tagliatela, M. (2006). Decreased Subunit Stability as a Novel Mechanism for Potassium Current Impairment by a KCNQ2 C Terminus Mutation Causing Benign Familial Neonatal Convulsions. *Journal of Biological Chemistry* Vol. 281, No. 1, pp. 418-428, ISSN 0021-9258
- Spampanato, J., Escayg, A., Meisler, M.H., & Goldin, A.L. (2003). Generalized epilepsy with febrile seizures plus type 2 mutation W1204R alters voltage-dependent gating of Na(v)1.1 sodium channels. *Neuroscience* Vol. 116, No. 1, pp. 37-48, ISSN 0306-4522
- Spampanato, J., Kearney, J.A., de Haan, G., McEwen, D.P., Escayg, A., Aradi, I., MacDonald, B.T., Levin, S.I., Soltesz, I., Benna, P., Montalenti, E., Isom, L.L., Goldin, A.L., & Meisler, M.H. (2004). A novel epilepsy mutation in the sodium channel SCN1A identifies a cytoplasmic domain for beta subunit interaction. *Journal of Neuroscience* Vol. 24, No. 44, pp. 10022-10034, ISSN 0270-6474
- Steinlein, O.K., Magnusson, A., Stoodt, J., Bertrand, S., Weiland, S., Berkovic, S.F., Nakken, K.O., Propping, P., & Bertrand, D. (1997). An insertion mutation of the CHRNA4

- gene in a family with autosomal dominant nocturnal frontal lobe epilepsy. *Human Molecular Genetics* Vol. 6, No. 6, pp. 943-947, ISSN 0964-6906
- Steinlein, O.K., Mulley, J.C., Propping, P., Wallace, R.H., Phillips, H.A., Sutherland, G.R., Scheffer, I.E., & Berkovic, S.F. (1995). A missense mutation in the neuronal nicotinic acetylcholine receptor [alpha]4 subunit is associated with autosomal dominant nocturnal frontal lobe epilepsy. *Nature Genetics* Vol. 11, No. 2, pp. 201-203, ISSN 1061-4036
- Sugawara, T., Tsurubuchi, Y., Agarwala, K.L., Ito, M., Fukuma, G., Mazaki-Miyazaki, E., Nagafuji, H., Noda, M., Imoto, K., Wada, K., Mitsudome, A., Kaneko, S., Montal, M., Nagata, K., Hirose, S., & Yamakawa, K. (2001). A missense mutation of the Na<sup>+</sup> channel alpha II subunit gene Nav1.2 in a patient with febrile and afebrile seizures causes channel dysfunction. *Proceedings of the National Academy of Sciences of the USA* Vol. 98, No. 11, pp. 6384-6389, ISSN 0027-8424
- Tang, B., Dutt, K., Papale, L., Rusconi, R., Shankar, A., Hunter, J., Tufik, S., Yu, F.H., Catterall, W.A., Mantegazza, M., Goldin, A.L., & Escayg, A. (2009). A BAC transgenic mouse model reveals neuron subtype-specific effects of a Generalized Epilepsy with Febrile Seizures Plus (GEFS plus ) mutation. *Neurobiology of Disease* Vol. 35, No. 1, pp. 91-102, ISSN 0969-9961
- Wallace, R.H., Wang, D.W., Singh, R., Scheffer, I.E., George, A.L., Phillips, H.A., Saar, K., Reis, A., Johnson, E.W., Sutherland, G.R., Berkovic, S.F., & Mulley, J.C. (1998). Febrile seizures and generalized epilepsy associated with a mutation in the Na<sup>+</sup>-channel  $\alpha 1$  subunit gene SCN1B. *Nature Genetics* Vol. 19, No. 4, pp. 366-370, ISSN 1061-4036
- Wan, Q., Xiong, Z.G., Man, H.Y., Ackerley, C.A., Branton, J., Lu, W.Y., Becker, L.E., MacDonald, J.F., & Wang, Y.T. (1997). Recruitment of functional GABAA receptors to postsynaptic domains by insulin. *Nature* Vol. 388, No. 6643, pp. 686-690, ISSN 0028-0836
- Watanabe, H., Nagata, E., Kosakai, A., Nakamura, M., Yokoyama, M., Tanaka, K., & Sasai, H. (2000). Disruption of the epilepsy KCNQ2 gene results in neural hyperexcitability. *Journal of Neurochemistry* Vol. 75, No. 1, pp. 28-33, ISSN 0022-3042
- Waxham, M.N. (2003). Neurotransmitter receptors. In: *Fundamental Neuroscience*, Squire, L.R., Bloom, F.E., McConnell, S.K., Roberts, J.L., Spitzer, N.C., & Zigmond, M.J., (Eds.) pp. 225-258, Academic Press, ISBN 0-12-660303-0, California, USA.
- Waxman, S.G. (2007). Channel, neuronal and clinical function in sodium channelopathies: from genotype to phenotype. *Nature Neuroscience* Vol. 10, No. 4, pp. 405-409, ISSN 1097-6256
- Weber, Y.G., Geiger, J., Kampchen, K., Landwehrmeyer, B., Sommer, C., & Lerche, H. (2006). Immunohistochemical analysis of KCNQ2 potassium channels in adult and developing mouse brain. *Brain Research* Vol. 1077, No. 1, pp. 1-6, ISSN 0006-8993
- Wickenden, A.D., Yu, W., Zou, A., Jegla, T., & Wagoner, P.K. (2000). Retigabine, A Novel Anti-Convulsant, Enhances Activation of KCNQ2/Q3 Potassium Channels. *Molecular Pharmacology* Vol. 58, No. 3, pp. 591-600, ISSN 0026-895X
- Yu, F.H., & Catterall, W.A. (2003). Overview of the voltage-gated sodium channel family. *Genome Biology* Vol. 4, No. 3, pp. 207, ISSN 1465-6906

Yu, F.H., Mantegazza, M., Westenbroek, R.E., Robbins, C.A., Kalume, F., Burton, K.A., Spain, W.J., McKnight, G.S., Scheuer, T., & Catterall, W.A. (2006). Reduced sodium current in GABAergic interneurons in a mouse model of severe myoclonic epilepsy in infancy. *Nature Neuroscience* Vol. 9, No. 9, pp. 1142-1149, ISSN 1097-6256

IntechOpen

IntechOpen



### **Novel Aspects on Epilepsy**

Edited by Prof. Humberto Foyaca-Sibat

ISBN 978-953-307-678-2

Hard cover, 338 pages

**Publisher** InTech

**Published online** 12, October, 2011

**Published in print edition** October, 2011

This book covers novel aspects of epilepsy without ignoring its foundation and therefore, apart from the classic issues that cannot be missing in any book about epilepsy, we introduced novel aspects related with epilepsy and neurocysticercosis as a leading cause of epilepsy in developing countries. We are looking forward with confidence and pride in the vital role that this book has to play for a new vision and mission. Therefore, we introduce novel aspects of epilepsy related to its impact on reproductive functions, oral health and epilepsy secondary to tuberous sclerosis, mitochondrial disorders and lisosomal storage disorders.

#### **How to reference**

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Sigrid Marie Blom and Henrik Sindal Jensen (2011). Epileptic Channelopathies and Dysfunctional Excitability - From Gene Mutations to Novel Treatments, Novel Aspects on Epilepsy, Prof. Humberto Foyaca-Sibat (Ed.), ISBN: 978-953-307-678-2, InTech, Available from: <http://www.intechopen.com/books/novel-aspects-on-epilepsy/epileptic-channelopathies-and-dysfunctional-excitability-from-gene-mutations-to-novel-treatments>

**INTECH**  
open science | open minds

#### **InTech Europe**

University Campus STeP Ri  
Slavka Krautzeka 83/A  
51000 Rijeka, Croatia  
Phone: +385 (51) 770 447  
Fax: +385 (51) 686 166  
[www.intechopen.com](http://www.intechopen.com)

#### **InTech China**

Unit 405, Office Block, Hotel Equatorial Shanghai  
No.65, Yan An Road (West), Shanghai, 200040, China  
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元  
Phone: +86-21-62489820  
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen