

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

185,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



The Impact of Bacteria-Induced Adaptive Immune Responses in Periodontal Disease

Vincent K. Tsiagbe^{1,2,3} and Daniel H. Fine^{1,2}

¹Department of Oral Biology, New Jersey Dental School,

²Graduate School of Biomedical Science,

³Department of Pathology and Laboratory Medicine, New Jersey Medical School,

University of Medicine and Dentistry of New Jersey,

Newark, NJ,

USA

1. Introduction

More than one microorganism causes periodontal disease, like many infectious diseases in humans. Because of the complexity of “polymicrobial infections”, their study requires a multidisciplinary approach, employing specific in vitro techniques, and various animal models (Bakaletz 2004). Inherently, no one approach or animal model can completely elucidate the mechanisms of periodontal disease. Notwithstanding these difficulties, animal models do provide critically important information regarding periodontal disease pathogenesis (Graves 2008). Another layer of complexity resides in the fact that different strains of bacteria, as in the case of *P. gingivalis* (*Pg*)-induced disease, cause different levels of disease in the same mouse strain (Baker and Roopenian 2002). Similarly, differences in disease susceptibility can result from the same bacterium strain, as in the case of *Aggregatibacter actinomycetemcomitans* (*Aa*) infected rodents (Fine et al. 2009). Furthermore, different strains of rodents exhibit different susceptibilities to challenge from the same strain of bacterium, such as *Pg* (Baker et al. 2000). This finding also holds true for *Aa* (Schreiner et al. 2011). These observations resemble findings in human disease, especially in the case of *Aa*-related disease where the JP2 strain of *Aa* appears to be more virulent than other *Aa* strains, and where individuals of African heritage appear to be more susceptible to *Aa*-induced periodontal disease than Caucasian individuals (Haubek et al. 2008). The similarities between rodent and human bacterial-induced periodontal diseases lend credence to the validity of the animal model designed to assess this disease (Fine 2009).

The virulence factors elaborated by pathogenic microorganisms and the host immunologic responses to such factors play a major role in disease induction and progression. It has been established that *Aa*, a gram-negative facultative capnophilic rod, is the causative agent in localized juvenile periodontitis (LJP) (Zambon 1985). This agent is also a key pathogen for localized progressing and severe forms of adult periodontitis (Dzink et al. 1985; Zambon et al. 1988). *Aa* possesses several virulence factors, including endotoxin and leukotoxin (Fives-Taylor et al. 1999). *Aa* secretes a protein toxin, Leukotoxin (LtxA), which helps the bacterium evade the host immune response during infection, by specifically targeting white blood cells

(WBCs) (Kachlany 2010). The ability of LtxA to bind WBCs from humans and Old World primates, by interacting with lymphocyte function antigen-1 (LFA-1) on susceptible cells, has opened a window of opportunity for the use of LtxA as a novel therapeutic agent in leukemia (Kachlany et al. 2010). *Aa* also produces cytolethal distending toxin (Cdt), which is a potent immunotoxin that induces G2 arrest in human lymphocytes (Shenker et al. 2007). The immunologic and systemic impact of these bacterial toxins in periodontal disease is yet to be clarified.

The binding of *Aa* to buccal epithelial cells (BEC) was shown to be mediated by two *Aa* autotransporter adhesions (ApiA and Aae), which work, in concert to modulate *Aa* binding to BEC, specifically in humans and Old World Monkeys (Yue et al. 2007). The type and extent of the immunologic response mounted in response to oral pathogen will undoubtedly depend on the particular microbial pathogen(s), the virulence factors invoked and the genetic background of the host. The immunologic reactions mounted in response to oral pathogens have a potential to precipitate other unforeseen systemic diseases of grave importance. The connection between immunologic responses to oral pathogens and systemic diseases is mostly unexplored, at present.

2. Immune responses to microbial pathogens

Our understanding of periodontal pathogenesis has evolved over the years, and has transformed from periodontitis being considered an almost ubiquitous condition in which the role of plaque was thought to be the sole aetiological factor to today where concepts of inflammation and individual susceptibility are considered (Preshaw and Taylor 2011). Neutrophils are a critical arm of the host defense in periodontitis, but bacterial evasion of neutrophil microbicidal machinery, together with delayed neutrophil apoptosis can transform neutrophil from defender to perpetrator (Nussbaum and Shapira 2011). In the recent Seventh European Workshop on Periodontology, aimed at understanding cellular and molecular mechanisms of host microbial interactions, a consensus was reached that "PMNs are important in the pathophysiology of periodontal disease but there is limited evidence on their much quoted destructive potential". Cytokine networks are enormously complex and we are really at the beginning of understanding their role in the disease process (Kinane, Preshaw and Loos 2011). Thus, there is an emerging appreciation for the complex role played by the adaptive immune system in responses to periodontal pathogens.

2.1 Humoral Immune responses to microbial pathogens

In early studies, significantly elevated serum immunoglobulin G (IgG) antibody levels to *B. gingivalis* were seen in adult and advanced destructive periodontitis patients, suggestive of distinctive host-parasite interactions in this disease (Ebersole and Cappelli 1994; Ebersole et al. 1986). Analysis of the proportion of various cell types present in gingival biopsies retrieved from subjects with severe chronic periodontitis showed that the proportion of B cells was larger than that of T cells, plasma cells and neutrophils. Furthermore, about 60% of the B cells were of the autoreactive B-1a sub-population (CD19⁺CD5⁺) (Donati et al. 2009). The plasma cells that developed were shown to derive from both B-2 cells (conventional B cells) and B-1a cells. There is strong evidence that B cells serve as antigen presenting cells in periodontitis (Gemmell et al. 2002; Mahanonda et al. 2002). Indeed, upregulation of the co-stimulatory molecule, CD86 (B7.2), and the dendritic cell marker, CD83, on B cells in

periodontal lesions, have been reported (Gemmell et al. 2002). Thus, it is likely that the B cells found in periodontal tissue might present bacterial antigens to host T cells, leading to the elaboration of a whole range of cytokines, the nature of which would depend on the type of bacteria, and the host.

Altered CD4/CD8 T-cell ratios and autologous mixed-lymphocyte reaction in LJP, suggested a potential regulatory role of T cells in periodontitis. Using immunohistochemical and *in situ* hybridization techniques, a higher frequency of CD4⁺CD45RO⁺ cells expressing IL-4 has been seen in lesions from individuals with chronic periodontitis compared to normal tissue (Yamazaki et al. 1994). Comparing two different compartments (peripheral blood vs. periodontal tissue), it was noted that even though mRNA for IL-12 and IL-13 were similar between the two compartments, the level of IFN- γ was higher in circulating cells than in gingival cells. Inversely, IL-10 expression was higher in the gingival cells (Yamazaki et al. 1997). Moreover, the frequency of IL-10 expressing CD14⁺ cells was higher in peripheral blood of chronic periodontitis, but not acute periodontitis patients, compared to healthy controls (Yamazaki et al. 1997).

In periodontal disease, the development of gingivitis involves Th1 cells, while in periodontitis, there is a shift toward Th2 cells (reviewed in (Berglundh and Donati 2005)). Autoimmune reactions do occur in periodontitis lesions; however, the role of autoantibodies in the regulation of host response in periodontitis needs to be clarified (Berglundh and Donati 2005). In studies conducted with *Aa*-induced periodontal disease rat model, we observed an early increase in serum IgG2a antibody 2-4 weeks post inoculation. This was accompanied by a concomitant increase in LtxA-specific IgG production, suggesting that the immune response was mediated by *Aa* (Li et al. 2010). An increase in B and CD4 T cell numbers in draining cervical and submandibular lymph nodes accompanied this *Aa*-specific antibody production. CD8 T cell numbers were not examined in this study (Li et al. 2010). In agreement with this observation, there was an increase in the expression of CD70 (TNFSF7) in B cells harvested from draining lymph nodes in rats infected by *Aa* (Li et al. 2010). CD70 has been shown to be expressed on a subpopulation of germinal center B cells (Hintzen et al. 1994).

2.2 Cytokines in periodontal disease

Innate immunity is mediated by macrophages, dendritic cells (DCs), neutrophils, monocytes, epithelial cells and endothelial cells that recognize and temporarily respond to pathogen associated molecular patterns (PAMPS), like LPS on gram-negative bacteria. The adaptive immune system, on the other hand, uses specific antigen recognition structures on T and B cell. Such responses are specific and maintained by the generation of memory. Various cytokines generated by macrophages and DCs create a milieu, which determines the differentiation of particular effector T-cell subsets as well as the class and subclass of immunoglobulin (Ig) antibodies synthesized. Cytokines act in concert with other signalling pathways and, especially, cell-to-cell interactions via antigen presentation and co-stimulatory molecules (Preshaw and Taylor 2011).

The role of inflammatory cytokines, such as interleukin (IL)-1 β , tumor necrosis factor- α , and IL-6, has been the most understood (reviewed in (Preshaw and Taylor 2011)). Inhibition of IL-1 and tumor necrosis factor (TNF) resulted in amelioration of bone loss in experimental periodontitis (Assuma et al. 1998; Graves et al. 1998). In our studies on *Aa*-induced periodontal disease, the early induction of *Aa*-specific IgG and IgG2a antibodies in *Aa*-fed rats is of interest since mRNA for Th1 cytokines TNF and lymphotoxin beta (LT β) (Abbas,

Murphy and Sher 1996) were upregulated early (2-4 weeks) in the inflammatory response, which could explain the significant switch in *Aa*-specific antibody production to IgG2a. This is consistent with the observation that Th1 cytokines drive isotype switching to IgG2a in inflammatory responses of atherosclerosis (Schulte, Sukhova and Libby 2008).

2.2.1 Th1/Th2 paradigm

Cytokines mediate and sustain the development and function of CD4⁺ Th cell subsets. In the original description of Th cell dichotomy, Th1 cells secrete interferon- γ (IFN- γ), and promote cell-mediated immunity by activating macrophages, natural killer (NK) cells and cytotoxic CD8⁺ T-cells, whereas Th2 cells secrete IL-4, IL-5 and IL-13 and regulate humoral (antibody-mediated) immunity and mast cell activity (Mosmann and Coffman 1989). It was conjectured that the dynamic interaction between T-cell subsets might result in fluctuations in disease activity and that a Th1 response (providing protective cell-mediated immunity) underlies a “stable” periodontal lesion, and a Th2 response (leading to activation of B-cells) mediates a destructive lesion possibly through enhanced B-cell-derived IL-1 β (Gemmell, Yamazaki and Seymour 2007; Seymour and Gemmell 2001). It is now becoming clearer that the Th1/Th2 model alone is inadequate to explain the role of T-cells in periodontal disease process (Gaffen and Hajishengallis 2008).

2.2.2 Role of Th17 cells

Th17 cells secrete the IL-17 cytokines (which have a number of pro-inflammatory activities in common with IL-1 β and TNF α) and IL-22, and are crucial for immunity against extracellular bacteria (Miossec, Korn and Kuchroo 2009). Th17 cells have been implicated in the pathogenesis of several autoimmune and inflammatory disorders, and *in vitro* polarization of human and mouse Th17 cells is under the influence of Notch1 (Keerthivasan et al. 2011). Studies have shown that IL-17A produced by Th17 cells stimulate the development of osteoclasts (osteoclastogenesis) in the presence of osteoblasts (Zhang et al. 2011), and expression of IL-17 has been observed in gingiva from patients with periodontitis (Cardoso et al. 2009).

In our studies on *Aa*-induced rat model for periodontal disease, we observed upregulation in IL-17 in CD4⁺ T cells (2.8 fold) and B cells (2 fold), in lymph nodes from *Aa*-infected rats, compared to control rats. This level of expression was below our stringent criterion of four-fold differential gene expression in this study. However, this finding is in conformity with the observation that IL-17 might be involved in inflammatory response and bone resorption in periodontal disease animal models (Oseko et al. 2009) (Xiong, Wei and Peng 2009). It should, however, be noted that T cells exhibit “functional plasticity” that is influenced by the cytokine milieu (Bluestone et al. 2009). For instance, Th17 cells can differentiate into Th1 cells, under the influence of IL-12 (Korn et al. 2009), and follicular T helper cells (Thf), present in the B cell follicles of lymph nodes, are dependent on IL-6 and IL-21 for their development, and are capable of secreting a cytokine profile corresponding to Th1, Th2 or Th17 cells (Korn et al. 2009).

2.2.3 Role of regulatory (Treg) cells

It has been established that naturally arising Foxp3⁺CD4⁺CD25⁺ (Treg) cells play a central role in the maintenance of immunological tolerance (Sakaguchi 2005). Treg cells secrete transforming growth factor- β (TGF- β) and IL-10 which are critical in regulating other T-cell

subsets and maintaining tolerance against self-antigens, thereby preventing autoimmunity (Josefowicz and Rudensky 2009). Gingival mononuclear cells from mice infected with *Pg* were found to exhibit increased levels of Treg cells 30 days post infection, suggesting that there are potential roles for Treg cells during the chronic stage of periodontitis in the regulation of gingival inflammation and alveolar bone loss (Kobayashi et al. 2011). FoxP3⁺CD8⁺ T cells, with suppressive function have recently been identified in simian immunodeficiency virus infected rhesus macaques, and in HIV-1 infected humans. Expansion of CD8⁺ Tregs correlated directly with acute phase viremia and inversely with the magnitude of antiviral T cell response (Nigam et al. 2010). Using transgenic OT-I mice, the administration of ovalbumin (OVA) enabled osteoclasts to cross-present OVA to Ag-specific CD8⁺ T cells to induce their proliferation, and secretion of IL-2, IL-6, and IFN- γ . CD8⁺ T cells activated by osteoclasts expressed FoxP3, CTLA4 and RANKL. Those CD8⁺ T cells were found to be anergic and suppressed dendritic cell priming of naive responder CD8⁺ T cells (Kiesel, Buchwald and Aurora 2009). The role of this novel group of CD8⁺ Treg cells in periodontal disease requires further examination.

2.2.4 Novel cytokine roles in periodontal disease

In our studies on *Aa*-induced periodontal disease rat model, we observed upregulation in mRNA for a number of cytokines, not normally ascribed to periodontal disease. IL-16 was upregulated in CD4 T cells in the early phase of the response (Li et al. 2010). IL-16 has been shown to be involved in the selective migration of CD4 T cells, and participates in inflammatory diseases (Akiyama et al. 2009). It was detected in gingival crevicular fluid (Sakai et al. 2006). IL-19, a novel cytokine of the IL-10 family, was also upregulated in CD4 T cells in response to *Aa*. IL-19 produced by synovial cell in Rheumatoid arthritis (RA) patients promotes joint inflammation (Sakurai et al. 2008). IL-21, which has recently been shown to induce receptor activator of nuclear factor kappaB ligand (RANKL) and was implicated in arthritis (Jang et al. 2009), was upregulated in B cells responding to *Aa*. There was also an induction of IL-24 by 12 weeks in CD4 T cells responding to *Aa*. Studies conducted on RA showed an increase in IL-24 in the synovium of RA patients, and this cytokine was implicated in recruitment of neutrophil granulocytes (Kragstrup et al. 2008). B-cell-activating-factor (BAFF, or TNFSF13B) and a proliferation-inducing ligand (APRIL), members of the TNF family, were upregulated in B cells and CD4 T cells, respectively, in response to *Aa* infection. Both of these factors were found to be upregulated in children with atopic dermatitis (Jee et al. 2009), and thus would represent factors that characterize *Aa*-induced periodontal disease.

IL-23, a proinflammatory cytokine composed of IL-23p19 and IL-12/23p40 subunits, is known to promote the differentiation of Th17 cells. Studies showed that IL-23 and IL-12 were expressed at significantly higher levels in periodontal lesions than in control sites, suggesting that IL-23-induced Th17 pathway is stimulated in inflammatory periodontal lesions (Ohyama et al. 2009). IL-33 is a new member of the IL-1 family, which plays a role in inflammatory response. Injection of TNF transgenic mice, overexpressing human TNF, with IL-33 or IL-33R agonistic antibody inhibited the development of spontaneous joint inflammation and cartilage destruction. Furthermore, *in vitro*, IL-33 directly inhibits mouse and human M-CSF/receptor activator for NF- κ B ligand-driven osteoclast differentiation, suggesting an important role for IL-33 as a bone-protecting cytokine with potential for treating bone resorption (Zaiss et al. 2011).

2.2.5 Role of RANKL and related molecules

RANKL plays a role in T cell-mediated bone resorption. Interference with RANKL by systemic administration of osteoprotegerin (OPG), the decoy receptor for (and inhibitor of) RANKL, was found to result in abrogation of periodontal bone resorption in a rat model (Taubman et al. 2005). Studies in humans have demonstrated that RANKL levels in gingival crevicular fluid (GCF) were low in health or gingivitis, but increased in periodontitis. On the other hand, OPG levels were higher in health than periodontitis, or gingivitis groups (Bostanci et al. 2007). Thus, GCF RANKL and OPG levels were oppositely regulated in periodontitis, but not gingivitis, resulting in an enhanced RANKL/OPG ratio. In our studies with *Aa*-induced periodontal disease rat model, while the bone resorption protein RANKL (TNFSF11) was induced in CD4 T cells from *Aa*-fed rats, its soluble decoy receptor OPG (TNFSF11b) was also induced in the CD4 T cells (Li et al. 2010). Developments in the field of osteoimmunology, which examine the crosstalk of immune cells and bone, have uncovered a novel role for the RANKL-RANK-OPG system in other processes such as in controlling autoimmunity or immune responses in the skin (Leibbrandt and Penninger 2010). Despite the sustained upregulation of OPG, bone resorption still occurred. The critical balance between osteoblast-mediated bone formation and osteoclast-mediated bone resorption has been described as “coupling” of bone formation to bone resorption (Parfitt 1982).

2.2.6 Role of BMPs and GDFs in periodontal disease

Bone morphogenic proteins (BMPs) and growth differentiation factors (GDFs) are members of the transforming growth factor- β (TGF- β) superfamily. They play important roles during development and organogenesis in delivering positional information in both vertebrates and invertebrates, and are involved in the development of hard as well as soft tissue (Herpin, Lelong and Favrel 2004).

BMPs can also act locally on target tissues to affect proliferation and survival (Rosen 2006). BMP2, even though dispensable for bone formation, is a necessary component of the signaling cascade that governs fracture repair (Tsuji et al. 2006). In our studies, BMP2 was induced in B cells early (week 4) of an inflammatory process, at the same time that RANKL was induced in CD4 T cells (Li et al. 2010). This suggests that bone repair mechanisms were induced early, well ahead of impending bone resorption. However, by 12 weeks of infection by *Aa*, BMP2 was shut down, as bone resorption proceeded. BMP3 was also upregulated at week 4 in B cells responding to *Aa*. BMP3 has been shown to be a negative regulator in the skeleton, as mice lacking BMP3 have increased bone mass. Transgenic mice over-expressing BMP3 had altered endochondral bone formation resulting in spontaneous rib fractures (Gamer et al. 2009). On the other hand, it has been suggested that BMP2 and BMP3 might be co-regulated. BMP-2 was found to enhance BMP-3 and -4 mRNA expressions in primary cultures of fetal rat calvarial osteoblasts. The enhancement of BMP-3 and -4 mRNA expressions by BMP-2 was associated with an increased expression of bone cell differentiation marker genes (Chen et al. 1997). It is of interest that BMP2 and BMP3 were upregulated in B cells at the same time (4 weeks post infection), and were shut down at 12 weeks, at which time bone resorption was evident.

In our studies with *Aa*- rat model for periodontal disease, we found that B cells responding to *Aa* upregulated BMP10 at all time points (Li et al. 2010). BMP10 has been shown to regulate myocardial hypertrophic growth (Chen et al. 2006), and may function as a tumor

suppressor and apoptosis regulator for prostate cancer (Ye, Kynaston and Jiang 2009). To our knowledge, our work is the first report on the production of BMP10 by B cells responding to infection. The expression pattern of BMP10 in our studies, suggests that it might be involved in inflammation, as well as in bone resorption. Furthermore, the involvement of BMP10 in cardiac hypertrophy and cancer, suggests that it might represent one of the possible “missing links” between periodontal disease and other systemic diseases like heart disease and cancer. Evidence for this is provided in the modeled biological interaction pathway depicted in Fig 1.




Fig. 1. Proposed biological interaction network of differentially expressed genes from B and CD4 T cells of *Aa*-fed rats at 12 weeks post infection by *Aa*, and their relationship to disease. Genes upregulated by at least four-fold (i.e. Log₂ fold greater than 2) in B and CD4 T cells derived from cervical and submandibular lymph nodes of *Aa*-fed rats, in comparison to B and CD4 T cells from control rats, were imported into Pathway Studio (Ariadne Genomics, Inc., Rockville, MD, USA) (Yuryev et al. 2006) for analyses. The picture shows interactions between upregulated genes in the expression data (shown as green highlights) and their interactions with related genes and diseases. The biological relationships revealed by the network are depicted in the pallets at the right of the figure. The relevance of the expression data to various diseases, as determined by the mining of the published Resnet 7 database in Pathway Studio, is indicated in the network. Reprinted with permission from Li Y *et al.* Molecular Oral Microbiology 2010; 25:275-292.

Growth differentiation factor 11 (GDF11) or BMP11, plays an important role in establishing embryonic axial skeletal patterns (McPherron, Lawler and Lee 1999). Transfection of GFF11 gene was found to stimulate a large amount of reparative dentin formation in amputated dental pulp of canine teeth in vivo (Nakashima et al. 2003). In our studies with *Aa*-induced periodontal disease rat model, GDF11 was upregulated at 12 weeks post infection, in both B and CD4 T cells, at the time of bone resorption. This suggests that GDF11 may have a novel role in bone resorption. The fact that GDF11 activation has been observed in cancer (Yokoe et al. 2007), may also provide another possible link between periodontal disease and cancer.

Growth differentiation factor 15 (GDF15), was upregulated in both B and CD4 T cells of *Aa*-infected rats at 12 week, coinciding with the time of bone resorption. However, there are conflicting reports on the role of GDF15 in bone resorption and other systemic diseases. Studies have shown that pure GDF15 and the GDF15-containing growth medium of 1,25(OH)₂-vitamin D₃-treated prostate adenocarcinoma LNCaP cells suppress osteoclast differentiation (Vanhara et al. 2009). In addition, elevation in GDF15 has been associated with cardiovascular disease (Kempf and Wollert 2009), and colorectal cancer metastasis (Xue et al.). Thus, GDF15 may also contribute another possible link between periodontal disease and systemic diseases.

3. Conclusions

The nature of the adaptive response to oral microbial insult is vastly dependent on the nature of the microbe, the host (including genetic background), as well as the milieu of prevailing cytokines and chemokines. The *Aa*-induced rat model and *Pg*-induced mouse model for periodontal disease have provided extensive knowledge about role of several previously uncharacterized genes in periodontal disease, however, much more work needs to be done. Therefore, examination of B and CD4 T cells from lymph nodes draining the oral cavity of *Aa*-fed rats showed that inflammatory processes are initially activated early (2-4 weeks) post infection. This, ultimately, leads to activation of bone resorption pathways that end in overt bone resorption by 12 weeks post infection. Apart from induction of known inflammatory cytokines (such as TNF α , IL-1 β , and LT β), other cytokines and TGF- β superfamily member genes, not previously associated with bone resorption, were found to be upregulated in B and/or CD4 T cells. Some of these genes have known effects on systemic diseases such as heart disease, cancer, autoimmune disease, and diabetes. The role of CD8 T cells in adaptive immune responses to periodontal pathogens is not yet clarified. This evidence suggests a subtle link between periodontal disease and other systemic diseases. In conclusion, animal studies have played an important role in unraveling key elements of our understanding of microbial pathogenesis in many human diseases (Shea et al. 2010). The availability of new and more complete data from mouse and rat genome studies coupled with the access to powerful tools that can uncover microbial and host expression can provide novel ways to examine periodontal disease pathogenesis. Application of these tools can allow for comparisons to common pathways with respect to other infectious diseases. This chapter has presented some data derived from the application of one of these new immune response pathway tools to microbial-induced periodontal disease in a rat model.

4. Acknowledgements

This work was supported by grants from the Foundation of University of Medicine and Dentistry of New Jersey (grant #36-08 and PC31-10), and by grant DE-016306 from the National Institute for Dental and Craniofacial Research.

5. References

- Abbas, A. K., K. M. Murphy, and A. Sher. 1996. "Functional diversity of helper T lymphocytes." *Nature* 383(6603):787-93.
- Akiyama, K., M. Karaki, R. Kobayshi, H. Dobashi, T. Ishida, and N. Mori. 2009. "IL-16 variability and modulation by antiallergic drugs in a murine experimental allergic rhinitis model." *Int Arch Allergy Immunol* 149(4):315-22.
- Assuma, R., T. Oates, D. Cochran, S. Amar, and D. T. Graves. 1998. "IL-1 and TNF antagonists inhibit the inflammatory response and bone loss in experimental periodontitis." *J Immunol* 160(1):403-9.
- Bakaletz, L. O. 2004. "Developing animal models for polymicrobial diseases." *Nat Rev Microbiol* 2(7):552-68.
- Baker, P. J., M. Dixon, R. T. Evans, and D. C. Roopenian. 2000. "Heterogeneity of Porphyromonas gingivalis strains in the induction of alveolar bone loss in mice." *Oral Microbiol Immunol* 15(1):27-32.
- Baker, P. J., and D. C. Roopenian. 2002. "Genetic susceptibility to chronic periodontal disease." *Microbes Infect* 4(11):1157-67.
- Berglundh, T., and M. Donati. 2005. "Aspects of adaptive host response in periodontitis." *J Clin Periodontol* 32 Suppl 6:87-107.
- Bluestone, J. A., C. R. Mackay, J. J. O'Shea, and B. Stockinger. 2009. "The functional plasticity of T cell subsets." *Nat Rev Immunol* 9(11):811-6.
- Bostanci, N., T. Ilgenli, G. Emingil, B. Afacan, B. Han, H. Toz, G. Atilla, F. J. Hughes, and G. N. Belibasakis. 2007. "Gingival crevicular fluid levels of RANKL and OPG in periodontal diseases: implications of their relative ratio." *J Clin Periodontol* 34(5):370-6.
- Cardoso, C. R., G. P. Garlet, G. E. Crippa, A. L. Rosa, W. M. Junior, M. A. Rossi, and J. S. Silva. 2009. "Evidence of the presence of T helper type 17 cells in chronic lesions of human periodontal disease." *Oral Microbiol Immunol* 24(1):1-6.
- Chen, D., M. A. Harris, G. Rossini, C. R. Dunstan, S. L. Dallas, J. Q. Feng, G. R. Mundy, and S. E. Harris. 1997. "Bone morphogenetic protein 2 (BMP-2) enhances BMP-3, BMP-4, and bone cell differentiation marker gene expression during the induction of mineralized bone matrix formation in cultures of fetal rat calvarial osteoblasts." *Calcif Tissue Int* 60(3):283-90.
- Chen, H., W. Yong, S. Ren, W. Shen, Y. He, K. A. Cox, W. Zhu, W. Li, M. Soonpaa, R. M. Payne, D. Franco, L. J. Field, V. Rosen, Y. Wang, and W. Shou. 2006. "Overexpression of bone morphogenetic protein 10 in myocardium disrupts cardiac postnatal hypertrophic growth." *J Biol Chem* 281(37):27481-91.
- Donati, M., B. Liljenberg, N. U. Zitzmann, and T. Berglundh. 2009. "B-1a cells and plasma cells in periodontitis lesions." *J Periodontal Res* 44(5):683-8.

- Dzink, J. L., A. C. Tanner, A. D. Haffajee, and S. S. Socransky. 1985. "Gram negative species associated with active destructive periodontal lesions." *J Clin Periodontol* 12(8):648-59.
- Ebersole, J. L., and D. Cappelli. 1994. "Gingival crevicular fluid antibody to *Actinobacillus actinomycetemcomitans* in periodontal disease." *Oral Microbiol Immunol* 9(6):335-44.
- Ebersole, J. L., M. A. Taubman, D. J. Smith, and D. E. Frey. 1986. "Human immune responses to oral microorganisms: patterns of systemic antibody levels to *Bacteroides* species." *Infect Immun* 51(2):507-13.
- Fine, D. H. 2009. "Of mice and men: animal models of human periodontal disease." *J Clin Periodontol* 36(11):913-4.
- Fine, D. H., H. Schreiner, C. Nasri-Heir, B. Greenberg, S. Jiang, K. Markowitz, and D. Furgang. 2009. "An improved cost-effective, reproducible method for evaluation of bone loss in a rodent model." *J Clin Periodontol* 36(2):106-13.
- Fives-Taylor, P. M., D. H. Meyer, K. P. Mintz, and C. Brissette. 1999. "Virulence factors of *Actinobacillus actinomycetemcomitans*." *Periodontol* 2000 20:136-67.
- Gaffen, S. L., and G. Hajishengallis. 2008. "A new inflammatory cytokine on the block: rethinking periodontal disease and the Th1/Th2 paradigm in the context of Th17 cells and IL-17." *J Dent Res* 87(9):817-28.
- Gamer, L. W., K. Cox, J. M. Carlo, and V. Rosen. 2009. "Overexpression of BMP3 in the developing skeleton alters endochondral bone formation resulting in spontaneous rib fractures." *Dev Dyn* 238(9):2374-81.
- Gemmell, E., C. L. Carter, D. N. Hart, K. E. Drysdale, and G. J. Seymour. 2002. "Antigen-presenting cells in human periodontal disease tissues." *Oral Microbiol Immunol* 17(6):388-93.
- Gemmell, E., K. Yamazaki, and G. J. Seymour. 2007. "The role of T cells in periodontal disease: homeostasis and autoimmunity." *Periodontol* 2000 43:14-40.
- Graves, D. 2008. "Cytokines that promote periodontal tissue destruction." *J Periodontol* 79(8 Suppl):1585-91.
- Graves, D. T., A. J. Delima, R. Assuma, S. Amar, T. Oates, and D. Cochran. 1998. "Interleukin-1 and tumor necrosis factor antagonists inhibit the progression of inflammatory cell infiltration toward alveolar bone in experimental periodontitis." *J Periodontol* 69(12):1419-25.
- Haubek, D., O. K. Ennibi, K. Poulsen, M. Vaeth, S. Poulsen, and M. Kilian. 2008. "Risk of aggressive periodontitis in adolescent carriers of the JP2 clone of *Aggregatibacter* (*Actinobacillus*) *actinomycetemcomitans* in Morocco: a prospective longitudinal cohort study." *Lancet* 371(9608):237-42.
- Herpin, A., C. Lelong, and P. Favrel. 2004. "Transforming growth factor-beta-related proteins: an ancestral and widespread superfamily of cytokines in metazoans." *Dev Comp Immunol* 28(5):461-85.
- Hintzen, R. Q., S. M. Lens, G. Koopman, S. T. Pals, H. Spits, and R. A. van Lier. 1994. "CD70 represents the human ligand for CD27." *Int Immunol* 6(3):477-80.
- Jang, E., S. H. Cho, H. Park, D. J. Paik, J. M. Kim, and J. Youn. 2009. "A positive feedback loop of IL-21 signaling provoked by homeostatic CD4⁺CD25⁻ T cell expansion is essential for the development of arthritis in autoimmune K/BxN mice." *J Immunol* 182(8):4649-56.

- Jee, H. M., K. W. Kim, J. Y. Hong, M. H. Sohn, and K. E. Kim. 2009. "Increased serum B cell-activating factor level in children with atopic dermatitis." *Clin Exp Dermatol*.
- Josefowicz, S. Z., and A. Rudensky. 2009. "Control of regulatory T cell lineage commitment and maintenance." *Immunity* 30(5):616-25.
- Kachlany, S. C. 2010. "Aggregatibacter actinomycetemcomitans leukotoxin: from threat to therapy." *J Dent Res* 89(6):561-70.
- Kachlany, S. C., A. B. Schwartz, N. V. Balashova, C. E. Hioe, M. Tuen, A. Le, M. Kaur, Y. Mei, and J. Rao. 2010. "Anti-leukemia activity of a bacterial toxin with natural specificity for LFA-1 on white blood cells." *Leuk Res* 34(6):777-85.
- Keerthivasan, S., R. Suleiman, R. Lawlor, J. Roderick, T. Bates, L. Minter, J. Anguita, I. Juncadella, B. J. Nickoloff, I. C. Le Poole, L. Miele, and B. A. Osborne. 2011. "Notch signaling regulates mouse and human th17 differentiation." *J Immunol* 187(2):692-701.
- Kempf, T., and K. C. Wollert. 2009. "Growth-differentiation factor-15 in heart failure." *Heart Fail Clin* 5(4):537-47.
- Kiesel, J. R., Z. S. Buchwald, and R. Aurora. 2009. "Cross-presentation by osteoclasts induces FoxP3 in CD8+ T cells." *J Immunol* 182(9):5477-87.
- Kinane, D. F., P. M. Preshaw, and B. G. Loos. 2011. "Host-response: understanding the cellular and molecular mechanisms of host-microbial interactions--consensus of the Seventh European Workshop on Periodontology." *J Clin Periodontol* 38 Suppl 11:44-8.
- Kobayashi, R., T. Kono, B. A. Bolerjack, Y. Fukuyama, R. S. Gilbert, K. Fujihashi, J. Ruby, K. Kataoka, M. Wada, and M. Yamamoto. 2011. "Induction of IL-10-producing CD4+ T-cells in chronic periodontitis." *J Dent Res* 90(5):653-8.
- Korn, T., E. Bettelli, M. Oukka, and V. K. Kuchroo. 2009. "IL-17 and Th17 Cells." *Annu Rev Immunol* 27:485-517.
- Kragstrup, T. W., K. Otkjaer, C. Holm, A. Jorgensen, M. Hokland, L. Iversen, and B. Deleuran. 2008. "The expression of IL-20 and IL-24 and their shared receptors are increased in rheumatoid arthritis and spondyloarthritis." *Cytokine* 41(1):16-23.
- Leibbrandt, A., and J. M. Penninger. 2010. "Novel Functions of RANK(L) Signaling in the Immune System." *Adv Exp Med Biol* 658:77-94.
- Li, Y., C. Messina, M. Bendaoud, D. H. Fine, H. Schreiner, and V. K. Tsiagbe. 2010. "Adaptive immune response in osteoclastic bone resorption induced by orally administered Aggregatibacter actinomycetemcomitans in a rat model of periodontal disease." *Mol Oral Microbiol* 25(4):275-92.
- Mahanonda, R., N. Sa-Ard-Iam, K. Yongvanitchit, M. Wisetchang, I. Ishikawa, T. Nagasawa, D. S. Walsh, and S. Pichyangkul. 2002. "Upregulation of co-stimulatory molecule expression and dendritic cell marker (CD83) on B cells in periodontal disease." *J Periodontal Res* 37(3):177-83.
- McPherron, A. C., A. M. Lawler, and S. J. Lee. 1999. "Regulation of anterior/posterior patterning of the axial skeleton by growth/differentiation factor 11." *Nat Genet* 22(3):260-4.
- Miossec, P., T. Korn, and V. K. Kuchroo. 2009. "Interleukin-17 and type 17 helper T cells." *N Engl J Med* 361(9):888-98.


- Mosmann, T. R., and R. L. Coffman. 1989. "TH1 and TH2 cells: different patterns of lymphokine secretion lead to different functional properties." *Annu Rev Immunol* 7:145-73.
- Nakashima, M., K. Tachibana, K. Iohara, M. Ito, M. Ishikawa, and A. Akamine. 2003. "Induction of reparative dentin formation by ultrasound-mediated gene delivery of growth/differentiation factor 11." *Hum Gene Ther* 14(6):591-7.
- Nigam, P., V. Velu, S. Kannanganat, L. Chennareddi, S. Kwa, M. Siddiqui, and R. R. Amara. 2010. "Expansion of FOXP3+ CD8 T cells with suppressive potential in colorectal mucosa following a pathogenic simian immunodeficiency virus infection correlates with diminished antiviral T cell response and viral control." *J Immunol* 184(4):1690-701.
- Nussbaum, G., and L. Shapira. 2011. "How has neutrophil research improved our understanding of periodontal pathogenesis?" *J Clin Periodontol* 38 Suppl 11:49-59.
- Ohyama, H., N. Kato-Kogoe, A. Kuhara, F. Nishimura, K. Nakasho, K. Yamanegi, N. Yamada, M. Hata, J. Yamane, and N. Terada. 2009. "The involvement of IL-23 and the Th17 pathway in periodontitis." *J Dent Res* 88(7):633-8.
- Oseko, F., T. Yamamoto, Y. Akamatsu, N. Kanamura, Y. Iwakura, J. Imanishi, and M. Kita. 2009. "IL-17 is involved in bone resorption in mouse periapical lesions." *Microbiol Immunol* 53(5):287-94.
- Parfitt, A. M. 1982. "The coupling of bone formation to bone resorption: a critical analysis of the concept and of its relevance to the pathogenesis of osteoporosis." *Metab Bone Dis Relat Res* 4(1):1-6.
- Preshaw, P. M., and J. J. Taylor. 2011. "How has research into cytokine interactions and their role in driving immune responses impacted our understanding of periodontitis?" *J Clin Periodontol* 38 Suppl 11:60-84.
- Rosen, V. 2006. "BMP and BMP inhibitors in bone." *Ann N Y Acad Sci* 1068:19-25.
- Sakaguchi, S. 2005. "Naturally arising Foxp3-expressing CD25+CD4+ regulatory T cells in immunological tolerance to self and non-self." *Nat Immunol* 6(4):345-52.
- Sakai, A., M. Ohshima, N. Sugano, K. Otsuka, and K. Ito. 2006. "Profiling the cytokines in gingival crevicular fluid using a cytokine antibody array." *J Periodontol* 77(5):856-64.
- Sakurai, N., T. Kuroiwa, H. Ikeuchi, N. Hiramatsu, A. Maeshima, Y. Kaneko, K. Hiromura, and Y. Nojima. 2008. "Expression of IL-19 and its receptors in RA: potential role for synovial hyperplasia formation." *Rheumatology (Oxford)* 47(6):815-20.
- Schreiner, H., K. Markowitz, M. Miryalkar, D. Moore, S. Diehl, and D. H. Fine. 2011. "Aggregatibacter actinomycetemcomitans-induced bone loss and antibody response in three rat strains." *J Periodontol* 82(1):142-50.
- Schulte, S., G. K. Sukhova, and P. Libby. 2008. "Genetically programmed biases in Th1 and Th2 immune responses modulate atherogenesis." *Am J Pathol* 172(6):1500-8.
- Seymour, G. J., and E. Gemmell. 2001. "Cytokines in periodontal disease: where to from here?" *Acta Odontol Scand* 59(3):167-73.
- Shea, P. R., K. Virtaneva, J. J. Kupko, 3rd, S. F. Porcella, W. T. Barry, F. A. Wright, S. D. Kobayashi, A. Carmody, R. M. Ireland, D. E. Sturdevant, S. M. Ricklefs, I. Babar, C. A. Johnson, M. R. Graham, D. J. Gardner, J. R. Bailey, M. J. Parnell, F. R. Deleo, and J. M. Musser. 2010. "Interactome analysis of longitudinal pharyngeal infection of cynomolgus macaques by group A Streptococcus." *Proc Natl Acad Sci U S A* 107(10):4693-8.

- Shenker, B. J., M. Dlakic, L. P. Walker, D. Besack, E. Jaffe, E. LaBelle, and K. Boesze-Battaglia. 2007. "A novel mode of action for a microbial-derived immunotoxin: the cytolethal distending toxin subunit B exhibits phosphatidylinositol 3,4,5-triphosphate phosphatase activity." *J Immunol* 178(8):5099-108.
- Taubman, M. A., P. Valverde, X. Han, and T. Kawai. 2005. "Immune response: the key to bone resorption in periodontal disease." *J Periodontol* 76(11 Suppl):2033-41.
- Tsuji, K., A. Bandyopadhyay, B. D. Harfe, K. Cox, S. Kakar, L. Gerstenfeld, T. Einhorn, C. J. Tabin, and V. Rosen. 2006. "BMP2 activity, although dispensable for bone formation, is required for the initiation of fracture healing." *Nat Genet* 38(12):1424-9.
- Vanhara, P., E. Lincova, A. Kozubik, P. Jurdic, K. Soucek, and J. Smarda. 2009. "Growth/differentiation factor-15 inhibits differentiation into osteoclasts--a novel factor involved in control of osteoclast differentiation." *Differentiation* 78(4):213-22.
- Xiong, H., L. Wei, and B. Peng. 2009. "Immunohistochemical localization of IL-17 in induced rat periapical lesions." *J Endod* 35(2):216-20.
- Xue, H., B. Lu, J. Zhang, M. Wu, Q. Huang, Q. Wu, H. Sheng, D. Wu, J. Hu, and M. Lai. "Identification of serum biomarkers for colorectal cancer metastasis using a differential secretome approach." *J Proteome Res* 9(1):545-55.
- Yamazaki, K., T. Nakajima, E. Gemmell, B. Polak, G. J. Seymour, and K. Hara. 1994. "IL-4- and IL-6-producing cells in human periodontal disease tissue." *J Oral Pathol Med* 23(8):347-53.
- Yamazaki, K., T. Nakajima, Y. Kubota, E. Gemmell, G. J. Seymour, and K. Hara. 1997. "Cytokine messenger RNA expression in chronic inflammatory periodontal disease." *Oral Microbiol Immunol* 12(5):281-7.
- Ye, L., H. Kynaston, and W. G. Jiang. 2009. "Bone morphogenetic protein-10 suppresses the growth and aggressiveness of prostate cancer cells through a Smad independent pathway." *J Urol* 181(6):2749-59.
- Yokoe, T., T. Ohmachi, H. Inoue, K. Mimori, F. Tanaka, M. Kusunoki, and M. Mori. 2007. "Clinical significance of growth differentiation factor 11 in colorectal cancer." *Int J Oncol* 31(5):1097-101.
- Yue, G., J. B. Kaplan, D. Furgang, K. G. Mansfield, and D. H. Fine. 2007. "A second *Aggregatibacter actinomycetemcomitans* autotransporter adhesin exhibits specificity for buccal epithelial cells in humans and Old World primates." *Infect Immun* 75(9):4440-8.
- Yuryev, A., Z. Mulyukov, E. Kotelnikova, S. Maslov, S. Egorov, A. Nikitin, N. Daraselia, and I. Mazo. 2006. "Automatic pathway building in biological association networks." *BMC Bioinformatics* 7:171.
- Zaiss, M. M., M. Kurowska-Stolarska, C. Bohm, R. Gary, C. Scholtysek, B. Stolarski, J. Reilly, S. Kerr, N. L. Millar, T. Kamradt, I. B. McInnes, P. G. Fallon, J. P. David, F. Y. Liew, and G. Schett. 2011. "IL-33 shifts the balance from osteoclast to alternatively activated macrophage differentiation and protects from TNF-alpha-mediated bone loss." *J Immunol* 186(11):6097-105.
- Zambon, J. J. 1985. "Actinobacillus actinomycetemcomitans in human periodontal disease." *J Clin Periodontol* 12(1):1-20.
- Zambon, J. J., T. Umemoto, E. De Nardin, F. Nakazawa, L. A. Christersson, and R. J. Genco. 1988. "Actinobacillus actinomycetemcomitans in the pathogenesis of human periodontal disease." *Adv Dent Res* 2(2):269-74.

Zhang, F., H. Tanaka, T. Kawato, S. Kitami, K. Nakai, M. Motohashi, N. Suzuki, C. L. Wang, K. Ochiai, K. Isokawa, and M. Maeno. 2011. "Interleukin-17A induces cathepsin K and MMP-9 expression in osteoclasts via celecoxib-blocked prostaglandin E2 in osteoblasts." *Biochimie* 93(2):296-305.

IntechOpen

IntechOpen



Periodontal Diseases - A Clinician's Guide

Edited by Dr. Jane Manakil

ISBN 978-953-307-818-2

Hard cover, 368 pages

Publisher InTech

Published online 03, February, 2012

Published in print edition February, 2012

"Periodontal diseases" is a web-based resource intended to reach the contemporary practitioners as well as educators and students in the field of periodontology. It is fully searchable and designed to enhance the learning experience. Within the book a description is presented of the current concepts presenting the complex interactions of microbial fingerprint, multiple genotypes, and host modulations. In addition, an overview is given of the clinical outcome of the disease's progression, as influenced by the epigenetic factors. Emerging concepts on periodontitis as a risk factor for various systemic diseases and as a bilateral modulating factor have been elucidated in detail as well.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Vincent K. Tsiagbe and Daniel H. Fine (2012). The Impact of Bacteria-Induced Adaptive Immune Responses in Periodontal Disease, Periodontal Diseases - A Clinician's Guide, Dr. Jane Manakil (Ed.), ISBN: 978-953-307-818-2, InTech, Available from: <http://www.intechopen.com/books/periodontal-diseases-a-clinician-s-guide/the-impact-of-bacteria-induced-adaptive-immune-responses-in-periodontal-disease>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen