

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Diagnostic Methods for *Mycobacterium tuberculosis* and Challenges in Its Detection in India

Shamsher S. Kanwar

Department of Biotechnology
Himachal Pradesh University, Summer Hill, Shimla
India

1. Introduction

Tuberculosis (TB) is one of the world's oldest and most important disseminating infectious diseases that still accounts for a high morbidity and mortality among adults. Despite high prevalence, case detection rates are low, posing major hurdles for TB control in developed and developing countries. Traditional diagnosis of TB bacilli depends upon smear positivity in sputum samples, culture and chest radiography. All these tests have known limitations. Conventional tests for detection of drug resistance are slow, tedious and difficult to perform in field conditions. For rapid diagnosis, new methods include newer versions of nucleic acid amplification tests, immune-based assays, skin patch test and rapid culture systems. For drug resistance analysis line-probe assays, bacteriophage-based assays, molecular beacons and microscopic observation drug susceptibility assay are available. An ideal test for TB is still not available and fast emergence of drug resistant tubercle strains aided by the ever-increasing HIV AIDS-epidemic in third-world countries has stressed the need of rapid diagnostic test(s) to show the presence of mycobacteria in the clinical samples. Microscopy and culture are still the major backbone for laboratory diagnosis of tuberculosis; new methods including molecular diagnostic tests have evolved over a period of time. The majority of molecular tests have been focused on: (i) detection of nucleic acids both DNA and RNA, which are specific to *Mycobacterium tuberculosis*, by amplification techniques such as polymerase chain reaction (PCR) focusing on detection and molecular epidemiology of *M. tuberculosis*; and (ii) detection of mutations in the genes which are associated with resistance to anti-tuberculosis drugs by sequencing or nucleic acid hybridization. The development and use of rapid diagnostic tools become increasingly important in addressing the emergence and treatment of multi-drug resistant (MDR) and extreme-drug (XDR) resistant *M. tuberculosis* strains.

Tuberculosis remains one of the most challenging bacterial diseases in spite of development of a realm of antibiotics and diagnostic molecular biology techniques. The tubercle bacillus was discovered more than two hundred years ago and substantial advancements have been made in our knowledge about the development of tuberculosis in human. The organism seems to evolve over a period of time in terms of its ability to survive the action of front line

anti mycobacterial antibiotics by developing appropriate antibiotic resistant mechanisms. The estimated mortality by World Health Organization reports over 1.7 million deaths in 2006 and 9 million new cases of tuberculosis [WHO 2008]. The economic burden of management of disease in patients in the prime of their age is enormous because of prolonged antibiotic treatment. In spite of availability of anti mycobacterial drugs, tuberculosis remains one of the major health problems facing mankind particularly in developing countries. Presently, about one third of world's population is infected with *Mycobacterium tuberculosis*. Currently, the number of people dying of tuberculosis is more than any other infectious diseases. Death from tuberculosis comprises 25% of all avoidable deaths in developing countries [Ramachandran and Parmasivan 2003]. Nearly 95% of all tuberculosis cases and 98% of deaths due to tuberculosis are in developing countries and 75% of tuberculosis cases are in the economically productive age group. Currently, more people die of tuberculosis than from any other infectious disease. In India, out of a total population of more than 1 billion, approximately 2 million develop active disease and up to half a million die of tuberculosis. It also imparts a financial burden on the economy in terms of out-put losses because of premature deaths and ill health. To add to the existing cost burden, the cumulative effect is seen because of ever increasing number of new TB cases associated with HIV patients and about 1.8 million of these are co-infected with TB [Ramachandran and Paramasivan 2003].

2. Traditional methods of tuberculosis detection, management and limitations

Robert Koch discovered the tubercle bacillus in 1882, and there after methods of staining these microorganisms were developed to assist the diagnosis of the disease. Early diagnosis of the tuberculosis in the patients is a challenging task especially in the pauci-bacillary and extra-pulmonary forms. The conventional methods that are still the mainstay of the diagnosis of TB like Tuberculin test/ Mantoux test, radiological examination and other imaging methods and sputum smear microscopy have their own limitations. Sputum smear microscopy requires 10,000 to 1,00,000 organisms/ ml and acid fast bacilli could be any pathogenic or saprophytic mycobacteria. Although smear microscopy may be made more convenient by using various fluorochromes (auramine, rhodamine, FITC etc.) but the scarce presence of tuber bacilli in the sputum has its own disadvantage. The smear positivity has to be supplemented with the culture positivity that has its own limitations because of failure of bacilli to grow or often become contaminated with other microbes. The slow growth of the tubercle bacilli on medium lingers on the confirmation of the causative organism. Histopathology is characteristic but there could be problems to get representative specimen, and non-specific features. Immunoassay based approaches are doubtful as the antibodies and the antigens may persist for some time after control of the clinical or sub-clinical disease. Thus Acid Fast Bacilli (AFB) staining of clinical material followed by smear microscopy remains the most cost effective, frequently used microbiological test for detection of TB. The major drawback of sputum smear microscopy is its poor sensitivity, especially to be ~70% in a recent review [Steingart et al., 2006]. Although the AFB staining is easy to perform in the field settings especially in the poor third world countries as well in the developing countries but the sensitivity of sputum smear microscopy is clearly less in many settings and may be sometimes as low as ~35% in some situations with high rates of TB and HIV co-infection [Khatri and Frieden 2002]. Compounding the poor test sensitivity is

in adequate or absent test quality assurance in some resource-constrained settings, further cut down the overall yield of the microscopy, driving up the laboratory workload as more sputum tests per patient are performed in an effort to reach a diagnosis, and increasing delay in diagnosis and patient's compliance to repeated follow-up [Dorman 2010]. Moreover drug-susceptibility status cannot be determined from the smear microscopy.

Unfortunately, the world's largest democracy India has over 1.2 billion people and this overpopulated country also has the highest burden of tuberculosis in the world. India accounts for about 20% incidence of tuberculosis besides a high incidence of global occurrence of multi drug resistant (MDR) tuberculosis. The poor sanitation conditions, thickly populated urban and rural area, scanty medical services in villages and rural area, malnourishment, insensitivity of private sector towards quick diagnosis, treatment and management of TB positive patients and higher cost of non-standard methods of diagnosis of TB are some of the important reasons of concern. It is obvious that any global effort towards control and eradication of TB and fast emerging MDR-TB is invariably dependent upon success of concerted efforts to contain the spread of this contagious disease. In India, the National TB Programme (NTP) was initiated in 1962. However, the poor infrastructure, inadequate funding, administrative lack-luster approach, irregular drug supply, non-standard and multiple anti-tuberculosis drug therapy, irregularity or non-compliance of patients to the clinician and a long treatment period had little effect on containing the spread of TB and controlling the emerging MDR-TB strains. The management and control of TB was further compounded and complicated with low rates of TB case detection, compliance of treatment regimen (30%), high rate of default (40-60%) and continuing high mortality (1: 2000) the NTP programme had little success rate. To overcome the deficiencies of NTP a Revised National Tuberculosis Control Programme (RNTCP) was launched by the Government of India in 1997, based on the global DOTS (Directly Observed Treatment, Short Course) approach that was used to exert an epidemiological impact by achieving 70% case detection and 85% cure rate. It was an encouraging sign that by 2002, 100% of the Indian population was covered by the India's own drug-delivery model - the DOTS programme, making this extended coverage as India's most significant public health accomplishment. The RNTCP thus achieved a pronounced success in cure rates (>80% in new infectious cases), substantial decline in mortality with low rate (<10%) of default [Khatri and Frieden 2002; TB India 2009; Bhargava et al., 2011].

3. Tuberculosis and HIV epidemic

In spite of improvement in public health system, participation of private sector in TB detection and management still the sputum smear microscopy test is most commonly used in public health settings to detect pulmonary TB. This method has roughly 50% chance of detection. Unfortunately, a significant number of people outside the public health sector, where the most common test is the serological (various types of ELISA for detection of *M. tuberculosis* antigens; or anti-*M. tuberculosis* IgG or IgM antibodies) that are expensive and means nothing. The global impact of converging dual epidemics of tuberculosis and human immunodeficiency virus (HIV) is one of the major challenges of the present time. In India, there are 2.5 million people living with HIV and AIDS at the end of 2007 while the incidence of TB was ~1.8 million cases per year [WHO 2008, WHO Global Tuberculosis Program 1992].

In a survey carried out among new TB patients by RNTCP in 2007, HIV sero-prevalence varied widely and ranged from 1-13.8% across the 15 districts [Swaminathan and Narendran 2008]. Pulmonary involvement occurs in about 75% of all HIV-infected patients with TB [Devivanayagam et al., 2001; Ahmad and Shameen 2005]. Moreover, the interaction between HIV and TB in persons co-infected with HIV and TB is bi-directional and synergistic. As HIV progresses, there is cutaneous anergy as well as impaired tissue containment of mycobacteria leading to widespread dissemination of mycobacteria. While TB can develop at any CD4 count, extra-pulmonary and disseminated forms of the diseases are more common as immunodeficiency increases. Thus HIV infection is associated not only with an increased incidence of TB but also with altered clinical manifestations especially in the advanced stages of the disease. The cost management of anti-HIV and anti-TB therapy in the patients becomes a daunting task that compromises the success rate of containment and spread of TB from HIV infected patients. Current guidelines recommend that irrespective of HIV status, TB management require a minimum of 6 months of treatment with four drugs (including rifampin) in the intensive phase and two drugs in the continuation phase. In India, under the RNTCP, patients with newly diagnosed TB receive a 6-month thrice-weekly regimen 9cat I – 2EHRZ₃/4RH₃) while those with relapse, default or failure receive an 8-months regimen (cat II – 2SEHRZ₃/1SEHRZ₃/5EHR₃). The lifetime risk of TB in immunocompetent persons is 5% to 10%, but HIV positive individuals; there is a 5% to 15% annual risk of developing active TB diseases [Swaminathan et al., 2000]. WHO estimated 9.2 million new cases of TB globally in 2006 (139 per 100,000); of whom 7,09,000 (7.7%) were HIV positive (WHO 2008). India, China, Indonesia, South Africa and Nigeria rank 1st to 5th in terms of incident TB cases.

The first and foremost step in the diagnosis of TB is its accurate and early detection. To achieve this objective a number of methods have been developed and reported (Table 1) that achieve early growth of *M. tuberculosis* [Katoch and Sharma 1997; Katoch 2004].

4. Anti-TB drug resistance

The overall pattern of drug resistance to first line anti TB therapy is similar in HIV positive and negative patients; however, MDR-TB is marginally higher (3-4%) in HIV positive patients with newly diagnosed tuberculosis status [Swaminathan 2005]. Rifampicin mono-resistance is more common in HIV infected patients and arises independently from mutations in drug susceptible strains. Treatment of MDR-TB should employ at least 3-4 new drugs. The regimen should include an aminoglycoside and be given under direct observation. Extensively drug resistant (XDR) TB strains have emerged and have been reported from India [Singh et al., 2007; Thomas et al., 2007]. Such strains appeared to be as an outcome of the mismanagement of TB. It seems that XDR-TB is practically untreatable and thus an attempt may be made to limit its spread by strengthening the TB control programme. Presently, there is no national policy regarding TB preventive therapy for HIV positive patients in India [Swaminathan and Narendran 2008]. A clinical trial conducted at the Tuberculosis Research Center, Chennai investigated two different regimens; a 6-month regimen of ethambutol and isoniazid vs. a 3-year regimen of isoniazid alone, in order to establish ideal duration of therapy. In a TB-endemic country like India, consideration shall be given to provide preventive therapy to HIV-infected persons. Line probe assays, a family

S. No.	Method	Concept	Reference(s)
1	BACTEC system	Generation and detection of radioactive CO ₂ from substrate palmitic acid. Used world-over, detection of growth in 5-7 d. Inclusion of (NAP: beta nitro alpha acetylamine beta hydroxy propiophenone) distinguishes <i>M. tuberculosis</i> [inhibition] from other mycobacteria.	Venkataram et al., 1998; Bemer et al., 2002
2	Mycobacteria growth indicator tube (MGIT)	Developed by Becton Dickinson, growth detection by non-radioactive fluorochrome detection; useful in drug screening, early detection of mycobacterium growth in 7-12 d.	Bemer et al., 2002; Tortoli et al., 1999
3	MB/BacT system	Developed by Organon Technika; colorimetric detection of bacterial growth in, cultures; also useful for drug susceptibility testing	Brunello and Fontana 2000
4	TK Medium	Developed by Salubris, Inc., MA, USA is a novel colorimetric system that indicates growth of mycobacteria by changing its color, also discriminates between mycobacteria and contamination, and enables drug susceptibility testing. Test is low cost and simple. Sensitivity of TK medium is comparable to the LJ-medium.	Kocagoz et al., 2004; Salubris, Inc.
5	Septi-Check	Bi-phase system developed by Roche. Consists of enriched selective broth and a slide having non-selective Middlebrook agar on one side and two sections on other side: one with NAP + egg-containing agar, and second with chocolate agar for detection of contaminating microbes.	Isenberg et al., 1991

S. No.	Method	Concept	Reference(s)
6	Reporter phages/ Bronx box	Use of mycobacterium-specific phage(s) and a reporter gene (luciferase) for detection of growth and drug-susceptibility to anti-TB drugs. Viability detection by either emission of light from microbe due to activation of luciferase gene or production of plaque on an indicator strain of mycobacteria; results availability in 2 d.	Riska et al., 1999; Wilson et al., 1997; Krishnamurthy et al., 2002
7	E-test	Use of gradient of drug on a paper-strip; useful for drug susceptibility testing of <i>M. tuberculosis</i> .	Kirk et al., 1998
8	Flow cytometry	Use of FACS for drug susceptibility testing, high cost of equipment and trained operator are the drawbacks.	Kakkar et al., 2000
9	Line-probe assay	A novel DNA strip-based test that uses PCR and reverse hybridization methods for rapid detection of mutations associated with drug resistance. Designed to identify <i>M. tuberculosis</i> complex and simultaneous detection of mutations associated with drug resistance.	Morgan et al., 2005

Table 1. Methods of early detection of *M. tuberculosis*.

of novel DNA strip-based tests use PCR and reverse hybridization methods for the rapid detection of mutations associated with drug resistance. These kits [INNO-LiPA Rif TB kit, Innogenetics NV, Gent, Belgium; GenoType MTBDR assay; Hain Life-science GmbH, Nehren, Germany] are not currently FDA approved for use in USA. Line-probe assays are designed to identify *M. tuberculosis* complex and simultaneously detect mutations associated with drug resistance. In June 2008, WHO announced a new policy statement endorsing the use of line probe assays for rapid screening of patients at risk of MRD-TB (<http://www.who.int/tb/en/>). However, the line probe assays are not recommended as a complete replacement for conventional culture and drug susceptibility testing. Culture is

still required for smear-negative specimens and conventional drug susceptibility testing is still necessary to confirm XDR-TB.

5. Ineffective TB diagnostics in India

Ineffective TB diagnostics are a lucrative market in India. Patients seeking TB care in the private medical institutes are commonly subjected to diagnostic tests i.e. the anti-body-based blood tests, including ELISA that are completely ineffective at detecting TB. This is because a large number of the world's population has anti-TB antibodies, though only 10% of them do develop the active form of the disease. Obviously if patients who do not have TB are misdiagnosed, they could undergo 6-months of nasty toxic anti-TB chemotherapy. If patients have active TB and the test misses it, the disease may worsen and they may continue to spread the disease in their community. According to a preliminary analysis of over 80 labs in India, it is estimated that patients undergo more than 1.5 million useless TB antibody tests each year (WHO recommends against inaccurate tuberculosis tests by Kelly Morris; www.thelancet.com vol 377 Jan 8, 2011). The absence of any regulatory mechanisms results in the import of these dubious diagnostics from France, UK, USA and other countries, where these tests are not approved for TB diagnosis. These tests generate at least US \$ 15 million. In a country that has ~100,000 labs this is probably a fraction of the enormous total market. Accurate diagnose is critical to the control of tuberculosis in India, particularly in view of the fact that India has set new targets as a part of RNTCP, which includes early detection of 90% of all TB cases by year 2015.

6. Molecular diagnosis of TB in Indian context

In India various institutes working on TB have sufficient technical expertise and financial affordability to use molecular diagnostic methods for detection of tubercle bacilli in the samples. For a laboratory with good sample load and which is using rapid methods for early growth detection the additional cost of sample analysis shall not be significant. If 10-25 isolates/ growths are assessed for identity simultaneously, additional cost for each isolate using a non-radioactive detection system like digoxigenin (DIG) should not be more than 200-250 rupees (~US \$ 4-5). Similarly for a PCR system using primers, which are not patented, cost should be in similar range as prices of primers and reagents have considerably been reduced during the last couple of years [Katoch 2004]. The nucleic acid amplification tests (NAATs) are designed to amplify nucleic acid regions specific to the *Mycobacterium tuberculosis* complex. Such tests may be used directly on clinical samples/ sputum samples. Nucleic acid amplification test (NAAT) commercial kits are available under various brands like Amplified *M. tuberculosis* Direct Test (MTD; Gen-Probe Inc., CA, USA), the Amplicor *M. tuberculosis* (MTB) tests (Roche Molecular Diagnostics, CA, USA and the BD ProbeTec ET assay (BD Biosciences, MD, USA. Besides in-house lab developed PCR assays vary widely in their protocols and vary from lab to lab. In-house NAAT are cheap and are often used in research in developing countries where commercial NAATs are quite expensive to test large number of samples and thus in-house PCR technique/ protocol(s) provide a cheap option (Pai 2004). However, all the detection methods including the conventional AFB-staining, skin tuberculin test and new generation NAAT tests have some advantages as well as limitations (Table 2).

Method	Use	Intended use	Advantages	Limitations
AFB smear microscopy	Rapid tubercle bacilli detection	Community	Needs moderate training, microscope and low investment	Low sensitivity
Culture on solid media	Mycobacterial growth and drug susceptibility assay	Referral lab	Good sensitivity; gold standard	Long time to detect growth of bacteria
Chest radiography	Pulmonary TB detection	Referral by clinician	Indications and use not restricted to TB	Low specificity & sensitivity, trained clinician needed
Tuberculin skin test	Detection of <i>M. tuberculosis</i>	Community	Extensive clinical and published experience	Sensitivity decreases in immuno-compromised persons, positive reaction in BCG vaccinees
γ -Interferon release assay	Detection of <i>M. tuberculosis</i> infection	Referral to reference lab	Highly specific for <i>M. tuberculosis</i>	Trained manpower, poor sensitivity especially in immunocompromised hosts
Automated, non-integrated NAAT	Pulmonary TB detection	Reference lab	High sensitivity, rapidity and detection of mutations in MDR-TB strains.	Moderately trained personnel and equipment, laborious and possible cross-contamination among specimens
Culture in liquid media [MGIT; BacT/ Alert and others]	TB detection and as a prerequisite to drug-susceptibility testing	Referral lab	High sensitivity (more sensitivity than liquid media)	Long time for detection; less than solid medium but high contamination rate in some settings

Method	Use	Intended use	Advantages	Limitations
Line probe assay	TB detection and drug susceptibility testing	Reference lab	Poor sensitivity in smear-negative samples, short analysis time	Labor intensive, potential for cross-contamination, requires extensive training
Strip-based <i>Mycobacterium</i> species identification	Species identification i.e. TB versus non-TB in cultures positive for mycobacterial growth	Referral lab	Accurate, requires minimal-training/ equipment/ consumables	Moderate training in handling of pathogenic microbes

Adapted from references: Pai et al., 2006; Perkins and Cunningham 2007; Nyendak et al., 2009; WHO and Stop TB Partnership 2009; Dorman 2010.

Table 2. Tuberculosis diagnostic methods in use, recently endorsed by WHO and under development.

7. Challenges in the TB care and control

At present a vigorous approach is needed to proactively detect the TB cases under RNTCP. The DOT service providers may be actively involved in identifying fresh potential TB patients in their community and getting them diagnosed for TB. Another possibility is contact tracing of both adults and children diagnosed to have TB. Such approach is currently being followed in HIV programs and could be considered to improve upon the case detection rate. Further prevalence studies in different parts of India may be conducted systematically to determine the existing burden of TB. The district level data will be quite helpful in achieving a realistic figure of TB cases.

Lab strengthening: In most cases among the poorest strata of people living in slums or rural areas the sputum transportation is difficult to reach populations is a major consideration and TB is quite commonly said to be poor man’s disease. Thus inadequate number of microscopic centers/ labs put the burden on existing microscopic centers that causes a delay in the reporting of the results of the sputum samples. Most labs conduct AFB testing and are ill equipped to perform culture of tubercle bacilli.

Migrant populations: Presently, there is no national level strategy and guidelines for tuberculosis care and control for the migrant population in India who move from one state or place to another one as a part of their jobs or in search of jobs. Millions of migrants are currently working in unorganized job sectors with no health facilities, insurance facilities and work place policy for disease care and control like TB, HIV etc. Such workers are solely dependent on the relatively expensive private health sector for their healthcare. Accessing those migrants at their residences and working places with the key messages of TB, DOTS

and RNTCP is extremely challenging because of the geographically scattered areas and huge number of migrants. Also women engaged in unorganized job sectors are particularly prone to tuberculosis due to continuous exploitation by the employers. The migrant workers should be mapped in the urban and peri-urban areas (construction sites, street dwellers, illegal residents along the railway tracts, brick kilns etc.) and should be provided RNTCP services (like sensitization on TB, identification of suspected TB cases, referral and tracking) through community-based programs as part of the extended TB monitoring and care program and activities. Moreover, the TB component may be introduced into the existing HIV programs for migrant workers after collaboration with National AIDS Control Programs (NACO).

New TB testing tools: For 100% detection of the TB cases new technologies and techniques shall be established that are reasonably cheap, rapid and easily available. Recently a new PCR based diagnostic kit has been developed through a partnership between Cepheid and Foundation for Innovative New diagnostics, the University of Medicine and Dentistry of New Jersey, the Bill and Melinda Gates Foundation and national Institute of Health (U.S.). The study demonstrated high sensitivity and specificity, identifying 98% of patients with TB and correctly identified 98% of bacteria resistant to rifampin.

Besides PCR, some novel tests involving use of beacons for the rapid detection of mutations associated with drug resistance have been reported [Varma-Basil et al., 2004]. Employing Xpert/RIF kit 1,700 patients were screened at five sites across the world including Mumbai, and using this PCR 98% of patients with TB and resistant to rifampin were correctly identified. This PCR needed about 100 minutes compared to current tests that may take up to 3 months to have results. Unfortunately, the PCR NAATs are performed in a few national labs and specialized private hospitals only. Facilities for culture and drug susceptibility testing of TB cultures in India are grossly inadequate [Bhargava 2011]. As of 2008, only 17 accredited facilities were doing culture and drug-susceptibility testing (~0.1 facility/ 10 million residents against 1/ 10 million). Efforts are underway to enhance the number of labs/ facilities to 43 to perform drug-susceptibility testing. The RNTCP has started to include rapid diagnostic methods to perform culture and drug-susceptibility testing so as to prevent delay in management of MDR-TB patients.

8. Conclusion

TB one of the most communicable diseases is still evading accurate diagnosis because of lack and development of cheap, less labor/ equipment intensive, highly specific and sensitive methods. Culture positivity in sputum positive samples is still considered to be a gold standard as this method provides a further lead in accessing the drug susceptibility of the grown mycobacterial cultures. Currently, most of the tools/ techniques in demonstration or late-stage validation are sputum based and thus are likely to result in incremental gains in rate of TB detection. Still there is an urgent need to develop and validate a mycobacterial culture based or NAAT-based technique that is close to 100% specificity and sensitivity. The need to fast develop such techniques is urgent because of development of MDR mycobacterial strains in third-world countries as these countries are also experiencing an increased burden of HIV-positive patients. The previous decade has shown how despite '100% coverage' and impressive case detection and cure rates, TB continues to be an

epidemic of magnanimous magnitude in India. Thus in India a staunch collaboration between RNTCP, NACO and private partners is the need of the hour to contain the fast spread of MRD *M. tuberculosis* strains.

9. References

- Ahmad Z and Shameem M (2005). Manifestations of tuberculosis in HIV infected patients. *JIACM* 6: 302-305.
- Bemer P, Palicova F, Rusch-Gerdes S, Dugeon HB and Pfyffer GE (2002). Multicentre evaluation of fully automatic BACTEC mycobacteria growth indicator tube 960 system for susceptibility testing of *Mycobacterium tuberculosis*. *J Clin Microbiol* 40: 150-154.
- Bhargava A, Pinto I and Pai M (2011). Mismanagement of tuberculosis in India: causes, consequences and the way forward. *Hypothesis* 9: 1-13.
- Brunello F and Fontana R (2000). Reliability of the MB-BactT system for testing susceptibility of *Mycobacterium tuberculosis* complex strains to anti-tuberculous drugs. *J Clin Microbiol* 38: 872-873.
- Deivanayagam CN, Rajasekaran S, Senthilnathan V, Krishnarajasekhar R, Raja K, Chandrasekar C, Palanisamy S, Samuel DA, Jothivel G and Elango SV (2001). Clinico-radiological spectrum of tuberculosis among HIV-sero-positives - a Tambaram study. *Indian J Tuberc* 49: 123-127.
- Dorman SE (2010). New diagnostic tests for tuberculosis: Bench, bedside and beyond. *Clin Infect Dis* 50: S173-S177.
- Isenberg HD, D'Amate RF, Heifets L, Murray PR, Scardamaglia M, Jacob MC et al., (1991). Collaborative feasibility study of a biphasic system (Roche Septi-Chek AFB) for rapid detection and isolation of mycobacteria. *J Clin Microbiol* 29: 1713-1722.
- Kakkar N, Sharma M, Ray P, Seth S and Kumar S (2000). Evaluation of E-test for susceptibility testing of mycobacteria to primary antitubercular drugs. *Indian J Med Res* 111: 168-171.
- Katoch VM (2004). Newer diagnostic techniques for tuberculosis. *Indian J Med Res* 120: 418-428.
- Katoch VM and Sharma VD (1997). Advances in the diagnosis of mycobacterial diseases. *Indian J Med Microbiol* 15: 49-55.
- Khatri GR and Frieden T (2002). Controlling tuberculosis in India. *N Engl J Med* 347: 1420-1425.
- Kirk SM, Schell RF, Moore AV, Callister SM and Mazurek GH (1998). Flow cytometric testing of susceptibilities of *Mycobacterium tuberculosis* isolates to ethambutol, isoniazid and rifampicin in 24 hours. *J Clin Microbiol* 36: 1568-1571.
- Kocagoz T, O'Brien R and Perkins M (2004). A new colorimetric culture system for the diagnosis of tuberculosis. *Int J Tuberc Lung Dis* 8: 1512-1513.
- Krishnamurthy A, Rodrigues C and Mehta AP (2002). Rapid detection of rifampicin resistance in *Mycobacterium tuberculosis* by phage assay. *Indian J Med Microbiol* 20: 211-214.
- Morgan M, Kalantri S, Flores L and Pai M (2005). A commercial line probe assay for the rapid detection of rifampicin resistance in *Mycobacterium tuberculosis*. A systemic review and meta-analysis. *BMC Infect Dis* 5: 62.
- Nyendak MR, Lewinsohn DA and Lewinsohn DM (2009). New diagnostic methods for tuberculosis. *Curr Opin Infect Dis* 22: 174-182.

- Pai M (2004). The accuracy and reliability of nucleic acid amplification tests in the diagnosis of tuberculosis. *Natl Med J India* 17: 233-236.
- Pai M, Kalantri S and Dheda K (2006). New tools and emerging technologies for the diagnosis of tuberculosis: Part II. Active tuberculosis and drug resistance. *Expert Rev Mol Diagn* 6: 423-432.
- Perkin MD and Cunningham J (2007). Facing the crisis: Improving the diagnosis of tuberculosis in the HIV era. *J Infect Dis* 196(Suppl 1): S15-S27.
- Ramachandran R and Parmasivan CN (2003). What is new in the diagnosis of tuberculosis? Part I: Techniques for diagnosis of tuberculosis. *Indian J Tuberculosis* 50: 133-141.
- Riska PF, Su Y, Bardarov S, Freundlich L, Sarkis G, Hatfull G et al., (1999). Rapid film-based determination of antibiotic susceptibility of *Mycobacterium tuberculosis* strains by using a luciferase reporter phage and the Bronx box. *J Clin Microbiol* 37: 1144-1149.
- Singh S, Sankar MM and Gopinath K (2007). High rate of extensively drug-resistant tuberculosis in Indian AIDS patients. *AIDS* 21: 2345-2347.
- Steingart KR, Henry M, Ng V et al., (2006). Fluorescence versus conventional smear microscopy for tuberculosis: A systemic review. *Lancet Infect Dis* 6: 570-581.
- Swaminathan S and Narendran G (2008). HIV and tuberculosis in India. *J Biosci* 33: 527-537.
- Swaminathan S, Paramasivan CN, Ponnuraja C, Iliayas S, Rajasekaran S and naryanan PR (2005). Anti-tuberculosis drug resistance tuberculosis in South India. *Int J Tuberc Lung Dis* 9: 896-900.
- Swaminathan S, Ramachandran R, Baskaran G and Paramasivan CN (2000). Risk of development of tuberculosis in HIV infected patients. *Int J Tuberc Lung Dis* 4: 839-844.
- TB India (2009). RNTCP status report.
- Thomas BE, Ramachandran R, Anitha S and Swaminathan S (2007). Feasibility of routine HIV testing among TB patients through a voluntary, counseling and testing center (VCTC). *Int J Tuberc Lung Dis* 11: 1296-1301.
- Tortoli E, Cichero P, Piersimoni C, Simonetti T, Gesu G and Nistta D (1999). Use of BACTEC MGIT for recovery of mycobacteria from clinical specimens: multicentric study. *J Clin Microbiol* 37: 3578-3582.
- Varma-Basil M, El-Hajj H, Colangeli R et al., (2004). Rapid detection of rifampin resistance in *Mycobacterium tuberculosis* isolates from India and Mexico by a molecular beacon assay. *J Clin Microbiol* 42: 5512-5516.
- Venkataraman P, Herbert D and Paramasivan CN (1998). Evaluation of BACTEC radiometric method in the early diagnosis of tuberculosis. *Indian J Med Res* 108: 1201-127.
- WHO (2008). Global tuberculosis control: surveillance, planning, financing. WHO report 2008. WHO/HTM/TB/2008.393. Geneva.
- WHO Global Tuberculosis Program (1992). Tuberculosis programme review, India, September 1992.
- Wilson S, Al-Suwaidi A, McNerny R, Porter J and Drobniewski F (1997). Evaluation of a new rapid bacteriophage based method for the drug susceptibility testing of *Mycobacterium tuberculosis*. *Nature Med* 3: 415-418.
- World Health Organization (1997). Prevalence and incidence of tuberculosis in India. A comprehensive review. WHO/TB/97, 231.
- World Health Organization Stop TB partnership (2009). New laboratory diagnosis tools for tuberculosis control. <http://www.apps.who.int/tdr>.



Understanding Tuberculosis - Global Experiences and Innovative Approaches to the Diagnosis

Edited by Dr. Pere-Joan Cardona

ISBN 978-953-307-938-7

Hard cover, 552 pages

Publisher InTech

Published online 15, February, 2012

Published in print edition February, 2012

Mycobacterium tuberculosis is a disease that is transmitted through aerosol. This is the reason why it is estimated that a third of humankind is already infected by Mycobacterium tuberculosis. The vast majority of the infected do not know about their status. Mycobacterium tuberculosis is a silent pathogen, causing no symptomatology at all during the infection. In addition, infected people cannot cause further infections. Unfortunately, an estimated 10 per cent of the infected population has the probability to develop the disease, making it very difficult to eradicate. Once in this stage, the bacilli can be transmitted to other persons and the development of clinical symptoms is very progressive. Therefore the diagnosis, especially the discrimination between infection and disease, is a real challenge. In this book, we present the experience of worldwide specialists on the diagnosis, along with its lights and shadows.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Shamsher S. Kanwar (2012). Diagnostic Methods for Mycobacterium tuberculosis and Challenges in Its Detection in India, Understanding Tuberculosis - Global Experiences and Innovative Approaches to the Diagnosis, Dr. Pere-Joan Cardona (Ed.), ISBN: 978-953-307-938-7, InTech, Available from: <http://www.intechopen.com/books/understanding-tuberculosis-global-experiences-and-innovative-approaches-to-the-diagnosis/diagnostic-methods-for-mycobacterium-tuberculosis-and-challenges-in-its-detection-in-india>

INTech
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen