We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists



186,000

200M



Our authors are among the

TOP 1% most cited scientists





WEB OF SCIENCE

Selection of our books indexed in the Book Citation Index in Web of Science™ Core Collection (BKCI)

Interested in publishing with us? Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected. For more information visit www.intechopen.com



Chemical Degradation of Chlorinated Organic Pollutants for *In Situ* Remediation and Evaluation of Natural Attenuation

Junko Hara

Institute for Geo-resources and Environment, National Institute of Advanced Industrial Science and Technology, Japan

1. Introduction

Chlorinated organic compounds, prevalent contaminants found in the geo-environment, pose an ecological risk even at trace concentrations. More volatile chlorinated compounds such as VOCs (volatile organic compounds) have been detected in urban areas and industrial zones because of the use of these compounds as components of industrial solvents and both raw and intermediate synthetic products. Chlorinated organic compounds quickly evaporate from surface water but remain in groundwater and soil for a long time. Recently, several remediation techniques have been developed that can entirely remediate chlorinated organic compounds to non-toxic materials. However, high molecular weight chlorinated organic compounds (e.g., polychlorinated biphenyls (PCBs) and dichloro-diphenyltrichloroethane (DDT)) are highly toxic chemicals that persist for long periods of time in the environment and bioaccumulate. They are categorized as persistent organic pollutants (POPs). Although the amount and date of use vary by country, POPs were widely used for pesticides and disease control in crop production and industrial processes during the period of industrial production after World War II around the globe. DDT, PCBs and dioxins are the best known POPs. DDT is used to control mosquitoes, which carry malaria, and PCBs were useful in electrical transformers and large capacitors. Among these POPs, PCBs, hexachlorobenzene (HCB), chlordanes, dichloro-diphenyl-dichloroethane (DDE) and dieldrin show significant ecological accumulation, such as in human milk, human blood and other biological media.

The Stockholm Convention is intended to protect human health and the environment, starting with the reduction or elimination of the production, use, and/or release of 12 species of POPs (PCBs, HCB, aldrin, dieldrin, endrin, DDT, chlordane, toxaphene, mirex, polychlorinated dibenzo-*p*-dioxins heptachlor, (dioxins), and polychlorinated dibenzofurans (furans). The 9 additional chemicals adopted in amendments to the Stockholm convention as new POPs are α-hexachlorocyclohexane, βhexachlorocyclohexane, chlordecone, hexabromobiphenyl, hexabromobiphenyl ether and heptabromobiphenyl ether, Lindane, pentachlorobenzene, perfluorooctane sulfonic acid and its salts and perfluorooctasulfonyl fluoride, tetrabromobiphenyl ether and pentabromobiphenyl ether in 2009. The Stockholm Convention has led to a general global

decline in the concentration of these chemicals in the environment. However, some individual POPs still persist and accumulate in fatty tissue and are present in higher concentrations at higher levels in the food chain, with long-range mobility through natural processes, because complete removal of POPs from the environment is difficult.

Experimental degradation methods for POPs at room temperature have been reported, using chemical catalysis, bacteria, UV, photocatalysis, Fenton reagent and other methods. Among these techniques, reductive dechlorination processes encounter difficulty in achieving complete dechlorination and degradation of the chemical structure. Powerful oxidative processes are assumed to show some possibility for complete degradation of POPs. Reports of complete degradation of POPs are few, even when oxidative processes are applied for on-site remediation over a very extensive polluted area.

This chapter reviews remediation methodology for chlorinated organic pollutants and chemical remediation methods using ferric sulphide compared with reaction with zero-valent iron, which widely used as a practical in-situ remediation method for soil and groundwater pollution by VOCs. Ferric sulphides also have outstanding ability to degrade chlorinated organic pollutants, but the dechlorination processes differs from that of zero-valent iron. The natural remediation capability, reaction products, and reaction mechanisms using ferric sulphide for trichloroethylene, dieldrin and chlorinated benzenes are reported in this chapter.

2. Reported remediation methods for chlorinated organic pollutants

Several remediation methodologies for chlorinated organic pollutants such as bioremediation, bioaugmentation, and chemical or physical remediation have been reported by many researchers. In these remediation methodologies, chlorinated organic contaminants can be transformed chemically, photochemically, or biochemically by oxidation or reduction in the soil and the groundwater environments.

Bioremediation technology has developed rapidly in the last 15 years, and the application of this technology has been extended to several species of contaminants, including volatile organic compounds (VOCs), polycyclic aromatic hydrocarbons (PAHs) and polychlorinated biphenyls (PCBs). Reductive dechlorination by microorganisms under anaerobic conditions is an advantageous process because reductive dechlorination allows the reoxidation of metabolic intermediates. For VOCs, PCE (tetrachloroethylene) or TCE (trichloroethylene), which is most widely reported chemicals, are capable of reductive dechlorination by dehalococcoides, dehalobacter and desulfuromonas (Hollinger et al., 1993; Gerritse et al., 1996, 1999; Loffler et al., 1999; Krumholz et al., 1996; Maymo-Gatell, 1997). These microorganisms dechlorinated the PCE/TCE to cis-dichloroethylene (cis-DCE) or vinyl chloride (VC). Dehalococcoides especially (including the culture) is reported to have achieved complete dechlorination of PCE/TCE to ethylene. In aerobic or oxidizing environments, chlorinated ethylenes are oxidized to CO₂ and the chlorinated ethylenes are co-metabolized to CO_2 via trichloroacetate, trichloroethanol, trichloroethylene epoxide, dichloroacetate, glyoxylate, formate, oxalate, etc. The reaction pathway and the number of intermediates differ according to the species of bacteria in the local environment (Newman and Wackett, 1991, 1995, 1997; Weightman et al., 1985, 1993; Li and Wackett, 1992; Kim et al., 2009; Rosenzweig et al., 1993; Oldenhuis et al., 1989; Motosugi et al., 1982). Oxygen served as the electron acceptor in the aerobic oxidation, and non-specific microbial oxygenase enzymes produced by aerobic microorganisms participated in co-metabolism reactions.

Anaerobic microbial communities in sediments also dechlorinate PCBs. The ease of dechlorination of positions on aromatic rings is usually meta > para > ortho, and on biphenyl rings, mono-ortho- and ortho-chlorobiphenyls were not degraded after one year of reaction (Teidje *et al.*, 1993). PCE was also oxidatively degraded by Pseudomonas strain sp. P2. In this reaction, PCE is metabolized to chlorobenzoic acids with one to three chlorine atoms. At the actual reaction site, reductive and oxidative biological reaction is estimated to occur because of conditions at the site. Each reaction is therefore assumed to proceed partly under anaerobic conditions and partly under aerobic conditions at one polluted site, and this collaboration of reactions under different sets of conditions leads to complete remediation at the natural site.

Many chemical remediation methods have been investigated for rapid in-situ or off-site remediation of soil and groundwater. Among the chemical and photochemical dechlorination methods, zero-valent iron (ZVI) has frequently been reported as the remediation technique for reductive dechlorination of chlorinated ethenes and ethanes. Among actual chemical remediation techniques, the ZVI methodology is widely used for in situ remediation for soil (e.g., injection by direct drilling) and groundwater (application in a reactive permeable barrier). The chlorinated organic materials degrade under anaerobic (reducing) conditions using ZVI. This reductive dechlorination is generally divided into hydrogenolysis and reductive elimination. Both reaction mechanisms are accompanied by a net transfer of two electrons. Although reductive dechlorination is occasionally referred to as hydrolytic reduction, this term is misleading because hydrolysis is incidental to the actual reductive remediation catalysts instead of ZVI. The reaction pathway for TCE using ZVI and other metals is the same as the reductive dechlorination process using bioremediation.

Although the application of ZVI for remediation of high molecular weight persistent organic compounds is difficult, zero-valent metals such as Pd, Pt, Ni and Cu are also used as catalysts in the ZVI methodology for dechlorination of PCBs. Metal catalysis enhances the reductive dechlorination capability of ZVI. In the case of persistent organic compounds, electrochemical methods using metal electrodes are used for reductive dechlorination.

Fenton reaction and photochemical reaction are the predominant oxidative dechlorination processes and can remediate contaminants rapidly. The Fenton reaction is the oxidation of organic substrates by iron(II) and hydrogen peroxides. The Fenton reagent is effective in treating various industrial wastewaters polluted by chlorinated organic compounds, aromatic amines, pesticides, and surfactants. This oxidation system is based on the formation of reactive oxidizing species able to degrade the contaminants effectively in wastewater. In the Fenton chemistry, a two-reaction pathway is advanced as the first step. Production of hydroxyl radicals and a non-radical pathway using ferric ion production have been reported (Barbusinski, 2009). The nature of the oxidizing species is still controversial. Some researchers showed that the hydroxyl radical is the major species in the Fenton mechanism, and other groups showed that this reaction includes the formation of reactive oxidizing species such as ferric ion. Considering that the Fenton reaction occurs chemically and biologically as well as in the natural environment, there is a possibility that both mechanisms coexist in the Fenton reaction. Several hazardous pollutants can be oxidized by the Fenton reaction; for example, chlorophenol is degraded to hydroxyacetic intermediates (Barbeni et al., 1987a) and perchloroethylene is transformed to dichloroacetic acid, formic acid and CO₂ (Leung et al., 1992). More detailed information about Fenton reaction is discussed in the chapter "Fenton's Process for the Treatment of Mixed Waste Chemicals".

In natural waters exposed to solar radiation, the Fenton reaction is often perceived as a possible source of hydroxyl radical in sunlit waters. The photolysis of nitrates, metal-toligand charge transfer reactions, and photoFenton reactions are included as other possibilities for the Fenton reaction, and H₂O₂ and Fe(II) are photochemically produced in these sunlit waters. Photochemical dechlorination is reported for persistent high molecular weight organic compounds such as DDT and PCBs (Mochizuki 1977; Van Beek et al, 1982; Shimakoshi et al., 2004). Photochemical PCBs dechlorination in alkaline isopropyl alcohol effectively degrades PCBs to biphenyl, chloride ions, acetone and water (Mochizuki, 1977). Dechlorination of DDT was catalyzed by hydrophobic vitamin B12 (heptamethyl cobyrinate perchlorate) with irradiation by visible light (Shimakoshi et al., 2004). The Co(I) species of cobalamin and the related cobalt complex are supernucleophiles and react with an alkyl halide to form alkylated complexes with dehalogenation (Shimakoshi H. et al., 2004). In this reaction, DDT is mono-dechlorinated to DDD. Photocatalytic treatment assisted by TiO₂ is also widely reported for TCE, DDT, chlorinated phenol, chloroform, etc. (Ahmed and Ollis, 1984; Barbeni et al., 1986, 1987b; Dible and Raupp, 1990, 1992; D'Oliveria et al., 1990; Kondo and Jardim, 1991; Borello et al., 1989) These oxidative degradations proceed by radicals arising from the photoreaction of other compounds present in the system.

In a similar reaction, UV/H_2O_2 and UV/Fenton are used for the remediation of chlorinated ethylenes and ethanes, chlorobenzenes, benzenes, and chlorinated phenols (Froclich, 1992; Moza *et al.*, 1988; Sundstrom *et al.*, 1986, 1989; Weie *et al.*, 1987). UV/H_2O_2 oxidation involves the dissociation of H_2O_2 to form the hydroxyl radical. The hydroxyl radical oxidizes toxic organic materials by abstraction of protons to produce organic radicals. Some toxic materials degrade to lower molecular weight acids and finally transform to CO_2 . However, many reports only estimate degradation ability and reaction kinetics and do not mention reaction products, especially when high molecular weight persistent organic compounds are involved.

These strong chemical oxidation technologies such as Fenton reagent, UV radiation, catalysis and photochemical treatment have the prospect of degradation of persistent organic pollutants to non-toxic compounds.

3. Remediation ability of ferric sulphides

The metal sulphides show a great diversity in electrical and magnetic properties. The sulphides of the transition metals can be considered as intermediate between the transition metal oxides. The small particles of metal sulphide are often superior in electrical and chemical properties, as well as in catalysis.

Pyrite is the most abundant metal sulphide mineral and is treated as industrial waste at many mining sites. An outcrop including pyrite leads to oxidation in aerobic weathering processes and causes acidification of environmental water. Pyrite is also distributed in acidic coastal sulphate soils, which are naturally formed under waterlogged anaerobic conditions. These acidic sulphide soils are located in tropical coastal areas in West Africa, South and Southeast Asia, and northeast South America. These acidic conditions are mainly developed in recent or semi-recent sediments close to the sea. In these environments, the sulphur in pyrite is derived from the sulphate in seawater, which is biologically reduced to sulphide in the anaerobic mud. The organic matter which serves as the energy source for the sulphate-reductive bacteria is usually abundant in plants in the coastal area. Ferrous iron is also derived from reduction of insoluble ferric compounds resulting from the weathering of clay.

The reactivity of pyrite when the surface is exposed to H₂O, O₂, and a mixture of H₂O and O₂, has been studied experimentally by Guevremont *et al.* (1997, 1998a, b, c, d). The pyrite surface exposed to H₂O vapour up to 1 bar was oxidized, although the reaction site is limited to the defect site on pyrite surface. The surface reaction with O₂ vapour showed no oxidation. Substantial surface reaction is observed in the reaction with the H₂O/O₂ mixture. The intermediate oxidation products, sulphur oxoanion and zero-valent sulphur, are also identified and removed by use of either the O₂ or H₂O (Kendelewicz *et al.*, 2004). The oxidation mechanism of pyrite with the H₂O/O₂ mixture involves competitive adsorption of O₂ and H₂O at the surface Fe site, oxidation of surface Fe sites by O₂, dissociation of co-adsorbed H₂O at Fe sites, and charge redistribution in surface S atoms (Rosso *et al.*, 1999). This pathway allows for the production of hydroxyl radicals from dissociated water and subsequent nucleophilic attack of these hydroxyl radicals at surface S sites. The oxygen in the final product sulphate arises from water molecules (Taylor *et al.*, 1984; Usher *et al.*, 2004, 2005). The hydroxyl radical from the pyrite oxidation process causes the degradation of

3.1 Degradation ability for major volatile organic compounds

chlorinated organic compounds.

Chlorinated ethylenes are widespread groundwater and soil contaminants. Due to the prevalent pollution and the efforts to treat these compounds, substantial research has been conducted to identify the mechanism of reaction under various environmental conditions. As mentioned above, the chemical dechlorination method using ZVI is in wide use as a practical in situ remediation technique for soil and groundwater pollution with TCE. This reductive dechlorination reaction process includes hydrogenolysis, β -elimination and a hydrogen addition reaction, and degrades chlorinated ethylenes and ethanes to ethylene and ethane. The dechlorination of TCE by pyrite (ferric sulphide) under anaerobic conditions is also reported to be reductive dechlorination like ZVI (Weerasooriya R., 2001; Lee and Batchelor, 2002, 2003). Other chlorinated ethylenes and ethanes are also reductively dechlorinated under anaerobic conditions (Kriegman-King and Reinhard, 1994; Lee and Batchelor, 2003). The reductive dechlorination of chlorinated compounds occurs by the transfer of electrons from the mineral surface.

Fig. 1 shows the dechlorination of TCE by metallic sulphides and zero- valent iron. In this reaction, 100 ppm of TCE is dechlorinated with powdered pyrite and Milli-Q-gradeultrapure water under closed aerobic conditions. TCE is detected by a headspace method using gas chromatography (GC) (GC17A, Shimadzu Co. Ltd.). Pyrite and chalcopyrite showed an outstanding dechlorination rate, and complete dechlorination was confirmed by the mass balance of chloride ion. According to the recent pyrite oxidation work, the most reactive surface component is S²⁻, and the second most reactive surface component is the surface atom of the first disulphide layer (S_2^{2-}) with sulphur atoms of the disulphide groups beneath the surface layer being least reactive. Oxidized iron (Fe³⁺) states are proposed to arise after fracturing of S-S bonds by electron transfer from Fe²⁺ to this S⁻ state, which then reacts rapidly to sulphate (Nesbitt et al., 1998; Schaufuss et al., 1998). If this oxidation reaction is similar with metallic sulphide, the S-S bonds in metallic sulphide oxidize chlorinated contaminants, as assumed from this oxidation model. Disulphide metallic minerals involving S-S bonds are therefore assumed to have greater degradation ability than the mono-sulphide metals. In our experimental estimation in Figure 1, the disulphide metallic minerals are only pyrite and chalcopyrite, which have superior degradation ability relative to the other mono-sulphide metals and ZVI.



Fig. 1. Dechlorination of trichloroethylene (TCE) by metallic sulphide and zero-valent iron under aerobic conditions. C_0 is the initial concentration of TCE (100 mg/L), C_t is the concentration of TCE at time t. The reaction was represented as the evaluation of the normalized remaining percentage of TCE with time.

TCE is also degraded by pyrite under aerobic conditions (Hoa *et al.*, 2008). TCE dechlorination in anaerobic and aerobic pyrite suspensions is observed with time, but there is no outstanding degradation under anaerobic conditions and the dechlorination rate of TCE is proportional to the increase in oxygen. Fig. 2 shows the disappearance of TCE, the reaction intermediates and the final product in the reaction of TCE with a pyrite suspension under aerobic conditions. Under aerobic conditions, 98 % of TCE was degraded after about 2 weeks. The TCE degrades to dichloroacetic acid, glyoxylic acid, formic acid, oxalic acid and CO_2 by oxidative processes (Hoa *et al.*, 2009). This degradation process is similar to the oxidative metabolic pathway of TCE (Kim *et al.*, 2009; Li *et al.*, 1992). Figure 3 shows the expected dechlorination pathway of TCE by pyrite under aerobic conditions. The dechlorination pathway of TCE is divided into three pathways. All pathways show direct dechlorination of three chlorine atoms in TCE. The main pathway is degradation of TCE to formic acid (Eq. (1)).

$$C_2HCl_3(TCE) + 4^{\bullet}OH + H_2O \rightarrow 2CH_2O_2(formic \ acid) + 3HCl + \frac{1}{2}O_2$$
(1)

$$CH_2O_2(formic acid) + 2^{\bullet}OH \rightarrow CO_2(g) + 2H_2O$$
 (2)

Formic acid has toxicological properties toward aquatic organisms, but the degradation rate of formic acid by pyrite suspensions is also high, similar to the dechlorination of TCE to formic acid, and formic acid continuously transforms to $CO_2(g)$ without accumulation (Eq. (2)).



Fig. 2. Transformation of trichloroethylene (TCE) and detected reaction products in pyrite suspension (Edited from Hoa *et al.* (2008)). The initial TCE concentration is 10 μ mol, the oxygen volume is 0.268 mmol/L, and pyrite is 10 m²/L. The left side of the Y-axis shows the molecular weight of TCE, CO₂ and organic acids (formic acid, oxalic acid, dichloroacetic acid, glyoxylic acid). The right side of the Y-axis shows the chloride molecular weight as degraded from TCE.



Fig. 3. Degradation pathway of trichloroethylene (TCE) using a pyrite suspension under aerobic conditions (Edited from Hoa *et al.* (2009)).

3.2 Degradation capability for dieldrin

Dieldrin is a cyclodiene pesticide which has persisted in the soil over decades in some agricultural fields. Some agricultural crops and animals accumulate dieldrin through the food chain. From the previous report of TCE dechlorination by pyrite, we see that TCE is able to be reductively degraded under anaerobic conditions and oxidatively degraded under aerobic conditions. The reductive dechlorination under anaerobic conditions is not observed using pyrite suspensions. Dieldrin has a higher persistence than TCE and is less amenable to

reductive dechlorination under anaerobic conditions. Remediation techniques for dieldrin (UV/Fenton reagent, UV/chemical reaction, Pd/C catalyst, solar photocatalysis, and bioremediation) have been reported (Books, 1980; Maule *et al.*, 1987; Bandala *et al.*, 2002; Kusvuran and Erbatur, 2004; Chiu *et al.*, 2005; Zinovyev *et al.*, 2005; Dureja *et al.*, 1987; Baczynski *et al.*, 2004), but dieldrin transforms to mono- or di- dechlorinated intermediates still having a bicyclic ring structure. A study of the use of Fenton reagent for aldrin reports the transformation from aldrin to oxalic acid, acetic acid, chlorohexanone, cyclohexanol, cis-2-hydroxy-cyclohexanone, cis-2-methyl-cyclohexanol, 4-hydroxycyclohexanone, cis-4-methylcyclohexanol, 1-cyclopropyl-1-hydroxyethylene, and trans-dihydroxycyclohexane (Kusvuran and Erbatur, 2004).



Fig. 4. Degradation profile of dieldrin using pyrite and zero-valent iron under aerobic conditions. The blank denotes the non-metallic catalysis condition. C_0 is the initial concentration of dieldrin (0.131 µmol), C_t is the concentration of dieldrin at time t. (Edited from Hara *et al.*, (2009)

The degradation of dieldrin using ZVI and pyrite under anaerobic conditions is shown in Fig. 4. ZVI and pyrite have similar dechlorination capabilities for dieldrin, but the reaction process differs completely for the two reagents (Hara *et al.*, 2009). In the case of ZVI, dieldrin partially transforms to nono-dechlorinated products ($C_{12}H_9Cl_5$, $C_{12}H_9Cl_5O$) and aldrin ($C_{12}H_8Cl_5$). This reaction is a reductive dechlorination proceeded by the generation of H+ and an electron arising from oxidation of ZVI. This reaction is stopped only this pathway. In contrast, the mono-dichlorinated reaction intermediates and aldrin are not detected in the reaction using pyrite. Dieldrin is oxidatively degraded to water-soluble reaction intermediates by pyrite.

Fig. 5 shows the dechlorination rate of dieldrin using pyrite under anaerobic to aerobic conditions, changing with oxygen concentration ($O_2 = 0 \sim 833 \mu mol$). Dieldrin was gradually degraded under every condition of oxygen concentration, except for the $O_2 = 0 \mu mol$ condition. In Fig. 5, 5 ppm (0.131 μmol) of dieldrin is used with a powdered pyrite suspension under each oxygen condition. Dieldrin is detected by GC/MS (Shimadzu Co. Ltd.) after solvent/solvent extraction using acetone and hexane. The water-soluble reaction intermediates are detected by IC-TOF/MS (ICS-3000 (Dionex) and JMS-T 100LP (JEOL). A little oxygen ($O_2 = 10 \mu mol$) accelerates the dechlorination of dieldrin in

comparison to the anaerobic conditions. Approximately 99 % of the dieldrin was degraded after about one month under the most enhanced reaction conditions ($O_2 = 10 \mu mol$), and approximately 43 % of dieldrin remained under $O_2 = 300 \mu mol$ in the same reaction time, which is the slowest dechlorination condition. In these reactions, chloride ion resulting from the dechlorination was detected in the aqueous phase. The mass balance of dechlorinated chloride and degraded dieldrin is 90 % in the most degraded condition ($O_2 = 10 \mu mol$). This discrepancy in mass balance is obvious in the anaerobic conditions, but the mass balance agrees well under aerobic conditions ($O_2 = 300 \mu mol$).



Fig. 5. Oxygen dependence on degradation of dieldrin in pyrite suspension (Hara, 2011). C_0 is the initial concentration of dieldrin (0.131 µmol), C_t is the concentration of dieldrin at time t. O_2 = 833 µmol denotes the air volume conditions in this experiment. The blank is without pyrite.

As reaction products, formic acid, oxalic acid, malonic acid, succinic acid, acetoxyacetyl, lactic acid, and pyruvic acid are detected as the main reaction products in the case of aerobic conditions ($O_2 = 833 \mu mol$). Acetic acid, glycolic acid, hydroxybutyric acid, glyoxylic acid, propionic acid, glutaric acid, levulinic acid and sulphur-containing organic acids such as methanesulphonic acid, sulphopropionic acid, etc. are also detected as minor reaction products. Formic acid is predominantly generated, as shown in (Eq. (3)).

$$C_{12}H_8Cl_6O(dieldrin) + 24^{\circ}OH + H_2O \rightarrow 12CH_2O_2(formic acid) + 6HCl + O_2$$
(3)

Formic acid is also detected in TCE dechlorination by a pyrite suspension under aerobic conditions, and the formic acid continuously degrades to CO_2 (Eq. (2)). Dieldrin is therefore assumed to finally transform to CO_2 as a main reaction pathway.

In the case of a low O_2 volume or anaerobic conditions, these low molecular weight organic acids generated under oxygen-rich conditions are much lower in abundance and some organic acids are not detected. Instead of the generation of organic acids, 3-chloro-4-methyl-2-pentanol (C₆H₁₃ClO) and dibutyl phthalate (C₁₆H₂₂O₄) are generated. One part of these intermediates is continuously degraded to a low molecular weight organic acid, but the residual volume is also higher than the volume observed under aerobic conditions.

The reaction pathway of dieldrin proceeds mostly by oxidative degradation under anaerobic and aerobic conditions. Although the oxidative ability is obviously different from the content of oxygen, the active oxidant to degrade the dieldrin is promoted from both the pyrite/H₂O and pyrite/O₂ interface reaction, due to oxidants produced under either aerobic or anaerobic condition. The difference of degradation ability in oxygen volume is assumed to depend on the difference of radical species arising form pyrite/H₂O and pyrite/O₂ interface.

Considering the reaction products, the degradation pathway of dieldrin results mainly in the direct production of organic acids, which are easy to produce under oxygen-rich conditions. Under anaerobic or micro-aerobic conditions, the pathway of ring opening and the addition reaction of low molecular weight organic acids arising from $C_{16}H_{22}O_4$ and the pathway of generation of chlorinated hydrocarbons arising from $C_{6}H_{13}CIO$ (which continuously transforms to formic acid and malonic acid) become predominant.

3.3 Degradation ability for chlorinated benzenes

Dieldrin having a bicyclic ring could be decomposed by pyrite. This section discusses the ability to dechlorinate chlorobenzenes whose main structure is the benzene ring.

Chlorobenzenes are divided into 12 species based on the number and configuration of chlorine in the molecule: monochlorobenzene (mono-CB), 3 types of dichlorobenzene (1,2-, 1,3-, 1,4- di-CB), 3 types of trichlorobenzene (1,2,3-, 1,2,4-, 1,3,5- tri-CB), 3 types of tetrachlorobenzene (1,2,3,4-, 1,2,3,5-, 1,2,4,5- tetraCB), pentachlorobenzene (penta-CB), and hexachlrobenzene (hexa-CB). The chlorobenzenes are used in pesticides, deodorants, or as intermediates in a chemical synthesis process. Some tri- or tetra-chlorinated benzenes are extensively used as insulating materials. The risk associated with chlorinated benzenes increases relative to the increase in the chlorine number, because of their increasing lack of volatility with increasing chlorine number, the highest level of risk is associated with the misuse or accidental release of the mono- to trichlorinated benzenes. These compounds spread into the atmosphere are reported to be photolyzed or chemically reacted, and that in groundwater and soils these compounds are mainly remediated by microbial degradation. However, the residence time is increased because of the organic constituents present in the soil and groundwater, which results in adsorption and accumulation in the soil ecosystems. The major remediation method for more the accumulation of chlorobenzenes with a higher chlorination level in soils is incineration at high temperatures for digging out of soils.

In the previous reports of electrochemical dehalogenation of chlorinated benzenes (Miyoshi et al., 2004; Mohammad and Dennis, 1997; Farwell et al., 1975; Kargina et al., 1997; Guena et al., 2000), the chlorine is eliminated step by step from the highly chlorinated benzenes to yield less-chlorinated benzenes and finally transform to benzene. Farwell et al. (1975) reported on chlorobenzenes and the main cathodic reaction pathway for hexachlorobenzenes as follows: hexachlorobenzene \rightarrow pentachlorobenzene \rightarrow 1,2,3,5tetrachlorobenzene \rightarrow 1,2,4-trichlorobenzene \rightarrow 1,4-dichlorobenzene \rightarrow monochlorobenzene \rightarrow benzene. This dechlorination pathway is promoted by the electrochemical reductive dechlorination. Benzene is detected as the final reaction product.

Oxidative dechlorination of chlorobenzenes has been reported using bacteria, Fenton reagent, UV/H₂O₂, and TiO₂-assisted photocatalysis. In the microbial metabolism reported by Reineke and Knackmuss (1984), chlorobenzene was gradually degraded to 3-chloro-cis - 1,2-dihydroxycylohexa-3,5-diene, 3-chlorocatechol, 2-chloro-cis,cis-muconate, trans-4-carboxymethylenebuten-4-olide, maleylacetate, and 3-oxoadipate. Fenton reagent effectively

www.intechopen.com

354

degrades chlorobenzene to chlorohydroxycyclohexadienyl radical in the first step, and the radical dimerizes to produce dichlorobiphenyls, with bimolecular disproportionation to produce chlorophenol and chlorobenzene under non-oxygen conditions or in the absence of other strong oxidants (Reineke W. and Knackmuss H-J, 1984). In the presence of oxygen or other strong oxidants, reactions of the oxidant (O_2) with chlorohydroxycyclohexadienyl radical is predominant and results in lower concentrations of dichlorobiphenyl, which decreases remarkably, and chlorobenzoquinone is formed. Chlorophenol isomers were further oxidized by hydroxyl radical and formed chlorinated and non-chlorinated diols. Chlorobenzenes were also dechlorinated by the UV/H₂O₂ treatment method (Sundstrom *et al.,* 1989), and chlorobenzene was degraded to phenol, biphenyl, chlorobiphenyl isomers, and benzaldehyde with only UV, and chlorobenzenes also transform to chlorophenol and various isomers of chlorobiphenyl and dichlorobiphenyl in the UV/H₂O₂ system.

Degradation of chlorobenzenes in pyrite suspensions is shown in Fig. 6 and in Table 1, with four volatile species: mono-chlorobenzene, 1,2-dichlorobenzene, 1,3-dichlorobenzene, and 1,2,4-trichlorobenzene. The concentrations of these compounds were determined by headspace methods. The experiments were conducted at 25 °C at 200 rpm in the pyrite suspension under anaerobic conditions. The initial concentrations of chlorobenzenes were set to 10 mg/L using the each pure chlorobenzenes dissolved in hexane water. The initial rate of degradation is 1,2,4-trichlorobenzene > 1,2-dichlorobenzene > monochlorobenzene > 1,3-dichlorobenzene, and more than 90 % of these chlorobenzenes were dechlorinated after merely 10 days. The degraded amount of chlorobenzenes is not sensitive to differences in the distribution of their isomers. However, the amount of chloride ion arising from dechlorination of chlorobenzenes is less than the entire chlorine content from each of the chlorobenzenes. This disagreement in mass balance also shows less than one chloride ion degraded from each chlorobenzene. The order of total dechlorination ability is 1,2dichlorobenzene > monochlorobenzene > 1,2,4-trichlorobenzene > 1,3-dichlorobenzene. Among the 4 species of chlorobenzenes estimated here, 1,3-dichlorobenzene is the hardest to degrade because it has a meta-site chloride configuration. The meta-site dechlorination is not easier than 1,2-(ortho) and 1,4-(para-) dichlorobenzenes.



Fig. 6. Degradation of monochlorobenzene, 1,2- and 1,3-dichlorobenzene, and 1,2,4trichlorobenzene in pyrite suspension under aerobic conditions. C_0 is the initial concentration of chlorobenzenes (10 mg/L), C_t is the concentration of chlorobenzenes at time t.

species	$\begin{array}{c} -\Delta C^t_{CBs} / C^0_{CBs} \\ [mol\%] \end{array}$	$+\Delta C_{Cl}^{t}/C_{CBs}^{0}$ [mol%]
MonoCB	88.1	76.5
1,2-DiCB	96.4	88.4
1,3-DiCB	89.1	56.6
1,2,4-TriCB	95.6	68.7

Table 1. The fraction of degraded chlorinated benzenes and chloride ion arising from chlorinated benzenes after 10 days.

Fig. 7 shows the ratio of the residual chlorinated chlorobenzenes relative to the degradation of all species of chlorinated benzenes by pyrite suspension. Table 2 is the fraction of dechlorinated chloride ion from each chlorobenzene and the initial chloride content of the chlorobenzenes. The volatile chlorinated benzenes, mono-, 1,2-di-, 1,3-di- and 1,2,4-trichlorobenzenes were analyzed by the headspace method in addition to solvent extraction analysis to determine their concentrations. The non-degraded volatile chlorobenzenes adsorbed on the pyrite surface are correctly estimated here. The reaction products were also analyzed by GC/MS after eluting in the organic solutions along with the other chlorinated benzenes.

The ratio of complete dechlorination is higher for low molecular weight chlorobenzenes, and 1,2-dichlorobenzene is easier to degrade than 1,3-dichlorobenzene. The greater electron deviation due to chlorine configuration allows the degradation of chlorobenzenes. More than 80 to 90 % of mono-, di-, tri-chlorobenzenes, and 1,2,3,4- and 1,2,3,5-tetrachlorobenzene are degraded from the original concentrations. These compounds are easy to degrade, but there is a small variation depending on chlorine configuration. The residual ratio is 1,2- < 1,3- < 1,4- among the dichlorobenzenes, and 1,2,3- < 1,2,4- < 1,3,5- among the trichlorobenzenes. There is a significant difference in dechlorination ability for the tetrachlorobenzene isomers. The symmetrical chlorine configuration, 1,2,4,5-, on the benzene ring is stable and hard to degrade, so 1,2,4,5-tetrachlorobenzene, tetra- and hexachlorobenzenes with the 1,2,4,5-chlorine configuration are extremely difficult to degrade. Considering the mass balance of persistent chlorobenzenes and chloride ions, di-, tri-, and tetra- (limited to 1,2,3,4- and 1,2,3,5-) chlorobenzenes are transformed into chlorinated intermediates because the detected concentration of chloride ion is only one mole of chloride ion per one mole of chlorobenzene and that corresponds to a decline of the initial concentration of each of the chlorobenzenes.

The GC/ MS analysis of the solvent-extracted samples after 10 days indicates the presence of one or more dechlorinated chlorobenzenes from the initial materials for each chlorobenzene, except for hexachlorobenzene. In the experimental dechlorination of hexachlorobenzene, pentachlorobenzene is detected after 10 days. Hexachlorobenzene is assumed to have a slow dechlorination rate so that a little of the compound could be detected. As an example of reaction products, Fig. 8 shows the analytical result of GC/MS and NMR analysis for the solvent extracted from the reaction system of monochlorobenzenes. C₉H₁₄(OH)Cl, C₁₃H₁₇(OH)₃, C₁₆H₂₃SO₂ are detected by MS together with cyclic sulphur molecules such as S₆, S₇, S₈. The other saturated or unsaturated straight-chain hydrocarbons, such as 3-decyne (C₁₀H₁₈), 8,10-dodecadienial (C₁₂H₂₀O) and 2,2,6-trimethyl-1,4-cyclohexanedione (C₉H₁₄O₂) are detected as one of the reaction products of tri- to hexachlorobenzenes. The NMR spectrum also shows the alcohol, ether, and aromatic methyl peaks ($-CH_2OH$, -O-, -COH, $C_6H_6-CH_3$) with the benzene and chlorobenzene peaks. One part of the benzene ring structure is assumed to be opening in this reaction process.



Fig. 7. Residual ratio of 12 species of chlorinated benzenes estimated under aerobic conditions in pyrite suspension after 10 days (Hara *et al.* (2006).

species	+ $\Delta C_{Cl-}^{t} / C_{Cl-}^{0}$ [mol%]
monochlorobenzene	89
1,2-dichlorobenzene	47
1,3-dichlorobenzene	45
1,4-dichlorobenzene	45
1,2,3-trichlorobenzene	30
1,2,4-trichlorobenzene	29
1,3,5-trichlorobenzene	30
1,2,3,4-tetrachlorobenzene	20
1,2,3,5-tetrachlorobenzene	18
1,2,4,5-tetrachlorobenzene	11
pentachlorobenzene	7.5
hexachlorobenzene	5.5

Table 2. The fraction of dechlorinated chloride ion from each chlorobenzene and initial chloride content of chlorobenzenes.



Fig. 8. The spectrum of reaction intermediates arising from transformation of monochlorobenzene in pyrite suspension. (a) MS spectrum. (b) MNR spectrum. In the NMR spectrum, 1. aromatic peak (benzene and monochlorobenzene), 2-4. alcohol and ether peak (–CH₂OH, -O-, -COH), 5. aromatic methyl (methylbenzene).

From the analytical results, the benzene ring could have been opened by the dechlorination process. One part of the benzene ring opened to straight chain unsaturated hydrocarbons and these hydrocarbons polymerized with other straight chain unsaturated hydrocarbons. Some polymerized straight chain hydrocarbons remain, and some hydrocarbons are combined with unsaturated ring structures. A ring-opening reaction of benzene is assumed to be a common reaction for every chlorobenzene under our experimental conditions, not only turning into cyclohexanes. These reactions are initiated when one chlorine site is dechlorinated by pyrite. Therefore, at least one mole of chlorine ions was dissolved in solution for every mole of chlorinated benzene molecules that reacted with pyrite. Furthermore, non-detected chlorine ions may still bind the carbon atoms that are produced by this reaction.

The reactions are proceeded by the oxidant arising on the pyrite interface. The polymerization and ring-opening reactions are similar to the dechlorination of Dieldrin under anaerobic conditions.

4. Discussion

Pyrite suspension shows degradation ability predominantly for TCE, Dieldrin and chlorobenzenes as shown in Section 3. These degradations of chlorinated organic compounds are initiated by the oxidant arising from the pyrite interface reaction.

The surface reaction of pyrite/H₂O and pyrite/O₂ produces the radical as follows (Cohn *et al.,* 2006). In the pyrite/H₂O interface under non-oxygen conditions, H₂O is hydrolysed to \bullet OH and H⁺ (Eq. (4) and Eq. (5))

$$Fe^{3+}(pyrite) + H_2O \rightarrow Fe^{2+}(pyrite) + {}^{\bullet}OH + H^+$$
 (4)

$$2(^{\bullet}OH) \to H_2O_2 \tag{5}$$

Commonly, the radical is apt to arise under aerobic conditions, but the hydroxyl radical can be generated under anaerobic conditions and it acts as an oxidant.

In the presence of oxygen (aerobic conditions), superoxide $(O_2 \bullet)$ - was detected as an intermediate from H_2O_2 formation. The superoxide $(O_2 \bullet)$ - is generated by the oxidation of iron on the pyrite surface as follows:

$$Fe^{2+}(pyrite) + O_2 \rightarrow Fe^{3+}(pyrite) + (O_2^{\bullet})^-$$
 (6)

$$Fe^{2+}(pyrite) + (O_2^{\bullet})^- + 2H^+ \to Fe^{3+}(pyrite) + H_2O_2$$
 (7)

The reactions of the pyrite/ O_2 interface generate hydroxyl radicals from hydrogen peroxide by the oxidation of surface iron (Eq. (6) and Eq. (7)), and hydrogen peroxide is also used for the reduction of ferric to ferrous ion in the reaction shown in Eq. (8). However, the hydroxyl radicals are not noticeably produced under aerobic condition.

$$Fe^{2+}(aq) + H_2O_2 \to {}^{\bullet}OH + OH^- + Fe^{3+}(aq)$$
(8)

Under aerobic conditions, superoxide and hydrogen peroxide are dominant oxidants used for degradation of chlorinated organic compounds.

Although there is a difference in oxidative ability among oxidants, oxidants arise from the pyrite interface under both aerobic and anaerobic conditions. The hydroxyl radical has an oxidant ability superior to superoxide and hydrogen peroxide, but the total amount of ferrous ion on the pyrite surface is limited under anaerobic conditions; therefore, the total volume of the hydroxyl radical is also limited under anaerobic conditions as opposed to aerobic conditions. The difference between reaction products under aerobic and anaerobic conditions is caused by thus radical character of the pyrite interface.

The oxidation and polymerization of chlorinated organic compounds are caused by several radicals arising from the pyrite/ H_2O , pyrite/ O_2 interface reaction.

5. Conclusion

This chapter reviews several remediation methods for chlorinated organic pollutants, and emphasizes the degradation ability of natural ferric sulphide (pyrite) for TCE, Dieldrin and chlorobenzenes. The transformation of TCE and Dieldrin under aerobic conditions becomes clear and the main reaction product is formic acid, which continuously degrades to CO₂. Although the 1,2,4,5-chloride configuration in the chlorobenzene and benzene ring structure is hard to degrade, one part of the benzene ring is transformed into several hydrocarbons.

The oxidative reaction of the pyrite suspension has great degradative ability not only for low molecular weight organic pollutants such as VOCs but also for high molecular weight persistent organic compounds with a stable benzene ring and bicyclic rings. Natural ferric sulphide has a remediation ability predominantly for chlorinated organic pollutants.

Iron and ferric sulphides (e.g., pyrite) are widely distributed in the subsurface layer in the global environment. The reductive degradation ability of iron and the oxidative degradation ability of ferric sulphides assist the innovative remediation techniques that are conducted at in situ normal temperature and pressure. The ferric ion and ferric sulphide latent in natural systems have a potential for natural attenuation of contamination.

Pyrite is also widely distributed at mining sites and soils in coastal regions of the Asian and the Pacific areas located in tropical, subtropical and temperate climate areas. For instance, the Mekong Delta in Vietnam is a vast acidic sulphate soil region that includes metallic sulphide. Garvalho *et al.* (2008) reported the agrochemical and polychlorobiphenyl residues in the Mekong River Delta. Several chlorinated compounds, such as DDT, HCH (hexacychlochlorohexane), PCBs and endosulfan were detected in sediment and biota. However, the concentrations of PCBs and pesticide residues in the aquatic environment in the Mekong River Delta are lower than the values reported in other regions of Vietnam and Asia. The aquatic environment of the Mekong Delta is endowed with natural ferric sulphide and water, and consequently the natural environment in this region is assumed to enhance PCB degradation. The acidic sulphate soils distributed in coastal area assumes to actually attenuate the contamination of chlorinated organic compound and prevent the expansion of pollution.

6. Acknowledgments

The author would like to thank Dr. Tsuji (Yamagata Environmental Research Institute) for assistance with the analysis and detection of reaction intermediates.

This work was made possible by a grant-in-aid for scientific research from the Japan Society for the Promotion of Science (B-17760656).

7. References

- Ahmed S. and Ollis D.F. (1984) Solar photoassisted catalytic decomposition of the chlorinated hydrocarbons trichloroethylene and trichloromethane. *Solar Energy*, Vol.32(5), 597-601.
- Baczynski T.P., Grotenhuis T. and Knipscheer P. (2004) The dechlorination of cyclodiene pesticide by metanogenic granular sludge. *Chemosphere*, Vol.55, 653-659.
- Bandala E.R., Gelover S. and Leal M.T. (2002) Solar Photocatalytic degradation of aldrin. *Catalysis Today*, Vol.76, 189-199.
- Barbeni M., Minero C. and Pelizzetti E. (1987a) Chemical degradation of chlorophenols with Fenton's Reagent. *Chemosphere*, Vol.16(10-12), 2225-2237.
- Barbeni M., Morello M., Pramauro E. and Pilizzetti E. (1987b) Sunlight Photodegradation of 2,4,5-trichlorophenoxy-acetic acid and 2,4,5-trichlorophenol on TiO₂. Identification of itermediates and degradation pathway. *Chemosphere*, Vol.16(6), 1165-1179.
- Barbeni M., Pramauro E. and Pilizzetti E. (1986) Photochemical degradation of chlorinated dioxins, biphenyls phenols and benzene on semiconductor dispersion. *Chemosphere*, Vol.15(9), 1913-1916.
- Barbusinski K. (2009) Fenton reaction controversy concerning the chemistry. *Ecological chemistry and engineering*, Vol.16(3), 347-258.
- Borello R., Minero C., Pramauru E., Pelizzetti E., Serpone N. and Hidaka H. (1989) Photocatalytic degradation of DDT mediated in aqueous semiconductor slurries by simulated sunlight. *Environmental Toxicology and chemistry*, Vol.8, 997-1002.
- Brooks G.T. (1980) The preparation of some reductively dechlorinated analogues of deildrin, endosulfan and isobenzan. *Journal of Pesticide Science*, Vol.5, 565-574.
- Chiu T., Yen J., Hsieh Y. and Wang Y. (2005) Reductive transformation of dieldrin under anaerobic sediment culture. *Chemosphere*, Vol.60, 1182-1189.

360

- Cohn C.A., Mueller S. Wimmer E., Leifer N., Greenbaum S., Strongin S.R. and Schoonen M.A.A. (2006) Pyrite-induced hydroxyl radical formation and its effect on nucleic acids. *Geochemical Transactions*, Vol.7(3), 1-11.
- Dibble L.A. and Raupp, G. B. (1990) Kinetic of the gas-solid heterogeneous photocatalytic oxidation of trichloroethylene by near UV illuminated titanium dioxide. *Catalysis Letters*, Vol.4, 345-354.
- Dibble L.A. and Raupp, G. B. (1992) Fluidized-Bed photocatalytic oxidation of trichloroethylene in contaminated airstreams. *Environmental Science and Technology*, Vol.26, 492-495.
- Dureja P., Walia S. and Mukerjee S. K. (1987) Superoxide mediated dehydrohalogenation of photodieldrin and photoaldrin. *Tetrahedron letters*, Vol.28(8), 895-896.
- D'Ollvelra J.-C., Al-Sayyed G. and Plchat P. (1990) Photodegradation of 2- and 3chlorophenol in TiO₂ Aqueous Suspensions. *Environmental Science and Technology*, Vol.24, 990-996.
- Farwell O., Beland F. A., and Geer R. D. (1975) Reduction pathways of organohologen compounds. Part 1. Chlorinated benzenes. *Electroanalytical Chemistry and Interfacial Electrochemistry*, Vol. 61, 303-313.
- Froelich E.M. (1992) Chemical Oxidation. *Technomic Publishing Company, Inc.* Eckenfelder W. W., Nowers A.R. and Roth J. A. (eds.) Lancaster, PA, 104-113.
- Garvallho F.P., Villeneuve J.P, Cattini C., Tolosa I., Dao Dinh Thuan and Dang Duc Nhan (2008) Agrochemical and polychlorobyphenyl (PCB) residues in the Mekong River delta, Vietnam. *Marine Pllution Bullutin*, Vol.56, 1476-1485.
- Gerritse J., Drzyzga O., Kloetstra G., Keijmel M., Wiersum L.P., Hutson R., Collins M.D. and Gottschal J.C. (1999) Influence of different electron donors and acceptors on dehalorespiration of tetrachloroethene by Desulfitobacterium frappieri TCE1. *Applied and Environmental Microbiology*, Vol.65(12), 5212-5221.
- Gerritse J., Renard V., Pedro Gomes T.M., Lawson P.A., Collins M.D. and Gottschal J.C. (1996) Desulfitobacterium sp. Strain PCE1, an anaerobic bacterium that can grow by reductive dechlorination of tetrachloroethene or ortho-chlorinated phenols. *Archives of Microbiology*, Vol.165(2), 132-140.
- Guena T., Wang L., Gattrell M. and MacDougall (2000) Mediated Approach for the electrochemical reduction of chlorobenzens in Nonaqueous media. *Journal of the Electrochemical Society*, Vol. 147, 248-255.
- Guevremont J.M., Elsetinow A.R., Strongin D.R., Bebie J.M. and Schoonen M.A.A. (1998b) Structure Sensitivity of pyrite oxidation: Comparison of the (100) and (111) planes. *American Mineralogist*, Vol.83, 1353-1356.
- Guevremont J.M., Strongin D.R. and Schoonen M.A.A. (1998c) Photoemission of adsorbed Xenon, X-ray photoemission spectroscopy, and temperature-programmed desorption studies of H₂O on FeS₂ (100). *Langmuir* 14: 1361-1366.
- Guevremont J.M., Strongin D.R. and Schoonen M.A.A. (1998d) Thermal chemistry of H₂S and H₂O on the (100) plane of pyrite: Unique reactivity of defect sites. *American mineralogist*, Vol.83, 1246-1255.
- Guevremont J.M., Bebie J., Elsetinow A.R., Strongin D.R. and Schoonen M.A.A. (1998a) Reactivity of the (100) plane of pyrite in oxidizing gaseous and aqueous environments: Effect of Surface imperfections. *Environmental Science and Technology*, Vol.32, 3743-3748.

- Guevremont J.M., Strongin D.R. and Schoonen M.A.A. (1997) Effects of surface imperfections on the binding of CH₃OH and H₂O on FeS₂ (100): Using adsorbed Xe as a probe of mineral surface structure. *Surface Science*, Vol.391, 109-124.
- Hara J. (2001) The effect of oxygen on chemical dechlorination of dieldrin using iron sulfides. *Chemosphere*, Vol.82, 1308-1313.
- Hara J., Kawabe Y., Komai T. and Inoue C. (2009) Chemical degradation of dieldrin using ferric sulfide and iron powder. *International Journal of Environmental Science and Engineering*, Vol.1-2, 91-96.
- Hara J., Inoue C., Chida T., Kawabe Y., Komai T. (2006) Dehalogenation of Chlorinated Benzenes by iron sulfide. *International Journal of Power and Energy Systems*, Vol.1(1), 239-243.
- Hoa T. P., Kitsuneduka M., Hara J., Suto K., Inoue C. (2008) Trichloroethylene Transformation by Natural Mineral Pyrite: The deciding role of oxygen. *Environmental Science and Technology*, Vol.42. 7470-7475.
- Hoa T.P., Suto K. and Inoue C. (2009) Trichloroethylene transformation in Aerobic pyrite suspension: pathways and Kinetic modeling. *Environmental Science and Technology*, Vol.43, 6744-6749.
- Holliger C., Schraa G., Stams A.J.M. and Zehnder A.J.B. (1993) A highly purified enrichment culture couples the reductive dechlorination of tetrachloroethene to growth. *Applied and Environmental Microbiology*, Vol.59 (9), 2991-2997.
- Kargina O., MacDougall B., Kargin Y. M. and Wang L. (1997) Dechlorination of monochlorobenzene using organic mediators. *Journal of the Electrochemical Society*, Vol.144, 3715-3721.
- Kendelewiz T., Dolyle C.S., Bostick B.C. and Brown G.E. (2004) Initial oxidation of fractured surface of pyrite (FeS₂(100)) by molecular oxygen, water vapor, and air. *Surface Science*, Vol.558, 80-88.
- Kim S., Kim D., Pollack G. M., Collins L. B. and Rusyn I. (2009) Pharmacokinetic analysis of trichloroethylene metabolism in male B6C3F1 mice: Formation and disposition of trichloroacetic acid, dichloroacetic acid, S-(1,2-dichlorovinyl) glutathione and S-(1,2-dichlorovinyl)-L-Cysteine. *Toxicology and applied pharmacology*, Vol.238, 90-99.
- Kondo M.M. and Jardim W.F. (1991) Photodegradation of chloroform and urea using Agloaded titanium dioxide as catalyst. *Water Research*, Vol.25(7), 823-827.
- Kriegman-King M.R. and Reinhard M. (1994) Degradation of carbon tetrachloride by pyrite in aqueous solution. *Environmental Science and Technology*, Vol.28, 692-700.
- Krumholz L.R., Sharp R. and Fishbain S.S. (1996) A freshwater anaerobe coupling acetate oxidation to tetrachloroethylene dehalogenation. *Applied and Environmental Microbiology*, Vol.62(11), 4108-4113.
- Kusveran E. and Erbatur O. (2004) Degradation of aldrin in adsorbed system using advanced oxidation processes: comparison of the treatment methods. *Journal of Hazardous Materials*, 106B, 115-125.
- Lee W. and Batchelor B. (2002) Abiotic reductive dechlorination of chlorinated ethyelens by iron-bearing soil minerals. 1. pyrite and magnetite. *Environmental Science and Technology*, Vol.36, 5147-5154.
- Lee W. and Batchelor B. (2003) Reductive capacity of natural reductants. *Environmental Science and Technology*, Vol.37, 535-541.

- Leung S.W., Watts R. J. and Miller G.C. (1992) Degradation of Perchloroethylene by Fenton's Reagent: Speciation and Pathway. *Journal of Environmental Quality*, Vol.21(3), 377-381.
- Li S. and Wackett L.P. (1992) Trichloroethylene oxidation by toluene dioxygenase. Biochemical and Biophysical research communications, Vol.185(1), 443-451.
- Löffler F.E., Sanford R.A. and Tiedje J.M. (1996) Initial characterization of a reductive dehalogenase from Desulfitobacterium chlororespirans Co23. *Applied and Environmental Microbiology*, Vol.62(10), 3809–3813.
- Maule A., Plyte S. and Quirk A.V. (1987) Dehalogenation of organochlorine insecticides by mixed anaerobic microbial populations. *Pesticide Biochemistry and Physiology*, Vol.27, 229-236.
- Maymó-Gatell X., Anguish T. and Zinder S.H. (1999) Reductive dechlorination of chlorinated ethenes and 1,2-dichloroethane by Dehalococcoides ethenogenes 195. *Applied and Environmental Microbiology*, Vol.65(7), 3108–3113.
- Miyoshi K., Kameyama Y. and Matsumura M. (2004) Electrochemical reduction of organohalogen compound by noble metal sintered electrode. *Chemosphere*, Vol.56, 187-193.
- Mochizuki S. (1977) Photochemical dechlorination of PCBs. *Chemical Engineering Science*, Vol.32, 1205-1210.
- Mohammad S. M. and Dennis G. P. (1997) Electrochemical reduction of di-, tri- and tetrahalobenzenes at carbon cathodes in dimethylformamide. Evidence for a halogen dance during the electrolysis of 1,2,4,5-tetrabromobenzene. *Journal of Electroanalytical Chemistry*, Vol.435, 47-53.
- Motosugi K., Esaki N. and Soda K. (1982) Purification and properties of a new enzyme, DL-2-haloacid dehalogenase, from Psurdomonas sp. *Journal of Bacteriology*, Vol.150, No.2, 522-527.
- Moza P. N., Fytianos K., Samanidou V. and Korte F. (1988) Photodecomposition of chlrophenols in aqueous medium in presence of hydrogen peroxide. *Bulletin of Environmental Contamination and toxicology*, Vol.41, 678-682.
- Nesbitt H.W. and Muir I.J. (1998) Oxidation states and speciation of secondary products on pyrite and arsenopyrite reacted with mine waste waters and air. *Mineralogy and petrology*, Vol.62, 123-144.
- Newman L.M. and Wackett L.P. (1991) Fate of 2,2,2-trichloroacetaldehyde (chloral hydrate) produced during trichloroethylene oxidation by methanotrophs. *Applied Environmental Microbiology*, vol.57 (8), 2399-2402.
- Newman L.M. and Wackett L.P. (1995) Purification and characterization of toluene 2monooxygenase from Burkhokderia cepacia G4. *Biochemistry*, Vol.34(43), 14066-14076.
 Newman L.M. and Wackett L.P. (1997) Journal of Bacteriology, No.179(1), 90-96.
- Oldenhuis R., Vink R.L., Janssen D.B., Witholt B. (1989) Degradation of chlorinated aliphatic
- hydrocarbons by methylosius trichosporium OB3b expressing soluble methane monooxygenase. *Applied Environmental Microbiology*, Vol.55(11), 2819-2826.
- Reineke W. and Knackmuss H.J. (1984) Microbial Metabolism of Haloaromatics: Isolation and properties of Chlorobenzene-Degrading Bacterium. *Applied and Environmental Microbiology*, Vol.47(2), 395-402.
- Rosenzweig A.C., Frederick C.A., Lippard S.J., Nordlund P. (1993) Crystal structure of bacterial non-haem iron hydroxylase that catalyses the biological oxidation of methane. *Nature*, Vol.366(6455), 537-43

- Rosso K.M., Becker U. and Hochella M. F. J. (1999) The interaction of pyrite{100} surface with O₂ and H₂O: Fundamental oxidation mechanisms. *American Mineralogist*, Vol.84, 1549-1561.
- Schaufuss A.G., Nesbitt H.W., Kartio I., Laajalehto K., Bancroft G.M. and Szargan R. (2000) Reactivity of surface chemical states on fractured pyrite. *Surface Science*, Vol.411, 321-328.
- Shimakoshi H., Tokunaga M. Baba T. and Hisaeda Y. (2004) Photochemical dechlorination of DDT catalyzed by a hydrophobic vitamin B12 and a photosensitizer under irradiation with visible light. *Chemical Communications*, 1806-1807.
- Sundstrom D.W., Klei H.E., Nalette T.A., Reidy D. J., Weir B. A. (1986) Destruction of Halogenated Aliphatics by Ultraviolet Catalyzed Oxidation with Hydrogen Peroxide. *Hazardous Waste and Hazardous Materials*, 3(1), 101-110.
- Sundstrom D.W., Weir B.A. and Klei H.E. (1989) Destruction of aromatic pollutions by UV light catalyzed oxidation with hydrogen peroxide. *Environmental Progress*, Vol.8 (1), 6-11.
- Taylor B. E., Wheeler M.C. and Nordstorm D.K. (1984) Stable isotope geochemistry of acid mine drainage: Experimental oxidation of pyrite. *Geochimica et Cosmochimica Acta*, Vol.48, 2669-2678.
- Tiedje, J. M., Quensen Ⅲ J. F., Chee-Sanford, J., Schimel J. P. and Boud S. A. (1993) Microbial reductive dechlorination of PCBs. *Biodegradation*, Vol.4, 231-240.
- Usher C.R., Cleveland C.A., Strongin D.R. and Schoonen M.A. (2004) Origin of oxygen in sulfate during pyrite oxidation with water and dissolved oxygen: An in situ horizontal attenuated total reflectance infrared spectroscopy isotope study. *Environmental Science and Technology*, Vol.38, 5604-5606.
- Usher C.R., Paul K.W., Narayansamy J., Kubicki J.D., Sparks D.L., Schoonen M.A.A. and Strongin D.R. (2005) Mechanistic aspects of pyrite oxidation in an oxidizing gaseous environment: An in situ HATR-IR isotope study. *Environmental Science and Technology*, Vol.39, 7576-7584.
- Van Beek H.C.A., Van der Stoep H.J., Van Oort H. and Van Leene J. (1982) Photochemical Radical Chain Dechlorination of DDT in 2-Propanol. *Industrial & Engineering Chemistry Product Research and Development*, Vol.12, 123-125.
- Weerasooriya R. and Dharmasena B. (2001) Pyrite-assisted degradation of trichloroethene (TCE). *Chemosphere*, Vol.42, 389-396.
- Weie B.A., Sundstrom D.W. and Klei H.E. (1987) Destruction of Benzene by Ultraviolet Light-Catalyzed oxidation with Hydrogen Peroxide. *Hazardous Waste and Hazardous Materials*, Vol.4(2), 165-176.
- Weightman A.J., Weightman A.L., Slater J.H. (1985) Toxic effects of chlorinated and brominated akanoic acids on Pseudomonas putida PP3: selection at high frequencies of mutations in genes encoding dehalogenases. *Applied Environmental microbiology*, Vol.49(6), 1494-1501.
- Weightman A.L., Weightman A.J. and Slater J.H. (1992) Microbial dehalogenation of trichloroacetic acid. *World journal of microbiology and biotechnology*, Vol.8, 512-518.
- Zinovyev S.S., Shinkova N.A., Perosa A. and Tundo P. (2005) Liquid phase hydrodechlorination of dieldrin and DDT over Pd/C and Raney-Ni. *Applied Catalysis B: Environemtal*, Vol.55, 39-8.



Organic Pollutants Ten Years After the Stockholm Convention -Environmental and Analytical Update Edited by Dr. Tomasz Puzyn

ISBN 978-953-307-917-2 Hard cover, 472 pages Publisher InTech Published online 24, February, 2012 Published in print edition February, 2012

Ten years after coming into force of the Stockholm Convention on Persistent Organic Pollutants (POPs), a wide range of organic chemicals (industrial formulations, plant protection products, pharmaceuticals and personal care products, etc.) still poses the highest priority environmental hazard. The broadening of knowledge of organic pollutants (OPs) environmental fate and effects, as well as the decontamination techniques, is accompanied by an increase in significance of certain pollution sources (e.g. sewage sludge and dredged sediments application, textile industry), associated with a potential generation of new dangers for humans and natural ecosystems. The present book addresses these aspects, especially in the light of Organic Pollutants risk assessment as well as the practical application of novel analytical methods and techniques for removing OPs from the environment. Providing analytical and environmental update, this contribution can be particularly valuable for engineers and environmental scientists.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Junko Hara (2012). Chemical Degradation of Chlorinated Organic Pollutants for In Situ Remediation and Evaluation of Natural Attenuation, Organic Pollutants Ten Years After the Stockholm Convention - Environmental and Analytical Update, Dr. Tomasz Puzyn (Ed.), ISBN: 978-953-307-917-2, InTech, Available from: http://www.intechopen.com/books/organic-pollutants-ten-years-after-the-stockholm-convention-environmental-and-analytical-update/chemical-degradation-of-chlorinated-organic-pollutants-for-in-situ-remediation-and-evaluation-of-nat



InTech Europe

University Campus STeP Ri Slavka Krautzeka 83/A 51000 Rijeka, Croatia Phone: +385 (51) 770 447 Fax: +385 (51) 686 166 www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai No.65, Yan An Road (West), Shanghai, 200040, China 中国上海市延安西路65号上海国际贵都大饭店办公楼405单元 Phone: +86-21-62489820 Fax: +86-21-62489821 © 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the <u>Creative Commons Attribution 3.0</u> <u>License</u>, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen