

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Role of NK Cells in Tuberculous Pleurisy as Innate Promoters of Local Type 1 Immunity with Potential Application on Differential Diagnosis

Pablo Schierloh, Silvia De La Barrera and Maria Sasiain
*Instituto de Investigaciones Hematologicas, Academia Nacional de Medicina
 Argentina*

1. Introduction

An efficient host immune response against pathogens encompasses both fast acting innate immunity as well as slower, but more specific, adaptive immunity. The innate immune system is diverse and comprises a variety of cells including natural killer (NK) cells, neutrophils, macrophages, dendritic cells (DCs), as well as soluble factors such as complement. The adaptive immune response is typified by antigen-specific T and B lymphocytes that provide long-lasting protection known as immunological memory. While these two systems are often discussed separately, neither arm of the immune system works in isolation (Medzhitov & Janeway, 1999). The succession of cells interacting with *Mycobacterium tuberculosis* (*Mtb*) comprises tissue macrophages (MΦ) and dendritic cells (DC) followed by chemokine-attracted immigrating neutrophils and monocytes, and then activation and recruitment of natural killer (NK) and $\gamma\delta$ T cells, followed by effector T lymphocytes primed in the draining lymph nodes (Ulrichs & Kaufmann, 2006).

Pleuritis is the most frequent clinical manifestation of extrapulmonary tuberculosis (TB) among young adults, and is normally considered a relatively benign form of disease since it may resolve without chemotherapy (Light, 2010). Tuberculous pleurisy is caused by a severe delayed-type hypersensitivity reaction in response to the rupture of a subpleural focus of *Mtb* infection, but it may also be developed as a complication of primary pulmonary TB infection (Antoniskis et al., 1990). The presence of mycobacterial antigens in the pleural space elicits an intense cellular immune response, initially characterized by abundant neutrophils and macrophages, followed by interferon (IFN)- γ -producing T-helper cell (TH) type 1 lymphocytes, resulting in lymphocyte-predominant exudative effusions (Aleman et al., 2005; Mitra et al., 2005; Porcel, 2009). The cellular trafficking is facilitated by homing surface markers and chemokine gradients. This intense but poorly understood local immune response is synonymous of Koch phenomenon and normally prevents the caseous evolution of lesions. The inflammatory process results in an increased pleural vascular permeability leading to the accumulation of fluid enriched in proteins and the recruitment of specific leukocytes into the pleural space, making this biological sample a physiologically relevant model of human tuberculosis infection (Kroegel & Antony 1997).

Given this effective local resistance, the few bacilli that enter into pleural cavity are rapidly destroyed turning impracticable the rapid diagnosis by direct microscopic observation (Porcel, 2009). This fact often results in the requirement of alternative diagnostics strategies like *Mtb* identification through cultures or PCR amplification of pleural effusions, histopathological examination of pleural biopsies, inflammatory related enzymatic activities and immunological based methods (Liang et al., 2008; Trajman et al., 2008). Among the last ones, detection of mycobacterial antigens, antimycobacterial specific antibodies (Ab), TH1 related biomarkers and *in vitro* evoked T cell responses are matter of active current research (Steingart et al., 2007; Budak et al., 2008; Dheda et al., 2009a; Supriya et al., 2008). Regarding the performance of T and B cell based assays in detection of active forms of disease; the most important trouble is the immunological memory background as result of previous Ag exposition or BCG vaccination (Dheda et al., 2009b; Hooper et al., 2009; Salazar-Lezama et al., 1997). In this line, for high antigen experienced populations, we hypothesized that an innate immune cell based diagnostic assay may circumvent those issues.

Among innate lymphocytes, Natural Killer (NK) cells display an important number of effector functions, including recognition and lysis of infected, stressed, or transformed cells and production of immunoregulatory cytokines, particularly IFN- γ (Vivier et al., 2011). Human NK cells account for 10–20% of peripheral blood lymphocytes and are defined by the presence of the CD56 and NKp46 molecules and the lack of CD3 and CD19 expression. Their activity is regulated by both positive and inhibitory signals from a wide range of germ line encoded cell surface receptors. Two major subsets of NK cells have been identified in humans according to CD56 and CD16 intensity expression and also in terms of chemokine receptors and adhesion molecules expression that differ in phenotype and function (Caligiuri, 2008). Functionally, CD56^{bright} cells are effective cytokine producers, whereas CD56^{dim} cells are efficient effectors of natural and antibody-dependent target cell lysis (Hanna & Mandelboim 2007). Together with the classical NK functions (i.e. cytotoxicity and cytokine production), novel skills have recently been described in niche-specific and *in vitro*-activated human NK cells. These unconventional capabilities include angiogenesis and tissue remodeling, immunological memory, functional cross-talk with T cells and direct pathogen recognition (Cooper et al., 2009; Di Santo, 2008; Vivier et al., 2011).

During the last years we have begun to characterize the phenotype and function of pleural NK from TB patients. In contrast to peripheral blood (PB) counterpart that are mostly composed by CD56^{dim}CD16⁺ resting cells, pleural NK population is enriched in activated CD56^{bright}CD16^{neg} cells that quickly and strongly respond to *Mtb* stimulation by producing IFN- γ (Schierloh et al., 2005a, 2007). Besides, *Mtb* stimulated IFN- γ production by NK shows Ag-specific features, given that pleural NK cells derived from non TB patients' lack of this response. According to these findings, we realized that pleural NK cells properties could be utilized in an innate immune cell based assay for differential diagnosis of tuberculous pleurisy (Schierloh et al., 2008).

2. Phenotype of pleural NK cells: Activated CD56^{bright} NK cells

Along several studies, phenotype of tuberculous pleural fluid derived NK cells have been extensively analyzed as starting point in the understanding of their role at the site of active *Mtb* infection (Alvarez et al., 2010; Fu et al., 2011; Okubo et al., 1986, 1987; Pokkali et al.,

2009; Schierloh et al., 2005a, 2007, 2009). The finding that immunoregulatory CD56^{bright} NK subset showing cellular activation features are strongly enriched among pleural NK cells was interesting given that these cells are known to link innate with adaptative immunity in a number of intracellular infections (Artavanis-Tsakonas et al., 2003; Culley, 2009; Fehniger et al., 2003). Along the present section we summarize these findings.

2.1 Enrichment of CD56^{bright} NK cells in tuberculous pleural fluid

In tuberculous patients, T helper (CD4⁺/CD3⁺) is the predominant cell population among pleural fluid derived mononuclear cells (PFMC), which has been associated with a selective recruitment of antigen specific TH1 effectors cells to the site of infection (Li et al., 2010; Mitra et al., 2005). This increased T helper abundance account for the reduced percentage of pleural fluid NK cells (CD3⁺/CD56⁺) close to 5% in TB patients (Schierloh et al., 2005a). In contrast, in pleural effusions caused by cancer or paraneumonic infections, the percentage of NK among PFMC tends to be constant compared to peripheral blood (Dalbeth et al., 2004). However, when we analyzed the composition of NK cell populations, a drastic change in the proportion of these subsets was found (Schierloh et al., 2005a). In the case of tuberculosis, the cytotoxic CD16⁺CD56^{dim} NK subset, which represents more than 95% in the circulation, is reduced to less than 50% in the pleural effusions with a concomitant enhancement of the immunoregulatory CD16^{dim}/CD56^{bright} NK subset.

Phenotypic differences between CD56^{bright} NK with CD56^{dim} include higher expression of the C-type lectin CD94/NKG2 family, weak expression of killer cell immunoglobulin (Ig)-like receptors (KIRs) and high levels of L-selectin (CD62L) and CCR7, both of which are involved in trafficking of immune cells to lymph nodes (Caligiuri, 2008). In agreement, the levels of CD94/NKG2A, CD62L and CCR7 were all augmented in tuberculous pleural NK cells (Schierloh et al., 2005a). On the other hand, the percentages of NK cells expressing the fractalkine receptor (CX₃CR1), the cytotoxic granular protein perforin and the HLA-C2 receptor KIR2DL1/S1, which are all specific markers for the cytotoxic CD16⁺CD56^{dim} NK subset, are reduced in the tuberculous pleural effusions (Figure 1).

2.2 Pleural NK cells exhibit activated phenotype

NK cells activation can be triggered via two primary mechanisms: cytokine stimulation and engagement of activating NK receptors. Together or in isolation, both activation signals can result in NK cell responses (Vivier et al. 2011). Because tuberculous pleural microenvironment is plenty of soluble mediators and cells with stimulatory potential (Shimokata et al., 1991; Valdés et al., 2009; Vankayalapati et al., 2000), we hypothesized that NK cells arriving to this site may turn activated. Indeed, we found an elevated percentage of NK cells expressing the early activation markers CD69 and HLA-DR together with enhanced expression of the lymphocyte function-associated antigen 1a integrin (LFA-1/CD11a) and its ligand, intercellular adhesion molecule-1 (ICAM-1, CD54) (Schierloh et al., 2005a, 2005b, 2009). Simultaneously, we observed a subpopulation of pleural NK cells expressing Toll-like Receptor 2 (TLR2), a molecule undetectable in resting NK cells but up-regulated in response to IL-12 and protozoan glycolipids (Becker et al., 2003; Lindgren et al., 2010; Schierloh et al., 2007). Furthermore, we and others recently identified a subpopulation of pleural NK that down-modulates the CD45RA and up-regulates CD45R0 isoform,

resembling the well known phenomenon that takes places during memory differentiation of T lymphocytes (Warren et al., 1994; Fu et al., 2011).




Fig. 1. Enrichment of NK CD56^{bright} cells with activated phenotype in tuberculous pleural effusions.

Immunophenotypic analysis of peripheral blood (PB) and pleural fluid (PF) NK cells (CD56⁺/CD3⁻). Dot plots are from one representative TB patient. Color scale indicates relative variations of several markers between PF and PB samples: invariant (white), augment (orange), strong augment (red), reduction (light green) and strong reduction (dark green).

2.3 Pleural fluid factors inducing CD56^{dim} apoptosis explain the altered NK subset ratio

In order to understand the causes of altered CD56^{bright}/CD56^{dim} subset ratio observed on pleural NK population, we hypothesize that soluble factors (i.e.: chemokines, cytokines, pathogen derived factors, immune complexes, etc.), present at the site of *Mtb* infection, may differentially affect the migration, the differentiation, the proliferation or the apoptosis of NK subsets. Our experimental approaches directed to test this were conducted by incubating peripheral resting NK cells with tuberculous cell-free-pleural fluid or purified factors (Schierloh et al., 2005a). Indeed, CD56^{dim}CD16⁺ cells show an increased susceptibility to pleural fluid induced caspase 9 dependent-apoptosis, explaining the predominance of the CD56^{bright} population. These finding were later confirmed in different experimental settings. These studies demonstrate that NK CD56^{bright} subset has larger resistance to oxidative stress (Harlin et al., 2007; Thorén et al., 2007). Our experiments directed to test if NK CD16⁺ cells could be differentiated to CD16⁻ NK cells by cytokines present at the site of infection gave negative results, in accordance with other group (Dalbeth et al., 2004). Additionally, a recent

report indicates that differential migration of NK cell subsets to the site of infection also take place (Pokkali et al., 2009).

3. Function of pleural NK cells: IFN- γ production and TH1 cell co-stimulation

Considering that type 1 cytokine and chemokine profile is a hallmark of tuberculous pleurisy (Dheda et al., 2009a; Kroegel & Antony 1997; Li et al., 2010; Mayanja-Kizza et al., 2009; Trajman et al., 2008), and given that pleural NK cells exhibit an endogenously induced activation state together with an enrichment of immunoregulatory NK CD56^{bright} subset, we asked whether these cells were polarized to the production of pro and/or anti-inflammatory cytokines. Also, we evaluated if these cells were capable to modulate other cells function. The answers to these questions gave us important clues for better understanding the immunopathogenesis of TB infection and will be discussed during the present section.




Fig. 2. Pleural NK cells are a major early source of IFN- γ upon ex vivo *Mtb*-stimulation.

PFMC were stimulated with *Mtb* for 24h. Then cells were gated on the basis of IFN- γ -positive (Ly IFN- γ ⁺) and IFN- γ -negative (Ly IFN- γ ^{neg}) cells and their surface phenotype were determined according to CD56 and CD3 expression by flow cytometry. Dot plots are from one representative TB patient. NK cells are highlighted in red. Pie charts indicate mean values of 30 TB patients.

3.1 Pleural NK cells strongly produce IFN- γ after *Mtb* stimulation

Experiments varying the quality and quantity of signaling input for NK cell activation have revealed a hierarchy in requirements for induction of chemokines and cytokines (Fauriat et al., 2010). Furthermore, recent data indicate that anatomical niche and developmental stage of NK cells strongly determine its cytokine profile production giving rise to NK1, NK2 and

NK22 cells (Di Santo, 2008; Spits & Di Santo, 2011). In this sense, pleural fluid CD56^{bright} NK did not produce significant levels of IL-10, IL-17A or TNF- α spontaneously or after γ -irradiated *Mtb*-stimulation. However, the percentage of IFN- γ ⁺ NK cells (NK IFN- γ ⁺) was strongly increased under the same experimental conditions (Schierloh et al., 2005a). It is interesting to note that, in spite of its reduced numbers in TB pleural fluid (~5% of PF lymphocytes), NK constitute a major source of IFN- γ together with CD3⁺ cells (Figure 2). *Mtb* derived culture filtrate proteins (CFP), TLR2 and 4 agonist and recombinant IL-12 also induced IFN- γ among pleural NK but to a lesser extent than *Mtb* does (Schierloh et al., 2007). Interestingly, pleural fluid NK cells derived from other etiologies (i.e: cancer, paraneumonic or helminthic infections) did not give *Mtb*-stimulated IFN- γ responses, suggesting a paradoxical “Ag specific response” induced in an innate immune cell (see Figure 4). This finding is in accordance with recent data provided by experimental mice models which clearly demonstrate the adaptative properties of NK cells (Cooper et al., 2009; Paust & von Andrian, 2011; Vivier et al., 2011). Instead, this apparent “Ag specific” NK cell responses may be produced by T cell-secreted IL-2 (Horowitz et al., 2010; Fehniger et al., 2003).

3.2 Pleural NK cells co-stimulate local TH1 response

Among novel skills described for human activated NK cells, it has been shown that they may stimulate T cells by cell contact-dependent mechanisms (Hanna & Mandelboim 2007). In this line, we were able to demonstrate that, in TB pleurisy, a functional ICAM-1-dependent cell to cell interaction among pleural fluid NK and T cells lead to T cell activation (Schierloh et al 2009). Likewise, peripheral blood human NK cells can instruct *in vitro* cytotoxic CD8⁺ T cells from PPD responsive donors to lyse *Mtb*-infected monocytes (Vankayalapati et al., 2004). Taken together, these findings suggest a previously unappreciated role of NK cells in the maintenance and/or activation of T cell functions during the immune response in tuberculosis.

4. Cellular and molecular factors controlling pleural NK cells functions

Having observed that NK cells were the main early source of IFN- γ within the pleural space, we investigate extracellular events and signaling pathways that drive this process. The mechanisms involved reveal classical and particular ways of NK cell activation and signaling. Three environmental signals act in concert to fulfill IFN- γ response in pleural NK cells: cytokines, activation ligands expressed on accessory cells and direct *Mtb* recognition (Schierloh et al., 2007).

4.1 Pleural NK response involve Ca²⁺ influx and Calcineurin, ERK and p38 MAPK signaling pathways

NK cells express on their surface an array of germ-line encoded inhibitory and activating receptor as well as cytokine receptors that, upon activation, mediate intracellular signaling pathways for IFN- γ together with other functional responses. It is well known that interaction of NK receptors cells with their activating ligands on target and/or Ag presenting cells (APC) induce a quick increase of cytoplasmic calcium (Ca²⁺) concentration, a universal second messenger (Maghazachi, 2005). Employing divalent cation chelators we

demonstrate a pivotal role for this process during *Mtb*-induced pleural NK IFN- γ response. Downstream the Ca^{2+} influx, signaling proteins and their target transcription factors are activated, including calcineurin, a calmodulin-dependent serine/threonine phosphatase, and its target NFAT (nuclear factor of activated T cells). By mean of Cyclosporin A treatment, we confirm the involvement of this pathway too. Similarly, using several protein kinase specific inhibitors and phospho-specific monoclonal antibodies, we also address the participation of other two phosphorylation cascades, p38 MAPK and ERK1/2 (Schierloh et al., 2007). Figure 3 summarize the most important events.




Fig. 3. Extracellular signals and intracellular pathways in IFN- γ production by pleural NK cells.

Picture describe interactions among activation signals (i.e: cytokines, APC expressed activating ligands and *Mtb*), modulating/inhibitory signals (i.e: APC expressed inhibitory

ligands and *Mtb* derived lipoglycans recognized by C-type lectins) and cellular receptors together with their corresponding intracellular signaling pathways. Blue arrows indicate positive interactions. Dotted blue arrows denote cross-talk between pathways. Black arrows denote inhibitory interactions. Pharmacological inhibitors used in our experiments are written in red within brackets under its target molecule.

4.2 Pleural NK response is dependent on IL-12 and accessory cell contact

Numerous *in vitro* and *in vivo* studies show that a wide variety of APC secreted cytokines, especially IL-12, IL-15, and IL-18, can activate NK cells to produce IFN- γ . By means of cytokine neutralization, we showed that IL-12 is necessary for *Mtb*-induced NK IFN- γ production in tuberculous pleurisy (Schierloh et al., 2007). This requirement is not surprising because IL-12 has been shown to mediate bystander activation of NK cells in response to a number of different pathogens (Artavanis-Tsakonas et al., 2007; Culley, 2009; Lindgren et al., 2011). However, IL-12 alone does not allow the same level of IFN- γ ⁺ NK cells than *Mtb* does. Indeed, costimulatory signals delivered by APC such as receptor-ligand interactions are required as demonstrated by Ab blockade and APC depletion experiments. Among activating ligands ICAM-1, CD86 and Vimentin has been shown to be engaged. At the same time, these signals could be counter-balanced by MHC class I, PD-L1 and PD-L2 inhibitory ligands (Alvarez et al., 2010; Garg et al., 2006; Schierloh et al., 2007).

At the level of pleural APC, we also observed that *Mtb* induced stimulatory signals mediated by TLR2 and TLR4 are counter-modulated by C-type lectin receptors like mannose receptor (MR) and DC-SIGN (Schierloh et al., 2007). Both molecules bind mycobacterial derived mannosylated lipoglycans and may indirectly inhibit NK cell response by limiting IL-12 production and/or down-modulating activating ligands expression (van Kooyk Y & Geijtenbeek, 2003).

4.3 Pleural NK cells directly recognize *Mtb*

Recent studies have pointed out the capacity of NK cells to bind and been directly activated by *Mycobacterium* species (Esin et al., 2004; Evans et al., 2011; Watkins et al., 2008). This recognition seems to be mediated, at least, by two putative activating receptors: TLR2 and NKp44 (Esin et al., 2008; Marcenaro et al., 2008). Interestingly, these receptors are both up-regulated among pleural NK cells (Figure 1). Consistently we observe that, compared with its PB counterparts, pleural NK cells have enhanced capacity to bind *Mtb* and that p38 MAPK phosphorylation on pleural NK occurs shortly after *Mtb*-NK coculture, independently on bystander cell derived signals (Schierloh et al., 2007). Furthermore, it has been observed that NK CD56^{bright} cells are more reactive to direct BCG stimulation than CD56^{dim} NK (Batoni et al., 2005). Altogether, these results strongly indicate that direct interaction between NK and *Mtb* play a significant role during functional response of pleural NK CD56^{bright} cells.

5. Pleural NK application: Immunodiagnosis of tuberculous pleurisy

Conventional diagnostic tests for pleural TB include microscopic examination of pleural fluid for acid-fast bacilli and differential cytology; mycobacterial culture of pleural fluid, sputum or pleural tissue; pleural fluid Adenosin Deaminase (ADA) activity as well as

determination and histopathological examination of pleural tissue looking for granulomatous inflammation (Porcel, 2009; Light, 2010). These tests have limitations for clinical use; however, in combination, they have been recognized as the best reference standard for evaluation of the accuracy of novel tests (Trajman et al., 2008). Although detection of serum antibodies against *Mtb* antigens is known to have poor and highly variable sensitivity and specificity, attempts have been made to detect antibodies in pleural fluid by ELISA. Even though, these tests show high specificity they are limited by the very poor sensitivity (Weldingh & Pai 2007). Several T cell based assays have been employed for diagnosis of TB pleurisy such as in vitro stimulation of lymphocytes with PPD or RD-1 encoded antigens leading to T-cell proliferation and/or IFN- γ release by ELISPOT and ELISA assays (TIGRAs) (Hooper et al., 2009). For example, using a commercially available *Mtb*-specific ELISPOT for peripheral blood mononuclear cells and pleural fluid mononuclear cells from patients with exudative pleurisy, its sensitivity in active tuberculosis was very high (95%); however, the specificity was suboptimal (76%) (Losi et al., 2007). The high coverage of BCG vaccination as well as the high prevalence of latent TB infection (PPD+ individuals) at the population level might impair the results when employing T and B cell based immunodiagnostic methods. On the other hand, immunocompromised or HIV+ infected patients may lead to false negative results due to the ablation or reduction of cellular immunity (Trajman et al., 2008). In this context, we thought that pleural IFN- γ + NK cells could be a promising target for immunodiagnostic method that circumvent these memory related problems in differential diagnosis of tuberculous pleurisy.

5.1 Preliminary trial for testing clinical value of NK cell based assay in Argentina

In order to provide evidence that support or reject the clinical diagnostic utility of NK IFN- γ + cell based assay in differential diagnosis of TB pleural effusion, we have performed a retrospective, single-center preliminary study in a reference center of the city of Buenos Aires, Argentina. To do this, PFMFC were stimulated with *Mtb* and the percentage of NK and T cells expressing IFN- γ were determined by flow cytometry as described in Figure 2. In this trial we included 40 consecutive patients with profuse exudative pleural effusion (TB n=28 and No-TB n=12; Cancer=6, Paraneumonic infection=5, Helmintic infection=1) admitted and diagnosed at the Tisioneumonolgy service of the Hospital Muñiz during 2006-2009. According to epidemiological data provided by medical staff, all the patients were at high relative risk of TB infection. As can be observed in table 1 and Figure 4, pleural NK shows

	Median Δ % IFN- γ + (25–75% percentil)	Cut off (Max. likelihood ratio)	Area under ROCurve (95% CI)	Sensitivity (95% CI)	Specificity (95% CI)
NK assay	30.69% (9.23–42.77%)	> 1.555% (11.57)	0.9643 (0.9002–1.028)	96.43% (81.65–99.91%)	91.67% (61.52–99.79%)
T assay	2.14% (1.04– 4.63%)	> 1.230% (8.14)	0.9435 (0.8670–1.020)	67.86% (47.65–84.12%)	91.67% (61.52–99.79%)

Table 1. ROC analysis for NK cell and T cell assays. Data values summarized above are derived from ROC analysis depicted in Fig.4.

better performance than pleural T cell based assay in discriminating TB and no-TB pleural effusions. These differences may reflect the polyclonal vs oligoclonal nature of IFN- γ response in NK compared to T cells (Schierloh et al., 2008).




Fig. 4. Mtb-induced IFN- γ response by NK and T cells in the diagnosis of tuberculous pleurisy.

PFMC were incubated 24h with medium alone (Control) or stimulated with *Mtb*. Then, *Mtb* specific IFN- γ response was obtained from gated NK (CD56 $^+$ /CD3 $^-$) and T cells (CD3 $^+$) by subtracting the percentage of spontaneous production: $\Delta \% \text{IFN-}\gamma^+ = \% \text{IFN-}\gamma^+ \text{ Mtb} - \% \text{IFN-}\gamma^+ \text{ control}$. Upper left scatter plots depict the results of the study population (TB patients n=28; No-TB patients n=12). Lower graph present ROC analysis for NK and T cell based assays. Flow cytometry dot plots analyses are from one representative TB patient and one patient with helminthic (*Echinococcus granulosus*) infection (No TB). Red dots are gated NK (CD56 $^+$ /CD3 $^-$) and purple dots are T cells (CD3 $^+$).

5.2 Rational design for NK based IFN- γ release assay: “KIGRA”

Flow cytometers are sophisticated equipments that may be not available in almost all public health laboratories. Therefore, in order to make our NK derived IFN- γ based assay more applicable for common clinical settings, we attempt to introduce experimental modifications that could direct the development of a NK based IFN- γ release assay.

Unlike flow cytometry, cytokine release detection devises did not allow the identification of secreting cell. This fact represents an important challenge in our particular case, where the cellular source of IFN- γ is the basis of the diagnostic improvement. One way to avoid this issue is blocking IFN- γ production by pleural T cells, leaving NK as the main producer cells. To do so, we employed anti-HLA class I and class II monoclonal antibodies (mAbs), which block Ag presentation to CD8 $^{+}$ and CD4 $^{+}$ T cells. As can be observed in Figure 5, most T IFN- γ^{+} cells were inhibited when both mAbs were present during *Mtb* stimulation; however, NK cells still remain expressing IFN- γ^{+} under the same treatment. Similarly, when we analyzed IFN- γ release by ELISA, the presence of mAbs diminished but not abolished the secretion of this biomarker. This result, together with other adjustments that are under current testing, may constitute the rational for a more accurate assay.

Furthermore, we think that in the context of diagnosis of tuberculous pleurisy, commercially available IGRA, which employ ELISPOT (Losi et al., 2007), ELISA (Losi et al., 2011) or immunochromatography (Corstjens et al., 2008), could be easily adapted in order to obtain a Natural Killer IFN-gamma release assay or “KIGRA”.




Fig. 5. Rational design for development of NK IFN- γ release assay.

PFMC were *Mtb*-stimulated as indicated in Fig.4 with or without the addition of anti-HLA-class I and/or anti-HLA-class DR mAbs. A schematic explanation of how mAbs interfere on antigen presentation to CD4⁺ and CD8⁺ T $\alpha\beta$ cells without affecting NK or T $\gamma\delta$ cells activation (upper cartoon). Flow cytometry dot plots analyses are from one representative TB patient. Red dots are gated NK (CD56⁺/CD3⁻) and purple dots are T (CD3⁺) cells. Bar graph of IFN- γ ELISA assay (TB n=3).

6. Conclusion

Tuberculous pleurisy, one of the most common extrapulmonary manifestations of tuberculosis among young adults, is characterized by strong delayed type hypersensitivity reaction mediated by effector lymphocytes. Our data demonstrate that a substantial part of these cells are indeed NK cells. Herein, we have discussed phenotypic and functional features of this local innate immune cell population and the factors involved in its regulation. Furthermore, we identify NK cells as the main source of IFN- γ , the most widely used TB biomarker, in the context of tuberculous pleurisy.

In the context of high antigen experienced population the diagnosis that allow the discrimination between tuberculous pleurisy from other exudative pleural effusions remain as an unresolved clinical issue (Dheda et al., 2009b; Hooper et al., 2009; Salazar-Lezama et al., 1997). Hence, we propose an NK cell IFN- γ based assay as complementary procedure. To our knowledge, no previous NK immunodiagnostic were reported for TB or any other infectious diseases. The advantageous characteristics of this functional assay are: (i) short time result output (1 day) and ii) very good performance in terms of specificity and sensitivity.

7. Acknowledgement

All authors are members of Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET) scientific staff. M.C.S, S. de la B and P.S are supported by grants from Agencia Nacional de Promoción Científica y Tecnológica (ANPCyT; PAE-TBC PICTS 2328 and 2329). P.S is beneficiary of grant from Roemmers Foundation.

The authors thank the medical staff of Division de Tisiopneumología, Hospital F. J. Muñoz, especially to Dr R.M. Musella and Dr. J. Castagnino for their great help in providing samples, clinical data and useful discussion.

8. References


- Aleman M, de la Barrera SS, Schierloh PL, Alves L, Yokobori N, Baldini M, Abbate E, Sasiain MC. (2005). In tuberculous pleural effusions, activated neutrophils undergo apoptosis and acquire a dendritic cell-like phenotype. *J Infect Dis.* 192(3):399-409.
- Alvarez IB, Pasquinelli V, Jurado JO, Abbate E, Musella RM, de la Barrera SS, García VE. (2010). Role played by the programmed death-1-programmed death ligand pathway during innate immunity against Mycobacterium tuberculosis. *J Infect Dis.* 202(4):524-32.
- Antoniskis D, Amin K, Barnes PF. (1990). Pleuritis as a manifestation of reactivation tuberculosis. *Am J Med.* 89(4):447-50.

- Artavanis-Tsakonas K, Eleme K, McQueen KL, Cheng NW, Parham P, Davis DM, Riley EM. (2003). Activation of a subset of human NK cells upon contact with *Plasmodium falciparum*-infected erythrocytes. *J Immunol.* 171(10):5396-405.
- Batoni G, Esin S, Favilli F, Pardini M, Bottai D, Maisetta G, Florio W, Campa M. (2005). Human CD56^{bright} and CD56^{dim} natural killer cell subsets respond differentially to direct stimulation with *Mycobacterium bovis* bacillus Calmette-Guérin. *Scand J Immunol.* 62(6):498-506.
- Becker I, Salaiza N, Aguirre M, Delgado J, Carrillo-Carrasco N, Kobeh LG, Ruiz A, Cervantes R, Torres AP, Cabrera N, González A, Maldonado C, Isibasi A. (2003). *Leishmania* lipophosphoglycan (LPG) activates NK cells through toll-like receptor-2. *Mol Biochem Parasitol.* 130(2):65-74.
- Budak F, Uzaslan EK, Cangür S, Göral G, Oral HB. (2008). Increased pleural soluble fas ligand (sFasL) levels in tuberculosis pleurisy and its relation with T-helper type 1 cytokines. *Lung.* 186(5):337-43.
- Caligiuri MA. (2008). Human natural killer cells. *Blood.* 112(3):461-9.
- Cooper MA, Colonna M, Yokoyama WM. 2009. Hidden talents of natural killers: NK cells in innate and adaptive immunity. *EMBO Rep.* 10(10):1103-10.
- Corstjens PL, Zuiderwijk M, Tanke HJ, van der Ploeg-van Schip JJ, Ottenhoff TH, Geluk A. (2008). A user-friendly, highly sensitive assay to detect the IFN-gamma secretion by T cells. *Clin Biochem.* 41(6):440-4.
- Culley FJ. 2009. Natural killer cells in infection and inflammation of the lung. *Immunology.* 128, 151-163.
- Dalbeth N, Gundle R, Davies RJ, Lee YC, McMichael AJ, Callan MF. (2004). CD56^{bright} NK cells are enriched at inflammatory sites and can engage with monocytes in a reciprocal program of activation. *J Immunol.* 173(10):6418-26.
- Di Santo JP. 2008 Natural killer cells: diversity in search of a niche. *Nat Immunol.* 9(5):473-5.
- Dheda K, van Zyl-Smit RN, Sechi LA, Badri M, Meldau R, Meldau S, Symons G, Semple PL, Maredza A, Dawson R, et al. (2009a). Utility of quantitative T-cell responses versus unstimulated interferon-gamma for the diagnosis of pleural tuberculosis. *Eur Respir J.* 34(5):1118-26.
- Dheda K, Van-Zyl Smit RN, Sechi LA, Badri M, Meldau R, Symons G, Khalfey H, Carr I, Maredza A, Dawson R, Wainright H, Whitelaw A, Bateman ED, Zumla A. (2009b). Clinical diagnostic utility of IP-10 and LAM antigen levels for the diagnosis of tuberculous pleural effusions in a high burden setting. *PLoS One.* 4(3):e4689.
- Esin S, Batoni G, Pardini M, Favilli F, Bottai D, Maisetta G, Florio W, Vanacore R, Wigzell H, Campa M. (2004). Functional characterization of human natural killer cells responding to *Mycobacterium bovis* bacille Calmette-Guérin. *Immunology.* 112(1):143-52.
- Esin S, Batoni G, Counoupas C, Stringaro A, Brancatisano FL, Colone M, Maisetta G, Florio W, et al. (2008). Direct binding of human NK cell natural cytotoxicity receptor NKp44 to the surfaces of mycobacteria and other bacteria. *Infect Immun.* 76(4):1719-27.
- Evans JH, Horowitz A, Mehrabi M, Wise EL, Pease JE, Riley EM, Davis DM. (2011). A distinct subset of human NK cells expressing HLA-DR expand in response to IL-2 and can aid immune responses to BCG. *Eur J Immunol.* 41(7):1924-33.

- Fauriat C, Long EO, Ljunggren HG, Bryceson YT. (2010). Regulation of human NK-cell cytokine and chemokine production by target cell recognition. *Blood*. 115(11):2167-76.
- Fehniger TA, Cooper MA, Nuovo GJ, Cella M, Facchetti F, Colonna M, Caligiuri MA. (2003) CD56^{bright} natural killer cells are present in human lymph nodes and are activated by T cell-derived IL-2: a potential new link between adaptive and innate immunity. *Blood*. 101(8):3052-7.
- Fu X, Liu Y, Li L, Li Q, Qiao D, Wang H, Lao S, Fan Y, Wu C. (2011). Human natural killer cells expressing the memory-associated marker CD45RO from tuberculous pleurisy respond more strongly and rapidly than CD45RO(-) natural killer cells following stimulation with interleukin-12. *Immunology*. doi: 10.1111/j.1365-2567.2011.03464.x.
- Garg A, Barnes PF, Porgador A, Roy S, Wu S, Nanda JS, Griffith DE, Girard WM, Rawal N, Shetty S, Vankayalapati R. (2006). Vimentin expressed on Mycobacterium tuberculosis-infected human monocytes is involved in binding to the NKp46 receptor. *J Immunol*. 177(9):6192-8.
- Hanna J, Mandelboim O. (2007). When killers become helpers. *Trends Immunol*. 28(5):201-6.
- Harlin H, Hanson M, Johansson CC, Sakurai D, Poschke I, Norell H, Malmberg KJ, Kiessling R. (2007) The CD16- CD56^{bright} NK cell subset is resistant to reactive oxygen species produced by activated granulocytes and has higher antioxidative capacity than the CD16⁺CD56^{dim} subset. *J Immunol*. 179(7):4513-9.
- Hooper CE, Lee YC, Maskell NA. (2009). Interferon-gamma release assays for the diagnosis of TB pleural effusions: hype or real hope? *Curr Opin Pulm Med*. 15(4):358-65.
- Horowitz A, Behrens RH, Okell L, Fooks AR, Riley EM. (2010). NK cells as effectors of acquired immune responses: effector CD4⁺ T cell-dependent activation of NK cells following vaccination. *J Immunol*. 185(5):2808-18.
- Kroegel C, Antony VB. (1997). Immunobiology of pleural inflammation: potential implications for pathogenesis, diagnosis and therapy. *Eur Respir J*. 10(10):2411-8.
- Liang QL, Shi HZ, Wang K, Qin SM, Qin XJ. (2008). Diagnostic accuracy of adenosine deaminase in tuberculous pleurisy: a meta-analysis. *Respir Med*. 102(5):744-54.
- Li L, Qiao D, Fu X, Lao S, Zhang X, Wu C. (2011). Identification of Mycobacterium tuberculosis-specific Th1, Th17 and Th22 cells using the expression of CD40L in tuberculous pleurisy. *PLoS One*. 6(5):e20165.
- Light RW. (2010). Update on tuberculous pleural effusion. *Respirology*. 15(3):451-8.
- Lindgren A, Pavlovic V, Flach CF, Sjöling A, Lundin S. (2011). Interferon-gamma secretion is induced in IL-12 stimulated human NK cells by recognition of Helicobacter pylori or TLR2 ligands. *Innate Immunol*. 17(2):191-203.
- Losi M, Bossink A, Codecasa L, Jafari C, Ernst M, Thijsen S, Cirillo D, Ferrarese M, Greinert U, Fabbri LM, Richeldi L, Lange C; European Tuberculosis Network TBNET. (2007). Use of a T-cell interferon-gamma release assay for the diagnosis of tuberculous pleurisy. *Eur Respir J*. 30(6):1173-9.
- Losi M, Bocchino M, Matarese A, Bellofiore B, Roversi P, Rumpianesi F, Alma MG, Chiaradonna P, Del Giovane C, Altieri AM, Richeldi L, Sanduzzi A. (2011). Role of the quantiferon-TB test in ruling out pleural tuberculosis: a multi-centre study. *Int J Immunopathol Pharmacol*. 24(1):159-65.
- Marcenaro E, Ferranti B, Falco M, Moretta L, Moretta A. (2008). Human NK cells directly recognize Mycobacterium bovis via TLR2 and acquire the ability to kill monocyte-derived DC. *Int Immunol*. 20(9):1155-67.

- Maghazachi AA. (2005). Insights into seven and single transmembrane-spanning domain receptors and their signaling pathways in human natural killer cells. *Pharmacol Rev.* 57(3):339-57.
- Mayanja-Kizza H, Wu M, Aung H, Liu S, Luzze H, Hirsch C, Toossi Z. (2009). The interaction of monocyte chemoattractant protein-1 and tumour necrosis factor- α in Mycobacterium tuberculosis-induced HIV-1 replication at sites of active tuberculosis. *Scand J Immunol.* 69(6):516-20.
- Medzhitov R, Janeway CA Jr. (1999). Innate immune induction of the adaptive immune response. *Cold Spring Harb Symp Quant Biol.* 64:429-35.
- Mitra DK, Sharma SK, Dinda AK, Bindra MS, Madan B, Ghosh B. (2005). Polarized helper T cells in tubercular pleural effusion: phenotypic identity and selective recruitment. *Eur J Immunol.* 35(8):2367-75.
- Okubo Y, Nakata M, Kuroiwa Y, Wada S, Kusama S. (1986). Phenotypic and functional study of natural killer cells in tuberculous pleurisy patients. *Jpn J Med.* 25(3):250-6.
- Okubo Y, Nakata M, Kuroiwa Y, Wada S, Kusama S. (1987). NK cells in carcinomatous and tuberculous pleurisy. Phenotypic and functional analyses of NK cells in peripheral blood and pleural effusions. *Chest.* 92(3):500-4.
- Pai M, Zwerling A, Menzies D. (2008). Systematic review: T-cell-based assays for the diagnosis of latent tuberculosis infection: an update. *Ann Intern Med* 149(3):177-84.
- Paust S, von Andrian UH. (2011). Natural killer cell memory. *Nat Immunol.* 12(6):500-8.
- Pokkali S, Das SD, Selvaraj A. Differential upregulation of chemokine receptors on CD56 NK cells and their transmigration to the site of infection in tuberculous pleurisy. (2009). *FEMS Immunol Med Microbiol.* 55(3):352-60.
- Porcel JM. (2009). Tuberculous pleural effusion. *Lung.* 187(5):263-70.
- Salazar-Lezama M, Quiroz-Rosales H, Bañales-Méndez JL, Sánchez-Guzmán M, Villarreal-Velarde H, et al. (1997). Diagnostic methods of primary tuberculous pleural effusion in a region with high prevalence of tuberculosis. A study in Mexican population. *Rev Invest Clin.* 49(6):453-6.
- Schierloh P, Yokobori N, Alemán M, Musella RM, Beigier-Bompadre M, Saab MA, Alves L, et al. (2005a). Increased susceptibility to apoptosis of CD56^{dim}CD16⁺ NK cells induces the enrichment of IFN- γ -producing CD56^{bright} cells in tuberculous pleurisy. *J Immunol.* 175(10):6852-60.
- Schierloh P, Alemán M, Yokobori N, Alves L, Roldán N, Abbate E, del C Sasiain M, de la Barrera S. (2005b). NK cell activity in tuberculosis is associated with impaired CD11a and ICAM-1 expression: a regulatory role of monocytes in NK activation. *Immunology.* 116(4):541-52.
- Schierloh P, Yokobori N, Alemán M, Landoni V, Geffner L, Musella RM, Castagnino J, et al. (2007). Mycobacterium tuberculosis-induced gamma interferon production by natural killer cells requires cross talk with antigen-presenting cells involving Toll-like receptors 2 and 4 and the mannose receptor in tuberculous pleurisy. *Infect Immun.* 75(11):5325-37.
- Schierloh P, Yokobori N, Geffner L, Balboa L, Romero MM, Musella RM, Alemán M, et al. (2009). NK cells from tuberculous pleurisy express high ICAM-1 levels and exert stimulatory effect on local T cells. *Eur J Immunol.* 39(9):2450-8.
- Schierloh, P. Musella R, Castagnino J. Sasiain M.C. (2008). Interferon (IFN)- γ -producing Natural Killer (NK) cells in differential diagnosis of tuberculous pleurisy. Virtues of a functional innate immune cell based diagnostic method. *Proceedings of*

- International Conference Immunodiagnosis of Tuberculosis: New Questions, New Tools.* Virginia Beach. USA.
- Shimokata K, Saka H, Murate T, Hasegawa Y, Hasegawa T. (1991). Cytokine content in pleural effusion. Comparison between tuberculous and carcinomatous pleurisy. *Chest*. 99(5):1103-7.
- Spits H, Di Santo JP. (2011) The expanding family of innate lymphoid cells: regulators and effectors of immunity and tissue remodeling. *Nat Immunol*. 12(1):21-7.
- Supriya P, Chandrasekaran P, Das SD. (2008). Diagnostic utility of interferon-gamma-induced protein of 10 kDa (IP-10) in tuberculous pleurisy. *Diagn Microbiol Infect Dis*. 62(2):186-92.
- Steingart KR, Henry M, Laal S, Hopewell PC, Ramsay A, Menzies D, Cunningham J, Weldingh K, Pai M. (2007). A systematic review of commercial serological antibody detection tests for the diagnosis of extrapulmonary tuberculosis. *Postgrad Med J*. 83(985):705-12.
- Thorén FB, Romero AI, Hermodsson S, Hellstrand K. (2007). The CD16⁺/CD56^{bright} subset of NK cells is resistant to oxidant-induced cell death. *J Immunol*. 179(2):781-5.
- Trajman A, Pai M, Dheda K, van Zyl Smit R, Zwerling AA, Joshi R, Kalantri S, Daley P, Menzies D. (2008). Novel tests for diagnosing tuberculous pleural effusion: what works and what does not? *Eur Respir J*. 31(5):1098-106.
- Ulrichs T, Kaufmann SH. (2006). New insights into the function of granulomas in human tuberculosis. *J Pathol*. 208(2):261-9.
- Valdés L, San José E, Alvarez Dobaño JM, Golpe A, Valle JM, Penela P, González Barcala FJ. (2009). Diagnostic value of interleukin-12 p40 in tuberculous pleural effusions. *Eur Respir J*. 33(4):816-20.
- van Kooyk Y, Geijtenbeek TB. (2003). DC-SIGN: escape mechanism for pathogens. *Nat Rev Immunol*. 3(9):697-709.
- Vivier E, Raulet DH, Moretta A, Caligiuri MA, Zitvogel L, Lanier LL, Yokoyama WM, Ugolini S. (2011). Innate or adaptive immunity? The example of natural killer cells. *Science*. 331(6013):44-9.
- Vankayalapati R, Wize B, Weis SE, Samten B, Girard WM, Barnes PF. (2000). Production of interleukin-18 in human tuberculosis. *J Infect Dis*. 182(1):234-9.
- Vankayalapati R, Klucar P, Wize B, Weis SE, Samten B, Safi H, Shams H, Barnes PF. (2004) NK cells regulate CD8⁺ T cell effector function in response to an intracellular pathogen. *J Immunol*. 172(1):130-7.
- Warren HS, Kinnear BF, Skipsey LJ, Pembrey RG. (1994). Differential expression of CD45RO on natural killer (NK) cells in patients with an NK lymphocytosis. *Immunol Cell Biol*. 72(6):500-7.
- Watkins ML, Semple PL, Abel B, Hanekom WA, Kaplan G, Ress SR. (2008). Exposure of cord blood to Mycobacterium bovis BCG induces an innate response but not a T-cell cytokine response. *Clin Vaccine Immunol*. 15(11):1666-73.
- Yokobori N, Schierloh P, Geffner L, Balboa L, Romero M, Musella R, Castagnino J, et al. (2009). CD3 expression distinguishes two gammadeltaT cell receptor subsets with different phenotype and effector function in tuberculous pleurisy. *Clin Exp Immunol*. 157(3):385-94.



Understanding Tuberculosis - Analyzing the Origin of Mycobacterium Tuberculosis Pathogenicity

Edited by Dr. Pere-Joan Cardona

ISBN 978-953-307-942-4

Hard cover, 560 pages

Publisher InTech

Published online 24, February, 2012

Published in print edition February, 2012

Mycobacterium tuberculosis in an attempt to understand the extent to which the bacilli has adapted itself to the host and to its final target. On the other hand, there is a section in which other specialists discuss how to manipulate this immune response to obtain innovative prophylactic and therapeutic approaches to truncate the intimal co-evolution between Mycobacterium tuberculosis and the Homo sapiens.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Pablo Schierloh, Silvia De La Barrera and Maria Sasiain (2012). Role of NK Cells in Tuberculous Pleurisy as Innate Promoters of Local Type 1 Immunity with Potential Application on Differential Diagnosis, Understanding Tuberculosis - Analyzing the Origin of Mycobacterium Tuberculosis Pathogenicity, Dr. Pere-Joan Cardona (Ed.), ISBN: 978-953-307-942-4, InTech, Available from: <http://www.intechopen.com/books/understanding-tuberculosis-analyzing-the-origin-of-mycobacterium-tuberculosis-pathogenicity/role-of-nk-cells-in-tuberculous-pleurisy-as-innate-promoters-of-local-type-immunity-with-potential-a>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen