

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Carbonic Anhydrase and Heavy Metals

Maria Giulia Lionetto, Roberto Caricato, Maria Elena Giordano,
Elisa Erroi and Trifone Schettino

*University of Salento - Dept. of Biological and Environmental Sciences and Technologies
Italy*

1. Introduction

Carbonic anhydrase (CA; EC 4.2.1.1) is a zinc metalloenzyme catalysing the reversible hydration of CO_2 to produce H^+ and HCO_3^- . Its activity is virtually ubiquitous in nature. The fundamental role of this biochemical reaction in diverse biological systems has driven the evolution of several distinct and unrelated families of CAs. Five CA families, referred as α -, β -, γ -CA, δ , and ζ -CAs have been identified in animals, plants and bacteria (Hewett-Emmett and Tashian, 1996; Supuran, 2010). These are the α -CAs, present in vertebrates, bacteria, algae and plants; the β -CAs, predominantly in bacteria, algae and plants; the γ -CAs, mainly present in archaea and some bacteria; the δ -CAs and ζ -CAs only found in some marine diatoms (Supuran, 2010).

The monomeric α -carbonic anhydrases are by far the best studied, being found in animals. In mammals at least 16 different CA isoforms were isolated and several novel isozymes have also been identified in non-mammalian vertebrates. The α -CA isoenzymes differ in their kinetic properties, their tissue distribution and subcellular localization, and their susceptibility to various inhibitors. In general, there are three distinct groups of CA isozymes within the α -CA gene family. One of these groups contains the cytoplasmic CAs, which includes mammalian CA I, II, III, V, VII and XIII. These isozymes are found in the cytoplasm of various tissues, with the exception of the mitochondrial confined CA V. Another group of isozymes, termed the membrane-bound CAs, consists of mammalian CA IV, IX, XII, XIV and XV (Esbaugh and Tufts, 2006). These isozymes are associated with the plasma membranes of many different tissue types. The final group contains several very intriguing isozymes, CA VIII, X and XI, which are termed the CA-related proteins (CA-RP; Tashian et al., 2000). These isozymes have lost classical CA activity - the hydration/dehydration of CO_2 - and have no known physiological function; however, their highly conserved nature does suggest a very important role in vertebrates (Tashian et al., 2000).

The β -carbonic anhydrases are dimers, tetramers, or octamers and include the majority of the higher plant CA isoforms (Kimber and Pai, 2000). The γ -carbonic anhydrase is a homotrimer that has been reported for the bacterium *Methanosarcina thermophila* (Alber and Ferry, 1994). The δ class has its prototype in the monomeric CA TWCA1 from the marine diatom *Thalassiosira weissflogii* (Roberts et al., 1997; Tripp et al., 2001). The ζ -CAs are probably monomer with three slightly different active sites on the same protein backbone (Xu et al 2008).

All CAs are metalloenzymes but whereas α -, β -, and δ -CAs use Zn(II) ions at the active site, the γ -CAs are probably Fe(II) enzymes (Ferry et al., 2010), but they are active also with bound Zn(II) or Co(II) ions, and the ζ -class uses also Cd(II) to perform the physiologic reaction catalysis (Lane et al., 2000; Lane et al., 2005).

CA plays key roles in a wide variety of physiological processes involving CO₂ and HCO₃⁻. In animals the various CA isozymes are found in many different tissues and are involved in a number of different physiological processes, including bone resorption, calcification, ion transport, acid-base transport, and a number of different metabolic processes such as biosynthetic reactions (gluconeogenesis, lipogenesis, and ureagenesis). In algae and plants they play an important role in photosynthesis (Ivanov et al., 2007; Zhang et al., 2010; Cannon Gordon et al., 2010).

Considerable advances towards a detailed understanding of the catalytic mechanism of the zinc enzyme carbonic anhydrase have been made during the past years as a result of the application of crystallographic and kinetic methods to wild-type and mutant enzymes. Moreover, a great amount of work has been performed on CA inhibitors, first of all sulfonamides, RSO₂NH₂, which represent the classical CA inhibitors (CAIs) and are in clinical use for more than 50 years as diuretics and systemically acting antiglaucoma drugs (Supuran, 2010).

The review focuses on one interesting but less investigated aspect of the biochemistry of this metalloenzyme, encompassing several areas of interest from human health to environmental science: the relationships between carbonic anhydrase and heavy metals. Heavy metals are chemical elements with a density higher than 5.0 g/cm³, characterized by high reactivity, redox behaviour, and complex formation based on the characteristic of the outer *d* electron shell. In the scientific literature the following elements are normally ascribed to the heavy metal groups: aluminium, iron, silver, barium, beryllium, manganese, mercury, molybdenum, nickel, lead, copper, tin, titanium, thallium, vanadium, zinc. Some metalloids, such as arsenic, bismuthum, and selenium, are also included in the heavy metals groups.

Heavy metals generally regarded as essential for animals in trace amounts include zinc, the known cofactor of CAs, iron, copper, manganese, chromium, molybdenum and selenium. They are essential because they form an integral part of one or more enzymes involved in a metabolic or biochemical process. Besides essential metals, a number of other heavy metals, such as arsenic, lead, cadmium, mercury, have no known function in the body and are referred as toxic metals. However, also essential metals become toxic when their levels in the body exceed the homeostatic capacity of the organism. The intracellular levels of essential metals are regulated by transporters (which translocate metal across the plasma membrane) as well as by metallothionein and other metal binding proteins (Maret and Wolfgang, 2011). The toxicity of heavy metals is generally ascribed to their high affinity for nucleophilic groups like sulfhydryls. In fact they are soft donors and will therefore readily bind to soft acceptors such as sulphhydryl groups.

Recently, a number body of evidence has emerged regarding the effect of several heavy metals on carbonic anhydrase catalytic activity and protein expression. These studies encompass a wide area of interest from human health to environmental sciences.

2. Heavy metals as carbonic anhydrase cofactors

CAs catalyze the reversible hydration of carbon dioxide to bicarbonate and protons by means of a metal-hydroxide ($\text{Lig}^3\text{M}^{2+}(\text{OH})^-$) mechanism, although the α -CAs possess other catalytic activities such as esterase, phosphatase, cyanate/cyanamide hydrazase, etc. (Supuran et al., 2003; Supuran and Scozzafava 2007; Innocenti et al., 2008). In the α -, γ , and δ -CA classes, Lig^3 is always constituted by three His residues. The metal (M) is ZnII for all classes. The zinc atom is in the +2 state and is located in a cleft near the center of the enzyme. The role of zinc in carbonic anhydrase is to facilitate the deprotonation of water with the formation of the nucleophilic hydroxide ion, which can attack carbonyl group of carbon dioxide to convert it into bicarbonate. This is obtained through the +2 charge of the zinc ion which attracts the oxygen of water, deprotonates water, thus converting it into a better nucleophile able to attack the carbon dioxide.

Water naturally deprotonates itself, but it is a rather slow process. Zinc deprotonates water by providing a positive charge for the hydroxide ion. The proton is donated temporarily to the surrounding amino acid residues, and then it is given to the environment, while allowing the reaction to continue. Zinc is able to help the deprotonation of water by lowering the pKa of water. Therefore, more water molecules are now able to deprotonate at a lower pH than normal, increasing the number of hydroxide ions available for the nucleophilic attack to carbon dioxide (Berg, 2007).

The affinity of carbonic anhydrase for zinc is in subpicomolar range, as assessed for studies on the α -class (Tripp et al., 2001). Cox et al. (2000) and Hunt et al. (1999) ascribed a role for hydrophobic core residues in human CA-II that are important for preorienting the histidine ligands in a geometry that favours zinc binding and destabilizes geometries that favour other metals. In particular, mutagenesis experiments demonstrated that substitutions of these amino acids at position 93, 95, and 97 decrease the affinity of zinc, thereby altering the metal binding specificity up to 10^4 -fold. Furthermore, the free energy of the stability of native CAII, determined by solvent-induced denaturation, correlates positively with increased hydrophobicity of the amino acids at positions 93, 95, and 97 as well as with zinc affinity (Hunt et al., 1999).

β -CAs, present in green plants and cyanobacteria, contain also Zn^{2+} in the active site but are differentiated from α -CAs by virtue of the fact that the active site is coordinated by a pair of cysteine residues and a single histidine residue, whereas the fourth ligand may be either a water molecule/hydroxide ion, or a carboxylate from a conserved aspartate residue in some β -CAs (Type II β -CAs) [Trip et al., 2001; Xu et al., 2008]. The metal hydroxide catalytic mechanism seems to be also valid for these enzymes [Supuran, 2008].

Besides zinc, other metals have demonstrated to be physiologically relevant cofactor for some CAs. In fact, in the γ -CAs metal may also be FeII (Ferry et al., 2010). Cam, the prototypic γ -class carbonic anhydrase, from the anaerobic methane producing Archaea species *Methanosarcina thermophila*, contains zinc in the active site when overproduced in *Escherichia coli* and purified aerobically [Alber et al., 1996], while it has 3-fold greater carbonic anhydrase activity and contains Fe^{2+} in the active site (Fe-Cam) when purified anaerobically from *E. coli* or overproduced in the closely related species *M. acetivorans* and purified anaerobically. Soluble Fe^{2+} is abundant in oxygen free environments and available to anaerobic microbes. The different results obtained in aerobic and anaerobic conditions is

explained by the fact that in aerobic conditions Fe^{3+} is oxidized and rapidly loss from CAM enzyme, substituted by Zn^{2+} contaminating buffers not treated with chelating agents. These results indicate Fe^{2+} as the physiologically relevant metal [MacAuley et al., 2009; Tripp et al., 2004] in the active site for CAM enzyme. Interestingly, evidence for the role of ferrous ion in CA has been obtained also for the α class. In fact carbonic anhydrase activity from duck erythrocytes is increased in the presence of iron in the incubation medium suggesting a role for iron in the active site (Wu et al., 2007).

The ζ -CA naturally uses Cd^{2+} as its catalytic metal in marine diatoms (Lane and Morel, 2000; Lane et al., 2005; Park et al., 2008). This cadmium-CA (CDCA1) consists of three tandem CA repeats (R1–R3), which share 85% identity in their primary sequences (Lane et al., 2005). Although CDCA1 was initially isolated as a Cd enzyme, it is actually a “cambialistic” enzyme since it can use either Zn or Cd for catalysis – and spontaneously exchanges the two metals (Xu et al., 2008). Kinetic data show that the replacement of Zn by Cd results nonetheless in a decrease in catalytic efficiency (Xu et al., 2008). In the active site, Cd is coordinated by three invariant residues in CDCA of all diatom species (Park et al., 2007): Cys 263, His 315 and Cys 325. The tetrahedral coordination of Cd is completed by a water molecule. The use of Cd in CDCA is thought to explain the nutrient-like concentration profile of Cd in the oceans, where the metal is impoverished at the surface by phytoplankton uptake and regenerated at depth by remineralization of sinking organic matter (Lane and Morel 2000). It is cycled in the water column like an algal nutrient. It is thought that the expression of a CDCA in diatoms, which are responsible for about 40% of net marine primary production, represents an adaptation to life in a medium containing vanishingly small concentrations of essential metals (Xu et al., 2008). As suggested by Xu et al. (2008) the remarkable ability to make use of cadmium, an element known for its toxicity, gave presumably a significant competitive advantage to diatoms in the oceans, poor in metals, with respect to other species, and could have contributed to the global ratiation of diatoms during the Cenozoic Era and to the parallel decrease in atmospheric CO_2 .

Moreover, Co(II) has been shown to replace Zn(II) in α -, β and γ -CA (Hoffmann et al., 2011). Cobalt ionic radius and polarizability are very similar to those of Zn(II). In contrast to Zn(II) (d^{10}), the d^7 electron configuration of Co(II) is accessible to electronic spectroscopic methods (, yielding information about the interactions protein-metal. As a result, spectroscopy of Co(II) substituted CA isozymes has been used to probe the environment of the metal ions in the active sites and get information on the nature of the first coordination sphere of the metal (Hoffmann et al., 2011). The Co-containing form of the enzyme generally shows a marked decrease in activity compared with the native Zn form (Tu and Silverman, 1985). The demonstration that Zn can be extracted from a protein and replaced with Co *in vitro* does not demonstrate that such metal substitution takes place *in vivo*. The evidence for *in vivo* Co substitution in a CA was for the first time provided by Morel et al (1994) and Yee and Morel (1996) in the diatoms *T. weissflogii*, who demonstrated ^{65}Zn and ^{57}Co bands to co-migrate with a single band of CA activity on a native gel of diatom proteins.

3. Heavy metals as inhibitors of carbonic anhydrase activity

Several heavy metals were demonstrated to *in vitro* inhibit CA activity in a variety of organisms, including fishes, crabs, bovines, and humans.

The early work of Christensen and Tucker (1976) demonstrated carbonic anhydrase inhibition by heavy metals for the first time in fish. The study was carried out on red blood cells CA of the teleost *Oncorhynchus mykiss*. Erythrocyte CA, which represents the most abundant pool of the enzyme in fish, appeared significantly *in vitro* inhibited by several heavy metals cations, such as Cd^{2+} , Cu^{2+} , Ag^{+} , and Zn^{2+} (Tab1).

In the intestine and gills of the European eel, *Anguilla anguilla*, Lionetto et al. (1998; 2000) found cadmium to significantly inhibit carbonic anhydrase activity. The inhibition appeared tissue specific (Lionetto et al., 1998; Lionetto et al., 2000). The gill CA was much more sensitive to the heavy metal as compared to the enzyme activity in the intestine, as observed by comparing the IC50 values (Tab1). In particular in the intestine the inhibitory effect of cadmium was more pronounced on the cytosolic than the membrane-bound CA, which revealed only a partial inhibition at high concentrations. Moreover CA activity inhibition showed a certain time-dependence, with a delay of at least 10 min and 30 min for the cytosolic isoform and the membrane bound isoform respectively. The authors attributed this behaviour to the time required by cadmium for displacing the metal (zinc) associated with the enzyme, giving an inactive Cd-substituted carbonic anhydrase. Cadmium is a bivalent metal, similar in many respects to zinc: both are in the same group of the periodic table, contain the same common oxidation state (+2), and when ionized have almost the same size. Due to these similarities, cadmium can replace zinc in many biological systems. Moreover, the delayed inhibition of membrane-bound CA with respect to the cytosolic isoform was explained by a more difficult access of cadmium to the active site of the enzyme bound to the membrane. In fact, it has to be considered that the membrane-bound CA is stabilised by disulfide bonds (Whitney and Briggles, 1982) which could contribute to a less sensitivity of the membrane bound CA to cadmium.

As suggested by Lionetto et al (2000), the observed *in vitro* inhibition of cadmium on CA activity could be useful in the understanding of the toxic effects that the heavy metal can elicits on fish physiology *in vivo*. The inhibitory effect on gill CA activity suggests that the heavy metal might interfere with a number of physiological functions in which gill CA is involved as gas exchanges (Randall and Daxbaeck, 1984), acid-base balance (Heisler, 1984), osmoregulation (Henry, 1984) and clearance of the waste products from nitrogenous metabolism (Evans and Cameron, 1986). Morgan et al (2004) directly demonstrated in *in vivo* exposure experiments on rainbow trout that inhibition of branchial CA was able to induce an early decline in the gill Na^{+} and Cl^{-} uptake. With regards to the intestine, the physiological role of the cytosolic CA is that of generating HCO_3^{-} from metabolic CO_2 while the role of the CA enzyme associated to the brush-border membrane should be that of mediating the environmental HCO_3^{-} uptake (Maffia et al., 1996). Therefore, the inhibitory effect of cadmium on intestinal CA isoforms should interfere with bicarbonate balance and in turn with systemic acid-base balance and osmoregulation in fish. In fact, as previously shown (Schettino et al., 1992), the HCO_3^{-} entry via the membrane-bound CA in the cell across the luminal membrane of the enterocytes seems to be essential for maintaining a steady intracellular HCO_3^{-} concentration and/or pHi ; as a consequence the salt transport in eel intestine occurs at a highest rate and the passive water loss is recovered, so solving in part the osmoregulatory problem in marine fish. Therefore, inhibition of CA enzymes by cadmium could alter $[\text{HCO}_3^{-}]_i$ and/or pHi leading to a reduction of salt absorption and consequently impairing the osmoregulation of marine fish.

More recently, Soyut et al (2008) demonstrated Co^{2+} , Cu^{2+} , Zn^{2+} , Ag^+ , and Cd^{2+} to be potent inhibitor for brain CA enzyme activity in Rainbow trout (*Oncorhynchus mykiss*), with the following sequence $\text{Co}^{2+} > \text{Zn}^{2+} > \text{Cu}^{2+} > \text{Cd}^{2+} > \text{Ag}^+$. They also demonstrated that Co^{2+} , Ag^+ , and Cd^{2+} inhibit the enzyme with competitive manner, Cu^{2+} inhibits with noncompetitive manner, and Zn^{2+} with uncompetitive manner.

Ceyhun et al., 2011 *in vitro* demonstrated Al^{+3} , Cu^{+2} , Pb^{+2} , Co^{+3} , Ag^{+1} , Zn^{+2} and Hg^{+2} to exert inhibitory effects on fish liver CA. Metal ions inhibited the enzyme activity at low concentrations. Al^{+3} and Cu^{2+} resulted the most potent inhibitors of the CA enzyme. All the metals inhibited CA in competitive manner and aluminium showed to be the best inhibitor for fish liver CA. Concerning the mechanism of inhibition, the authors argued a possible interaction of the metal with the histidines exposed on the surface of the molecule and/or other aminoacids around the active site.

In invertebrates Vitale et al (1999) demonstrated cadmium, copper and zinc to *in vitro* inhibit CA activity in the gills of the estuarine crabs *Chasmagnathus granulata* (Tab.1). The inhibitory potentials of the three metals on CA was in the following sequence: $\text{Cu}^{2+} > \text{Zn}^{2+} > \text{Cd}^{2+}$. The observed inhibitory effect *in vitro* was confirmed by a corresponding inhibitory effect *in vivo*.

In the euryhaline crabs *Callinectes sapidus* and *Carcinus maenas* Skaggs et al (2002) also documented a significant *in vitro* inhibition of gill CA by Ag^+ , Cd^{2+} , Cu^{2+} and Zn^{2+} . The binding affinities of the metals were one thousand times weaker for cytoplasmic CA from the gills of *C. maenas* than that from *C. sapidus*. The large differences in K_i values (Tab.1) suggests the presence of two different CA isoforms in the gills of these species, with *Callinectes sapidus* possessing a highly metal-sensitive CA isoform and *Carcinus maenas* having a metal-resistant isoform. Interestingly, heavy metal inhibition of CA from the gills of another euryhaline crab, *Chasmagnathus granulata*, (as reported by Vitale et al., 1999, see above) appears to be intermediate between that found in the other two species. Moreover, in *Callinectes sapidus* CA isolated from the cytoplasmic pool of gill homogenates was much more sensitive to heavy metal inhibition than was CA from the microsomal fraction, which is believed to be anchored to the basolateral membrane, and as such, it exists within a lipid-rich environment. The authors argued that metal could be sequestered in the lipid component of the microsomal fraction and, therefore, higher amounts of metals are required to achieve an effective concentration of free metals available for CA inhibition. However, the authors did not considered the time-dependence of the inhibition, which can be an important aspect to be taken into account (see Lionetto et al., 2000) in the analysis of membrane bound vs cytosolic isoform CA inhibition.

In humans Ekinici et al (2007) demonstrated the inhibition of two human carbonic anhydrase isozymes *in vitro*, the cytosolic HCA I and II by lead, cobalt and mercury. Lead was a noncompetitive inhibitor for HCA-I and competitive for HCA-II, cobalt was competitive for HCA-I and noncompetitive for HCA-II and mercury was uncompetitive for both HCA-I and HCA-II. Lead was the best inhibitor for both HCA-I and HCA-II.

In tab.1 the K_i , IC_{50} values and the type of inhibition for several heavy metals on CA from different vertebrate and invertebrate species is summarized. A great variability among species, tissues and metals can be observed. This suggests that the inhibitory mechanisms through which heavy metals exert their effect on carbonic anhydrase activity could be different for different isoenzymes and that also small structural differences between CA isoforms could result in different metal binding affinities.

Metal	Average value of Ki (M)	IC50 (M)	Type of inhibition	Tissue	Species	Ref
Cd ²⁺	n.d.	9.979 10 ⁻⁶	n.d.	gills	<i>Anguilla anguilla</i>	Lionetto et al 2000
	n.d.	3.64 10 ⁻⁵	n.d.	Intestine (cytosolic isoform)	<i>Anguilla anguilla</i>	Lionetto et al 2000
	n.d.	2.15 10 ⁻⁵	n.d.	gills	<i>Chasmagnathus granulata</i>	Vitale et al., 1999
	n.d.	9.00 10 ⁻⁴	n.d.	Red blood cells	<i>Ictalurus punctatus</i>	Christensen and Tucker, 1976
	94.16 10 ⁻³ M	8.25 ± 10 ⁻²	Competitive	brain	<i>Oncorhynchus mykiss</i>	Soyut et al., 2008
	5.0 10 ⁻⁷	n.d.	n.d.	Gills (cytoplasmic isoform)	<i>Callinectes sapidus</i>	Skaggs and Hery, 2002
	6.0 -25.0 10 ⁻⁴	n.d.	n.d.	Gills (cytoplasmic isoform)	<i>Carcinus maenas</i>	Skaggs and Hery, 2002
Ag ⁺	193.8 10 ⁻³ M	1.59 10 ⁻¹	Competitive	brain	<i>Oncorhynchus mykiss</i>	Soyut et al., 2008
	6.40 10 ⁻⁴	3.79 10 ⁻⁴	Competitive	liver	<i>Dicentrarchus labrax</i>	Ceyhun et al., 2011
	n.d.	3.50 10 ⁻⁵	n.d.	Red blood cells	<i>Ictalurus punctatus</i>	Christensen and Tucker, 1976
	5.0–0.10 10 ⁻⁸	n.d.	n.d.	Gills (cytoplasmic isoform)	<i>Callinectes sapidus</i>	Skaggs and Hery, 2002
	6.0 -25.0 10 ⁻⁴	n.d.	n.d.	Gills (cytoplasmic isoform)	<i>Carcinus maenas</i>	Skaggs and Hery, 2002
Zn ²⁺	2.15 10 ⁻³ M	3.10 10 ⁻⁴	Uncompetitive	brain	<i>Oncorhynchus mykiss</i>	Soyut et al., 2008
	7.21 10 ⁻⁴	3.90 10 ⁻⁴	Competitive	liver	<i>Dicentrarchus labrax</i>	Ceyhun et al., 2011
	n.d.	7.00 10 ⁻⁴	n.d.	Red blood cells	<i>Ictalurus punctatus</i>	Christensen and Tucker, 1976
	n.d.	1.62 10 ⁻⁵	n.d.	gills	<i>Chasmagnathus granulata</i>	Vitale et al., 1999
	6.0 -25.0 10 ⁻⁴	n.d.	n.d.	Gills (cytoplasmic isoform)	<i>Carcinus maenas</i>	Skaggs and Hery, 2002

Metal	Average value of Ki (M)	IC50 (M)	Type of inhibition	Tissue	Species	Ref
Cu ²⁺	27.6 10 ⁻³ M	3.00 10 ⁻²	Non competitive	brain	<i>Oncorhynchus mykiss</i>	Soyut et al., 2008
	1.75 10 ⁻⁵	7.15 10 ⁻⁵	Competitive	liver	<i>Dicentrarchus labrax</i>	Ceyhun et al., 2011
	n.d.	6.50 10 ⁻⁵	n.d.	Red blood cells	<i>Ictalurus punctatus</i>	Christensen and Tucker, 1976
	n.d.	3.75 10 ⁻⁶	n.d.	gills	<i>Chasmagnathus granulata</i>	Vitale et al., 1999
	3.60 10 ⁻⁷	n.d.	n.d.	Gills (cytoplasmic isofom)	<i>Callinectes sapidus</i>	Skaggs and Hery, 2002
	6.0 -25.0 10 ⁻⁴	n.d.	n.d.	Gills (cytoplasmic isofom)	<i>Carcinus maenas</i>	Skaggs and Hery, 2002
Co ²⁺	5 10 ⁻⁵ M	1.40 10 ⁻⁵	competitive	brain	<i>Oncorhynchus mykiss</i>	Soyut et al., 2008
	5.32 10 ⁻⁴	3.16 10 ⁻⁴	competitive	liver	<i>Dicentrarchus labrax</i>	Ceyhun et al., 2011
	3.91 10 ⁻³	n.d.	competititve	Erythrocytes (CAI)	<i>Homo sapiens</i>	Ekinci et al., 2007
	1.7 10 ⁻³	n.d.	non competitive	Erythrocytes (CAII)	<i>Homo sapiens</i>	Ekinci et al., 2007
Al ³⁺	1.48 10 ⁻⁴	6.92 10 ⁻⁵	competitive	liver	<i>Dicentrarchus labrax</i>	Ceyhun et al., 2011
Pb ²⁺	2.42 10 ⁻⁴	1.13 10 ⁻⁴	competitive	liver	<i>Dicentrarchus labrax</i>	Ceyhun et al., 2011
	9.90 10 ⁻⁴	n.d.	Non competitive	Erythrocytes (CAI)	<i>Homo sapiens</i>	Ekinci et al., 2007
	5.6 10 ⁻⁵	n.d.	uncompetitive	Erythrocytes (CAII)	<i>Homo sapiens</i>	Ekinci et al., 2007
Hg ²⁺	7.68 10 ⁻⁴	4.48 10 ⁻⁴	competitive	liver	<i>Dicentrarchus labrax</i>	Ceyhun et al., 2011
	1.42 10 ⁻³	n.d.	uncompetitive	Erythrocytes (CAI)	<i>Homo sapiens</i>	Ekinci et al., 2007
	3.12 10 ⁻⁴	n.d.	uncompetitive	Erythrocytes (CAII)	<i>Homo sapiens</i>	Ekinci et al., 2007

Table 1. Ki, IC50 and type of inhibition for several heavy metals in different species and tissues as assessed in *in vitro* studies.

Concerning the mechanisms of inhibition some heavy metals are believed to bind to CA not at the specific catalytic site of CO₂ hydration but nearby in a pocket, the so called ‘proton

shuttle' as demonstrated for human CAII (Tu et al., 1981). His-64 is a proton shuttle in catalysis, where it accepts the proton product (via the bridging solvent molecules) from zinc-bound water as zinc-bound hydroxide is regenerated; subsequently, the proton product is passed along to buffer (Liang et al, 1988; Tu et al., 1989; Vedani et al., 1989). The mechanism of inhibition of heavy metals on proton shuttle has been elucidated for copper on human CA II. Cu^{2+} is believed to competitively inhibit CAII by binding to the imidazole side chain of His-64, blocking its role in proton transfer from the zinc-bound water molecule to buffer molecules located outside of the active site region [Tu et al., 1981]. However, the knowledge of the mechanism of action of other metals on different CA isoforms is lacking. It cannot be excluded the CA binding to other different parts of the protein, possibly cysteine residues, as demonstrated in studies with other enzymes for silver and mercury.

4. Heavy metals as modulators of carbonic anhydrase activity and expression

If it has been widely demonstrated *in vitro* that heavy metals are able to inhibit CA activity in a variety of organisms, on the contrary little is known about the *in vivo* effects of trace metals on the activity and the expression of this metalloenzyme. The major information regards Zn^{2+} , while very few is known about other metals.

In humans early studies demonstrated that dietary zinc deficiency significantly reduces zinc concentrations of serum and in turn CA activity in erythrocytes (Hove, 1940; Rahman et al., 1961; Kirchgessner et al., 1975) suggesting a possible influence of Zn^{2+} on CA protein expression. These early data have been more recently confirmed by Lukaski (2005) who demonstrated zinc concentration of serum and erythrocyte to be positively correlated to CA activity *in vivo*. Low dietary zinc decreases erythrocyte carbonic anhydrase activity and, in turn, impairs cardiorespiratory function in men during exercise (Lukaski et al., 2005). In ducks Zn^{2+} at a low level (up to $1.25 \mu\text{M}$ Zn) induced the rise of CA activity in erythrocytes (Wu et al., 2007). In parotid saliva of patients with CAVI deficiency Zn^{2+} treatment was able to stimulate synthesis/secretion of CAVI (Henkin et al., 1999), probably through stimulation of CAIV gene. In rats Zn^{2+} deficiency significantly reduced CAII protein expression in the submandibular gland (Goto et al., 2008).

As regards other metals Grimes et al (1997) reported the depression of CAIII mRNA and, in turn, CAIII protein in the mouse mutant 'toxic milk' (tx) liver following copper accumulation, Kuhara et al (2011) found CAIII suppression by copper accumulation during carcinogenesis, while Wu et al (2007) found iron at low levels to induce a rise in CA activity in duck erythrocytes.

Recently, Caricato et al (2010) demonstrated for the first time CA activity and protein expression to be enhanced by the exposure to the trace element cadmium in animals, opening new perspective in the comprehension of the functioning and regulation of this enzyme. Digestive gland CA activity showed a weak sensitivity to *in vitro* cadmium exposure since only high concentrations of CdCl_2 (from 10^{-5} to 10^{-3} M) were able to exert a significant inhibition. On the contrary digestive gland CA activity showed a significant increment in cadmium exposed animals (about 40% after two week of exposure). This was the first time that CA activity appears to be increased by cadmium in animals. Carbonic anhydrases from the microalgae *Chlamydomonas reinhardtii* (Wang et al., 2005) and *Thalassiosira weissflogii* (Morel et

al., 1994; Lee et al., 1995) are the only other examples reported in nature of CA activity increase induced by cadmium exposure. Evidence of *in vivo* utilization of Cd in CA has been found in microalgae (Price and Morel, 1990; Morel et al., 1994; Lee et al. 1995, Xu et al., 2008). In these organisms the ability of Cd to substitute for Zn at the active site of the enzyme is reflected in the regulation of the enzyme expression. In *Thalassiosira weissflogii* a cadmium-containing CA was found to be expressed during zinc limitation (Lane and Morel, 2000; Lane et al., 2005). This cadmium CA (CDCA1) which naturally uses Cd as its catalytic metal (Trip et al., 2001; Lane et al., 2005) has been ascribed to a novel ζ -CA class (see above). Genes coding for similar proteins have been identified in other cultured diatoms (Park et al., 2007). In mussel digestive gland western blotting analysis clearly demonstrated the enhancement of CA protein expression following cadmium exposure, according to the enzymatic activity data (Caricato et al., 2010). Laboratory experimental results were confirmed by a field experiment. Mussels exposed for 30 days to an anthropogenic impacted site showed a significant increase in CA activity and protein expression with respect to animals exposed for 30 days in a control site. If the new synthesized enzyme is a Cd-CA is not possible to say at the moment. If it was the case, then the increase in CA would not be a direct adaptive response to Cd pollution; rather, Cd could remove any limitations placed on CA synthesis by the availability of Zn. However, future studies will be needed to clarify this intriguing aspect of the research.

5. Carbonic anhydrase and heavy metals interactions: Potential applications

In the last years the interactions between carbonic anhydrase and heavy metals have found a number of applications in environmental and health fields, including the development of biomarkers of pollution exposure, *in vitro* bioassays, and biosensors.

5.1 Carbonic anhydrase sensitivity to heavy metals and development of biomarkers of pollution exposure

Pollution by trace metals is a world-wide problem due to the persistency and continuing accumulation of metals in the environment (de Mora et al. 2004; Hwang et al 2006). Heavy metals may enter the organisms through food, water, air, or absorption through the skin. As a result of mining, waste disposal and fuel combustion the environment is becoming increasingly contaminated with heavy metals.

In recent years the increasing sensibility to pollution problems has promoted the development of environmental "diagnostic" tools for early warning detection of pollution. Pollution monitoring has been increasingly concerned with the use of biological responses to pollutants at molecular and cellular level for evaluating biological hazard of toxic chemicals. Methods based on biological effects and their underlying mechanisms can complement the use of analytical chemistry in environmental monitoring. The major advantages of such biological, mechanism-based methods are their toxicological specificity, rapidity, and low cost. Toxicological specificity refers to the relationship between the assay response and the toxic potential rather than simply the contaminant concentrations (provided by chemical analysis) of the sample being analyzed. Moreover, biological assays provide rapid, sensitive, easily learnt and readily interpretable new useful tools for environmental biomonitoring and risk assessment. They include biomarkers, and *in vivo* and *in vitro* bioassays. It is known that the harmful effects of pollutants are typically first manifested at lower levels of biological organization before disturbances are realized at

population, community and ecosystem levels (Adams, 1990). This is the reason why in recent years the study of molecular and cellular effects of pollutants has given important advancement in the developing of biologically-based methodologies useful for environmental biomonitoring and risk assessment. Enzymatic inhibition studies have been a very fruitful field for environmental monitoring application as biomarker of exposure/effect. Biomarkers are defined as pollutant induced variation in cellular or biochemical components occurring in organisms as a result of natural exposure to contaminants in their environment (Depledge, 1994). As reported by several authors, the evaluation of biomarkers in bioindicator organisms sampled in one or more areas suspected of chemical contamination and their comparison with organisms sampled in a control area can allow the evaluation of the potential risk of toxicological exposure of the studied community (Lionetto et al., 2003; Lionetto et al., 2004).

Carbonic anhydrase sensitivity to heavy metal exposure has been recently explored for its possible applications as biomarker of exposure to heavy metal pollution (Lionetto et al. 2006; Caricato et al, 2010b.) in “sentinel” organisms. Lionetto et al., (2006) investigated CA activity inhibition by heavy metals in the filter feeding *Mytilus galloprovincialis*, widely used in pollution monitoring programs as sentinel organism (Jernelov et al., 1996). Following *in vitro* and *in vivo* exposure to cadmium, mantle CA activity was significantly inhibited. The inhibitory effect of cadmium on mantle CA activity can explain results previously obtained by Soto et al. (2000), who observed a significant decrease in shell growth in *M. galloprovincialis* exposed to heavy metals. The sensitivity of CA to heavy metals in mussels appears to be tissue-specific. In fact, as reported above, in mussel’s digestive gland CA activity and expression was found to increase following Cd exposure (Caricato et al., 2010). Because of the widely application of *M. galloprovincialis* in environmental quality monitoring and assessment, data on tissue specific sensitivity of carbonic anhydrase to heavy metals represent a starting point for future potential application of CA activity changes as biomarker of exposure to heavy metals in the sentinel organism *M. galloprovincialis*.

Other studies carried out on corals have suggested alteration in CA activity as potential biomarker of exposure to environmental chemical stress. CA activity has been demonstrated to be inhibited by heavy metal exposure in anemones and corals (Gilbert and Guzman, 2001), where the enzyme plays a key role in the calcification process. Coral growth has been shown to be an effective indicator of the overall health of a coral reef ecosystem and reduced growth can reflect impaired photosynthetic output of the zooxanthellae and/or changes in enzyme activity (Moya et al., 2008). In an era of climate change and ocean acidification, where factors impacting growth and resilience factors are becoming important, understanding the biological effects of metal exposure to these keystone tropical organisms may be critical (Bielmyer et al., 2010).

5.2 Carbonic anhydrase based bioassay

Bioassays use biological systems to detect the presence of toxic chemicals in the environmental matrices (water, sediment, sewage, soil, etc.). In recent years, *in vitro* bioassays, employing cultured cells or cellular extracts, are increasingly being developed and used to detect the presence of contaminants. Examples include assays that measure enzyme inhibition, receptor-binding, or changes in gene expression in *in vitro* systems. Although *in vitro* assay is not a substitute for biomarker approach, it can be used as an

adjunct model to whole-animal *in vivo* exposure and to ecotoxicological evaluation of the potential risk of trace pollutants in aquatic environments. They are rapid, low cost and simple tools to be utilized in combination with chemical analysis, for the pre-screening of the environmental samples that should be analyzed. Lionetto et al (2005; 2006) explored the possible application of heavy metal CA inhibition for the development of an *in vitro* bioassay applicable to the determination of the toxicity of environmental aqueous samples. They developed rapid and sensitive chemical hazard detection system for standardizing rapid, sensitive, and low cost CA based *in vitro* bioassay (Schettino et al., 2008).

6. Carbonic anhydrase-based biosensing of metal ions

In the last years the affinity of carbonic anhydrase for metal ions has been applied for the development of fluorescence based biosensors for determination of free metal ions in solution using variants of human carbonic anhydrase (apoCA). In particular, Cu^{2+} , Co^{2+} , Zn^{2+} , Cd^{2+} , and Ni^{2+} have been determined at concentration down the picomolar range (Fierke and Thompson, 2001; Thompson and Jones, 1993; Mey et al., 2011) by changes in fluorescence emission (Thompson et al., 2000) and excitation wavelength ratios (Thompson et al., 2002), lifetimes (Thompson and Patchan, 1995), and anisotropy (polarization) (Elbaum et al., 1996; Thompson et al., 2000). The sensitivity, selectivity, analyte binding, kinetics and stability of the biosensors have been improved by subtle modification of the protein structure by directed mutagenesis (Kiefer et al., 1995; Hunt et al., 1999; DiTusa et al., 2001; McCall et al., 2004; Burton et al., 2000). These studies have hallowed the development of highly selective and sensitive fluorescence-based biosensors for Zn^{2+} e Cu^{2+} , which have been shown to be viable approach in some important applications. In fact, the CA-based Cu^{2+} biosensor has been used to obtain real-time measurement of free Cu(II) at picomolar concentrations in seawater (Zeng et al., 2003), while the CA-base Zn^{2+} biosensor has been used for measurement of free Zn ion at picomolar levels in cultured cells (Bozym et al, 2004).

7. Conclusions

Although carbonic anhydrase represents one of the most investigated metalloenzyme in nature, its interaction with heavy metals has been only partially elucidated to date and some issues still remains to be explored. An intriguing aspect that needs more investigation is the *in vivo* effect of heavy metals on CA expression. From the few studies available in literature some metals appear to be important modulator of the expression of this protein. The understanding of the underlying mechanisms could open new perspective in the comprehension of the functioning and regulation of this enzyme. Another intriguing aspect of the biochemistry of CA is the inhibition by heavy metals. It has been documented for some species and some metals, but the mechanisms behind the inhibition, its metal specificity and isoform specificity remains still unknown. These aspects merits in depth examination and open new perspective for drug design and biomarkers development.

8. References

- Adams, S.M.; Crumby, W.D.; Greeley, M.S.; Ryon, M.G. & Schilling, E.M. (1990). Relationship between Physiological and Fish Population Responses in a Contaminated Stream. *Environmental Toxicology and Chemistry*, Vol. 11, Issue 11, (November 1992), pp. 1549-1557, ISSN: 0730-7268

- Alber, B.E. & Ferry, J.G. (1994). A carbonic anhydrase from the archaeon *Methanosarcina thermophila*. *Proceedings of the National Academy of Sciences of the United States of America*, Vol.91, (July 1994), pp. 6909-6913, ISSN 0027-8424
- Alber, B.E. & Ferry, J.G. (1996). Characterization of heterologously produced carbonic anhydrase from *Methanosarcina thermophila*, *The Journal of Bacteriology*, Vol.178, No.11, (June 1996), pp.3270-3274, ISSN 0021-9193
- Alber, B.E.; Colangelo, C.M.; Dong, J.; Staalhandske, C.M.V.; Baird, T.T.; Tu, C.; Fierke, C.A.; Silverman, D.N.; Scott, R.A. & Ferry, J.G. (1999). Kinetic and Spectroscopic Characterization of the Gamma-Carbonic Anhydrase from the Methanoarchaeon *Methanosarcina thermophil.* *Biochemistry*, Vol.38, Issue 40, (October 1999), pp. 13119-13128, ISSN 0006-2960
- Berg, J.M. (2007). *Biochemistry*, 6th Ed., Sara Tenney. ISBN0-7167-8724-5
- Bertini, I. & Luchinat, C. (1984). High spin cobalt(II) as a probe for the investigation of metalloproteins. *Advances in inorganic biochemistry*, Vol. 6, pp. 71-111, ISSN 0190-0218
- Bielmyer, G.K.; Grosell, M.; Bhagooli, R.; Baker, A.C.; Langdon, C.; Gillette, P. & Capo, T.R. (2010). Differential effects of copper on three species of scleractinian corals and their algal symbionts (*Symbiodinium* spp.). *Aquatic Toxicology*, Vol. 97, No. 2, (April 2010), pp. 125-133, ISSN 0166-445X
- Bozym, R.A.; Zeng, H.H.; Cramer, M.; Stoddard, A.; Fierke, C.A. & Thompson, R.B. (2004). In vivo and intracellular sensing and imaging of free zinc ion. In: *Proceedings of the SPIE Conference on Advanced Biomedical and Clinical Diagnostic Systems II*, Cohn, G.E., Grundfest, W.S., Benaron, D.A.; and Vo-Dinh, T., eds. Bellingham, WA: SPIE, 2004.
- Burton, R.E.; Hunt, J.A.; Fierke, C.A. & Oas, T.G. (2000). Novel disulfide engineering in human carbonic anhydrase II using the PAIRWISE side-chain geometry database. *Protein Science*, Vol. 9, Issue 4, (April 2000), pp. 776-785, ISSN 0961-8368
- Cannon, G.C.; Heinhorst, S. & Kerfeld C.A. (2010). Carboxysomal carbonic anhydrases: Structure and role in microbial CO₂ fixation. *Biochimica et Biophysica Acta-Proteins and Proteomics*, Vol. 1804, Issue 2, (February 2010), pp. 382-392, ISSN: 1570-9639
- Caricato, R.; Lionetto, M.G.; Dondero, F.; Viarengo, A. & Schettino, T. (2010a) Carbonic anhydrase activity in *Mytilus galloprovincialis* digestive gland: sensitivity to heavy metal exposure. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*, Vol.152C, Issue 3, (September 2010), pp.241-247, ISSN 1532-0456
- Caricato, R.; Lionetto, M.G. & Schettino, T. (2010b). Seasonal variation of biomarkers in *Mytilus galloprovincialis* sampled inside and outside Mar Piccolo of Taranto (Italy). *Chemistry and Ecology*, Vol.26, supplement 1, (June 2010), pp.143-153, ISSN 0275-7540
- Ceyhun, S.B.; Şentürk, M.; Yerlikaya, E.; Erdoğan, O.; Küfrevioğlu, Ö.I. & Ekinçi, D. (2011). Purification and characterization of carbonic anhydrase from the teleost fish *Dicentrarchus labrax* (European Seabass) liver and toxicological effects of metals on enzyme activity. *Environmental Toxicology and Pharmacology*. Vol.32, Issue 1, (July 2011), pp. 69-74, ISSN 1382-6689
- Christensen, G.M. & Tucker, J.H. (1976). Effects of selected water toxicants on the in vitro activity of fish carbonic anhydrase. *Chemico Biological Interactions*, Vol.13, Issue 2, (May 1976), pp.181-92, ISSN 0009-2797

- Cox, J.D.; Hunt, J.A.; Compher, K.M.; Fierke, C.A. & Christianson, D.W. (2000). Structural Influence of Hydrophobic Core Residues on Metal Binding and Specificity in Carbonic Anhydrase II. *Biochemistry*, Vol.39, No.45, (November 2000), pp.13687–13694, ISSN 0006-2960
- De Mora, S.; Fowler, S.W.; Wyse, E. & Azemard, S. (2004). Distribution of heavy metals in marine bivalves, fish and coastal sediments in the Gulf and Gulf of Oman, *Marine Pollution Bulletin*, Vol. 49, Issue 5-6, (September 2004), pp.410-424, ISSN 0025-326X
- Depledge, M.H. (1994). The rational basis for the use of biomarkers as ecotoxicological tools. In: *Nondestructive Biomarkers in Vertebrates*, M.C., Fossi, C., Leonzio (Eds.), pp. 271–295, Lewis Publisher, ISBN 978-0873716482, Boca Raton, USA
- DiTusa, C.A.; McCall, K.A.; Chritensen, T.; Mahapatro, M.; Fierke, C.A. & Toone, E.J. (2001). Thermodynamics of metal ion binding. 2. Metal ion binding by carbonic anhydrase variants. *Biochemistry*, Vol.40, Issue 18, (May 2001), pp.5345-5351, ISSN 0006-2960
- Ekinci, D.; Beydemir, Ş. & Küfrevioğlu Ö.İ. (2007). In vitro inhibitory effects of some heavy metals on human erythrocyte carbonic anhydrases. *Journal of Enzyme Inhibition and Medicinal Chemistry*, Vol.22, Issue 6, pp.745–750, ISSN 1475-6366
- Elbaum, D.; Nair, S.K.; Patchan, M.W.; Thompson, R.B. & Christianson, D.W. (1996). Structure-based design of a sulfonamide probe for fluorescence anisotropy detection of zinc with a carbonic anhydrase-based biosensor. *Journal of the American Chemical Society*, Vol. 118, Issue 35, (September 1996), pp. 8381-8387, ISSN 0002-7863
- Esbaugh, A.J. & Tufts, B.L. (2006). The structure and function of carbonic anhydrase isozymes in the respiratory system of vertebrates. *Respiratory Physiology & Neurobiology*, Vol.154, Issue 1-2, (November 2006), pp. 185–198, ISSN 1569-9048
- Evans, D.H. & Cameron, J.N. (1986). Gill ammonia transport. *Journal of Experimental Zoology*, Vol.239, Issue 1, (July 1986), pp. 17–23, ISSN 0022-104X
- Ferry, J.F. (2010). The gamma class of carbonic anhydrases. *Biochimica et Biophysica Acta (BBA) - Proteins & Proteomics*, Vol.1804, Issue 2, (February 2010), pp. 374-38, ISSN 1570-9639
- Fierke, C.A. & Thompson, R.B. (2001). Fluorescence-based biosensing of zinc using carbonic anhydrase. *Biometals*, Vol. 14, Issue 3-4, (September 2001), pp. 205-222, ISSN 0966-0844
- Gilbert, A.L. & Guzman, H.M. (2001). Bioindication potential of carbonic anhydrase activity in anemones and corals. *Marine Pollution Bulletin*, Vol. 42, Issue 9, (September 2001), pp. 742–744, ISSN 0025-326X
- Goto, T.; Shirakawa, H.; Furukawa Y. & Komai, M. (2008). Decreased expression of carbonic anhydrase isozyme II, rather than of isozyme VI, in submandibular glands in long-term zinc-deficient rats. *British Journal of Nutrition*, Vol.99, Issue 2, (February 2008), pp. 248-53, ISSN 0007-1145
- Grimes, A.; Paynter, J.; Walker, I.D.; Bhave, M. & Mercer, J.F.B. (1997). Decreased carbonic anhydrase III levels in the liver of the mouse mutant “toxic milk” (tx) due to copper accumulation, *Biochemical Journal*, Vol. 321, Part. 2, (January 1997), pp. 341–346, ISSN 0264-6021
- Heisler, N. (1984). Acid-base regulation in fishes. In: *Hoar, W.S., Randall, D.J. (Eds.), Fish Physiology*, vol. 10A. Academic Press, New York, pp. 315–401.

- Henkin, R.; Martin, B.M. & Agarwal, R. (1999). Efficacy of exogenous oral zinc in treatment of patients with carbonic anhydrase VI deficiency. *American Journal of the Medical Science*, Vol. 318, Issue 6, (December 1999), pp. 392-405, ISSN 0002-9629
- Henry, R.P. (1984). The role of carbonic anhydrase in blood ion and acid-base regulation. *American Zoologist*. Vol. 24(1), pp. 241-253, 0003-1569.
- Hewett-Emmett, D. & Tashian, R.E. (1996). Functional diversity, conservation, and convergence in the evolution of the α -, β -, and γ -carbonic anhydrase gene families. *Molecular Phylogenetics and Evolution*, Vol. 5, Issue 1, (February 1996), pp. 50-77, ISSN 1055-7903
- Hoffmann, K.M.; Samardzic, D.; van den Heever, K. & Rowlett, R.S. (2011). Co(II)-substituted *Haemophilus influenzae* β -carbonic anhydrase: Spectral evidence for allosteric regulation by pH and bicarbonate ion. *Archives of Biochemistry and Biophysics*, Vol. 511, Issue 1-2, (July 2011), pp. 80-87, ISSN 0003-9861
- Hove, C.; Elvehjem, C.A. & Hart, E.B. (1940). The relation of zinc to carbonic anhydrase. *J Biol Chem* 136:425-434
- Hunt, J.A. & Fierke, C.A. (1997). Selection of carbonic anhydrase variants displayed on phage. Aromatic residues in zinc binding site enhance metal affinity and equilibration kinetics. *Journal of Biological Chemistry*, Vol.272, Issue 33, (August 1997), pp. 20364-20372, ISSN 0021-9258
- Hunt, J.A.; Ahmed, M. & Fierke, C.A. (1999). Metal Binding Specificity in Carbonic Anhydrase is Influenced by Conserved Hydrophobic Amino Acids. *Biochemistry*, Vol. 38, Issue 28, (July 1999), pp. 9054-9060, ISSN 0006-2960
- Hwang, H.; Green, P.G.; Higashi, R.M. & Young, T.M. (2006). Tidal salt marsh sediment in California, USA. Part 2: Occurrence and anthropogenic input of trace metals, *Chemosphere*, Vol. 64, Issue 11, (September 2006), pp.1899-1909, ISSN: 0045-6535
- Innocenti, A.; Scozzafava, S.; Parkkila, L.; Puccetti, G.; De Simone, G. & Supuran, C.T. (2008). Investigations of the esterase, phosphatase, and sulfatase activities of the cytosolic mammalian carbonic anhydrase isoforms I, II, and XIII with 4-nitrophenyl esters as substrates. *Bioorganic & Medicinal Chemistry Letters*, Vol. 18, Issue 7, (April), pp.2267-2271, ISSN 0960-894X
- Ivanov, B.N.; Ignatova, L.K. & Romanova, A.K. (2007). Diversity in Forms and Functions of Carbonic Anhydrase in Terrestrial Higher Plants. *Russian Journal of Plant Physiology*, Vol. 54, No. 2, (March-April 2007), pp. 143-162, ISSN 1021-4437
- Jernelov, A. (1996). The international mussel watch: a global assessment of environmental levels of chemical contaminants. *Science of the Total Environment*, Vol.188, Supplement 1, (September 1996), pp. 37-44, ISSN: 0048-9697
- Kiefer L.L.; Paterno S.A. & Fierke, C.A. (1995). Hydrogen-Bond Network in the Metal-Binding Site of Carbonic Anhydrase Enhances Zinc Affinity and Catalytic Efficiency. *Journal of the American Chemical Society*, Vol. 117, Issue 26, (July 1995), pp. 6831-6837, ISSN 0002-7863
- Kimber, M.S. & Pai, E.F. (2000). The active site architecture of *Pisum sativum* β -carbonic anhydrase is a mirror image of that of α -carbonic anhydrases. *Embo Journal*, Vol. 19, Issue 7, (April 2000), pp.1407-1418, ISSN 0261-4189
- Kirchgessner M, Stadler AE, Roth HP (1975) Carbonic anhydrase activity and erythrocyte count in the blood of zinc-deficient rats. *Bioinorganic chemistry*, Vol. 5, Issue 1, pp. 33-38, ISSN: 0006-3061

- Kuhara, M.; Wang, J.; Flores, M.J.; Qiao, Z.; Koizumi, Y.; Koyota, S.; Taniguchi, M. & Sugiyama, T. (2011). Sexual dimorphism in LEC rat liver: suppression of carbonic anhydrase III, by copper accumulation during hepatocarcinogenesis. *Biomedical Research*, Vol. 32 (2), (April 2011), pp. 111-117, ISSN 0388-6107
- Lane, T.W. & Morel, F.M.M. (2000). A biological function for cadmium in marine diatoms. *Proceedings of the National Academy of Sciences of the United States of America*, Vol.97, Issue 9, (April 2000), pp. 4627-4631, ISSN 0027-8424
- Lane, T.W.; Saito, M.A.; George, G.N.; Pickering, I.J.; Prince, R.C. & Morel, F.M.M. (2005). A cadmium enzyme from a marine diatom. *Nature*, Vol. 435, Issue 7038, (May 2005), pp.42-42, - ISSN 0028-0836
- Lee, J.G.; Roberts, S.B. & Morel, F.M.M. (1995). Cadmium a nutrient for the marine diatom. *Limnology and Oceanography*, Vol. 40, Issue 6, (September 1995), pp. 1056-1063, ISSN 0024-3590
- Liang, J.Y. & Lipscomb, W. N (1988). Hydration of CO₂ By Carbonic-Anhydrase - Intramolecular Proton Transfer between Zn²⁺-Bound H₂O and Histidine-64 in Human Carbonic Anhydrase-II. *Biochemistry*, Vol. 27, Issue 23, (November 1988), pp. 8676-8682, ISSN 0006-2960
- Lionetto, M.G.; Maffia, M.; Cappello, M.S.; Giordano, M.E.; Storelli, C. & Schettino, T. (1998). Effect of cadmium on carbonic anhydrase and Na⁺-K⁺-ATPase in eel, *Anguilla anguilla*, intestine and gills. *Comparative Biochemistry and Physiology A- Molecular and Integrative Physiology*, Vol.120, Issue 1, (May 1998), pp.89-91, ISSN 1095-6433
- Lionetto, M.G.; Giordano, M.E.; Vilella, S. & Schettino, T. (2000). Inhibition of eel enzymatic activities by cadmium. *Aquatic Toxicology*, Vol. 48, Issue 4, (April 2000), pp. 561-571, ISSN: 0166-445X
- Lionetto, M.G.; Caricato, R.; Giordano, M.E.; Pascariello, M.F.; Marinosci, L. & Schettino, T. (2003). Integrated use of biomarkers (acetylcholinesterase and antioxidant enzymatic activities) in *Mytilus galloprovincialis* and *Mullus barbatus* in an Italian coastal marine area. *Marine Pollution Bulletin*, Vol. 46, Issue 3, (March 2003), pp. 324-330, ISSN 0025-326X
- Lionetto, M.G.; Caricato, R.; Erroi, E.; Giordano, M.E. & Schettino, T. (2005). Carbonic anhydrase based environmental bioassay. *International Journal of Environmental Analytical Chemistry*, Vol.85, Issue 12-13, (October-November 2005), pp. 895-903, ISSN: 0306-7319
- Lionetto, M.G.; Caricato, R.; Erroi, E.; Giordano, M.E. & Schettino, T. (2006). Potential application of carbonic anhydrase activity in bioassay and biomarker studies. *Chemistry and Ecology*, Vol. 22, Supplement 1, pp. 119-125, ISSN: 0275-7540
- Lukaski, H.C. (2005). Low dietary zinc decreases erythrocyte carbonic anhydrase activities and impairs cardiorespiratory function in men during exercise. *American Journal of Clinical Nutrition*, Vol 81, Issue 5, (May 2005), pp.1045-1051, ISSN 0002-9165
- MacAuley, S.R.; Zimmerman, S.A.; Apolinario, E.E.; Evilia, C.; Hou, Y.; Ferry, J.G. & Sowers, K.R. (2009). The archetype γ -class carbonic anhydrase (Cam) contains iron when synthesized in vivo, *Biochemistry*, Vol. 48, Issue 5, (February 2009) pp 817-819, ISSN: 0006-2960
- Maffia, M.; Trischitta, F.; Lionetto, M.G.; Storelli, C. & Schettino, T. (1996). Bicarbonate absorption in eel intestine: Evidence for the presence of membrane-bound carbonic


- anhydrase on the brush border membranes of the enterocyte. *Journal of Experimental Zoology*, Vol: 275, Issue 5, (August 1996), pp. 365-373, ISSN 0022-104X
- Maret, W. (2011). Metals on the move: zinc ions in cellular regulation and in the coordination dynamics of zinc proteins. *Biometals*, Vol. 24, Issue 3, (June 2011), pp. 411-418, ISSN: 0966-0844
- Marouan, R.; Cecchi, A.; Montero, J.L.; Innocenti, A.; Vullo, D.; Scozzafava, A.; Winum, J.Y. & Supuran, C.T. (2008) Carbonic Anhydrase Inhibitors: Design of Membrane-Impermeant Copper(II) Complexes of DTPA-, DOTA-, and TETA-Tailed Sulfonamides Targeting the Tumor-Associated Transmembrane Isoform IX. *ChemMedChem*, Vol. 3, Issue 11, (November 2008), pp. 1780 – 1788, ISSN 1860-7179
- McCall, K.A. & Fierke, C.A. (2004). Probing Determinants of the Metal Ion Selectivity in Carbonic Anhydrase Using Mutagenesis. *Biochemistry*, Vol. 43, Issue 13, (April 2004), pp. 3979-3986, ISSN 0006-2960
- Mei, Y.J.; Frederickson, C.J.; Giblin, L.J.; Weiss, J.H.; Medvedeva, Y. & Bentley, PA (2011). Sensitive and selective detection of zinc ions in neuronal vesicles using PYDPY1, a simple turn-on dipyrin. *Chemical Communications*, Vol. 47, Issue 25, pp. 7107-7109, ISSN 1359-7345
- Morel, F.M.M.; Reinfelder, J.R.; Roberts, S.B.; Chamberlain, C.P.; Lee, J.G. & Yee, D. (1994). Zinc and carbon co-limitation of marine phytoplankton. *Nature*, Vol. 369, Issue 6483, (June 1994), pp. 740–742, ISSN: 0028-0836
- Morgan, I.J.; Henry, R.P. & Wood, C.M. (1997). The mechanism of acute silver nitrate toxicity in freshwater rainbow trout (*Oncorhynchus mykiss*) is inhibition of gill Na⁺ and Cl⁻ transport. *Aquatic Toxicology*, Vol.38, Issue 1-3, (May 1997), pp. 145–63, ISSN: 0166-445X
- Morgan, T.P.; Grosell, M.; Gilmour, K.M.; Playle, R.C. & Wood, C.M. (2004). Time course analysis of the mechanism by which silver inhibits active Na⁺ and Cl⁻ uptake in gills of rainbow trout. *American Journal of Physiology-Regulatory Integrative and Comparative Physiology*, Vol. 287, Issue 1, (July 2004), ISSN 0363-6119
- Moya, A.; Ferrier-Pages, C.; Furla, P.; Richier, S.; Tambutte, E.; Allemand, D. & Tambutte, S. (2008). Calcification and associated physiological parameters during a stress event in the scleractinian coral *Stylophora pistillata*. *Comparative Biochemistry and Physiology A – Molecular & Integrative Physiology*, Vol. 151, Issue 1, (September 2008), pp. 29–36, ISSN 1095-6433
- Park, H.; McGinn, P.J. & Morel, F.M.M. (2008). Expression of cadmium carbonic anhydrase of diatoms in seawater. *Aquatic Microbial. Ecology*, Vol.51, Issue2, (May 2008), pp. 183–193, ISSN: 0948-3055
- Park, H.; Song, B. & Morel, F.M.M. (2007). Diversity of the cadmium-containing carbonic anhydrase in marine diatoms and natural waters. *Environmental Microbiology*, Vol. 9, Issue 2, (Feb 2007), pp. 403–413, ISSN: 1462-2912
- Price, N.M. & Morel, F.M.M. (1990). Cadmium and cobalt substitution for zinc in a marine diatom. *Nature*, Vol. 344, Issue 6267, (April 1990), pp. 658–660, ISSN 0028-0836
- Rahman, M.W.; Davies, R.E.; Deyoe, C.W.; Reid, B.L. & Couch, J.R. (1961). Role of zinc in the nutrition of growing pullets. *Poultry Science*, Vol. 40, pp. 195-200, ISSN 0032-5791.

- Randall, D.J. & Daxbaeck, C. (1984). Oxygen and carbon dioxide transfer across fish gills. In: Hoar, W.S., Randall, D.J. (Eds.), *Fish Physiology*, vol. 10A. Academic Press, New York, pp. 263–314, ISBN 0-12-350430-9
- Roberts, S.B.; Lane, T.W. & Morel, F.M.M. (1997). Carbonic anhydrase in the marine diatom *Thalassiosira weissflogii* (Bacillariophyceae). *Journal of Phycology*, Vol. 33, Issue 5, (October 1997), pp. 845–850, ISSN 0022-3646
- Rowlett, R.S. (2010). Structure and catalytic mechanism of the β -carbonic anhydrases. *Biochimica et Biophysica Acta (BBA) - Proteins & Proteomics*, Vol.1804, Issue 2, (February 2010), pp. 362-373, ISSN: 1570-9639
- Schettino, T.; Trischitta, F.; Denaro, M.G.; Faggio, C. & Fucile, I. (1992). Requirement of HCO_3^- For Cl^- -Absorption in Seawater-Adapted Eel Intestine. *Pflugers Archiv-European Journal of Physiology*, Vol. 421, Issue 2-3, (June 1992), pp. 146-154, ISSN 0031-6768
- Schettino, T.; Lionetto, M.G. & Erroi, E. (2008). Enzymatic method for the detection of the toxicity of aqueous environmental matrices. Patent n. MI2008A008813, PCT/EP2008/064703
- Skaggs, H.S. & Henry, R.P. (2002). Inhibition of carbonic anhydrase in the gills of two euryhaline crabs, *Callinectes sapidus* and *Carcinus maenas*, by heavy metals. *Comparative Biochemistry and Physiology C-Toxicology & Pharmacology*, Vol. 133, Issue 4, (December 2002), pp. 605-612, ISSN 1532-0456
- Soto, M.; Ireland, M.P. & Marigómez, I. (2000). Changes in mussel biometry on exposure to metals: implications in estimation of metal bioavailability in “Mussel-Watch” programmes. *The Sciences of the Total Environment*, Vol. 247, Issue 2-3, (March 2000), pp. 175-187, ISSN 0048-9697
- Soyut, H.; Beydemir, Ş. & Hisar, O. (2008). Effects of Some Metals on Carbonic Anhydrase from Brains of Rainbow Trout. *Biological Trace Element Research*, Vol.123, Issue 1-3, (June 2008), pp.179–190, ISSN 0163-4984
- Supuran, C.T.; Scozzafava, A. & Casini, A. (2003). Carbonic anhydrase inhibitors. *Medicinal Research Reviews*, Vol.23, Issue 2, (March 2003), pp. 146– 189, ISSN: 0198-6325
- Supuran, C.T. & Scozzafava, A. (2007). Carbonic anhydrases as targets for medicinal chemistry. *Bioorganic & Medicinal Chemistry*, Vol. 15, Issue 13, (July 2007), pp. 4336–4350, ISSN: 0968-0896
- Supuran C.T., (2008). Carbonic anhydrases: novel therapeutic applications for inhibitors and activators. *Nature Reviews Drug Discovery*, Vol.7, Issue 2, (February), pp.168–181, ISSN 1474-1776
- Supuran, C.T. (2010). Carbonic anhydrase inhibitor. *Bioorganic & Medicinal Chemistry Letters*, Vol. 20, Issue 12, (June), pp. 3467-3474, ISSN: 0960-894X
- Tashian, R.E.; Hewett-Emmett, D.; Carter, N.D. & Bergenhem, N.C.H. (2000). Carbonic anhydrase (CA)-related proteins (CA-RPs) and transmembrane proteins with CA or CA-RP domains. In: Chegwiddden, W.R.; Carter, N.D. & Edwards, Y.H. (Eds.), *Carbonic anhydrase (CA)-related proteins (CA-RPs) and transmembrane proteins with CA or CA-RP domains. The Carbonic Anhydrases: New Horizons*. Birkhauser, Basel, pp. 105–120, ISBN 3-7643-5670-7
- Thompson, R.B. & Jones, E.R. (1993). Enzyme-based fiber optic zinc biosensor. *Analytical Chemistry*, Vol.65, Issue 6, March 1993), pp. 730-734, ISSN 0003-2700

- Thompson, R.B. & Patchan, M.W. (1995). Lifetime-Based Fluorescence Energy-Transfer Biosensing of Zinc. *Analytical Biochemistry*, Vol. 227, Issue 1, (May 1995), pp. 123-128, ISSN 0003-2697
- Thompson, R.B.; Maliwal, B.P. & Zeng H.H. (2000). Zinc biosensing with multiphoton excitation using carbonic anhydrase and improved fluorophores. *Journal of Biomedical Optics*, Vol. 5, Issue 1, (January 2000), pp. 17-22, ISSN 1083-3668
- Thompson, R.B.; Thompson, R.B. & Meisinger, J.J. (2002). Fluorescent zinc indicators for neurobiology. *Journal of Neuroscience Methods*, Vol. 118, Issue 1, (July 2002), pp. 63-75, ISSN 0047-2425
- Tripp, B.C.; Bell, C.B.; Cruz, F.; Krebs, C. & Ferry, J.G. (2004). A role for iron in an ancient carbonic anhydrase, *Journal of Biological Chemistry*, Vol. 279, Issue 20, (May), pp. 21677-21677, ISSN 0021-9258
- Tripp, B.C.; Smith K.S. & Ferry J.G. (2001)., *Journal of Biological Chemistry*, Vol.276, Issue 52, (December), pp. 48615- 48618, ISSN: 0021-9258
- Tu, C.; Silverman, D. N.; Forsman, C.; Jonsson, B.-H. & Lindskog, S. (1989). Role of Histidine-64 in the Catalytic Mechanism of Human Carbonic Anhydrase-II Studied with a Site-Specific Mutation. *Biochemistry*, Vol.28, Issue 19, (September 1989), pp. 7913-7918, ISSN 0006-2960
- Tu, C.K. & Silverman, D.N. (1985). Catalysis by cobalt(II)- substituted carbonic anhydrase II of the exchange of oxygen-18 between CO₂ and H₂O. *Biochemistry*, Vol.24, Issue 21, pp. 5881-5887, ISSN: 0006-2960
- Tu, C; Wynns, G.C. & Silverman, D.N. (1981). Inhibition by cupric ions of ¹⁸O exchange catalyzed by human carbonic anhydrase II. Relation to the Interaction Between Carbonic Anhydrase and Hemoglobin. *The Journal of Biological Chemistry*, Vol. 256, No.18, (September 1981), pp. 9466-9470, ISSN 0021-9258
- Vedani, A.; Huhta, D. W. & Jacober, S. P. J. (1989). Metal Coordination, H-Bound Network Formation, and Protein-Solvent Interactions in Native and Complexed Human Carbonic Anhydrase-I - a Molecular Mechanism Study. *Journal of the American Chemical Society*, Vol.111, Issue 11, (May 1989), pp. 4075-4081, ISSN: 0002-7863
- Viarengo, A.; Ponzano, E.; Dondero, F. & Fabbri, R. (1997). A simple spectrophotometric method for metallothionein evaluation in marine organisms: an application to Mediterranean and Antarctic molluscs. *Marine Environmental Research*, Vol. 44, Issue 1, (July 1997), pp. 69-84, ISSN 0141-1136
- Vitale, A.M.; Monserrat, J.M.; Casthilo, P. & Rodriguez, E.M. (1999). Inhibitory effects of cadmium on carbonic anhydrase activity and ionic regulation of the estuarine crab, *Chasmagnathus granulata* (Decapoda, Grapsidae). *Comparative Biochemistry and physiology C-Toxicology & Pharmacology*, Vol. 122, Issue 1, (January 1999), pp. 121-129, ISSN 1532-0456
- Wang, B.; Liu, C.Q. & Wu, Y. (2005). Effect of Heavy Metals on the Activity of External Carbonic Anhydrase of Microalga *Chlamydomonas reinhardtii* and Microalgae from Karst Lakes. *Bulletin of Environmental Contamination and Toxicology*, Vol. 74, Issue 2, (February 2005), pp. 227-233, ISSN 0007-4861
- Whitney, P.L. & Briggler, T.V. (1982). Membrane-associated carbonic anhydrase purified from bovine lung. *The Journal of Biological Chemistry*, Vol.257, (October 1982), pp. 12056-12059, ISSN 0021-9258

- Wu, Y.; Zhao, X.; Li, P. & Huang, H. (2007). Impact of Zn, Cu, and Fe on the Activity of Carbonic Anhydrase of Erythrocytes in Ducks. *Biological Trace Element Research*, Vol. 118, No. 3, (September 2007), pp. 227-232, ISSN 0163-4984
- Xu, Y.; Feng, L.; Jeffrey, P.D.; Shi, Y. & Morel, F.M. (2008). Structure and metal exchange in the cadmium carbonic anhydrase of marine diatoms. *Nature*, Vol. 452, Issue 7183, (March 2008), pp.56-U3, ISSN 0028-0836
- Yee, D. & Morel, F.M.M. (1996). In vivo substitution of zinc by cobalt in carbonic anhydrase of a marine diatom. *Limnology and Oceanography*, Vol.41, Issue 3, (May 1996), pp. 573-577, ISSN 0024-3590
- Zeng, H.H.; Thompson, R.B.; Maliwal, B.P.; Fones, G.R.; Moffet, J.W. & Fierke, C.A. (2003). Real-time determination of picomolar free Cu(II) in seawater using a fluorescence-based fiber optic biosensor. *Analytical Chemistry*, Vol.75, Issue 24, (December 2003), pp. 6807-6812, ISSN 0003-2700
- Zhang, B.Y.; Yang, F.; Wang, G.C. & Peng, G. (2010) Cloning and Quantitative Analysis of the Carbonic Anhydrase Gene from *Porphyra yezoensis*. *Journal of Phycology*, Vol. 46, Issue 2, (April 2010), pp.290-296, ISSN 0022-3646

IntechOpen



Biochemistry

Edited by Prof. Deniz Ekinçi

ISBN 978-953-51-0076-8

Hard cover, 452 pages

Publisher InTech

Published online 02, March, 2012

Published in print edition March, 2012

Over the recent years, biochemistry has become responsible for explaining living processes such that many scientists in the life sciences from agronomy to medicine are engaged in biochemical research. This book contains an overview focusing on the research area of proteins, enzymes, cellular mechanisms and chemical compounds used in relevant approaches. The book deals with basic issues and some of the recent developments in biochemistry. Particular emphasis is devoted to both theoretical and experimental aspect of modern biochemistry. The primary target audience for the book includes students, researchers, biologists, chemists, chemical engineers and professionals who are interested in biochemistry, molecular biology and associated areas. The book is written by international scientists with expertise in protein biochemistry, enzymology, molecular biology and genetics many of which are active in biochemical and biomedical research. We hope that the book will enhance the knowledge of scientists in the complexities of some biochemical approaches; it will stimulate both professionals and students to dedicate part of their future research in understanding relevant mechanisms and applications of biochemistry.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Maria Giulia Lionetto, Roberto Caricato, Maria Elena Giordano, Elisa Erroi and Trifone Schettino (2012). Carbonic Anhydrase and Heavy Metals, Biochemistry, Prof. Deniz Ekinçi (Ed.), ISBN: 978-953-51-0076-8, InTech, Available from: <http://www.intechopen.com/books/biochemistry/carbonic-anhydrase-and-heavy-metals>

INTeCH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen