

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

185,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Sustentable Use of the Wetting Agents in Protected Horticulture

Carlos Guillén and Miguel Urrestarazu
*Universidad of Almería,
Spain*

1. Introduction

The wetting agents in agriculture are materials that are classified within a larger group identified as adjuvants (Cahn and Lynn, 2000; Hazen, 2000; Kosswig, 2000; Thacker, 2003; Young, 2003; Lynn y Bory, 2005). According to their chemical nature, adjuvants are grouped into four families: surfactants, oils, inorganic salts and non-traditional adjuvants (figure 1). Nowadays some of the wetting agentes can be considered inside of called Green Chemistry (Carrasco and Urrestarazu, 2010).




Fig. 1. Classification of adjuvants according to their chemical nature and function

1.1 Surfactant

A common mistake is the use of adjuvant and surfactant terms interchangeably, which arises from the same discrepancy between producers (Young, 2003). The truth is that surfactants are a chemical family of adjuvants that improve emulsion, dispersion, foam, wet or other properties of a fluid to alter the characteristics of surface or interface and surface tension (Hazen, 2000, Thacker, 2003; Young, 2003). The term surfactant is derived from the contraction of three words Surface Active Agents (Thacker, 2003; Rosen, 2004, RAE, 2001).

Global consumption of surfactants in agriculture is 250 thousand tons per year, 180 thousand of which are incorporated in the formulation of phytosanitary products. 70 000 tonnes are used as tank mix adjuvants (Thacker, 2003). Besides agriculture surfactants are used in various products such as motor oils for cars, pharmaceuticals, in detergents, in materials used in oil exploration and floating agents in mining, in recent decades has extended its application to electronic printing, magnetic recording, biotechnology and viral research (Rosen, 2004).

Surfactants are molecules of low to middle molecular weight, it is amphipathic nature, it is containing a hydrophobic or lipophilic part (carbon chain) and a hydrophilic or lipophobic (Malmsten, 2002, Rosen, 2004; Tadros, 2005).

The hydrophobic groups are alkyl and alkylaryl groups mostly hydrocarbon, hydrophilic groups can be ionic or nonionic (Kosswig, 2000). According to this type ionic surfactants may be anionic, cationic, amphoteric and nonionic surfactants (Figure 2). In an anionic surfactant hydrophilic segment of the molecule forms an anion when dissolved in water. The opposite occurs in the cationic surfactant in which the active portion of the molecule in the hydrophilic segment is only a cation when dissolved in water. An amphoteric surfactant is capable of forming anions in aqueous solution depending on pH or cations. The nonionic surfactant is a surface active agent without ionic polar group is not ionized in aqueous solution (American Society for Testing and Materials E 1519, 1999, Cahn and Lynn, 2000; Kosswig, 2000; Malmsten, 2002; Rosen, 2004, Young, 2003, Lynn and Bory, 2005).




Fig. 2. Diagram of a humectant molecules

Reference	Physical properties of substrate			Growth and yield		Uptake of nutrients	Phytotoxicity
	Wettability	AV	WHC	EAW			
Sheldrake and Matkin (1971)							
Airhart <i>et al.</i> (1980)			increase				according surf. according dose
Wilson (1985)		increase					
Powell (1986)	increase	increase					according dose
Milkset <i>et al.</i> (1989)		increase	decrease				
Bhat <i>et al.</i> (1990)					better post-harvest	increase Ca ²⁺	
Bhat <i>et al.</i> (1992)							
Handreck (1992)			increase				according plant
Handreck and Black (1994)			increase				according dose
Elliot (1992)	increase		non effect				
Blodgett <i>et al.</i> (1993)	increase		non effect	increase			
Cid <i>et al.</i> (1993)					increase		
Blodgett <i>et al.</i> (1995)							
Riviere <i>et al.</i> (1996)	increase		increase	increase			according plant
Michel <i>et al.</i> (1997)	increase						
Reinikainen and Herranen (1997)	increase	increase					
Bilderback and Lorscheider (1997)		decrease	increase		increase		
Cid <i>et al.</i> (1998)			increase		increase	increase Ca ²⁺	
Guillen and Urrestarazu (2006)					increase	increase	
Urrestarazu <i>et al.</i> (2008)	increase		increase	increase			according dose

AV: air volume, WHC: water holding capacity, EAW: easily available water

Table 1. Summary of research on the effect of application of wetting agent in ornamental and horticultural cultures on substrate

The nonionic surfactants are chemically less active, but also less phytotoxic (Powell, 1986; Bures, 1997; Reinikainen and Herranen, 1997;) and less irritating than anionic and cationic (Malmnsten, 2002). They can with stand hard water and soluble in water and organic solvents, but can be sensitive to high temperatures (Rosen, 2004). Besides its critical concentration of micelle formation is twice lower than the anionic surfactants (Tadros, 2005). These nonionic surfactants are excellent wetting at very low concentrations (Thacker, 2003). They are the most dominant adjuvants of surfactants commercially available for the application of herbicides, they are less toxic to mammals (Young, 2003).

1.2 Wetting agent

Wetting agent is a substance used to reduce surface tension (Figure 3) and lead to better contact of a solution or suspension with a surface (WSSA Herbicide Handbook, 1994). Generally wetting agent is a surfactant, whose effectiveness is measured by the increase of spread of a liquid on a surface or the contact angle of liquid with the surface (Shurtleff and Averre, 1998).




Fig. 3. Contact angle and contact area for a drop water with and without a wetting agent on a hydrophobic surface

The wetting agents have been successfully used to improve the uniformity of wetting, improve relationships air: water of organic substrates and to improve nutrient availability to plants, without altering the physical characteristics of the substrate (Powell, 1986; Baht et al., 1992, Cid et al., 1992, Blodgett et al., 1993).

Crabtree and Gilkes (1999) indicate that the wetting agent also apply water repellent soils and have several benefits for the soil-plant system because it improves the soil wet releases

the fertilizer applied in the soil solution increases the mineralization of organic matter, wet consistency, improves water infiltration, reduces the flooded surface, reduces evaporation and increases the efficiency of water use in farming.

In trade, de Linan (2006), we found in the group of adjuvants to 30 products, among which are 17 active substances to wetting (Table 2).

Material	Coadjuvant kind
1 Alkyl ethoxylated fatty amines 73.5% w/v. SL	Surfactant nonionic
2 Alkyl ethoxylated fatty amine / propoxylated 48.7% w/v. SL	Surfactant nonionic
3 Alquil aminas grasas etoxiladas/propoxiladas 92% p/v. EC	Surfactant nonionic
4 Alquilpoliglicol 20% w/v. EC	Nonionic adjuvant
5 Alquilpoliglicol 44% + 1.5% sodium dioctyl w/v. SL	Surfactant nonionic
6 Sodium Alquiletersulfato 29% w/v. SL	Surfactant nonionic
7 Ethylene Alquilfenilhidroxipolioxo + 10.1% synthetic latex 45.45% w/v. EC	Surfactant nonionic
8 Montan wax 20% w/v. EW	Adherent-adjuvant and non-ionic wetting
9 ammonium dodecylbenzenesulfonate 20% w/v. SL	Adjuvant anionic wetting power
10 Ammonium dodecylbenzenesulfonate 20% w/v. EC	Adjuvant anionic wetting power
11 Sodium dodecylbenzenesulfonate 5% + nonilfeniloxietilenatosulfonado 2% w/v. SL	With wetting power
12 Methyl oleate / methyl palmitate 34.5% w/v. EC	It reduces the surface tension
13 Surfactant nonionic 20% p/v. SL	Surfactant nonionic
14 EterNonilfenolpolietilenglicolr 20% p/v. SL	Surfactant nonionic
15 Eter Nonilfenolpolietilenglicol 30% + inorganic acid 30% w/v. SL	With wetting power
16 Copper phthalate, are grouped a series of copper salts of fatty acids a dozen	With wetting power
17 Copper phthalate 66.5% w/v. EC	Adjuvant anionic wetting power

Source: De Liñán (2006)
EC: Emulsifiable concentrate, EW: Emulsion oil-water; SL: soluble concentrate, p/v: weight/volume

Table 2. Wetting adjuvant for agricultural use in Spain

2. Application of wetting agent in crop production

2.1 Assessment of toxicity

It is important before using a new wetting agent, evaluate their toxicity to plants (Handreck and Black, 1994). The bioassays used to assess the phytotoxicity of substrates (Zucconi et al., 1981, Ortega et al., 1996) were used to evaluate the toxicity of some products used in soilless culture (Urrestarazu and Mazuela, 2005). Urrestarazu et al. (2006) reported an adequate concentration of a surfactant with bioassays of tomato and cress.

2.2 Improving substrates

2.2.1 Wettability and water holding of growing media

The wettability of growing media (figure 4) is an important characteristic that could be limiting due to alternations of humectation and desiccation or drying accidental events that could modify considerably and reversibly or not the properties of the substrate during growing crops (Da Silva et al., 1993; Otten et al., 1999, Chambers and Urrestarazu, 2004, Lemaire et al., 2005).

Reduction of wettability contributes to vertical flow and goes against horizontal flow and water retention in growing media (Handreck and Black, 1994; Beeson and Haydu, 1995; Dekker and Ritsema, 1994, 1996 and 2000; Salas and Urrestarazu, 2004; Dekker et al., 2005).

Greater water retention capacity at low matrix potential is very important for optimal plant growth (Plaut et al., 1973, Plaut and Zieslin, 1974, Feigin et al., 1988; Raviv et al., 2002, Sahin et al., 2002, Chambers and Urrea, 2004). There have been several experiments to correlate the growth of plants with water retention and air capacity of the substrates and these has been found that the plant growth is highly correlated with water retention and low air capacity is associated with a low crop growth (Allaire et al., 1996). These problems of hydrophobia and low holding water in substrates can be overcome by application of wetting agent.



Fig. 4. Water repellency of dry organic substrates

It has been shown that surfactants facilitate the movement of water into and through the substrate, control the infiltration of water distribution and drainage, because it affects reserves of moisture, nutrient availability and aeration, with an optimal concentration of these surfactants improve rewetting potential of substrates and reduce root stress related problems, allowing greater control of plant growth (Powell, 1986, Bhat et al., 1990).




Fig. 5. The square of Humectant delete de red line behind, this must be like de red line as below

2.2.2 Improving physical properties of substrates

There is a considerable amount of articles about application of surfactants in ornamental and horticultural crops on different substrates. Table 1 shows some research on application of wetting agent on crop on substrate. Surfactants applied to organic growing media such as peat and pine fiber can improve absorption of water (wettability) of substrate (Handreck, 1992, Elliot, 1992; Blodgett et al., 1993, Riviere et al., 1996; Michelet al., 1997; Reinikainen and Herranen, 1997), also can increase air capacity, raising the drain (Wilson, 1985; Powell, 1986; Milksetal, 1989; Reinikainen and Herranen, 1997) or can improve the water holding capacity (Airhart et al., 1980; Handreck, 1992; Handreck and Black, 1994, Elliot, 1992; Bilderback and Lorscheider, 1997, Cid et al., 1998), increasing the water held in micropores. May also increase the water content readily available, without increasing the capacity of container (Blodgett et al., 1993) or increase the total available water content with increased water holding capacity (Riviere et al., 1996.). Urrestarazu et al (2008) reported that the effect of wetting agent added through fertigation is directly dependant on the substrate type evaluated and it can reduce the available water and increase the easily available water and total water holding capacity.

2.3 Increasing efficiency in ornamentals

Also can improve growth and production of ornamental species if applied at rates not phytotoxic (Bhat et al., 1990, Cid et al., 1993; Bilderback and Lorscheider, 1997, Cid et al., 1998), even allow to increase the availability and absorption of nutrients, especially calcium absorption (Bath et al., 1990, Cid et al., 1998). And apparently the application in fertigation could be the best method of application, less phytotoxic than when initially mixed with the substrate (Bhat et al., 1990).

2.4 Increasing efficiency in vegetables

In application by fertigation, the surfactant increased nutrients uptake, especially potassium ($P = 0.08$), nitrate ($P = 0.15$) and phosphate ($P = 0.25$), in melon crop on coir waste. Also it was observed a decrease in the percentage of drainage and reducing the emission of nitrates, phosphates, potassium and calcium ($P \leq 0.2$) in melon crop, showing its usefulness in lower environmental pollution (Guillen and Urrestarazu, 2006).

This observed in coir *mojantes* *mejoran* shows that the properties of the substrate water quality by reducing the hydrophobic effect acquired especially for use. But we have discussed in the previous section that this property is related to the concentration of *mojante* used, hence the application by fertigation should be adjusted to find the desired effect. In short, the goal is to improve efficiency, since the increased incorporation counterpart brings in reducing polluting emissions.

In tomato trial, the first year was observed higher yield in crop on reused coir waste and greater efficiency in water use ($P < 0.10$), proving useful for improving the use of water when a wetting agent is applied by fertigation. Also was observed a greater efficiency in the second year in reused coir waste ($P = 0.07$) and reused rockwool ($P = 0.03$) (Guillen and Urrestarazu, 2006).

It is inferred that the wetting by fertigation is a useful tool to adjust the water relations of substrates used, which is reflected in higher production volume of water consumed.




Fig. 6. Significant increases in the incorporation of potassium in melon cultivation of coconut fiber used. T0 = Control, T1 with humectant. FCN: New coir fiber, FCU: Used fiber coir. LRN: New rockwool, FRU: Used rockwool.




Fig. 7. Significant reduction in the emission of nutrients in a melon crop in reused coco fiber as a result of a wetting agent applied in fertigation. T0 = Control, T1 with humectant. FCN: New coir fiber, FCU: Used fiber coir. LRN: New rockwool, FRU: Used rockwool.




Fig. 8. Efficient water use in different substrates with and without wetting. T0= Testigo, T1 with humectant. FCN: New coir fiber, FCU: Used fiber coir. LRN: New rockwool, FRU: Used rockwool.

3. Acknowledgements

The authors acknowledge the collaboration of Ministry of Science and Innovation Spanish through the project FEDER AGL2010-18391.

4. References

- Airhart, D.L.; Ntarella, N.J. & Pokorny, F.A., (1980). Wetting a milled pine bark potting medium with surfactants. *Forest Products Journal* 30 (12), 30-33.
- Allaire, S.; Caron, J.; Duchense, I., Parent, L. & Rioux, J.A., (1996). Air filled porosity, gas relative diffusivity, and tortuosity, indices of *prunasxcistena* sp. Growth in peat substrates. *Journal of the American Society for Horticultural Science* 121(2), 236-242.
- American Society for Testing and Materials E 1519, (1999). Annual Book of ASTM Standards, Vol. 11.05. Designation E 1519-95, Standard Terminology Relating to Agricultural Tank Mix Adjuvants. pp. 905-906.
- Beeson, R.C. & Haydu, J. (1995). Cyclic microirrigation in container-grown land-scape plants improves plant growth and water conservation. *Journal of Environmental Horticulture* 13, 6-11.
- Bhat, N.R.; Tayama, H.K.; Prince, T.L., Prince, T.A. & Carver, S.A., (1990). Effect of Aqua-Gro wetting agent on the growth, flowering, and postproduction quality of potted chrysanthemum, *Ohio Florists' Association Bulletin* 723, 3-6.
- Bhat, N.R.; Prince, T.L.; Tayama, H.K. & Carver, S.A. (1992). Rooted cutting establishment in media containing wetting agent. *HotScience* 27(1), 78.
- Bilderback, T.E. & Lorscheider, M.R. (1997). Wetting agents used in container substrates are they BMP'S?. *ActaHorticulturae* 450, 313-319.
- Blodgett, A.M.; Beattie, D.J.; White, J.W. & Elliot, G.C. (1993). Hydrophilic polymers and wetting agents affect absorption and evaporative water loss. *HortScience* 28(6), 633-635.
- Burés, S. (1997). *Substratos*. Ediciones Agrotécnicas, Madrid, pp. 342. ISBN Madrid, Spain. 342 pg. ISBN 84-87480-75-6
- Cahn, A. & Lynn, J.L. (2000). Surfactants and deterative systems. *Kirk-Othmer Encyclopedia of Chemical Technology*, pp. 1142-1146.
- Carrasco G. & Urrestarazu M. (2010). Green Chemistry in Protected Horticulture: The Use of Peroxyacetic Acid as a Sustainable Strategy. *International Journal of Molecular Sciences*, 11(5), 1999-2009.
- Cid, M.C.; Socorro, A.R. & Pérez, R.L. (1993). Root growth and quality rating of *Schefflera* "Golden Capella" and *Ficus* "Starlight" on several peat-based substrates. *Acta Horticulturae* 342, 307-311.
- Cid, M.C.; Muñoz, C.R.; Socorro, A.R. and Gonzáles, T.G. (1998). Wetting agent effects on peat properties related to nutrient solution losses and plant growth. *Acta Horticulturae* 458, 161-165.
- Crabtree, W. & Gilkes, R. (1999). Banded wetting agent and compaction improve barley production on a water-repellent sand. *Agronomy Journal* 91, 463-467.
- Da Silva, F.F.; Wallach, R. & Chen, Y. (1993). Hydraulic properties of sphagnum peat moss and tuff (scoria) and their potential effects on water availability. *Plant Soil* 154, 119-126.
- Dekker, L.W. & Ritsema C.J. (1994). How water moves in a water repellent sandy soil 1, potencial and actual water repellency. *Water Resources Research* 30, 2507-2517.
- Dekker, L.W. & Ritsema, C.J. (1996). Preferential flow paths in a water repellent clay soil with grass cover. *Water Resources Research* 32(5), 1239-1249.


- Dekker L.W. & Ritsema, C.J. (2000). Wetting patterns and moisture variability in water repellent Dutch soils. *Journal of Hydrology* 231-232, 148-164.
- Dekker, L.; Oostinde, K. & Ritsema, C. (2005). Exponential increase of publications related to oil water repellency. *Australian Journal of Soil Research* 43, 403-441.
- De Liñán, C. (2005), *Vademécum de productos fitosanitarios y nutricionales*, 21^a ed. Ediciones Agrotécnicas S.L., pp 286-288. ISBN 84-87480-37-3, Madrid, Spain.
- Elliot, G.C. (1992). Imbibition of water by rookwool-peat container media amended with hydrophilic gel or wetting agent. *Journal of the American Society for Horticultural Science* 117(5), 757-761.
- Elliot, G.C. (1992). Imbibition of water by rookwool-peat container media amended with hydrophilic gel or wetting agent. *Journal of the American Society for Horticultural Science* 117(5), 757-761.
- Feigin, A.; Gammore, R. & Gilead, S. (1988). Response of rose plants to NO₃ or Cl salinity under different CO₂ atmosphere. *Soilless Culture* IV, 66.
- Handreck, K.A. & Black, N.D. (1994). *Growing Media for Ornamental Plants and Turf*. New South Wales University Press, pp. 401. ISBN 0-86840-333-4, Kensington, Australia
- Handreck, K. (1992). Wetting agents, how to achieve the best results. *Australian Horticulture*, August.
- Hazen, J.L. (2000). Adjuvants-Terminology, Classification, and Chemistry. *Weed Technology* 14, 773-784.
- Kosswig, K. (2000). Surfactants. *Ullmann's Encyclopedia of Industrial Chemistry*, pp. 293-368.
- Lemaire, F.; Dartigues, A.; Riviere, L.M.; Charpentier, S. & Morel, P. (2005). *Cultivos en macetas y contenedores principios agronómicos y aplicación*. INRA - MundiPrensa, pp. 210. ISBN: 84-8476-201-7, Madrid, Spain.
- Lynn, J.L. & Bory, B.H. (2005). Surfactants. *Van Nostrand's Encyclopedia of Chemistry*, pp. 1583-1586.
- Malmsten, M. (2002). *Surfactants and Polymers in Drug Delivery*. pp. 1-17. Marcel Dekker Incorporated, ISBN: 0-8247-0804-0, New York, USA.
- Michel, J.C.; Riviere, L.M.; Bellon-Fontaine, M.N. & Allaire, C. (1997). Effects of Wetting agents on the wettability of air-dried Sphagnum peats. *Peat in Horticulture* 2-7 Nov, 74-79.
- Milks, R.; Fonteno, W. & Larson, R. (1989). Hydrology of horticultural substrates, III. Predicting air and water content of limited-volume plug cells. *Journal of the American Society for Horticultural Science* 114(1), 57-61.
- Ortega, M.C.; Moreno, M.T.; Ordovás, J. & Aguado, M.T. (1996). Behaviour of different horticultural species in phytotoxicity bioassays of bark substrates. *Scientia Horticulturae* 66, 125-132.
- Otten, W.; Raats, P.A.C.; Challa, H. & Kabat, P. (1999). Spacial and temporal dynamics of water in the root environment of potted plants on a flooded bench fertigation system. *Netherlands Journal of Agricultural Science* 47, 51-65
- Plaut, Z.; Zieslin, N. & Arnon, Y. (1973). The influence of moisture regime on greenhouse rose production in various growth media. *Scientia Horticulturae* 1, 239-250.
- Plaut, Z. & Zieslin, N. (1974). Productivity of greenhouse roses following changes in soil moisture and soil air regimes. *Scientia Horticulturae* 2, 137-143.

- Powell, D. (1986). Wetting agents Tools to control water movement. Ohio Florists Assosiation Bulletin 681, 6-8.
- RAE (Real Academia Española), (2001). Diccionario de la Lengua Española, 22^a edición. La Real Academia Española, Madrid.
- Raviv, M.;Wallach, R.; Silber, A. & Bar-Tal, A. (2002). Substrates and their analysis. in: Savvas, D., Passam, H. (Eds.), Hydroponic Production of Vegetables and Ornamentals, Embryo Publications, ISBN 960-8002-12-5, Athens, Greece.
- Reinikainen, O. & Herranen, M. (1997). The influence of wetting agent on physical properties of peat. *Acta Horticulturae* 450, 375-379.
- Riviere, L.M. & Nicolas, H. (1987). Conduite de l'irrigation des cultures hors sol sur substrants, contraintes liees au choix des substrats. Milieux poreux et transferts hydriques. *Bull GFHN* 22, 47-70.
- Riviere, L.M. & Caron, J. (2001). Research on substrates: state of the art and need for the coming 10 years. *Acta Horticulturae* 548, 29-14.
- Rosen, M.J. (2004). Surfactants and interfacial phenomena, 3^a ed., John Wiley and Sons, Incorporated, pp, 1-33. ISBN 0471478180, Hoboken, NJ, USA.
- Sahin, U.; Anapali, O. & Ercisli, S. (2002). Physico-chemical and physical properties of some substrates used in horticultura. *Gartenbauwissenschaft* 67(2), 55-60.
- Salas, M.C. & Urrestarazu, M. (2004). Métodos de riego y fertirrigación en cultivo sin suelo. In: Urrestarazu, M., Tratado de cultivo sin suelo, 3^a edición, Ed. Mundi-Prensa, ISBN 84-8476-139-8, Madrid, España.
- Shurtleff, M.C. & Averre, C.W. (1998). Glossary of plant-pathological terms. APS Press, pp. 361. ISBN 0-89054-176-0, Minesota, USA.
- Tadros, T.F. (2005). Applied surfactants, principles and applications. Wiley- VCH Verlag GmbH & Co. KGaA, pp. 634. ISBN: 3-527-30629-3, Weinheimen, Germany.
- Thacker, J.R. (2003). Pesticide adjuvants. *Enciclopedia of Agrochemicals*, pp. 1199-1205.
- Urrestarazu, M. (2004), Tratado de cultivos sin suelo, 3^a ed. Mundi-Prensa, Servicio de Publicaciones de la Universidad de Almería, Madrid, España, pp. 914. ISBN 84-8476-139-8,, Madrid, Spain.
- Urrestarazu, M. & Mazuela, P.C. (2005a). Effect of slow-release oxygen supplí by fertigation on horticultural crops under soilless culture. *Scientia Horticulturae* 106, 484-490.
- Urrestarazu, M. & Mazuela, P.C. (2005b). La pujanza de los cultivos sin suelo en la horticultura protegida. *Vida Rural* 205, 25-28.
- Urrestarazu, M. & Guillén, C. (2006). Application of wetting agent by fertigation in horticultural crops on substrate. Tesis de Doctorado. Universidad de Almería, Spain.
- Urrestarazu, M.; Guillén, C.; Mazuela, P.C. & Carrasco, G. (2008). Wetting agent effect on physical properties on new and reused rockwool and coconut coir waste. *Scientia Horticulturae*, 116 (1) 104-108.
- Wilson, G. (1985), Effects of additives to peat on the air and water capacity, *Acta Horticulturae* 172, 207-209.
- WSSA Herbicide Handbook, 7th ed. (1994). Champaign, IL, Weed Science Society of America. pp. 313.
- Young, B.G. (2003). Herbicide adjuvants. *Enciclopedia of Agrochemicals*, pp. 707-718.

Zucconi, F.; Forte, M. & De Bertoldo, M. (1981). Biological evaluation of compost maturity. *BioCycle* (July/August), 27-29.

IntechOpen

IntechOpen



Horticulture

Edited by Prof. Alejandro Isabel Luna Maldonado

ISBN 978-953-51-0252-6

Hard cover, 172 pages

Publisher InTech

Published online 16, March, 2012

Published in print edition March, 2012

This book is about the novel aspects and future trends of the horticulture. The topics covered by this book are the effect of the climate and soil characteristics on the nitrogen balance, influence of fertilizers with prolongation effect, diversity in grapevine gene pools, growth and nutrient uptake for tomato plants, post-harvest quality, chemical composition and antioxidant activity, local botanical knowledge and agrobiodiversity, urban horticulture, use of the humectant agents in protected horticulture as well as post-harvest technologies of fresh horticulture produce. This book is a general reference work for students, professional horticulturalists and readers with interest in the subject.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Carlos Guillén and Miguel Urrestarazu (2012). Sustentable Use of the Wetting Agents in Protected Horticulture, Horticulture, Prof. Alejandro Isabel Luna Maldonado (Ed.), ISBN: 978-953-51-0252-6, InTech, Available from: <http://www.intechopen.com/books/horticulture/use-of-the-humectant-agents-in-protected-horticulture>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen