

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



The Type I and Type II Receptor Complexes for IL-4 and IL-13 Differentially Regulate Allergic Lung Inflammation

Nicola M. Heller, Preeta Dasgupta, Nicolas J. Dorsey,
Svetlana P. Chapoval and Achsah D. Keegan
*University of Maryland School of Medicine, Baltimore, MD,
USA*

1. Introduction

Approximately 300 million (M) people worldwide currently suffer from asthma; this number is projected to reach 400M by 2025 (Bahadori, et al., 2009). During the allergic immune response, inhaled allergens first stimulate epithelial cells, basophils, mast cells, and macrophages. This priming leads to the generation of allergen specific T-cells. Atopic asthma is strongly correlated with a robust CD4⁺ Th2 effector response, which results in elevated levels of the cytokines IL-4, 5, and 13. These cytokines act on multiple cells types to initiate and propagate the hallmark features of asthma such as pulmonary inflammation, periodic narrowing of airways, and mucus hypersecretion. Two of these cytokines, IL-4 and IL-13, share receptor chains and signaling proteins. In this chapter we discuss the cells that produce IL-4 and IL-13 including CD4⁺ T-cells and cells of the innate immune system, the structure of their receptors, their binding potency and kinetics, and their signal transduction. Furthermore, we present evidence for the differential effects of IL-4 and IL-13 acting via these receptor complexes on features of allergic lung inflammation. Finally, we discuss their contribution to the control of negative regulatory mechanisms that act to suppress allergic inflammation.

2. Cells that produce IL-4 and IL-13

Interleukin-4 and Interleukin-13 are closely related cytokines (Chomarat & Banchereau, 1998) critical to the development of T cell-mediated humoral immune responses, which are associated with allergy and asthma. Both cytokines display many overlapping functions (Chomarat & Banchereau, 1998). However, studies in cytokine-knockout mice and the use of blocking antibodies *in vivo* have shown that IL-4 is a critical cytokine for Th2 development whereas IL-13 plays critical roles in allergic asthmatic response (Grunig, et al., 1998, Wills-Karp, et al., 1998). Both cytokines can be secreted by many cell types but Th2 cells are considered to be the major producers. In this section, we will focus on cells capable of producing either IL-4, or IL-13, or both.

2.1 T cells

Upon antigen receptor stimulation, naive CD4⁺ T cells can differentiate into several different types, including T helper type 2 cells (Th2) depending on the cytokine milieu

present during the priming. IL-4 itself is important for the differentiation of naïve CD4⁺ T cells into Th2 cells capable of producing large amounts of IL-4, IL-13, and IL-5 (Zhu&Paul, 2008). Recent evidence points to basophils (van Panhuys, et al., 2011), mast cells (Plaut, et al., 1989), natural killer (NK) T cells (Akbari, et al., 2003), and γ/δ T cells (Ferrick, et al., 1995) as early producers of IL-4 in the innate immune response necessary for the optimal priming of the Th2 adaptive response. Therefore, in addition to CD4⁺ T cells, other T cell subpopulations such as NKT cells and γ/δ T cells are capable of making IL-4.

IL-4 and IL-13 producing NKT cells have been shown to be essential for the development and progression of allergic airway inflammation (Akbari, et al., 2003). NKT cell-deficient (Cd1d1^{-/-}) mice showed reduced Th2 responses after allergen challenge including allergic airway inflammation and airway hyperreactivity. The abrogated Th2 response in these mice could be restored by the adoptive transfer of purified NKT cells producing IL-4 and IL-13, but not by IL-4-deficient and IL-13-deficient NKT cells. IL-13 instillation to the mice could restore allergic airway responses. This led the authors to conclude that IL-4 and IL-13 produced by NKT cells potentiate the development of Th2 response in the lung. However, other studies suggest that NKT cells do not play an important role in allergic lung inflammation models (Das, et al., 2006).

Gamma/delta T cells also can differentiate into cells producing Th2 cytokines (Wen, et al., 1998). When WT mice were infected with *Nippostrongylus brasiliensis*, an extracellular parasite known to induce Th2 responses, γ/δ T cells from these mice produced IL-4 (Ferrick, et al., 1995). The ability of human γ/δ T cells to differentiate into Th2 cytokine producing cells was tested using stimulation of peripheral blood-derived γ/δ T cells with phosphoantigen isopentenyl pyrophosphate (Wesch, et al., 2001). When these cells were stimulated with Ag under Th2 priming conditions, they developed into cells producing IL-4. CD8⁺ T cells have been shown to produce IL-4 under specific *in vitro* stimulation (Seder, et al., 1992). When mouse CD8⁺ T cells were stimulated with immobilized CD3 in the presence of IL-2 and IL-4, they became high IL-4 producers. Moreover, both CD4⁺ T cells and CD8⁺ T cells separated from bulk lymph node T cell cultures with anti-CD3 plus IL-2 and IL-4 were equally effective in secretion of IL-4 after restimulation with anti-CD3 plus IL-2. The authors suggested that in certain *in vivo* conditions, CD8⁺ T cells could be major IL-4 producers. Human CD8⁺ T cell clones capable of IL-4 secretion were identified previously (Paliard, et al., 1988).

2.2 Granulocytes (Mast cells, basophils, and eosinophils)

Mouse non-T, non-B cells derived from spleen produce IL-4 and IL-13 in response to FcR crosslinkage (Ben-Sasson, et al., 1990). The same phenomenon was described for human bone marrow non-B and non-T cells in response to stimulation through either Fc ϵ or Fc γ receptors (Piccinni, et al., 1991). It has been suggested then that those IL-4-secreting cells could be either mast cells or basophils, or both. Indeed, the ability of mast cells to secrete both cytokines has been reported utilizing human cord blood derived mast cells (Toru, et al., 1998). These cells can generate both cytokines after stimulation with PMA or Fc ϵ R crosslinking. However, these cells do not produce Th2 cytokines spontaneously. Similarly, the ability of mouse mast cells to secrete IL-4 and IL-13 in response to specific immunological stimulation has been reported for fetal liver derived mast cell lines, bone marrow derived mast cells, and the mast cell line C1.MC/C57.1 (Brown, et al., 1987, Burd, et al., 1995). Of note, mast cells secrete relatively high IL-13 but low IL-4 levels.

As noted above, basophils were among three major cell populations of IL-4 producers in the lung under inflammatory conditions (Voehringer, et al., 2004). The ability of basophils to secrete both cytokines and other mediators of inflammatory response was extensively reviewed by Min and Paul in 2008 (Min & Paul, 2008). Importantly, it has been shown that peripheral blood basophils in asthmatic patients are the main producers of IL-4 and IL-13 (Schroeder, et al., 1995), suggesting a role in asthma exacerbation. IL-4 was detected in cultures of human basophils treated with diesel exhaust particles (Devouassoux, et al., 2002) suggesting that environmental exposure can predispose basophils to initiate Th2 responses. Eosinophils can also make IL-4 under certain circumstances. When mice were injected with *Schistosoma mansoni* eggs intraperitoneally, there were high levels of IL-4 in the peritoneal exudate cell cultures (Sabin, et al., 1996). IL-5 and eosinophils were necessary for the observed IL-4 production as suggested from similar experiments using egg-immunized IL-5^{-/-} mice or anti-IL-5 treated mice (Kopf, et al., 1996). Interestingly, these eosinophils were found to produce IL-4 early after immunization. The authors suggested that *Schistosoma mansoni* induced an early IL-5 production by mast cells that attracted eosinophils which, in their turn, produced IL-4 thus stimulating the development of antigen-specific Th2 cells. The ability of eosinophils to make Th2 cytokines was also tested in a more recent study (Voehringer, et al., 2004). The authors characterized IL-4 producing cells in the inflamed lungs by immunohistochemistry, flow cytometry, and microarray in mice with a bicistronic knock-in IL-4 gene linked via internal ribosomal entry site (IRES) with enhanced green fluorescent protein (eGFP). Eosinophils, basophils, and Th2 cells were reported as three cell populations producing IL-4 in these mice.

2.3 Myeloid cells

Macrophages have been shown to produce IL-4 or IL-13 in response to certain stimuli. A strong expression of IL-13 in the lung was observed in the experimental model of particle inhalation-induced inflammation (Kang, et al., 2005). Immunostaining of lung tissues of TiO₂-exposed mice demonstrated that alveolar macrophages are major producers of IL-13 and IL-25 in the inflamed lungs (Kang, et al., 2005). It has also been shown that human alveolar macrophages can produce IL-4 (Pouliot, et al., 2005). It has been shown that infectious pathogens including *Francisella tularensis* and respiratory syncytial virus (RSV) induce lung and peritoneal macrophages to produce IL-4 and IL-13 (Shirey, et al. 2008, Shirey, et al., 2010).

3. Receptor structure, ligand binding properties, and signal transduction

3.1 The structure of the IL-4 and IL-13 receptors

IL-4 and IL-13 elicit a wide variety of cellular responses by binding to high affinity receptor complexes expressed on the surface of cells. The IL-4/IL-13 receptor system is complex (**Figure 1**). The IL-4 specific receptor is composed of the IL-4R α chain paired with the common γ chain, or γ C, forming the Type I IL-4 receptor complex. The IL-4R α chain can also pair with the IL-13R α 1 chain, forming the Type II receptor (**Figure 1**). Type I receptors are activated by the binding of IL-4 to the ligand-binding IL-4R α chain and Type II receptors can be activated by either IL-4 or IL-13, with the IL-4R α or IL-13R α 1 acting as the initial ligand-binding chain, respectively. IL-13 can also engage another kind of IL-13 receptor, the IL-13R α 2 chain.




Fig. 1. **The IL-4 and IL-13 Receptor System.** A functional IL-4 receptor is composed of two transmembrane proteins. IL-4R α chain binds IL-4 with high affinity, leading to dimerization with either common gamma chain (γ C) or IL-13R α 1, forming the Type I or Type II receptor complex, respectively. IL-13 binds to IL-13R α 1 with lower affinity, followed by heterodimerization with IL-4R α to form a high affinity complex. IL-13 also binds to IL-13R α 2 (the so-called “decoy receptor”) at the cell surface, or in soluble form, but this interaction fails to activate the JAK/STAT pathway and generally it is thought to be inhibitory. Soluble forms of IL-4R α (sIL-4R α), IL-13R α 1 and IL-13R α 2 exist that can also bind ligand. Following ligand binding and heterodimerization, receptor-associated Janus Kinases (JAKs) are activated.

3.1.1 IL-4R α

The IL-4R α chain (CD124) is a 140 kDa protein. Human and mouse IL-4 receptors show a broad distribution on hematopoietic and non-hematopoietic cells (Park, et al., 1987, Lowenthal, et al., 1988), generally expressed at low levels (20-4000 receptors per cell). The IL-4R α cDNA was cloned from the mouse cytotoxic T-lymphocytic (CTLL-2) cell line (Mosley, et al., 1989) and from the human myeloid cell line, TF-1 (Galizzi, et al., 1990). Sequence analysis demonstrated that the IL-4R α belongs to the hematopoietin receptor superfamily. There are two extracellular structural features that characterize this family: type III fibronectin (FN) repeats and a membrane-proximal WSXWS motif.

The 2.3 Å resolution crystal structure of human IL-4 complexed to the extracellular domain of the human IL-4R α determined the overall shape of the two, linked type III FN-like domains, D1 and D2 (Hage, et al., 1999). Five IL-4R α peptide loops protrude from these two D domains and interact with IL-4. Binding of IL-4 to IL-4R α occurs through interaction of IL-4 with two clusters or sites (I and II) within the receptor, with Y183 and D72 at the center, surrounded by a shell of hydrophobic residues.

3.1.2 γ C

The involvement of the IL-2R γ subunit, or common γ chain, in forming heterodimeric IL-4 receptors was recognized by three groups in the early 90's (Kondo, et al., 1993, Leonard, et

al., 1994, Russell, et al., 1993). The γ C chain (CD132), which dimerizes with the IL-4R α chain to form a functional type I IL-4 receptor, is a 60 kDa protein. The extracellular domain of γ C possesses the tandem FN-III domains with four Cys residues and membrane-proximal WSXWS motif creating the classical CHR. Mouse γ C chain was cloned in 1993 (Kumaki, et al., 1993) and human γ C in 1992 (Takeshita, et al., 1992). Both human and mouse γ C genes map to the X-chromosome in humans (Noguchi, et al., 1993) and mouse (Cao, et al., 1993). The γ C subunit participates in the formation of many other cytokine receptor complexes including the IL-2, IL-7, IL-9, IL-15, and IL-21 receptors. Thus, mutations that either diminish or eliminate γ C expression or prevent association of the JAK3 kinase with γ C impair activity of these cytokines important for the development and proliferation of many different cells of the immune system, resulting in X-linked severe combined immunodeficiency (XSCID) in humans (Leonard, et al., 1994). γ C-deficient mice have no NK cells and severely diminished T- and B-cells, virtually absent lymph nodes and spontaneously develop inflammatory bowel lesions (Cao, et al., 1995).

3.1.3 IL-13 receptor α 1 (IL-13R α 1)

The IL-13R α 1 chain (CD213a1) is a 65-70 kDa glycosylated protein, encoded on the X chromosome in mice and humans. The mouse IL-13R α 1 cDNA was cloned in 1996 (Hilton, et al., 1996), followed by characterization of the human gene (Aman, et al., 1996), revealing the characteristic WSXWS motif and four conserved Cys residues. IL-13R α 1 can act either as a ligand-binding chain for IL-13 or as a dimerization partner to the type II receptor's IL-4-IL-4R α ternary complex (Zurawski, et al., 1993). The IL-13R α 1 chain is widely expressed on the surface of many hematopoietic and non-hematopoietic cells. It is through this Type II receptor complex that IL-4 and IL-13 mediate their effects on non-hematopoietic cells, which generally lack γ C, and therefore Type I receptor expression. IL-13R α 1 surface expression is absent on resting mouse and human T-cells and on mouse B-cells (Ogata, et al., 1998, Umeshita-Suyama, et al., 2000), although recent studies suggested inducible expression on mouse and human CD4⁺ T-cells (Newcomb, et al., 2009, Newcomb, et al., 2011). Related evolutionarily to γ C, IL-13R α 1 has acquired a third extra Ig-like domain, D1, allowing extra contacts with IL-13 in Type II receptor complexes (LaPorte, et al., 2008). This D1 domain is required for IL-13 binding and the formation of a functional Type II receptor, while it is not required for IL-4 binding or its activation of the Type II receptor (Ito, et al., 2009).

3.1.4 IL-13 receptor α 2 (IL-13R α 2)

IL-13 binds to a second IL-13 receptor, IL-13R α 2 (CD213a2, IL13BP) that was cloned in humans (Caput, et al., 1996) and from mice (Donaldson, et al., 1998). The gene is also found on the X-chromosome. IL-13R α 2 (~65 kDa) is inducibly expressed on fibroblasts, keratinocytes, epithelial cells, macrophages, and certain tumor cells and requires STAT6 for its expression (David, et al., 2003). The soluble form can be generated by proteolytic cleavage of the membrane-bound form by matrix metalloproteinases (MMPS) (Matsumura, et al., 2007) or by alternative splicing (Tabata, et al., 2006). Interestingly, sIL-13R α 2 is detected in serum from mice but not humans (Chen, et al., 2009). Treatment of cells with IL-4 or IL-13 in combination with TNF- α upregulated IL-13R α 2 cell surface expression (Zheng, et al., 2003). In contrast to IL-13R α 1, IL-13R α 2 binds to IL-13 with very high affinity 10⁻¹¹ M (Andrews, et al., 2002) one of the highest measured protein-protein interactions, possibly

due to an interlocking IL-13-binding interface (Lupardus, et al., 2010). The IL-13R α 2 is proposed to act as a “decoy receptor” for IL-13 (Yoshikawa, et al., 2003) and, more recently, for IL-4 signaling (Rahaman, et al., 2002). Consistent with this model, mice deficient in IL-13R α 2 have exaggerated IL-13 responses, such as severe liver fibrosis following *S. mansoni* infection (Chiaramonte, et al., 2003, Mentink-Kane, et al., 2004), reversible by soluble IL-13R α 2-Fc. A signaling role was hypothesized, however, in TNBS-induced colitis, tumor surveillance, and cancer (Strober, et al., 2009) and in monocytic cell lines through AP-1 (Fichtner-Feigl, et al., 2006).

3.2 Ligand binding properties

IL-4 binds to the IL-4 receptor α chain with high affinity with a K_d ranging between 20 - 300 pM (Lowenthal, et al., 1988), allowing ligand binding at low IL-4 concentrations, as would be present in the initiating phase of an allergic inflammatory response. There is species specificity for the IL-4:IL-4R α interaction (Park, et al., 1987), yet dimerization of the binary IL-4:IL-4R α complex with the γ C chain is not species specific (Idzerda, et al., 1990). Dimerization with the γ C chain forming Type I IL-4 receptors increases the affinity of IL-4 binding approximately three-fold (Russell, et al., 1993). The affinity of interaction of the binary IL-4:IL-4R α complex with either of its dimerization partners, γ C or IL-13R α 1, is low (559 nM for γ C with IL-4:IL-4R α (Andrews, et al., 2006, LaPorte, et al., 2008, Zhang, et al., 2002) and 487 nM for IL-13R α 1 (LaPorte, et al., 2008)). In contrast, IL-13 binding to IL-13R α 1 is a relatively low affinity interaction ($K_d \sim 30$ nM (Andrews, et al., 2002, LaPorte, et al., 2008)) that is not species-specific (Andrews, et al., 2001). Dimerization of IL-13:IL-13R α 1 with IL-4R α to form the ternary Type II complex is a high affinity, species-specific interaction (Andrews, et al., 2001, LaPorte, et al., 2008).

The crystal structures of the three ternary complexes were solved in 2008 (LaPorte, et al., 2008) revealing that the IL-4/IL-13 receptor system was unique in that when forming a Type II receptor the “binder” (i.e. ligand-binding) and “trigger” (i.e. dimerizing partner) chains were switched depending on whether IL-4 or IL-13 is the ligand. Furthermore, there was a $\sim 8^\circ$ angle difference in the position of the IL-4R α chain relative to the IL-13R α 1 chain between the two Type II structures. The impact of these subtle differences in initial binding affinities, order of chain assembly, and 3-dimensional structure of the extracellular domains of the ternary complexes on responsiveness to IL-4 and IL-13 is not clear. A recent study suggests that the relative abundance of the two receptor types and ratio of the ligand-binding chain to the trigger chain can fine-tune sensitivity to these cytokines (Junttila et al., 2008).

3.3 Signal transduction pathways activated by Type I and Type II receptor engagement

3.3.1 Janus kinases

Ligand-induced dimerization of the Type I or Type II IL-4 receptor activates receptor-associated kinases of the Janus kinase (JAK) family. JAK1 was activated by IL-4 and IL-13 (Welham, et al., 1995). IL-13 did not induce JAK3 activation, as the IL-13R α 1 does not recruit JAK3 (Keegan, et al., 1995, Welham, et al., 1995). JAK2 appeared to be constitutively associated with the IL-4R α chain in human monocytes and stimulation of the Type II receptor complex with IL-13 enhanced the interaction (Roy, et al., 2002). IL-13 predominantly activates TYK2 (Murata & Puri, 1997), as well as JAK2 (Murata, et al., 1996). JAK1 is a substrate for other non-JAK kinases that can affect the activation of IL-4-induced

signaling. PKC ζ is required for full IL-4-induced JAK1 activation (Martin, et al., 2005). Mutational studies on cytoplasmic domain of IL-4R α have revealed the presence of a membrane-proximal, proline-rich “box 1” motif (aa262 – 267 for hIL-4R α and 263 – 271 for mIL-4R α) to which JAK1 can bind (Fujiwara, et al., 1997, Russell, et al., 1994). The cytoplasmic domain of the γ C also has a “box 1” motif :Pro-X-Pro and a preceding cluster of hydrophobic amino acids (aa286 – 294 that binds JAK3, Murakami, et al., 1991)).

The proline-rich region in the IL-13R α 1 (aa 373 – 378 in mouse and aa 376 – 381 in human) and the next six amino acids downstream mediate the interaction between IL-13R α 1 and the associated JAKs (TYK2, JAK1 in a transfected FDCP-1 cell line (Orchansky, et al., 1999)). The cytoplasmic domain of the IL-13R α 1 chain is shorter than that of the IL-4R α or γ C. It contains the box 1 motif and two tyrosines, Y402 and Y405, which act as STAT3 binding motifs (Orchansky, et al., 1999, Umeshita-Suyama, et al., 2000). Engagement of the receptor chain by IL-13 activates JAK1 and TYK2 and triggers a variety of signaling cascades which will be discussed below.

Tyrosine residues within the cytoplasmic domains of the Type I and Type II receptor subunits are targets for rapid phosphorylation by the JAKs (**Figure 2**). Both IL-4 and IL-13 induce tyrosine phosphorylation of IL-4R α (Wang, et al., 1992). Tyrosine phosphorylation of IL-13R α 1 was not detected in immunoprecipitated lysates from IL-13-stimulated FD-5 cells transfected with the IL-13R α 1 subunit (Orchansky, et al., 1999) although mutational studies suggested that IL-13R α 1 tyrosines would indeed become phosphorylated (Umeshita-Suyama, et al., 2000). Studies using deletion, Y-to-F mutants, and chimeric forms of the IL-4R α (Deutsch, et al., 1995, Keegan, et al., 1994, Koettnitz & Kalthoff, 1993, Seldin & Leder, 1994) have characterized distinct regions of the IL-4R α cytoplasmic domain containing five essential tyrosine residues (Y1-Y5). These phosphotyrosines become docking sites for SH2- and PTB-containing proteins: Y1 (Y497) recruits IRS proteins, Y2-Y4 (Y575, Y603, Y631) recruit STAT6 and Y5 (Y713) is bound by the phosphatases, SHP-1, -2 and SHIP (Hanson, et al., 2003, Kashiwada, et al, 2001).

3.3.2 IRS proteins

IL-4 strongly induced the tyrosine phosphorylation of a ~170 kDa protein in the mouse IL-3-dependent hematopoietic cell line, FDCP-2 (Wang, et al., 1992), named IL-4-induced phosphotyrosine substrate (4PS) and it associated with PI3-K and PI3-K activity. This protein was identical to that tyrosine phosphorylated in response insulin and IGF-I in FDC cells (Wang, et al., 1993). 4PS was cloned from myeloid progenitor cells and renamed IRS-2 due to similarity to IRS-1 (Sun, et al., 1995). IRS-1 is also tyrosine phosphorylated in response to IL-4 stimulation and whether IRS-1 or IRS-2 or both are tyrosine phosphorylated after IL-4 stimulation depends on cellular expression of each protein and the surface expression of Type I or II receptors (Sun, et al., 1997). IRS-2 is predominantly found in hematopoietic cells and IRS-1 in non-hematopoietic cells: studies in 32D cells, which express neither IRS protein, revealed the role of both IRS-1 and IRS-2 in IL-4-induced cellular proliferation (Wang, et al., 1993).

The sequence of amino acids important for IRS binding to the IL-4 receptor was determined by truncation mutational analysis (between aa437 and 557). Within this interval, there is a homologous sequence that binds IRS proteins in the insulin and IGF-I receptor, known as the insulin and IL-4 receptor (I4R) motif, whose sequence is ⁴⁸⁸PL-(X)₄-NPXYXSXSD⁵⁰². The central tyrosine when phosphorylated is critical for association of the PTB domain of IRS proteins with the I4R motif of the IL-4R α (Keegan, et al., 1994, Zhou, et al., 1996).




Fig. 2. **Signaling by the Type I and Type II receptors.** The signaling pathways activated by the Type I and Type II receptor complexes are shown in cartoon form. The *font size* is an indication of the *relative strength* of activation of IRS-2, STAT6, and SHP1 by the IL-4 or IL-13 receptor complexes. Differences in signaling by the receptor complexes are highlighted. A. Type I Receptor. IL-4 (yellow), IL-4Rα (blue) and γc (red). IL-4Rα binds JAK1 (pink) while γc binds JAK3 (rust). γc associates with Nedda4. B. Type II Receptor bound to IL-4. IL-4 (yellow), IL-4Rα (blue), and IL-13Rα1 (green). The D1 domain of IL-13Rα1 is shown. IL-13Rα1 binds to TYK2 (green). IL-13Rα1 recruits STAT3 (orange). C. Type II Receptor bound to IL-13. IL-13 (orange), IL-13Rα1 (green), IL-4Rα (blue). The relative ability of these receptor complexes to activate Shc or the MAP kinase pathways is unclear.

The IRS proteins are tyrosine phosphorylated in response to engagement of the IL-4 receptors. JAK1 is required for this to occur (Wang, et al., 1997). Once phosphorylated, tyrosines that are part of typical SH2-binding motifs (Sun, et al., 1991) provide docking sites for a variety of different SH2-domain-containing downstream molecules, such as the p85 subunit of PI3-K and Grb2 (Pruett, et al., 1995). There are three tyrosines, part of classical YXXM motifs, that act as p85 binding sites in IRS-1 and two in IRS-2 (White, 2002). Binding of p85 can activate PI3-K thus allowing IL-4 to initiate a large number of downstream signaling cascades. The IRS proteins can also bind SHP-2 following IL-4 stimulation and IRS-2 can recruit PLC-γ in response to IL-13 (Sozzani, et al., 1998). The IRS proteins also interact with the SOCS proteins that are negative regulators of IL-4 signaling and will be discussed in more detail in a later section.

The contribution of IRS-2 to allergy and asthma is not well understood. Transgenic overexpression of IRS-2 enhanced IgE production *in vivo*, and increased IL-5 secretion from *in vitro* differentiated CD4⁺ Th2 cells (Kelly-Welch, et al., 2004). Early studies of T-cells isolated from IRS2^{-/-} mice found reduced T-cell proliferation and IL-5 production by Th2 cells compared to wildtype T-cells (Wurster, et al., 2002). Surprisingly, mice with a mutation in the IRS-2 docking site of the IL-4Rα (Y500F) demonstrated enhanced allergic inflammation, suggesting a significant contribution of this region of the IL-4Rα to inflammation control *in vivo* (Blaeser, et al., 2003). Activation of the IRS-2 pathway was abrogated but this Y500 region of the IL-4Rα also recruits a number of other signaling molecules including Shc, FRIP1, p62DOK, and p85β (Nelms, et al., 1998) that may negatively regulate the pathway (described below).

3.3.3 Activation of PI3-K

The p85 or regulatory subunit of PI3-K binds phosphotyrosines on the IRS protein via SH2-domains and the resulting conformational change releases inhibition of the enzymatic activity of the p110 (catalytic) subunit and allows it to translocate to the plasma membrane. Activation of PI3K activity in response to IL-4 was first demonstrated in hematopoietic FDCP cells inducing mitogenic signals (Wang, et al., 1992). The kinase transfers a phosphate group from ATP to phosphoinositol (PI) to rapidly form PIP3, activating a myriad of downstream pathways. PIP3 has the potential to activate protein kinase C (PKC) and protein kinase B (PKB)/Akt. Activation of Akt in response to IL-4 has been shown in human eosinophils (Coffer, et al., 1998), although we found no induction of phosphorylation on Akt^{Ser473} in mouse eosinophils by IL-4 or IL-13 (Heller et al, under review).

3.3.4 Signal Transducers and Activators of Transcription (STATs)

IL-4 receptor engagement can activate a number of members of the STAT family. STAT6 is the predominantly activated STAT but other members of the STAT family can also be activated to a lesser degree. The human STAT6 gene was cloned in 1994 and the same group determined that STAT6 (IL-4 Stat) directly interacted with the IL-4 receptor cytoplasmic domain, homodimerized via its SH2-domain and characterized the DNA binding motif recognized by STAT6 (Hou, et al., 1994). STAT6 docks via its highly conserved SH2-domain to the “gene regulation domain” (aa 557 – 657) encompassing three of the five conserved tyrosines of human IL-4R α (Ryan, et al., 1996). STAT6 becomes tyrosine phosphorylated on Y641 and forms homodimers through pY641-SH2 interactions (Mikita et al., 1996). The C-terminus of STAT6 contains the transcriptional activation domain (Goenka, et al., 1999). In addition to tyrosine phosphorylation, the STAT6 protein can be post-translationally modified in other ways to affect its function: methylation (Chen, et al., 2003), serine phosphorylation (Pesu, et al., 2000) on S756 (Wang, et al., 2004) and S707 (Shirakawa, et al., 2011) and acetylation (Shankaranarayanan, et al., 2001).

Once in the nucleus, STAT6 homodimers bind to consensus DNA motifs in STAT6-responsive genes (reviewed in Goenka & Kaplan, 2011). The preferred DNA binding motif recognized by STAT6 is a dyad symmetric recognition element TTC-GAA separated by four nucleotides although STAT6 can also bind the dyad element separated by three nucleotides. STAT6 often co-operates with other transcription factors and co-activators to activate transcription including NK- κ B, CEBP β , CBP and p300 and p160 steroid receptor nuclear coactivator (NCoA-1).

STAT6 deficiency (Takeda, et al., 1996) is protective in many different *in vivo* models of allergy including allergic airway disease, food allergy, eosinophilic esophagitis and atopic dermatitis. In contrast, mice expressing constitutively active STAT6 are predisposed to an allergic phenotype (Sehra, et al., 2008, Sehra, et al., 2010). Hyperactive STAT6 can lead to cellular transformation and various cancers, due to dysregulated p27^{Kip}/cell cycle progression (Bruns, et al., 2003). Mice deficient in STAT6 have compromised expulsion of helminth parasites: they cannot produce Th2-cells, mount an effective IgE response, produce mucus or chemokines (Kaplan, et al., 1998).

While STAT6 is clearly the dominant STAT family member activated by IL-4 and IL-13, there are reports of activation of several other STATs to varying degrees. Some STAT1 α activation was documented in the mouse T-helper cell line, HT-2, in response to IL-4 (Brunn, et al., 1995) and by IL-4 and IL-13 in five primary human cell types generally to a lesser

degree than STAT6 (Wang, et al., 2004). Primary human monocytes respond to IL-13 with tyrosine phosphorylation of STAT1a (Roy, et al., 2002). Furthermore, STAT5 phosphorylation was detected in human B-cells in response to IL-4 and IL-13 (Rolling, et al., 1996) and in primary human monocytes by IL-13 (Roy, et al., 2002). The role of STAT1 and STAT5 in IL-4- or IL-13-induced responses is unknown.

The cytoplasmic domain of IL-13R α 1 contains two STAT3 binding motifs (Y402 and Y405, (Orchansky, et al., 1999)). STAT3 activation by IL-4 and IL-13 is dependent upon expression of the IL-13R α 1 (Orchansky, et al., 1999) and occurs in response to IL-4 and IL-13 in human B-cells (Rolling, et al., 1996) and weakly in HMVEC-L and NHLF cells (Wang, et al., 2004). STAT3 phosphorylation was induced by IL-13 in primary human monocytes (Roy, et al., 2002) and by IL-4 in keratinocytes (Wery-Zennaro, et al., 1999). We have observed the relatively weak induction of STAT3 phosphorylation by IL-4 and IL-13 in murine bone-marrow-derived macrophages and the human lung adenocarcinoma cell line A549 (LaPorte, et al., 2008). The specific function of STAT3 in mediating responses to IL-4 and IL-13 is unclear. A recent report demonstrated that STAT3 played a role in Th2 differentiation, however the cytokines responsible for STAT3 activation in that setting were thought to be IL-6 or IL-21 (Stritesky, et al., 2011).

3.3.5 Other pathways activated by IL-4R α (Ras/MAPK, Shc, Dok)

Activation of the Ras/MAPK pathway is not generally observed in response to IL-4 despite IRS activation leading to interaction with the adapter molecule, Grb2 (Pruett, et al., 1995, Wang, et al., 1995). Since Grb2 is constitutively associated with the guanine nucleotide exchange protein, SOS, that catalyzes exchange of GDP bound to Ras for GTP, it is often assumed that IL-4 will trigger the phosphorylation cascade of Raf/MEK/ERK-1/-2. However, IL-4 does not activate p21ras (Duronio, et al., 1992, Satoh, et al., 1991, Welham, et al., 1994), Raf1 or ERK1/2 (Welham, et al., 1992, Welham, et al., 1994). We too have been unable to detect ERK-1/2 activation in response to IL-4 or IL-13 in primary mouse bone marrow-derived macrophages (Heller, et al., 2008) or peripheral blood eosinophils. Activation of the Ras/MAPK pathway was demonstrated to enhance IL-4 signaling, possibly through MEK phosphorylation of JAK1 and STAT6 (Yamashita, et al., 1999).

Activation of Shc can be linked to Ras activation, via the Grb2 adapter-Sos interaction. There are three widely-expressed Shc proteins (~46, 52 and 66 kDa) containing a C-terminal SH2 domain, regions homologous to the α 1 chain of collagen (Pelicci, et al., 1992) and an N-terminal PTB domain, similar in structure to IRS protein PTB domain (Zhou, et al., 1996). Shc moves to the plasma membrane and docks to the phosphorylated I4R motifs on activated IL-4 receptors via its SH2 and PTB domains (Wolf, et al., 1995). Phospho-Shc is then bound by the SH2/3 motifs of the Grb2 adaptor and bound Sos is activated. Shc activation in response to IL-4 appears to be dependent on cell type (Crowley, et al., 1996, Wery, et al., 1996, Welham, et al., 1994).

Two hematopoietically-expressed members of the Dok proteins play a role in IL-4R-induced responses. The N-terminal PH and PTB domains of the Dok proteins suggest that they could bind phosphotyrosines in membrane-localized receptors (Mashima, et al., 2009). Indeed, Dok-2 (also known as FRIP, Dok-R or p56^{dok}) has been shown to interact with the I4R motif of the IL-4R α (Nelms, et al., 1998). T-cells from Dok-2-deficient mice (the *hairless* allele, *hr/hr*) have an increased proliferative response to IL-2 and IL-4 (Nelms, et al., 1998). Studies of Dok-1 (p62^{dok})-deficient cells suggest this protein plays a positive role in sustaining IL-4

signaling responses (IL-4-induced T-cell proliferation and CD23 expression and IgE class switching in B-cells (Inoue, et al., 2007)). The Dok proteins have multiple docking sites for SH2-containing proteins in their C-terminus and thus act as adaptor proteins. Because both Dok proteins bind RasGAP, which inactivates Ras by hydrolysis of GTP, it is thought that they might be negative regulators of the Ras/MAPK pathway. Recruitment of the Dok proteins to the IL-4R α may explain the lack of Ras activation by IL-4 in some cell types.

3.4 Negative regulation of receptor signaling

3.4.1 SHP-1 and SHP-2

Protein tyrosine phosphatases (PTP) that remove phosphate groups from phosphotyrosine residues that are activating signals can downregulate the signaling cascades initiated by IL-4. SHP-1 and SHP-2 are PTP molecules containing two N-terminal SH2-domains, a single central phosphatase domain and C-termini with two potential tyrosine phosphorylation sites that affect activity. The SH2-domain targets the phosphatase to a particular cellular location and binds to and inhibits the catalytic domain when the phosphatase is not bound to substrate. The expression of SHP-1 is restricted to hematopoietic cells (Yi, et al., 1992) and low expression is also found in epithelial cells. SHP-2, on the other hand, is widely expressed including in cells that express SHP-1.

SHP-1 may positively (White, et al., 2001) or negatively (Imani, et al., 1997) regulate IL-4 responses, depending upon the cell type (Huang, et al., 2005). Interestingly, SHP-1 constitutively associates with the IL-4R α chain even in resting lymphocytes (Huang & Paul, 2000). Subsequent studies have indicated that SHP-1 also negatively regulates signaling responses to IL-13 by downregulating the phosphorylation of STAT6 (Haque, et al., 1998).

SHP SH2-domains bind to ITIM sequences [I/V/L]xY(p)xx[I/V/L] in activated receptors (Ravetch & Lanier, 2000). The cytoplasmic domain of the IL-4R α possesses a putative ITIM surrounding the fifth tyrosine, Y713 (**Figure 2**) and a variety of studies have indicated this motif may interact with SHP-1, SHP-2, Shc and SHIP (Kashiwada, et al., 2001, Hershey, et al., 1997, Kruse, et al., 2002). Our group showed that Y713 in human IL-4R α mediated recruitment of SHP-1 (Hanson, et al., 2003) and SHIP (Zamorano & Keegan, 1998) to the IL-4 receptor complex. Knock-in of a Y713F mutant of IL-4R α in mice resulted in enhanced STAT6 phosphorylation, IgE production, and allergic lung inflammation (Tachdjian, et al., 2010). Loss of Y713 had a more dramatic effect on the magnitude of responses to IL-13 *vs.* IL-4. Taken together these results suggest that IL-4 and IL-13 signaling is modulated by SHP1 or other phosphatases capable of binding to Y713, and that this modulation may be more profound for the Type II receptor complex (**Figure 2**).

While tyrosine phosphorylation of SHP-1 increases its phosphatase activity, tyrosine phosphorylation of SHP-2 has been proposed to allow this molecule to function as an adaptor protein by providing docking sites for other SH2-domain-containing proteins (Lorenz, 2009). Conflicting data describe that SHP-2 was (Kruse, et al., 2002, Wang, et al., 1999) or was not (Gadina, et al., 1999) tyrosine phosphorylated in response to IL-4 stimulation. IL-13 stimulation of PBMC also induced tyrosine phosphorylation of SHP-2 (Kruse, et al., 2002). SHP-2 interacts with IRS-1 (Xiao, et al., 2002), JAK1, JAK3 and coprecipitates with Grb-2 and p85 after cytokine stimulation (Gadina, et al., 1998, Kuhne, et al., 1993). Peptides derived from the IL-4R α (aa 545–558) were able to pull down SHP-2 from lysates of IL-13-stimulated PBMC (Kruse, et al., 2002).

Hematopoietically-expressed SH2-domain-containing inositol 5'-phosphatase (SHIP) dephosphorylates PIP3, the product of the PI3-K enzyme, to form PIP2. IL-4 stimulation can induce tyrosine phosphorylation of SHIP, suggesting that SHIP can dock to multiple sites in the IL-4R α (Zamorano & Keegan, 1998). SHIP-1-deficient mice spontaneously developed allergic lung inflammation, have increased mast cells that spontaneously released histamine indicating a potential homeostatic role for SHIP-1 in regulating Th2-responses *in vivo* (Oh, et al., 2007). Recent exciting data have defined a role for SHIP-1 in the skewing of macrophage phenotype (Rauh, et al., 2005).

3.4.2 Suppressors of Cytokine Signaling (SOCS)

The suppressor of cytokine signaling (SOCS) proteins are a family of cytokine-induced negative regulators of cytokine signaling (Starr, et al., 1997, Yoshimura, et al., 2003). The general structure of the SOCS protein includes a central SH2-domain, critical for binding to their tyrosine phosphorylated substrates, and a C-terminal SOCS box that mediates ubiquitin-dependent proteolysis. SOCS-1, -3 and CIS are induced by IL-4 and SOCS-1 and -3 were shown to inhibit IL-4 signaling transduction (Haque, et al., 2000, Losman, et al., 1999). SOCS-1 and SOCS-3, both of which are induced by IL-4, have an additional kinase inhibitory region that functions as a pseudosubstrate to inhibit JAK activity (Yasukawa, et al., 1999). Another mechanism of action is by SOCS interaction with the phosphorylated tyrosines within the cytoplasmic domains of the receptor. SOCS-3 directly interacted with IL-4 α (O'Connor, et al., 2007). SOCS proteins can also target activated signaling intermediates to the proteasome. SOCS-1 is able to regulate the half-life of JAKs and insulin/IGF-I-induced IRS-2 (Rui, et al., 2001) in this manner.

4. Differential roles for IL-4 and IL-13 acting via the Type I or Type II receptors on features of allergic lung inflammation

Recent evidence suggests that even though IL-4 and IL-13 share receptor components and signaling proteins, and elicit overlapping responses *in vitro*, they can elicit different functional responses *in vivo*. IL-4 is primarily responsible for regulating Th2 development and inflammation while IL-13 is responsible for effector activities such as airway hypersensitivity, collagen production, and mucus hypersecretion (Gavett, et al., 1994, Pernis & Rothman, 2002, Wills-Karp, et al., 1998). The molecular basis for this variation is not understood clearly, since both IL-4 and IL-13 use the Type II receptor complex. It has been postulated that differences in the relative abundance of the Type I or Type II receptor subunits in different cell types may be responsible for the differences in responses elicited by IL-4 versus IL-13. Certainly the presence or absence of individual receptor subunits and appropriate Janus kinases in each cell determines whether a cell can respond to IL-4 or IL-13. However, many of the cell types involved in the effector activities express the Type II receptor that is activated by both IL-4 and IL-13.

4.1 Different receptor signaling pathways utilized by IL-4 and IL-13 via the Type I and Type II receptors

One reason proposed for differential functions of these cytokines is the observed differences in the amounts of IL-4 and IL-13 produced in tissues during Type II inflammation. Various reports have shown that IL-13 is secreted by large number of cell types and in much greater quantities than IL-4 during Th2 responses in both asthma patients (Huang, et al., 1995) as

well as mouse models of this disease (Munitz, et al., 2008). However, analysis of the binding affinities of the Type I and Type II receptors with their respective ligands have shown that the relative amounts of each cytokine does not necessarily explain the functional differences between IL-4 and IL-13.

As discussed above, LaPorte and colleagues have shown that although IL-4 binding to the IL-4R α chain occurs with high affinity, complex formation of IL-4: IL-4R α with γ C or IL-13R α 1 is quite unstable and inefficient (LaPorte, et al., 2008). On the other hand, IL-13 binds to IL-13R α 1 with low affinity, but the interaction of IL-13: IL-13R α 1 with IL-4R α is more favorable and stable. As a result even at very low concentrations, IL-4 is able to mediate efficient and rapid STAT6 phosphorylation via Type I and Type II receptors, while cells have to be stimulated with much higher concentrations of IL-13 and for a longer time to obtain similar responses via the Type II receptor. Since the IL-4 bound complexes are less stable, LaPorte *et. al.* proposed that when expression of receptor chains in cells become limiting, IL-4 responses would be limited, while IL-13 responses would take over.

However, experiments using transgenic overexpression of large quantities of IL-4 or IL-13 still showed differences in the pathophysiology elicited by these two cytokines (Rankin, et al., 1996, Zhu, et al., 1999). These results suggest that there are real signaling differences between IL-4 and IL-13. To analyze potential signaling differences, we undertook a careful, side-by-side comparison of primary cells and cell lines that expressed either both Type I and II receptors or Type II receptors only. IL-4 stimulated tyrosine phosphorylation of STAT6 in the human airway epithelial cell line, A549, and the human B-cell line, Ramos, at significantly lower doses than IL-13. We demonstrated that IL-4 signaling through the Type I receptor induced robust tyrosine phosphorylation of the downstream adaptor protein IRS-2 and greater expression of the mRNAs for a subset of alternatively activated macrophage genes in primary mouse bone marrow-derived macrophages (BMM) (Heller, et al., 2008). This was in contrast to IL-4/IL-13 signaling through the Type II receptor which resulted in weaker tyrosine phosphorylation of IRS-2 and less mRNA for the AAM genes studied. This marked difference in IRS-2 phosphorylation and AAM gene expression induced by IL-4 was dependent upon expression of the γ C subunit.

4.2 Differential functions of IL-4 and IL-13 in allergic lung inflammation

In both humans and mice, IL-4 and IL-13 signaling through the Type I and Type II receptors play a critical role in inducing asthma. The hallmark features of this disease include excessive pulmonary inflammation, periodic narrowing of airways, airway hyperresponsiveness (AHR) and enhanced mucus secretion. IL-4 and IL-13 have differential roles in asthma pathogenesis. Studies using IL-4R α ^{-/-} and STAT6^{-/-} mice in our lab and by other investigators have suggested that many of the asthma symptoms mentioned above are regulated by IL-4R α and STAT6 (Cohn, et al., 1997, Corry, et al., 1998, Grunig, et al., 1998, Kelly-Welch, et al., 2004, Kuperman, et al., 1998, Mathew, et al., 2001, Wills-Karp, et al., 1998). However, since IL-4/IL-13 binding to either the Type I or Type II receptor activates STAT6, the contributions of these individual pathways in inducing the pathophysiology associated with this disease was unclear, until recently.

It is known that IL-4 is predominantly required for Th2 cell differentiation and proliferation (Kaplan, et al., 1996). Since most naive T cells lack the Type II receptor, they are unresponsive to IL-13. IL-4 signaling through the Type I IL-4R/STAT6 axis upregulates GATA3, the Th2 master transcription factor (reviewed in (Zhu & Paul, 2008)). STAT6-deficient T cells cannot differentiate into Th2 cells *in vitro* (Kaplan, et al., 1996). Recent

studies have shown that STAT6 is not required for *in vivo* differentiation, although it is required for stabilization of Th2 cells and generating memory responses (reviewed in (Chapoval, et al., 2010)). Activated Th2 cells then secrete large quantities of IL-4, IL-5 and IL-13 which can act on many different cell types. IL-4 induces expression of MHC Class II in resting B cells and also causes antibody class switching from IgM to IgE and IgG1 (reviewed in Nelms, et al., 1999). Treatment with anti-IL-4 antibody blocks both primary and secondary IgE responses *in vivo*, when administered at the time of antigenic challenge (Finkelman, et al., 1988). IL-13 on the other hand is thought to be responsible for causing AHR, excessive mucus production and lung fibrosis. Neutralization of IL-13 was able to completely reverse allergen induced airway resistance and abolished mucus production by airway epithelial cells seen in control mice (Grunig, et al., 1998, Wills-Karp, et al., 1998).

Apart from its action on lymphocytes, IL-4 also activates mast cells. This cytokine enhances surface expression of FcεRI, the high affinity receptor for IgE (Toru, et al., 1996). Binding of IgE to FcεRI causes crosslinking of the cytoplasmic Fc domain of this receptor and triggers degranulation (release of mast cell granules). This process causes rapid release of many inflammatory mediators such as histamine, leukotrienes and prostaglandins (reviewed in (Weller, et al., 2011)). Histamine increases blood circulation and permeability of blood vessels, causing increased recruitment of inflammatory cells, including eosinophils, T cells, dendritic cells and monocytes. Leukotrienes and prostaglandins promote bronchoconstriction and stimulate epithelial cell induced mucus production. The importance of FcεRI in allergic responses has been demonstrated in studies using a soluble form of FcεRI and mice lacking the α chain of this receptor (Dombrowicz, et al., 1993, Ra, et al., 1993). In both cases IgE-mediated allergic responses were abrogated.

In addition to mast cells, eosinophils are closely associated with asthma pathogenesis. Increased numbers of eosinophils in the lung and other tissues in asthmatic patients usually correlate with disease severity and it is thought to be the central effector cell involved in airway inflammation (reviewed in (Hogan, et al., 2008)). IL-5 plays an important role in eosinophil development, proliferation and survival in the bone marrow. It is also required for migration of eosinophils into the blood and subsequently the lung. Recruitment of eosinophils to the peribronchial regions of the lung is thought to be mediated by secretion of various eotaxins (eotaxin 1, 2 and 3) by airway epithelial cells. Moreover, IL-5 and the eotaxins cooperate to induce tissue eosinophilia. Various eosinophilic granule components such as major basic protein (MBP) and eosinophilic cationic protein (ECP) have been implicated in initiating and propagating many features of asthma including pulmonary inflammation, airway hyperresponsiveness and bronchoconstriction. Eosinophils express both the Type I and Type II receptors. We have shown that IL-4, but not IL-13, can enhance chemotaxis of eosinophils to eotaxin 1 *in vitro* through the Type I receptor (Heller and Keegan, unpublished). Studies using IL-13Rα1^{-/-} mice have shown that while eotaxin production and secretion by epithelial cells was completely dependent on IL-13 signaling through the Type II receptor, recruitment of eosinophils into the lungs was not (Munitz, et al., 2008, Ramalingam, et al., 2008). Therefore, it is possible that the Type I receptor is compensating for the absence of the Type II receptor. Alternatively, IL-5 may be playing a role in this response. Unlike hematopoietic cells, epithelial, endothelial and smooth muscle cells contain only the Type II receptor. Although both IL-4 and IL-13 can bind to this receptor complex, IL-13 is considered to be the main effector cytokine responsible for AHR, excessive mucus production and lung fibrosis.

The unique contributions of the Type II receptor in allergic lung inflammation have been studied using IL-13R α 1^{-/-} mice. Mucus secretion, airway resistance, eotaxin production and induction of pro-fibrotic mediators such as TGF β were completely dependent on the IL-13R α 1 chain, and thus the Type II receptor (Munitz, et al., 2008, Ramalingam, et al., 2008). However, the authors showed that Th2 cell differentiation, IgE secretion in response to T cell dependent antigens (such as ovalbumin) and recruitment of eosinophils and other inflammatory cells into the lungs could occur independently of IL-13R α 1. In addition, DNA microarray analysis of cells isolated from allergen or IL-4 treated WT or IL-13R α 1^{-/-} mice indicated that several AAM genes were differentially regulated. Munitz *et.al.* showed that allergen and IL-4 induced *Retnla* expression levels were similar in both WT and IL-13R α 1^{-/-} mice, but induction of chitinase (*Chia*) was completely dependent on IL-13R α 1 (Munitz, et al., 2008). Interestingly enough, allergen induced arginase 1 expression required the Type II receptor, but IL-4 induced arginase 1 expression did not. Thus, it appears that IL-4 utilizes both the Type I and Type II receptors to stimulate AAM development in the lung.

Studies conducted by us as well as other groups have shown that IL-4 preferentially induces robust AAM gene expression, while IL-13 does so only weakly (Heller, et al., 2008, Munitz, et al., 2008, Ramalingam, et al., 2008). Intriguingly, mutation of the ITIM motif in the IL-4R α chain resulted in increased sensitivity of macrophages to IL-13 mediated AAM activation. As demonstrated earlier, the Y709 (WT) BMs treated with IL-4 led to significantly higher expression of AAM genes (*Arginase1*, *Chi3l3*) and also *Ccl11* in contrast to IL-13. Mutation of Y709 residue to F709 resulted in dramatic amplification of *Arginase1*, *Chi3l3* and *Ccl11* genes in response to IL-13, while leaving the IL-4 induced responses intact or slightly enhanced (Tachdjian, et al., 2010). IL-13 but not IL-4 induced similar responses in primary lung fibroblasts. These results suggest that there is a disproportionate increase in AAM activation induced by IL-13 signaling via the Type II receptor. The authors hinted that differential recruitment of SHP-1 by the Type I and Type II receptors may be the reason behind these observations (**Figure 2**). Although much progress has been made in understanding the mechanisms by which IL-4 and IL-13 may elicit different responses in different cell types and in the lung during allergic diseases and asthma, many more questions remain unanswered. Future research in this area will shed light on the molecular basis of the separation of functions of IL-4 and IL-13 and their consequences *in vivo*.

4.3 Single nucleotide polymorphisms leading to amino acid changes are commonly found in the IL-4R α

Commonly occurring genetic polymorphisms leading to amino acid changes in the IL-4R α have been linked to susceptibility to asthma and/or to asthma severity (Hershey, et al., 1997, Ober, et al., 2000, Risma, et al., 2002, Shirakawa, et al., 2000). The E400A, Q576R and the S503P polymorphisms reside in the cytoplasmic domain while the I50V resides in the extracellular domain of the IL-4R α chain. Two of the polymorphisms located in the cytoplasmic domain, S503 to P and the Q576 to R, are frequently linked (Kruse, et al., 1999). This double mutation (S503P/Q576R) was associated with lower total IgE concentrations, similar to the single S503P polymorphism, and an increase in the phosphorylation of IRS2. E400A is especially prevalent in African-American populations and was associated with severe asthma exacerbations (Wenzel, et al., 2007). An I⁵⁰ in the extracellular domain of the IL-4R α chain was linked with enhanced signal transduction culminating in an increase in the production of IgE (Mitsuyasu, et al., 1998, Mitsuyasu, et al., 1999). On the other hand, several other studies reported no correlation of this polymorphism with enhanced IgE levels

in patients (Khoo, et al., 2006, Noguchi, et al., 1999). Furthermore, the V⁵⁰ polymorphism has been linked with enhanced CD23 expression, an increase in atopic asthma, and an increase in allergic bronchopulmonary aspergillosis (Knutson, et al., 2006, Risma, et al., 2002). More recently the V⁵⁰ polymorphism reduced the ability of IL-4 to suppress IL-17 production by human peripheral mononuclear cells (Wallis, et al., 2011). Because many of these polymorphisms are located in the cytoplasmic domain of the IL-4R α , many investigators have hypothesized that they modulate signal transduction. However, experiments designed to analyze the direct effects of these polymorphisms on receptor signaling have lead to contradictory reports (Franjkovic, et al., 2005, Kruse, et al., 1999, Mitsuyasu, et al., 1998, Mitsuyasu, et al., 1999, Prots, et al., 2006, Risma, et al., 2002, Stephenson, et al., 2004). We analyzed the impact of the Q576R, S503P, and I50V polymorphisms on signal transduction by the Type I receptor complex *in vitro*. While the R⁵⁷⁶ and P⁵⁰³ polymorphisms had no effect on STAT6 or IRS2 activation induced by IL-4 (Wang, et al., 1999), we found that the V⁵⁰ polymorphism located in the extracellular domain of the IL-4R α mediated a prolonged STAT6 signaling induced by IL-4 through the Type I receptor (Ford, et al., 2009). This was associated with a prolonged expression of the SOCS family member *Cis*. The effect of this polymorphism on signaling by the Type II receptor is unknown.

Using mouse knock-in strategies, the murine IL-4R α -Q⁵⁷⁶ was replaced with IL-4R α expressing the R⁵⁷⁶ polymorphism (Tachdjian, et al., 2009). This change was shown to enhance allergic asthma *in vivo*. The IL-4R α -R⁵⁷⁶ enhanced Th2 differentiation and IgE production; both of these responses are Type I receptor dependent in the mouse model. Furthermore, it enhanced the production of CCL11 by BMM, fibroblasts, and tracheal epithelial cells (TEC) in response to IL-4 or IL-13. These results suggest the R⁵⁷⁶ polymorphism affects STAT6-dependent responses down stream of both the Type I and Type II receptors. However, there was no apparent effect of the R⁵⁷⁶ on the tyrosine phosphorylation of STAT6 (Tachdjian, et al., 2009), consistent with our study in cell lines (Wang, et al., 1999). Furthermore, there was no effect on the tyrosine phosphorylation of Shc. The effect of the R⁵⁷⁶ on IRS2 phosphorylation was not reported. Interestingly, the longevity of Erk1,2 phosphorylation in TEC was dramatically enhanced by R⁵⁷⁶, however the mechanism by which R⁵⁷⁶ leads to Erk phosphorylation is still unclear. It will be important to understand the effects of the polymorphisms and their interactions on both IL-4 and IL-13 signaling since they are linked to different clinical phenotypes in human populations (Wenzel, et al., 2007).

5. Contribution of Type I and Type II receptor complexes to the control of regulatory mechanisms that act to modulate allergic inflammation

During Th2-driven allergic lung inflammation, a number of effector and regulatory mechanisms are orchestrated by IL-4 and IL-13 through Type I and Type II receptors. Several of these regulatory mechanisms allow for the amplification of Th2 differentiation and function, while others function as part of a negative feed-back loop to limit Th2-driven inflammation (**Figure 3**). Just as IL-4 and IL-13 are differentially involved in promoting various features of allergic lung disease, these cytokines utilize separate mechanisms to negatively regulate the signaling pathways activated by each other. These regulatory mechanisms will need to be considered in the design of inhibitors of the IL-4/IL-13 system.




Fig. 3. **Differential control of regulatory mechanisms by the Type I and II Receptors.** IL-4 signaling through the Type I receptor on Th2 cell surfaces renders them resistant to control by T regulatory cells. Furthermore, IL-4 signaling STAT6 activation through the Type I receptor on naive CD4+ T-cells inhibits the differentiation of iTregs and Th1 cells. These responses lead to enhanced allergic inflammation. On the other hand, IL-4 signals through the Type I receptor on NK cell surfaces to induce IFN γ production. IFN γ in turn suppresses Th2 differentiation and inhibits signaling by both Type I and Type II receptors by inducing SOCS family members. IL-13 signals through the Type II receptor that is induced on the surface of Th17 cells and decreases IL-17 production. This suppression of IL-17 production by Th17 cells would limit the influx of neutrophils that are present in steroid resistant asthma.

5.1 Type I IL-4 receptor modulation of regulatory T-cells

Signaling through the Type I IL-4 receptor antagonizes the differentiation and function of regulatory T cells (Tregs). Tregs are a subset of T lymphocytes that regulate immune responses and prevent excessive immune system activation (Sakaguchi, et al., 2008, Shevach, 2009). The most studied Tregs are CD4⁺CD25⁺FoxP3⁺, which have been found to play a role in allergic disease. FoxP3 is a transcription factor important for Treg development and function. Mice lacking FoxP3 expression (Scurfy mice) or humans with mutations in their *foxP3* gene (X-linked autoimmunity-allergic dysregulation syndrome) develop widespread autoimmune disease with a Th2-mediated allergic component (Khatttri, et al., 2003). There are many mechanisms elicited by Tregs to suppress effector T cells, such as: (1) cell-mediated presentation of TGF-β or galectin-1 (Shevach, 2009), (2) secretion of the immunosuppressive cytokines IL-10 or TGF-β (Bettini & Vignali, 2009), or (3) consumption of IL-2, a limiting growth factor for T cells (Scheffold, et al., 2007). Th2 cells utilize several

methods to inhibit immunosuppression by Tregs. Gata3 is the master transcriptional regulator of Th2 cells and is activated by signal transduction through the Type I IL-4 receptor. Using co-immunoprecipitation assays, Dardalhon and colleagues have shown physical interaction between Gata3 and FoxP3 in transiently transfected human embryo kidney 293 cells. The authors also observed IL-4 inhibition of FoxP3 induction in Ag-specific adaptive (inducible) Tregs, which was dependent on STAT6 expression (Dardalhon, et al., 2008). Our recent findings are in support of this antagonistic relationship between STAT6 and Tregs and IL-4-induced, STAT6-dependent inhibition of FoxP3. We have found that STAT6^{-/-} mice have higher numbers of Tregs than wildtype mice (Chapoval, et al., 2011). This observation has also been confirmed by Takaki and colleagues who identified a STAT6-binding site in the silencer region in the FoxP3 mRNA transcript. STAT6 binding to this site reduced TGF- β 1 induction of FoxP3 transcriptional activation (Takaki, et al., 2008).

Additionally, IL-4 was found to serve as a survival factor for CD4⁺CD25⁻ T helper (Th) cells and to aid in their protection from immunosuppression by CD4⁺CD25⁺ Tregs (Pace, et al., 2005). IL-13 failed to have this same effect. Therefore, this phenomenon was found to be dependent on Th cell surface expression of IL-4R α chain and a functional Type I IL-4 receptor complex, as IL-4R α ^{-/-} Th cells were not protected by IL-4 from Treg immunosuppression. IL-4 activation of the Type I IL-4 receptor, but not the Type II receptor, maintains anti-apoptotic and pro-proliferative processes in Th cells and protects them from Treg-induced perturbed cell growth and proliferation. Surprisingly, CFSE-labeled cocultures of IL-4R α ^{-/-} Th cells and IL-4R α ^{+/+} Tregs revealed that although IL-4 has an effect on Treg immunosuppression of IL-4R α ^{-/-} Th cells, it promoted proliferation of Tregs *in-vitro* (Pace, et al., 2006). This could be a direct effect of IL-4 that would further complicate the role of this Th2 cytokine in modulating Treg immunosuppression.

5.2 Role of Type I receptors in control of Th1 responses that act to suppress allergic inflammation

Th1 cell programming antagonizes Th2 cell differentiation and could serve as a regulatory mechanism to suppress Th2-mediated allergic response. Gavett and colleagues showed that IL-12 can inhibit antigen induced-airway hyperreactivity and inflammation and to also reduce Th2 cytokine production (Gavett, et al., 1995). Th2 cells inhibit Th1 cell induction during the allergic response. Gata3 is activated downstream of STAT6 phosphorylation, which is induced by signaling through the Type I receptor (Kurata, et al., 1999, Nelms, et al., 1999). Not only does it serve as the master regulator transcription factor (TF) for Th2 cells, but Gata3 has also been shown to inhibit Th1 cell-specific factors. Gata3-deficient cell clones produced high levels of the Th1 cytokine, IFN γ , and had enhanced expression of T-bet (the Th1 cell master regulator TF) (Zhu, et al., 2006). Therefore, activation of the Type I IL-4 receptor could lead to activation of Gata3, which in turn inhibits Th1-inducing factors during the Th2-mediated allergic response. Th1 cells have also been shown to play a potentially stimulatory role in airway inflammation. Hansen et al. found that Th1 cells decreased airway eosinophilia, but failed to reduce airway hyperreactivity in ovalbumin-immunized Balb/c mice (Hansen, et al., 1999). But this may be a late phenomenon in airway inflammation during which additional non-Th2 inflammatory cells aid in amplifying chronic lung inflammation. This would be in direct contrast to the initiation of allergic airway inflammation which is characterized by strong Th2 immune responses and can be inhibited by non-Th2-promoting immune cells.

5.3 Modulation of NK cells

Although IL-4 and IFN γ are antagonistic towards each other during T cell differentiation, IL-4 can increase IL-2 and IL-12 induced IFN γ secretion by Natural Killer (NK) cells. Bream and co-authors also found that the increase in IFN γ production caused by IL-4 in conjunction with IL-2 was STAT6 dependent, while IL-4 synergy with IL-12 was independent of STAT6 activation (Bream, et al., 2003, Morris, et al., 2006). Further studies have shown that IL-13 was unable to cause a similar increase in NK cell derived IFN γ release (Morris, et al., 2006). This result is in agreement with the fact that IL-13 signals through the Type II receptor, and this receptor complex is absent in NK cells. IL-4 stimulated IFN γ production by NK cells has significant implications in the context for allergic lung inflammation. One group has shown that Sendai virus infection of mice suppressed NK cell derived IFN γ secretion. This led to enhanced Th2 responses and subsequent development of exacerbated allergic lung disease (Kaiko, et al., 2010). IFN γ production would suppress Th2 differentiation and inhibit signaling by both Type I and Type II receptors by inducing SOCS family members.

The role of NK cells in human allergic disease has been extensively examined; because of their ability to produce cytokines, NK cells have the potential to heavily influence the adaptive allergic immune response. Based on cytokine production, NK cells can be divided into 2 classes: (1) NK1 cells produce Th1 cytokines, such as IFN γ (Romagnani, 1992) and (2) NK2 cells produce Th2 cytokines IL-4, 5, and 13 (Hoshino, et al., 1999, Peritt, et al., 1998, Warren, et al., 1995). In a recent study analyzing NK cell populations in healthy and allergic patients, the authors found a predominance of NK2 cells in the peripheral blood of allergic patients. These NK2 cells produced high amounts of Th2 cytokines that could promote allergic inflammation (Timonen & Stenius-Aarniala, 1985, Wei, et al., 2005).

5.4 IL-13 inhibition of Th17 cells via the Type II receptor in severe asthma

It has long been thought that naïve T cells do not express the IL-13R α 1 chain of the Type II receptor complex and therefore cannot be regulated by IL-13. Due to restricted expression of IL-13R α 1 on non-hematopoietic cells, Type II receptor signaling has been limited to those cells and not seen in T cells. But this central dogma has been recently challenged by observations that Th17 cells are able to induce surface expression of IL-13R α 1 chain (Newcomb, et al., 2009, Newcomb, et al., 2011).

Th17 cells are a distinct population of CD4⁺ T cells, whose differentiation is induced by IL-6 or IL-21 and TGF β (McGeachy, et al., 2007). Th17 cells produce IL-17, IL-6, and tumor necrosis factor. They have been shown to play a role in autoimmune diseases including the experimental autoimmune encephalitis (EAE) model of multiple sclerosis (Langrish, et al., 2005). Th17 cells also provide protection from some extracellular pathogens, such as *Klebsiella pneumoniae* infection of the lung (Happel, et al., 2005).

A statistically higher number of IL-17⁺ cells can be found in the sputum and BAL of asthmatic patients compared to controls and the greater expression of IL-17A in the lungs was associated with increased asthma severity. (Jatakanon, et al., 1999, Molet, et al., 2001). IL-17A induces neutrophil recruitment to the airway and augments the pathogenesis of steroid-resistant, severe asthma (McKinley, et al., 2008). Th17 cells alone cannot induce eosinophilic infiltration into the airway following immunization and challenge, but in the presence of Th2 cells, antigen-specific Th17 cells can enhance the eosinophil-activating properties of Th2 cells (Wakashin, et al., 2008).

Th17 polarized cells from mouse spleens were shown to have increased mRNA and protein levels of IL-13R α 1 after stimulation *in vitro*. When added to Th17 cell cultures, IL-13 reduced IL-17A production by Th17 cells and decreased the percentage of CD4⁺ Th17 cells. Additionally, IL-13 caused a reduction in the expression of ROR γ t, the master regulator transcription factor for Th17 cells (Newcomb, et al., 2009). This phenomenon of IL-13 suppression of IL-17A production by Th17 cells was also observed *in vitro* using human CD4⁺ T cells (Newcomb, et al., 2011). Therefore, activated Th17 cells upregulate their surface expression of IL-13R α 1 chain and this allows for IL-13 to signal through a functional Type II receptor complex to decrease IL-17 production by Th17 cells. Thus, paradoxically, IL-13, a major effector cytokine of atopic asthma, inhibits the Th17 component of severe asthma. This concept could also explain the observation that IL-25-induced production of IL-13 inhibited Th17-mediated EAE disease progression (Kleinschek, et al., 2007). This is a unique mechanism whereby a Th2 immune-mediated illness is prevented from becoming more severe by IL-13, a Th2 cytokine. Consequently, efforts to suppress IL-13 function to treat allergic asthma may lead to Th17 induction and severe and persistent asthma in susceptible individuals.

6. Conclusion

The importance of the Type I IL-4 receptor in regulating T cells to become Th2 cells has been well documented. Furthermore, numerous studies have indicated the IL-4R α expressed on lung epithelium is necessary for goblet cell differentiation and mucus hypersecretion. In addition, the IL-4R α is expressed on many cell types that could contribute to the overall pathology and severity of asthma. The relative role of the Type I and Type II receptors on these cells has not yet been fully delineated. Using mice lacking one or the other complex (i.e. γ C^{-/-} or IL-13R α 1^{-/-}) several groups have recently delineated interesting differences in their contributions to lung pathology. The Type I receptor is the major regulator of eosinophilic inflammation and the alternative activation of macrophages while the Type II receptor controls mucus hypersecretion and airway hyperresponsiveness. These receptors differentially regulate potential regulatory pathways including the control of T regulatory cells and the production of cytokines by NK cells; both of these responses are controlled by the Type I receptor complex. The Type II receptor complex can be induced on Th17 cells and allows IL-13 to down regulate Th17 differentiation. This could be of clinical importance for severe, steroid resistant forms of human asthma that may be mediated by Th17 cells. In this scenario, inhibiting IL-13 could be detrimental to the patient and illustrates the need to stratify patients prior to treatment.

As mentioned above, the intricate differences between the Type I and Type II receptor complexes could impact the therapeutic effectiveness of agents designed to target these receptors in asthma. Indeed initial trials to inhibit IL-4 using the soluble IL-4R α in asthmatic patients were largely unsuccessful and possibly detrimental (Borish, et al., 2001, Wenzel, et al., 2007). The ability of IL-4 to suppress TNF α production, a highly pro-inflammatory cytokine, was suggested to be part of a negative regulatory mechanism that was unintentionally blocked by the therapy (Borish, 2010). Studies using soluble IL-13R α 2 to inhibit IL-13 signaling are ongoing. However, blocking one cytokine at a time may not prove beneficial. Since both cytokines can elicit effector functions, blocking only one could actually exacerbate disease because of the loss of a negative regulatory pathway. The most promising approach thus far has been to use a mutant IL-4 (Pitrakinra) that binds to the IL-4R α , and blocks dimerization with either the γ C or the IL-13R α 1 (Wenzel, et al., 2007). Thus, this single agent can prevent the formation of both the Type I and the Type II receptor

complexes. Further understanding of the complex IL-4/IL-13 receptor system and its contribution to various features of allergic asthma will be essential to fine-tune therapeutic strategies for the treatment of asthma.

7. Acknowledgement

The authors acknowledge their financial support from the National Institutes of Health: RO1 AI038985 (ADK); T32HL06798 and K99/R00 HL096897 (NMH); Meyerhoff Graduate Research Fellowship, Minority Biomedical Research Support – Initiative for Maximizing Student Diversity, NIGMS, T32HL06798 and AI007540 (NJD); and R21AI076736 (SPC).

8. References

- Akbari, O., Stock, P., Meyer, E., Kronenberg, M., Sidobre, S., Nakayama, T., Taniguchi, M., Grusby, M. J., DeKruyff, R. H., & Umetsu, D. T. (2003). Essential role of NKT cells producing IL-4 and IL-13 in the development of allergen-induced airway hyperreactivity. *Nature Medicine*, Vol. 9, No. 5, (May, 2003), pp. 582-588.
- Aman, M. J., Tayebi, N., Obiri, N. I., Puri, R. K., Modi, W. S., & Leonard, W. J. (1996). cDNA cloning and characterization of the human interleukin 13 receptor alpha chain. *Journal of Biological Chemistry*, Vol. 271, No. 46, (Nov 15, 1996), pp. 29265-29270.
- Andrews, A. L., Holloway, J. W., Holgate, S. T., & Davies, D. E. (2006). IL-4 receptor alpha is an important modulator of IL-4 and IL-13 receptor binding: implications for the development of therapeutic targets. *Journal of Immunology*, Vol. 176, No. 12, (Jun 15, 2006), pp. 7456-7461.
- Andrews, A. L., Holloway, J. W., Puddicombe, S. M., Holgate, S. T., & Davies, D. E. (2002). Kinetic analysis of the interleukin-13 receptor complex. *Journal of Biological Chemistry*, Vol. 277, No. 48, (Nov 29, 2002), pp. 46073-46078.
- Andrews, A. L., Nasir, T., Bucchieri, F., Holloway, J. W., Holgate, S. T., & Davies, D. E. (2006). IL-13 receptor alpha 2: a regulator of IL-13 and IL-4 signal transduction in primary human fibroblasts. *Journal of Allergy and Clinical Immunology*, Vol. 118, No. 4, (Oct, 2006), pp. 858-865.
- Andrews, R., Rosa, L., Daines, M., & Khurana Hershey, G. (2001). Reconstitution of a functional human type II IL-4/IL-13 receptor in mouse B cells: demonstration of species specificity. *Journal of Immunology*, Vol. 166, No. 3, (Feb 1, 2001), pp. 1716-1722.
- Bahadori, K., Doyle-Waters, M. M., Marra, C., Lynd, L., Alasaly, K., Swiston, J., & FitzGerald, J. M. (2009). Economic burden of asthma: a systematic review. *BMC Pulm Med*, Vol. 9, No., (2009), pp. 24, 1471-2466.
- Ben-Sasson, S. Z., Le Gros, G., Conrad, D. H., Finkelman, F. D., & Paul, W. E. (1990). Cross-linking Fc receptors stimulate splenic non-B, non-T cells to secrete interleukin 4 and other lymphokines. *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 87, No. 4, (Feb, 1990), pp. 1421-1425.
- Bettini, M., & Vignali, D. A. (2009). Regulatory T cells and inhibitory cytokines in autoimmunity. *Current Opinion in Immunology*, Vol. 21, No. 6, (Dec, 2009), pp. 612-618.
- Blaeser, F., Bryce, P. J., Ho, N., Raman, V., Dedeoglu, F., Donaldson, D. D., Geha, R. S., Oettgen, H. C., & Chatila, T. A. (2003). Targeted inactivation of the IL-4 receptor alpha chain I4R motif promotes allergic airway inflammation. *Journal of Experimental Medicine*, Vol. 198, No. 8, (Oct 20, 2003), pp. 1189-1200.

- Borish, L. (2010). IL-4 and IL-13 dual antagonism: a promising approach to the dilemma of generating effective asthma biotherapeutics. *American Journal of Respiratory and Critical Care Medicine*, Vol. 181, No. 8, (Apr 15, 2010), pp. 769-770.
- Borish, L. C., Nelson, H. S., Corren, J., Bensch, G., Busse, W. W., Whitmore, J. B., & Agosti, J. M. (2001). Efficacy of soluble IL-4 receptor for the treatment of adults with asthma. *Journal of Allergy and Clinical Immunology*, Vol. 107, No. 6, (Jun, 2001), pp. 963-970.
- Bream, J. H., Curiel, R. E., Yu, C. R., Egwuagu, C. E., Grusby, M. J., Aune, T. M., & Young, H. A. (2003). IL-4 synergistically enhances both IL-2- and IL-12-induced IFN-gamma expression in murine NK cells. *Blood*, Vol. 102, No. 1, (Jul 1, 2003), pp. 207-214.
- Brown, M. A., Pierce, J. H., Watson, C. J., Falco, J., Ihle, J. N., & Paul, W. E. (1987). B cell stimulatory factor-1/interleukin-4 mRNA is expressed by normal and transformed mast cells. *Cell*, Vol. 50, No. 5, (Aug 28, 1987), pp. 809-818.
- Bruns, H. A., Schindler, U., & Kaplan, M. H. (2003). Expression of a constitutively active Stat6 in vivo alters lymphocyte homeostasis with distinct effects in T and B cells. *Journal of Immunology*, Vol. 170, No. 7, (Apr 1, 2003), pp. 3478-3487.
- Burd, P. R., Thompson, W. C., Max, E. E., & Mills, F. C. (1995). Activated mast cells produce interleukin 13. *Journal of Experimental Medicine*, Vol. 181, No. 4, (Apr 1, 1995), pp. 1373-1380.
- Cao, X., Kozak, C. A., Liu, Y. J., Noguchi, M., O'Connell, E., & Leonard, W. J. (1993). Characterization of cDNAs encoding the murine interleukin 2 receptor (IL-2R) gamma chain: chromosomal mapping and tissue specificity of IL-2R gamma chain expression. *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 90, No. 18, (Sep 15, 1993), pp. 8464-8468.
- Cao, X., Shores, E. W., Hu-Li, J., Anver, M. R., Kelsall, B. L., Russell, S. M., Drago, J., Noguchi, M., Grinberg, A., Bloom, E. T., & et al. (1995). Defective lymphoid development in mice lacking expression of the common cytokine receptor gamma chain. *Immunity*, Vol. 2, No. 3, (Mar, 1995), pp. 223-238.
- Caput, D., Laurent, P., Kaghad, M., Lelias, J. M., Lefort, S., Vita, N., & Ferrara, P. (1996). Cloning and characterization of a specific interleukin (IL)-13 binding protein structurally related to the IL-5 receptor alpha chain. *Journal of Biological Chemistry*, Vol. 271, No. 28, (Jul 12, 1996), pp. 16921-16926.
- Chapoval, S., Dasgupta, P., Dorsey, N. J., & Keegan, A. D. (2010). Regulation of the T helper cell type 2 (Th2)/T regulatory cell (Treg) balance by IL-4 and STAT6. *Journal of Leukocyte Biology*, Vol. 87, No. 6, (Jun, 2010), pp. 1011-1018.
- Chapoval, S. P., Dasgupta, P., Smith, E. P., DeTolla, L. J., Lipsky, M. M., Kelly-Welch, A. E., & Keegan, A. D. (2011). STAT6 expression in multiple cell types mediates the cooperative development of allergic airway disease. *Journal of Immunology*, Vol. 186, No. 4, (Feb 15, 2011), pp. 2571-2583.
- Chen, W., Daines, M. O., & Khurana Hershey, G. K. (2003). Arginine 27, a conserved methylation site in Stat6, is essential for IL-4-induced tyrosine phosphorylation, DNA binding activity, and nuclear translocation of Stat6. *Journal of Allergy and Clinical Immunology*, Vol. 111, No. 2, (2003), pp. S855.
- Chen, W., Sivaprasad, U., Tabata, Y., Gibson, A. M., Stier, M. T., Finkelman, F. D., & Hershey, G. K. (2009). IL-13R alpha 2 membrane and soluble isoforms differ in humans and mice. *Journal of Immunology*, Vol. 183, No. 12, (Dec 15, 2009), pp. 7870-7876.

- Chiaramonte, M. G., Mentink-Kane, M., Jacobson, B. A., Cheever, A. W., Whitters, M. J., Goad, M. E., Wong, A., Collins, M., Donaldson, D. D., Grusby, M. J., & Wynn, T. A. (2003). Regulation and function of the interleukin 13 receptor alpha 2 during a T helper cell type 2-dominant immune response. *Journal of Experimental Medicine*, Vol. 197, No. 6, (Mar 17, 2003), pp. 687-701.
- Chomarat, P., & Banchereau, J. (1998). Interleukin-4 and interleukin-13: their similarities and discrepancies. *International Reviews of Immunology*, Vol. 17, No. 1-4, (1998), pp. 1-52, 0883-0185.
- Coffer, P. J., Schweizer, R. C., Dubois, G. R., Maikoe, T., Lammers, J. W., & Koenderman, L. (1998). Analysis of signal transduction pathways in human eosinophils activated by chemoattractants and the T-helper 2-derived cytokines interleukin-4 and interleukin-5. *Blood*, Vol. 91, No. 7, (Apr 1, 1998), pp. 2547-2557.
- Cohn, L., Homer, R. J., Marinov, A., Rankin, J., & Bottomly, K. (1997). Induction of airway mucus production By T helper 2 (Th2) cells: a critical role for interleukin 4 in cell recruitment but not mucus production. *Journal of Experimental Medicine*, Vol. 186, No. 10, (Nov 17, 1997), pp. 1737-1747.
- Corry, D. B., Grunig, G., Hadeiba, H., Kurup, V. P., Warnock, M. L., Sheppard, D., Rennick, D. M., & Locksley, R. M. (1998). Requirements for allergen-induced airway hyperreactivity in T and B cell-deficient mice. *Molecular Medicine*, Vol. 4, No. 5, (May, 1998), pp. 344-355.
- Crowley, M. T., Harmer, S. L., & DeFranco, A. L. (1996). Activation-induced association of a 145-kDa tyrosine-phosphorylated protein with Shc and Syk in B lymphocytes and macrophages. *Journal of Biological Chemistry*, Vol. 271, No. 2, (Jan 12, 1996), pp. 1145-115.
- Dardalhon, V., Awasthi, A., Kwon, H., Galileos, G., Gao, W., Sobel, R. A., Mitsdoerffer, M., Strom, T. B., Elyaman, W., Ho, I. C., Khoury, S., Oukka, M., & Kuchroo, V. K. (2008). IL-4 inhibits TGF-beta-induced Foxp3⁺ T cells and, together with TGF-beta, generates IL-9⁺ IL-10⁺ Foxp3(-) effector T cells. *Nature Immunology*, Vol. 9, No. 12, (Dec, 2008), pp. 1347-1355.
- Das, J., Eynott, P., Jupp, R., Bothwell, A., Van Kaer, L., Shi, Y., & Das, G. (2006). Natural killer T cells and CD8⁺ T cells are dispensable for T cell-dependent allergic airway inflammation. *Nature Medicine*, Vol. 12, No. 12, (Dec, 2006), pp. 1345-1346; author reply 1347.
- David, M. D., Bertoglio, J., & Pierre, J. (2003). Functional characterization of IL-13 receptor alpha2 gene promoter: a critical role of the transcription factor STAT6 for regulated expression. *Oncogene*, Vol. 22, No. 22, (2003), pp. 3386-3394.
- Deutsch, H. H., Koettnitz, K., Chung, J., & Kalthoff, F. S. (1995). Distinct sequence motifs within the cytoplasmic domain of the human IL-4 receptor differentially regulate apoptosis inhibition and cell growth. *Journal of Immunology*, Vol. 154, No. 8, (Apr 15, 1995), pp. 3696-3703.
- Devouassoux, G., Saxon, A., Metcalfe, D. D., Prussin, C., Colomb, M. G., Brambilla, C., & Diaz-Sanchez, D. (2002). Chemical constituents of diesel exhaust particles induce IL-4 production and histamine release by human basophils. *Journal of Allergy and Clinical Immunology*, Vol. 109, No. 5, (May, 2002), pp. 847-853.
- Dombrowicz, D., Flamand, V., Brigman, K. K., Koller, B. H., & Kinet, J. P. (1993). Abolition of anaphylaxis by targeted disruption of the high affinity immunoglobulin E receptor alpha chain gene. *Cell*, Vol. 75, No. 5, (Dec 3, 1993), pp. 969-976.

- Donaldson, D. D., Whitters, M. J., Fitz, L. J., Neben, T. Y., Finnerty, H., Henderson, S. L., O'Hara, R. M., Jr., Beier, D. R., Turner, K. J., Wood, C. R., & Collins, M. (1998). The murine IL-13 receptor alpha 2: molecular cloning, characterization, and comparison with murine IL-13 receptor alpha 1. *Journal of Immunology*, Vol. 161, No. 5, (Sep 1, 1998), pp. 2317-2324.
- Duronio, V., Welham, M. J., Abraham, S., Dryden, P., & Schrader, J. W. (1992). p21ras activation via hemopoietin receptors and c-kit requires tyrosine kinase activity but not tyrosine phosphorylation of p21ras GTPase-activating protein. *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 89, No. 5, (Mar 1, 1992), pp. 1587-1591.
- Ferrick, D. A., Schrenzel, M. D., Mulvania, T., Hsieh, B., Ferlin, W. G., & Lepper, H. (1995). Differential production of interferon-gamma and interleukin-4 in response to Th1- and Th2-stimulating pathogens by gamma delta T cells in vivo. *Nature*, Vol. 373, No. 6511, (Jan 19, 1995), pp. 255-257.
- Fichtner-Feigl, S., Strober, W., Kawakami, K., Puri, R. K., & Kitani, A. (2006). IL-13 signaling through the IL-13alpha2 receptor is involved in induction of TGF-beta1 production and fibrosis. *Nature Medicine*, Vol. 12, No. 1, (Jan, 2006), pp. 99-106.
- Finkelman, F. D., Katona, I. M., Urban, J. F., Jr., Holmes, J., Ohara, J., Tung, A. S., Sample, J. V., & Paul, W. E. (1988). IL-4 is required to generate and sustain in vivo IgE responses. *Journal of Immunology*, Vol. 141, No. 7, (Oct 1, 1988), pp. 2335-2341.
- Ford, A. Q., Heller, N. M., Stephenson, L., Boothby, M. R., & Keegan, A. D. (2009). An Atopy-Associated Polymorphism in the Ectodomain of the IL-4R{alpha} Chain (V50) Regulates the Persistence of STAT6 Phosphorylation. *Journal of Immunology*, Vol. 183, No. 3, (August 1, 2009, 2009), pp. 1607-1616.
- Franjkovic, I., Gessner, A., Konig, I., Kissel, K., Bohnert, A., Hartung, A., Ohly, A., Ziegler, A., Hackstein, H., & Bein, G. (2005). Effects of common atopy-associated amino acid substitutions in the IL-4 receptor alpha chain on IL-4 induced phenotypes. *Immunogenetics*, Vol. 56, No. 11, (Feb, 2005), pp. 808-817.
- Fujiwara, H., Hanissian, S. H., Tsytsykova, A., & Geha, R. S. (1997). Homodimerization of the human interleukin 4 receptor alpha chain induces Cepsilon germline transcripts in B cells in the absence of the interleukin 2 receptor gamma chain. *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 94, No. 11, (May 27, 1997), pp. 5866-5871.
- Gadina, M., Stancato, L. M., Bacon, C. M., Lerner, A. C., & O'Shea, J. J. (1998). Involvement of SHP-2 in multiple aspects of IL-2 signaling: evidence for a positive regulatory role. *Journal of Immunology*, Vol. 160, No. 10, (May 15, 1998), pp. 4657-4661.
- Gadina, M., Sudarshan, C., & O'Shea, J. J. (1999). IL-2, but not IL-4 and other cytokines, induces phosphorylation of a 98-kDa protein associated with SHP-2, phosphatidylinositol 3'-kinase, and Grb2. *Journal of Immunology*, Vol. 162, No. 4, (Feb 15, 1999), pp. 2081-2086.
- Galizzi, J. P., Zuber, C. E., Harada, N., Gorman, D. M., Djossou, O., Kastelein, R., Banchereau, J., Howard, M., & Miyajima, A. (1990). Molecular cloning of a cDNA encoding the human interleukin 4 receptor. *International Immunology*, Vol. 2, No. 7, (1990), pp. 669-675.
- Gavett, S. H., Chen, X., Finkelman, F., & Wills-Karp, M. (1994). Depletion of murine CD4+ T lymphocytes prevents antigen-induced airway hyperreactivity and pulmonary

- eosinophilia. *American Journal of Respiratory Cell and Molecular Biology*, Vol. 10, No. 6, (Jun, 1994), pp. 587-593.
- Gavett, S. H., O'Hearn, D. J., Li, X., Huang, S. K., Finkelman, F. D., & Wills-Karp, M. (1995). Interleukin 12 inhibits antigen-induced airway hyperresponsiveness, inflammation, and Th2 cytokine expression in mice. *The Journal of experimental medicine*, Vol. 182, No. 5, (Nov 1, 1995), pp. 1527-1536
- Goenka, S., & Kaplan, M. H. (2011). Transcriptional regulation by STAT6. *Immunologic Research*, Vol. 50, No. 1, (May, 2011), pp. 87-96.
- Goenka, S., Youn, J., Dzurek, L. M., Schindler, U., Yu-Lee, L. Y., & Boothby, M. (1999). Paired Stat6 C-terminal transcription activation domains required both for inhibition of an IFN-responsive promoter and trans-activation. *Journal of Immunology*, Vol. 163, No. 9, (Nov 1, 1999), pp. 4663-4672.
- Grunig, G., Warnock, M., Wakil, A. E., Venkayya, R., Brombacher, F., Rennick, D. M., Sheppard, D., Mohrs, M., Donaldson, D. D., Locksley, R. M., & Corry, D. B. (1998). Requirement for IL-13 independently of IL-4 in experimental asthma. *Science*, Vol. 282, No. 5397, (Dec 18, 1998), pp. 2261-2263.
- Haan, C., Rolvering, C., Raulf, F., Kapp, M., Druckes, P., Thoma, G., Behrmann, I., & Zerwes, H. G. (2011). Jak1 has a dominant role over Jak3 in signal transduction through gammac-containing cytokine receptors. *Chemistry and Biology*, Vol. 18, No. 3, (Mar 25, 2011), pp. 314-323.
- Hage, T., Sebald, W., & Reinemer, P. (1999). Crystal structure of the IL-4/IL-4 receptor alpha chain complex reveals a mosaic binding interface. *Cell*, Vol. 97, No. 2, (April 16, 1999), pp 271-281.
- Hansen, G., Berry, G., DeKruyff, R. H., & Umetsu, D. T. (1999). Allergen-specific Th1 cells fail to counterbalance Th2 cell-induced airway hyperreactivity but cause severe airway inflammation. *The Journal of Clinical Investigation*, Vol. 103, No. 2, (Jan, 1999), pp. 175-183.
- Hanson, E. M., Dickensheets, H., Qu, C. K., Donnelly, R. P., & Keegan, A. D. (2003). Regulation of the dephosphorylation of Stat6. Participation of Tyr-713 in the interleukin-4 receptor alpha, the tyrosine phosphatase SHP-1, and the proteasome. *Journal of Biological Chemistry*, Vol. 278, No. 6, (2003), pp. 3903-3911.
- Happel, K. I., Dubin, P. J., Zheng, M., Ghilardi, N., Lockhart, C., Quinton, L. J., Odden, A. R., Shellito, J. E., Bagby, G. J., Nelson, S., & Kolls, J. K. (2005). Divergent roles of IL-23 and IL-12 in host defense against *Klebsiella pneumoniae*. *The Journal of experimental medicine*, Vol. 202, No. 6, (Sep 19, 2005), pp. 761-769.
- Haque, S. J., Harbor, P., Tabrizi, M., Yi, T., & Williams, B. R. (1998). Protein-tyrosine phosphatase Shp-1 is a negative regulator of IL-4- and IL-13-dependent signal transduction. *Journal of Biological Chemistry*, Vol. 273, No. 51, (1998), pp. 33893-33896.
- Haque, S. J., Harbor, P. C., & Williams, B. R. (2000). Identification of critical residues required for suppressor of cytokine signaling-specific regulation of interleukin-4 signaling. *Journal of Biological Chemistry*, Vol. 275, No. 34, (2000), pp. 26500-26506.
- Heller, N. M., Qi, X., Junttila, I. S., Shirey, K. A., Vogel, S. N., Paul, W. E., & Keegan, A. D. (2008). Type I IL-4Rs selectively activate IRS-2 to induce target gene expression in macrophages. *Sci Signal*, Vol. 1, No. 51, (2008), pp. ra17.
- Hershey, G. K., Friedrich, M. F., Esswein, L. A., Thomas, M. L., & Chatila, T. A. (1997). The association of atopy with a gain-of-function mutation in the alpha subunit of the

- interleukin-4 receptor. *New England Journal of Medicine*, Vol. 337, No. 24, (Dec 11, 1997), pp. 1720-1725.
- Hilton, D. J., Zhang, J. G., Metcalf, D., Alexander, W. S., Nicola, N. A., & Willson, T. A. (1996). Cloning and characterization of a binding subunit of the interleukin 13 receptor that is also a component of the interleukin 4 receptor. *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 93, No. 1, (Jan 9, 1996), pp. 497-501.
- Hogan, S. P., Rosenberg, H. F., Moqbel, R., Phipps, S., Foster, P. S., Lacy, P., Kay, A. B., & Rothenberg, M. E. (2008). Eosinophils: biological properties and role in health and disease. *Clinical and Experimental Allergy*, Vol. 38, No. 5, (May, 2008), pp. 709-750.
- Hoshino, T., Winkler-Pickett, R. T., Mason, A. T., Ortaldo, J. R., & Young, H. A. (1999). IL-13 production by NK cells: IL-13-producing NK and T cells are present in vivo in the absence of IFN-gamma. *Journal of immunology (Baltimore, Md: 1950)*, Vol. 162, No. 1, (Jan 1, 1999), pp. 51-59, 0022-1767; 0022-1767.
- Hou, J., Schindler, U., Henzel, W. J., Ho, T. C., Brasseur, M., & McKnight, S. L. (1994). An interleukin-4-induced transcription factor: IL-4 Stat. *Science*, Vol. 265, No. 5179, (1994), pp. 1701-1706.
- Huang, H., & Paul, W. E. (2000). Protein tyrosine phosphatase activity is required for IL-4 induction of IL-4 receptor alpha-chain. *Journal of Immunology*, Vol. 164, No. 3, (Feb 1, 2000), pp. 1211-1215.
- Huang, S. K., Xiao, H. Q., Kleine-Tebbe, J., Paciotti, G., Marsh, D. G., Lichtenstein, L. M., & Liu, M. C. (1995). IL-13 expression at the sites of allergen challenge in patients with asthma. *Journal of Immunology*, Vol. 155, No. 5, (Sep 1, 1995), pp. 2688-2694.
- Huang, Z., Coleman, J. M., Su, Y., Mann, M., Ryan, J., Shultz, L. D., & Huang, H. (2005). SHP-1 regulates STAT6 phosphorylation and IL-4-mediated function in a cell type-specific manner. *Cytokine*, Vol. 29, No. 3, (Feb 7, 2005), pp. 118-124.
- Idzerda, R. L., March, C. J., Mosley, B., Lyman, S. D., Vanden Bos, T., Gimpel, S. D., Din, W. S., Grabstein, K. H., Widmer, M. B., Park, L. S., & et al. (1990). Human interleukin 4 receptor confers biological responsiveness and defines a novel receptor superfamily. *Journal of Experimental Medicine*, Vol. 171, No. 3, (Mar 1, 1990), pp. 861-873.
- Imani, F., Rager, K. J., Catipovic, B., & Marsh, D. G. (1997). Interleukin-4 (IL-4) induces phosphatidylinositol 3-kinase (p85) dephosphorylation. Implications for the role of SHP-1 in the IL-4-induced signals in human B cells. *Journal of Biological Chemistry*, Vol. 272, No. 12, (Mar 21, 1997), pp. 7927-7931.
- Inoue, A., Yasuda, T., Yamamoto, T., & Yamanashi, Y. (2007). Dok-1 is a positive regulator of IL-4 signalling and IgE response. *J Biochem*, Vol. 142, No. 2, (Aug, 2007), pp. 257-263.
- Ito, T., Suzuki, S., Kanaji, S., Shiraishi, H., Ohta, S., Arima, K., Tanaka, G., Tamada, T., Honjo, E., Garcia, K. C., Kuroki, R., & Izuhara, K. (2009). Distinct structural requirements for interleukin-4 (IL-4) and IL-13 binding to the shared IL-13 receptor facilitate cellular tuning of cytokine responsiveness. *Journal of Biological Chemistry*, Vol. 284, No. 36, (Sep 4, 2009), pp. 24289-24296.
- Izuhara, K., & Harada, N. (1993). Interleukin-4 (IL-4) induces protein tyrosine phosphorylation of the IL-4 receptor and association of phosphatidylinositol 3-kinase to the IL-4 receptor in a mouse T cell line, HT2. *Journal of Biological Chemistry*, Vol. 268, No. 18, (Jun 25, 1993), pp. 13097-13102.

- Jatakanon, A., Uasuf, C., Maziak, W., Lim, S., Chung, K. F., & Barnes, P. J. (1999). Neutrophilic inflammation in severe persistent asthma. *American Journal of Respiratory and Critical Care Medicine*, Vol. 160, No. 5 Pt 1, (Nov, 1999), pp. 1532-1539.
- Junttila, I. S., Mizukami, K., Dickensheets, H., Meier-Schellersheim, M., Yamane, H., Donnelly, R. P., Paul, W. E. (2008). Tuning sensitivity to IL-4 and IL-13: Differential expression of IL-4R α , IL-13R α 1 and γ c regulates relative cytokine sensitivity. *Journal of Experimental Medicine*, Vol., No., (2008), pp.
- Kaiko, G. E., Phipps, S., Angkasekwinai, P., Dong, C., & Foster, P. S. (2010). NK cell deficiency predisposes to viral-induced Th2-type allergic inflammation via epithelial-derived IL-25. *Journal of Immunology*, Vol. 185, No. 8, (Oct 15, 2010), pp. 4681-4690.
- Kang, C. M., Jang, A. S., Ahn, M. H., Shin, J. A., Kim, J. H., Choi, Y. S., Rhim, T. Y., & Park, C. S. (2005). Interleukin-25 and interleukin-13 production by alveolar macrophages in response to particles. *American Journal of Respiratory Cell and Molecular Biology*, Vol. 33, No. 3, (Sep, 2005), pp. 290-296.
- Kaplan, M. H., Schindler, U., Smiley, S. T., & Grusby, M. J. (1996). Stat6 is required for mediating responses to IL-4 and for development of Th2 cells. *Immunity*, Vol. 4, No. 3, (1996), pp. 313-319.
- Kaplan, M. H., Whitfield, J. R., Boros, D. L., & Grusby, M. J. (1998). Th2 cells are required for the *Schistosoma mansoni* egg-induced granulomatous response. *Journal of Immunology*, Vol. 160, No. 4, (Feb 15, 1998), pp. 1850-1856.
- Kashiwada, M., Giallourakis, C. C., Pan, P. Y., & Rothman, P. B. (2001). Immunoreceptor tyrosine-based inhibitory motif of the IL-4 receptor associates with SH2-containing phosphatases and regulates IL-4-induced proliferation. *Journal of Immunology*, Vol. 167, No. 11, (Dec 1, 2001), pp. 6382-6387.
- Keegan, A. D., Johnston, J. A., Tortolani, P. J., McReynolds, L. J., Kinzer, C., O'Shea, J. J., & Paul, W. E. (1995). Similarities and differences in signal transduction by interleukin 4 and interleukin 13: analysis of Janus kinase activation. *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 92, No. 17, (Aug 15, 1995), pp. 7681-7685.
- Keegan, A. D., Nelms, K., White, M., Wang, L. M., Pierce, J. H., & Paul, W. E. (1994). An IL-4 receptor region containing an insulin receptor motif is important for IL-4-mediated IRS-1 phosphorylation and cell growth. *Cell*, Vol. 76, No. 5, (Mar 11, 1994), pp. 811-820.
- Kelly-Welch, A. E., Melo, M. E., Smith, E., Ford, A. Q., Haudenschild, C., Noben-Trauth, N., & Keegan, A. D. (2004). Complex role of the IL-4 receptor alpha in a murine model of airway inflammation: expression of the IL-4 receptor alpha on nonlymphoid cells of bone marrow origin contributes to severity of inflammation. *Journal of Immunology*, Vol. 172, No. 7, (Apr 1, 2004), pp. 4545-4555.
- Kelly-Welch, A. E., Wang, H. Y., Wang, L. M., Pierce, J. H., Jay, G., Finkelman, F., & Keegan, A. D. (2004). Transgenic expression of insulin receptor substrate 2 in murine B cells alters the cell density-dependence of IgE production in vitro and enhances IgE production in vivo. *Journal of Immunology*, Vol. 172, No. 5, (Mar 1, 2004), pp. 2803-2810.
- Khattari, R., Cox, T., Yasayko, S. A., & Ramsdell, F. (2003). An essential role for Scurfin in CD4+CD25+ T regulatory cells. *Nature Immunology*, Vol. 4, No. 4, (Apr, 2003), pp. 337-342.
- Khoo, S. K., Zhang, G., Backer, V., Porsbjerg, C., Nepper-Christensen, S., Creegan, R., Baynam, G., de Klerk, N., Rossi, G. A., Hagel, I., Di Prisco, M. C., Lynch, N., Britton, J., Hall, I., Musk, A. W., Goldblatt, J., & Le Souef, P. N. (2006). Associations of a

- novel IL4RA polymorphism, Ala57Thr, in Greenlander Inuit. *Journal of Allergy and Clinical Immunology*, Vol. 118, No. 3, (Sep, 2006), pp. 627-634.
- Kleinschek, M. A., Owyang, A. M., Joyce-Shaikh, B., Langrish, C. L., Chen, Y., Gorman, D. M., Blumenschein, W. M., McClanahan, T., Brombacher, F., Hurst, S. D., Kastelein, R. A., & Cua, D. J. (2007). IL-25 regulates Th17 function in autoimmune inflammation. *The Journal of experimental medicine*, Vol. 204, No. 1, (Jan 22, 2007), pp. 161-170.
- Knutsen, A. P., Kariuki, B., Consolino, J. D., & Warrier, M. R. (2006). IL-4 alpha chain receptor (IL-4Ralpha) polymorphisms in allergic bronchopulmonary aspergillosis. *Clin Mol Allergy*, Vol. 4, No., (2006), pp. 3.
- Koettnitz, K., & Kalthoff, F. S. (1993). Human interleukin-4 receptor signaling requires sequences contained within two cytoplasmic regions. *European Journal of Immunology*, Vol. 23, No. 4, (Apr, 1993), pp. 988-991.
- Kondo, M., Takeshita, T., Ishii, N., Nakamura, M., Watanabe, S., Arai, K., & Sugamura, K. (1993). Sharing of the interleukin-2 (IL-2) receptor gamma chain between receptors for IL-2 and IL-4. *Science*, Vol. 262, No. 5141, (Dec 17, 1993), pp. 1874-1877.
- Kopf, M., Brombacher, F., Hodgkin, P. D., Ramsay, A. J., Milbourne, E. A., Dai, W. J., Ovington, K. S., Behm, C. A., Kohler, G., Young, I. G., & Matthaei, K. I. (1996). IL-5-deficient mice have a developmental defect in CD5+ B-1 cells and lack eosinophilia but have normal antibody and cytotoxic T cell responses. *Immunity*, Vol. 4, No. 1, (Jan, 1996), pp. 15-24.
- Kruse, S., Braun, S., & Deichmann, K. A. (2002). Distinct signal transduction processes by IL-4 and IL-13 and influences from the Q551R variant of the human IL-4 receptor alpha chain. *Respir Res*, Vol. 3, No., (2002), pp. 24.
- Kruse, S., Japha, T., Tedner, M., Sparholt, S. H., Forster, J., Kuehr, J., & Deichmann, K. A. (1999). The polymorphisms S503P and Q576R in the interleukin-4 receptor alpha gene are associated with atopy and influence the signal transduction. *Immunology*, Vol. 96, No. 3, (Mar, 1999), pp. 365-371.
- Kuhne, M. R., Pawson, T., Lienhard, G. E., & Feng, G. S. (1993). The insulin receptor substrate 1 associates with the SH2-containing phosphotyrosine phosphatase Syp. *Journal of Biological Chemistry*, Vol. 268, No. 16, (Jun 5, 1993), pp. 11479-11481.
- Kumaki, S., Kondo, M., Takeshita, T., Asao, H., Nakamura, M., & Sugamura, K. (1993). Cloning of the mouse interleukin 2 receptor gamma chain: demonstration of functional differences between the mouse and human receptors. *Biochemical and Biophysical Research Communications*, Vol. 193, No. 1, (May 28, 1993), pp. 356-363.
- Kuperman, D., Schofield, B., Wills-Karp, M., & Grusby, M. J. (1998). Signal transducer and activator of transcription factor 6 (Stat6)-deficient mice are protected from antigen-induced airway hyperresponsiveness and mucus production. *Journal of Experimental Medicine*, Vol. 187, No. 6, (1998), pp. 939-948.
- Kurata, H., Lee, H. J., O'Garra, A., & Arai, N. (1999). Ectopic expression of activated Stat6 induces the expression of Th2-specific cytokines and transcription factors in developing Th1 cells. *Immunity*, Vol. 11, No. 6, (Dec, 1999), pp. 677-688.
- Langrish, C. L., Chen, Y., Blumenschein, W. M., Mattson, J., Basham, B., Sedgwick, J. D., McClanahan, T., Kastelein, R. A., & Cua, D. J. (2005). IL-23 drives a pathogenic T cell population that induces autoimmune inflammation. *The Journal of experimental medicine*, Vol. 201, No. 2, (Jan 17, 2005), pp. 233-240.

- LaPorte, S. L., Juo, Z. S., Vaclavikova, J., Colf, L. A., Qi, X., Heller, N. M., Keegan, A. D., & Garcia, K. C. (2008). Molecular and structural basis of cytokine receptor pleiotropy in the interleukin-4/13 system. *Cell*, Vol. 132, No. 2, (Jan 25, 2008), pp. 259-272.
- Leonard, W. J., Noguchi, M., & Russell, S. M. (1994). Sharing of a common gamma chain, gamma c, by the IL-2, IL-4, and IL-7 receptors: implications for X-linked severe combined immunodeficiency (XSCID). *Advances in Experimental Medicine and Biology*, Vol. 365, No., (1994), pp. 225-232.
- Lorenz, U. (2009). SHP-1 and SHP-2 in T cells: two phosphatases functioning at many levels. *Immunological Reviews*, Vol. 228, No. 1, (Mar, 2009), pp. 342-359.
- Losman, J., Chen, X. P., Jiang, H., Pan, P. Y., Kashiwada, M., Giallourakis, C., Cowan, S., Foltényi, K., & Rothman, P. (1999). IL-4 signaling is regulated through the recruitment of phosphatases, kinases, and SOCS proteins to the receptor complex. *Cold Spring Harbor Symposia on Quantitative Biology*, Vol. 64, No., (1999), pp. 405-416.
- Lowenthal, J. W., Castle, B. E., Christiansen, J., Schreurs, J., Rennick, D., Arai, N., Hoy, P., Takebe, Y., & Howard, M. (1988). Expression of high affinity receptors for murine interleukin 4 (BSF-1) on hemopoietic and nonhemopoietic cells. *Journal of Immunology*, Vol. 140, No. 2, (Jan 15, 1988), pp. 456-464.
- Lupardus, P. J., Birnbaum, M. E., & Garcia, K. C. (2010). Molecular basis for shared cytokine recognition revealed in the structure of an unusually high affinity complex between IL-13 and IL-13Ralpha2. *Structure*, Vol. 18, No. 3, (Mar 10, 2010), pp. 332-342.
- Martin, P., Villares, R., Rodriguez-Mascarenhas, S., Zaballos, A., Leitges, M., Kovac, J., Sizing, I., Rennert, P., Marquez, G., Martinez, A. C., Diaz-Meco, M. T., & Moscat, J. (2005). Control of T helper 2 cell function and allergic airway inflammation by PKCzeta. *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 102, No. 28, (Jul 12, 2005), pp. 9866-9871.
- Mashima, R., Hishida, Y., Tezuka, T., & Yamanashi, Y. (2009). The roles of Dok family adapters in immunoreceptor signaling. *Immunological Reviews*, Vol. 232, No. 1, (Nov, 2009), pp. 273-285.
- Mathew, A., MacLean, J. A., DeHaan, E., Tager, A. M., Green, F. H., & Luster, A. D. (2001). Signal transducer and activator of transcription 6 controls chemokine production and T helper cell type 2 cell trafficking in allergic pulmonary inflammation. *Journal of Experimental Medicine*, Vol. 193, No. 9, (2001), pp. 1087-1096.
- Matsumura, M., Inoue, H., Matsumoto, T., Nakano, T., Fukuyama, S., Matsumoto, K., Takayama, K., Saito, M., Kawakami, K., & Nakanishi, Y. (2007). Endogenous metalloprotease solubilizes IL-13 receptor alpha2 in airway epithelial cells. *Biochemical and Biophysical Research Communications*, Vol. 360, No. 2, (Aug 24, 2007), pp. 464-469.
- McGeachy, M. J., Bak-Jensen, K. S., Chen, Y., Tato, C. M., Blumenschein, W., McClanahan, T., & Cua, D. J. (2007). TGF-beta and IL-6 drive the production of IL-17 and IL-10 by T cells and restrain T(H)-17 cell-mediated pathology. *Nature Immunology*, Vol. 8, No. 12, (Dec, 2007), pp. 1390-1397.
- McKinley, L., Alcorn, J. F., Peterson, A., Dupont, R. B., Kapadia, S., Logar, A., Henry, A., Irvin, C. G., Piganelli, J. D., Ray, A., & Kolls, J. K. (2008). TH17 cells mediate steroid-resistant airway inflammation and airway hyperresponsiveness in mice. *Journal of immunology (Baltimore, Md: 1950)*, Vol. 181, No. 6, (Sep 15, 2008), pp. 4089-4097.
- Mentink-Kane, M. M., Cheever, A. W., Thompson, R. W., Hari, D. M., Kabatereine, N. B., Vennervald, B. J., Ouma, J. H., Mwatha, J. K., Jones, F. M., Donaldson, D. D.,

- Grusby, M. J., Dunne, D. W., & Wynn, T. A. (2004). IL-13 receptor alpha 2 down-modulates granulomatous inflammation and prolongs host survival in schistosomiasis. *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 101, No. 2, (Jan 13, 2004), pp. 586-590.
- Mikita, T., Campbell, D., Wu, P., Williamson, K., & Schindler, U. (1996). Requirements for interleukin-4-induced gene expression and functional characterization of Stat6. *Molecular and Cellular Biology*, Vol. 16, No. 10, (1996), pp. 5811-5820.
- Min, B., & Paul, W. E. (2008). Basophils and type 2 immunity. *Current Opinion in Hematology*, Vol. 15, No. 1, (Jan, 2008), pp. 59-63.
- Mitsuyasu, H., Izuhara, K., Mao, X. Q., Gao, P. S., Arinobu, Y., Enomoto, T., Kawai, M., Sasaki, S., Dake, Y., Hamasaki, N., Shirakawa, T., & Hopkin, J. M. (1998). Ile50Val variant of IL4R alpha upregulates IgE synthesis and associates with atopic asthma. *Nature Genetics*, Vol. 19, No. 2, (Jun, 1998), pp. 119-120.
- Mitsuyasu, H., Yanagihara, Y., Mao, X. Q., Gao, P. S., Arinobu, Y., Ihara, K., Takabayashi, A., Hara, T., Enomoto, T., Sasaki, S., Kawai, M., Hamasaki, N., Shirakawa, T., Hopkin, J. M., & Izuhara, K. (1999). Cutting edge: dominant effect of Ile50Val variant of the human IL-4 receptor alpha-chain in IgE synthesis. *Journal of Immunology*, Vol. 162, No. 3, (Feb 1, 1999), pp. 1227-1231.
- Molet, S., Hamid, Q., Davoine, F., Nutku, E., Taha, R., Page, N., Olivenstein, R., Elias, J., & Chakir, J. (2001). IL-17 is increased in asthmatic airways and induces human bronchial fibroblasts to produce cytokines. *The Journal of Allergy and Clinical Immunology*, Vol. 108, No. 3, (Sep, 2001), pp. 430-438.
- Morris, S. C., Orekhova, T., Meadows, M. J., Heidorn, S. M., Yang, J., & Finkelman, F. D. (2006). IL-4 induces in vivo production of IFN-gamma by NK and NKT cells. *Journal of Immunology*, Vol. 176, No. 9, (May 1, 2006), pp. 5299-5305.
- Mosley, B., Beckmann, M. P., March, C. J., Idzerda, R. L., Gimpel, S. D., VandenBos, T., Friend, D., Alpert, A., Anderson, D., Jackson, J., & et al. (1989). The murine interleukin-4 receptor: molecular cloning and characterization of secreted and membrane bound forms. *Cell*, Vol. 59, No. 2, (Oct 20, 1989), pp. 335-348.
- Munitz, A., Brandt, E. B., Mingler, M., Finkelman, F. D., & Rothenberg, M. E. (2008). Distinct roles for IL-13 and IL-4 via IL-13 receptor alpha1 and the type II IL-4 receptor in asthma pathogenesis. *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 105, No. 20, (May 20, 2008), pp. 7240-7245.
- Murakami, M., Narazaki, M., Hibi, M., Yawata, H., Yasukawa, K., Hamaguchi, M., Taga, T., & Kishimoto, T. (1991). Critical cytoplasmic region of the interleukin 6 signal transducer gp130 is conserved in the cytokine receptor family. *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 88, No. 24, (Dec 15, 1991), pp. 11349-11353.
- Murata, T., Noguchi, P. D., & Puri, R. K. (1996). IL-13 induces phosphorylation and activation of JAK2 Janus kinase in human colon carcinoma cell lines: similarities between IL-4 and IL-13 signaling. *Journal of Immunology*, Vol. 156, No. 8, (Apr 15, 1996), pp. 2972-2978.
- Murata, T., & Puri, R. K. (1997). Comparison of IL-13- and IL-4-induced signaling in EBV-immortalized human B cells. *Cellular Immunology*, Vol. 175, No. 1, (Jan 10, 1997), pp. 33-40.

- Nelms, K., Keegan, A. D., Zamorano, J., Ryan, J. J., & Paul, W. E. (1999). The IL-4 receptor: signaling mechanisms and biologic functions. *Annual Review of Immunology*, Vol. 17, No., (1999), pp. 701-738.
- Nelms, K., Snow, A. L., Hu-Li, J., & Paul, W. E. (1998). FRIP, a hematopoietic cell-specific rasGAP-interacting protein phosphorylated in response to cytokine stimulation. *Immunity*, Vol. 9, No. 1, (Jul, 1998), pp. 13-24.
- Nelson, B. H., Lord, J. D., & Greenberg, P. D. (1996). A membrane-proximal region of the interleukin-2 receptor gamma c chain sufficient for Jak kinase activation and induction of proliferation in T cells. *Molecular and Cellular Biology*, Vol. 16, No. 1, (Jan, 1996), pp. 309-317.
- Newcomb, D. C., Boswell, M. G., Zhou, W., Huckabee, M. M., Goleniewska, K., Sevin, C. M., Hershey, G. K., Kolls, J. K., & Peebles, R. S., Jr. (2011). Human TH17 cells express a functional IL-13 receptor and IL-13 attenuates IL-17A production. *Journal of Allergy and Clinical Immunology*, Vol. 127, No. 4, (Apr, 2011), pp. 1006-1013.
- Newcomb, D. C., Zhou, W., Moore, M. L., Goleniewska, K., Hershey, G. K., Kolls, J. K., & Peebles, R. S., Jr. (2009). A functional IL-13 receptor is expressed on polarized murine CD4⁺ Th17 cells and IL-13 signaling attenuates Th17 cytokine production. *Journal of Immunology*, Vol. 182, No. 9, (May 1, 2009), pp. 5317-5321.
- Noguchi, E., Shibasaki, M., Arinami, T., Takeda, K., Yokouchi, Y., Kobayashi, K., Imoto, N., Nakahara, S., Matsui, A., & Hamaguchi, H. (1999). No association between atopy/asthma and the ILe50Val polymorphism of IL-4 receptor. *American Journal of Respiratory and Critical Care Medicine*, Vol. 160, No. 1, (Jul, 1999), pp. 342-345.
- Noguchi, M., Adelstein, S., Cao, X., & Leonard, W. J. (1993). Characterization of the human interleukin-2 receptor gamma chain gene. *Journal of Biological Chemistry*, Vol. 268, No. 18, (Jun 25, 1993), pp. 13601-13608.
- Noguchi, M., Yi, H., Rosenblatt, H. M., Filipovich, A. H., Adelstein, S., Modi, W. S., McBride, O. W., & Leonard, W. J. (1993). Interleukin-2 receptor gamma chain mutation results in X-linked severe combined immunodeficiency in humans. *Cell*, Vol. 73, No. 1, (Apr 9, 1993), pp. 147-157.
- O'Connor, J. C., Sherry, C. L., Guest, C. B., & Freund, G. G. (2007). Type 2 diabetes impairs insulin receptor substrate-2-mediated phosphatidylinositol 3-kinase activity in primary macrophages to induce a state of cytokine resistance to IL-4 in association with overexpression of suppressor of cytokine signaling-3. *Journal of Immunology*, Vol. 178, No. 11, (Jun 1, 2007), pp. 6886-6893.
- Ober, C., Leavitt, S. A., Tsalenko, A., Howard, T. D., Hoki, D. M., Daniel, R., Newman, D. L., Wu, X., Parry, R., Lester, L. A., Solway, J., Blumenthal, M., King, R. A., Xu, J., Meyers, D. A., Bleecker, E. R., & Cox, N. J. (2000). Variation in the interleukin 4-receptor alpha gene confers susceptibility to asthma and atopy in ethnically diverse populations. *American Journal of Human Genetics*, Vol. 66, No. 2, (Feb, 2000), pp. 517-526.
- Ogata, H., Ford, D., Kouttab, N., King, T. C., Vita, N., Minty, A., Stoeckler, J., Morgan, D., Girasole, C., Morgan, J. W., & Maizel, A. L. (1998). Regulation of interleukin-13 receptor constituents on mature human B lymphocytes. *Journal of Biological Chemistry*, Vol. 273, No. 16, (Apr 17, 1998), pp. 9864-9871.
- Oh, S. Y., Zheng, T., Bailey, M. L., Barber, D. L., Schroeder, J. T., Kim, Y. K., & Zhu, Z. (2007). Src homology 2 domain-containing inositol 5-phosphatase 1 deficiency leads to a

- spontaneous allergic inflammation in the murine lung. *Journal of Allergy and Clinical Immunology*, Vol. 119, No. 1, (Jan, 2007), pp. 123-131.
- Orchansky, P. L., Kwan, R., Lee, F., & Schrader, J. W. (1999). Characterization of the cytoplasmic domain of interleukin-13 receptor-alpha. *Journal of Biological Chemistry*, Vol. 274, No. 30, (Jul 23, 1999), pp. 20818-20825.
- Pace, L., Pioli, C., & Doria, G. (2005). IL-4 modulation of CD4+CD25+ T regulatory cell-mediated suppression. *Journal of Immunology*, Vol. 174, No. 12, (Jun 15, 2005), pp. 7645-7653.
- Pace, L., Rizzo, S., Palombi, C., Brombacher, F., & Doria, G. (2006). Cutting edge: IL-4-induced protection of CD4+CD25- Th cells from CD4+CD25+ regulatory T cell-mediated suppression. *Journal of Immunology*, Vol. 176, No. 7, (Apr 1, 2006), pp. 3900-3904.
- Paliard, X., de Waal Malefijt, R., Yssel, H., Blanchard, D., Chretien, I., Abrams, J., de Vries, J., & Spits, H. (1988). Simultaneous production of IL-2, IL-4, and IFN-gamma by activated human CD4+ and CD8+ T cell clones. *Journal of Immunology*, Vol. 141, No. 3, (Aug 1, 1988), pp. 849-855.
- Park, L. S., Friend, D., Grabstein, K., & Urdal, D. L. (1987). Characterization of the high-affinity cell-surface receptor for murine B-cell-stimulating factor 1. *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 84, No. 6, (Mar, 1987), pp. 1669-1673.
- Park, L. S., Friend, D., Sassenfeld, H. M., & Urdal, D. L. (1987). Characterization of the human B cell stimulatory factor 1 receptor. *Journal of Experimental Medicine*, Vol. 166, No. 2, (Aug 1, 1987), pp. 476-488.
- Pelicci, G., Lanfranccone, L., Grignani, F., McGlade, J., Cavallo, F., Forni, G., Nicoletti, I., Pawson, T., & Pelicci, P. G. (1992). A novel transforming protein (SHC) with an SH2 domain is implicated in mitogenic signal transduction. *Cell*, Vol. 70, No. 1, (Jul 10, 1992), pp. 93-104.
- Peritt, D., Robertson, S., Gri, G., Showe, L., Aste-Amezaga, M., & Trinchieri, G. (1998). Differentiation of human NK cells into NK1 and NK2 subsets. *Journal of Immunology* Vol. 161, No. 11, (Dec 1, 1998), pp. 5821-5824.
- Pesu, M., Takaluoma, K., Aittomaki, S., Lagerstedt, A., Saksela, K., Kovanen, P. E., & Silvennoinen, O. (2000). Interleukin-4-induced transcriptional activation by stat6 involves multiple serine/threonine kinase pathways and serine phosphorylation of stat6. *Blood*, Vol. 95, No. 2, (Jan 15, 2000), pp. 494-502.
- Piccinni, M. P., Macchia, D., Parronchi, P., Giudizi, M. G., Bani, D., Alterini, R., Grossi, A., Ricci, M., Maggi, E., & Romagnani, S. (1991). Human bone marrow non-B, non-T cells produce interleukin 4 in response to cross-linkage of Fc epsilon and Fc gamma receptors. *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 88, No. 19, (Oct 1, 1991), pp. 8656-8660.
- Plaut, M., Pierce, J. H., Watson, C. J., Hanley-Hyde, J., Nordan, R. P., & Paul, W. E. (1989). Mast cell lines produce lymphokines in response to cross-linkage of Fc epsilon RI or to calcium ionophores. *Nature*, Vol. 339, No. 6219, (May 4, 1989), pp. 64-67.
- Pouliot, P., Turmel, V., Gelinas, E., Laviolette, M., & Bissonnette, E. Y. (2005). Interleukin-4 production by human alveolar macrophages. *Clinical and Experimental Allergy*, Vol. 35, No. 6, (Jun, 2005), pp. 804-810.

- Prots, I., Skapenko, A., Wendler, J., Mattyasovszky, S., Yone, C. L., Spriewald, B., Burkhardt, H., Rau, R., Kalden, J. R., Lipsky, P. E., & Schulze-Koops, H. (2006). Association of the IL4R single-nucleotide polymorphism I50V with rapidly erosive rheumatoid arthritis. *Arthritis and Rheumatism*, Vol. 54, No. 5, (May, 2006), pp. 1491-1500.
- Pruett, W., Yuan, Y., Rose, E., Batzer, A. G., Harada, N., & Skolnik, E. Y. (1995). Association between GRB2/Sos and insulin receptor substrate 1 is not sufficient for activation of extracellular signal-regulated kinases by interleukin-4: implications for Ras activation by insulin. *Molecular and Cellular Biology*, Vol. 15, No. 3, (Mar, 1995), pp. 1778-1785.
- Ra, C., Kuromitsu, S., Hirose, T., Yasuda, S., Furuichi, K., & Okumura, K. (1993). Soluble human high-affinity receptor for IgE abrogates the IgE-mediated allergic reaction. *International Immunology*, Vol. 5, No. 1, (Jan, 1993), pp. 47-54.
- Rahaman, S. O., Sharma, P., Harbor, P. C., Aman, M. J., Vogelbaum, M. A., & Haque, S. J. (2002). IL-13R(alpha)2, a decoy receptor for IL-13 acts as an inhibitor of IL-4-dependent signal transduction in glioblastoma cells. *Cancer Research*, Vol. 62, No. 4, (Feb 15, 2002), pp. 1103-1109.
- Ramalingam, T. R., Pesce, J. T., Sheikh, F., Cheever, A. W., Mentink-Kane, M. M., Wilson, M. S., Stevens, S., Valenzuela, D. M., Murphy, A. J., Yancopoulos, G. D., Urban, J. F., Jr., Donnelly, R. P., & Wynn, T. A. (2008). Unique functions of the type II interleukin 4 receptor identified in mice lacking the interleukin 13 receptor alpha1 chain. *Nat Immunol*, Vol. 9, No. 1, (Jan, 2008), pp. 25-33.
- Rankin, J. A., Picarella, D. E., Geba, G. P., Temann, U. A., Prasad, B., DiCosmo, B., Tarallo, A., Stripp, B., Whitsett, J., & Flavell, R. A. (1996). Phenotypic and physiologic characterization of transgenic mice expressing interleukin 4 in the lung: lymphocytic and eosinophilic inflammation without airway hyperreactivity. *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 93, No. 15, (Jul 23, 1996), pp. 7821-7825.
- Rauh, M. J., Ho, V., Pereira, C., Sham, A., Sly, L. M., Lam, V., Huxham, L., Minchinton, A. I., Mui, A., & Krystal, G. (2005). SHIP represses the generation of alternatively activated macrophages. *Immunity*, Vol. 23, No. 4, (Oct, 2005), pp. 361-374.
- Ravetch, J. V., & Lanier, L. L. (2000). Immune inhibitory receptors. *Science*, Vol. 290, No. 5489, (Oct 6, 2000), pp. 84-89.
- Reichel, M., Nelson, B. H., Greenberg, P. D., & Rothman, P. B. (1997). The IL-4 receptor alpha-chain cytoplasmic domain is sufficient for activation of JAK-1 and STAT6 and the induction of IL-4-specific gene expression. *Journal of Immunology*, Vol. 158, No. 12, (Jun 15, 1997), pp. 5860-5867.
- Risma, K. A., Wang, N., Andrews, R. P., Cunningham, C. M., Ericksen, M. B., Bernstein, J. A., Chakraborty, R., & Hershey, G. K. (2002). V75R576 IL-4 receptor alpha is associated with allergic asthma and enhanced IL-4 receptor function. *Journal of Immunology*, Vol. 169, No. 3, (Aug 1, 2002), pp. 1604-1610.
- Rolling, C., Treton, D., Pellegrini, S., Galanaud, P., & Richard, Y. (1996). IL4 and IL13 receptors share the gamma c chain and activate STAT6, STAT3 and STAT5 proteins in normal human B cells. *FEBS Letters*, Vol. 393, No. 1, (1996), pp. 53-56.
- Romagnani, S. (1992). Induction of TH1 and TH2 responses: a key role for the 'natural' immune response? *Immunology Today*, Vol. 13, No. 10, (Oct, 1992), pp. 379-381.

- Roy, B., Bhattacharjee, A., Xu, B., Ford, D., Maizel, A. L., & Cathcart, M. K. (2002). IL-13 signal transduction in human monocytes: phosphorylation of receptor components, association with Jaks, and phosphorylation/activation of Stats. *Journal of Leukocyte Biology*, Vol. 72, No. 3, (Sep, 2002), pp. 580-589.
- Rui, L., Fisher, T. L., Thomas, J., & White, M. F. (2001). Regulation of insulin/insulin-like growth factor-1 signaling by proteasome-mediated degradation of insulin receptor substrate-2. *Journal of Biological Chemistry*, Vol. 276, No. 43, (Oct 26, 2001), pp. 40362-40367.
- Russell, S. M., Johnston, J. A., Noguchi, M., Kawamura, M., Bacon, C. M., Friedmann, M., Berg, M., McVicar, D. W., Witthuhn, B. A., Silvennoinen, O., & et al. (1994). Interaction of IL-2R beta and gamma c chains with Jak1 and Jak3: implications for XSCID and XCID. *Science*, Vol. 266, No. 5187, (Nov 11, 1994), pp. 1042-1045.
- Russell, S. M., Keegan, A. D., Harada, N., Nakamura, Y., Noguchi, M., Leland, P., Friedmann, M. C., Miyajima, A., Puri, R. K., Paul, W. E., & et al. (1993). Interleukin-2 receptor gamma chain: a functional component of the interleukin-4 receptor. *Science*, Vol. 262, No. 5141, (Dec 17, 1993), pp. 1880-1883.
- Ryan, J. J., McReynolds, L. J., Keegan, A., Wang, L. H., Garfein, E., Rothman, P., Nelms, K., & Paul, W. E. (1996). Growth and gene expression are predominantly controlled by distinct regions of the human IL-4 receptor. *Immunity*, Vol. 4, No. 2, (Feb, 1996), pp. 123-132.
- Sabin, E. A., Kopf, M. A., & Pearce, E. J. (1996). *Schistosoma mansoni* egg-induced early IL-4 production is dependent upon IL-5 and eosinophils. *Journal of Experimental Medicine*, Vol. 184, No. 5, (Nov 1, 1996), pp. 1871-1878.
- Sakaguchi, S., Yamaguchi, T., Nomura, T., & Ono, M. (2008). Regulatory T cells and immune tolerance. *Cell*, Vol. 133, No. 5, (May 30, 2008), pp. 775-787.
- Satoh, T., Nakafuku, M., Miyajima, A., & Kaziro, Y. (1991). Involvement of ras p21 protein in signal-transduction pathways from interleukin 2, interleukin 3, and granulocyte/macrophage colony-stimulating factor, but not from interleukin 4. *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 88, No. 8, (Apr 15, 1991), pp. 3314-3318.
- Scheffold, A., Murphy, K. M., & Hofer, T. (2007). Competition for cytokines: T(reg) cells take all. *Nature Immunology*, Vol. 8, No. 12, (Dec, 2007), pp. 1285-1287.
- Schroeder, J. T., Kagey-Sobotka, A., & Lichtenstein, L. M. (1995). The role of the basophil in allergic inflammation. *Allergy*, Vol. 50, No. 6, (Jun, 1995), pp. 463-472.
- Seder, R. A., Boulay, J. L., Finkelman, F., Barbier, S., Ben-Sasson, S. Z., Le Gros, G., & Paul, W. E. (1992). CD8+ T cells can be primed in vitro to produce IL-4. *Journal of Immunology*, Vol. 148, No. 6, (Mar 15, 1992), pp. 1652-1656.
- Sehra, S., Bruns, H. A., Ahyi, A. N., Nguyen, E. T., Schmidt, N. W., Michels, E. G., von Bulow, G. U., & Kaplan, M. H. (2008). IL-4 is a critical determinant in the generation of allergic inflammation initiated by a constitutively active Stat6. *Journal of Immunology*, Vol. 180, No. 5, (Mar 1, 2008), pp. 3551-3559.
- Sehra, S., Yao, Y., Howell, M. D., Nguyen, E. T., Kansas, G. S., Leung, D. Y., Travers, J. B., & Kaplan, M. H. (2010). IL-4 regulates skin homeostasis and the predisposition toward allergic skin inflammation. *Journal of Immunology*, Vol. 184, No. 6, (Mar 15, 2010), pp. 3186-3190.

- Seldin, D. C., & Leder, P. (1994). Mutational analysis of a critical signaling domain of the human interleukin 4 receptor. *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 91, No. 6, (Mar 15, 1994), pp. 2140-2144.
- Shankaranarayanan, P., Chaitidis, P., Kuhn, H., & Nigam, S. (2001). Acetylation by histone acetyltransferase CREB-binding protein/p300 of STAT6 is required for transcriptional activation of the 15-lipoxygenase- 1 gene. *Journal of Biological Chemistry*, Vol. 276, No. 46, (2001), pp. 42753-42760.
- Shevach, E. M. (2009). Mechanisms of foxp3+ T regulatory cell-mediated suppression. *Immunity*, Vol. 30, No. 5, (May, 2009), pp. 636-645.
- Shirakawa, I., Deichmann, K. A., Izuhara, I., Mao, I., Adra, C. N., & Hopkin, J. M. (2000). Atopy and asthma: genetic variants of IL-4 and IL-13 signalling. *Immunology Today*, Vol. 21, No. 2, (Feb, 2000), pp. 60-64.
- Shirakawa, T., Kawazoe, Y., Tsujikawa, T., Jung, D., Sato, S., & Uesugi, M. (2011). Deactivation of STAT6 through serine 707 phosphorylation by JNK. *Journal of Biological Chemistry*, Vol. 286, No. 5, (Feb 4, 2011), pp. 4003-4010.
- Shirey, K., Cole, L. E., Keegan, A. D., Elkins, K. E., Vogel, S. N. (2008). *Francisella tularensis* LVS induces Macrophage Alternative Activation as a Survival Mechanism. *Journal of Immunology*, Vol., No., (2008), pp.
- Shirey, K. A., Pletneva, L. M., Puche, A. C., Keegan, A. D., Prince, G. A., Blanco, J. C., & Vogel, S. N. (2010). Control of RSV-induced lung injury by alternatively activated macrophages is IL-4R alpha-, TLR4-, and IFN-beta-dependent. *Mucosal Immunol*, Vol. 3, No. 3, (May, 2010), pp. 291-300.
- Sozzani, P., Hasan, L., Seguelas, M. H., Caput, D., Ferrara, P., Pipy, B., & Cambon, C. (1998). IL-13 induces tyrosine phosphorylation of phospholipase C gamma-1 following IRS-2 association in human monocytes: relationship with the inhibitory effect of IL-13 on ROI production. *Biochemical and Biophysical Research Communications*, Vol. 244, No. 3, (Mar 27, 1998), pp. 665-670.
- Starr, R., Willson, T. A., Viney, E. M., Murray, L. J., Rayner, J. R., Jenkins, B. J., Gonda, T. J., Alexander, W. S., Metcalf, D., Nicola, N. A., & Hilton, D. J. (1997). A family of cytokine-inducible inhibitors of signalling. *Nature*, Vol. 387, No. 6636, (Jun 26, 1997), pp. 917-921.
- Stephenson, L., Johns, M. H., Woodward, E., Mora, A. L., & Boothby, M. (2004). An IL-4R alpha allelic variant, I50, acts as a gain-of-function variant relative to V50 for Stat6, but not Th2 differentiation. *Journal of Immunology*, Vol. 173, No. 7, (Oct 1, 2004), pp. 4523-4528.
- Stritesky, G. L., Muthukrishnan, R., Sehra, S., Goswami, R., Pham, D., Travers, J., Nguyen, E. T., Levy, D. E., & Kaplan, M. H. (2011). The transcription factor STAT3 is required for T helper 2 cell development. *Immunity*, Vol. 34, No. 1, (Jan 28, 2011), pp. 39-49.
- Strober, W., Kitani, A., Fichtner-Feigl, S., & Fuss, I. J. (2009). The signaling function of the IL-13Ralpha2 receptor in the development of gastrointestinal fibrosis and cancer surveillance. *Curr Mol Med*, Vol. 9, No. 6, (Aug, 2009), pp. 740-750.
- Sun, X. J., Pons, S., Wang, L. M., Zhang, Y., Yenush, L., Burks, D., Myers, M. G., Jr., Glasheen, E., Copeland, N. G., Jenkins, N. A., Pierce, J. H., & White, M. F. (1997). The IRS-2 gene on murine chromosome 8 encodes a unique signaling adapter for insulin and cytokine action. *Molecular Endocrinology*, Vol. 11, No. 2, (Feb, 1997), pp. 251-262.


- Sun, X. J., Rothenberg, P., Kahn, C. R., Backer, J. M., Araki, E., Wilden, P. A., Cahill, D. A., Goldstein, B. J., & White, M. F. (1991). Structure of the insulin receptor substrate IRS-1 defines a unique signal transduction protein. *Nature*, Vol. 352, No. 6330, (Jul 4, 1991), pp. 73-77.
- Sun, X. J., Wang, L. M., Zhang, Y., Yenush, L., Myers, M. G., Jr., Glasheen, E., Lane, W. S., Pierce, J. H., & White, M. F. (1995). Role of IRS-2 in insulin and cytokine signalling. *Nature*, Vol. 377, No. 6545, (Sep 14, 1995), pp. 173-177.
- Suzuki, K., Nakajima, H., Kagami, S., Suto, A., Ikeda, K., Hirose, K., Hiwasa, T., Takeda, K., Saito, Y., Akira, S., & Iwamoto, I. (2002). Proteolytic processing of Stat6 signaling in mast cells as a negative regulatory mechanism. *Journal of Experimental Medicine*, Vol. 196, No. 1, (Jul 1, 2002), pp. 27-38.
- Tabata, Y., Chen, W., Warriar, M. R., Gibson, A. M., Daines, M. O., & Hershey, G. K. (2006). Allergy-driven alternative splicing of IL-13 receptor alpha2 yields distinct membrane and soluble forms. *Journal of Immunology*, Vol. 177, No. 11, (Dec 1, 2006), pp. 7905-7912.
- Tachdjian, R., Al Khatib, S., Schwinglshackl, A., Kim, H. S., Chen, A., Blasioli, J., Mathias, C., Kim, H. Y., Umetsu, D. T., Oettgen, H. C., & Chatila, T. A. (2010). In vivo regulation of the allergic response by the IL-4 receptor alpha chain immunoreceptor tyrosine-based inhibitory motif. *Journal of Allergy and Clinical Immunology*, Vol. 125, No. 5, (May, 2010), pp. 1128-1136.
- Tachdjian, R., Mathias, C., Al Khatib, S., Bryce, P. J., Kim, H. S., Blaeser, F., O'Connor, B. D., Rzymkiewicz, D., Chen, A., Holtzman, M. J., Hershey, G. K., Garn, H., Harb, H., Renz, H., Oettgen, H. C., & Chatila, T. A. (2009). Pathogenicity of a disease-associated human IL-4 receptor allele in experimental asthma. *Journal of Experimental Medicine*, Vol. 206, No. 10, (Sep 28, 2009), pp. 2191-2204.
- Takaki, H., Ichiyama, K., Koga, K., Chinen, T., Takaesu, G., Sugiyama, Y., Kato, S., Yoshimura, A., & Kobayashi, T. (2008). STAT6 Inhibits TGF-beta1-mediated Foxp3 induction through direct binding to the Foxp3 promoter, which is reverted by retinoic acid receptor. *The Journal of biological chemistry*, Vol. 283, No. 22, (May 30, 2008), pp. 14955-14962.
- Takeda, K., Tanaka, T., Shi, W., Matsumoto, M., Minami, M., Kashiwamura, S., Nakanishi, K., Yoshida, N., Kishimoto, T., & Akira, S. (1996). Essential role of Stat6 in IL-4 signalling. *Nature*, Vol. 380, No. 6575, (Apr 18, 1996), pp. 627-630.
- Takeshita, T., Asao, H., Ohtani, K., Ishii, N., Kumaki, S., Tanaka, N., Munakata, H., Nakamura, M., & Sugamura, K. (1992). Cloning of the gamma chain of the human IL-2 receptor. *Science*, Vol. 257, No. 5068, (Jul 17, 1992), pp. 379-382.
- Timonen, T., & Stenius-Aarniala, B. (1985). Natural killer cell activity in asthma. *Clinical and Experimental Immunology*, Vol. 59, No. 1, (Jan, 1985), pp. 85-90.
- Toru, H., Pawankar, R., Ra, C., Yata, J., & Nakahata, T. (1998). Human mast cells produce IL-13 by high-affinity IgE receptor cross-linking: enhanced IL-13 production by IL-4-primed human mast cells. *Journal of Allergy and Clinical Immunology*, Vol. 102, No. 3, (Sep, 1998), pp. 491-502.
- Toru, H., Ra, C., Nonoyama, S., Suzuki, K., Yata, J., & Nakahata, T. (1996). Induction of the high-affinity IgE receptor (Fc epsilon RI) on human mast cells by IL-4. *International Immunology*, Vol. 8, No. 9, (Sep, 1996), pp. 1367-1373.

- Umeshita-Suyama, R., Sugimoto, R., Akaiwa, M., Arima, K., Yu, B., Wada, M., Kuwano, M., Nakajima, K., Hamasaki, N., & Izuhara, K. (2000). Characterization of IL-4 and IL-13 signals dependent on the human IL-13 receptor alpha chain 1: redundancy of requirement of tyrosine residue for STAT3 activation. *International Immunology*, Vol. 12, No. 11, (Nov, 2000), pp. 1499-1509.
- van Panhuys, N., Prout, M., Forbes, E., Min, B., Paul, W. E., & Le Gros, G. (2011). Basophils are the major producers of IL-4 during primary helminth infection. *Journal of Immunology*, Vol. 186, No. 5, (Mar 1, 2011), pp. 2719-2728.
- Voehringer, D., Shinkai, K., & Locksley, R. M. (2004). Type 2 immunity reflects orchestrated recruitment of cells committed to IL-4 production. *Immunity*, Vol. 20, No. 3, (Mar, 2004), pp. 267-277.
- Wakashin, H., Hirose, K., Maezawa, Y., Kagami, S., Suto, A., Watanabe, N., Saito, Y., Hatano, M., Tokuhisa, T., Iwakura, Y., Puccetti, P., Iwamoto, I., & Nakajima, H. (2008). IL-23 and Th17 cells enhance Th2-cell-mediated eosinophilic airway inflammation in mice. *American Journal of Respiratory and Critical Care Medicine*, Vol. 178, No. 10, (Nov 15, 2008), pp. 1023-1032.
- Wallis, S. K., Cooney, L. A., Endres, J. L., Lee, M. J., Ryu, J., Somers, E. C., & Fox, D. A. (2011). A polymorphism in the interleukin-4 receptor affects the ability of interleukin-4 to regulate Th17 cells: a possible immunoregulatory mechanism for genetic control of the severity of rheumatoid arthritis. *Arthritis Res Ther*, Vol. 13, No. 1, (Feb 4, 2011), pp. R15.
- Wang, H. Y., Shelburne, C. P., Zamorano, J., Kelly, A. E., Ryan, J. J., & Keegan, A. D. (1999). Cutting edge: effects of an allergy-associated mutation in the human IL-4R alpha (Q576R) on human IL-4-induced signal transduction. *Journal of Immunology*, Vol. 162, No. 8, (Apr 15, 1999), pp. 4385-4389.
- Wang, H. Y., Zamorano, J., & Keegan, A. D. (1998). A role for the insulin-interleukin (IL)-4 receptor motif of the IL-4 receptor alpha-chain in regulating activation of the insulin receptor substrate 2 and signal transducer and activator of transcription 6 pathways. Analysis by mutagenesis. *Journal of Biological Chemistry*, Vol. 273, No. 16, (Apr 17, 1998), pp. 9898-9905.
- Wang, H. Y., Zamorano, J., Yoerkie, J. L., Paul, W. E., & Keegan, A. D. (1997). The IL-4-induced tyrosine phosphorylation of the insulin receptor substrate is dependent on JAK1 expression in human fibrosarcoma cells. *Journal of Immunology*, Vol. 158, No. 3, (Feb 1, 1997), pp. 1037-1040.
- Wang, I. M., Lin, H., Goldman, S. J., & Kobayashi, M. (2004). STAT-1 is activated by IL-4 and IL-13 in multiple cell types. *Molecular Immunology*, Vol. 41, No. 9, (Jul, 2004), pp. 873-884.
- Wang, L. M., Keegan, A. D., Li, W., Lienhard, G. E., Pacini, S., Gutkind, J. S., Myers, M. G., Jr., Sun, X. J., White, M. F., Aaronson, S. A., & et al. (1993). Common elements in interleukin 4 and insulin signaling pathways in factor-dependent hematopoietic cells. *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 90, No. 9, (May 1, 1993), pp. 4032-4036.
- Wang, L. M., Keegan, A. D., Paul, W. E., Heidaran, M. A., Gutkind, J. S., & Pierce, J. H. (1992). IL-4 activates a distinct signal transduction cascade from IL-3 in factor-dependent myeloid cells. *EMBO Journal*, Vol. 11, No. 13, (Dec, 1992), pp. 4899-4908.

- Wang, L. M., Michieli, P., Lie, W. R., Liu, F., Lee, C. C., Minty, A., Sun, X. J., Levine, A., White, M. F., & Pierce, J. H. (1995). The insulin receptor substrate-1-related 4PS substrate but not the interleukin-2R gamma chain is involved in interleukin-13-mediated signal transduction. *Blood*, Vol. 86, No. 11, (Dec 1, 1995), pp. 4218-4227.
- Wang, L. M., Myers, M. G., Jr., Sun, X. J., Aaronson, S. A., White, M., & Pierce, J. H. (1993). IRS-1: essential for insulin- and IL-4-stimulated mitogenesis in hematopoietic cells. *Science*, Vol. 261, No. 5128, (Sep 17, 1993), pp. 1591-1594.
- Wang, X. Y., Gelfanov, V., Sun, H. B., Tsai, S., & Yang, Y. C. (1999). Distinct actions of interleukin-9 and interleukin-4 on a hematopoietic stem cell line, EMLC1. *Experimental Hematology*, Vol. 27, No. 1, (Jan, 1999), pp. 139-146.
- Wang, Y., Malabarba, M. G., Nagy, Z. S., & Kirken, R. A. (2004). Interleukin 4 regulates phosphorylation of serine 756 in the transactivation domain of Stat6. Roles for multiple phosphorylation sites and Stat6 function. *Journal of Biological Chemistry*, Vol. 279, No. 24, (Jun 11, 2004), pp. 25196-25203.
- Warren, H. S., Kinnear, B. F., Phillips, J. H., & Lanier, L. L. (1995). Production of IL-5 by human NK cells and regulation of IL-5 secretion by IL-4, IL-10, and IL-12. *Journal of Immunology*, Vol. 154, No. 10, (May 15, 1995), pp. 5144-5152.
- Wei, H., Zhang, J., Xiao, W., Feng, J., Sun, R., & Tian, Z. (2005). Involvement of human natural killer cells in asthma pathogenesis: natural killer 2 cells in type 2 cytokine predominance. *The Journal of Allergy and Clinical Immunology*, Vol. 115, No. 4, (Apr, 2005), pp. 841-847.
- Welham, M. J., Duronio, V., Leslie, K. B., Bowtell, D., & Schrader, J. W. (1994). Multiple hemopoietins, with the exception of interleukin-4, induce modification of Shc and mSos1, but not their translocation. *Journal of Biological Chemistry*, Vol. 269, No. 33, (Aug 19, 1994), pp. 21165-21176.
- Welham, M. J., Duronio, V., Sanghera, J. S., Pelech, S. L., & Schrader, J. W. (1992). Multiple hemopoietic growth factors stimulate activation of mitogen-activated protein kinase family members. *Journal of Immunology*, Vol. 149, No. 5, (Sep 1, 1992), pp. 1683-1693.
- Welham, M. J., Duronio, V., & Schrader, J. W. (1994). Interleukin-4-dependent proliferation dissociates p44erk-1, p42erk-2, and p21ras activation from cell growth. *Journal of Biological Chemistry*, Vol. 269, No. 8, (Feb 25, 1994), pp. 5865-5873.
- Welham, M. J., Learmonth, L., Bone, H., & Schrader, J. W. (1995). Interleukin-13 signal transduction in lymphohemopoietic cells. Similarities and differences in signal transduction with interleukin-4 and insulin. *Journal of Biological Chemistry*, Vol. 270, No. 20, (May 19, 1995), pp. 12286-12296.
- Weller, C. L., Collington, S. J., Williams, T., & Lamb, J. R. (2011). Mast cells in health and disease. *Clin Sci (Lond)*, Vol. 120, No. 11, (Jun, 2011), pp. 473-484.
- Wen, L., Barber, D. F., Pao, W., Wong, F. S., Owen, M. J., & Hayday, A. (1998). Primary gamma delta cell clones can be defined phenotypically and functionally as Th1/Th2 cells and illustrate the association of CD4 with Th2 differentiation. *Journal of Immunology*, Vol. 160, No. 4, (Feb 15, 1998), pp. 1965-1974.
- Wenzel, S., Wilbraham, D., Fuller, R., Getz, E. B., & Longphre, M. (2007). Effect of an interleukin-4 variant on late phase asthmatic response to allergen challenge in asthmatic patients: results of two phase 2a studies. *Lancet*, Vol. 370, No. 9596, (Oct 20, 2007), pp. 1422-1431.

- Wenzel, S. E., Balzar, S., Ampleford, E., Hawkins, G. A., Busse, W. W., Calhoun, W. J., Castro, M., Chung, K. F., Erzurum, S., Gaston, B., Israel, E., Teague, W. G., Curran-Everett, D., Meyers, D. A., & Bleecker, E. R. (2007). IL4R alpha mutations are associated with asthma exacerbations and mast cell/IgE expression. *American Journal of Respiratory and Critical Care Medicine*, Vol. 175, No. 6, (Mar 15, 2007), pp. 570-576.
- Wery-Zennaro, S., Letourneur, M., David, M., Bertoglio, J., & Pierre, J. (1999). Binding of IL-4 to the IL-13Ralpha(1)/IL-4Ralpha receptor complex leads to STAT3 phosphorylation but not to its nuclear translocation. *FEBS Letters*, Vol. 464, No. 1-2, (Dec 24, 1999), pp. 91-96.
- Wery, S., Letourneur, M., Bertoglio, J., & Pierre, J. (1996). Interleukin-4 induces activation of mitogen-activated protein kinase and phosphorylation of shc in human keratinocytes. *Journal of Biological Chemistry*, Vol. 271, No. 15, (Apr 12, 1996), pp. 8529-8532.
- Wesch, D., Glatzel, A., & Kabelitz, D. (2001). Differentiation of resting human peripheral blood gamma delta T cells toward Th1- or Th2-phenotype. *Cellular Immunology*, Vol. 212, No. 2, (Sep 15, 2001), pp. 110-117.
- White, E. D., Andrews, R. P., & Hershey, G. K. (2001). Sulfhydryl-2 domain-containing protein tyrosine phosphatase-1 is not a negative regulator of interleukin-4 signaling in murine mast cells. *Journal of Leukocyte Biology*, Vol. 69, No. 5, (2001), pp. 825-830.
- White, M. F. (2002). IRS proteins and the common path to diabetes. *Am J Physiol Endocrinol Metab*, Vol. 283, No. 3, (Sep, 2002), pp. E413-422.
- Wills-Karp, M., Luyimbazi, J., Xu, X., Schofield, B., Neben, T. Y., Karp, C. L., & Donaldson, D. D. (1998). Interleukin-13: central mediator of allergic asthma. *Science*, Vol. 282, No. 5397, (Dec 18, 1998), pp. 2258-2261.
- Witthuhn, B. A., Silvennoinen, O., Miura, O., Lai, K. S., Cwik, C., Liu, E. T., & Ihle, J. N. (1994). Involvement of the Jak-3 Janus kinase in signalling by interleukins 2 and 4 in lymphoid and myeloid cells. *Nature*, Vol. 370, No. 6485, (Jul 14, 1994), pp. 153-157.
- Wolf, G., Trub, T., Ottinger, E., Groninga, L., Lynch, A., White, M. F., Miyazaki, M., Lee, J., & Shoelson, S. E. (1995). PTB domains of IRS-1 and Shc have distinct but overlapping binding specificities. *Journal of Biological Chemistry*, Vol. 270, No. 46, (Nov 17, 1995), pp. 27407-27410.
- Wu, L., Bijian, K., & Shen, S. H. (2009). CD45 recruits adapter protein DOK-1 and negatively regulates JAK-STAT signaling in hematopoietic cells. *Molecular Immunology*, Vol. 46, No. 11-12, (Jul, 2009), pp. 2167-2177.
- Wurster, A. L., Rodgers, V. L., White, M. F., Rothstein, T. L., & Grusby, M. J. (2002). Interleukin-4-mediated protection of primary B cells from apoptosis through Stat6-dependent up-regulation of Bcl-xL. *Journal of Biological Chemistry*, Vol. 277, No. 30, (Jul 26, 2002), pp. 27169-27175.
- Wymann, M. P., & Pirola, L. (1998). Structure and function of phosphoinositide 3-kinases. *Biochimica et Biophysica Acta*, Vol. 1436, No. 1-2, (Dec 8, 1998), pp. 127-150.
- Yamashita, M., Kimura, M., Kubo, M., Shimizu, C., Tada, T., Perlmutter, R. M., & Nakayama, T. (1999). T cell antigen receptor-mediated activation of the Ras/mitogen-activated protein kinase pathway controls interleukin 4 receptor

- function and type-2 helper T cell differentiation. *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 96, No. 3, (Feb 2, 1999), pp. 1024-1029.
- Yi, T. L., Cleveland, J. L., & Ihle, J. N. (1992). Protein tyrosine phosphatase containing SH2 domains: characterization, preferential expression in hematopoietic cells, and localization to human chromosome 12p12-p13. *Molecular and Cellular Biology*, Vol. 12, No. 2, (Feb, 1992), pp. 836-846.
- Yoshikawa, M., Nakajima, T., Tsukidate, T., Matsumoto, K., Iida, M., Otori, N., Haruna, S., Moriyama, H., & Saito, H. (2003). TNF-alpha and IL-4 regulate expression of IL-13 receptor alpha2 on human fibroblasts. *Biochemical and Biophysical Research Communications*, Vol. 312, No. 4, (Dec 26, 2003), pp. 1248-1255.
- Yoshimura, A., Mori, H., Ohishi, M., Aki, D., & Hanada, T. (2003). Negative regulation of cytokine signaling influences inflammation. *Current Opinion in Immunology*, Vol. 15, No. 6, (Dec, 2003), pp. 704-708.
- Zamorano, J., & Keegan, A. D. (1998). Regulation of apoptosis by tyrosine-containing domains of IL-4R alpha: Y497 and Y713, but not the STAT6-docking tyrosines, signal protection from apoptosis. *Journal of Immunology*, Vol. 161, No. 2, (Jul 15, 1998), pp. 859-867.
- Zhang, J. L., Buehner, M., & Sebald, W. (2002). Functional epitope of common gamma chain for interleukin-4 binding. *European Journal of Biochemistry*, Vol. 269, No. 5, (Mar, 2002), pp. 1490-1499.
- Zheng, T., Liu, W., Oh, S. Y., Zhu, Z., Hu, B., Homer, R. J., Cohn, L., Grusby, M. J., & Elias, J. A. (2008). IL-13 receptor alpha2 selectively inhibits IL-13-induced responses in the murine lung. *Journal of Immunology*, Vol. 180, No. 1, (Jan 1, 2008), pp. 522-529.
- Zheng, T., Zhu, Z., Liu, W., Lee, C. G., Chen, Q., Homer, R. J., & Elias, J. A. (2003). Cytokine regulation of IL-13Ralpha2 and IL-13Ralpha1 in vivo and in vitro. *Journal of Allergy and Clinical Immunology*, Vol. 111, No. 4, (Apr, 2003), pp. 720-728.
- Zhou, M. M., Huang, B., Olejniczak, E. T., Meadows, R. P., Shuker, S. B., Miyazaki, M., Trub, T., Shoelson, S. E., & Fesik, S. W. (1996). Structural basis for IL-4 receptor phosphopeptide recognition by the IRS-1 PTB domain. *Nature Structural Biology*, Vol. 3, No. 4, (Apr, 1996), pp. 388-393.
- Zhu, J., & Paul, W. E. (2008). CD4 T cells: fates, functions, and faults. *Blood*, Vol. 112, No. 5, (Sep 1, 2008), pp. 1557-1569.
- Zhu, J., Yamane, H., Cote-Sierra, J., Guo, L., & Paul, W. E. (2006). GATA-3 promotes Th2 responses through three different mechanisms: induction of Th2 cytokine production, selective growth of Th2 cells and inhibition of Th1 cell-specific factors. *Cell Research*, Vol. 16, No. 1, (Jan, 2006), pp. 3-10.
- Zhu, Z., Homer, R. J., Wang, Z., Chen, Q., Geba, G. P., Wang, J., Zhang, Y., & Elias, J. A. (1999). Pulmonary expression of interleukin-13 causes inflammation, mucus hypersecretion, subepithelial fibrosis, physiologic abnormalities, and eotaxin production. *Journal of Clinical Investigation*, Vol. 103, No. 6, (Mar, 1999), pp. 779-788.
- Zurawski, S. M., Vega, F., Jr., Huyghe, B., & Zurawski, G. (1993). Receptors for interleukin-13 and interleukin-4 are complex and share a novel component that functions in signal transduction. *EMBO Journal*, Vol. 12, No. 7, (Jul, 1993), pp. 2663-2670.



Allergic Diseases - Highlights in the Clinic, Mechanisms and Treatment

Edited by Prof. Celso Pereira

ISBN 978-953-51-0227-4

Hard cover, 554 pages

Publisher InTech

Published online 14, March, 2012

Published in print edition March, 2012

The present Edition "Allergic diseases - highlights in the clinic, mechanisms and treatment" aims to present some recent aspects related to one of the most prevalent daily clinical expression disease. The effort of a group of outstanding experts from many countries reflects a set of scientific studies very promising for a better clinical care and also to the treatment and control of the allergy. This book provides a valuable reference text in several topics of the clinical allergy and basic issues related to the immune system response. The inflammatory reaction understanding in allergic disease is clearly evidenced, as well as new strategies for further researches.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Nicola M. Heller, Preeta Dasgupta, Nicolas J. Dorsey, Svetlana P. Chapoval and Achsah D. Keegan (2012). The Type I and Type II Receptor Complexes for IL-4 and IL-13 Differentially Regulate Allergic Lung Inflammation, Allergic Diseases - Highlights in the Clinic, Mechanisms and Treatment, Prof. Celso Pereira (Ed.), ISBN: 978-953-51-0227-4, InTech, Available from: <http://www.intechopen.com/books/allergic-diseases-highlights-in-the-clinic-mechanisms-and-treatment/the-type-i-and-type-ii-receptor-complexes-for-il-4-and-il-13-differentially-regulate-allergic-lung-i>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen