

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Reassessing the Value of the Exercise Electrocardiogram in the Diagnosis of Stable Chest Pain

Peter Bourdillon
*Imperial College,
UK*

1. Introduction

“Do not use exercise ECG to diagnose or exclude angina for people without known coronary artery disease” - a quote from the summary (Cooper et al, 2010) of the guidelines of the United Kingdom’s National Institute for Health and Clinical Excellence (NICE) on the diagnosis of discomfort of suspected cardiac origin (NICE, 2010). Amongst other things, NICE recommends that the probability of coronary artery disease (pCAD) is determined from the person’s symptoms and risk factors, and that the subsequent management depends on whether the pCAD is less than 10%, 10 to 30%, 30 to 60%, 60 to 90% or greater than 90%: if the pCAD is less than 10% or greater than 90% further testing is not required; if the pCAD is between 60 and 90%, the patient is a candidate for coronary arteriography; if the pCAD is between 10 and 60% non-invasive testing is indicated.

Ever since a meta-analysis (Gianrossi et al., 1989) of the exercise electrocardiogram (ECG), in which the sensitivity was assessed as being 68% and the specificity as 77%, the test’s role in the diagnosis of coronary artery disease has been questioned. However, in 1979 Diamond & Forrester demonstrated that the greater the ST segment shift on exercise ECG, the greater the post-exercise probability of coronary artery disease (pCAD) whatever the pre-exercise pCAD. Furthermore, the Duke Treadmill Score (Mark et al., 1991) demonstrates that the greater the ST segment depression the worse the prognosis.

The purpose of this study is to show, using the data obtained from a West London population and NICE’s methodology, the different impact the exercise ECG has on subsequent management, if Diamond & Forrester’s method of analysing exercise ECG data is used instead of NICE’s.

2. Methods

Patients with a history of chest pain referred direct from primary care to the rapid access chest pain clinic of the Hammersmith Hospital, London, between January 1999 and March 2010 were studied. Each person was asked to complete a questionnaire (Table 1) which combines questions from two previously published studies (Joswig et al., 1985; Pryor et al., 1993).

		Yes	No
1	Have you had a coronary bypass operation or a coronary angioplasty?		
2	Have you ever been told that you have had a heart attack?		
3	Have you smoked 10 or more cigarettes a day in the last 5 years?		
4	Have you ever had diabetes?		
5	Has your blood cholesterol ever been more than 6.2 or have you ever been told that your cholesterol is high?		
6	Do you get <i>breathless</i> with the pain or discomfort?		
7	Do you feel <i>sick</i> with the pain or discomfort?		
8	Is the pain or discomfort <i>burning</i> in nature?		
9	Is the pain or discomfort <i>prickling</i> in nature?		
10	Do you feel the pain or discomfort in the <i>middle</i> of the chest?		
11	Does the pain or discomfort go into <i>both</i> arms?		
12	Do you feel the pain or discomfort <i>deep</i> of the chest?		
13	Is the pain or discomfort made <i>worse</i> by moving your arms or twisting your body?		
14	Is the pain or discomfort brought on by <i>exertion</i> such as climbing stairs or walking uphill?		
15	Is the pain or discomfort brought on by <i>sexual intercourse</i> ?		
16	How long does the pain or discomfort <i>usually</i> last if you stop doing whatever started it?		
	less than 30 seconds		
	between 30 seconds and 10 minutes		
	more than 10 minutes		
		Yes	No
17	Do you use glyceryl trinitrate (the spray or the small white tablet under the tongue used to treat chest pain)?		
18	How long does the pain or discomfort <i>usually</i> last if you use glyceryl trinitrate?		
	less than 30 seconds		
	between 30 seconds and 10 minutes		
	more than 10 minutes		

Table 1. The questionnaire each patient was asked to complete.

The questionnaire was translated into 5 languages commonly used in the local community, viz. Farsi, Hindi, Polish, Punjabi and Urdu. The raw data from the questionnaire were stored in a bespoke database. For a symptomatic diagnosis of Typical Angina (chest pain symptom score 3 points) the following criteria had to be met: a) 'Yes' to both question 10 and question 12 and b) 'Yes' to either question 14 or question 15 and c) 'between 30 seconds and 10 minutes' to question 16 or 'between 30 seconds and 10 minutes' to question 18 if the answer to question 17 was 'Yes'. Atypical Angina was diagnosed if any 2 of criteria a), b) and c) were met (chest pain symptom score 2 points); otherwise Non-Anginal Chest Pain was diagnosed (chest pain symptom score either 1 or 0 points).

From the questionnaire data, pCAD was calculated (Pryor et al., 1993) for routine clinical use (pCAD_DUKE). To assess the impact of NICE's guidelines, NICE's modification of pCAD_DUKE that omits the history of myocardial infarction and omits ECG data, was used to recalculate pCAD as pCAD_NICE. The doctor or nurse assessing the patient had the pCAD_DUKE available when deciding whether or not to exercise the patient.

Treadmill exercise ECGs were performed following the Bruce protocol (Bruce et al., 1949). The Mason-Likar electrode positions (Mason & Likar, 1966) were used. The maximum ST segment shift 60-80 ms after the end of the QRS was noted; it was considered less than 50 μV if the ST segment was upsloping and the end of the QRS depressed. ST segment shifts due to rhythm disturbances were ignored. From the exercise ECG the post-exercise pCAD was calculated using Bayes' Theorem of conditional probability

$$\text{post-exercise pCAD} = [100 * \text{pre-exercise pCAD} * \text{sensitivity}] / [(\text{pre-exercise pCAD} * \text{sensitivity}) + (100 - \text{pre-exercise pCAD}) * (100 - \text{specificity})]$$

and using the sensitivities and specificities in Table 2 for different levels of ST segment shift (Diamond & Forrester, 1979). Also calculated were

$$\text{the positive likelihood ratio} = \text{sensitivity} / (100 - \text{specificity}) \text{ and}$$

$$\text{the negative likelihood ratio} = (100 - \text{sensitivity}) / \text{specificity}$$

ST segment shift (μV)	Sensitivity (%)	Specificity (%)	Positive Likelihood Ratio	Negative Likelihood Ratio
0 - 50	14.3	37.5	0.23	2.29
51 - 100	20.8	77.3	0.92	1.02
101 - 150	23.3	89.0	2.12	0.86
151 - 200	8.8	97.9	4.19	0.93
201 - 250	13.3	98.8	11.08	0.88
> 250	19.5	99.5	39.00	0.81

Table 2. The 6 combinations of sensitivity and specificity corresponding to the maximum ST segment shift on exercise (Diamond & Forrester, 1979). NICE's assumption of a sensitivity of 67% and a specificity of 69% for the single threshold point of 100 μV has a positive likelihood ratio of 2.16 and a negative likelihood ratio of 0.48

The post-exercise pCAD was also calculated using NICE's assumption of sensitivity 67% and specificity 69% for a ST segment shift threshold of 100 μV. NICE's sensitivity and specificity were obtained from table 14 of Mowatt et al., 2008, in which the sensitivities and specificities obtained from Kuntz et al., 1999, and from Mowatt et al., 2004, were averaged.

3. Statistics

Multivariate linear regression analysis (StatsDirect version 2.7.8; StatsDirect Ltd, Altrincham, WA14 4QA, UK) was used to perform analysis of variance (ANOVA) to determine the impact of age, sex and race on the likelihood of a patient being referred for an exercise ECG. Dummy variables were generated for sex and race. Multivariate linear regression analysis was also used to determine the impact of age, sex, race on chest pain

symptom score and on exercise workload. Chi-square analysis was used to compare the numbers of exercised patients in each pCAD category with the numbers not exercised.

4. Results

All 7739 patients referred to the rapid access chest pain clinic completed the questionnaire. Following clinical assessment 5157 of the 7739 (66.6%) were exercised. 359 of 2582 (13.9%) who were not exercised had either left ventricular hypertrophy or an intraventricular conduction defect on 12 lead ECG, while 513 of 5157 (9.9%) who were exercised had either left ventricular hypertrophy or an intraventricular conduction defect ($\chi^2 = 26.9$ P < 0.0001).

Age	Black	South Asian	Miscellaneous	Oriental	White	Females Total
10 - 19	0 (1)	1.5 (2)	0 (2)		1 (2)	0.71 (7)
20 - 29	1.45 (11)	1.88 (9)	0.5 (6)	2.5 (2)	1.41 (29)	1.43 (57)
30 - 39	1.41 (36)	1.16 (12)	1.5 (16)	0.5 (4)	1.10 (58)	1.23 (126)
40 - 49	1.17 (47)	1.47 (46)	1.29 (24)	1.38 (13)	1.32 (129)	1.32 (259)
50 - 59	1.15 (51)	1.36 (66)	1.5 (18)	1.25 (8)	1.26 (178)	1.28 (321)
60 - 69	1.53 (43)	1.55 (58)	1.55 (29)	1.5 (8)	1.24 (161)	1.38 (299)
70 - 79	1.42 (33)	1.48 (35)	1.22 (9)	1.33 (6)	1.36 (176)	1.38 (259)
80 - 89	1.28 (7)	1.33 (6)	1 (2)	2 (1)	1.4 (90)	1.38 (106)
90 - 99		2 (1)			1.14 (7)	1.25 (8)
Total	1.32 (229)	1.46 (235)	1.34 (106)	1.35 (42)	1.3 (830)	1.33 (1442)

Age	Black	South Asian	Miscellaneous	Oriental	White	Males Total
10-19		0 (1)	1 (1)		0.66 (6)	0.62 (8)
20-29	0.25 (8)	0.9 (10)	1.66 (6)	2 (1)	1.06 (32)	1 (57)
30-39	0.9 (20)	1.34 (29)	1.06 (16)	0.5 (4)	0.87 (94)	0.96 (163)
40-49	0.92 (28)	1.12 (48)	1.27 (33)	1.2 (5)	1.04 (107)	1.08 (221)
50-59	1.12 (8)	1.53 (39)	1.48 (27)	0.8 (5)	1.19 (129)	1.28 (208)
60-69	1.30 (23)	1.47 (40)	1.41 (12)	1.33 (3)	1.37 (127)	1.39 (205)
70-79	1.07 (28)	1.53 (32)	1 (8)	1 (1)	1.35 (116)	1.32 (185)
80-89	1.57 (7)	1 (10)	2 (3)		1.41 (72)	1.40 (92)
90-99					2 (1)	2 (1)
Total	1.03 (122)	1.33 (209)	1.33 (106)	1 (19)	1.20 (684)	1.21 (1140)

Table 3. The average chest pain symptom score of females (above) and males (below) not exercised by race and decade of age. Typical angina = 3, atypical angina = 2, non-anginal chest pain = 1 or 0. The numbers in brackets are the numbers of patients.

Table 3 shows the distribution of age, sex and race of those not exercised, while table 4 shows the distribution of age, sex and race of those exercised. The distributions are similar. Black men had similar chest pain symptom scores to White men and Black men to Black women, but the South Asian, Miscellaneous and Oriental women had higher chest pain

Age	Black	South Asian	Miscellaneous	Oriental	White	Females Total
20 - 29	1.5 (6) 9.83	1.5 (6) 10.3	1.16 (6) 10.3		1.27 (18) 12.8	1.33 (36) 11.5
30 - 39	1.52 (17) 9.05	1.53 (28) 8.07	1.54 (11) 7.72	2 (4) 9.25	1.47 (76) 11.2	1.51 (136) 9.99
40 - 49	1.72 (84) 8.10	1.81 (103) 8.14	1.79 (64) 8.18	1.41 (17) 10.0	1.53 (236) 9.48	1.65 (504) 8.83
50 - 59	1.41 (87) 7.90	1.6 (135) 7.8	1.75 (68) 8.07	1.89 (19) 8.73	1.45 (373) 8.49	1.52 (682) 8.24
60 - 69	1.44 (90) 6.86	1.70 (98) 6.71	1.56 (41) 7.48	1.63 (19) 7.52	1.46 (363) 7.69	1.50 (611) 7.39
70 - 79	1.46 (43) 6.16	1.36 (25) 5.84	1.73 (15) 6.73	1 (1) 4	1.35 (234) 6.43	1.38 (318) 6.35
80 - 89	2 (2) 2.5	1 (1) 4			1.42 (42) 4.85	1.44 (45) 4.73
90 - 99					1 (1) 7	1 (1) 7
Total	1.51 (329) 7.50	1.65 (396) 7.54	1.69 (205) 7.94	1.66 (60) 8.68	1.45 (1343) 8.19	1.52 (2333) 7.97

Age	Black	South Asian	Miscellaneous	Oriental	White	Males Total
10-19					0 (1) 16	0 (1) 16
20-29	1.25 (8) 14.6	0.83 (12) 13.0	1.12 (8) 13.6		1 (26) 13.2	1.01 (54) 13.4
30-39	1.6 (30) 12.3	1.22 (84) 11.7	1.20 (43) 12.5	1.5 (6) 12.6	1.17 (173) 13.0	1.23 (336) 12.5
40-49	1.38 (68) 11.3	1.13 (143) 10.9	1.46 (84) 11.0	0.90 (11) 11.2	1.22 (396) 11.6	1.24 (702) 11.4
50-59	1.37 (43) 9.62	1.28 (146) 9.60	1.32 (92) 10.0	0.91 (12) 11.3	1.19 (472) 10.0	1.23 (765) 9.94
60-69	1.28 (53) 8.05	1.50 (111) 7.54	1.30 (43) 8.93	2 (3) 10	1.21 (419) 8.65	1.28 (629) 8.43
70-79	1.38 (42) 6.42	1.37 (51) 7.09	1.53 (13) 8.07	0.75 (4) 8.75	1.36 (188) 6.96	1.36 (298) 6.98
80-89	1.5 (4) 6.75	1 (2) 5.5	2 (2) 5.5		1.38 (31) 5.51	1.41 (39) 5.64
Total	1.38 (248) 9.66	1.27 (549) 9.69	1.35 (285) 10.5	1.08 (36) 11.1	1.22 (1706) 10.0	1.25 (2824) 9.98

Table 4. The average chest pain symptom score of females (above) and males (below) exercised and average exercise workload achieved (metabolic equivalents) by race and decade of age. Typical angina = 3, atypical angina = 2, non-anginal chest pain = 1 or 0. The numbers in brackets are the numbers of patients and the third number in a cell is the average exercise workload

symptom scores than the Black and White women. Analysis of Variance (ANOVA) showed that a higher chest pain symptom score was found in females and in those who were South Asians or in those having a Miscellaneous race (compared to being White). However, the descriptors explained only 2.4% of the variance in chest pain symptom score. ANOVA also showed that the younger the age, being male and a higher symptom score all made being exercised more likely. The descriptors explained 1.6% of the variance in the likelihood of being exercised. Table 4 also shows the workload achieved on exercise. For the 5157 patients who were exercised, younger males with low symptom scores were those who achieved the highest workload. Whites and Orientals were the races that achieved the highest workload and South Asians the lowest. The descriptors explained 36.4% of the variance.

Table 5 shows a comparison of the pre-test pCADs of the patients who were not exercised with those who were exercised. Chi-square analysis showed a significant difference in the proportions exercised and the proportions not exercised ($\chi^2=120.4$ with 4 degrees of freedom; $P < 0.0001$). Those with a pCAD below 10% were less likely to be exercised than those with pCAD between 10 and 90%.

pre-test pCAD	not exercised	exercised	total
< 10%	853	1243 (59.3%)	2096
10 – 30%	697	1676 (70.6%)	2373
30 – 60%	470	1234 (72.4%)	1704
60 – 90%	411	834 (67.0%)	1245
> 90%	151	170 (53.0%)	321
Total	2582	5157 (66.6%)	7739

pCAD = probability of coronary artery disease

Table 5. The pre-test pCAD of the patients according to whether or not they were exercised. The “exercised” column includes the percentage (in brackets) of the patients for a given pre-test pCAD who were exercised. Patients with a pre-test pCAD of less than 10% were less likely to be exercised than those with pCAD between 10 and 90%

Table 6 shows the change in pCAD categories following exercise. In the top part of Table 6 the numbers in each cell show how the pCAD changes when 6 combinations of sensitivity and specificity are used to calculate the post-exercise pCAD. 3296 of 5157 (63.9%) changed category: 535 of the 3296 increased pCAD while 2761 reduced pCAD. Had only the 2910 patients with pre-test pCAD between 10 and 60% been exercised, 1383 (47.5%) would have had CAD ruled out and 48 (1.6%) CAD ruled in. Another 170 (5.8%) would have become candidates for coronary arteriography. Figure 1 illustrates the change in pCAD. The bottom part of Table 6 shows the change in pCAD categories following exercise when NICE’s assumption of one combination of sensitivity (67%) and specificity (69%) is used to calculate the post-exercise pCAD. 2271 of 5157 (44.0%) changed category. Of those with a pre-exercise pCAD of between 10% and 60%, 793 of 2910 (27.3%) had CAD ruled out and 128 (4.4%) became candidates for coronary arteriography. Figure 2 illustrates the change in pCAD.

Post-exercise pCAD% 6 Sensitivities and Specificities						
Pre-exercise pCAD%	< 10	10 - 30	30 - 60	60 - 90	> 90	Total
< 10	1146	73	21	3		1243
10 - 30	1287	225	119	38	7	1676
30 - 60	96	795	170	132	41	1234
60 - 90		113	393	207	121	834
> 90				77	93	170
Total	2529	1206	703	457	262	5157

Post-exercise pCAD% Sensitivity 67% Specificity 69%						
Pre-exercise pCAD%	< 10	10 - 30	30 - 60	60 - 90	> 90	Total
< 10	1162	81				1243
10 - 30	793	760	123			1676
30 - 60		664	442	128		1234
60 - 90			371	417	46	834
> 90				65	105	170
Total	1955	1505	936	610	151	5157

pCAD = probability of coronary artery disease

Table 6. The top part of the table shows how the numbers of patients in each category changed following exercise using the 6 combinations of sensitivity and specificity as in table 2. The bottom part of the table shows how the numbers of patients in each category changed following exercise using a single combination of sensitivity (67%) and specificity (69%).




Fig. 1. The probability (%) of CAD after exercise plotted against the probability of CAD before exercise when 6 combinations (5 thresholds of ST shift) of sensitivity and specificity were used. The graph shows the discrimination that can be obtained by using several thresholds of ST shift and how CAD can be ruled out if the ST shift is less than 50 μ V. The units of the thresholds in the legend are μ V.




Fig. 2. The probability (%) of CAD after exercise plotted against the probability of CAD before exercise using a single threshold of 100 μ V.

5. Discussion

The purpose of this study has been to try and demonstrate the superior diagnostic value of taking the amount of ST segment shift into account compared with using a single threshold point when interpreting the exercise ECG. The publication of the method used by NICE to obtain the pre-exercise probability of CAD (pCAD) made this comparison and the testing of NICE's recommendations feasible. NICE recommends firstly, that symptoms and risk factors should be used to derive pCAD and secondly, that the subsequent management depends on whether the pCAD is less than 10%, between 10 and 30%, between 30 and 60%, between 60 and 90%, or greater than 90%. Those with a pCAD of less than 10% have CAD ruled out and need no further investigation for CAD. Those with a pCAD of more than 90% have CAD ruled in. Those with a pCAD between 60 and 90% should be considered for coronary arteriography. NICE recommends that those with a pCAD between 10 and 60% need further investigation, but not exercise ECG.

In this study 5157 of 7739 consecutive patients referred to the rapid access chest pain clinic had an exercise ECG. Table 6 shows that using NICE's methodology and a sensitivity of 67% and a specificity of 69% for an ST segment shift of 100 μ V, the exercise ECG rules out CAD in 793 (27.3%) of the 2910 patients with a pre-exercise pCAD between 10 and 60%, while another 128 (4.4%) become candidates for coronary arteriography. However, modifying the calculation of the post-exercise pCAD to account for increasing ST segment shift during exercise has a dramatic effect: of the 2910 patients who had a pre-test pCAD between 10 and 60% and who were exercised, 1383 (47.5%) would have had CAD ruled out and 48 (1.6%) CAD ruled in. Another 170 (5.8%) would have become candidates for coronary arteriography. The exercise ECG would therefore have defined the subsequent management of more than half the patients that had a pre-test pCAD between 10 and 60%.

The distributions by age, sex and race of patients referred for exercise ECG were similar to those of the whole cohort. Women were more likely to have angina than men. As recommended by NICE, patients referred for exercise testing were less likely to have a

pCAD below 10% than those with a pCAD between 10 and 30%. The workload achieved during exercise was greater in men than in women and declined with age.

Table 2 reproduces part of table 4 of Diamond & Forrester, 1979 and adds corresponding positive and negative likelihood ratios. Diamond & Forrester reviewed 31 published papers to derive the table. From these papers they were able to derive a sensitivity and a specificity for each of the 5 thresholds. The assumption is that these are mutually exclusive events, each having their own receiver operating characteristic curve. Table 2 shows that increasing the ST segment shift above 100 μV steadily increases the likelihood of CAD, while an ST segment shift of less than 50 μV reduces the likelihood of CAD.

Table 7 shows the effect of varying the combination of sensitivity and specificity on a single ST segment shift with a threshold of 100 μV . NICE assumed for calcium scoring a sensitivity

		Post-exercise pCAD% Sensitivity 89% Specificity 43%					
Pre-exercise pCAD%	< 10	10 - 30	30 - 60	60 - 90	> 90	Total	
< 10	1199	44				1243	
10 - 30	1473	137	66			1676	
30 - 60	21	1001	144	68		1234	
60 - 90		66	511	233	24	834	
> 90				90	80	170	
Total	2693	1248	721	391	104	5157	

		Post-exercise pCAD% Sensitivity 86% Specificity 64%					
Pre-exercise pCAD%	< 10	10 - 30	30 - 60	60 - 90	> 90	Total	
< 10	1157	86				1243	
10 - 30	1473	65	138			1676	
30 - 60	151	871	73	139		1234	
60 - 90		151	459	171	53	834	
> 90				93	77	170	
Total	2781	1173	670	403	130	5157	

		Post-exercise pCAD% Sensitivity 68% Specificity 77%					
Pre-exercise pCAD%	< 10	10 - 30	30 - 60	60 - 90	> 90	Total	
< 10	1144	99				1243	
10 - 30	987	522	167			1676	
30 - 60		764	280	190		1234	
60 - 90			407	351	76	834	
> 90				70	100	170	
Total	2131	1385	854	611	176	5157	

Table 7. Three parts of the table show the effect on the data of using different combinations of sensitivity and specificity. At the top is that for a sensitivity of 89% and a specificity of 43% (NICE's combination for Calcium Scoring). In the middle is that for a sensitivity of 86% and a specificity of 64% (NICE's combination for Myocardial Perfusion Scanning). Below is that for a sensitivity of 68% and a specificity of 77% (the combination from the meta-analysis of Gianrossi et al., 1989).

of 89% and a specificity of 43%; this is the combination in the top part. NICE assumed for myocardial perfusion scanning with single photon emission computed tomography a sensitivity of 86% and a specificity of 64%; this is the combination in the middle part. From their meta-analysis Gianrossi et al., 1989 obtained a sensitivity of 68% and a specificity of 77%; this is the combination in the bottom part. For ruling out CAD when the pre-exercise pCAD is between 10 and 60%, table 7 and the bottom part of table 6 indicate that NICE's combinations of sensitivity and specificity used with calcium scoring (51.3% ruled out) and with myocardial perfusion scanning (55.8% ruled out) are both superior to NICE's combination used with exercise ECG (27.3% ruled out) and to the combination used by Gianrossi et al (33.9% ruled out). However, NICE recommends using myocardial perfusion scanning only if the pre-exercise pCAD is between 30 and 60%. Consequently, only 151 of 1234 (12.2%) would have had CAD ruled out. Only the combination of 6 sensitivities and specificities resulted in any of those with a pre-test pCAD between 10 and 60% having CAD ruled in.

As NICE, 2010, assumed costs of \$100 (£66), \$450 (£293) and \$160 (£103) for exercise ECG, for myocardial perfusion scanning and for calcium scoring respectively, the explanation for the recommendation not to use the exercise ECG may be partly due to the assumptions about the proportions of indeterminate tests. NICE assumed 24% of exercise ECGs were indeterminate, 6% of myocardial perfusion scans were indeterminate and 2% of calcium scores were indeterminate. The 24% was NICE's modification of the unsubstantiated 30% in Kuntz et al., 1999. In the present study no exercise ECG was indeterminate, although it is accepted that interpretation of ST shift in the presence of left ventricular hypertrophy or of intraventricular conduction defect is unreliable. Ten per cent of the ECGs of the patients exercised showed either left ventricular hypertrophy or an intraventricular conduction defect.

6. Study limitations

Firstly, the sensitivities and specificities of each of the 5 thresholds of ST shift used were obtained by a review of the literature (Diamond & Forrester, 1979) rather than by a formal meta-analysis. Secondly, the patients were not asked about their race. The members of the ECG team assessed a person's race. Even though there were only 5 different races from which to choose, it is likely that misclassification occurred. Thirdly, being in paper format, the answers to the questionnaire were subject to transcription errors when entered into the database. Fourthly, measurement of the maximum ST segment shift on exercise was made by the member of the ECG team supervising the test. Until June 2006 all exercise tests were checked by the author. After June 2006 a weekly tutorial session with the author enabled the ECG team to discuss ST segment shifts that were difficult to assess.

7. Conclusions

It is concluded firstly, that the exercise ECG does indeed have limited diagnostic value if a single threshold is used for the diagnosis of CAD; secondly, that the exercise ECG has considerable diagnostic value if several thresholds are used; and thirdly, that NICE's recommendation "Do not use exercise ECG to diagnose or exclude angina for people without known coronary artery disease" seems to be partly the result of an unjustified assumption about the proportion of indeterminate exercise ECGs.

8. Acknowledgements

I thank the members of the ECG Team who entered the data into the bespoke database. Some of the data presented has previously been published as a letter (Bourdillon, 2010).

9. References


- Bourdillon, P. (2010). Exercise ECG useful in finding coronary artery disease. *British Medical Journal* Vol. 340, c1971, ISSN 0959-8138
- Bruce, R. Lovejoy, F Jr. Pearson, R. Yu, P. Brothers, G. & Velasquez, T. (1949). Normal respiratory and circulatory pathways of adaptation in exercise. *Journal of Clinical Investigation*, Vol. 28, No. 6, pp. 1423-1430, ISSN 0021-9738
- Cooper, A. Timmis, A. & Skinner J. (2010). Assessment of recent onset chest pain or discomfort of suspected cardiac origin: summary of NICE guidance. *British Medical Journal*, Vol. 340, No. 7749, c1118, ISSN 0959-8138
- Diamond, G. & Forrester, J. (1979). Analysis of probability as an aid in the clinical diagnosis of coronary-artery disease. *New England Journal of Medicine*, Vol. 300, No. 24, pp. 1350-8, ISSN 0028-4793
- Gianrossi, R. Detrano, R. Mullvihill, D. Lehmann, K. Dubach, P. Colombo, A. McArthur, D. & Froelicher, V. (1989). Exercise-induced ST depression in the diagnosis of coronary artery disease. A meta-analysis. *Circulation*, Vol. 80, No. 1, pp. 87-98, ISSN 0009-7322
- Joswig, B. Glover, M. Nelson, D. Handler, J. & Henderson, J. (1985). Analysis of historical variables, risk factors and the resting electrocardiogram as an aid in the clinical diagnosis of recurrent chest pain. *Computers in Biology and Medicine*, Vol. 15, No. 2, pp. 71-80, ISSN: 0010-4825
- Kuntz, K. Fleischmann, K. Hunink, M. & Douglas, P. (1999). Cost-effectiveness of diagnostic strategies for patients with chest pain. *Annals of Internal Medicine*, Vol. 130, No. 9, pp. 709-718, ISSN 0003-4819
- Mark, D. Shaw, L. Harrell, F Jr. Hlatky, M. Lee, K. Bengtson, F. McCants, C. Califf, R. & Pryor D. (1991). Prognostic value of a treadmill exercise score in outpatients with suspected coronary artery disease. *New England Journal of Medicine*, Vol. 325, No.12, pp. 849-853, ISSN 0028-4793
- Mason, R. & Likar, I. (1966). A new system of multiple-lead exercise electrocardiography. *American Heart Journal* Vol. 71, No. 2, pp. 196-205, ISSN 0002-8703
- Mowatt, G. Vale, L. Brazzelli, M. Hernandez, R. Murray, A. Scott, N. Fraser, C. McKenzie, L. Gemmell, H. Hillis, G. & Metcalfe, M. (2004). Systematic review of the effectiveness and cost-effectiveness and economic evaluation of myocardial perfusion scintigraphy for the diagnosis and management of angina and myocardial infarction. *Health Technology Assessment*, Vol. 8, No. 30, pp. 1-207, ISSN 1366-5278
- Mowatt, G. Cummins, E. Waugh, N. Walker, S. Cook, J. Jia, X. Hillis, G. & Fraser, C. (2008). Systematic review of the clinical effectiveness and cost-effectiveness of 64-slice or higher computed tomography angiography as an alternative to invasive coronary angiography in the investigation of coronary artery disease. *Health Technology Assessment*, Vol. 12, No. 17, pp. 1-143, ISSN 1366-5278

National Institute for Health and Clinical Excellence. (10 June 2010). *Chest pain of recent onset: assessment and diagnosis of recent onset chest pain or discomfort of suspected cardiac origin (clinical guideline 95)*. Retrieved from www.nice.org.uk/guidance/cg95

Pryor, D. Shaw, L. McCants, C Lee, K. Mark, D. Harrell, F Jr. Muhlbaier, L. & Califf, R. (1993) Value of the history and physical in identifying patients at increased risk for coronary artery disease. *Annals of Internal Medicine*, Vol. 118, No. 2, pp. 81-90, ISSN 0003-4819

IntechOpen

IntechOpen



Coronary Artery Disease - Current Concepts in Epidemiology, Pathophysiology, Diagnostics and Treatment

Edited by Dr. David Gaze

ISBN 978-953-51-0262-5

Hard cover, 272 pages

Publisher InTech

Published online 16, March, 2012

Published in print edition March, 2012

Cardiovascular disease is ranked as the leading cause of death world wide, responsible for 17.1 million deaths globally each year. Such numbers are often difficult to comprehend. Heart disease kills one person every 34 seconds in the USA alone. Although the leading killer, the incidence of cardiovascular disease has declined in recent years due to a better understanding of the pathology, implementation of lipid lowering therapy new drug regimens including low molecular weight heparin and antiplatelet drugs such as glycoprotein IIb/IIIa receptor inhibitors and acute surgical intervention. The disease burden has a great financial impact on global healthcare systems and major economic consequences for world economies. This text aims to deliver the current understanding of coronary artery disease and is split into three main sections: 1. Epidemiology and pathophysiology of coronary artery disease 2. Coronary artery disease diagnostics and 3. Treatment regimens for coronary artery disease

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Peter Bourdillon (2012). Reassessing the Value of the Exercise Electrocardiogram in the Diagnosis of Stable Chest Pain, Coronary Artery Disease - Current Concepts in Epidemiology, Pathophysiology, Diagnostics and Treatment, Dr. David Gaze (Ed.), ISBN: 978-953-51-0262-5, InTech, Available from:
<http://www.intechopen.com/books/coronary-artery-disease-current-concepts-in-epidemiology-pathophysiology-diagnostics-and-treatment/reassessing-the-value-of-the-exercise-electrocardiogram-in-the-diagnosis-of-stable-chest-pain>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen