

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

185,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



## Screening of some Traditionally Used Plants for Their Hepatoprotective Effect

Saleh I. Alqasoumi<sup>1,2</sup> and Maged S. Abdel-Kader<sup>1</sup>

<sup>1</sup>*Department of Pharmacognosy, College of Pharmacy, Salman Bin Abdulaziz University,*

<sup>2</sup>*Department of Pharmacognosy, College of Pharmacy, King Saud University,  
Kingdom of Saudi Arabia*

### 1. Introduction

Liver is one of the largest organs in human body and the chief site for intense metabolism and excretion (Ram, 2001). It has a surprising role in the maintenance, performance and regulating homeostasis of the body. It is involved with almost all the biochemical pathways responsible for growth, fight against disease, nutrient supply, energy provision and reproduction (Ward & Daly, 1999). The major functions of the liver are carbohydrate, protein and fat metabolism, detoxification, blood coagulation, immunomodulation, secretion of bile and storage of vitamin.

Two major types of reactions occur in the liver in the presence of exogenous substances. The first involve chemical modification of function groups by oxidation, reduction, hydroxylation, sulfonation and dealkylation. Various enzymes including mixed oxidases, cytochromes P-450, and the glutathione S-acyltransferases are involved in such biochemical transformations that usually lead to inactivation of drugs. This step is usually followed by conversion of the resulted metabolites into more water-soluble derivatives that are excreted in the bile or urine via coupling with glucuronate, sulfate, acetate, taurine or glycine moieties (Ram, 2001).

Liver damage inflicted by hepatotoxic agents is of grave consequences (Subramoniam & Pushpangadan, 1999). Liver ailments represent a major global health problem (Baranisrinivasan et al., 2009). Liver cirrhosis is the ninth leading cause of death in the USA (Kim et al., 2002). Toxic chemicals, xenobiotics, alcohol consumption, malnutrition, anaemia, medications, autoimmune disorders (Marina, 2006), viral infections (hepatitis A, B, C, D, etc.) and microbial infections (Sharma & Ahuja, 1997) are harmful and cause damage to the hepatocytes. Hepatotoxic chemicals cause damage to the liver cells mainly by inducing lipid peroxidation and other oxidative events (Dianzani et al., 1991).

In spite of the tremendous advances in modern medicine, no effective drugs are available, which stimulate liver functions and/or offers protection to the liver from damage or help to regenerate hepatic cells (Chatterjee, 2000). In the absence of reliable liver protective drugs in modern medicine, there exists a challenge for pharmaceutical scientists to explore the potential of hepatoprotective activity of plants based on traditional use (Witte et al., 1983). A large number of medicinal preparations are recommended for the treatment of liver

disorders (Chatterjee, 2000) and quite often claimed to offer significant relief. Study of many traditional plants used for liver problems led to the discovery of active compounds yet developed to successful drugs. Silymarin (Morazzoni & Bombardelli, 1995), schisandrin B (Cyang et al., 2000), phyllanthin, hypophyllanthin, picroside I and kutkoside (Ram, 2001) are examples of natural antihepatotoxic compounds derived from traditional herbs. About 600 commercial preparations with claimed liver protecting activity are available all over the world. About 100 Indian medicinal plants belonging to 40 families are components of liver herbal formulation (Handa et al., 1986). The effectiveness of most of these plant products must be scientifically verified to identify new medicaments for the management of liver disorders.

## 2. Silymarin, the standard antihepatotoxic drug

Silymarin, the collective name for an extract from milk thistle, *Silybum marianum* (L.) Gaertneri, is a naturally occurring flavonolignan. Silymarin is a mixture of stereoisomers mainly silybin (also called silybinin, silibin or silibinin) representing 80%, w/w of silymarin. Other minor stereoisomers include isosilybin, dihydrosilybin, silydianin and silychristin (Wagner et al., 1968). Silymarin protects experimental animals against the hepatotoxin  $\alpha$ -amanitin (Hahn et al., 1968) and has a strong antioxidant property (Comoglio et al., 1990). Other reported biological properties of silymarin include inhibition of LOX (Fiebrich & Koch, 1979a) and PG synthetase (Fiebrich & Koch, 1979b). For decades, silymarin has been used clinically in Europe for the treatment of alcoholic liver disease and as antihepatotoxic agent (Salmi & Sarna, 1982). Silymarin is well tolerated and largely free of adverse effects (Comoglio et al., 1990). Silymarin act as an antioxidant by scavenging preoxidant free radicals and by increasing the intracellular concentration of glutathione (GSH). It also exhibits a regulatory action of cellular membrane permeability and increase its stability against xenobiotics injury, increasing the synthesis of ribosomal RNA by stimulating DNA polymerase-I, exerting a steroid like regulatory action on DNA transcription and stimulation of protein synthesis and regeneration of liver cells (Dehmlow et al., 1996; Saller et al., 2007). Silymarin efficacy is not limited to the treatment of toxic and metabolic liver damage; it is also effective in acute, chronic hepatitis and in inhibiting fibrotic activity (Saller et al., 2007). It acts as inhibitor of the transformation of stellate hepatocytes into myofibroblasts, this process is responsible for the deposition of collagen fibres leading to cirrhosis (Fraschini et al., 2002).

## 3. Induction of liver toxicity in experimental animals

In order to study the hepatoprotective effect of plant extracts or pure isolates it is necessary to induce liver toxicity in experimental animal models. The reported protocols for induction of liver toxicity varying greatly in terms of the used liver toxin, doses, duration and route of administration. Below is a collection of the most common experimental procedures use by different groups.

### 3.1 Carbon tetrachloride-induced liver toxicity

Liver damage induced by carbon tetrachloride is the most commonly used model for the screening of hepatoprotective drugs (Slater, 1965). The rise in serum levels of Glutamic

Pyruvate Transaminase (SGPT), Glutamic Oxaloacetic Transaminase (SGOT) and cholesterol following carbon tetrachloride has been attributed to the damaged structural integrity of the liver cells. These components are cytoplasmic in location and released into circulation after cellular damages (Sallie et al., 1991). Carbon tetrachloride also plays a significant role in inducing triacylglyceral accumulation, depletion of GSH, depression of protein synthesis and loss of enzymes activity (Recknagel et al., 1989). Carbon tetrachloride induces hepatotoxicity in rats following its metabolic activation in the hepatocytes. Therefore, it selectively causes toxicity in the liver cells while maintaining semi-normal metabolic function. Carbon tetrachloride is metabolically activated by the cytochrome P-450 dependent mixed oxidase in the endoplasmic reticulum to form trichloromethyl free radical ( $\cdot\text{CCl}_3$ ) and  $\cdot\text{Cl}_3\text{COO}$  which combined with critical cellular macromolecules, cellular lipids and proteins in the presence of oxygen to induce lipid peroxidation (Snyder & Andrews, 1996). Some of the lipid peroxidation products are reactive aldehydes, e.g., 4-hydroxynonenal, which can form adducts with proteins (Weber et al., 2003). These consequences lead to changes in the structures of the endoplasmic reticulum and other membranes hence to increase in plasma membrane permeability to  $\text{Ca}^{2+}$  resulting in a severe disturbances of calcium homeostasis and consequently necrotic cell death (Weber et al., 2003). The loss of metabolic enzyme activation, reduction of protein synthesis and loss of glucose-6-phosphatase activation, all lead to liver injury (Recknagel & Glende, 1973; Azri et al., 1992). In addition to the intracellular events, Kupffer cell activation can contribute to liver injury (elSisi et al., 1993). Kupffer cells are resident macrophages of the liver which constitute approximately 80% of the fixed macrophages in the body (McCuskey, 2006). They may enhance liver injury by oxidant stress (elSisi et al., 1993) or  $\text{TNF-}\alpha$  generation, which may lead to apoptosis (Shi et al., 1998). In more than 70% of the reviewed published data liver toxicity were experimentally induced by Carbon tetrachloride. However, the experimental procedures were considerably different.

### 3.1.1 Single dose carbon tetrachloride-induced liver toxicity

Acute liver toxicity can be induced by a single dose of carbon tetrachloride. However, the route of administration and dose are different from one research group to another. Intraperitoneal injection seems to be the most commonly used method for carbon tetrachloride administration due to the ease of handling and rapid onset of action. Rats are usually a popular experimental animal model and the reported doses were 3 ml/kg (Jamshidzadeh et al., 2005), 2.5 ml/kg (Sen et al., 1993; Nishigaki et al., 1992), 2 ml/kg (Channabasavaraj et al., 2008), 1.5 ml/kg (Bhadauria et al., 2009), 1.25 ml/kg (Rafatullah et al., 2008) or 0.5 ml/kg (Rao et al., 1993). In most cases, carbon tetrachloride is diluted with oils (1:1). The large variation in doses may arise from whether the stated volumes represented the volume of pure carbon tetrachloride or the total volume of the mixture. If this is the case, the wide range of doses from 3- 0.5 ml will shrink to 1 ml (1.5- 0.5 ml). In case of using mice as the experimental animal model the reported carbon tetrachloride doses were much less (0.01, 0.016, 0.02 and 0.03 ml/kg) (Amat et al., 2010; Suzuki et al., 1990; Zhou et al., 2010; Wang et al., 2008).

Induction of liver toxicity via oral routes was also reported. The used doses were 1ml/kg (Harish & Shivanandappa, 2006) 1.25 ml/Kg (Aktay et al., 2000) and 1.5 ml/kg (Gilani, & Janbaz, 1995). Subcutaneous route for administration of carbon tetrachloride was also used

and the reported doses were 0.3 ml/kg (Kumar et al., 2009), 1 ml/kg (Ahmed et al., 2001) or 1.25 ml/kg (Mohamed et al., 2005).

### 3.1.2 Multi doses carbon tetrachloride-induced liver toxicity

The most popular multi-dose protocol for induction of liver toxicity is the subcutaneous administration of 2 ml of carbon tetrachloride/olive oil mixture (1:1) in days 2 and 3 of a five days long experiment (Zafar & Ali, 1998). In another protocol, carbon tetrachloride/olive oil (1:1) mixture was given daily via intraperitoneal injection in a 7 days long experiment. The doses were 0.5 ml (Maheswari & Rao, 2005), 0.8 ml (Özbek et al, 2004) or 1 ml/kg (Somasundaram et al., 2010). In 14 days experiment carbon tetrachloride/liquid paraffin mixture (1:1) was administered intraperitoneal every 72 hours (Christian et al., 2006). Chronic reversible cirrhosis were induced in rats by oral administration of mixture of 20% carbon tetrachloride in corn oil at 0.5mL/ kg body weight doses twice a week (Monday and Thursday) for 6 weeks (Hernandez-Munoz et al., 2001). Subcutaneous injection of 50% carbon tetrachloride in liquid paraffin (3 mL/kg) every other day for four weeks was also used to induce chronic liver toxicity (Chun-ching & Wei-Chih, 1995).

### 3.2 Paracetamol-induced liver toxicity

Paracetamol (acetaminophen) is a well-known antipyretic and analgesic agent. Therapeutic doses of paracetamol are safe, however, toxic doses can produce fatal hepatic necrosis in man, rats and mice (Mitchell et al., 1973). Paracetamol is eliminated mainly as sulfate and glucuronide (Eriksson et al., 1992) when administered in the regular therapeutic doses. Only 5% of the dose is converted into *N*-acetyl-*p*-benzoquinimine (NAPQI). However, upon administration of toxic doses of paracetamol the sulfation and glucuronidation routes become saturated and hence, higher percentage of paracetamol molecules are oxidized to highly reactive NAPQI by cytochrome p-450 enzymes. Semiquinone radicals, obtained by one electron reduction of NAPQI, has an extremely short half-life and is rapidly conjugated with glutathione (GSH), a sulphydryl donor which results in the depletion of liver GSH pool (Remirez et al., 1995). Under conditions of excessive NAPQI formation or reduced of glutathione store, NAPQI covalently binds to vital proteins, the lipid bilayer of hepatocyte membranes and increases the lipid peroxidation. The result is hepatocellular death and centrilobular liver necrosis (McConnachie et al., 2007). Due to liver injury caused by paracetamol overdose, the transport function of the hepatocytes gets disturbed resulting in leakage of plasma membrane (Zimmerman & Seeff, 1970), thus causing an increase in serum enzyme levels.

When rats are used as experimental animal model a single oral dose of 2 gm/kg paracetamol was used to induce liver damage (Chattopadhyay, 2003). The dose in case of using mice was 250 mg/kg (Sabir & Rocha, 2008). Intraperitoneal route of administration was also utilized. A doses of 750 mg/kg (Bhakta et al., 2001) or 835 mg/kg (Yen et al., 2007) were administered to produce liver intoxication in rats, while lower doses of 300 mg/kg (Yuan et al., 2010) were used for induction of liver damage in mice.

### 3.3 D-Galactosamine-induced liver toxicity

Exogenous administration of D-galactosamine (D-GalN) has been found to induce liver damage closely resembles human viral hepatitis (Decker & Keppler, 1972). A single injection

of D-GalN can decrease the uracil nucleotides in the liver and heart (Wills & Asha, 2006). D-GalN markedly depletes hepatic UDP-glucuronic acid whereas extrahepatic UDP-glucuronic is minimally affected. This suggests that D-GalN predominantly inhibits hepatic glucuronidation. It disrupts the synthesis of essential uridylate nucleotides resulting in organelle injury. Depletion of these nucleotides ultimately impairs the synthesis of protein and glycoprotein, leads to progressive damage of cellular membranes. These consequences lead to change in cellular membrane permeability which leads to enzyme leakage to the circulation (Keppler et al., 1970; Abdul-Hussain & Mehendale, 1991). In addition, increased production of reactive oxygen species (ROS) has been reported in primary culture of rat hepatocytes treated with D-GalN (Quintero et al., 2002). Oxygen-derived free radicals released from activated hepatic-macrophages are also one of the primary causes of D-GalN-induced liver damage (Shiratori et al., 1988; Hu & Chen, 1992).

Experimentally induced liver damage was achieved in rats by a single dose of D-GalN 400 mg/kg (Ferenčíková et al., 2003; Kmiec et al., 2000) or 200 mg/kg (Decker & Keppler, 1974) via intrapretoneal injection. For induction of liver toxicity in mice a single dose of 15 mg D-GalN in 0.3mL saline per 20 g by intraperitoneal injection (Wang et al., 2000) was used. The use of D-GalN as liver toxin with a very small concentration of lipopolysaccharide (LPS) (10 µg/kg) was also reported (Tiegs et al., 1989).

### 3.4 Other methods for induction of hepatotoxicity

These methods include the use of some drugs with known side effects target the liver upon prolonged use. Rifampicin (1 g/kg in rats) (Anusuya et al., 2010), menadione (60 mg/kg in mice) (Ip et al., 2004) and anti-tubercular drugs were applied for induction of experimental liver toxicity (Tandon et al., 2008). Ethanol induced liver damage is a major cause of morbidity and mortality worldwide (Purohit et al., 2009). Consequently, ethanol was used as a liver damaging agent in some experiments (Noh et al., 2011; Sathaye et al., 2010). Natural toxins such as aflatoxins are known to have toxic effect on liver. Aflatoxin B1, under the influence of microsomal cytochrome p-450 mediated oxidation, is biotransformed into aflatoxin 8-9-epoxide, which is a reactive intermediate and highly toxic (Iyer et al., 1994). The use of aflatoxin B1 and other aflatoxins as hepatotoxic agent in experimental animal was reported (Banu et al., 2009; Naaz et al., 2007). Chemicals such as trichloroacetic acid (Celik et al., 2009), nitrobenzene (Rathi et al., 2010), thioacetamide (Khatri et al., 2009) and heavy metals such as Cadmium (Obioha et al., 2009) were also utilized to induce experimental liver injury.

Away from the use of chemicals, the hepatoprotective effect of some plant extracts was challenged against liver fibrosis caused by bile duct ligation (Fursule & Patil, 2010).

## 4. Assesment of hepatoprotective activity

The experimental animals are usually treated with the plant extract under investigation for specified period of time. The hepatotoxic agent is usually administered near the end of the experimental period for induction of acute toxicity or in several doses during the course of the experiment for chronic toxicity. The hepatoprotective power of the tested material is assessed by measuring certain biochemical parameters, liver tissue parameters and comparing their levels with normal animals, group receiving standard drug in addition to

the hepatotoxic agent and group receiving only hepatotoxic agent. The most common measured parameters are summarized below.

#### **4.1 Serum biochemical parameters**

##### **4.1.1 Transaminases**

Alanine transaminase (ALT), also called Serum Glutamic Pyruvate Transaminase (SGPT) or Alanine aminotransferase (ALAT) is an enzyme present in hepatocytes. Upon cell damaged, the enzyme leaks into the blood. SGPT level rises dramatically in acute liver damage, such as viral hepatitis or paracetamol overdose (Zimmerman & Seeff, 1970). Elevations are often measured in multiples of the upper limit of normal values. Aspartate transaminase (AST) also called Serum Glutamic Oxaloacetic Transaminase (SGOT) or aspartate aminotransferase (ASAT) is another enzyme associated with liver parenchymal cells. The level of SGOT is raised in acute liver damage, however, it is not specific as it is also present in red blood cells, heart, kidney and skeletal muscle. The ratio of SGOT to SGPT is sometimes useful in differentiating between liver damage and other conditions that elevate the levels of transaminases (Nyblom et al., 2004; Feild et al., 2008). Effective hepatoprotective agents must decrease the elevated levels of transaminases and bring them closer to the normal values as a signe for liver healing.

##### **4.1.2 Alkaline phosphatase**

Alkaline phosphatase (ALP) catalysis the hydrolysis of phosphate esters, and is found in biliary epithelium and the bile canalicular region of hepatocytes. Its function is not well established, but is thought to involve in metabolite transport across cell membranes. Elevation of the level of ALP can suggest intrahepatic, extrahepatic biliary obstruction, or infiltrative diseases of the liver (Feild et al., 2008). Agents that can lower ALP levels will be considered as useful hepatoprotective agents.

##### **4.1.3 Bilirubin**

Bilirubin (Bil) is the breakdown product of normal haem -a part of haemoglobin in red blood cells- catabolism of aged erythrocytes. Bilirubin, loosely bound to albumin in plasma to form a soluble species taken up from the Disse spaces of liver sinusoids into hepatocytes, where it is esterified at its propionyl sites with glucuronic acid under the catalytic activity of uridinediphosphoglucuronate 1A1 transferase enzymes. Esterified bilirubin is excreted into bile as water-soluble bilirubin diglucuronide. Serum concentration of bilirubin is a marker of the liver's ability to take up bilirubin from the plasma into the hepatocyte, conjugate it with glucuronic acid, and excrete bilirubin glucuronides into bile. Elevated level of serum conjugated bilirubin implies regurgitation of bilirubin glucuronides from hepatocytes back into plasma, usually because of intrahepatic or extrahepatic obstruction to bile outflow and cholestasis. The liver has substantial reserve capacity, and normal serum bilirubin levels can be maintained until there is enough injury to reduce the liver's capacity to clear bilirubin from plasma. Serum concentration of bilirubin is very specific for potentially serious liver damage, and is an important indicator of the loss of liver function (Feild et al., 2008). Reduction in the level of serum bilirubin is a strong indication of restoring normal liver function.

#### 4.1.4 Gamma glutamyl transpeptidase (GGT)

Serum Gamma glutamyl transpeptidase (GGT) (also Gamma-glutamyl transferase) is specific to liver injury and more sensitive marker for cholestatic damage than ALP. GGT may be elevated with even minor, sub-clinical levels of liver dysfunction. GGT is raised in alcohol toxicity following several days of moderate ingestion. Rifampin, phenytoin, or barbiturates all resulted in elevation of GGT level. An isolated GGT elevation in these situations does not indicate hepatocellular injury. The GGT level will return to normal after discontinuation of the offending agent. Hepatic dysfunction should be considered if the GGT elevation is associated with other abnormalities in liver biochemistry (Owvns& Evans, 1975). Hepatoprotective agents will reduce the elevated level of GGT.

#### 4.1.5 Total protein & albumin (Alb)

One of the most important liver functions is protein synthesis. Albumin is a major part of the total protein (TP) made specifically by the liver. Liver damage causes disruption and disassociation of polyribosomes on endoplasmic reticulum and thereby reducing the biosynthesis of protein. The TP levels including Alb levels will be depressed in hepatotoxic conditions due to defective protein biosynthesis in liver. Restoring the normal levels of TP including Alb is an important parameter for liver recovery (Navarro & Senior, 2006).

### 4.2 Liver tissue parameters

#### 4.2.1 Glutathione and antioxidant enzymes

Glutathione (GSH) and its related enzymes are playing a vital role as intracellular antioxidants. GSH prevents damage to important cellular components caused by reactive oxygen species such as free radicals and peroxides (Pompella et al., 2003). Glutathione exists in both reduced (GSH) and oxidized (GSSG) states as well. In the reduced state, the thiol group of cysteine is able to donate a reducing equivalent ( $H^{++} e^{-}$ ) to other unstable molecules, such as ROS. In donating an electron, GSH itself becomes reactive, but readily reacts with another reactive GSH to form glutathione disulfide (GSSG). Such a reaction is possible due to the relatively high concentration of glutathione in cells (up to 5 mM in the liver). GSH can be regenerated from GSSG by the enzyme glutathione reductase (GSR or GR) (Boyer, 1989; Tandogan & Ulus, 2006). In healthy cells and tissues, more than 90% of the total glutathione pool is in the reduced form (GSH) and less than 10% exists in the disulfide form (GSSG). An increased GSSG-to-GSH ratio is considered indicative of oxidative stress (Pastore et al., 2003). Another protection from oxidative damage is assured by Glutathione peroxidase (GPx), an enzyme family with peroxidase activity. GPx reduce lipid hydroperoxides to their corresponding alcohols and breakdown hydrogen peroxide into water and oxygen (Castro & Freeman, 2001).

The glutathione S-transferase (GSTs) family are composed of many cytosolic, mitochondrial, and microsomal proteins. GSTs catalyze a variety of reactions and accept endogenous and xenobiotic substrates as well (Udomsinprasert et al., 2005). GSTs catalyse the conjugation of reduced glutathione - via a sulfhydryl group - to electrophilic centres on a wide variety of substrates (Douglas, 1987). This activity detoxifies endogenous compounds such as peroxidised lipids, and enable the breakdown of xenobiotics. GSTs may also bind toxins and serve as transport proteins (Leaver & George, 1998).

Plant	Dose	SGOT		SGPT		ALP		T.Bil		Ref.
		Test	St	Test	St	Test	St	Test	St	
<i>Abutilon indicum</i>	200	64.6	81.5	69.2	90.5	46.9	62.7	54.6	73.8	Porchezhian & Ansari, 2005
<i>Abutilon indicum</i> <sup>a</sup>	200	32.7	68.5	78.8	83.8	52.3	72.4	60.7	70.8	Porchezhian & Ansari, 2005
<i>Adhatoda vasica</i>	100	53.3	62.8	56.0	59.4	-	-	-	-	Bhattacharyya et al., 2005
<i>Anisochilus carnosus</i>	400	52.8	54.9	29.9	30.1	28.2	28.7	13.2	10.5	Venkatesh et al., 2011
<i>Arachniodes exilis</i>	750	71.9	75.8	41.7	49.3	-	-	-	-	Zhou et al., 2010
<i>Artemisia absinthium</i>	200	64.7	70.9	60.1	61.4	-	-	-	-	Amat et al., 2010
<i>Azadirachta indica</i> <sup>a</sup>	500	30.6	36.4	26.9	59.0	28.0	45.5	40.4	53.6	Gomase et al., 2011
<i>Balanites aegyptiaca</i>	500	28.7	56.9	29.9	64.5	21.5	42.8	38.4	52.0	Abdel-Kader & Alqasoumi, 2008
<i>Bixa orellana</i>	500	57.37	-	52.08	-	-	-	21.15	-	Ahsan et al., 2009
<i>Butea monosperma</i>	800	52.1	53.2	78.1	87.1	-	-	-	-	Sharma & Shukla, 2010
<i>Byrsocarpus coccineus</i> <sup>*</sup>	400	44.3	42.9	68.4	55.9	49.0	38.5	46.6	51.0	Akindele et al., 2010
<i>Cajanus cajan</i>	500	56.53	-	50.22	-	-	-	25.0	-	Akindele et al., 2010
<i>Calotropis procera</i> <sup>a</sup>	400	62.2	66.5	69.1	73.5	58.4	61.9	68.7	69.6	Setty et al., 2007
<i>Cassia fistula</i> <sup>a, **</sup>	400	54.0	64.3	46.4	63.1	53.9	58.2	54.1	66.5	Bhakta et al., 2001
<i>Castanea crenata</i> <sup>c</sup>	150	54.1	-	70.6	-	-	-	-	-	Noh et al., 2011
<i>Carduus nutans</i>	500	44.16	-	64.68	-	-	-	-	-	Aktay et al., 2000
<i>Chamomile capitula</i> <sup>a</sup>	400	-	-	-	-	82.6	-	74.2	-	Gupta & Misra, 2006
<i>Cichorium intybus</i>	500	81.9	78.1	56.1	84.3	40.8	47.3	-	-	Ahmed et al., 2003
<i>Cistanche tubulosa</i>	1000	91.6	-	89.7	-	-	-	-	-	Morikawa et al., 2010
<i>Clerodendrum inerme</i>	200	31.6	40.1	83.0	85.6	88.4	89.0	-	-	Gopal & Sengottuvelu, 2008
<i>Commiphora berryi</i>	200	45.6	44.4	65.8	62.4	61.1	65.7	56.3	73.6	Shankar et al., 2008
<i>C. opobalsamum</i>	500	66.2	-	75.6	-	33.0	-	37.3	-	Al-Howiriny et al., 2004
<i>Coptidis rhizoma</i>	600	93.9	-	82.5	-	-	-	-	-	Ye et al., 2009
<i>Cordia macleodii</i>	200	84.8	86.5	77.5	82.2	63.3	60.6	40.6	42.2	Qureshi et al., 2009
<i>Cuscuta chinensis</i> <sup>a</sup>	250	86.8	-	81.6	-	31.0	-	-	-	Yen et al., 2007
<i>Encicostemma axillare</i> <sup>b</sup>	200	91.1	92.2	45.3	20.4	49.2	40.3	33.3	18.6	Jaishree & Badami, 2010
<i>Ephedra foliata</i>	500	42.6	55.1	39.5	66.1	21.2	39.6	46.2	63.5	Alqasoumi et al., 2008b
<i>Euphorbia fusiformis</i>	500	43.1	43.7	30.2	31.7	34.7	37.1	99.9	65.9	Anusuya et al., 2010
<i>Ficus glomerata</i>	500	44.0	30.4	72.8	57.2	68.5	74.6	-	-	Channabasavaraj et al., 2008
<i>Filipendula ulmaria</i>	100	58.7	-	81.9	-	-	-	-	-	Shilova et al., 2008
<i>Fumaria indica</i> <sup>b</sup>	400	72.6	-	79.0	-	68.2	-	79.7	-	Rathi et al., 2008
<i>F. vailantii</i>	500	60.75	-	66.93	-	-	-	-	-	Aktay et al., 2000
<i>Ganoderma lucidum</i> <sup>b</sup>	180	76.6	-	83.6	-	-	-	-	-	Shi et al., 2008
<i>Gentiana olivieri</i>	500	69.57	-	86.39	-	-	-	-	-	Aktay et al., 2000
<i>Hibiscus sabdariffa</i>	500	44.5	66.5	37.1	65.02	21.0	50.6	35.0	69.6	Alqasoumi et al., 2008b
<i>Halenia elliptica</i>	200	57.2	48.8	58.0	47.6	39.0	49.4	46.6	48.6	Huang et al., 2010b
<i>Hedyotis corymbosa</i> <sup>a</sup>	200	59.6	60.8	75.8	66.9	81.0	79.2	37.8	72.4	Sadasivan et al., 2006
<i>Helminthostachys zeylanica</i>	300	64.3	65.0	77.7	78.2	44.3	45.1	66.4	74.0	Suja et al., 2004
<i>Hygrophila auriculata</i>	150	40.8	43.2	24.9	23.5	28.1	26.2	53.0	60.2	Shanmugasundara & Venkataraman, 2006
<i>Kyllinga nemoralis</i>	200	45.6	42.5	68.3	65.5	61.3	66.0	46.6	51.1	Somasundaram et al., 2010
<i>Laggera pterodonta</i>	100	39.6	7.3	31.3	12.5	-	-	-	-	Wu et al., 2007
<i>Laggera pterodonta</i> <sup>b,***</sup>	100	41.9	9.3	35.0	7.5	-	-	-	-	Wu et al., 2007
<i>Mollugo pentaphylla</i>	200	37.2	43.6	53.0	63.9	55.1	55.6	32.6	36.0	Valarmathi et al., 2010
<i>Momordica balsamina</i>	500	37.5	57.6	39.1	57.1	23.2	37.5	52.7	62.0	Alqasoumi et al., 2009b
<i>M. dioica</i>	200	44.7	64.0	54.3	54.6	51.9	59.7	31.1	58.5	Jain et al., 2008
<i>Nelumbo nucifera</i>	500	82.8	82.0	76.5	74.3	39.7	42.0	46.5	50.5	Huang et al., 2010a
<i>Phyllanthus amarus</i> <sup>d</sup>	75	16.6	21.9	28.1	31.8	-	-	-	-	Pramyothin, et al., 2007
<i>Pittosporum neelgherrense</i>	200	70.6	71.9	54.1	55.5	-	-	-	-	Shyamal et al., 2006
<i>Pittosporum neelgherrense</i> <sup>b</sup>	200	65.5	65.4	58.4	59.1	-	-	-	-	Shyamal et al., 2006
<i>Propolis</i>	500	29.4	53.4	37.3	60.2	25.5	43.5	30.1	57.0	Alqasoumi et al., 2008a
<i>Premna corymbosa</i>	400	-	-	-	-	61.6	52.4	-	-	Karthikeyan & Deepa, 2010
<i>Rubus aleaefolius</i>	35	12.7	-	-	-	-	-	-	-	Hong et al., 2010
<i>Sida acuta</i> <sup>a</sup>	100	61.9	60.7	67.7	66.9	79.6	79.2	67.3	72.4	Sreedevi et al., 2009
<i>Smilax regelii</i>	500	13.5	-	47.0	-	-	-	56.4	-	Rafatullah et al., 1991

<sup>a</sup> Paracetamol, <sup>b</sup> D-galactosamine, <sup>c</sup> ethanol, <sup>d</sup> aflatoxin, <sup>e</sup> nitrobenzene, <sup>f</sup> thioacetamide- induced liver toxicity, otherwise CCl<sub>4</sub> was used

\* Livolin, \*\* Liver tonic, \*\*\* Silibinin were used as hepatoprotective standard, otherwise silymarin was use

Table 1. Effect of selected plants on serum biochemical parameters

Treatment of animals with hepatotoxic agents lead to depletion of GSH, reduction in the non-protein sulfhydryl moiety (NP-SH), GPx, GSR activities and ultration of GSTs activity (Naaz et al., 2007; Mitchell et al., 1973; Abdel-Kader et al., 2010; Alqasoumi et al., 2009).

Another part of the antioxidant systems in the bodies is the enzymes Superoxide dismutases (SOD) and Catalase (CAT) (Scott et al., 1991). They are an important antioxidant defence containing heavy metals in nearly all cells exposed to oxygen. SOD catalyzes the dismutation of superoxide into oxygen and hydrogen peroxide. SOD is the most efficient catalytic enzyme; its activity is only limited by the frequency of collision between itself and superoxide (Fredovich, 1997).

CAT catalyzes the decomposition of hydrogen peroxide to water and oxygen (Chelikani et al., 2004). CAT has one of the highest turnover numbers of all enzymes; one CAT enzyme can convert 40 million molecules of hydrogen peroxide to water and oxygen per second (RCSB Protein Data Bank, 2007).

Effective hepatoprotective agents will be able to restore the normal levels of these systems in liver tissue.

#### 4.2.2 Harmful peroxidation products

Malonaldehyde (MDA) is the main end-product of polyunsaturated fatty acid peroxidation (PUFA) following Reactive oxygen species (ROS) insult (Esterbauer et al., 1991). PUFA are essential part of biological membranes (Vaca et al., 1988). MDA is a reactive aldehyde and is one of many reactive electrophile species that cause toxic stress in cells and form covalent protein adducts (Farmer & Davoine, 2007). The production of this aldehyde is used as a biomarker to measure the level of oxidative stress in an organism (Del Rio et al., 2005). The increase in liver MDA levels induced by hepatotoxic agents suggests enhanced lipid peroxidation, leading to hepatic tissue damage and failure of endogenous antioxidant defence mechanisms to prevent formation of excessive free radicals (Souza et al., 1997).

ThioBarbituric Acid Reactive Substances (TBARS) are another harmful substances formed by lipid peroxidation. TBARS are one of the end products formed during the decomposition of lipids by ROS (Olinescu et al., 1994). The tissue concentration of TBARS increase with induced liver toxicity (Sabir & Rocha, 2008).

Physiological amounts of nitric oxide (NO) in the liver has protective effect against damage induced by tumour necrosis factor- $\alpha$  or Fas-dependent apoptosis (Fiorucci et al., 2001). The production of high levels of NO within the liver, via inducible NO synthase (iNOS) promote damage via interference with mitochondrial respiration (Moncada & Erusalimsky, 2002). Hepatocytes of experimental animals produce NO during chronic hepatic inflammation (Billiar et al., 1990 a, b). Human hepatocytes were also stimulated to produce NO by the same combination of endotoxin and cytokines as rat hepatocytes (Palmer et al., 1988; Nussler et al., 1992).

ROS are known to convert amino groups of protein to carbonyl moieties (Perry et al. 2000). Oxidative modification of protein leads to increased recognition and degradation by proteases and loss of enzymatic activity (Rivett & Levine, 1990). Accumulation of carbonyl derivatives of proteins (protein carbonyl) is taken as a biomarker of oxidative protein damage in aging and in various diseases (Dalle-Donne et al., 2003).

Plant	Dose	SOD		CAT		GSH		GPx		MDA		TBAR	
<i>Abutilon indicum</i>	200	-	-	-	-	81.4	82.8	-	-	-	-	-	-
<i>Abutilon indicum</i> <sup>a</sup>	200	-	-	-	-	75.9	77.3	-	-	-	-	-	-
<i>Adhatoda vasica</i>	100	-	-	-	-	-	-	-	-	-	-	55.4	5
<i>Arachniodes exilis</i>	750	88.4	91.4	-	-	-	-	-	-	50.0	55.4	-	-
<i>Artemisia absinthium</i>	200	88.0	90.1	-	-	-	-	71.7	89.0	54.9	58.6	-	-
<i>Butea monosperma</i>	800	-	-	-	-	92.3	92.3	-	-	-	-	-	-
<i>Byrsocarpus coccineus</i> <sup>*</sup>	400	92.8	73.6	93.7	73.6	104.5	75.5	92.2	96.0	82.7	87.3	-	-
<i>Calotropis procera</i> <sup>a</sup>	400	-	-	-	-	58.4	60.5	-	-	-	-	-	-
<i>Castanea crenata</i> <sup>c</sup>	150	104.8	-	89.0	-	-	-	38.6	-	-	-	21.8	-
<i>Carduus nutans</i>	500	-	-	-	-	-	-	-	-	34.8	-	-	-
<i>Chamomile capitula</i> <sup>a</sup>	400	-	-	-	-	91.0	-	-	-	-	-	50.2	-
<i>Clerodendrum inerme</i>	200	-	-	-	-	92.9	88.2	-	-	-	-	-	-
<i>C. opobalsamum</i>	500	-	-	-	-	92.6	-	-	-	-	-	-	-
<i>Coptidis rhizoma</i>	600	97.5	-	-	-	-	-	-	-	-	-	-	-
<i>Cuscuta chinensis</i> <sup>a</sup>	250	87.4	-	86.6	-	-	-	95.6	-	55.4	-	-	-
<i>Ficus glomerata</i>	500	62.5	106.3	105.0	101.7	-	-	-	-	-	-	44.1	4
<i>Filipendula ulmaria</i>	100	66.0	56.1	92.9	73.2	-	-	-	-	-	-	48.4	3
<i>Fumaria indica</i> <sup>b</sup>	400	-	-	-	-	-	-	89.8	93.2	48.6	55.1	-	-
<i>F. vailantii</i>	500	-	-	-	-	-	-	-	-	33.7	-	-	-
<i>Ganoderma lucidum</i> <sup>b</sup>	180	98.5	-	-	-	96.1	-	-	-	51.6	-	-	-
<i>Gentiana olivieri</i>	500	-	-	-	-	-	-	-	-	32.3	-	-	-
<i>Kyllinga nemoralis</i>	200	66.1	51.4	81.3	87.3	59.0	57.2	-	-	51.1	52.6	-	-
<i>Memordica dioica</i>	200	76.8	84.5	83.0	91.5	85.3	88.0	-	-	36.0	48.8	-	-
<i>Nelumbo nucifera</i>	500	85.4	76.5	87.9	55.7	126.0	78.7	-	-	-	-	31.0	3
<i>Phyllanthus amarus</i> <sup>d</sup>	75	85.4	-	100.1	-	202.3	-	108.0	-	-	-	48.3	-
<i>Rubus aleaeifolius</i>	35	89.8	-	-	-	-	-	-	-	20.1	-	-	-
<i>Tecomella undulata</i> <sup>e</sup>	1000	-	-	-	-	134.9	150.9	-	-	50.7	69.4	-	-
<i>Tephrosia purpurea</i> <sup>e</sup>	500	-	-	-	-	139.0	150.9	-	-	65.0	69.4	-	-
<i>Trichosanthes cucumerina</i>	500	-	-	-	-	93.8	94.9	-	-	51.2	54.4	-	-
<i>Spermacoce hispida</i>	200	85.1	-	82.9	-	-	-	89.9	-	51.7	-	-	-
<i>Uvaria chamae</i> <sup>a</sup>	60	-	-	-	-	-	-	-	-	-	-	-	-
<i>Zanthoxylum armatum</i>	400	93.7	99.7	90.9	97.9	89.0	94.2	-	-	67.3	70.0	-	-

<sup>a</sup> Paracetamol, <sup>b</sup> D-galactosamine, <sup>c</sup> ethanol, <sup>d</sup> aflatoxin, <sup>e</sup> thioacetamide- induced liver toxicity, otherwise CCl<sub>4</sub> was used

<sup>\*</sup> Livolin was used as hepatoprotective standard, otherwise silymarin was used

Table 2. Effect of selected plants on liver tissue parameters

### 4.3 Barbiturates sleep time

Short acting barbiturates such as hexobarbiton are metabolized almost exclusively in the liver. Duration of barbiturates induced sleep in intact animals is considered as a reliable index for the activity of hepatic metabolism (Vogel, 1977). Pre-existing liver damage will result in prolongation of the sleeping time after a given dose of barbiturates due to decrease in the amount of the hypnotic broken down per unit time as a result of decreased availability of CYP2E1 contents (Singh et al., 2001). Extracts that can shorten this prolongation of barbiturates sleep time exert protective effect on CYP2E1 system.

### 4.4 Histopathological study of liver tissue

The histological appearance of the hepatocytes reflects their damage conditions (Prophet, et al., 1994). Exposure of hepatocytes to toxic agents such as carbon tetrachloride leads to histopathological changes from the normal histological appearance (Fig. 1). The hepatocytes of rat treated with carbon tetrachloride, showed centrilobular necrosis and extensive fatty changes observed on the midzonal or entire lobe 24 h after treatment (Fig. 1B). Liver tissues of rats treated with carbon tetrachloride and the standard drugs like silymarin showed no necrosis or fatty deposition but had only minimal portal inflammation (Fig.1C). The protective effect of tested extracts or pure materials will be expressed in histopathology as the ability to improve the histological appearance of hepatocytes and bring it closer to the normal hepatocytes of healthy liver (Fig.1A).




Fig. 1. Histological appearance of normal liver (1A), rat liver treated with carbon tetrachloride (1B), rat liver treated with silymarin+ carbon tetrachloride (1C)(H & E,  $\times 200$ ).

## 5. Literature results for some promising hepatoprotective plants

Many plant extracts were tested for their hepatoprotective activity against various liver toxins. All the above mentioned parameters were used to evaluate the protective power of such plants against liver cells damage. In the following Tables some of the most promising results are presented. For handy comparative evaluation of the results percentage of protection is presented even if it was not calculate in the original article. For values where normal levels are increase due to destruction of hepatocytes integrity ( SGOT, SGEPT) or over production of harmful oxidation products (MDA, TBARS) calculations were preformed based on control groups receiving the hepatotoxic agents. Percentage of reduction in the elevated values was compared to protection achieved by silymarin or any other

hepatoprotective agents used in the original articles. On the other hand when experimental liver toxicity resulted in decrease in the normal levels of the measured parameters like the case of normal protective antioxidant enzymes (SOD, CAT), the protection efficacy is presented as percent recovery relative to the negative control values.

## 6. Summary, conclusions and future directions

Liver diseases represent a major global health problem that still has no cure in modern medicine. Some of the traditionally used plants for liver disorders provided useful therapeutic agents. A large number of such plants lack the scientific evidences supporting their effectiveness. Many groups of researchers worldwide were involved in studying the protective effects of plant extracts against experimentally induced liver toxicity. Rats and mice are the animals of choice in such experiments. Carbon tetrachloride followed by paracetamol and D-galactosamine (D-GalN) were used to induce liver toxicity in addition to ethanol and some other drugs affecting the liver on prolonged use. In most cases, positive control groups received silymarin as standard drug for liver protection. Both serum biochemical parameters, liver tissue parameters, barbiturates sleeping time and histopathological examination were used to access liver protection. The most promising data were presented in tables 1 and 2. These biological studies are extremely important to discriminate between useful and useless plants claimed to have liver healing properties. Scientific examination of all traditionally used plants for liver problems is a great goal still to be achieved. This collection of data can be a helpful guide for Phytochemists to explore the constituents of the most promising plants in order to discover new useful natural drugs for the management of liver disorders.

## 7. Reference

- Abdel-Kader, M. S. & Alqasoumi, S. I. (2008). Evaluation of the hepatoprotective effect of the ethanol extracts of *Solanum nigrum*, *Cassia fistula*, *Balanites aegyptiaca* and *Carthamus tinctorius* against experimentally induced liver Injury in Rats. *Alexandria Journal of Pharmaceutical Sciences*, Vol 22, No.1, pp. 47- 50, ISSN 1110-1792
- Abdel-Kader, M. S.; Alqasoumi, S. I.; Hefnawy, M. M. & AlSheikh, A. M. (2010). Hepatoprotective effect and safety studies of *Cleome droserifolia*. *Alexandria Journal of Pharmaceutical Sciences*, Vol 24, No.1, pp. 13- 21, ISSN 1110-1792
- Abdul-Hussain, S. K. & Mehendale, H. M. (1991). Studies on the age dependant effect of galactosamine in primary rat hepatocyte cultures. *Toxicology and Applied Pharmacology*, Vol 107, pp. 504-513, ISSN 0041-008X
- Al-Howiriny, T. A.; Al-Sohaibani, M. O.; Al-Said, M. S.; Al-Yahya, M. A.; El-Tahir, K. H. & Rafatullah, S. (2004). Hepatoprotective properties of *Commiphora opobalsamum* (Balessan, A traditional medicinal plant of Saudi Arabia. *Drugs Under Experimental & Clinical Research*, Vol XXX, No.5/6, pp. 213- 220, ISSN 0378-6501
- Ahmed, B.; Alam, T. & Khan, S. A.(2001). Hepatoprotective activity of *Luffa echinata* fruits. *Journal of Ethnopharmacology*, Vol 76, No.2 ,pp. 187-189, ISSN 0378-8741
- Ahmed, B.; Al-Howiriny, T. A. & Siddiqui, A. B. (2003). Antihepatotoxic activity of seeds of *Cichorium intybus*. *Journal of Ethnopharmacology*, Vol 87, No. 2-3, pp. 237-240, ISSN 0378-8741

- Ahsan, R., Islam, K. M.; Musaddik, A. & Haque, E. (2009). Hepatoprotective activity of methanol extract of some medicinal plants against carbon tetrachloride induced hepatotoxicity in albino rats. *Global Journal of Pharmacology*, Vol 3, No.3, pp. 116-122, ISSN 1992-0075
- Akindele; A. J.; Ezenwanebe, K. O.; Anunobi, C. C. & Adeyemi, O. O. (2010). Hepatoprotective and *in vivo* antioxidant effects of *Byrsocarpus coccineus* Schum. and Thonn. (Connaraceae). *Journal of Ethnopharmacology*, Vol 129, No.1, pp. 46-52, ISSN 0378-8741
- Aktay, G., Deliorman, D.; Ergun, E.; Ergun, F.; Yeşilada, E. & Çevik, C. (2000). Hepatoprotective effects of Turkish folk remedies on experimental liver injury. *Journal of Ethnopharmacology*, Vol 73, pp. 121-129, ISSN 0378-8741
- Alqasoumi, S. I.; El Tahir, K. E. H.; AlSheikh, A. M. & Abdel-Kader, M. S. (2009a). Hepatoprotective Effect and Safety Studies of *Juniperus phoenicea*. *Alexandria Journal of Pharmaceutical Sciences*, Vol 23, No.2, pp. 81- 88, ISSN 1110-1792
- Alqasoumi, S. I.; Al-Howiriny, T. A. & Abdel-Kader, M. S. (2008a). Evaluation of the hepatoprotective effect of *Aloe Vera*, *Clematis hirsute*, *Cucumis prophetarum* and Bee Propolis against experimentally induced liver injury in rats. *International Journal of Pharmacology*, Vol 4, No.3, pp. 213- 217, ISSN 1811-7775
- Alqasoumi, S. I.; Al-Dosari, M. S.; AlSheikh, A. M. & Abdel-Kader, M. S. (2009b). Evaluation of the hepatoprotective effect of *Fumaria parviflora* and *Momordica balsamina* from Saudi Folk medicine against experimentally induced liver injury in rats. *Research Journal of Medicinal Plant*, Vol 3, No.1, pp. 9- 15, ISSN 1996-0875
- Alqasoumi, S. I.; Al-Rehaily, A. J.; Abdulmalik M. AlSheikh & Abdel-Kader, M. S. (2008b). Evaluation of the hepatoprotective effect of *Ephedra foliate*, *Alhagi maurorum*, *Capsella bursa-pastoris* and *Hibiscus sabdariffa* against experimentally induced liver injury in rats. *Natural Product Sciences*, Vol 14, No.2, pp. 95-99, ISSN 1226-3907
- Amat, N.; Upur, H. & Blažeković, B. (2010). *In vivo* hepatoprotective activity of the aqueous extract of *Artemisia absinthium* L. against chemically and immunologically induced liver injuries in mice. *Journal of Ethnopharmacology*, Vol 131, No.2, pp. 478-484, ISSN 0378-8741
- Anusuya, N.; Raju, K. & Manian, S. (2010). Hepatoprotective and toxicological assessment of an ethnomedicinal plant *Euphorbia fusiformis* Buch.-Ham.ex D.Don. *Journal of Ethnopharmacology*, Vol 127, No.2, pp. 463-467, ISSN 0378-8741
- Azri, S.; Mat, H. P.; Reid, L. L.; Gandlofi, A. J. & Brendel, K. (1992). Further examination of the selective toxicity of CCl<sub>4</sub> rat liver silices. *Toxicology and Applied Pharmacology*, Vol 112, No.1, pp. 81- 86, ISSN 0041-008X
- Banu, G. S.; Kumar, G. & Murugesan, A. G. (2009). Effect of ethanolic leaf extract of *Trianthema portulacastrum* L. on aflatoxin induced hepatic damage in rats. *Indian Journal of Clinical Biochemistry*, Vol 24, No.4, pp. 414-418, ISSN 0970-1915
- Baranisrinivasan, P.; Elumalai, E. K.; Sivakumar, C.; Therasa, S. V. & David, E. (2009). Hepatoprotective effect of *Enicostemma littorale* blume and *Eclipta alba* during ethanol induced oxidative stress in albino rats. *International Journal of Pharmacology*, Vol 5, No.4, pp. 268-272, ISSN 1811-7775
- Bhadauria, M.; Nirala, S. K.; Shrivastava, S.; Sharma, A.; Johri, S.; Chandan, B. K.; Singh, B.; Saxena, A. K. & Shukla, S. (2009). Emodin reverses CCl<sub>4</sub> induced

- hepaticcytochrome P450 (CYP) enzymatic and ultra structural changes: the *in vivo* evidence. *Hepatology Research*, Vol 39, No.3, 290-300, ISSN 1386-6346
- Bhakta, T.; Banerjee, S.; Mandal, S. C.; Maity, T. K.; Saha, B. P. & Pal, M. (2001). Hepatoprotective activity of *Cassia fistula* leaf extract. *Phytomedicine*, Vol. 8, No.3, pp. 220-224, ISSN 0944-7113
- Bhattacharyya, D.; Pandit, S.; Jana, U.; Sen, S. & Sur, T. S. (2005). Hepatoprotective activity of *Adhatoda vasica* aqueous leaf extract on d-galactosamine-induced liver damage in rats. *Fitoterapia*, Vol 76, No.2, 223– 225, ISSN 0367-326X
- Billiar, T.R.; Curran, R. D.; Harbrecht, B. G.; Stuehr, D. J.; Demetris, A. J. & Simmons R. L. (1990a). Modulation of nitrogen oxide synthesis *in vivo*: NG-monomethyl-L-arginine inhibits endotoxin-induced nitrate/nitrite biosynthesis while promoting hepatic damage. *Journal of Leukocyte Biology*, Vol 48, No.6, pp. 568-569, ISSN 0741-5400
- Billiar, T. R.; Curran, R. D.; Stuer, D. J.; Stadler, J.; Simmons, R. L. & Murray S. A. (1990b). Inducible cytosolic enzyme activity for the production of nitrogen oxide from L-arginine in hepatocytes. *Biochemical and Biophysical Research Communications*, Vol 168, No.3, pp. 1034-1040, ISSN 0006-291X
- Boyer, T. D. (1989). The glutathione S-transferases: an update. *Hepatology*, Vol 9, No.3, pp. 486-496, ISSN 0270-9139
- Castro, L. & Freeman, B. A. (2001). Reactive oxygen species in human health and disease. *Nutrition*, Vol 17, No.2, pp. 161-165, ISSN: 0899-9007
- Celik, I.; Temur A. & Isik, I. (2009). Hepatoprotective role and antioxidant capacity of pomegranate (*Punica granatum*) flowers infusion against trichloroacetic acid-exposed in rats. *Food and Chemical Toxicology*, Vol 47, No.1, pp. 145-149, ISSN 0278-6915
- Channabasavaraj, K. P.; Badami, S. & Bhojraj, S. (2008). Hepatoprotective and antioxidant activity of methanol extract of *Ficus glomerata*. *Journal of Natural Medicines*, Vol 62, No.3, pp. 379-383, ISSN 1340-3443
- Chatterjee, T. K. (2000). *Medicinal Plants with Hepatoprotective Properties in Herbal Opinions*, Vol III. 135, Books and Allied (P) Ltd., Calcutta.
- Chattopadhyay, R. R. (2003). Possible mechanism of hepatoprotective activity of *Azadirachta indica* leaf extract. Part II. *Journal of Ethnopharmacology*, Vol 89, No.2-3, pp. 217-219, ISSN 0378-8741
- Chelikani, P.; Fita, I. & Loewen, P. C. (2004). Diversity of structures and properties among catalases. *Cellular and Molecular Life Sciences*, Vol 61, No.2, pp. 192-208, ISSN 1420-682X
- Christian, A. J.; Saraswathy, G. R.; Robert, S. J.; Kothai, R.; Chidambaranathan, N.; Nalini, G. & Therasal R. L. (2006). Inhibition of CCl<sub>4</sub> induced liver fibrosis by *Piper longum* Linn. *Phytomedicine*, Vol 13, pp. 196- 198, ISSN 0944-7113
- Chun-Ching, L. & Wei-Chih, L. (1995). Anti-inflammatory and hepatoprotective effects of *Ventilago leiocarpa*. *Phytotherapy Research*, Vol 9, No.1, pp. 11-15. ISSN 0951-418X
- Comoglio, A.; Leonarduzzi, G.; Carini, R.; Busolin, D.; Basaga, H.; Albano, E.; Tomasi, A.; Poli G.; Morazzoni, P. & Magistretti M. J. (1990). Studies on the antioxidant and free radical scavenging properties of IdB 1016: a new flavanolignan complex. *Free*

- Radical Research Communications*, Vol 11, No.1-3, pp. 109–115, Sons. ISSN 0951-418X
- Cyong, J. C.; Ki, S. M.; Iijima, K.; Kobayashi, T. & Furuya, M. (2000). Clinical and pharmacological studies on liver diseases treated with kampo herbal medicine. *The American Journal of Chinese Medicine*, Vol 28, No.3-4, pp. 351–360, ISSN 0192-415X
- Dalle-Donne, I.; Rossi, R.; Giustarini, D.; Aldo Milzani, A. & Colombo, R. (2003). Protein carbonyl groups as biomarkers of oxidative stress. *Clinica Chimica Acta*, Vol 329, No.1-2, pp. 23-38, ISSN 0009-8981
- Decker, K. & Keppler, D. (1972). Galactosamine induced liver injury, In: *Progress in Liver Disease*, H. Popper; F., Schaffner (Eds.) 183–199, Grune and Stratton, New York.
- Decker, K. & Keppler, D. (1974). Galactosamine hepatitis: key role of the nucleotide deficiency period in the pathogenesis of cell injury and cell death. *Reviews of Physiology, Biochemistry and Pharmacology*, Vol 71, pp. 77- 106, ISSN 0303-4240
- Dehmlow, C.; Erhard, J. & de Groot, H. (1996). Inhibition of Kupffer cell functions as an explanation for the hepatoprotective properties of silibinin. *Hepatology*, Vol 23, No. 4, pp. 749–754, ISSN 0270-9139
- Del Rio, D.; Stewart, A. J. & Pellegrini, N. (2005). A review of recent studies on malonaldehyde as toxic molecule and biological marker of oxidative stress. *Nutrition, Metabolism & Cardiovascular Diseases*, Vol 15, No.4, pp. 316 28, ISSN 0939-4753
- Dianzani, M. U.; Muzia, G.; Biocca, M. E. & Canuto, R. A. (1991). Lipid peroxidation in fatty liver induced by caffeine in rats. *International Journal of Tissue Reaction*, Vol 13, No.1, pp. 79–85, ISSN 0250-0868
- Douglas, K. T. (1987). Mechanism of action of glutathione-dependent enzymes. *Advances in Enzymology & Related Areas of Molecular Biology*, Vol 59, pp. 103–167, ISSN 0065258X
- elSisi, A. E.; Earnest, D. L. & Sipes, I. G. (1993). Vitamin A potentiation of carbon tetrachloride hepatotoxicity: role of liver macrophages and active oxygen species. *Toxicology and Applied Pharmacology*, Vol 119, No.2, pp. 295–301, ISSN 0041-008X
- Eriksson, L.; Broome, U., Kahn, M. & Lindholm, M. (1992). Hepatotoxicity due to repeated intake of low doses of paracetamol. *Journal of Internal Medicine*, Vol 231, No.5, pp. 567-570, ISSN 0041-008X
- Esterbauer, H.; Schaur, R. J. & Zollner, H. (1991). Chemistry and biochemistry of 4-hydroxynonenal, malonaldehyde and related aldehydes. *Free Radical Biology and Medicine*, Vol 11, No.1, pp. 81–128, ISSN 0891-5849
- Farmer, E. E. & Davoine, C. (2007). *Reactive electrophile species*. *Current Opinion in Plant Biology*, Vol 10, No.4, pp. 380- 386, ISSN 1369-5266
- Ferenčíková, R.; Červinková, Z. & Drahotka, Z. (2003). Hepatotoxic effect of D-galactosamine and protective role of lipid emulsion. *Physiological research*, Vol 52, No.1, pp. 73- 78, ISSN 0862-8408
- Fiebrich, F. & Koch, H. (1979a). Silymarin, an inhibitor of lipoxygenase. *Experientia*, Vol 35, No.12, pp. 1548–1550, ISSN: 0014-4754
- Fiebrich, F. & Koch, H. (1979b). Silymarin, an inhibitor of prostaglandin synthetase. *Experientia*, Vol 35, No.12, pp. 1550–1552, ISSN: 0014-4754
- Field, K. M.; Dow, C. & Michael, M. (2008). Liver function in oncology: biochemistry and beyond. *The Lancet Oncology*, Vol 9, No.11, pp. 1092–1101, ISSN 0140-6736

- Fiorucci, S.; Mencarelli, A.; Palazzetti, B.; Del Soldato, P.; Morelli, A. & Ignarro, L.J. (2001). An NO-derivative of ursodeoxycholic acid protects against Fas-mediated liver injury by inhibiting caspase activity. *Proceedings of the National Academy of Sciences of the United States of America*, Vol 98, No.5, pp. 2652- 2657, ISSN 0027-8424
- Fraschini, F.; Demartini, G. & Esposti, D. (2002). Pharmacology of silymarin. *Clinical Drug Investigation*, Vol 22, No.1, pp. 51-65, ISSN 1173-2563
- Fredovich, I. (1997). Superoxide anion radical (O<sub>2</sub><sup>-</sup>), superoxide dismutases, and related matters. *The Journal of Biological Chemistry*, Vol 272, No.30, pp. 18515- 18517, ISSN 0021-9258
- Fursule, R. A. & Patil, S. D. (2010). Hepatoprotective and antioxidant activity of Phaseolus trilobus, Ait on bile duct ligation induced liver fibrosis in rats. *Journal of Ethnopharmacology*, Vol 129, No.3 , pp. 416-419, ISSN 0378-8741
- Gilani, A. H. & Janbaz, K. H. (1995). Preventive and curative effects of *Artemisia absinthium* on acetaminophen and CCl<sub>4</sub>-induced hepatotoxicity. *General Pharmacology*, Vol 26, No.2, pp. 309-315, ISSN 0306-3623
- Gomase, P. V.; Rangari V. D. & Verma P. R. (2011). Phytochemical evaluation and hepatoprotective activity of fresh juice of young stem (tender) bark of *Azadirachta indica* A. JUSS. *International Journal of Pharmacy and Pharmaceutical Sciences*, Vol 3, Suppl 2, pp. 55- 59, ISSN 0975-1491
- Gopal, N. & Sengottuvelu, S. (2008). Hepatoprotective activity of Clerodendrum inerme against CCL<sub>4</sub> induced hepatic injury in rats. *Fitoterapia*, Vol 79, No.1, pp. 24- 26, ISSN 0367-326X
- Gupta, A. K. & Misra, N. (2006). Hepatoprotective Activity of Aqueous Ethanolic Extract of *Chamomile capitula* in Paracetamol Intoxicated Albino Rats. *American Journal of Pharmacology and Toxicology*, Vol 1, No.1, pp. 17-20, ISSN 1557-4962
- Hahn, G.; Lehmann, H. D.; Kurten, M.; Ubel, H. & Vogel, G. (1968). Zur Pharmakologie und Toxikologie von Silymarin, des antihepatotoxischen Wirkprinzipes aus *Silybum marianum* (L.) Gaertn. *Arzneimittelforschung*, Vol 18, pp. 698- 704, ISSN 0004-4172
- Handa, S. S.; Sharma, A. & Chakarborti, K. K. (1986). Natural products and plants as liver protecting drugs. *Fitoterapia*, Vol 57, No.5, pp. 307-351, ISSN 0367-326X
- Harish, R. & Shivanandappa T. (2006). Antioxidant activity and hepatoprotective potential of *Phyllanthus niruri*. *Food Chemistry*, Vol 95, No.2 ,pp. 180- 185, ISSN 0308-8146
- Hernandez-Munoz, R.; Diaz-Munoz, M.; Suarez-Cuenca, J. A.; Trejo-Solis, C.; Lopez, V.; Sanchez-Sevilla, L.; Yanez, L. & De Sanches, V. C. (2001). Adenosine reverses a preestablished CCl<sub>4</sub>-induced micronodular cirrhosis through enhancing collagenolytic activity and stimulating hepatocyte cell proliferation in rats. *Hepatology*, Vol 34, No.4, pp. 677- 687, ISSN 0270-9139
- Hong, Z.; Chen, W.; Zhao, J.; Wu, Z.; Zhou, J. H.; Li, T. & Hu J. (2010). Hepatoprotective effects of *Rubus aleaefolius* Poir. And identification of its active constituents. *Journal of Ethnopharmacology*, Vol 129, No.2, pp. 267- 272, ISSN 0378-8741
- Hu, H. L. & Chen, R. D. (1992). Changes in free radicals, trace elements, by d-galactosamine. *Biological Trace Element Research*, Vol 34, No.1 ,pp. 19- 25, ISSN 0163-4984
- Huang, B.; Ban, X.; He, J.; Tong, H.; Zhang, J.; Tian, J. & Wang, Y. (2010a). Hepatoprotective and antioxidant activity of ethanolic extracts of edible lotus (*Nelumbo nucifera* Gaertn.) leaves. *Food Chemistry*, Vol 120, No.3, pp. 873-878, ISSN 0308-8146

- Huang, B.; Ban, X.; He, J.; Zeng, H.; Zhang, P. & Wang, Y. (2010b). Hepatoprotective and antioxidant effects of the methanolic extract from *Halenia elliptica*. *Journal of Ethnopharmacology*, Vol 131, No.2, pp. 276– 281, ISSN 0378-8741
- Ip, S.-P.; Woo, K.-Y.; Lau, O.-W. & Che1, C.-T. (2004). Hepatoprotective Effect of *Sabina przewalskii* against Menadione-induced Toxicity. *Phytotherapy Research*, Vol 18, No.4, pp. 329–331, ISSN 0951-418X
- Iyer, R.S.; Voehler, M.W. & Harris, T.M. (1994). Adenine adduct of Aflatoxin B1 epoxide. *J. Am. Chem. Soc.*, Vol 116, No.20, pp. 8863–8869, ISSN 0002-7863
- Jain, A.; Soni, M.; Deb, L.; Jain, A.; Rout, S. P.; Gupta, V. B. & Krishna, K. L. (2008). Antioxidant and hepatoprotective activity of ethanolic and aqueous extracts of *Momordica dioica* Roxb. Leaves. *Journal of Ethnopharmacology*, Vol 115, No.1, pp. 61–66, ISSN 0378-8741
- Jaishree, V. & Badami, S. (2010). Antioxidant and hepatoprotective effect of swertiamarin from *Enicostemma axillare* against d-galactosamine induced acute liver damage in rats. *Journal of Ethnopharmacology*, Vol 130, No.1, pp. 103– 106, ISSN 0378-8741
- Jamshidzadeh, A.; Fereidooni, F.; Salehi, Z. & Niknahad, H. (2005). Hepatoprotective activity of *Gundelia tourenfortii*. *Journal of Ethnopharmacology*, Vol 101, No.1-3, pp. 233-237, ISSN 0378-8741
- Karthikeyan, M. & Deepa, K. (2010). Hepatoprotective effect of *Premna corymbosa* (Burn. F.) Rottl. & Willd. Leaves extract on CCl<sub>4</sub> induced hepatic damage in Wistar albino rats. *Asian Pacific Journal of Tropical Medicine*, Vol 3, No.1, pp. 17- 20, ISSN 16879686
- Keppler, D. O.; Rudigier, J. F.; Bischoff, E. & Decker, K. (1970). The trapping of uridine phosphates by d-galactosamine, d-glucosamine, and 2-deoxy-d-galactose. A study on the mechanism of galactosamine hepatitis. *European Journal of Biochemistry*, Vol 17, No.2, pp. 246– 253, ISSN 0014-2956
- Kim, W. R.; Brown, R. S.; Terrault, N. A. & El-Serag, H. (2002). Burden of liver disease in the United States: Summary of work-Shop. *Hepatology*, Vol 36, No.1, pp. 227-242, ISSN 0270-9139
- Khatri, A.; Garg, A. & Agrawal, S. A. (2009). Evaluation of hepatoprotective activity of aerial parts of *Tephrosia purpurea* L. and stem bark of *Tecomella undulate*. *Journal of Ethnopharmacology*, Vol 122, No.1, pp. 1–5, ISSN 0378-8741
- Kmieć, Z.; Ryszard Smoleński, R. T.; Zyśh, M. & Myśliwski, A. (2000). The effects of galactosamine on UTP levels in the livers of young, adult and old rats. *Acta Biochimica Polonica*, Vol 47, No.2, pp. 349- 353, ISSN 0001-527X
- Kumar, S. S.; Kumar, B. R. & Mohan G. K. (2009). Hepatoprotective effect of *Trichosanthes cucumerina* Var *cucumerina* L. on carbon tetrachloride induced liver damage in rats. *Journal of Ethnopharmacology*, Vol 123, No.2, pp. 347–350, ISSN 0378-8741
- Leaver, M. J. & George, S. G. (1998). A piscine glutathione S-transferase which efficiently conjugates the end-products of lipid peroxidation. *Marine Environmental Research*, Vol 46, No.1-5, pp. 71–74, ISSN 0141-1136
- Madubunyi, I. I. (2010). Hepatoprotective activity of *Uvaria chamae* root bark methanol extract against acetaminophen-induced liver lesions in rats. *Comparative Clinical Pathology*, In Press, ISSN 1618-5641
- Maheswari, M. U. & Rao, P. G. M. (2005). Anti hepatotoxic effect of grape seed oil. *Indian Journal of Pharmacology*, Vol 37, No.3, pp. 179- 182, ISSN 0253-7613

- Marina, N. (2006). Hepatotoxicity of antiretrovirals: incidence, mechanisms and management. *Journal of Hepatology*, Vol 44, No.1, pp. S132–S139, ISSN 0168-8278
- McConnachie, L. A.; Mohar, I.; Hudson, F. N.; Ware, C. B.; Ladiges, W. C.; Fernandez, C.; Chatterton-Kirchmeier, S.; White, C. C.; Pierce, R. H. & Kavanagh, T. J. (2007). Glutamate cysteine ligase modifier subunit deficiency and gender as determinants of acetaminophen-induced hepatotoxicity in mice. *Toxicological Sciences*, Vol 99, No.2, pp. 628–636, ISSN 1096-6080
- McCuskey, R. S. (2006). Anatomy of the liver. In: *Zakim and Boyer's Hepatology*, T. D. Boyer; T. L. Wright; M. Manns, (Eds.), 37, 5th ed., Saunders–Elsevier, Philadelphia.
- Mitchell, J. R.; Jollow, D. J.; Potter, W. Z.; Gillette, J. R. & Brrodle, B.N. (1973). Acetaminophen induced hepatic necrosis. I. Role of drug metabolism. *Journal of Pharmacology and Experimental Therapeutics*, Vol 187, No.1, pp. 185–194, ISSN 0022-3565
- Mohamed, M. A.; Marzouk, M. S. A.; Moharram, F. A.; El-Sayed, M. M. & Baiuomy A. R. (2005). Phytochemical constituents and hepatoprotective activity of *Viburnum tinus*. *Phytochemistry*, Vol 66, No.23, pp. 2780–2786, ISSN 1819-3471
- Moncada, S. & Erusalimsky, J. D. (2002). Does nitric oxide modulate mitochondrial energy generation and apoptosis? *Nature Reviews Molecular Cell Biology*, Vol 3, pp. 214–220, ISSN 1471-0072
- Morazzoni, P. & Bombardelli, E. (1995). *Silybum marianum*. *Fitoterapia*, Vol 66, No.1, pp. 3–42, ISSN 0367-326X
- Morikawa, T.; Pan, Y.; Ninomiya, K.; Imura, K.; Matsuda, H.; Yoshikawa, M.; Yuan, D. & Muraoka, O. (2010). Acylated phenylethanoid oligoglycosides with hepatoprotective activity from the desert plant *Cistanche tubulosa*. *Bioorganic & Medicinal Chemistry*, Vol 18, No.5, pp. 1882–1890, ISSN 0960-894X
- Naaz, F.; Javed, S. & Abdin, M. Z. (2007). Hepatoprotective effect of ethanolic extract of *Phyllanthus amarus* Schum. et Thonn. on aflatoxin B1-induced liver damage in mice. *Journal of Ethnopharmacology*, Vol 113, No.3, pp. 503–509, ISSN 0378-8741
- Navarro, V. J. & Senior, J. R. (2006). Drug-related hepatotoxicity. *New England Journal of Medicine*, Vol 35, No.4, pp. 731–739, ISSN 0028-4793
- Nishigaki, I.; Kuttan, R.; Oku, H.; Ashoori, F.; Abe, H. & Yagi, K. (1992). Suppressive effect of curcumin on lipid peroxidation induced in rats by CCl<sub>4</sub> or <sup>60</sup>Co irradiation. *Journal of Clinical Biochemistry and Nutrition*, Vol 13, pp. 23–30, ISSN 0912-0009
- Noh J.-R.; Kim, Y.-H.; Gang, G.-T.; Hwang, J. H.; Lee, H.-S.; Ly, S.-Y.; Oh, W.-K.; Song, K.-S. & Lee, C.-H. (2011). Hepatoprotective effects of chestnut (*Castanea crenata*) inner shell extract against chronic ethanol-induced oxidative stress in C57BL/6 mice. *Food and Chemical Toxicology*, In Press, ISSN 0278-6915
- Nussler, A.; DiSilvio, M.; Billiar, T. R.; Hoffman, R. A.; Geller, D. A.; Selby, R.; Madariaga, J. & Simmons R. L. (1992). Stimulation of the nitric oxide synthase pathway in human hepatocytes by cytokines and endotoxin. *Journal of Experimental Medicine*, Vol 176, pp. 261–264, ISSN 0022-1007
- Nyblom, H.; Berggren, U.; Balldin, J. & Olsson, R. (2004). High AST/ALT ratio may indicate advanced alcoholic liver disease rather than heavy drinking. *Alcohol & Alcoholism*, Vol 39, No.4, pp. 336–339, ISSN 0735-0414

- Obioha, U. E.; Suru, S. M.; Ola-Mudathir, K. F. & Faremi, T. Y. (2009). Hepatoprotective potentials of Onion and Garlic extracts on Cadmium-Induced oxidative damage in rats. *Biological Trace Element Research*, Vol 129, No.1-3, pp. 143–156, ISSN 0163-4984
- Olinescu, R.; Alexandrescu, R.; Hulea, S. A. & Kummerow, F. A. (1994). Tissue lipid peroxidation may be triggered by increased formation of bilirubin *in vivo*. *Research Communications in Chemical Pathology & Pharmacology*, Vol 84, No.1, pp. 27-34, ISSN 0034-5164
- Owvans, D. & Evans, J. (1975). Population studies on Gilbert's syndrome. *Journal of Chemical Genetics*, Vol 12, pp. 152- 156, ISSN 0104-6632
- Özbek, H.; Çitoğlu, G. S.; Dülger, H.; Uğraş, S. & Sever, B. (2004). Hepatoprotective and anti-inflammatory activities of *Ballota glandulosissima*. *Journal of Ethnopharmacology*, Vol 95, No.2-3, pp. 143–149, ISSN 0378-8741
- Palmer, R. M. J.; Ashton, D. S.; & Moncada, S. (1988). Vascular endothelial cells synthesis nitric oxide from L-arginine. *Nature*, Vol 333, No.6174, pp. 664-666, ISSN 0028-0836
- Pastore, A.; Piemonte, F.; Locatelli, M.; Russo, A. L.; Gaeta, L. M.; Tozzi, G. & Federici, G. (2003). Determination of blood total, reduced, and oxidized glutathione in pediatric subjects. *Clinical Chemistry*, Vol 47, No.8, pp. 1467–1469, ISSN 0009-9147
- Perry, G.; Raina, A. K.; Nunomura, A.; Wataya T.; Sayre, L. M. & Smith M. A. (2000). How important is oxidative damage? Lessons from Alzheimer's disease. *Free Radical Biology and Medicine*, Vol 28, No.5, pp. 831-834, ISSN 0891-5849
- Pompella, A.; Visvikis, A.; Paolicchi, A.; de Tata, V. & Casini, A. F. (2003). The changing faces of glutathione, a cellular protagonist. *Biochemical Pharmacology*, Vol 66, No.8, pp. 1499–503, ISSN 0006-2952
- Porchezian, E. & Ansari, S. H. (2005). Hepatoprotective activity of *Abutilon indicum* on experimental liver damage in rats. *Phytomedicine*, Vol 12, No.1-2, pp. 62–64, ISSN 0944-7113
- Pramyothin, P.; Ngamtin, C.; Pongshompoo, S. & Chaichantipyuth, C. (2007). Hepatoprotective activity of *Phyllanthus amarus* Schum. et Thonn. extract in ethanol treated rats: *In vitro* and *in vivo* studies. *Journal of Ethnopharmacology*, Vol 114, No.2, pp. 169–173, ISSN 0378-8741
- Prophet, E. P.; Mills, B.; Arrington, J. B. & Sobin, L. H. (1994). *Laboratory Methods in Histology*. 1<sup>st</sup> Ed, American Registry of Pathology, Washington, D.C.
- Purohit, V.; Gao, B. & Song, B.-J. (2009). Molecular Mechanisms of Alcoholic Fatty Liver. *Alcohol.: Clin. Exp. Res.*, Vol 33, No.2, pp. 191–205, ISSN 1530-0277
- Quintero, A.; Pedraza, C. A.; Siendones, E.; Kamal, E. A. M.; Colell, A.; García -Ruiz, C.; Montero, J. L.; Dela, M. M.; Fernández-Checa, J. C.; Miño, G.; Muntané, J.; Kamal, E. A. M. & Colell, A. (2002). PGE1 protection against apoptosis induced by D-galactosamine is not related to the modulation of intracellular free radical production in primary culture of rat hepatocytes. *Free Radical Research*, Vol 36, No.3, pp. 345–355, ISSN 1071-5762
- Qureshi, N. N.; Kuchekar, B. S.; Logade, N. A. & Haleem, M. A. (2009). Antioxidant and hepatoprotective activity of *Cordia macleodii* leaves. *Saudi Pharmaceutical Journal*, Vol 17, No.4, pp. 299– 302, ISSN 1319-0164
- Rafatullah, S.; Al-Sheikh, A.; Alqasoumi, S.; Al-Yahya, M.; El-Tahir, K. & Galal, A. (2008). Protective effect of fresh radish juice (*Raphanus sativus* L.) against carbon


- tetrachloride-induced hepatotoxicity. *International Journal of Pharmacology*, Vol 4, No.2, pp. 130- 134. ISSN 1811-7775.
- Rafatullah, S.; Mossa, J. S.; Ageel, A. M.; Al-Yahya, M. A. & Tariq, M. (1991). Hepatoprotective and safety evaluation studies on Sarsaparilla. *International Journal of Pharmacognosy (Now Pharmaceutical Biology)*, Vol 29, No.4, pp. 296- 301, ISSN 1388-0209
- Ram, V. J. (2001). Herbal preparations as a source of hepatoprotective agents. *Drug News and Perspective*, Vol 14, No.6, pp. 353-363. ISSN 0214-0934
- Ranawat, L.; Bhatt, J. & Patel J. (2010). Hepatoprotective activity of ethanolic extracts of bark of *Zanthoxylum armatum* DC in CCl<sub>4</sub> induced hepatic damage in rats. *Journal of Ethnopharmacology*, Vol 127, No.3, pp. 777-780, ISSN 0378-8741
- Rao, P. G. M.; Rao, S. G. & Kumar, V. (1993). Effects of hepatogard against carbon tetrachloride induced liver damage in rats. *Fitoterapia*, Vol 64, No.2, pp. 108-113, ISSN 0367-326X
- Rathi, M. A.; Thirumoorthi, L.; Sunitha, M.; Meenakshi, P.; Gurukumar, D. & Gopalakrishnan, V. K. (2010). Hepatoprotective activity of *Spermacoce hispida* Linn. extract against nitrobenzene induced hepatotoxicity in rats. (2010). *Journal of Herbal Medicine and Toxicology*, Vol 4, No.2, pp. 201-205, ISSN 0973-4643.
- Rathi, A., Srivastava, A. K.; Shirwaikar, A.; Rawat, A. K. S. & Mehrotra S. (2008). Hepatoprotective potential of *Fumaria indica* Pugsley whole plant extracts, fractions and an isolated alkaloid protopine. *Phytomedicine*, Vol 15, pp. 470- 477, ISSN 0944-7113
- RCSB Protein Data Bank. Catalase. *Molecule of the Month*. (2004-09-01). Retrieved on 2007-02-11, <http://www.pdb.org/pdb/101/motm.do?momID=57>
- Recknagel, R. O. & Glende, E. A. Jr. (1973). Carbon tetrachloride hepatotoxicity: an example of lethal cleavage. *CRC (Critical Reviews in Toxicology)*. Vol. 2, No.3, pp. 263-297, ISSN 1040-8444
- Recknagel, R. O.; Glonde, E. A.; Dolak, J. A. & Walter, R. L. (1989). Mechanism of carbon tetrachloride toxicity. *Pharmacology and Therapeutics*, Vol 43, No.1, pp. 139-154, ISSN 0163-7258
- Remirez, D.; Commandeur, J. N. M.; Ed Groot, E. & Vermeulen, N. P. E. (1995). Mechanism of protection of *Lobenzarti* against paracetamol-induced toxicity in rat hepatocytes. *European Journal of Pharmacology: Environmental Toxicology and Pharmacology*, Vol 293, No.4, pp. 301- 308, ISSN 0926-6917
- Rivett, A. J. & Levine, R. L. (1990): Metal-catalyzed oxidation of *Escherichia coli* glutamine synthetase: correlation of structural and functional changes. *Archives of Biochemistry and Biophysics*. Vol 278, No.1, pp. 26- 34, ISSN 0003-9861
- Sabir, S. M. & Rocha, J. B. T. (2008). Antioxidant and hepatoprotective activity of aqueous extract of *Solanum fastigiatum* (false “Jurubeba”) against paracetamol-induced liver damage in mice. *Journal of Ethnopharmacology*, Vol 120, No.2, pp. 226- 232, ISSN 0378-8741
- Sadasivan, S.; Latha, P. G.; Sasikumar, J. M.; Rajashekar, S.; Shyamal, S. & Shine, V. J. (2006). Hepatoprotective studies on *Hedyotis corymbosa* (L.) Lam. *Journal of Ethnopharmacology*, Vol 106, No.2, pp. 245- 249, ISSN 0378-8741

- Saller, R.; Melzer, J.; Reichling, J.; Brignoli, R. & Meier, R. (2007). An updated systematic review of the pharmacology of silymarin. *Forschende Komplementmedizin (Research in Complementary Medicine)*, Vol 14, No.2, pp. 70- 80, ISSN 1661-4119
- Sallie, R.; Tredger, J. M. & William, R. (1991). *Drugs and the liver. Part I. Testing liver function. Biopharmaceutics & Drug Disposition*, Vol 12, No.4, pp. 251- 259, ISSN 0142-2782
- Salmi, H. A. & Sarna S. (1982). Effect of silymarin on chemical, functional, and morphological alterations of the liver. A double-blind controlled study. *Scandinavian Journal of Gastroenterology*, Vol17, No.4, pp. 517- 521, ISSN 0036-5521
- Sathaye, S., Bagul, Y.; Gupta, S.; Kaur, H. & Roopali Redkar, R. (2010). Hepatoprotective effects of aqueous leaf extract and crude isolates of *Murraya koenigii* against *in vitro* ethanol-induced hepatotoxicity model. *Experimental and Toxicologic Pathology*, In Press, ISSN 0940-2993
- Scott, M. D.; Lubin, B. H.; Zuo, L. & Kuypers, F. A. (1991). Erythro-cyte defense against hydrogen peroxide: preeminent importance of catalase. *Journal of Laboratory and Clinical Medicine*, Vol 118, No.1, 7- 16, ISSN 0022-2143
- Sen, T.; Basu, A.; Ray, R. N. & Nag Shaudhuri, A. K. (1993). Hepatoprotective effects of *Pluchea indica* (Lees) extract in experimental acute liver damage in rodents. *Phytotherapy Research*, Vol 7, No.5, pp. 352-355, ISSN 0951-418X
- Setty, S. R.; Quereshi, A. A.; Swamy, A. H. M. V.; Patil, T.; Prakash, T.; Prabhu, K. & Gouda, A. V. (2007). Hepatoprotective activity of *Calotropis procera* flowers against paracetamol-induced hepatic injury in rats. *Fitoterapia*, Vol 78, pp. 451- 454, ISSN 0367-326X
- Shankar, N. L. G.; Manavalan, R.; Venkappayya, D. & Raj, C. D. (2008). Hepatoprotective and antioxidant effects of *Commiphora berryi* (Arn) Engl bark extract against CCl<sub>4</sub>-induced oxidative damage in rats. *Food and Chemical Toxicology*, Vol 46, No.9, pp. 3182- 3185, ISSN 0278-6915
- Shanmugasundaram, P. & Venkataraman, S. (2006). Hepatoprotective and anti oxidant effects of *Hygrophila Auriculta* (K. Schum) Heine Acanthaceae root extract. *Journal of Ethnopharmacology*, Vol 104, No.1-2, pp. 124- 128, ISSN 0378-8741
- Sharma, N. & Shukla, S. (2010). Hepatoprotective potential of aqueous extract of *Butea monosperma* against CCl<sub>4</sub> induced damage in rats. *Experimental and Toxicologic Pathology*, In Press, ISSN 0940-2993
- Shi, J.; Aisaki, K.; Ikawa, Y. & Wake, K., (1998). Evidence of hepatocyte apoptosis in rat liver after administration of carbon tetrachloride. *American Journal of Pathology*, Vol 153, No.2, 515- 525, ISSN 0002-9440
- Shi, Y.; Sun J.; He, H.; Guo, H. & Zhang, S. (2008). Hepatoprotective effects of *Ganoderma lucidum* peptides against d-galactosamine-induced liver injury in mice. *Journal of Ethnopharmacology*, Vol 117, No.3, pp. 415- 419, ISSN 0378-8741
- Shilova, I. V.; Zhavoronok, T. V.; Souslov, N. I.; Novozheeva, T. P.; Mustafin, R. N. & Losseva, A. M. (2008). Hepatoprotective Properties of Fractions from Meadowsweet Extract during Experimental Toxic Hepatitis. *Bulletin of Experimental Biology and Medicine, Pharmacology and Toxicology*, Vol. 146, No.1, pp. 49- 51, ISSN 1535-3702
- Shiratori, Y.; Kawase, T.; Shiina, S.; Okano, K.; Sugimoto, T.; Teraoka, H.; Matano, S.; Matsumoto, K. & Kamii, K. (1988). Modulation of hepatotoxicity by macrophages in the liver. *Hepatology*, Vol 8, No.4, pp. 815- 821, ISSN 0270-9139

- Shyamal, S.; Latha, P. G.; Shine, V. J.; Suja, S. R.; Rajasekharan S. & Devi, T. G. (2006). Hepatoprotective effects of *Pittosporum neelgherrense* Wight & Arn., a popular Indian ethnomedicine. *Journal of Ethnopharmacology*, Vol 107, No.1, pp. 151– 155, ISSN 0378-8741
- Singh, B; Saxena, A. K.; Chandan, B. K.; Agrawal, S. G. & Anand, K. K. (2001). In vivo hepatoprotective activity of active fraction from ethanolic extract of *Eclipta alba* leaves. *Indian Journal of Physiology and Pharmacology*, Vol 45, No.4, pp. 435– 441, ISSN 0019-5499
- Slater, T. F. (1965). *Biochemical mechanism of liver injury*. London: Academic Press.
- Snyder, R. & Andrews, L. S. (1996). Toxic effects of solvents and vapors. In: *Toxicology: the Basic Science of Poisons*, C. D. Klassen, (Ed.), 737–772, McGraw-Hill, New York.
- Somasundaram, A.; Karthikeyan, R.; Velmurugan, V.; Dhandapani, B. & Raja, M. (2010). Evaluation of hepatoprotective activity of *Kyllinga nemoralis* (Hutch & Dalz) rhizomes. *Journal of Ethnopharmacology*, Vol 127, No.2, pp. 555– 557, ISSN 0378-8741
- Souza, M. F.; Rao, V. S. N. & Silveira, E. R. (1997). Inhibition of lipid peroxidation by ternatin, a tetramethoxyflavone from *Egletes viscosa* L. *Phytomedicine*, Vol 4, pp. 25– 9, ISSN 0944-7113
- Sreedevi, C. D.; Latha, P. G.; Ancy, P.; Suja, S. R.; Shyamal, S.; Shine, V. J.; Sini, S.; Anuja, G. I. & Rajasekharan, S. (2009). Hepatoprotective studies on *Sida acuta* Burm. f. *Journal of Ethnopharmacology*, Vol 124, No.2, pp. 171– 175, ISSN 0378-8741
- Subramoniam, A. & Pushpangadan, P. (1999). Development of phytomedicine for liver diseases. *Indian Journal of Pharmacology*, Vol 31, No.3, pp. 166–175, ISSN 0253-7613
- Suja, S. R.; Latha, P. G.; Pushpangadan, P. & Rajasekharan, S. (2004). Evaluation of hepatoprotective effects of *Helminthostachys zeylanica* (L.) Hook against carbon tetrachloride-induced liver damage in Wistar rats. *Journal of Ethnopharmacology*, Vol 92, No.1, pp. 61– 66, ISSN 0378-8741
- Suzuki, I.; Tanaka, H.; Yajima, H.; Fukuda, H.; Sezaki, H.; Koga, K.; Hirobe, M. & Nakajima, T. (1990). *Pharmaceutical Research and Development*, Vol 19, 227, Hirokawa Publishing, Tokyo.
- Tandogan, B. & Ulus, N. N. (2006). Kinetic Mechanism and Molecular Properties of Glutathione Reductas, *FABAD Journal of Pharmaceutical Sciences*, Vol 31, pp. 230– 237. ISSN 1300-4182.
- Tandon, V. R.; Khajuria, V.; Kapoor, B.; Kour, D. & Gupta, S. (2008). Hepatoprotective activity of *Vitex negundo* leaf extract against anti-tubercular drugs induced hepatotoxicity. *Fitoterapia*, Vol 79, pp. 533– 538, ISSN 0367-326X
- Tiegs, G., Wolter, M. & Wendel, A. (1989). Tumor necrosis factor is a terminal mediator in galactosamine/endotoxin-induced hepatitis in mice. *Biochemical Pharmacology*, Vol 38, No.4, pp. 627– 631, ISSN 0006-2952
- Udomsinprasert, R.; Pongjaroenkit, S.; Wongsantichon, J.; Oakley, A. J.; Prapanthadara, L. A.; Wilce, M. C. & Ketterman, A. J. (2005). Identification, characterization and structure of a new Delta class glutathione transferase isoenzyme. *Biochemical Journal*, Vol 388, No.3, pp. 763 – 771, ISSN 0264-6021
- Vaca, C. E.; Wilhelm, J. & Harms-Rihdsahl, M. (1988). Interaction of lipid peroxidation product with DNA. A review. *Mutation Research-Reviews in Genetic Toxicology*, Vol 195, No.2, pp. 137– 149, ISSN 0165-1110

- Valarmathi, R.; Rajendran, A.; Gopal, V.; Senthamarai, R.; Akilandeswari, S. & Srileka, B. (2010). Protective Effect of the whole plant of *Mollugo pentaphylla* Linn. against Carbon Tetrachloride induced Hepatotoxicity in Rats. *International Journal of PharmTech Research*, Vol 2, No.3, pp. 1658-1661, ISSN 0974-4304
- Venkatesh, P.; Dinakar, A. & Senthilkumar, N. (2011). Hepatoprotective activity of an ethanolic extract of stems of *Anisochilus carnosus* against carbon tetrachloride induced hepatotoxicity in rats. *International Journal of Pharmacy and Pharmaceutical Sciences*, Vol 3, No.1, pp. 243- 245, ISSN- 0975-1491
- Vogel, G. (1977). Natural substances with effects on the liver, In: *New natural products and plant drugs with pharmacological, biological or therapeutic activity*. H. Wagner & P. Wolff (eds), pp 249-265, Springer-Verlag, Berlin, Heidelberg, New York.
- Wagner, H.; Seligmann, O.; Horhammer, L. & Munster, R. (1968). The chemistry of silymarin (silybin), the active principle of the fruits of *Silybum marianum* (L.) Gaertn. [*Carduus marianus* (L.)]. *Arzneimittelforschung*, Vol 18, No.6, pp. 688- 696, ISSN 0004-4172
- Wang, M. Y.; Liu, Q.; Che, Q. M. & Lin, Z. B. (2000). Effects of triterpenoids from *Ganoderma lucidum* (Leyss. Ex fr.) karst on three different experimental liver injury models in mice. *Acta Pharmaceutica Sinica*, Vol 35, No.5, pp. 326- 329, ISSN 0513-4870
- Wang, N.; Li, P.; Wang, Y.; Peng, W.; Wu, Z.; Tan, S.; Liang, S.; Shen, X. & Su, W. (2008). Hepatoprotective effect of *Hypericum japonicum* extract and its fractions. *Journal of Ethnopharmacology*, Vol 116, No.1, pp. 1- 6, ISSN 0378-8741
- Ward, F. M. & Daly, M. J. (1999). Hepatic Disease. In: *Clinical Pharmacy and Therapeutics*, R. Walker & C. Edwards (Eds.), 195- 212, Churchill Livingstone, New York.
- Weber, L.W.; Boll, M. & Stampfl, A. (2003). Hepatotoxicity and mechanism of action of haloalkanes: carbon tetrachloride as a toxicological model. *Critical Reviews in Toxicology*, Vol 33, No.2, pp. 105- 136, ISSN 1040-8444
- Wills, P. J. & Asha, V. V. (2006). Protective effect of *Lygodium exuosum* (L.) Sw. (Lygodiaceae) against d-galactosamine induced liver injury in rats. *Journal of Ethnopharmacology*, Vol 108, No.1, pp. 116-123, ISSN 0378-8741
- Witte, I.; Berlin, J.; Wray, V.; Schubert, W.; Kohl, W.; Hofle, G. & Hammer, J. (1983). Mono and diterpins from cell culture of *Thuja Occidenalis*. *Planta Medica*, Vol 49, pp. 216-221, ISSN 00320943
- Wu, Y.; Yang, L.; Wang, F.; Wu, X.; Zhou, C.; Shi, S.; Mo, J. & Zhao, Y. (2007). Hepatoprotective and antioxidative effects of total phenolics from *Laggetera pterodonta* on chemical-induced injury in primary cultured neonatal rat hepatocytes. *Food and Chemical Toxicology*, Vol 45, No.8, pp. 1349- 1355, ISSN 0278-6915
- Ye, X.; Feng, Y.; Tong, Y.; Ng, K.-M.; Tsao, S.; Lau, G. K. K.; Sze, C.; Zhang, Y.; Tang, J.; Shen, J. & Kobayashi, S. (2009). Hepatoprotective effects of *Coptidis rhizoma* aqueous extract on carbon tetrachloride-induced acute liver hepatotoxicity in rats. *Journal of Ethnopharmacology*, Vol 124, No.1, pp. 130- 136, ISSN 0378-8741
- Yen, F.-L.; Wu, T.-H.; Lin, L.-T. & Lin, C.-C. (2007). Hepatoprotective and antioxidant effects of *Cuscuta chinensis* against acetaminophen-induced hepatotoxicity in rats. *Journal of Ethnopharmacology*, Vol 111, No.1, pp. 123-128, ISSN 0378-8741
- Yuan, H.-D., Jin, G.-Z. & Piao, G.-C. (2010). Hepatoprotective effects of an active part from *Artemisia sacrorum* Ledeb. against acetaminophen-induced toxicity in mice. *Journal of Ethnopharmacology*, Vol 127, No.2, pp. 528- 533, ISSN 0378-8741

- Zafar, R. & Ali, S. M. (1998). Anti-hepatotoxic effects of root and root callus extracts of *Cichorium intybus* L. *Journal of Ethnopharmacology*, Vol 63, No.3, pp. 227-231, ISSN 0378-8741
- Zhou, D.; Ruan, J.; Cai, Y.; Xiong, Z.; Fu, W. & Wei, A. (2010). Antioxidant and hepatoprotective activity of ethanol extract of *Arachniodes exilis* (Hance) Ching. *Journal of Ethnopharmacology*, Vol 129, No.2, pp. 232- 237, ISSN 0378-8741
- Zimmerman, H. J. & Seeff, L. B. (1970). Enzymes in hepatic disease. In: *Diagnostic Enzymology*. E. I. Goodly (Ed.), 1, Lea and Febiger, Philadelphia, USA.



## **Phytochemicals as Nutraceuticals - Global Approaches to Their Role in Nutrition and Health**

Edited by Dr Venketeshwer Rao

ISBN 978-953-51-0203-8

Hard cover, 278 pages

**Publisher** InTech

**Published online** 23, March, 2012

**Published in print edition** March, 2012

Phytochemicals are biologically active compounds present in plants used for food and medicine. A great deal of interest has been generated recently in the isolation, characterization and biological activity of these phytochemicals. This book is in response to the need for more current and global scope of phytochemicals. It contains chapters written by internationally recognized authors. The topics covered in the book range from their occurrence, chemical and physical characteristics, analytical procedures, biological activity, safety and industrial applications. The book has been planned to meet the needs of the researchers, health professionals, government regulatory agencies and industries. This book will serve as a standard reference book in this important and fast growing area of phytochemicals, human nutrition and health.

### **How to reference**

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Saleh I. Alqasoumi and Maged S. Abdel-Kader (2012). Screening of some Traditionally Used Plants for Their Hepatoprotective Effect, *Phytochemicals as Nutraceuticals - Global Approaches to Their Role in Nutrition and Health*, Dr Venketeshwer Rao (Ed.), ISBN: 978-953-51-0203-8, InTech, Available from:

<http://www.intechopen.com/books/phytochemicals-as-nutraceuticals-global-approaches-to-their-role-in-nutrition-and-health/review-screening-of-some-traditionally-used-plants-for-their-hepatoprotective-effect>

**INTECH**  
open science | open minds

### **InTech Europe**

University Campus STeP Ri  
Slavka Krautzeka 83/A  
51000 Rijeka, Croatia  
Phone: +385 (51) 770 447  
Fax: +385 (51) 686 166  
[www.intechopen.com](http://www.intechopen.com)

### **InTech China**

Unit 405, Office Block, Hotel Equatorial Shanghai  
No.65, Yan An Road (West), Shanghai, 200040, China  
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元  
Phone: +86-21-62489820  
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen