

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

185,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Mucosal Immunity and Evasion Strategies of *Neisseria gonorrhoeae*

Mónica Imarai¹, Claudio Acuña¹, Alejandro Escobar²,
Kevin Maisey¹ and Sebastián Reyes-Cerpa¹

¹Department of Biology, Faculty of Chemistry and Biology,
Universidad de Santiago de Chile

²Department of Basic Sciences, Faculty of
Dentistry, University of Chile

^{1,2}Chile

1. Introduction

Neisseria gonorrhoeae, the gonococcus, is a gram-negative diplococcus which causes the sexually transmitted disease gonorrhea (Figure 1). The contagious nature of gonococcal infection remains a major global health problem and represents 88 million new cases every year (WHO, 2011). *N. gonorrhoeae* is transmitted by human to human contact and is highly adapted to the genital tract, surviving poorly outside the human body. However, gonococcus develops resistance to antimicrobials, antigenic variability and mechanisms of immune evasion by which it evades host defenses, thus persisting and often causing undetected asymptomatic infection (Tapsall, 2001).




Fig. 1. *Neisseria gonorrhoeae*. (A) Colonies on agar, (B) Gram-staining, (C) Transmission electron microscopy, (D) Confocal microscopy of the bacteria (in green).

The symptoms of gonorrhoea are similar to those caused by other agents, most notably *Chlamydia trachomatis*. *N. gonorrhoeae* causes infections mainly of the urethra in men and the endocervix in women, although it may also infect extragenital mucosal sites, including the oropharynx and anorectum. Ocular infections also occur, and in neonates could cause blindness (Tapsall, 2001). Genital infection in men usually causes urethritis and epididimitis with purulent urethral exudates (Apicella et al., 1996), however an important proportion of infected men never develop symptoms and more than half of infected women develop an asymptomatic silent infection (Farley et al., 2003; Handsfield et al., 1974; John and Donald, 1978). Genital tract gonorrhea gives rise to well recognized complications including upper reproductive tract infections in women, such as pelvic inflammatory disease (PID), a condition that affects between 10% and 20% of infected women in the third world. PID encompasses a range of inflammatory conditions of the upper reproductive tract and has several potential sequelae including infertility, ectopic pregnancy, among others (Hoyme, 1990; Tapsall, 2001).

The purulent exudates produced by infected men and the cervical secretions of women with gonorrhoeae contain bacteria attached to and within polymorphonuclear leukocytes (PMNs) (Apicella et al., 1996; Ovcinnikov and Delektorskij, 1971), which are the primary innate immune responders to bacterial and fungal infection and are capable of phagocytosing and killing a variety of microorganisms (Borregaard, 2010). Yet, in spite of the numerous PMNs at the site of gonorrheal infection, viable gonococci can be cultured from the exudates of infected individuals suggesting that the PMNs driven innate immune response to *N. gonorrhoeae* are ineffective at clearing a gonorrheal infection (Johnson and Criss, 2011). Considering further that the humoral immune response against *N. gonorrhoeae* is extremely low in serum and in secretions of the human (male and female) during infection (Hedges et al., 1999; Hedges et al., 1998), the persistence of gonococcus can be explained by the presence of an ineffective immune response which facilitates the long-term colonization of hosts, creating enhanced opportunity for dissemination and transmission of gonorrhea.

The aim of this chapter is to review advances in our understanding of the immunity against *N. gonorrhoeae* and those mechanisms of evasion that seem to be responsible for the restricted immune response frequently observed.

2. Gonococcal membrane proteins and early steps of infection

Infection of genital epithelial cells by *N. gonorrhoeae* is a multi-step process, consisting of adherence, invasion, intracellular survival and exocytosis. These events are initiated and mediated by multiple interactions between gonococcal surface molecules and epithelial cells. These interactions activate signalling cascades in host cells and trigger the reorganization of the actin cytoskeleton, which is required for the entry of the bacteria into the cells. Pilus retraction from adherent gonococci on the epithelial cell surface activates calcium flux (Ayala et al., 2005), the PI3K/ Akt pathway (Lee et al., 2005) and the MAP kinase ERK pathway (Howie et al., 2008). The interaction of opacity protein (Opa) with heparin sulfate proteoglycans (HSPG) activates phosphatidylcholine-specific phospholipase C (PLC) and the acid sphingomyelinase (Grassme et al., 1997). Opa also can trigger integrin-mediated protein kinase C (PKC) activation through binding to the serum-derived extracellular matrix proteins, fibronectin and vitronectin (Dehio et al., 1998a; Dehio et al., 1998b; Dehio et al., 1998c).

2.1 Type IV pili

Although there are many potential cell surface proteins of *N. gonorrhoeae* having a role in cell-host interaction, attention of most studies has focused on a few of them. This is the case of the gonococcal type IV pili which are composed of a major structural subunit, the pilin or PilE protein, assembled into helical pilus fibers (Parge et al., 1995). *In vitro* studies indicate that antigenic variation of this protein can affect pilus-mediated adherence to human cells (Jonsson et al., 1994; Long et al., 1998; Rudel et al., 1992). The binding of *N. gonorrhoeae* pili to host cells is thought to involve the complement regulatory protein CD46 (membrane cofactor receptor) (Kallstrom et al., 2001). In fact, the association of pili with CD46 in cervical carcinoma cells results in a cytoplasmic calcium flux derived from intracellular calcium stores (Kallstrom et al., 1998). However, not all studies support this role for CD46. Accordingly, no binding of piliated gonococci was observed on CD46-transfected cells and furthermore, specific down-regulation of CD46 expression in human epithelial cell lines did not alter the binding of piliated gonococci (Kirchner et al., 2005; Kirchner and Meyer, 2005; Tobiasson and Seifert, 2001). Thus the topic remains controversial. There are other potential receptors for pili attachment, i.e., integrins containing a domain known as I-domain. Edwards et al. demonstrated that pili bind to I-domain-containing integrins on primary cell cultures derived from cervical and male urethral epithelia (Edwards and Apicella, 2005; Edwards et al., 2002). In this regard, the complement receptor 3 (CR3; integrin $\alpha M\beta 2$ or CD11b/CD18) serves as the key receptor mediating gonococcal adherence to human cervical epithelia, *in vivo*, as well as *ex vivo* (Figure 2) (Edwards et al., 2001).




Fig. 2. Adhesion and internalization of *N. gonorrhoeae* in the host cell.

Pilus engagement has also been demonstrated to play a role in host cell cytoskeletal rearrangements (Edwards et al., 2000; Grassme et al., 1996; Griffiss et al., 1999; Merz et al., 1999; Merz and So, 1997). Membrane ruffles are induced in response to gonococci; epithelial cell invasion occur primarily in an actin-dependent manner, but it does not appear to require *de novo* protein synthesis by either the bacterium or the host cervical cell, eliciting substantial rearrangements of filamentous actin in the host cell cortex directly beneath sites of bacterial contact (Merz et al., 1999). Engagement of CR3 on primary cervical epithelial cells results in vinculin- and ezrin-enriched focal complex formation before membrane ruffle formation, bacteria reside within macropinosomes, and an accumulation of actin-associated protein occur in response to gonococcal infection (Edwards et al., 2000). Finally a signal transduction cascade that is dependent upon the activation of phosphatidylinositol 3-kinase or MAP kinases and Rho GTPases is activated (Edwards et al., 2001).

2.2 Opacity-associated (Opa) adhesin proteins

Opacity-associated adhesion proteins located in the gonococcal outer membrane facilitate the interaction of bacteria with a number of host cell types, including epithelial cells on mucosal surfaces and various immune cells, indicating a direct effect on the immune response. Receptor tropism of Opa proteins can be broadly divided into two categories, those that bind to members of the carcinoembryonic antigen cell adhesion molecule (CEACAM) family (Figure 2) and those that bind to HSPGs (Bos et al., 1997; Chen et al., 1995; Chen and Gotschlich, 1996; Chen et al., 1997; Gray-Owen et al., 1997a; Gray-Owen et al., 1997b; Popp et al., 1999; van Putten and Paul, 1995; Virji et al., 1999; Virji et al., 1996a; Virji et al., 1996b). These categories are represented by Opa50, i.e., Opa proteins that recognize HSPG, and Opa52, i.e., Opa proteins that recognize members of CEACAM family. Vitronectin and fibronectin function as required bridging molecules between the gonococcus and its target HSPG receptor(s) (Duensing and Putten, 1998; Duensing and van Putten, 1997; Gomez-Duarte et al., 1997; van Putten et al., 1998). Association with an integrin coreceptor ($\alpha 3$ or $\alpha 5$ for vitronectin-mediated adherence or $\alpha 1$ for fibronectin-mediated adherence) triggers a signaling cascade within the target cell that is dependent upon the activation of PKC (Dehio et al., 1998c).

The binding sites of Opa proteins reside on the amino-terminal domains of the CEACAM family, which are largely conserved and therefore allow one or more Opa proteins to target distinct CEACAMs (Chen and Gotschlich, 1996; Muenzner et al., 2000; Virji et al., 1996b). As CEACAMs may contain immunoreceptor tyrosine-based inhibitory motifs (ITIMs) or immunoreceptor tyrosine-based activation motifs (ITAMs) (Hammarstrom, 1999), the consequences of downstream signaling following bacterial ligation depend on the receptor and target cell involved. From studies so far, it can be concluded that Opa-CEACAM interactions result in cellular invasion (Muenzner et al., 2000; Virji et al., 1999). The CEACAM receptor-mediated phagocytosis of Opa(52)-expressing *N. gonorrhoeae* results in a rapid activation of the acid sphingomyelinase. Furthermore, the CEACAM receptor-initiated stimulation activates a cascade via Src-like protein tyrosine kinases, Rac1 and PAK to Jun-N-terminal kinases (Hauck et al., 2000; Hauck et al., 1998).

The gonococcal opacity proteins are a well-studied example of phase-variable surface structures. Gonococcal strains express several antigenically distinct Opa proteins that are encoded by separate chromosomal alleles (Bhat et al., 1991; Connell et al., 1990; Dempsey et

al., 1991). Each *opa* gene undergoes phase variation via frameshift mutations that cause changes in pentameric repeats in the *opa* structural gene (Murphy et al., 1989). Bacteria that express no Opa proteins, bacteria that express one Opa protein, and bacteria that express multiple Opa proteins simultaneously result from these reversible mutations. This is one of the earliest described mechanisms of immune evasion found in gonococcus.

2.3 Porin, the outer membrane protein channel

Porin, is membrane channel through which small molecules traverse the gonococcal outer membrane, is thought to play multiple roles in potentiating disease caused by *N. gonorrhoeae*. Porin molecules trigger different responses within host cells depending upon the particular porin and the host cell type. A feature of gonococcal porin is its ability to translocate into eukaryotic cell membranes (Lynch et al., 1984; Weel and van Putten, 1991), where it acts as voltage-gated channel that is modulated by host cell ATP and GTP (Rudel et al., 1996). It has also been demonstrated that *N. gonorrhoeae* infection of epithelial cells results in selective porin transport to the mitochondria (Muller et al., 2000; Muller et al., 2002). Within the cell mitochondrial membrane, porin initiates apoptosis by inducing a calcium influx and, consequently, calpain and caspase activity within these cells (Muller et al., 1999). In contrast to the porin-induced apoptosis observed in HeLa cells, gonococcal infection of primary human male urethral epithelial cells results in antiapoptotic events (Binnicker et al., 2003; Binnicker et al., 2004). Finally, porin also acts as an actin-nucleating protein in epithelial cells (Wen et al., 2000) facilitating the cytoskeletal rearrangements required for actin-mediated entry of the gonococcus into its target host cell.

2.4 Lipooligosaccharide (LOS)

Lipooligosaccharide is a major antigenic and immunogenic component of pathogenic *Neisseria* species. LOS produced by these bacteria consists of an oligosaccharide (OS) moiety and lipid A, and structural variation of the OS leads to LOS heterogeneity (Preston et al., 1996). Studies using primary male urethral epithelial cells demonstrated an association between the urethral epithelium and the gonococcus through the interaction of the asialoglycoprotein receptor (ASGP-R) and gonococcal LOS (Harvey et al., 2001). In primary cell culture, engagement of the ASGP-R by the gonococcus results in pedestal formation beneath the bacterium (Harvey et al., 1997). Pedestal formation is also observed in microscopic analysis of exudates collected from men with naturally acquired gonococcal urethritis (Apicella et al., 1996; Harvey et al., 1997). Evidence suggests that endocytosis occurs primarily by actin-dependent (Giardina et al., 1998) and clathrin-dependent processes (Harvey et al., 1997).

3. Immune response against *N. gonorrhoeae*

The immunity against bacterial infections is achieved by many levels of defense that are triggered depending on the type, number and virulence of the bacteria that enter to the body (Figure 3). If the infection is mild, the tissue phagocytes are able to kill the bacteria in the phagolysosomes by reactive oxygen and nitrogen intermediates and proteolytic enzymes. In addition, these cells are able to secrete pro-inflammatory cytokines such as interleukin (IL)-1, IL-6, IL-8 and Tumoral Necrosis Factor (TNF) which initiates

inflammatory response. On the other hand, antigen processing and presentation, which is a crucial biological process for the initiation of adaptive immune response, might initiate antigen-specific immune response and a long lasting memory. Dendritic cells (DCs) are a subgroup of high-power antigen presenting cells (APC) with a unique ability to attract and interact with naive T cells to induce a primary immune response (Sabatos et al., 2008; Verhagen et al., 2008). Immature DCs (iDCs) reside in most peripheral tissues where they act as sentinels for incoming pathogens (Rowell and Wilson, 2009). After exposure to pro-inflammatory cytokines and microbial products, iDCs undergo a process termed maturation, which involve up-regulation of MHC molecules, co-stimulatory molecules, secretion of pro-inflammatory cytokines and migration to draining lymph node where mature DCs activate T cells (Xu et al., 2007). Following antigen stimulation by DCs, T cells begin an expansion process, as a result of extensive division. Under a control of DCs, helper T cells (Th) acquire the ability to respond to infection through the production of powerful cytokines like interferon gamma (IFN- γ), which is able to activate macrophages resisting infections by facultative and obligate intracellular microbes (Th1 cells) (Napolitani et al., 2005), or IL-17, which mobilize phagocytes at body surfaces to resist extracellular bacteria (Th17 cells) (LeibundGut-Landmann et al., 2007). Th17 cells represent a distinct lineage that originates mainly in the presence of TGF- β 1 and IL-6 and need the presence of IL-23 for their expansion and/or maintenance (Annunziato et al., 2007). IL-23 is secreted by macrophages and DCs in response to microbial products and inflammatory cytokines (Langrish et al., 2004). Once differentiated, Th17 cells are able to secrete preferentially IL-17A, IL-17F and IL-22, a particular set of cytokines not secreted by the other Th cells (Ouyang et al., 2008). IL-17 plays a particularly significant role in regulating neutrophils (PMNn) recruitment and granulopoiesis via the production of IL-8 and MIP-2 (CXCL2) (Laan et al., 1999). In early stages of infections, interactions between *N. gonorrhoeae* and epithelial mucosa trigger immune response with release of IL-1, IL-6, IL-8 and TNF which serve to recruit and activate PMNn to the site of infection and promoting their bactericidal activity limiting bacterial penetration into submucosal tissues (Fisette et al., 2003; Maisey et al., 2003; McGee et al., 1999). In men, PMNs appear in urethral swabs and urine several days after infection and immediately prior to the onset of symptoms (Cohen and Cannon, 1999). In women with gonorrhea, the cervical secretions also contain PMNn (Evans, 1977) and bacteria are attached to and within PMNn (Johnson and Criss, 2011). In spite of the numerous PMNn at the site of gonorrheal infection, viable gonococci can be cultured from the exudates of infected individuals (Wiesner and Thompson, 1980). When bacteria crosses the layer of epithelial cells, obtaining access to submucosa, they have the first encounter with macrophages and DCs (Rarick et al., 2006). DCs interaction with *N. gonorrhoeae* surface factors like Pili and Opa is mediated by members of the CD66 family, CD46 and CR3. However, C-type lectins (macrophage galactose-type lectin -MLG- and DC-SIGN) constitute the main DCs expressed receptor for *N. gonorrhoeae* (Astier et al., 2010).

Most studies that have investigated antigenococcal immune responses have focused predominantly on humoral responses (Table 1). The hallmark of humoral immune response against *N. gonorrhoeae* is the extremely low levels of anti-gonococcal antibodies found in serum and secretions of the human (male and female) during infection (Hedges et al., 1999; Hedges et al., 1998), indicating that humoral immunity against gonococcus is highly limited (Hedges et al., 1999; Song et al., 2008). Antigenic heterogeneity is a major consideration for humoral immunity in gonococcal infection studies. Meanwhile Pili, protein I (PI), protein II

(PII), H.8, IgA1 protease and LOS are quantitatively the most important antigens in generating antibody responses in gonococcal infection. It is clear that patients make antibodies against the pili of the infecting gonococcal strain. In women, pili appeared to be the predominant antigen in the immune response unlike men, that have higher levels of antibodies to other antigens than pili (Brooks and Lammel, 1989). Thus, antibodies generated are directed against other major membrane molecules, such as Opa proteins and Porin protein (Brooks and Lammel, 1989; Hedges et al., 1999; Hedges et al., 1998; Plummer et al., 1993; Plummer et al., 1994; Zheng, 1997). Although some of these have bactericidal activity, they are not protective and seem to be blocked by immunoglobulins against the outer membrane protein 3 (RmP). In fact, women with these antibodies were at increased risk to gonococcal infection (Plummer et al., 1993). The absence of induction of humoral response results on limited or no protection against re-infection with *N. gonorrhoeae* despite the generation of serum antibody responses against antigens produced by several prototype gonococcal vaccines (Boslego et al., 1991; McChesney et al., 1982; Tramont et al., 1981).




Fig. 3. Protective immune responses against intracellular bacteria.

Antigen	Effects on immune response	References
<i>N. gonorrhoeae</i>	IgG>IgM and sIgA>IgA	(Tapchaisri and Sirisinha, 1976)
Outer membrane protein I	IgA>IgG	(Jeurissen et al., 1987)
Outer membrane protein I	IgA>IgG	(Kohl et al., 1992)
Porin	Up-regulated B7-2 expression of B lymphocytes	(Wetzler et al., 1996)
Heat labile and fixed <i>N. gonorrhoeae</i>	IgG>IgA>IgM serum	(Cohen, 1967)
Pili and outer membrane protein II	IgG serum	(Lammel et al., 1985)
Outer membrane protein II	IgA serum	(Lammel et al., 1985)
Outer membrane protein I and II and pili	IgG and IgA vaginal fluid	(Lammel et al., 1985)
LPS	IgG>IgA in serum	(Ison et al., 1986)
Outer membrane protein III	IgG in serum	(Rice et al., 1986)
<i>N. gonorrhoeae</i> MS11	Slight increase of IgG and IgA in serum	(Hedges et al., 1999)
<i>N. gonorrhoeae</i> mice infection	Increased of neutrophils and macrophage in site of infection	(Song et al., 2008)
<i>N. gonorrhoeae</i> mice infection	IgG>IgA>IgM in vaginal fluid	(Song et al., 2008)
Whole-protein extractes of <i>N. gonorrhoeae</i> P9-17	Low titers of IgG in serum	(Imarai et al., 2008)
<i>N. gonorrhoeae</i>	Th17 profile immune response and increased IL-17	(Feinen et al., 2010)
IgA1 protease	Th1 pro-inflammatory profile with increased IFN γ and TNF α	(Tsirpouchtsidis et al., 2002)
IgA1 protease	IL-1 β , IL-6, TNF α and IL-8	(Lorenzen et al., 1999)
Porin	Th2 pro-inflammatory profile with increased of IL-4	(Simpson et al., 1999)
Porin	B lymphocytes activation and up-regulation of CD40	(Snapper et al., 1997)
Pili	Activation and proliferation of CD4 ⁺ T Lymphocyte	(Plant and Jonsson, 2006)
Opa	Arrest of activation and proliferation of CD4 ⁺ T lymphocytes binding CEACAM1	(Boulton and Gray-Owen, 2002)

Table 1. Anti-gonococcus immune responses.

Similarly, in the murine model of infection, no antibody induced response has been detected. In Balb/c mice, *N. gonorrhoeae* is able to reach the upper genital organs and to invade uterine tissue, as it occurs in humans (Imarai et al., 2008; Jerse, 1999). Interestingly, studies in this model revealed several features of infection that mimic a good spectrum of the characteristics of the human disease. Some of them seem highly valuable, (i) the bacteria introduced in the vagina not only invade and colonize the lower genital tract, but also spread to the upper organs (uterus and oviducts) (Inaba et al., 1992), (ii) infection occurs with no signs of the disease, just as it occurs in a high percentage of women during natural infection (Farley et al., 2003; Inaba et al., 1992) and (iii) the bacteria remains alive within the internal compartments of the infected cells, invading uterine tissues up to 22 days post inoculation (Imarai et al., 2008).

On the other hand, CD4⁺ T cell responses also occur with gonococcal antigens. Antigenic stimulation with Porin induces an increase in the percentage of IL-4 producing CD4⁺ T cells, but no production of other cytokines such as IL-2, IL-10, IFN- γ , or TNF occurs, indicating that infected individuals produce a Th2 response against Porin (Simpson et al., 1999).

Moreover, gonococcal pili interaction with CD4⁺ T cells induces the activation and proliferation of lymphocytes and stimulates the secretion of IL-10 (Plant and Jonsson, 2006). In contrast, Opa proteins mediate binding to CEACAM-1 expressed by CD4⁺ T cells and suppress activation and proliferation of naive lymphocytes (Boulton and Gray-Owen, 2002; Lee et al., 2008). Rarick et al. showed a differential activation of peripheral blood mononuclear cells by *N. gonorrhoeae* that includes the early induction (96 hours) of IL-2, IL-12 and IFN- γ (Th1 cytokines), IL-4 (Th2 cytokine), IL-10 (immunosuppressive cytokine) and MCP-1, MCP-2 (chemotactic cytokines). The cytokine response observed in this study indicate that distinct gonococcal components produce antagonistic signaling and cytokines suggesting that *N. gonorrhoeae* infection cannot be initially categorized as Th1 or Th2 response (Rarick et al., 2006). Recent studies in the murine model of gonococcal genital tract infection showed that Th17 cells are involved in the immune response to *N. gonorrhoeae*. This response leads to IL-17 dependent secretion of IL-6, LIX and MIP2 α and subsequent recruitment of PMNn, which is delayed in the presence of IL-17A blocking antibodies or deletion of IL-17RA prolonging infection (Feinen et al., 2010).

Overall, most known gonococcal antigens are able to induce humoral and cellular immune responses in the human, but a protective immune response has not been identified. Data suggest that the bacteria have mechanisms to evade destruction by PMNn, antibodies and T cells.

4. Mechanisms of immune evasion by *Neisseria gonorrhoeae*

Several studies suggest the promotion of direct inhibitory mechanisms by gonococcus to escape the immune response. Data indicate that bacteria induces transient reduction of CD4⁺ and CD8⁺T cells during acute gonococcal infection in HIV-1-positive woman, with increase in plasma HIV-1 RNA copy number and plasma concentration of IL-4, IL-6, IL-10 and TNF- α soluble receptor (Anzala et al., 2000). Gonococcal infection also correlates with suppression of activation and proliferation of CD4⁺ T lymphocytes (Boulton and Gray-Owen, 2002). In this respect, inhibitory effect is mediated by the ligation of gonococcus membrane component Opa52 present in blebs of outer membrane (OMV) to CEACAM1 (also known as CD66a or BGP) on lymphocyte (Lee et al., 2007). The interaction triggers the

phosphorylation of the ITIM impeding the normal expression of early activation marker CD69 and the subsequent proliferation of T cells. This strategy represents an effective means by which to create a “zone of immunosuppression” surrounding the infected site (Lee et al., 2008). Opa-CEACAM1-induced immunosuppression might also control the development of a humoral response, decreasing the T cell help for B cell activation or targeting the CEACAM1-expressing B cells causing cell death (Pantelic et al., 2005) and subsequent inhibition of antibody production.

On the other hand, piliated gonococci enhances T-cell activation and proliferation, regardless of whether this is mediated by the pilus itself or is due to the act of binding to a pilus receptor such as CD46 or integrins. Upon pili-CD46 ligation, the IL-10-secreting type 1 regulatory T (Tr1) cells are elicited (Plant and Jonsson, 2006). Tr1 cells are able to suppress the activation of bystander T cells via induction of IL-10 (Jonuleit et al., 2001). In addition, IL-10 suppresses the production of pro-inflammatory cytokines thereby inhibiting the ability of APC to induce differentiation of Th1 cells, just the type of response associated to protection against intracellular pathogens such as gonococcus (McGuirk et al., 2002).

In regard to the role of regulatory T (Treg) cells, we showed that in Balb/c, the mouse model of gonococcus infections, *N. gonorrhoeae* induces transforming growth factor β 1 (TGF- β 1)+ CD4⁺ T cells in the mucosal lymph nodes, including a subset of CD25⁺Foxp3⁺ Tregs, indicating that type of response may avoid the host mechanisms of protection (Imarai et al., 2008). Mature CD4⁺ CD25⁻ T cells can be converted in Treg cells in peripheral tissues under immunosuppressive conditions, such as exposure to IL-10 or TGF- β 1 made by antigen presenting cells (Sakaguchi, 2004). Treg cells inhibit the function of effectors T cells through the secretion of suppressive cytokines IL-10 and TGF- β 1, or by a cell contact-dependent mechanism via CTLA4 or GITR molecules (Bending et al., 2009; O'Connor et al., 2008; Sakaguchi, 2004). Considering that, the source of TGF- β 1 could be stromal and epithelial cells of reproductive organs (Nocera and Chu, 1995; Srivastava et al., 1996; Taylor et al., 2000), which are target of gonococcus infection, it is possible that the cytokine milieu found in reproductive tract subsidized the inductions of Tregs by *N.gonorrhoeae*. In fact, we must consider that the genital tract is an immune-privileged site with expression of regulatory cytokines which might induce a tolerogenic response against gonococcus (Nocera and Chu, 1995; Srivastava et al., 1996; Taylor et al., 2000). For instance, epithelia and stromal cells of the reproductive organs of the mouse and human, express high levels of TGF- β 1 and other molecules involved in conditioning immune privilege sites (Chegini et al., 1994; Grant and Wira, 2003; Jin et al., 1992; Wada et al., 1996). Moreover, antigen presenting cells such as macrophages and DCs, regularly present in the reproductive tissues might also contribute to regulatory response, since they could promote T regulatory cells differentiation through production of IL-10 and TGF- β 1 after infection (Givan et al., 1997; Stagg et al., 1998).

The effects of interaction of different stable gonococcal LOS phenotypes with human DCs were evaluated (van Vliet et al., 2009). Interestingly, this study revealed that LOS variants result in alterations of cytokine secretion profiles of DCs and in the induction of distinct adaptive CD4⁺ T helper responses. Gonococcus significantly increased IL-10 production, as well as pro-inflammatory cytokines, such as TNF- α , IL-6, IL-8 and IL-12. However, only IL-10 production was modulated by LOS variation. Supporting the anti-inflammatory or regulatory effects of gonococcus on APCs, recently we found that the bacterium was unable to induce significant up-regulation of cell surface co-stimulatory molecule CD86 in

macrophages, as well in DCs (unpublished data). This suggests that, although *N. gonorrhoeae* is actually phagocytosed by macrophages and DCs, the bacteria can deteriorate antigen presenting function. Moreover, gonococcus was unable to induce up-regulation of MHC class II in macrophages, this molecule is involved in antigen processing and presentation, therefore this implies that macrophages do not have a proper antigen presenting function. In addition, gonococcus induced IL-10 and TGF- β 1 in dendritic cells reaching an anti-inflammatory or regulatory response.

Altogether, studies so far show that *N. gonorrhoeae* might control immune response by inducing (1) suppression of activation and proliferation of CD4⁺ T lymphocytes, (2) secretion of tolerogenic-type cytokines, (3) inhibitory T cells (Tr1 and Tregs), and (4) by deteriorating antigen presenting function. This wide variety of immune evasion mechanism may explain the frequent appearance of persistent and asymptomatic infection (Figure 4).




Fig. 4. *N. gonorrhoeae* interferes with immune responses.

5. Conclusion

Neisseria gonorrhoeae infects the reproductive tract of the human causing gonorrhea. Mechanisms of infection involved the early attachment of gonococcal membrane components, such as Pili, Opa, Porin and LOS, to cell host receptors. Frequently, the bacteria develop persistence and asymptomatic disease, which seem to be associated to the gonococcus ability to evade immune response. Several lines of research revealed that *N.*

gonorrhoeae is able to subvert polymorphonuclear cell phagocytic destruction, induces a weak humoral immune response, decreases antigen presenting properties of macrophages and dendritic cells, inhibits T cell proliferation and induces anti-inflammatory cytokines and T inhibitory cells. On the whole, this seems to be a complex network of immune evasion mechanisms responsible for the restricted immune response frequently observed during gonococcal infection.

6. Acknowledgment

We thanks the support of DICYT 021043IB, PBCT to CA and CONICYT Fellowship to KM and SR.

7. References

- Annunziato F., Cosmi L., Santarlasci V., Maggi L., Liotta F., Mazzinghi B., Parente E., Fili L., Ferri S., Frosali F., Giudici F., Romagnani P., Parronchi P., Tonelli F., Maggi E. and Romagnani S., 2007. Phenotypic and functional features of human Th17 cells. *J Exp Med* 204, 1849-61.
- Anzala A. O., Simonsen J. N., Kimani J., Ball T. B., Nagelkerke N. J., Rutherford J., Ngugi E. N., Bwayo J. J. and Plummer F. A., 2000. Acute sexually transmitted infections increase human immunodeficiency virus type 1 plasma viremia, increase plasma type 2 cytokines, and decrease CD4 cell counts. *J Infect Dis* 182, 459-66.
- Apicella M. A., Ketterer M., Lee F. K., Zhou D., Rice P. A. and Blake M. S., 1996. The pathogenesis of gonococcal urethritis in men: confocal and immunoelectron microscopic analysis of urethral exudates from men infected with *Neisseria gonorrhoeae*. *J Infect Dis* 173, 636-46.
- Astier A. L., Beriou G., Eisenhaure T. M., Anderton S. M., Hafler D. A. and Hacohen N., 2010. RNA interference screen in primary human T cells reveals FLT3 as a modulator of IL-10 levels. *J Immunol* 184, 685-93.
- Ayala P., Wilbur J. S., Wetzler L. M., Tainer J. A., Snyder A. and So M., 2005. The pilus and porin of *Neisseria gonorrhoeae* cooperatively induce Ca(2+) transients in infected epithelial cells. *Cell Microbiol* 7, 1736-48.
- Bending D., De La Pena H., Veldhoen M., Phillips J. M., Uyttenhove C., Stockinger B. and Cooke A., 2009. Highly purified Th17 cells from BDC2.5NOD mice convert into Th1-like cells in NOD/SCID recipient mice. *J Clin Invest* 119, 565-72.
- Bhat K. S., Gibbs C. P., Barrera O., Morrison S. G., Jahnig F., Stern A., Kupsch E. M., Meyer T. F. and Swanson J., 1991. The opacity proteins of *Neisseria gonorrhoeae* strain MS11 are encoded by a family of 11 complete genes. *Mol Microbiol* 5, 1889-901.
- Binnicker M. J., Williams R. D. and Apicella M. A., 2003. Infection of human urethral epithelium with *Neisseria gonorrhoeae* elicits an upregulation of host anti-apoptotic factors and protects cells from staurosporine-induced apoptosis. *Cell Microbiol* 5, 549-60.
- Binnicker M. J., Williams R. D. and Apicella M. A., 2004. Gonococcal porin IB activates NF-kappaB in human urethral epithelium and increases the expression of host antiapoptotic factors. *Infect Immun* 72, 6408-17.
- Borreagaard N., 2010. Neutrophils, from marrow to microbes. *Immunity* 33, 657-70.

- Bos M. P., Grunert F. and Belland R. J., 1997. Differential recognition of members of the carcinoembryonic antigen family by Opa variants of *Neisseria gonorrhoeae*. *Infect Immun* 65, 2353-61.
- Boslego J. W., Tramont E. C., Chung R. C., McChesney D. G., Ciak J., Sadoff J. C., Piziak M. V., Brown J. D., Brinton C. C., Jr., Wood S. W. and et al., 1991. Efficacy trial of a parenteral gonococcal pilus vaccine in men. *Vaccine* 9, 154-62.
- Boulton I. C. and Gray-Owen S. D., 2002. Neisserial binding to CEACAM1 arrests the activation and proliferation of CD4+ T lymphocytes. *Nat Immunol* 3, 229-36.
- Brooks G. F. and Lammel C. J., 1989. Humoral immune response to gonococcal infections. *Clin Microbiol Rev* 2 Suppl, S5-10.
- Cohen I. R., 1967. Natural and immune human antibodies reactive with antigens of virulent *Neisseria gonorrhoeae*: immunoglobulins G, M, And A. *J Bacteriol* 94, 141-8.
- Cohen M. S. and Cannon J. G., 1999. Human experimentation with *Neisseria gonorrhoeae*: progress and goals. *J Infect Dis* 179 Suppl 2, S375-9.
- Connell T. D., Shaffer D. and Cannon J. G., 1990. Characterization of the repertoire of hypervariable regions in the Protein II (opa) gene family of *Neisseria gonorrhoeae*. *Mol Microbiol* 4, 439-49.
- Chegini N., Zhao Y., Williams R. S. and Flanders K. C., 1994. Human uterine tissue throughout the menstrual cycle expresses transforming growth factor-beta 1 (TGF beta 1), TGF beta 2, TGF beta 3, and TGF beta type II receptor messenger ribonucleic acid and protein and contains [125I]TGF beta 1-binding sites. *Endocrinology* 135, 439-49.
- Chen T., Belland R. J., Wilson J. and Swanson J., 1995. Adherence of pilus- Opa+ gonococci to epithelial cells in vitro involves heparan sulfate. *J Exp Med* 182, 511-7.
- Chen T. and Gotschlich E. C., 1996. CGM1a antigen of neutrophils, a receptor of gonococcal opacity proteins. *Proc Natl Acad Sci U S A* 93, 14851-6.
- Chen T., Grunert F., Medina-Marino A. and Gotschlich E. C., 1997. Several carcinoembryonic antigens (CD66) serve as receptors for gonococcal opacity proteins. *J Exp Med* 185, 1557-64.
- Dehio C., Freissler E., Lanz C., Gomez-Duarte O. G., David G. and Meyer T. F., 1998a. Ligation of cell surface heparan sulfate proteoglycans by antibody-coated beads stimulates phagocytic uptake into epithelial cells: a model for cellular invasion by *Neisseria gonorrhoeae*. *Exp Cell Res* 242, 528-39.
- Dehio C., Gray-Owen S. D. and Meyer T. F., 1998b. The role of neisserial Opa proteins in interactions with host cells. *Trends Microbiol* 6, 489-95.
- Dehio M., Gomez-Duarte O. G., Dehio C. and Meyer T. F., 1998c. Vitronectin-dependent invasion of epithelial cells by *Neisseria gonorrhoeae* involves alpha(v) integrin receptors. *FEBS Lett* 424, 84-8.
- Dempsey J. A., Litaker W., Madhure A., Snodgrass T. L. and Cannon J. G., 1991. Physical map of the chromosome of *Neisseria gonorrhoeae* FA1090 with locations of genetic markers, including opa and pil genes. *J Bacteriol* 173, 5476-86.
- Duensing T. D. and Putten J. P., 1998. Vitronectin binds to the gonococcal adhesin OpaA through a glycosaminoglycan molecular bridge. *Biochem J* 334 (Pt 1), 133-9.
- Duensing T. D. and van Putten J. P., 1997. Vitronectin mediates internalization of *Neisseria gonorrhoeae* by Chinese hamster ovary cells. *Infect Immun* 65, 964-70.

- Edwards J. L. and Apicella M. A., 2005. I-domain-containing integrins serve as pilus receptors for *Neisseria gonorrhoeae* adherence to human epithelial cells. *Cell Microbiol* 7, 1197-211.
- Edwards J. L., Brown E. J., Ault K. A. and Apicella M. A., 2001. The role of complement receptor 3 (CR3) in *Neisseria gonorrhoeae* infection of human cervical epithelia. *Cell Microbiol* 3, 611-22.
- Edwards J. L., Brown E. J., Uk-Nham S., Cannon J. G., Blake M. S. and Apicella M. A., 2002. A co-operative interaction between *Neisseria gonorrhoeae* and complement receptor 3 mediates infection of primary cervical epithelial cells. *Cell Microbiol* 4, 571-84.
- Edwards J. L., Shao J. Q., Ault K. A. and Apicella M. A., 2000. *Neisseria gonorrhoeae* elicits membrane ruffling and cytoskeletal rearrangements upon infection of primary human endocervical and ectocervical cells. *Infect Immun* 68, 5354-63.
- Evans B. A., 1977. Ultrastructural study of cervical gonorrhea. *J Infect Dis* 136, 248-55.
- Farley T. A., Cohen D. A. and Elkins W., 2003. Asymptomatic sexually transmitted diseases: the case for screening. *Prev Med* 36, 502-9.
- Feinen B., Jerse A. E., Gaffen S. L. and Russell M. W., 2010. Critical role of Th17 responses in a murine model of *Neisseria gonorrhoeae* genital infection. *Mucosal Immunol* 3, 312-21.
- Fisette P. L., Ram S., Andersen J. M., Guo W. and Ingalls R. R., 2003. The Lip lipoprotein from *Neisseria gonorrhoeae* stimulates cytokine release and NF-kappaB activation in epithelial cells in a Toll-like receptor 2-dependent manner. *J Biol Chem* 278, 46252-60.
- Giardina P. C., Williams R., Lubaroff D. and Apicella M. A., 1998. *Neisseria gonorrhoeae* induces focal polymerization of actin in primary human urethral epithelium. *Infect Immun* 66, 3416-9.
- Givan A. L., White H. D., Stern J. E., Colby E., Gosselin E. J., Guyre P. M. and Wira C. R., 1997. Flow cytometric analysis of leukocytes in the human female reproductive tract: comparison of fallopian tube, uterus, cervix, and vagina. *Am J Reprod Immunol* 38, 350-9.
- Gomez-Duarte O. G., Dehio M., Guzman C. A., Chhatwal G. S., Dehio C. and Meyer T. F., 1997. Binding of vitronectin to opa-expressing *Neisseria gonorrhoeae* mediates invasion of HeLa cells. *Infect Immun* 65, 3857-66.
- Grant K. S. and Wira C. R., 2003. Effect of mouse uterine stromal cells on epithelial cell transepithelial resistance (TER) and TNFalpha and TGFbeta release in culture. *Biol Reprod* 69, 1091-8.
- Grassme H., Gulbins E., Brenner B., Ferlinz K., Sandhoff K., Harzer K., Lang F. and Meyer T. F., 1997. Acidic sphingomyelinase mediates entry of *N. gonorrhoeae* into nonphagocytic cells. *Cell* 91, 605-15.
- Grassme H. U., Ireland R. M. and van Putten J. P., 1996. Gonococcal opacity protein promotes bacterial entry-associated rearrangements of the epithelial cell actin cytoskeleton. *Infect Immun* 64, 1621-30.
- Gray-Owen S. D., Dehio C., Haude A., Grunert F. and Meyer T. F., 1997a. CD66 carcinoembryonic antigens mediate interactions between Opa-expressing *Neisseria gonorrhoeae* and human polymorphonuclear phagocytes. *EMBO J* 16, 3435-45.

- Gray-Owen S. D., Lorenzen D. R., Haude A., Meyer T. F. and Dehio C., 1997b. Differential Opa specificities for CD66 receptors influence tissue interactions and cellular response to *Neisseria gonorrhoeae*. *Mol Microbiol* 26, 971-80.
- Griffiss J. M., Lammel C. J., Wang J., Dekker N. P. and Brooks G. F., 1999. *Neisseria gonorrhoeae* coordinately uses Pili and Opa to activate HEC-1-B cell microvilli, which causes engulfment of the gonococci. *Infect Immun* 67, 3469-80.
- Hammarstrom S., 1999. The carcinoembryonic antigen (CEA) family: structures, suggested functions and expression in normal and malignant tissues. *Semin Cancer Biol* 9, 67-81.
- Handsfield H. H., Lipman T. O., Harnisch J. P., Tronca E. and Holmes K. K., 1974. Asymptomatic gonorrhea in men. Diagnosis, natural course, prevalence and significance. *N Engl J Med* 290, 117-23.
- Harvey H. A., Jennings M. P., Campbell C. A., Williams R. and Apicella M. A., 2001. Receptor-mediated endocytosis of *Neisseria gonorrhoeae* into primary human urethral epithelial cells: the role of the asialoglycoprotein receptor. *Mol Microbiol* 42, 659-72.
- Harvey H. A., Ketterer M. R., Preston A., Lubaroff D., Williams R. and Apicella M. A., 1997. Ultrastructural analysis of primary human urethral epithelial cell cultures infected with *Neisseria gonorrhoeae*. *Infect Immun* 65, 2420-7.
- Hauck C. R., Grassme H., Bock J., Jendrossek V., Ferlinz K., Meyer T. F. and Gulbins E., 2000. Acid sphingomyelinase is involved in CEACAM receptor-mediated phagocytosis of *Neisseria gonorrhoeae*. *FEBS Lett* 478, 260-6.
- Hauck C. R., Meyer T. F., Lang F. and Gulbins E., 1998. CD66-mediated phagocytosis of Opa52 *Neisseria gonorrhoeae* requires a Src-like tyrosine kinase- and Rac1-dependent signalling pathway. *EMBO J* 17, 443-54.
- Hedges S. R., Mayo M. S., Mestecky J., Hook E. W., 3rd and Russell M. W., 1999. Limited local and systemic antibody responses to *Neisseria gonorrhoeae* during uncomplicated genital infections. *Infect Immun* 67, 3937-46.
- Hedges S. R., Sibley D. A., Mayo M. S., Hook E. W., 3rd and Russell M. W., 1998. Cytokine and antibody responses in women infected with *Neisseria gonorrhoeae*: effects of concomitant infections. *J Infect Dis* 178, 742-51.
- Howie H. L., Shiflett S. L. and So M., 2008. Extracellular signal-regulated kinase activation by *Neisseria gonorrhoeae* downregulates epithelial cell proapoptotic proteins Bad and Bim. *Infect Immun* 76, 2715-21.
- Hoyme U. B., 1990. Pelvic inflammatory disease and associated sexually transmitted diseases. *Curr Opin Obstet Gynecol* 2, 668-74.
- Imarai M., Candia E., Rodriguez-Tirado C., Tognarelli J., Pardo M., Perez T., Valdes D., Reyes-Cerpa S., Nelson P., Acuna-Castillo C. and Maisey K., 2008. Regulatory T cells are locally induced during intravaginal infection of mice with *Neisseria gonorrhoeae*. *Infect Immun* 76, 5456-65.
- Inaba K., Steinman R. M., Pack M. W., Aya H., Inaba M., Sudo T., Wolpe S. and Schuler G., 1992. Identification of proliferating dendritic cell precursors in mouse blood. *J Exp Med* 175, 1157-67.
- Ison C. A., Hadfield S. G., Bellinger C. M., Dawson S. G. and Glynn A. A., 1986. The specificity of serum and local antibodies in female gonorrhoea. *Clin Exp Immunol* 65, 198-205.

- Jerse A. E., 1999. Experimental gonococcal genital tract infection and opacity protein expression in estradiol-treated mice. *Infect Immun* 67, 5699-708.
- Jeurissen S. H., Sminia T. and Beuvery E. C., 1987. Induction of mucosal immunoglobulin A immune response by preparations of *Neisseria gonorrhoeae* porin proteins. *Infect Immun* 55, 253-7.
- Jin L. W., Inaba K. and Saitoh T., 1992. The involvement of protein kinase C in activation-induced cell death in T-cell hybridoma. *Cell Immunol* 144, 217-27.
- John J. and Donald W. H., 1978. Asymptomatic urethral gonorrhoea in men. *Br J Vener Dis* 54, 322-3.
- Johnson M. B. and Criss A. K., 2011. Resistance of *Neisseria gonorrhoeae* to neutrophils. *Front Microbiol* 2, 77.
- Jonsson A. B., Ilver D., Falk P., Pepose J. and Normark S., 1994. Sequence changes in the pilus subunit lead to tropism variation of *Neisseria gonorrhoeae* to human tissue. *Mol Microbiol* 13, 403-16.
- Jonuleit H., Schmitt E., Stassen M., Tuettenberg A., Knop J. and Enk A. H., 2001. Identification and functional characterization of human CD4(+)CD25(+) T cells with regulatory properties isolated from peripheral blood. *J Exp Med* 193, 1285-94.
- Kallstrom H., Blackmer Gill D., Albiger B., Liszewski M. K., Atkinson J. P. and Jonsson A. B., 2001. Attachment of *Neisseria gonorrhoeae* to the cellular pilus receptor CD46: identification of domains important for bacterial adherence. *Cell Microbiol* 3, 133-43.
- Kallstrom H., Islam M. S., Berggren P. O. and Jonsson A. B., 1998. Cell signaling by the type IV pili of pathogenic *Neisseria*. *J Biol Chem* 273, 21777-82.
- Kirchner M., Heuer D. and Meyer T. F., 2005. CD46-independent binding of neisserial type IV pili and the major pilus adhesin, PilC, to human epithelial cells. *Infect Immun* 73, 3072-82.
- Kirchner M. and Meyer T. F., 2005. The PilC adhesin of the *Neisseria* type IV pilus-binding specificities and new insights into the nature of the host cell receptor. *Mol Microbiol* 56, 945-57.
- Kohl P. K., Kratochiel M., Gorner R., Kunze F., Senf-Blum A., Eggert-Kruse W., Gerhard I., Hoferer E. and Petzoldt D., 1992. [Local and systemic humoral immune response to protein I of *Neisseria gonorrhoeae*]. *Hautarzt* 43, 352-8.
- Laan M., Cui Z. H., Hoshino H., Lotvall J., Sjostrand M., Gruenert D. C., Skoogh B. E. and Linden A., 1999. Neutrophil recruitment by human IL-17 via C-X-C chemokine release in the airways. *J Immunol* 162, 2347-52.
- Lammel C. J., Sweet R. L., Rice P. A., Knapp J. S., Schoolnik G. K., Heilbron D. C. and Brooks G. F., 1985. Antibody-antigen specificity in the immune response to infection with *Neisseria gonorrhoeae*. *J Infect Dis* 152, 990-1001.
- Langrish C. L., McKenzie B. S., Wilson N. J., de Waal Malefyt R., Kastelein R. A. and Cua D. J., 2004. IL-12 and IL-23: master regulators of innate and adaptive immunity. *Immunol Rev* 202, 96-105.
- Lee H. S., Boulton I. C., Reddin K., Wong H., Halliwell D., Mandelboim O., Gorringer A. R. and Gray-Owen S. D., 2007. Neisserial outer membrane vesicles bind the coinhibitory receptor carcinoembryonic antigen-related cellular adhesion molecule 1 and suppress CD4+ T lymphocyte function. *Infect Immun* 75, 4449-55.

- Lee H. S., Ostrowski M. A. and Gray-Owen S. D., 2008. CEACAM1 dynamics during *Neisseria gonorrhoeae* suppression of CD4⁺ T lymphocyte activation. *J Immunol* 180, 6827-35.
- Lee S. W., Higashi D. L., Snyder A., Merz A. J., Potter L. and So M., 2005. PilT is required for PI(3,4,5)P₃-mediated crosstalk between *Neisseria gonorrhoeae* and epithelial cells. *Cell Microbiol* 7, 1271-84.
- LeibundGut-Landmann S., Gross O., Robinson M. J., Osorio F., Slack E. C., Tsoni S. V., Schweighoffer E., Tybulewicz V., Brown G. D., Ruland J. and Reis e Sousa C., 2007. Syk- and CARD9-dependent coupling of innate immunity to the induction of T helper cells that produce interleukin 17. *Nat Immunol* 8, 630-8.
- Long C. D., Madraswala R. N. and Seifert H. S., 1998. Comparisons between colony phase variation of *Neisseria gonorrhoeae* FA1090 and pilus, pilin, and S-pilin expression. *Infect Immun* 66, 1918-27.
- Lorenzen D. R., Dux F., Wolk U., Tsirpouchtsidis A., Haas G. and Meyer T. F., 1999. Immunoglobulin A1 protease, an exoenzyme of pathogenic *Neisseriae*, is a potent inducer of proinflammatory cytokines. *J Exp Med* 190, 1049-58.
- Lynch E. C., Blake M. S., Gotschlich E. C. and Mauro A., 1984. Studies of Porins: Spontaneously Transferred from Whole Cells and Reconstituted from Purified Proteins of *Neisseria gonorrhoeae* and *Neisseria meningitidis*. *Biophys J* 45, 104-7.
- Maisey K., Nardocci G., Imarai M., Cardenas H., Rios M., Croxatto H. B., Heckels J. E., Christodoulides M. and Velasquez L. A., 2003. Expression of Proinflammatory Cytokines and Receptors by Human Fallopian Tubes in Organ Culture following Challenge with *Neisseria gonorrhoeae*. *Infect. Immun.* 71, 527-532.
- McChesney D., Tramont E. C., Boslego J. W., Ciak J., Sadoff J. and Brinton C. C., 1982. Genital antibody response to a parenteral gonococcal pilus vaccine. *Infect Immun* 36, 1006-12.
- McGee Z. A., Jensen R. L., Clemens C. M., Taylor-Robinson D., Johnson A. P. and Gregg C. R., 1999. Gonococcal infection of human fallopian tube mucosa in organ culture: relationship of mucosal tissue TNF- α concentration to sloughing of ciliated cells. *Sex Transm Dis* 26, 160-5.
- McGuirk P., McCann C. and Mills K. H., 2002. Pathogen-specific T regulatory 1 cells induced in the respiratory tract by a bacterial molecule that stimulates interleukin 10 production by dendritic cells: a novel strategy for evasion of protective T helper type 1 responses by *Bordetella pertussis*. *J Exp Med* 195, 221-31.
- Merz A. J., Enns C. A. and So M., 1999. Type IV pili of pathogenic *Neisseriae* elicit cortical plaque formation in epithelial cells. *Mol Microbiol* 32, 1316-32.
- Merz A. J. and So M., 1997. Attachment of piliated, Opa- and Opc- gonococci and meningococci to epithelial cells elicits cortical actin rearrangements and clustering of tyrosine-phosphorylated proteins. *Infect Immun* 65, 4341-9.
- Muenzner P., Dehio C., Fujiwara T., Achtman M., Meyer T. F. and Gray-Owen S. D., 2000. Carcinoembryonic antigen family receptor specificity of *Neisseria meningitidis* Opa variants influences adherence to and invasion of proinflammatory cytokine-activated endothelial cells. *Infect Immun* 68, 3601-7.
- Muller A., Gunther D., Brinkmann V., Hurwitz R., Meyer T. F. and Rudel T., 2000. Targeting of the pro-apoptotic VDAC-like porin (PorB) of *Neisseria gonorrhoeae* to mitochondria of infected cells. *EMBO J* 19, 5332-43.

- Muller A., Gunther D., Dux F., Naumann M., Meyer T. F. and Rudel T., 1999. Neisserial porin (PorB) causes rapid calcium influx in target cells and induces apoptosis by the activation of cysteine proteases. *EMBO J* 18, 339-52.
- Muller A., Rassow J., Grimm J., Machuy N., Meyer T. F. and Rudel T., 2002. VDAC and the bacterial porin PorB of *Neisseria gonorrhoeae* share mitochondrial import pathways. *EMBO J* 21, 1916-29.
- Murphy G. L., Connell T. D., Barritt D. S., Koomey M. and Cannon J. G., 1989. Phase variation of gonococcal protein II: regulation of gene expression by slipped-strand mispairing of a repetitive DNA sequence. *Cell* 56, 539-47.
- Napolitani G., Rinaldi A., Bertonni F., Sallusto F. and Lanzavecchia A., 2005. Selected Toll-like receptor agonist combinations synergistically trigger a T helper type 1-polarizing program in dendritic cells. *Nat Immunol* 6, 769-76.
- Nocera M. and Chu T. M., 1995. Characterization of latent transforming growth factor-beta from human seminal plasma. *Am J Reprod Immunol* 33, 282-91.
- O'Connor R. A., Prendergast C. T., Sabatos C. A., Lau C. W., Leech M. D., Wraith D. C. and Anderton S. M., 2008. Cutting edge: Th1 cells facilitate the entry of Th17 cells to the central nervous system during experimental autoimmune encephalomyelitis. *J Immunol* 181, 3750-4.
- Ouyang W., Kolls J. K. and Zheng Y., 2008. The biological functions of T helper 17 cell effector cytokines in inflammation. *Immunity* 28, 454-67.
- Ovcinnikov N. M. and Delektorskij V. V., 1971. Electron microscope studies of gonococci in the urethral secretions of patients with gonorrhoea. *Br J Vener Dis* 47, 419-39.
- Pantelic M., Kim Y. J., Bolland S., Chen I., Shively J. and Chen T., 2005. *Neisseria gonorrhoeae* kills carcinoembryonic antigen-related cellular adhesion molecule 1 (CD66a)-expressing human B cells and inhibits antibody production. *Infect Immun* 73, 4171-9.
- Parge H. E., Forest K. T., Hickey M. J., Christensen D. A., Getzoff E. D. and Tainer J. A., 1995. Structure of the fibre-forming protein pilin at 2.6 Å resolution. *Nature* 378, 32-8.
- Plant L. J. and Jonsson A. B., 2006. Type IV pili of *Neisseria gonorrhoeae* influence the activation of human CD4+ T cells. *Infect Immun* 74, 442-8.
- Plummer F. A., Chubb H., Simonsen J. N., Bosire M., Slaney L., Maclean I., Ndinya-Achola J. O., Waiyaki P. and Brunham R. C., 1993. Antibody to Rmp (outer membrane protein 3) increases susceptibility to gonococcal infection. *J Clin Invest* 91, 339-43.
- Plummer F. A., Chubb H., Simonsen J. N., Bosire M., Slaney L., Nagelkerke N. J., Maclean I., Ndinya-Achola J. O., Waiyaki P. and Brunham R. C., 1994. Antibodies to opacity proteins (Opa) correlate with a reduced risk of gonococcal salpingitis. *J Clin Invest* 93, 1748-55.
- Popp A., Dehio C., Grunert F., Meyer T. F. and Gray-Owen S. D., 1999. Molecular analysis of neisserial Opa protein interactions with the CEA family of receptors: identification of determinants contributing to the differential specificities of binding. *Cell Microbiol* 1, 169-81.
- Preston A., Mandrell R. E., Gibson B. W. and Apicella M. A., 1996. The lipooligosaccharides of pathogenic gram-negative bacteria. *Crit Rev Microbiol* 22, 139-80.
- Rarick M., McPheeters C., Bright S., Navis A., Skefos J., Sebastiani P. and Montano M., 2006. Evidence for cross-regulated cytokine response in human peripheral blood

- mononuclear cells exposed to whole gonococcal bacteria in vitro. *Microb Pathog* 40, 261-70.
- Rice P. A., Vayo H. E., Tam M. R. and Blake M. S., 1986. Immunoglobulin G antibodies directed against protein III block killing of serum-resistant *Neisseria gonorrhoeae* by immune serum. *J Exp Med* 164, 1735-48.
- Rowell E. and Wilson C. B., 2009. Programming perpetual T helper cell plasticity. *Immunity* 30, 7-9.
- Rudel T., Schmid A., Benz R., Kolb H. A., Lang F. and Meyer T. F., 1996. Modulation of *Neisseria* porin (PorB) by cytosolic ATP/GTP of target cells: parallels between pathogen accommodation and mitochondrial endosymbiosis. *Cell* 85, 391-402.
- Rudel T., van Putten J. P., Gibbs C. P., Haas R. and Meyer T. F., 1992. Interaction of two variable proteins (PilE and PilC) required for pilus-mediated adherence of *Neisseria gonorrhoeae* to human epithelial cells. *Mol Microbiol* 6, 3439-50.
- Sabatos C. A., Doh J., Chakravarti S., Friedman R. S., Pandurangi P. G., Tooley A. J. and Krummel M. F., 2008. A synaptic basis for paracrine interleukin-2 signaling during homotypic T cell interaction. *Immunity* 29, 238-48.
- Sakaguchi S., 2004. Naturally arising CD4⁺ regulatory T cells for immunologic self-tolerance and negative control of immune responses. *Annu. Rev. Immunol.* 22, 531-562.
- Simpson S. D., Ho Y., Rice P. A. and Wetzler L. M., 1999. T lymphocyte response to *Neisseria gonorrhoeae* porin in individuals with mucosal gonococcal infections. *J Infect Dis* 180, 762-73.
- Snapper C. M., Rosas F. R., Kehry M. R., Mond J. J. and Wetzler L. M., 1997. Neisserial porins may provide critical second signals to polysaccharide-activated murine B cells for induction of immunoglobulin secretion. *Infect Immun* 65, 3203-8.
- Song W., Condron S., Mocca B. T., Veit S. J., Hill D., Abbas A. and Jerse A. E., 2008. Local and humoral immune responses against primary and repeat *Neisseria gonorrhoeae* genital tract infections of 17 β -estradiol-treated mice. *Vaccine* 26, 5741-51.
- Srivastava M. D., Lippes J. and Srivastava B. I., 1996. Cytokines of the human reproductive tract. *Am J Reprod Immunol* 36, 157-66.
- Stagg A. J., Tuffrey M., Woods C., Wunderink E. and Knight S. C., 1998. Protection against ascending infection of the genital tract by *Chlamydia trachomatis* is associated with recruitment of major histocompatibility complex class II antigen-presenting cells into uterine tissue. *Infect Immun* 66, 3535-44.
- Tapchaisri P. and Sirisinha S., 1976. Serum and secretory antibody responses to *Neisseria gonorrhoeae* in patients with gonococcal infections. *Br J Vener Dis* 52, 374-80.
- Tapsall J., 2001. Antimicrobial resistance in *Neisseria gonorrhoeae*. World Health Organization Department of Communicable Disease Surveillance and Response.
- Taylor B. N., Saavedra M. and Fidel P. L., Jr., 2000. Local Th1/Th2 cytokine production during experimental vaginal candidiasis: potential importance of transforming growth factor-beta. *Med Mycol* 38, 419-31.
- Tobiason D. M. and Seifert H. S., 2001. Inverse relationship between pilus-mediated gonococcal adherence and surface expression of the pilus receptor, CD46. *Microbiology* 147, 2333-40.
- Tramont E. C., Sadoff J. C., Boslego J. W., Ciak J., McChesney D., Brinton C. C., Wood S. and Takafuji E., 1981. Gonococcal pilus vaccine. Studies of antigenicity and inhibition of attachment. *J Clin Invest* 68, 881-8.

- Tsirpouchtsidis A., Hurwitz R., Brinkmann V., Meyer T. F. and Haas G., 2002. Neisserial immunoglobulin A1 protease induces specific T-cell responses in humans. *Infect Immun* 70, 335-44.
- van Putten J. P., Duensing T. D. and Cole R. L., 1998. Entry of OpaA+ gonococci into HEp-2 cells requires concerted action of glycosaminoglycans, fibronectin and integrin receptors. *Mol Microbiol* 29, 369-79.
- van Putten J. P. and Paul S. M., 1995. Binding of syndecan-like cell surface proteoglycan receptors is required for *Neisseria gonorrhoeae* entry into human mucosal cells. *EMBO J* 14, 2144-54.
- van Vliet S. J., Steeghs L., Bruijns S. C., Vaezirad M. M., Snijders Blok C., Arenas Busto J. A., Deken M., van Putten J. P. and van Kooyk Y., 2009. Variation of *Neisseria gonorrhoeae* lipooligosaccharide directs dendritic cell-induced T helper responses. *PLoS Pathog* 5, e1000625.
- Verhagen J., Sabatos C. A. and Wraith D. C., 2008. The role of CTLA-4 in immune regulation. *Immunol Lett* 115, 73-4.
- Virji M., Evans D., Hadfield A., Grunert F., Teixeira A. M. and Watt S. M., 1999. Critical determinants of host receptor targeting by *Neisseria meningitidis* and *Neisseria gonorrhoeae*: identification of Opa adhesiotopes on the N-domain of CD66 molecules. *Mol Microbiol* 34, 538-51.
- Virji M., Makepeace K., Ferguson D. J. and Watt S. M., 1996a. Carcinoembryonic antigens (CD66) on epithelial cells and neutrophils are receptors for Opa proteins of pathogenic neisseriae. *Mol Microbiol* 22, 941-50.
- Virji M., Watt S. M., Barker S., Makepeace K. and Doyonnas R., 1996b. The N-domain of the human CD66a adhesion molecule is a target for Opa proteins of *Neisseria meningitidis* and *Neisseria gonorrhoeae*. *Mol Microbiol* 22, 929-39.
- Wada K., Nomura S., Morii E., Kitamura Y., Nishizawa Y., Miyake A. and Terada N., 1996. Changes in levels of mRNAs of transforming growth factor (TGF)-beta1, -beta2, -beta3, TGF-beta type II receptor and sulfated glycoprotein-2 during apoptosis of mouse uterine epithelium. *J Steroid Biochem Mol Biol* 59, 367-75.
- Weel J. F. and van Putten J. P., 1991. Fate of the major outer membrane protein P.IA in early and late events of gonococcal infection of epithelial cells. *Res Microbiol* 142, 985-93.
- Wen K. K., Giardina P. C., Blake M. S., Edwards J., Apicella M. A. and Rubenstein P. A., 2000. Interaction of the gonococcal porin P.IB with G- and F-actin. *Biochemistry* 39, 8638-47.
- Wetzler L. M., Ho Y. and Reiser H., 1996. Neisserial porins induce B lymphocytes to express costimulatory B7-2 molecules and to proliferate. *J Exp Med* 183, 1151-9.
- WHO, 2011. Emergence of multi-drug resistant *Neisseria gonorrhoeae* - Threat of global rise in untreatable sexually transmitted infections.
- Wiesner P. J. and Thompson S. E., 3rd, 1980. Gonococcal diseases. *Dis Mon* 26, 1-44.
- Xu L., Kitani A., Fuss I. and Strober W., 2007. Cutting edge: regulatory T cells induce CD4+CD25-Foxp3- T cells or are self-induced to become Th17 cells in the absence of exogenous TGF-beta. *J Immunol* 178, 6725-9.
- Zheng H., 1997. Analysis of the antigen-antibody specificity in the semen of patients with *Neisseria gonorrhoeae*. *Chin Med Sci J* 12, 47-9.



Sexually Transmitted Infections

Edited by Prof. Nancy Malla

ISBN 978-953-51-0258-8

Hard cover, 244 pages

Publisher InTech

Published online 21, March, 2012

Published in print edition March, 2012

Sexually transmitted infections (STIs) are infections that are spread primarily through person to person sexual contact. There are more than 30 different sexually transmissible bacteria, viruses and parasites. STIs lead to high morbidity and complications. This book entitled as Sexually Transmitted Infections is not a text book but provides useful information for general reference work for physicians, researchers and students interested in the subject. Each chapter is abundant in tips useful to general readers as well. It also includes the Introductory chapter providing an overview with special emphasis on syndromic approach to the management of STIs in clinical setting.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Mónica Imarai, Claudio Acuña, Alejandro Escobar, Kevin Maisey and Sebastián Reyes-Cerpa (2012). Mucosal Immunity and Evasion Strategies of Neisseria gonorrhoeae, Sexually Transmitted Infections, Prof. Nancy Malla (Ed.), ISBN: 978-953-51-0258-8, InTech, Available from: <http://www.intechopen.com/books/sexually-transmitted-infections/mucosal-immunity-and-evasion-strategies-of-neisseria-gonorrhoeae>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen