

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Genetics of Hearing Loss

Nejat Mahdieh^{1,2}, Bahareh Rabbani¹ and Ituro Inoue¹

¹*Division of Human Genetics, National Institute of Genetics, Mishima, Shizuoka,*

²*Medical Genetic Group, Faculty of Medicine, Ilam University of Medical Sciences, Ilam,*

¹*Japan*

²*Iran*

1. Introduction

Hearing loss (HL) is the most common sensory defect in human beings, affecting 1.86 in 1000 newborns around the *world* which half of it is due to genetic causes (Morton & Nance, 2006). HL can be syndromic or nonsyndromic. Individuals affected with syndromic form have additional clinical signs whereas nonsyndromic HL is not associated with other clinical signs and symptoms. All Mendelian patterns of inheritance have been observed in nonsyndromic HL (NSHL) including autosomal dominant (AD), autosomal recessive (AR), X-linked inheritance (XL) and mitochondrial inheritance (MT); autosomal recessive is the main form of NSHL, i.e. 75-85 % of NSHL show AR pattern in affected pedigrees.

As known, ear is the organ of hearing and balance. Hearing is dependent on a series of complex events. The ear has three anatomical parts including outer, middle and inner ear. The external ear which is composed of the auricle, ear canal and eardrum membrane collects sound waves and transmits them to the eardrum. Three tiny bones of middle ear (the ossicles) act as levers and conduct the sounds to the oval window, and finally through the cochlea (a snail-shaped organ) which has the auditory receptors (the organ of Corti) in the inner ear [Raphael & Altschuler 2003]. A collagen-based extracellular matrix, called tectorial membrane on top of the hair cells is vibrated by sound waves [Richardson et al., 2008]. Within the organ of Corti, physical vibrations produce a mechano-electrical transduction which is detected by hair cells and these cells respond by producing electrical impulses. Nerves transmit these impulses to the brain where they are interpreted. Different sound frequencies stimulate the hair cells in different parts of the organ of Corti and lead to perception of different sound frequencies. Sounds are processed in both sides of the brain but the interpretation of the sounds takes place at the left side of brain. Sounds are heard at normal hearing thresholds between 0-20 dB across the 125-8000 Hz range while loss of more than 20 dB, is said to have hearing loss which is confirmed by measuring pure tone average (PTA) (average hearing sensitivity at 500, 1000 and 2000 Hz).

Nearly one hundred and twenty million people suffer from hearing impairment around the world. History of some important events about human hereditary HL is shown in table 1 [Nance & Sweeney, 1975; Wallis et al., 1988; Kimberling et al., 1990; Leon et al., 1992;

Guilford et al., 1994; Kelsell et al., 1997; Lynch et al., 1997; Eudy et al., 1998; Gorlin, 2004; Dror & Avraham 2009].

Time	Event
Sixteenth century	Reports indicating the prevention of the deaf from marrying
Seventeenth century	The mode of recessive and dominant HL
Nineteenth century	"The most frequent causes of congenital deafness are hereditary.."
1968	One of the genetic forms of deafness was described
1975	The forms of HL, the need for research and genetic counseling were described
1988	An X-linked form of deafness was mapped in a large Mauritian family
1990	The first locus for syndromic HL, <i>USH2A</i> , was mapped
1992	The first locus for ADNSHL*, <i>DFNA1</i> , was mapped in a Costa Rican family
1994	The first and second loci for ARNSHL▪, <i>DFNB1</i> and <i>DFNB2</i> , and <i>DFNA2</i> loci were mapped
1997	<i>GJB2</i> and <i>DIAPH1</i> genes were discovered for <i>DFNB1</i> and <i>DFNA1</i> loci, respectively

*Autosomal dominant non syndromic hearing loss, ▪Autosomal recessive nonsyndromic hearing loss

Table 1. Chronological events regarding hereditary HL.

One of the main programs of the World Health Organization (WHO) is to encourage countries for the prevention of deafness [Emery, 2003]. Understanding the molecular and genetic mechanisms of HL may lead to development of new therapy and treatment approaches. Here, we review major causes leading to either syndromic or nonsyndromic HL.

2. Classification of hearing loss

Approximately two-thirds of the HL affected children show the problem at birth and unfortunately it may not be diagnosed before the age of 3 years. HL has several classification criteria which are important for diagnosis, prognosis and treatment [Mahdiah et al., 2010a]. These criteria are summarized in table 2.

2.1 Etiology

Based on the cause of sensorineural deafness, HL is categorized into three major forms as acquired, genetic and unknown. There are many causes for hearing loss:

1. Acquired HL: infectious and pharmaceutical agents known as teratogens would affect the sense of hearing. HL could occur by physiological, biochemical or infectious factors. However, genetic background has an important effect on its occurrence. The risk factors that may affect the hearing process are as follows:

- a. factors before birth including congenital infections (e.g. toxoplasmosis, measles, syphilis, smallpox, cytomegalovirus, herpes virus), congenital deformities of auricular and ear duct [Willems, 2004; Shin et al., 2011].
 - b. factors during birth including prematurity and low birth weight (less than 1500gr) and increased blood bilirubin [Willems, 2004].
 - c. factors affecting after birth including infections and bacterial meningitis, mumps, otitis media, blood infection and autotoxic drugs such as aminoglycosides, head injury or skull fracture which lead to anesthesia [Willems, 2004].
2. Genetic HL: the genetic basis of HL is known for more than 100 years. In the early decade of 1800s, the Irish physician William Wild explained the inheritance of HL. His theory differentiated between dominant and recessive inheritance. He also observed that men showed more X-linked transmission [Willems, 2004].

2.2 Severity

Intensity of the sound is calculated in units of decibel (dB), which is logarithm intensity of the sound wave to a reference sound intensity divided by ten [Willems, 2004]. The normal hearing threshold is 15 dB. A regular conversation occurs at level of 45 to 60 dB.

2.3 Position of damage

- a. Conductive HL: Factors affecting sound transmission including auricular, ear canal, eardrum, outer and middle ear bones to the cochlea cause conductive HL. The most common causes of conductive HL are the external and middle ear congenital abnormalities such as atrophy and dysplasia, duct obstruction, impacted cerumen, otitis, middle ear and Tympanic membrane problems.
- b. Sensorineural HL (SNHL): The disorder occurs in the auditory nerve or the cochlea. In other words, the abnormalities occur some place between the hair cells and auditory brain regions. The most common causes of SNHL are:
 - b1. congenital causes such as rubella, syphilis, Usher syndrome, Alport Syndrome, Waardenburg syndrome and autosomal dominant and recessive sensorineural deafness [Friedman et al., 2003].
 - b2. acquired factors: Infections such as measles, cytomegalovirus, bacterial meningitis, autotoxicity of drug consumption, noise pollution including long-term exposure, presbycusis and sudden idiopathic HL [Willems, 2004].
- c. Mixed HL: in this type of hearing loss, conductive and sensorineural problems are observed simultaneously. Infections such as tuberculosis, some syndromes and skull fractures may also cause mixed HL.

2.4 Age of onset

On the basis of the age of onset, HL is divided into the following types:

- a. Prelingual: Loss of hearing occurs before speech is acquired. If a child has a congenital hearing impairment, he would not be able to speak normally.
- b. Postlingual: Loss of hearing occurs after speech is developed.
- c. Presbycusis or age-related HL (ARHL): Epidemiologic studies show that nearly 25 % of 60 year olds and more than 50 % of 80 year ages undergo ARHL [Dror & Avraham, 2009; Huang & Tang, 2010].

2.5 Signs

HL may be associated with other physical problems which are called syndromic HL. Genetic HLs without any other complications is called non-syndromic genetic hearing loss [Willems, 2004]. HL loci are named with the prefix DFN, followed by the mode of inheritance which is indicated by B, A, X and Y for autosomal recessive (DFNB), autosomal dominant (DFNA), X-linked (DFNX) and Y-linked (DFNY), respectively. The order in which loci have been described is denoted by a number after these letters, e.g. DFNB1 is the first identified locus causing autosomal recessive HL [Guilford et al., 1994].

Criterion	Class	Definition and example
Age of onset	Prelingual HL	HL occurs before language acquisition
	Postlingual HL	HL occurs after language acquisition
	Presbycusis	Age-related HL
Etiology	Acquired	Caused by environmental agents such as viral and bacterial infections (prenatal, e.g., CMV, toxoplasmosis, rubella; postnatal, e.g. meningitis), hyperbilirubinemia, head trauma, anoxia, noise exposure and ototoxic drugs
	Genetic	Caused by gene mutation
	Idiopathic	Unexplained cause
Clinical phenotypes	Syndromic	associated with other symptoms
	Nonsyndromic	Deafness is the only defect
Position of damage	Conductive HL	Caused by a problem transferring sound waves through the external ear, tympanic membrane or middle ear
	Sensorineural HL	Caused by damage to the inner ear (vestibulocochlear nerve)
	Mixed HL	Caused by a combination of sensorineural and conductive HL
Severity	Mild	Difficulty in hearing of 26–40 dB sounds
	Moderate	41–55 dB
	Moderately severe	56–70 dB
	Severe	71–90 dB
	Profound	>90 dB
Mode of inheritance	Autosomal dominant	DFNA loci (DFNA1-64)
	Autosomal recessive	DFNB loci (DFNB1-96)
	Sex-linked	DFNX loci (DFN1-8)
	Y-linked	DFNY loci (DFNY1)
	Mitochondrial	12SrRNA (MT-RNR1), tRNA ^{Ser(UCN)} (MT-TS1)

Table 2. Various criteria for the classification of hearing loss.

3. The frequency of genetic HL

Genetic HL occurs 1 in 2000 to 1 in 650 live births [Morton & Nance, 2006]. About 70% of the cases are nonsyndromic [Tekin et al., 2001]. Studies show that 75% of nonsyndromic HL

are inherited as autosomal recessive [Tekin et al., 2001]. 10-20% of cases are inherited as autosomal dominant and 1-5% are X-linked recessive. Approximately, 1% of human genes, i.e. 200 to 250 genes are responsible for hereditary HL [Finsterer & Fellingner, 2005]. So far, more than one hundred loci and 55 genes have identified which are involved in nonsyndromic HL (<http://hereditaryhearingloss.org>).

4. Non-syndromic HL

A high frequency of genetic HL occurs without any abnormality in other organs classified as non-syndromic HL. Different patterns of inheritance have been observed in NSHL.

4.1 Different types of NSHL

Variety of protein coding genes such as gap junctions (connexin encoding genes), motor proteins (myosins) cytoskeletal (e.g. actin), ion channels, structural proteins (Tectorin alpha, Otoancorin, Stereocilin, etc), transcription factors (POU3F4, POU4F3 and Eyes absent 4 or EYA4), and additionally microRNA genes are involved in HL [Willems, 2004; Mencia et al., 2009; Mahdieh et al., 2010a]. *GJB2* mutations are seen in 50% of autosomal recessive HL in the Caucasians [Kelsell et al., 1997; Tekin et al., 2001]. Some genes e.g. *GJB2* gene is expressed in a variety of organs of the body while others such as *OTO* Ancorin is only expressed in the inner ear.

4.1.1 Autosomal recessive non-syndromic HL

Autosomal recessive non-syndromic HL (ARNSHL) was first described in 1846. It is the severest form of congenital HL in which there is a defect in cochlea in nearly all cases. Loci of ARNSHL are designated as the DFNB; DF stands for Deafness and B indicates the autosomal recessive pattern of inheritance. Up to date, 46 genes and nearly 100 loci have been identified for HL (Table 3). Regarding different studies, connexin 26 gene mutations differ depending on geographical place and ethnicity [Zelante et al., 1997; Morell et al., 1998; Mahdieh & Rabbani, 2009]. Here, we discuss the most common genes causing ARNSHL.

4.1.1.1 *GJB2* and *GJB6* genes and connexins

The first locus of ARNSHL designated as DFNB1 was identified by Guilford and colleagues in 1994. These researches confirmed linkage to chromosome 13q12-q13 in two consanguineous families [Guilford et al., 1994]. More consanguineous families of different ethnic groups were linked to the DFNB1 locus [Morle et al., 2000]. Phenotypic differences were observed within different families which indicated that allelic heterogeneity may be present in the locus DFNB1.

GJB2 is a small gene encompassing 5.5 Kb. It has two exons encoding a 4.2Kb mRNA and a protein of 226 amino acids. A six repeat of G is located at position 30 to 35 of coding region of *GJB2* gene from which deletion of one G is known as 35delG or 30delG (Figure 1) [Kelley et al., 1998]. 35delG is the most common mutation in the Caucasians and may cause up to 70% of all *GJB2* gene mutations. Profound HL caused by *GJB2* gene mutations is found in 50% of the cases; 30% are severe, 20% moderate and 1-2% are mild cases [Smith & Hone, 2003]. Other *GJB2* mutations have been reported with higher frequencies in some ethnic

groups [Morell et al., 1998; Mahdieh & Rabbani, 2009]. A large number of studies have been reported about *GJB2* mutations including genotype-phenotype correlations, phenotypic variability, de novo mutations, dominant mutations, ethnic-specific distribution of mutations, digenic inheritance and allelic heterogeneity [del Castillo et al., 2002; Smith & Hone, 2003; Mahdieh et al., 2009; 2010b, 2010c]. Also, a modifier gene has been suggested because of intrafamilial phenotypic variability of the cases [Higert et al., 2009a; Mahdieh et al., 2010b].

GJB2 and *GJB6* genes are about 35 kb apart from each other. *GJB6* gene encodes a protein called Connexin 30 (MIM 604418) which has 261 amino acids. Connexin 30 is produced in different tissues of the body such as the cochlea, brain and thyroid [Grifa et al., 1999]. The importance of this gene was evident when some families had a mutated allele of *GJB2* and the second mutant allele was in the *GJB6* (digenic inheritance) [del Castillo et al., 2002].

Locus	Location	Gene	references
X-Linked			
DFNX1	Xq22	<i>PRPS1</i>	Liu et al., 2010
DFNX2	Xq21.1	<i>POU3F4</i>	De Kok et al., 1995
DFNX4	Xp22	<i>SMPX</i>	del Castillo et al., 1996
Autosomal Dominant			
DFNA1	5q31	<i>DIAPH1</i>	Lynch et al., 1997
DFNA2A	1p34	<i>KCNQ4</i>	Kubisch et al., 1999
DFNA2B	1p35.1	<i>GJB3</i>	Xia et al., 1998
DFNA3A	13q11-q12	<i>GJB2</i>	Kelsell et al., 1997
DFNA3B	13q12	<i>GJB6</i>	Grifa et al., 1999
DFNA4	19q13	<i>MYH14</i>	Donaudy et al, 2004
DFNA5	7p15	<i>DFNA5</i>	Van Laer et al., 1998
DFNA6	4p16.3	<i>WFS1</i>	Bespalova et al., 2001
DFNA9	14q12-q13	<i>COCH</i>	Robertson et al., 1998
DFNA10	6q22-q23	<i>EYA4</i>	Wayne et al., 2001
DFNA11	11q12.3-q21	<i>MYO7A</i>	Liu et al., 1997
DFNA12	11q22-24	<i>TECTA</i>	Verhoeven et al., 1998
DFNA13	6p21	<i>COL11A2</i>	McGuirt et al., 1999
DFNA15	5q31	<i>POU4F3</i>	Vahava et al., 1998
DFNA17	22q	<i>MYH9</i>	Lalwani et al., 2000
DFNA20	17q25	<i>ACTG1</i>	Zhu et al., 2003,
DFNA22	6q13	<i>MYO6</i>	Melchionda et al.,
DFNA28	8q22	<i>GRHL2</i>	Peters et al., 2002
DFNA36	9q13-q21	<i>TMC1</i>	Kurima et al., 2002
DFNA39	4q21.3	<i>DSPP</i>	Xiao et al., 2001
DFNA44	3q28-29	<i>CCDC50</i>	Modamio-Hoybjor et al., 2007
DFNA48	12q13-q14	<i>MYO1A</i>	Donaudy et al., 2003
DFNA50	7q32.2	<i>MIR96</i>	Mencia et al., 2009
DFNA51	9q21	<i>TJP2</i>	Walsh et al., 2010
DFNA64	12q24.31-12q24.32	<i>SMAC/DIABLO</i>	Cheng et al., 2011

Locus	Location	Gene	references
Autosomal Recessive			
DFNB1	13q12	<i>GJB2</i>	Kelsell et al., 1997
DFNB2	11q13.5	<i>MYO7A</i>	Liu et al., 1997
DFNB3	17p11.2	<i>MYO15A</i>	Wang et al., 1998
DFNB4	7q31	<i>SLC26A4</i>	Li et al., 1998
DFNB6	3p14-p21	<i>TMIE</i>	Naz et al, 2002
DFNB7/11	9q13-q21	<i>TMC1</i>	Kurima et al., 2002
DFNB8/10	21q22	<i>TMPRSS3</i>	Scott et al., 2001
DFNB9	2p22-p23	<i>OTOF</i>	Yasunaga et al., 1999
DFNB12	10q21-q22	<i>CDH23</i>	Bork et al., 2001
DFNB15	19p13	<i>GIPC3</i>	Charizopoulou et al., 2011
DFNB16	15q21-q22	<i>STRC</i>	Verpy et al., 2001
DFNB18	11p14-15.1	<i>USH1C</i>	Ouyang et al., 2002
DFNB21	11q	<i>TECTA</i>	Mustapha et al., 1999
DFNB22	16p12.2	<i>OTOA</i>	Zwaenepoel et al ., 2002
DFNB23	10p11.2-q21	<i>PCDH15</i>	Ahmed et al, 2003
DFNB24	11q23	<i>RDX</i>	Khan et al., 2007
DFNB25	4p13	<i>GRXCR1</i>	Schraders et al., 2010
DFNB28	22q13	<i>TRIOBP</i>	Riazuddin et al, 2006
DFNB29	21q22	<i>CLDN14</i>	Wilcox et al., 2001
DFNB30	10p11.1	<i>MYO3A</i>	Walsh et al., 2002
DFNB31	9q32-q34	<i>WHRN</i>	Mburu et al., 2003
DFNB32	1p13.3-22.1	<i>GPSM2</i>	Walsh et al., 2010
DFNB35	14q24.1-24.3	<i>ESRRB</i>	Collin et al., 2008
DFNB36	1p36.3	<i>ESPN</i>	Naz et al., 2004
DFNB37	6q13	<i>MYO6</i>	Ahmed et al., 2003
DFNB39	7q21.1	<i>HGF</i>	Schultz et al., 2009
DFNB42	3q13.31-q22.3	<i>ILDR1</i>	Borck et al., 2011
DFNB49	5q12.3-q14.1.	<i>MARVELD2</i>	Riazuddin et al., 2006
DFNB53	6p21.3	<i>COL11A2</i>	Chen et al., 2005
DFNB59	2q31.1-q31.3	<i>PJVK</i>	Delmaghani et al., 2006
DFNB61	7q22.1	<i>SLC26A5</i>	Liu et al., 2003
DFNB63	11q13.2-q13.4	<i>LRTOMT/ COMT2</i>	Ahmed et al., 2008
DFNB66	6p21.2-22.3	<i>LHFPL5</i>	Shabbir et al., 2006
DFNB72	19p13.3	<i>GIPC3</i>	Rehman et al., 2011
DFNB73	1p32.3	<i>BSND</i>	Riazuddin et al., 2009
DFNB74	12q14.2-q15	<i>MSRB3</i>	Ahmed et al., 2011
DFNB77	18q12-q21	<i>LOXHD1</i>	Grillet et al., 2009
DFNB79	9q34.3	<i>TPRN</i>	Rehman et al., 2010
DFNB84	12q21.2	<i>PTPRQ</i>	Schraders er al., 2010
DFNB91	6p25	<i>SERPINB6</i>	Sirmaci et al., 2010
DFNB95	19p13	<i>GIPC3</i>	Charizopoulou et al., 2011

Table 3. Non-syndromic genes responsible for HL up to 2011.




Fig. 1. Schematic structure and domains of Connexin 26 protein, Connexon and Gap Junction channel. A) The most common mutations in specific populations (35delG, 167delT, 235delC, R143W and W24X mutations in the Caucasian, Ashkenazi Jewish, Japanese, Ghanian and Indian populations, respectively) are shown. 35delG, W24X, 167delT, 235delC and R143W located on NT, TM1, EC1, TM2 and TM3 domains, respectively. TM1-TM4 denotes transmembrane domains, EC1-2 denotes extracellular domains, IC denotes cytoplasmic domain, NT denotes amino (NH₂) terminus and CT denotes carboxyl (COOH) terminus. B) Six connexins can oligomerize to form hemichannels named connexons. Connexons then pass throughout the membrane to make the gap junction channels. Homomeric and heteromeric channels can be formed as connexins selectively interact with each other.

A few point mutations have also been reported in *GJB6* as the cause of ARNSHL. *GJB6* [del Castillo et al., 2005; Pallares-Ruiz et al., 2002]. Later studies determined that *GJB6* mutations in cis state, not in trans, would destroy the *GJB2* expression. Therefore, the digenic hypothesis may not be correct. Four large deletions have been recognized in *GJB6* gene including del(*GJB6*-D13S1830), del(*GJB6*-D13S1854), del(chr13:19,837,344-19,968,698) and 920 Kb deletion [del Castillo et al., 2002, 2005; Wilch et al., 2010]. The deletions may include more than 10% of DFNB1 alleles [Stevenson et al., 2003]. So far, del (*GJB6*-D13S1830) has not been seen in many populations [Mahdiah et al., 2004, 2011]. The del (*GJB6*-D13S1830) and del (*GJB6*-D13S1854) mutations not only truncate the synthesis of *GJB6* gene but also destroy *GJB2* gene expression.

Connexins encoded by GJ genes are members of transmembrane family proteins that have 20 members in humans [Holms & Steel, 1999]. These proteins were classified in three groups of alpha, beta and gamma proteins. Common nomenclature system is based on molecular weight of proteins e.g. Cx26 and Cx32. Despite the differences in the size and primary amino acid composition, connexins have similar topology. These proteins have four transmembrane domains which are connected by two extracellular and one intracellular loop. The carboxyl and amino terminals are located at the cytoplasmic side. Most cells express more than one type of connexin. Gap junctions show different permeability and conductance which may create channels with specific characteristics. Also, in order to compensate for the decrease in the expression of some of the connexins, other connexins may be produced at an enhanced rate [Kumar & Kilula, 1996]. Hemi-channels (connexons) are composed of six connexin subunits and two hemi-channels make the channel forming the gap junctions [Kelley et al., 1998]. The important role of these channels is transportation of potassium ions [Kelley et al., 1998] and glutamate released from hair cells to initiate action potential. Different connexins may be made up of hemi channels with homomer or heteromer subunits.

4.1.1.2 *MYO15A* gene in DFNB3 locus

In 1995, a report showed that 2% of rural individuals in the north coast of Bali, Indonesia were affected with a profound sensorineural non-syndromic HL. Due to high percentage of deaf people in this village a local sign language had been created for communication [Wang et al., 1998].

The locus was mapped on chromosome 17p11.2 by whole genome study. *MYO15A* gene has 66 exons and 71097 bp, encoding a 11863 bp transcript [Liang et al., 1999]. Myosin gene was identified by positional and functional cloning approaches [Wang et al., 1998]. Mutations in the gene are responsible for 5% of severe to profound deafness cases in Pakistan [Friedman et al., 2003]. *MYO15A* gene mutations were reported in families from Turkey, Brazil and India [Kalay et al., 2007; Nal et al., 2007; Lezirovitz et al., 2008]. The role of myosin filaments can be traced in a variety of cellular functions including cell motility, muscle contraction, synaptic transmission, cytokinesis, endocytosis, exocytosis and probably in gene expression as a modulator [Craig & Woodhead, 2006; Loikkanen et al., 2009]. As the organism gets more complex, there may be more myosin isoforms found in the organism [Oliver et al., 1999; Friedman et al., 1999]. The heavy chain of XV myosin has 3531 amino acids. There is a unique proline-rich region at the amino terminal of myosin which weighs 140KDa and has no similarity to any of the known proteins. Next to this domain exists a motor domain and a tail domain [Belyantseva et al., 2003]. In addition to the

sensory cells of cochlear, myosin is expressed in the pituitary gland, neuroendocrine cells, parathyroid and pancreas [Llyod et al., 2001]. It is also found in stereocilia of hair cells [Belyantseva et al., 2003].

4.1.1.3 *SLC26A4* gene in DFNB4 locus

DFNB4 locus, located at chromosome 7q31, was first reported to be linked to recessive non-syndromic deafness in a large Middle-Eastern Druze family. In 1997, the *SLC26A4* (Pendrin coding protein) was identified by positional cloning at the pendred syndrome locus (Everett et al. 1997) and was later also shown to be the gene mutated in DFNB4 [Li et al., 1998]. Pendred syndrome was identified in 1896 as neurosensory HL and goiter. HL in Pendred syndrome is the most common cause of deafness due to defect of cochlea such as dilation sac and duct of endolymph and enlarged vestibular duct [Everett et al., 1997].

Mutations of *SLC26A4* gene are the second leading cause of ARNSHL. So far, more than 140 mutations have been reported for Pendred syndrome. Phenotypic spectrum of *SLC26A4* gene mutations varies from Pendred syndrome to nonsyndromic HL. Four mutations are common in northern Europeans i.e L236P, T416P, E384G, IVS8 +1 G> A) [Hilgert et al., 2008]. In a study conducted in Spanish population 27% had homozygous *SLC26A4* mutations [Pera et al., 2008]. Mutations of *SLC26A4* gene have been observed in several ethnic populations [Albert et al., 2006; Hu et al., 2007; Yoon et al., 2008]. The prevalence of *SLC26A4* gene mutation is about 40% in Caucasians of which 24% are bi-allelic [Albert et al., 2006].

SLC26A4 gene has 21 exons within 57175 bp of DNA sequence. Its transcript is about 5 Kb encoding into a 87KDa protein having 780 amino acids. The gene is expressed in lining cells of endolymph duct as well as non-sensory cells of utricle, saccule, kidney and thyroid. Various models have been reported for the structure of Pendrin protein. New model suggests that pendrin protein is a transmembrane protein traversing fifteen times throughout the membrane [Dossena et al., 2009]. The protein is involved in anion exchange of HCO⁻, Cl⁻, I⁻ and OH⁻ ions [Mount & Romero, 2004].

4.1.1.4 *TMC1* gene in DFNB7/11 locus

DFNB7 and DFNB11 were determined as the cause of HL on chromosome 9q13-q21 in two Indian and two inbred Israeli families, respectively [Jian et al., 1995]. In 2002, eight different mutations in *TMC1* gene were linked to one DFNA36 family and eleven DFNB7/11 families [Kurima et al., 2002]. More than twenty different point mutations and two deletions have been identified in different families. It seems that c.100C>T mutation includes approximately 40% of all *TMC1* mutations in Turkey [Hilgert et al., 2008, 2009b]. In a survey of 51 Turkish families, 5 patients had mutations of *TMC1* gene [Hilgert et al., 2008]. Mutations of *TMC1* are responsible for at least 6% of all cases of ARNSHL in northeast and eastern part of Turkey [Kalay et al., 2005]. Three mutations c.100C> T (R34X), c.77611G> A and g.94615A> C have been reported in Iranian families [Hilgert et al., 2009b].

Based on sequence homology studies, eight TMC genes exist in vertebrates. *TMC1* gene has 24 exons and encodes a 3201 nucleotide RNA. It expresses a complete transmembrane protein with six membrane passing domain which has no similarity to proteins of known function. Mouse ortholog transcript (*TMC1*) is expressed in cochlea and vestibular hair cells [Kurima et al., 2002].

4.1.1.5 *TMPRSS3* gene in DFNB8/10 locus

DFNB8/10 locus was separately mapped on chromosome 21q22.3 in two consanguineous Pakistani (DFNB8) and Palestinian families (DFNB10) [Bonné-Tamir et al., 1996; Veske et al., 1996]. Haplotype analysis and sequencing analysis of the families resulted in detection of mutations in *TMPRSS3* [Scott et al., 2001]. The gene belongs to a subfamily of transmembrane serine proteases type III protein [Szabo et al., 2003] expressed in supporting cells of the organ of Corti [Guipponi et al., 2002]. Although, the specific role of *TMPRSS3* protein in growth, development and survival of auditory apparatus has not been found but it activates the epithelial sodium channel (ENaC) in vitro [Guipponi et al., 2002]. The mutated alleles of the gene may inactivate the serine protease catalytic activity. Therefore, *TMPRSS3* proteolytic function may be important during the development of inner ear [Guipponi et al., 2002, 2008].

TMPRSS3 gene has 13 exons within 24 Kb, encoding a 2468 bp mRNA which encodes a protein with 454 amino acids [Guipponi et al., 2008]. In 2009, 16 mutations in *TMPRSS3* have been reviewed and reported by a study [Hilgert et al., 2009b]. From 25 studied Turkish families, three had mutations of *TMPRSS3* gene [Wattenhofer et al., 2005; Sahin-Calapoglu et al., 2005]. Mutations of *TMPRSS3* gene account for 1% of hearing loss in Caucasian children with non-syndromic HL [Wattenhofer et al., 2005]. Mutations of *TMPRSS3* gene have been reported in 4 of 290 Pakistani families [Ahmed et al., 2004].

4.1.1.6 *OTOF* gene in DFNB9 locus

OTOF gene contains 48 exons encoding a 1997 amino acid polypeptide called otoferlin which is member of Ferlin family of proteins [Mirghomizadeh et al., 2002]. Ferlin family of proteins have a domain called C2. These proteins contain a transmembrane C-terminal domain [Yasunaga et al., 1999]. C2 domain is a structural domain in some proteins that are involved in directing proteins to the cell membrane [Davletov & Südhof, 1993].

Otoferlin is expressed in the brain and cochlea. This protein plays an important role in releasing neurotransmitters in the auditory nerve cells [Yasunaga et al., 1999]. Mutations of the gene can lead to auditory neuropathy in which the sound from inner ear is not transferred to the brain. Q829X mutation is very common in the Hispanic which is the third cause of ARNSHL [Migliosi et al., 2002]. Mutations of the gene have been found in families of Lebanese origin [Yasunaga et al., 1999]. Varga *et al.* reported 8 mutations in 65 studied families with ARNSHL [Varga et al., 2006]. *OTOF* mutations have been found in Pakistani families; gene mutations may account for deafness in 2.3% of this population [Choi et al., 2009].

4.1.1.7 *CDH23* gene in DFNB12 locus

The superfamily of cadherin has about 100 members with a variety of roles in cell adhesion, growth and developmental signaling, maintenance and function of the tissues [Jamora & Fuchs, 2002; Nelson & Nusse, 2004; Gumbiner, 2005; Halbleib & Nelson, 2006]. Cadherin 23 protein has a role in connection of developing stereocilia [Siemens et al., 2004]. In 1996, DFNB12 was mapped to chromosome 10q21-q22 in a consanguineous Syrian family [Chaib et al., 1996]. Usher syndrome type 1 D (USH1D) was also mapped to the same position. Allelic mutations of the *CDH23* gene encoding cadherin 23 cause DFNB12 HL and USH1D [Bolz et al., 2001; Bork et al., 2001]. Missense mutations usually cause DFNB12 HL

but nonsense and premature stop codon mutations cause Usher syndrome type 1D although this relationship is not definite. No single gene mutation is common in this gene [Hilgert 2009b]. In 64 Japanese families, five mutations were found in *CDH23* [Wagatsuma et al., 2007].

4.1.1.8 *TMHS* or *LHFPL5* genes in *DFNB67* locus

Non syndromic HL in a Pakistani family linked to a new region on chromosome 6p21.1-p22.3 defining a new locus, *DFNB67* in 2006; *TMHS* or *LHFPL5* gene was mapped in this region [Shabbir et al., 2006]. *LHFPL5* has 4 exons and encodes a 2162 nucleotide mRNA and translates into a protein of 219 amino acids. The proposed structure of the protein is a four pass transmembrane domain. Mutations of this gene have been reported in patients from Pakistan and Turkey (C161F, Y127C, P83fsX84). *TMHS* is important for the transmission of sound.

4.1.2 Autosomal dominant non- syndromic hearing loss

Late onset, mild and progressive forms of HL are the usual phenotypes associated with autosomal dominant form of deafness. About 25 genes and more than 60 loci have been reported for autosomal dominant non- syndromic hearing loss (ADNSHL). There is no frequent gene mutated in ADNSHL but mutations in some genes including *WFS1*, *KCNQ4*, *COCH* and *GJB2* have been suggested to be common (Kelsell et al., 1997; Nie 2008; Higert et al., 2009).

4.1.2.1 *WFS1* gene and its protein

The *WFS1* (Wolfram) gene at *DFNA6* locus, located on 4p16, consists of 8 exons and has a length of about 33.4 kb and a 3.6 kb transcript. It codes for a polypeptide of 890 amino acids [Hofmann et al., 2003]. The Wolframin protein is a resident component of the endoplasmic reticulum (ER) and may be involved in membrane trafficking, processing and/or regulation of ER calcium homeostasis [Fonseca et al., 2010]. In the inner ear, however, this protein may be helpful to maintain the appropriate levels of calcium ions and/or other charged particles required for hearing process [Cryns et al., 2003].

Dominant mutations in *WFS1* can cause a characteristic type of HL which affects the low frequencies and less loss in hearing in the high frequencies [Bespalova et al., 2001; Fukuoka et al., 2007]. It has been shown that dominant mutations are usually located in the C-terminal domain. The recessive Wolfram syndrome is caused by numerous mutations distributed along the entire gene. Two mutations c.424_425ins16 and c.1362_1377del16 have a high frequency in some specific populations including Spanish patients and Italians, respectively [Gómez-Zaera et al., 2001; Colosimo et al., 2003]. It is hypothesized that, inactivating mutations may lead to Wolfram syndrome and missense mutations occurring in the C-terminal domain can cause the characteristic low-frequency ADNSHL [Cryns et al., 2003].

4.1.2.2 *KCNQ4* gene and its protein

The *KCNQ4* gene at *DFNA2* locus, located on 1p34, consists of 14 exons and codes for a polypeptide of 695 amino acids, a voltage-gated potassium channel. It is a member of the KCNQ voltage-gated K⁺ channel family [Coucke et al., 1999]. It has an important role in K⁺

secretion into the endolymph by stria marginal cells. Ten missense mutations, two small deletions and one splice mutation in *KCNQ4* have been reported so far. It is believed that a dominant-negative effect of the missense mutations in this gene lead to interference of the mutant protein with the normal channel subunit, affecting the pore structure of the channels. Therefore, hearing loss with a lower age of onset is observed at all frequencies [Coucke et al., 1999; Akita et al., 2001]. Deletion mutations which have a haploinsufficiency effect lead to a milder HL with an older age of onset at high frequencies [Coucke et al., 1999; Akita et al., 2001].

4.1.2.3 *COCH* gene and its protein

The DFNA9 causative gene, *COCH* located on 14q12-q13, consists of 11 exons and encodes a 550 amino acid extracellular matrix protein named cochlin. This protein has several domains including two von Willebrand factor A-like domains (vWFA1 and 2) and a LCCL domain (a region homologous to a domain in factor C of *Limulus*). To date, eleven missense mutations and one small deletion in *COCH* gene have been reported. Most of the missense mutations are located within exon 4 and 5 which encode the LCCL domain (Figure 2) [Robertson et al., 1998; Collin et al., 2006].




Fig. 2. Schematic structure of cochlin and distribution of the mutations along its domains. The NT signal peptide is followed by a LCCL domain and two vWF domains. S indicates several cysteine residues, NT denotes amino (NH₂) terminus and CT denotes carboxyl (COOH) terminus.

4.1.3 X and Y linked HL

There are fewer X-linked forms of HL (DFNX) than ARNSHL and ADNSHL. X-linked form of deafness has been reported as prelingual or progressive in different families. Five loci and three genes (*POU3F4*, *SMPX* and *PRPS1*) have been reported for X-linked HL (<http://hereditaryhearingloss.org/>).

To date, only one locus has been linked to chromosome Y (DFNY1) that was found in a very large Chinese family (seven generations) [Wang et al., 2004]. They reported that the ages of onset for the patrilineal relatives were from 7 to 27 years. *PCDH11Y*, encoding a protocadherin, was suggested to be the causality [Wang et al., 2004].

4.1.4 Mitochondrial HL

In healthy individuals, only one type of mitochondrial DNA genotype (homoplasmy) exists, but in many mitochondrial diseases, mitochondrial genome has mixed genotype (heteroplasmy). Heteroplasmy differs from one tissue to another and can even differ within the cells of a tissue. A few genes contribute to mitochondrial HL [Fischel-Ghodsian, 2003]. Due to the important function of mitochondria in producing chemical energy through oxidative phosphorylation, mitochondrial DNA mutations can cause systemic neuromuscular disorders such as HL. mtDNA mutations may be inherited or acquired (Table 4); the inherited mitochondrial mutations can cause many clinical features including myopathy, neuropathy, diabetes mellitus and sensorineural HL [Finsterer & Fellingner, 2005; Guan 2011]. Acquired mitochondrial mutations may be associated with aging and age-related HL or presbycusis [Fischel-Ghodsian, 1999, 2003]. Multiorganic mitochondrial syndromes are often lethal in homoplasmic state. Mitochondrial homoplasmy exists in LHON (Leber Hereditary Optic Neuropathy) and maternal inherited HL [Fischel-Ghodsian, 2003]. Myoclonic epilepsy and ragged red fibers (MERRF), Kearns-Sayre syndrome (KSS) and mitochondrial encephalomyopathy with lactic acidosis and stroke-like episodes are associated with progressive HL [Zeviani et al., 1998; Goto et al., 1990].

Gene	Mutation	Phenotype	Reference
<i>MTRNR1</i> (12S rRNA)	1555A->G	NSHL/ Aminoglycoside induced/worsened	Estivill et al., 1998
	1494C->T	NSHL/ Aminoglycoside induced/worsened	Zhao et al., 2004
	961 (mutations)	NSHL/ Aminoglycoside induced/worsened	Bacino et al., 1995
	1095T>C	NSHL/ Aminoglycoside induced/ parkinsonism, and neuropathy	Zhao et al., 2004
	827A>G	NSHL/ Aminoglycoside induced	Li et al., 2005
<i>MTTS1</i> (tRNA ^{Ser} (UCN))	7444G>A	NSHL/ Aminoglycoside induced	Pandya et al., 1999;
	7445A->G	NSHL/ Palmoplantar keratoderma	Fischel-Ghodsian, 2003
	7472insC	NSHL/ Neurological dysfunction, including ataxia, dysarthria and myoclonus	Jaksch et al., 1998
	7510T->C	NSHL/ no additional symptoms	Hutchin et al.2000
	7511T->C	NSHL/ no additional symptoms	Friedman et al., 1999
	7512T>C	HL/ Progressive myoclonic epilepsy and ataxia	Jaksch et al., 1998
<i>MTTL1</i> (tRNA ^{Leu} (UUR))	3243A>G	maternally inherited diabetes and deafness/MELAS	Goto et al., 1990
<i>tRNA^{Lys}</i>	8296A>G	maternally inherited diabetes and deafness	Kameoka et al., 1998
	8332A>G	dystonia, stroke-like episodes and HL	Gal et al., 2010
<i>tRNA^{Glu}</i>	14709T>C	maternally inherited diabetes and deafness	Rigoli et al., 2001

Table 4. Identified mitochondrial DNA mutations in HL.

5. Age-related HL

Biological changes accumulate in people during life as individuals age. About one hundred thousand individuals die each day of age-related causes around the world [de Grey, 2007]. Age-related HL (ARHL) or presbycusis is the most frequent sensory defect in the elderly people. It occurs due to accumulation of environmental and genetic changes i.e. gradual deleterious changes in the ear gives rise hearing impairment in older people. Approximately 25 % of 60 year olds and more than 50 % of 80 year ages suffer from ARHL [Dror & Avraham, 2009; Huang & Tang, 2010]. Many heterogeneous factors including family history, exposure to loud noises, ototoxic medication, exposure to chemicals, free radical (reactive oxygen species) chronic medical conditions, malnutrition, mtDNA mutations, alcohol abuse and smoking etc. may cause this type of HL [Van Eyken et al., 2007b; Huang & Tang, 2010].

Some common deletions and acquired mtDNA point mutations due to reactive oxygen species (ROS) have also been suggested to cause presbycusis. Although genetic studies on ARHL are increasing in the recent years, there is a little information about the role of genes to its etiology. Two basic approaches have been used to identify susceptibility genes for ARHL: the linkage study and the association study [Van Eyken et al., 2007b]. Several single nucleotide polymorphisms (SNPs) have been reported to correlate with presbycusis; variants in *GRHL2*, *GRM7*, *KCNQ4* and N-acetyltransferase 2 are involved (Table 5) [Van Eyken et al., 2006, 2007a; Van Laer et al., 2008; Friedman et al., 2009]. Mutations in cadherin 23 coded by *CDH23* gene may also cause ARHL [Johnson et al., 2010]. More recently, a genome-wide association scan was conducted on ARHL in the genetically isolated Finnish Saami population. This study confirmed, and also provided further evidence for the role of the previous reported gene, *GRM7* in ARHL. *IQGAP2* gene was also proposed to be involved in presbycusis [van Laer et al., 2010]. Mechanism of ARHL is not well understood. However, new promising technology and strategies may help to discover the exact role of genetic mutations in presbycusis. Finding of the genetic variants causing ARHL will ultimately lead to discovery of new pharmaceutical interventions and the development of new approaches to identify at risk individuals.

SNP (RS number)	Gene	Protein or Function	
SNP9 (rs727146) SNP12 (rs2149034) SNP18 (rs12143503)	<i>KCNQ4</i>	Potassium channel (voltage-gated)	Van Eyken et al., 2006
NAT2*6A (rs1799930)	N-acetyltransferase 2	metabolism of cytotoxic, carcinogenic compounds and ROS	Unal et al., 2005
36738A>G (rs10955255) 42731C>T (rs2127034) 53110C>T (rs1981361)	<i>GRHL2</i>	transcription factor cellular promoter 2-like 3	Van Laer et al., 2008
7155702T>A (rs11928865)	<i>GRM7</i>	glutamate receptor, metabotropic,7	Friedman et al., 2009
75920972A>G (rs457717) 75922504C>T (rs1697845)	<i>IQGAP2</i>	IQ motif-containing GTPase-activating-like protein	Van Laer et al., 2010

Table 5. SNPs associated with ARHL.

6. Syndromic genetic deafness

More than 400 syndromes have been described in OMIM. Here, genetic aspects of common syndromes which are associated with HL are briefly explained.

Usher Syndrome: Usher syndrome, named after Charles Usher (1914) a British ophthalmologist, is the most prevalent cause of autosomal recessive HL, accounting for nearly 3-5 per 100,000 in the general population and 1-10% among profoundly deaf children [Boughman et al., 1983]. Several clinical subtypes have been distinguished based on its characterized features i. e. severity of the HL and the onset of retinitis pigmentosa [Yan & Liu, 2010]. Type 1 patients have profound HL, vestibular dysfunction and the onset of retinitis pigmentosa in childhood [Hope et al., 1997]. The type 2 patients have normal vestibular response, mild to moderate HL and RP begins in the second decade of life [Hope et al., 1997]. Progressive HL and variable vestibular response characterize type 3 patients and the onset of retinitis pigmentosa is variable as well [Smith et al., 1995]. Usher syndrome has a heterogeneous causality (Table 6); to date, 12 different loci and 10 genes have been reported (<http://hereditaryhearingloss.org/>). One of these identified genes, *MYO7A*, encoding myosin 7A, is a unique molecular motor for hair cells [Weil et al., 1995]. Cadherin 23, an adhesion molecule, coded by *CDH23* gene may have an important role in crosslinking of stereocilia [Bolz et al., 2001; Bork et al., 2001].

Locus	Gene	Ref.
USH1B (11q13.5)	<i>MYO7A</i>	Weil et al.,1995
<i>USH1C</i> (11p15.1)	<i>USH1C</i>	Smith et al., 1992
USH1D (10q22.1)	<i>CDH23</i>	Bork et al., 2001
USH1F (10q21-22)	<i>PCDH15</i>	Ahmed et al., 2001
USH1G (17q24-25)	<i>SANS</i>	Mustapha et al., 2002
<i>USH2A</i> (1q41)	<i>USH2A</i>	Kimberling et al., 1990
USH2C (5q14.3-q21.3)	<i>VLGR1</i>	Weston et al., 2004
USH2D (9q32)	<i>WHRN</i>	Ebermann et al., 2007
USH3 (3q21-q25)	<i>USH3A</i>	Joensuu et al., 2001
10q24.31	<i>PDZD7</i>	Ebermann et al., 2010

Table 6. Reported genes for Usher syndrome.

Pendred syndrome: Pendred syndrome, named after Vaughan Pendred (1896) a British physician, is the most common syndromic form of HL and associated with abnormal iodine metabolism (goiter). It is an autosomal recessive disorder which accounts for 4-10% of deaf cases [Fraser 1965]. The defective organic binding of iodine in the thyroid gland may distinguished by a positive potassium perchlorate discharge test; however the test is not specific and its sensitivity is unclear. HL is usually bilateral, severe to profound and may be present at birth, and sloping in the higher frequencies [Kopp et al., 2008]. The casual gene is *SLC26A4* (PDS) on chromosome 7q31 encoding a protein named pendrin (Figure 3). It regulates transportation of iodine and chloride/ bicarbonate ions in the inner ear, thyroid, and kidney. Mutations of this gene can cause NSHL DFNB4 and enlarged vestibular aqueduct syndrome as well [Everett et al., 1997].




Fig. 3. Hypothetic structure and domains of Pendrin protein. The most common mutations (L236P, IVS8+1G>A, T416P, and H723R) accounting for approximately 60% of the total PS genetic load are shown. TM1-TM12 denotes transmembrane domains, EC1-6 denotes extracellular domains, IC denotes cytoplasmic domain, NT denotes amino (NH₂) terminus and CT denotes carboxyl (COOH) terminus.

Alport syndrome: Alport syndrome, a hereditary disorder of basement membranes, is characterized by renal abnormalities including glomerulonephritis, hematuria (“red diaper”) and renal failure, and ocular problems as well as progressive sensorineural HL [Wester et al., 1995]. Mutations in various genes encoding type 4 collagen (COL4A3, COL4A4 and COL4A5) have been reported to cause Alport syndrome [Lemmink et al., 1994; Hudson et al., 2003]; nearly 85% of the cases are due to COL4A5 mutations [Hudson et al., 2003]. These collagens are components of the basilar membranes, the spiral ligament and stria vascularis. X-linked pattern of inheritance is observed in the majority (80 %); the remaining shows autosomal recessive [Lemmink et al., 1994] and autosomal dominant [van der Loop et al., 2000], inheritance patterns. It is estimated that 10% to 15% of X-linked patients represent de novo mutations in COL4A5 [Gubler et al., 2007]. Since uremia leads to death in males prior to 30 years of age, it is essential to diagnose it early in men. Symptoms are usually more severe than women. The progressive sensorineural HL usually begins in the adolescent years [Wester et al., 1995]. The mechanism of HL has not been explained exactly yet, although the basement membrane damages are suggested to affect adhesion of the cells of the organ of Corti and basilar membrane leading to HL [Merchant et al., 2004].

Waardenburg syndrome: Waardenburg disease, named after Petrus Johannes Waardenburg (1886-1979), accounts for 1-3% of congenital HL [Read & Newton, 1997]. In addition, the disease shows other clinical features. Four types of syndrome can be distinguished on the basis of accompanying abnormalities [Read & Newton, 1997]: In type 1, patients show dystopia canthorum, iris heterochromy, brilliant blue eyes, broad nasal root, premature

graying of hair, white forelock, and vestibular dysfunction. Type 2 patients have similar phenotype but not dystopia canthorum. In type 3 (so called Klein-Waardenburg syndrome) [Klein, 1983], upper extremity abnormalities other Type 1 clinical features and dystopia canthorum and are observed. In type 4 (so called Shah-Waardenburg syndrome) [Shah et al., 1981] patients demonstrate all findings shown in Type 2 with the addition of pigmentation abnormalities and Hirschsprung's disease. Sensorineural hearing loss is observed in 60 % and 90 % of type 1 and type 2 patients, respectively [Newton, 1990].

Types 1 and 3 of Waardenburg syndrome occur due to mutations in the *PAX3* gene encoding a DNA-binding protein essential for determining the fate of neural crest cells [Baldwin et al., 1994]. Type 2 is due to mutations in *MITF* gene [Tassabehji et al., 1994]. Mutations in three genes, *EDN3*, *SOX10* and *EDNRB* genes, can lead to Type 4 [Edery et al., 1996; Hofstra et al., 1996; Pingault et al., 1998]. *SOX10* mutations, account for approximately half of type 4 patients and are likely responsible for about 15% of Type 2 as well [Bondurand et al., 2007]. *In vitro* studies have shown that *EDN3* plays as a stimulation factor of proliferation and melanogenesis of neural crest cells. *EDNRB* is suggested to have an important role in the development of epidermal melanocytes and enteric neurons. *SOX10* is a DNA-binding transcription factor and involved in promoting cell survival prior to lineage commitment [Kapur, 1999]. There is a wide range of variation in HL phenotype so that some patients may not exhibit HL.

Branchio-oto-renal syndrome: Branchio-oto-renal syndrome (BOR) is an autosomal dominant disorder, accounting for 2% of profoundly deaf children and is characterized by branchial derived anomalies, otologic anomalies (Mondini's dysplasia and stapes fixation) and renal malformation. HL may affect 70-93% of the BOR patients but there is a high variability in age of onset and severity [Chen et al., 1995]. HL can be sensorineural, conductive or mixed, stable or progressive and mild or profound. Mutations in *EYA1* gene have been identified to cause BOR syndrome (BOR1) [Abdelhak et al., 1997]. It has been shown that this gene has a role in development of the inner ear and kidney [Abdelhak et al., 1997]. Studies of transgenic mice have indicated that *EYA1* homozygous knockouts have not developed ears and kidneys. In addition to *EYA1*, mutations in two genes named *SIX1* and *SIX5* have been reported to cause BOR3 and BOR2, respectively [Ruf et al., 2004; Hoskins et al., 2007].

Stickler Syndrome: Stickler Syndrome (STL), named after Stickler (1965), follows an autosomal dominant pattern of inheritance and is characterized by progressive sensorineural HL, cleft palate, abnormal development of the epiphysis, vertebral abnormalities and osteoarthritis. On the basis of clinical features, four types of STL exist. Type 1 patients have typical features of the disease including progressive myopia leading to retinal detachment, midface hypoplasia, cleft palate, variable sensorineural HL and vitreoretinal degeneration. Mutations in *COL2A1* gene encoding a fibrillar collagen type 2 subunitA1 can cause the classic phenotype [Ahmad et al., 1991]. There is no retinal detachment in Type 2 and the phenotype is caused by *COL11A1* gene mutations [Richards et al., 1996]. Facial abnormalities seen in Type 1 are not observed in Type 3. Mutations of *COL11A2* lead to STL Type 3 [Vikkula et al., 1995]. Recently, mutations in *COL9A1* have been identified to cause an autosomal recessive form of STL, Type 4 [Van Camp et al., 2006].

7. Genetic evaluation

The main problem in the diagnosis of disorders such as deafness is its heterogeneity. Genetic study of HL has considerable benefits for patients which are as follows:

- a. Identifying the medical and non medical decisions e.g cochlear implant
- b. Carrier testing and prenatal diagnosis
- c. Prediction for the progressive state of the disease
- d. Eliminating unnecessary tests and investigations
- e. Providing appropriate genetic counseling before marriage, especially when they have heterogeneous conditions that carry different mutated genes.

Genetic evaluation should be considered for children with newly diagnosed loss of hearing especially if no specific cause is determined. For example, there is no need for genetic evaluation of the family of a child with HL due to meningitis; although, they may need assurance of not transmitting the disease to the next generation. Genetic evaluation includes several steps:

1. Reviewing the complete history of prenatal, neonatal and medical history of growth and development
2. Complete physical examination of patients and other family members
3. Evaluating the genetics, molecular and cellular diagnosis

Based on previous studies, deaf people have positive assortive marriage; it is estimated that 90% of deaf individuals marry deaf. Depending on the pattern of inheritance they might have a deaf child. For example if both parental recessive alleles are similar, there is 100% chance of having a deaf child; and if one of the parents carry a dominant form of HL and the other carry the recessive form of HL the chance would be 50% for the dominant gene.

Early diagnosis of HL is important in gaining speech progression and social skills of the children which would lead to better life of these individuals and would later help them in cochlea implant. Hereditary or genetic understanding of the causes of HL is important. The benefits of this understanding and knowledge, not only allows physicians to help the families of at risk but also may help in treatment and control of HL. Sometimes it is possible to prevent hearing loss from worsening. HL may be one of the clinical signs of a syndrome and if the genetic cause of HL is determined it may help to predict and treat other clinical complications [Extivill et al., 1998].

8. Conclusion

HL is the most common sensory defect affecting human beings. It is categorized on the basis of several criteria. Genetic factors can be traced in half of the cases. Nonsyndromic HL can follow any of the Mendelian inheritance patterns, but the majority are ARNSHL. Approximately fifty genes have been reported to be involved in HL, and based on an estimation nearly 200 to 250 genes may cause HL. Genetic understanding of the causes of HL and finding the molecular mechanism of hearing process are valuable for genetic counseling, prevention and development of new therapeutic approaches. Many studies have been published about finding new genes causing prelingual nonsyndromic HL. Presbycusis is very common among elderly people and research on this phenotype needs more attention.

New technology and strategies such as next generation sequencing can help to discover new genes for deafness in future.

9. References

- Abdelhak S, Kalatzis V, Heilig R, Compain S, Samson D, Vincent C, Levi-Acobas F, Cruaud C, Le Merrer M, Mathieu M, König R, Vigneron J, Weissenbach J, Petit C & Weil D. Clustering of mutations responsible for branchio-oto-renal (BOR) syndrome in the eyes absent homologous region (*eyaHR*) of *EYA1*. *Hum Mol Genet.* 1997;6(13):2247-55.
- Ahmad NN, Ala-Kokko L, Knowlton RG, Jimenez SA, Weaver EJ, Maguire JL, Tasman W & Prockop DJ. Stop codon in the procollagen II gene (*COL2A1*) in a family with the Stickler syndrome (arthro-ophthalmopathy). *Proc Natl Acad Sci U S A.* 1991;88(15):6624-7.
- Ahmed ZM, Li XC, Powell SD, Riazuddin S, Young TL, Ramzan K, Ahmad Z, Luscombe S, Dhillon K, MacLaren L, Ploplis B, Shotland LI, Ives E, Riazuddin S, Friedman TB, Morell RJ & Wilcox ER. Characterization of a new full length *TMPRSS3* isoform and identification of mutant alleles responsible for nonsyndromic recessive deafness in Newfoundland and Pakistan. *BMC Med Genet.* 2004;5:24.
- Ahmed ZM, Masmoudi S, Kalay E, Belyantseva IA, Mosrati MA, Collin RW, Riazuddin S, Hmani-Aifa M, Venselaar H, Kawar MN, Tlili A, van der Zwaag B, Khan SY, Ayadi L, Riazuddin SA, Morell RJ, Griffith AJ, Charfedine I, Caylan R, Oostrik J, Karaguzel A, Ghorbel A, Riazuddin S, Friedman TB, Ayadi H & Kremer H. Mutations of *LRTOMT*, a fusion gene with alternative reading frames, cause nonsyndromic deafness in humans. *Nat Genet.* 2008;40(11):1335-40.
- Ahmed ZM, Riazuddin S, Ahmad J, Bernstein SL, Guo Y, Sabar MF, Sieving P, Riazuddin S, Griffith AJ, Friedman TB, Belyantseva IA & Wilcox ER. *PCDH15* is expressed in the neurosensory epithelium of the eye and ear and mutant alleles are responsible for both USH1F and DFNB23. *Hum Mol Genet.* 2003;12(24):3215-23.
- Ahmed ZM, Riazuddin S, Bernstein SL, Ahmed Z, Khan S, Griffith AJ, Morell RJ, Friedman TB, Riazuddin S & Wilcox ER. Mutations of the protocadherin gene *PCDH15* cause Usher syndrome type 1F. *Am J Hum Genet.* 2001;69(1):25-34.
- Ahmed ZM, Yousaf R, Lee BC, Khan SN, Lee S, Lee K, Husnain T, Rehman AU, Bonneux S, Ansar M, Ahmad W, Leal SM, Gladyshev VN, Belyantseva IA, Van Camp G, Riazuddin S, Friedman TB & Riazuddin S. Functional null mutations of *MSRB3* encoding methionine sulfoxide reductase are associated with human deafness DFNB74. *Am J Hum Genet.* 2011;88(1):19-29.
- Akita J, Abe S, Shinkawa H, Kimberling WJ & Usami S. Clinical and genetic features of nonsyndromic autosomal dominant sensorineural hearing loss: *KCNQ4* is a gene responsible in Japanese. *J Hum Genet.* 2001;46(7):355-61.
- Albert S, Blons H, Jonard L, Feldmann D, Chauvin P, Loundon N, Sergent-Allaoui A, Houang M, Joannard A, Schmerber S, Delobel B, Leman J, Journel H, Catros H, Dollfus H, Eliot MM, David A, Calais C, Drouin-Garraud V, Obstoy MF, Tran Ba Huy P, Lacombe D, Duriez F, Francannet C, Bitoun P, Petit C, Garabédian EN, Couderc R, Marlin S & Denoyelle F. *SLC26A4* gene is frequently involved in nonsyndromic hearing impairment with enlarged vestibular aqueduct in Caucasian populations. *Eur J Hum Genet.* 2006;14(6):773-9.

- Bacino C, Prezant TR, Bu X, Fournier P & Fischel-Ghodsian N. Susceptibility mutations in the mitochondrial small ribosomal RNA gene in aminoglycoside induced deafness. *Pharmacogenetics*. 1995 Jun;5(3):165-72.
- Baldwin CT, Lipsky NR, Hoth CF, Cohen T, Mamuya W & Milunsky A. Mutations in *PAX3* associated with Waardenburg syndrome type I. *Hum Mutat*. 1994;3(3):205-11.
- Baldwin CT, Weiss S, Farrer LA, De Stefano AL, Adair R, Franklyn B, Kidd KK, Korostishevsky M & Bonn -Tamir B. Linkage of congenital, recessive deafness (DFNB4) to chromosome 7q31 and evidence for genetic heterogeneity in the Middle Eastern Druze population. *Hum Mol Genet*. 1995;4(9):1637-42.
- Belyantseva IA, Boger ET & Friedman TB. Myosin XVa localizes to the tips of inner ear sensory cell stereocilia and is essential for staircase formation of the hair bundle. *Proc Natl Acad Sci U S A*. 2003;100(24):13958-63.
- Bespalova IN, Van Camp G, Bom SJ, Brown DJ, Cryns K, DeWan AT, Erson AE, Flothmann K, Kunst HP, Kurnool P, Sivakumaran TA, Cremers CW, Leal SM, Burmeister M & Lesperance MM. Mutations in the Wolfram syndrome 1 gene (*WFS1*) are a common cause of low frequency sensorineural hearing loss. *Hum Mol Genet*. 2001;10(22):2501-8.
- Bolz H, von Brederlow B, Ram rez A, Bryda EC, Kutsche K, Nothwang HG, Seeliger M, del C-Salced  Cabrera M, Vila MC, Molina OP, Gal A & Kubisch C. Mutation of *CDH23*, encoding a new member of the cadherin gene family, causes Usher syndrome type 1D. *Nat Genet*. 2001;27(1):108-12.
- Bondurand N, Dastot-Le Moal F, Stanchina L, Collot N, Baral V, Marlin S, Attie-Bitach T, Giurgea I, Skopinski L, Reardon W, Toutain A, Sarda P, Echaieb A, Lackmy-Port-Lis M, Touraine R, Amiel J, Goossens M & Pingault V. Deletions at the *SOX10* gene locus cause Waardenburg syndrome types 2 and 4. *Am J Hum Genet*. 2007;81(6):1169-85.
- Bonn -Tamir B, DeStefano AL, Briggs CE, Adair R, Franklyn B, Weiss S, Korostishevsky M, Frydman M, Baldwin CT & Farrer LA. Linkage of congenital recessive deafness (gene DFNB10) to chromosome 21q22.3. *Am J Hum Genet*. 1996;58(6):1254-9.
- Borck G, Ur Rehman A, Lee K, Pogoda HM, Kakar N, von Ameln S, Grillet N, Hildebrand MS, Ahmed ZM, N rnberg G, Ansar M, Basit S, Javed Q, Morell RJ, Nasreen N, Shearer AE, Ahmad A, Kahrizi K, Shaikh RS, Ali RA, Khan SN, Goebel I, Meyer NC, Kimberling WJ, Webster JA, Stephan DA, Schiller MR, Bahlo M, Najmabadi H, Gillespie PG, N rnberg P, Wollnik B, Riazuddin S, Smith RJ, Ahmad W, M ller U, Hammerschmidt M, Friedman TB, Riazuddin S, Leal SM, Ahmad J & Kubisch C. Loss-of-function mutations of *ILDR1* cause autosomal-recessive hearing impairment DFNB42. *Am J Hum Genet*. 2011;88(2):127-37.
- Bork JM, Peters LM, Riazuddin S, Bernstein SL, Ahmed ZM, Ness SL, Polomeno R, Ramesh A, Schloss M, Srisailpathy CR, Wayne S, Bellman S, Desmukh D, Ahmed Z, Khan SN, Kaloustian VM, Li XC, Lalwani A, Riazuddin S, Bitner-Glindzicz M, Nance WE, Liu XZ, Wistow G, Smith RJ, Griffith AJ, Wilcox ER, Friedman TB & Morell RJ. Usher syndrome 1D and nonsyndromic autosomal recessive deafness DFNB12 are caused by allelic mutations of the novel cadherin-like gene *CDH23*. *Am J Hum Genet*. 2001;68(1):26-37.
- Boughman JA, Vernon M & Shaver KA. Usher syndrome: definition and estimate of prevalence from two high-risk populations. *J Chronic Dis*. 1983;36(8):595-603.

- Chaib H, Place C, Salem N, Dodé C, Chardenoux S, Weissenbach J, el Zir E, Loiselet J & Petit C. Mapping of DFNB12, a gene for a non-syndromal autosomal recessive deafness, to chromosome 10q21-22. *Hum Mol Genet.* 1996;5(7):1061-4.
- Charizopoulou N, Lelli A, Schraders M, Ray K, Hildebrand MS, Ramesh A, Srisailapathy CR, Oostrik J, Admiraal RJ, Neely HR, Latoche JR, Smith RJ, Northup JK, Kremer H, Holt JR & Noben-Trauth K. *GIPC3* mutations associated with audiogenic seizures and sensorineural hearing loss in mouse and human. *Nat Commun.* 2011;2:201.
- Chen A, Francis M, Ni L, Cremers CW, Kimberling WJ, Sato Y, Phelps PD, Bellman SC, Wagner MJ & Pembrey M. Phenotypic manifestations of branchio-oto-renal syndrome. *Am J Med Genet.* 1995;58(4):365-70.
- Chen W, Kahrizi K, Meyer NC, Riazalhosseini Y, Van Camp G, Najmabadi H & Smith RJ. Mutation of *COL11A2* causes autosomal recessive non-syndromic hearing loss at the DFNB53 locus. *J Med Genet.* 2005 Oct;42(10):e61.
- Cheng J, Zhu Y, He S, Lu Y, Chen J, Han B, Petrillo M, Wrzeszczynski KO, Yang S, Dai P, Zhai S, Han D, Zhang MQ, Li W, Liu X, Li H, Chen ZY & Yuan H. Functional mutation of *SMAC/DIABLO*, encoding a mitochondrial proapoptotic protein, causes human progressive hearing loss DFNA64. *Am J Hum Genet.* 2011;89(1):56-66.
- Choi BY, Ahmed ZM, Riazuddin S, Bhinder MA, Shahzad M, Husnain T, Riazuddin S, Griffith AJ & Friedman TB. Identities and frequencies of mutations of the otoferlin gene (*OTOF*) causing DFNB9 deafness in Pakistan. *Clin Genet.* 2009;75(3):237-43.
- Collin RW, Kalay E, Tariq M, Peters T, van der Zwaag B, Venselaar H, Oostrik J, Lee K, Ahmed ZM, Caylan R, Li Y, Spierenburg HA, Eyupoglu E, Heister A, Riazuddin S, Bahat E, Ansar M, Arslan S, Wollnik B, Brunner HG, Cremers CW, Karaguzel A, Ahmad W, Cremers FP, Vriend G, Friedman TB, Riazuddin S, Leal SM & Kremer H. Mutations of *ESRRB* encoding estrogen-related receptor beta cause autosomal-recessive nonsyndromic hearing impairment DFNB35. *Am J Hum Genet.* 2008;82(1):125-38.
- Collin RW, Pauw RJ, Schoots J, Huygen PL, Hoefsloot LH, Cremers CW & Kremer H. Identification of a novel *COCH* mutation, G87W, causing autosomal dominant hearing impairment (DFNA9). *Am J Med Genet A.* 2006;140(16):1791-4.
- Colosimo A, Guida V, Rigoli L, Di Bella C, De Luca A, Briuglia S, Stuppia L, Salpietro DC & Dallapiccola B. Molecular detection of novel *WFS1* mutations in patients with Wolfram syndrome by a DHPLC-based assay. *Hum Mutat.* 2003;21(6):622-9.
- Coucke PJ, Van Hauwe P, Kelley PM, Kunst H, Schatteman I, Van Velzen D, Meyers J, Ensink RJ, Verstreken M, Declau F, Marres H, Kastury K, Bhasin S, McGuirt WT, Smith RJ, Cremers CW, Van de Heyning P, Willems PJ, Smith SD & Van Camp G. Mutations in the *KCNQ4* gene are responsible for autosomal dominant deafness in four DFNA2 families. *Hum Mol Genet.* 1999;8(7):1321-8.
- Craig R & Woodhead JL. Structure and function of myosin filaments. *Curr Opin Struct Biol.* 2006;16(2):204-12.
- Cryns K, Sivakumaran TA, Van den Ouweland JM, Pennings RJ, Cremers CW, Flothmann K, Young TL, Smith RJ, Lesperance MM & Van Camp G. Mutational spectrum of the *WFS1* gene in Wolfram syndrome, nonsyndromic hearing impairment, diabetes mellitus, and psychiatric disease. *Hum Mutat.* 2003;22(4):275-87.
- Davletov BA & Südhof TC. A single C2 domain from synaptotagmin I is sufficient for high affinity Ca²⁺/phospholipid binding. *J Biol Chem* 1993;268(35):26386-90.

- de Grey ADNJ. Life Span Extension Research and Public Debate: Societal Considerations. *Studies in Ethics, Law, and Technology* 1 2007.doi:10.2202/1941-6008.1011.
- de Kok YJ, van der Maarel SM, Bitner-Glindzicz M, Huber I, Monaco AP, Malcolm S, Pembrey ME, Ropers HH & Cremers FP. Association between X-linked mixed deafness and mutations in the POU domain gene *POU3F4*. *Science* 1995;267(5198):685-8.
- del Castillo FJ, Rodríguez-Ballesteros M, Alvarez A, Hutchin T, Leonardi E, de Oliveira CA, Azaiez H, Brownstein Z, Avenarius MR, Marlin S, Pandya A, Shahin H, Siemering KR, Weil D, Wuyts W, Aguirre LA, Martín Y, Moreno-Pelayo MA, Villamar M, Avraham KB, Dahl HH, Kanaan M, Nance WE, Petit C, Smith RJ, Van Camp G, Sartorato EL, Murgia A, Moreno F & del Castillo I. A novel deletion involving the connexin-30 gene, del(*GJB6*-d13s1854), found in trans with mutations in the *GJB2* gene (connexin-26) in subjects with DFNB1 non-syndromic hearing impairment. *J Med Genet*. 2005;42(7):588-94.
- del Castillo I, Villamar M, Moreno-Pelayo MA, del Castillo FJ, Alvarez A, Tellería D, Menéndez I & Moreno F. A deletion involving the connexin 30 gene in nonsyndromic hearing impairment. *N Engl J Med*. 2002;346(4):243-9.
- del Castillo I, Villamar M, Sarduy M, Romero L, Herraiz C, Hernández FJ, Rodríguez M, Borrás I, Montero A, Bellón J, Tapia MC & Moreno F. A novel locus for non-syndromic sensorineural deafness (DFN6) maps to chromosome Xp22. *Hum Mol Genet*. 1996;5(9):1383-7.
- Delmaghani S, del Castillo FJ, Michel V, Leibovici M, Aghaie A, Ron U, Van Laer L, Ben-Tal N, Van Camp G, Weil D, Langa F, Lathrop M, Avan P & Petit C. Mutations in the gene encoding pejvakin, a newly identified protein of the afferent auditory pathway, cause DFNB59 auditory neuropathy. *Nat Genet*. 2006;38(7):770-8.
- Di Palma F, Pellegrino R & Noben-Trauth K. Genomic structure, alternative splice forms and normal and mutant alleles of cadherin 23 (*CDH23*). *Gene*. 2001;281(1-2):31-41.
- Donaudy F, Ferrara A, Esposito L, Hertzano R, Ben-David O, Bell RE, Melchionda S, Zelante L, Avraham KB & Gasparini P. Multiple mutations of *MYO1A*, a cochlear-expressed gene, in sensorineural hearing loss. *Am J Hum Genet*. 2003;72(6):1571-7.
- Donaudy F, Snoeckx R, Pfister M, Zenner HP, Blin N, Di Stazio M, Ferrara A, Lanzara C, Ficarella R, Declau F, Pusch CM, Nürnberg P, Melchionda S, Zelante L, Ballana E, Estivill X, Van Camp G, Gasparini P & Savoia A. Nonmuscle myosin heavy-chain gene *MYH14* is expressed in cochlea and mutated in patients affected by autosomal dominant hearing impairment (DFNA4). *Am J Hum Genet*. 2004;74(4):770-6.
- Dossena S, Rodighiero S, Vezzoli V, Nofziger C, Salvioni E, Boccazzi M, Grabmayer E, Bottà G, Meyer G, Fugazzola L, Beck-Peccoz P & Paulmichl M. Functional characterization of wild-type and mutated pendrin (*SLC26A4*), the anion transporter involved in Pendred syndrome. *J Mol Endocrinol*. 2009;43(3):93-103.
- Dror AA & Avraham KB. Hearing loss: mechanisms revealed by genetics and cell biology. *Annu Rev Genet*. 2009;43:411-37.
- Ebermann I, Phillips JB, Liebau MC, Koenekoop RK, Schermer B, Lopez I, Schäfer E, Roux AF, Dafinger C, Bernd A, Zrenner E, Claustres M, Blanco B, Nürnberg G, Nürnberg P, Ruland R, Westerfield M, Benzing T & Bolz HJ. *PDZD7* is a modifier of retinal disease and a contributor to digenic Usher syndrome. *J Clin Invest*. 2010;120(6):1812-23.

- Ebermann I, Scholl HP, Charbel Issa P, Becirovic E, Lamprecht J, Jurklies B, Millán JM, Aller E, Mitter D & Bolz H. A novel gene for Usher syndrome type 2: mutations in the long isoform of whirlin are associated with retinitis pigmentosa and sensorineural hearing loss. *Hum Genet.* 2007;121(2):203-11.
- Edery P, Attié T, Amiel J, Pelet A, Eng C, Hofstra RM, Martelli H, Bidaud C, Munnich A & Lyonnet S. Mutation of the endothelin-3 gene in the Waardenburg-Hirschsprung disease (Shah-Waardenburg syndrome). *Nat Genet.* 1996;12(4):442-4.
- Emery A. Hereditary. Rimoin DL, Connor JM, Pyeritz RE, Korf BR. In: Emery and Rimoin's principles and practice of Medical Genetics. Churchill Livingstone, 5th edition. 2003; 1:243-244.
- Estivill X, Fortina P, Surrey S, Rabionet R, Melchionda S, D'Agruma L, Mansfield E, Rappaport E, Govea N, Milà M, Zelante L & Gasparini P. Connexin-26 mutations in sporadic and inherited sensorineural deafness. *Lancet.* 1998;351: 394-398.
- Eudy JD, Weston MD, Yao S, Hoover DM, Rehm HL, Ma-Edmonds M, Yan D, Ahmad I, Cheng JJ, Ayuso C, Cremers C, Davenport S, Moller C, Talmadge CB, Beisel KW, Tamayo M, Morton CC, Swaroop A, Kimberling WJ & Sumegi J. Mutation of a gene encoding a protein with extracellular matrix motifs in Usher syndrome type IIa. *Science.* 1998;280(5370):1753-7.
- Everett LA, Glaser B, Beck JC, Idol JR, Buchs A, Heyman M, Adawi F, Hazani E, Nassir E, Baxevanis AD, Sheffield VC & Green ED. Pendred syndrome is caused by mutations in a putative sulphate transporter gene (PDS). *Nat Genet.* 1997;17(4):411-22.
- Finsterer J & Fellingner J. Nuclear and mitochondrial genes mutated in nonsyndromic impaired hearing. *Int J Pediatr Otorhinolaryngol.* 2005;69(5):621-47.
- Fischel-Ghodsian N. Mitochondrial deafness mutations reviewed. *Hum Mutat.* 1999;13:261-270.
- Fonseca SG, Ishigaki S, Osowski CM, Lu S, Lipson KL, Ghosh R, Hayashi E, Ishihara H, Oka Y, Permutt MA, Urano F. Wolfram syndrome 1 gene negatively regulates ER stress signaling in rodent and human cells. *J Clin Invest.* 2010;120(3):744-55.
- Fraser Gr. Association of congenital deafness with gitre (Pendred's syndrome) a study of 207 families. *Ann Hum Genet.* 1965;28:201-49.
- Friedman RA, Van Laer L, Huentelman MJ, Sheth SS, Van Eyken E, Corneveaux JJ, Tembe WD, Halperin RF, Thorburn AQ, Thys S, Bonneux S, Fransen E, Huyghe J, Pyykkö I, Cremers CW, Kremer H, Dhooge I, Stephens D, Orzan E, Pfister M, Bille M, Parving A, Sorri M, Van de Heyning PH, Makmura L, Ohmen JD, Linthicum FH Jr, Fayad JN, Pearson JV, Craig DW, Stephan DA & Van Camp G. *GRM7* variants confer susceptibility to age-related hearing impairment. *Hum Mol Genet.* 2009;18(4):785-96
- Friedman TB, Schultz JM, Ben-Yosef T, Pryor SP, Lagziel A, Fisher RA, Wilcox ER, Riazuddin S, Ahmed ZM, Belyantseva IA & Griffith AJ. Recent advances in the understanding of syndromic forms of hearing loss. *Ear Hear* 2003;24(4):289-302.
- Friedman TB, Sellers JR & Avraham KB. Unconventional myosins and the genetics of hearing loss. *Med Genet.* 1999;89:147-157.
- Fukuoka H, Kanda Y, Ohta S & Usami S. Mutations in the *WFS1* gene are a frequent cause of autosomal dominant nonsyndromic low-frequency hearing loss in Japanese. *J Hum Genet.* 2007;52(6):510-5.

- Gal A, Pentelenyi K, Remenyi V, Pal Z, Csanyi B, Tomory G, Rasko I & Molnar MJ. Novel heteroplasmic mutation in the anticodon stem of mitochondrial tRNA(Lys) associated with dystonia and stroke-like episodes. *Acta Neurol Scand.* 2010;122(4):252-6.
- Gómez-Zaera M, Strom TM, Rodríguez B, Estivill X, Meitinger T & Nunes V. Presence of a major *WFS1* mutation in Spanish Wolfram syndrome pedigrees. *Mol Genet Metab.* 2001;72(1):72-81.
- Gorlin RJ. Genetic hearing loss—a brief history. Toriello HV, Reardon W, Gorlin RJ. In: *Hereditary Hearing Loss and Its Syndromes. Second Edition*, Oxf University Press, Baltimore 2004.
- Goto Y, Nonaka I & Horai S. A mutation in the tRNA^{Leu}(UUR) gene associated with the MELAS subgroup of mitochondrial encephalomyopathies. *Nature.* 1990; 348:651-653.
- Greinwald JH Jr, Scott DA, Marietta JR, Carmi R, Manaligod J, Ramesh A, Zbar RI, Kraft ML, Elbedour K, Yairi Y, Musy M, Skvorak AB, Van Camp G, Srisailapathy CR, Lovett M, Morton CC, Sheffield VC & Smith RJ. Construction of P1-derived artificial chromosome and yeast artificial chromosome contigs encompassing the DFNB7 and DFNB11 region of chromosome 9q13-21. *Genome Res.* 1997;7(9):879-86.
- Grifa A, Wagner CA, D'Ambrosio L, Melchionda S, Bernardi F, Lopez-Bigas N, Rabionet R, Arbones M, Monica MD, Estivill X, Zelante L, Lang F & Gasparini P. Mutations in *GJB6* cause nonsyndromic autosomal dominant deafness at DFNA3 locus. *Nat Genet.* 1999;23(1):16-8.
- Grillet N, Schwander M, Hildebrand MS, Sczaniecka A, Kolatkar A, Velasco J, Webster JA, Kahrizi K, Najmabadi H, Kimberling WJ, Stephan D, Bahlo M, Wiltshire T, Tarantino LM, Kuhn P, Smith RJ & Müller U. Mutations in *LOXHD1*, an evolutionarily conserved stereociliary protein, disrupt hair cell function in mice and cause progressive hearing loss in humans. *Am J Hum Genet.* 2009;85(3):328-37.
- Guan MX. Mitochondrial 12S rRNA mutations associated with aminoglycoside ototoxicity. *Mitochondrion.* 2011;11(2):237-45.
- Gubler MC, Heidet L & Antignac C. Alport's syndrome, thin basement membrane nephropathy, nail-patella syndrome, and type III collagen glomerulopathy. In: Jennette JC, Olson JL, Schwartz MM, Silva FG, eds. *Heptinstall's Pathology of the Kidney*. 6th ed. Philadelphia, Pa: LippincottWilliams & Wilkins; 2007:487-515.
- Guilford P, Ben Arab S, Blanchard S, Levilliers J, Weissenbach J, Belkahia A & Petit C. A non-syndrome form of neurosensory, recessive deafness maps to the pericentromeric region of chromosome 13q. *Nat Genet.* 1994;6(1):24-8.
- Guipponi M, Antonarakis SE & Scott HS. *TMPRSS3*, a type II transmembrane serine protease mutated in non-syndromic autosomal recessive deafness. *Front Biosci.* 2008;13:1557-67.
- Guipponi M, Vuagniaux G, Wattenhofer M, Shibuya K, Vazquez M, Dougherty L, Scamuffa N, Guida E, Okui M, Rossier C, Hancock M, Buchet K, Reymond A, Hummler E, Marzella PL, Kudoh J, Shimizu N, Scott HS, Antonarakis SE & Rossier BC. The transmembrane serine protease (*TMPRSS3*) mutated in deafness DFNB8/10 activates the epithelial sodium channel (ENaC) in vitro. *Hum Mol Genet.* 2002;11(23):2829-36.

- Gumbiner BM. Regulation of cadherin-mediated adhesion in morphogenesis. *Nat Rev Mol Cell Biol* 2005;6:622-634.
- Hilgert N, Alasti F, Dieltjens N, Pawlik B, Wollnik B, Uyguner O, Delmaghani S, Weil D, Petit C, Danis E, Yang T, Pandelia E, Petersen MB, Goossens D, Favero JD, Sanati MH, Smith RJ & Van Camp G. Mutation analysis of *TMC1* identifies four new mutations and suggests an additional deafness gene at loci DFNA36 and DFNB7/11. *Clin Genet*. 2008;74(3):223-32.
- Hilgert N, Huentelman MJ, Thorburn AQ, Fransen E, Dieltjens N, Mueller-Malesinska M, Pollak A, et al. Phenotypic variability of patients homozygous for the GJB2 mutation 35delG cannot be explained by the influence of one major modifier gene. *Eur J Hum Genet*. 2009a Apr;17(4):517-24.
- Hilgert N, Smith RJ & Van Camp G. Forty-six genes causing nonsyndromic hearing impairment: which ones should be analyzed in DNA diagnostics? *Mutat Res* 2009b;681(2-3):189-96.
- Hofmann S, Philbrook C, Gerbitz KD & Bauer MF. Wolfram syndrome: structural and functional analyses of mutant and wild-type wolframin, the *WFS1* gene product. *Hum Mol Genet*. 2003;12(16):2003-12.
- Hofstra RM, Osinga J, Tan-Sindhunata G, Wu Y, Kamsteeg EJ, Stulp RP, van Ravenswaaij-Arts C, Majoor-Krakauer D, Angrist M, Chakravarti A, Meijers C, Buys CH. A homozygous mutation in the endothelin-3 gene associated with a combined Waardenburg type 2 and Hirschsprung phenotype (Shah-Waardenburg syndrome). *Nat Genet*. 1996;12(4):445-7.
- Holms RH & Steel KP. Genes involved in deafness. *Curr Opin Gene Dev*. 1999; 9:309-314.
- Hope CI, Bunday S, Proops D & Fielder AR. Usher syndrome in the city of Birmingham--prevalence and clinical classification. *Br J Ophthalmol*. 1997;81(1):46-53.
- Hoskins BE, Cramer CH, Silvius D, Zou D, Raymond RM, Orten DJ, Kimberling WJ, Smith RJ, Weil D, Petit C, Otto EA, Xu PX & Hildebrandt F. Transcription factor SIX5 is mutated in patients with branchio-oto-renal syndrome. *Am J Hum Genet*. 2007;80(4):800-4.
- Hu H, Wu L, Feng Y, Pan Q, Long Z, Li J, Dai H, Xia K, Liang D, Niikawa N & Xia J. Molecular analysis of hearing loss associated with enlarged vestibular aqueduct in the mainland Chinese: a unique *SLC26A4* mutation spectrum. *J Hum Genet*. 2007;52(6):492-7.
- Huang Q & Tang J. Age-related hearing loss or presbycusis. *Eur Arch Otorhinolaryngol*. 2010;267(8):1179-91.
- Hudson BG, Tryggvason K, Sundaramoorthy M & Neilson EG. Alport's syndrome, Goodpasture's syndrome, and type IV collagen. *N Engl J Med*. 2003;348(25):2543-56.
- Jaksch M, Klopstock T, Kurlmann G, et al. Progressive myoclonus epilepsy and mitochondrial myopathy associated with mutations in the tRNASer(UCN) gene. *Ann Neurol*. 1998; 44:635- 640.
- Jamora C & Fuchs E. Intercellular adhesion, signalling and the cytoskeleton. *Nat Cell Biol*. 2002;4(4):E101-8.
- Joensuu T, Hämäläinen R, Yuan B, Johnson C, Tegelberg S, Gasparini P, Zelante L, Pirvola U, Pakarinen L, Lehesjoki AE, de la Chapelle A & Sankila EM. Mutations in a novel

- gene with transmembrane domains underlie Usher syndrome type 3. *Am J Hum Genet.* 2001;69(4):673-84.
- Johnson KR, Yu H, Ding D, Jiang H, Gagnon LH & Salvi RJ. Separate and combined effects of *Sod1* and *CDH23* mutations on age-related hearing loss and cochlear pathology in C57BL/6J mice. *Hear Res.* 2010;268(1-2):85-92.
- Kalay E, Uzumcu A, Krieger E, Caylan R, Uyguner O, Ulubil-Emiroglu M, Erdol H, Kayserili H, Hafiz G, Başerer N, Heister AJ, Hennies HC, Nürnberg P, Başaran S, Brunner HG, Cremers CW, Karaguzel A, Wollnik B & Kremer H. *MYO15A* (DFNB3) mutations in Turkish hearing loss families and functional modeling of a novel motor domain mutation. *Am J Med Genet A.* 2007;143A(20):2382-9.
- Kameoka K, Isotani H, Tanaka K, et al. Novel mitochondrial DNA mutation in tRNA(Lys) (8296A→G) associated with diabetes. *Biochem Biophys Res Commun.* 1998;245:523-527.
- Kapur RP. Early death of neural crest cells is responsible for total enteric aganglionosis in *SOX10(Dom)/SOX10(Dom)* mouse embryos. *Pediatr Dev Pathol.* 1999;2(6):559-69.
- Kelley PM, Harris DJ, Comer BC, Askew JW, Fowler T, Smith SD & Kimberling WJ. Novel mutations in the connexin 26 gene (*GJB2*) that cause autosomal recessive (DFNB1) hearing loss. *Am J Hum Genet.* 1998; 62: 792-799.
- Kelsell DP, Dunlop J, Stevens HP, Lench NJ, Liang JN, Parry G, Mueller RF & Leigh IM. Connexin 26 mutations in hereditary non-syndromic sensorineural deafness. *Nature.* 1997;387(6628):80-3.
- Khan SY, Ahmed ZM, Shabbir MI, Kitajiri S, Kalsoom S, Tasneem S, Shayiq S, Ramesh A, Srisailpathy S, Khan SN, Smith RJ, Riazuddin S, Friedman TB & Riazuddin S. Mutations of the *RDX* gene cause nonsyndromic hearing loss at the DFNB24 locus. *Hum Mutat.* 2007;28(5):417-23.
- Kimberling WJ, Weston MD, Möller C, Davenport SL, Shugart YY, Priluck IA, Martini A, Milani M & Smith RJ. Localization of Usher syndrome type II to chromosome 1q. *Genomics.* 1990;7(2):245-9.
- Klein D. Historical background and evidence for dominant inheritance of the Klein-Waardenburg syndrome (type III). *Am J Med Genet.* 1983;14(2):231-9.
- Kopp P, Pesce L & Solis-S JC. Pendred syndrome and iodide transport in the thyroid. *Trends Endocrinol Metab.* 2008;19(7):260-8.
- Kubisch C, Schroeder BC, Friedrich T, Lütjohann B, El-Amraoui A, Marlin S, Petit C & Jentsch TJ. *KCNQ4*, a novel potassium channel expressed in sensory outer hair cells, is mutated in dominant deafness. *Cell* 1999;96(3):437-46.
- Kumar NM & Gilula NB. The gap Junction communication channel. *Cell.* 1996; 84:381-388.
- Kurima K, Peters LM, Yang Y, Riazuddin S, Ahmed ZM, Naz S, Arnaud D, Drury S, Mo J, Makishima T, Ghosh M, Menon PS, Deshmukh D, Oddoux C, Ostrer H, Khan S, Riazuddin S, Deininger PL, Hampton LL, Sullivan SL, Battey JF Jr, Keats BJ, Wilcox ER, Friedman TB & Griffith AJ. Dominant and recessive deafness caused by mutations of a novel gene, *TMC1*, required for cochlear hair-cell function. *Nat Genet.* 2002;30(3):277-84.
- Lalwani AK, Goldstein JA, Kelley MJ, Luxford W, Castelein CM & Mhatre AN. Human nonsyndromic hereditary deafness DFNA17 is due to a mutation in nonmuscle myosin *MYH9*. *Am J Hum Genet.* 2000;67(5):1121-8.

- Lemmink HH, Mochizuki T, van den Heuvel LP, Schröder CH, Barrientos A, Monnens LA, van Oost BA, Brunner HG, Reeders ST & Smeets HJ. Mutations in the type IV collagen alpha 3 (*COL4A3*) gene in autosomal recessive Alport syndrome. *Hum Mol Genet.* 1994;3(8):1269-73.
- Leon PE, Raventos H, Lynch E, Morrow J & King MC. The gene for an inherited form of deafness maps to chromosome 5q31. *Proc Natl Acad Sci U S A.* 1992;89(11):5181-4.
- Lezirovitz K, Pardon E, de Mello Auricchio MTB, de Carvalho e Silva FL, Lopes JJ, Abreu-Silva RS, Romanos J, Batisso AC & Mingroni-Netto RC. Unexpected genetic heterogeneity in a large consanguineous Brazilian pedigree presenting deafness. *Eur J Hum Genet.* 2008;16:89-96.
- Li XC, Everett LA, Lalwani AK, Desmukh D, Friedman TB, Green ED & Wilcox ER. A mutation in *PDS* causes non-syndromic recessive deafness. *Nat Genet.* 1998;18(3):215-7.
- Li Z, Li R, Chen J, et al. Mutational analysis of the mitochondrial 12S rRNA gene in Chinese pediatric subjects with aminoglycoside-induced and nonsyndromic hearing loss. *Hum Genet.* 2005; 117:9-15.
- Liang Y, Wang A, Belyantseva IA, Anderson DW, Probst FJ, Barber TD, Miller W, Touchman JW, Jin L, Sullivan SL, Sellers JR, Camper SA, Lloyd RV, Kachar B, Friedman TB & Fridell RA. Characterization of the human and mouse unconventional myosin XV genes responsible for hereditary deafness *DFNB3* and *shaker 2*. *Genomics.* 1999;61:243-258.
- Liu X, Han D, Li J, Han B, Ouyang X, Cheng J, Li X, Jin Z, Wang Y, Bitner-Glindzicz M, Kong X, Xu H, Kantardzhieva A, Eavey RD, Seidman CE, Seidman JG, Du LL, Chen ZY, Dai P, Teng M, Yan D & Yuan H. Loss-of-function mutations in the *PRPS1* gene cause a type of nonsyndromic X-linked sensorineural deafness, *DFN2*. *Am J Hum Genet.* 2010;86(1):65-71.
- Liu XZ, Ouyang XM, Xia XJ, Zheng J, Pandya A, Li F, Du LL, Welch KO, Petit C, Smith RJ, Webb BT, Yan D, Arnos KS, Corey D, Dallos P, Nance WE & Chen ZY. Prestin, a cochlear motor protein, is defective in non-syndromic hearing loss. *Hum Mol Genet.* 2003;12(10):1155-62.
- Liu XZ, Walsh J, Tamagawa Y, Kitamura K, Nishizawa M, Steel KP & Brown SD. Autosomal dominant non-syndromic deafness caused by a mutation in the myosin VIIA gene. *Nat Genet.* 1997;17(3):268-9.
- Loikkanen I, Toljamo K, Hirvikoski P, Väisänen T, Paavonen TK & Vaarala MH. Myosin VI is a modulator of androgen-dependent gene expression. *Oncol Rep.* 2009;22(5):991-5.
- Lynch ED, Lee MK, Morrow JE, Welch PL, León PE & King MC. Nonsyndromic deafness *DFNA1* associated with mutation of a human homolog of the *Drosophila* gene *diaphanous*. *Science.* 1997;278(5341):1315-8.
- Mahdieh N & Rabbani B. Statistical study of 35delG mutation of *GJB2* gene: a meta-analysis of carrier frequency. *Int J Audiol.* 2009;48(6):363-70.
- Mahdieh N, Bagherian H, Shirkavand A, Sharafi M & Zeinali S. High level of intrafamilial phenotypic variability of non-syndromic hearing loss in a Lur family due to delE120 mutation in *GJB2* gene. *Int J Pediatr Otorhinolaryngol.* 2010b;74:1089-1091.
- Mahdieh N, Nishimura C, Ali-Madadi K, Riazalhosseini Y, Yazdan H, Arzhanghi S, Jalalvand K, Ebrahimi A, Kazemi S, Smith RJ & Najmabadi H. The frequency of *GJB2*

- mutations and the Delta (*GJB6-D13S1830*) deletion as a cause of autosomal recessive non-syndromic deafness in the Kurdish population. *Clin Genet*. 2004;65(6):506-8.
- Mahdieh N, Rabbani B, Shirkavand A, Bagherian H, Movahed ZS, Fouladi P, Rahiminejad F, Masoudifard M, Akbari MT & Zeinali S. Impact of consanguineous marriages in *GJB2*-related hearing loss in the Iranian population: a report of a novel variant. *Genet Test Mol Biomarkers*. 2011;15(7-8):489-93.
- Mahdieh N, Rabbani B, Wiley S, Akabari MT & Zeinali S. Genetic causes of nonsyndromic hearing loss in Iran in comparison with other populations. *J Hum Genet*. 2010a;55(10):639-48.
- Mahdieh N, Shirkavand A, Raeisi M, Akbari MT & Zeinali S. Unexpected heterogeneity due to recessive and de novo dominant mutations of *GJB2* in an Iranian family with nonsyndromic hearing loss: implication for genetic counseling. *Biochem Biophys Res Commun* 2010c;402(2):305-7.
- Mburu P, Mustapha M, Varela A, Weil D, El-Amraoui A, Holme RH, Rump A, Hardisty RE, Blanchard S, Coimbra RS, Perfettini I, Parkinson N, Mallon AM, Glenister P, Rogers MJ, Paige AJ, Moir L, Clay J, Rosenthal A, Liu XZ, Blanco G, Steel KP, Petit C & Brown SD. Defects in whirlin, a PDZ domain molecule involved in stereocilia elongation, cause deafness in the whirler mouse and families with DFNB31. *Nat Genet*. 2003;34(4):421-8.
- McGuirt WT, Prasad SD, Griffith AJ, Kunst HP, Green GE, Shpargel KB, Runge C, Huybrechts C, Mueller RF, Lynch E, King MC, Brunner HG, Cremers CW, Takanosu M, Li SW, Arita M, Mayne R, Prockop DJ, Van Camp G & Smith RJ. Mutations in *COL11A2* cause non-syndromic hearing loss (DFNA13). *Nat Genet*. 1999 Dec;23(4):413-9.
- Melchionda S, Ahituv N, Bisceglia L, Sobe T, Glaser F, Rabionet R, Arbones ML, Notarangelo A, Di Iorio E, Carella M, Zelante L, Estivill X, Avraham KB & Gasparini P. *MYO6*, the human homologue of the gene responsible for deafness in Snell's waltzer mice, is mutated in autosomal dominant nonsyndromic hearing loss. *Am J Hum Genet*. 2001;69(3):635-40.
- Mencia A, Modamio-Høybjør S, Redshaw N, Morín M, Mayo-Merino F, Olavarrieta L, Aguirre LA, del Castillo I, Steel KP, Dalmay T, Moreno F & Moreno-Pelayo MA. Mutations in the seed region of human miR-96 are responsible for nonsyndromic progressive hearing loss. *Nat Genet*. 2009;41(5):609-13.
- Merchant SN, Burgess BJ, Adams JC, Kashtan CE, Gregory MC, Santi PA, Colvin R, Collins B & Nadol JB Jr. Temporal bone histopathology in alport syndrome. *Laryngoscope*. 2004;114(9):1609-18.
- Migliosi V, Modamio-Høybjør S, Moreno-Pelayo MA, Rodríguez-Ballesteros M, Villamar M, Tellería D, Menéndez I, Moreno F & Del Castillo I. Q829X, a novel mutation in the gene encoding *OTOFerlin (OTOF)*, is frequently found in Spanish patients with prelingual non-syndromic hearing loss. *J Med Genet*. 2002;39(7):502-6.
- Mirghomizadeh F, Pfister M, Apaydin F, Petit C, Kupka S, Pusch CM, Zenner HP & Blin N. Substitutions in the conserved C2C domain of *OTOFerlin* cause DFNB9, a form of nonsyndromic autosomal recessive deafness. *Neurobiol Dis*. 2002;10(2):157-64.
- Modamio-Hoybjør S, Mencia A, Goodyear R, del Castillo I, Richardson G, Moreno F & Moreno-Pelayo MA. A mutation in *CCDC50*, a gene encoding an effector of

- epidermal growth factor-mediated cell signaling, causes progressive hearing loss. *Am J Hum Genet.* 2007;80(6):1076-89.
- Morell RJ, Kim HJ, Hood LJ, Goforth L, Friderici K, Fisher R, Van Camp G, Berlin CI, Oddoux C, Ostrer H, Keats B & Friedman TB. Mutations in the connexin 26 gene (*GJB2*) among Ashkenazi Jews with nonsyndromic recessive deafness. *N Engl J Med.* 1998 Nov 19;339(21):1500-5.
- Morle L, Bozon M, Alloisio N, Latour P, Vandenberghe A, Plauchu H, Collet L, Edery P, Godet J & Lina-Granade G. A novel C202F mutation in the connexin26 gene (*GJB2*) associated with autosomal dominant isolated hearing loss. *J Med Genet.* 2000; 37: 368-370.
- Morton CC & Nance WE. Newborn hearing screening--a silent revolution. *N Engl J Med.* 2006;354(20):2151-64.
- Mount DB & Romero MF. The SLC26 gene family of multifunctional anion exchangers. *Pflugers Archiv* 2004;447:710-721.
- Mustapha M, Chouery E, Torchard-Pagnez D, Nouaille S, Khrais A, Sayegh FN, Mégarbané A, Loiselet J, Lathrop M, Petit C & Weil D. A novel locus for Usher syndrome type I, *USH1G*, maps to chromosome 17q24-25. *Hum Genet.* 2002;110(4):348-50.
- Mustapha M, Weil D, Chardenoux S, Elias S, El-Zir E, Beckmann JS, Loiselet J & Petit C. An alpha-tectorin gene defect causes a newly identified autosomal recessive form of sensorineural pre-lingual non-syndromic deafness, *DFNB21*. *Hum Mol Genet.* 1999;8(3):409-12.
- Nal N, Ahmed ZM, Erkal E, Alper OM, Lüleci G, Dinç O, Waryah AM, Ain Q, Tasneem S, Husnain T, Chattaraj P, Riazuddin S, Boger E, Ghosh M, Kabra M, Riazuddin S, Morell RJ & Friedman TB. Mutational spectrum of *MYO15A*: the large N-terminal extension of myosin XVA is required for hearing. *Hum Mutat.* 2007;28(10):1014-9.
- Nance WE & Sweeney A. Symposium on sensorineural hearing loss in children: early detection and intervention. Genetic factors in deafness of early life. *Otolaryngol Clin North Am.* 1975;8(1):19-48.
- Naz S, Giguere CM, Kohrman DC, Mitchem KL, Riazuddin S, Morell RJ, Ramesh A, Srisailpathy S, Deshmukh D, Riazuddin S, Griffith AJ, Friedman TB, Smith RJ & Wilcox ER. Mutations in a novel gene, *TMIE*, are associated with hearing loss linked to the *DFNB6* locus. *Am J Hum Genet.* 2002;71(3):632-6.
- Naz S, Griffith AJ, Riazuddin S, Hampton LL, Battey JF Jr, Khan SN, Riazuddin S, Wilcox ER & Friedman TB. Mutations of *ESPN* cause autosomal recessive deafness and vestibular dysfunction. *J Med Genet.* 2004;41(8):591-5.
- Nelson WJ & Nusse R. Convergence of Wnt, beta-catenin, and cadherin pathways. *Science.* 2004;303(5663):1483-7.
- Newton V. Hearing loss and Waardenburg's syndrome: implications for genetic counselling. *J Laryngol Otol.* 1990;104(2):97-103.
- Nie L. *KCNQ4* mutations associated with nonsyndromic progressive sensorineural hearing loss. *Curr Opin Otolaryngol Head Neck Surg.* 2008;16(5):441-4.
- Oliver TN, Berg JS & Cheney RE. Tails of unconventional myosins. *Cell Mol Life Sci.* 1999;56:243-257.
- Ouyang XM, Xia XJ, Verpy E, Du LL, Pandya A, Petit C, Balkany T, Nance WE & Liu XZ. Mutations in the alternatively spliced exons of *USH1C* cause non-syndromic recessive deafness. *Hum Genet.* 2002 Jul;111(1):26-30.

- Pallares-Ruiz N, Blanchet P, Mondain M, Claustres M & Roux AF. A large deletion including most of *GJB6* in recessive non syndromic deafness: a digenic effect? *Eur J Hum Genet.* 2002;10(1):72-6.
- Pandya A, Xia XJ, Erdenetungalag R, et al. Heterogenous point mutations in the mitochondrial tRNA^{Ser}(UCN) precursor coexisting with the A1555G mutation in deafness students from Mongolia. *Am J Hum Genet.* 1999;65:1803-1806.
- Peters LM, Anderson DW, Griffith AJ, Grundfast KM, San Agustin TB, Madeo AC, Friedman TB & Morell RJ. Mutation of a transcription factor, *TFCP2L3*, causes progressive autosomal dominant hearing loss, *DFNA28*. *Hum Mol Genet.* 2002;11(23):2877-85.
- Pingault V, Bondurand N, Kuhlbrodt K, Goerich DE, Préhu MO, Puliti A, Herbarth B, Hermans-Borgmeyer I, Legius E, Matthijs G, Amiel J, Lyonnet S, Ceccherini I, Romeo G, Smith JC, Read AP, Wegner M & Goossens M. *SOX10* mutations in patients with Waardenburg-Hirschsprung disease. *Nat Genet.* 1998;18(2):171-3.
- Raphael Y & Altschuler RA. Structure and innervation of the cochlea. *Brain Res Bull.* 2003;60(5-6):397-422.
- Read AP & Newton VE. Waardenburg syndrome. *J Med Genet* 1997;34:656-665.
- Rehman AU, Gul K, Morell RJ, Lee K, Ahmed ZM, Riazuddin S, Ali RA, Shahzad M, Jaleel AU, Andrade PB, Khan SN, Khan S, Brewer CC, Ahmad W, Leal SM, Riazuddin S & Friedman TB. Mutations of *GIPC3* cause nonsyndromic hearing loss *DFNB72* but not *DFNB81* that also maps to chromosome 19p. *Hum Genet.* 2011 Jun 10. [Epub ahead of print]
- Rehman AU, Morell RJ, Belyantseva IA, Khan SY, Boger ET, Shahzad M, Ahmed ZM, Riazuddin S, Khan SN, Riazuddin S & Friedman TB. Targeted capture and next-generation sequencing identifies *C9orf75*, encoding taperin, as the mutated gene in nonsyndromic deafness *DFNB79*. *Am J Hum Genet.* 2010;86(3):378-88.
- Riazuddin S, Ahmed ZM, Fanning AS, Lagziel A, Kitajiri S, Ramzan K, Khan SN, Chattaraj P, Friedman PL, Anderson JM, Belyantseva IA, Forge A, Riazuddin S & Friedman TB. Tricellulin is a tight-junction protein necessary for hearing. *Am J Hum Genet.* 2006;79(6):1040-51.
- Riazuddin S, Anwar S, Fischer M, Ahmed ZM, Khan SY, Janssen AG, Zafar AU, Scholl U, Husnain T, Belyantseva IA, Friedman PL, Riazuddin S, Friedman TB & Fahlke C. Molecular basis of *DFNB73*: mutations of *BSND* can cause nonsyndromic deafness or Bartter syndrome. *Am J Hum Genet.* 2009;85(2):273-80.
- Riazuddin S, Khan SN, Ahmed ZM, Ghosh M, Caution K, Nazli S, Kabra M, Zafar AU, Chen K, Naz S, Antonellis A, Pavan WJ, Green ED, Wilcox ER, Friedman PL, Morell RJ, Riazuddin S & Friedman TB. Mutations in *TRIOBP*, which encodes a putative cytoskeletal-organizing protein, are associated with nonsyndromic recessive deafness. *Am J Hum Genet.* 2006;78(1):137-43.
- Richards AJ, Yates JR, Williams R, Payne SJ, Pope FM, Scott JD & Snead MP. A family with Stickler syndrome type 2 has a mutation in the *COL11A1* gene resulting in the substitution of glycine 97 by valine in alpha 1 (XI) collagen. *Hum Mol Genet.* 1996;5(9):1339-43.
- Richardson GP, Lukashkin AN & Russell IJ. The tectorial membrane: one slice of a complex cochlear sandwich. *Curr Opin Otolaryngol Head Neck Surg.* 2008;16(5):458-64.


- Rigoli L, Prisco F, Caruso RA, Iafusco D, Ursomanno G, Zuccarello D, Ingenito N, Rigoli M & Barberi I. Association of the T14709C mutation of mitochondrial DNA with maternally inherited diabetes mellitus and/or deafness in an Italian family. *Diabet Med.* 2001;18(4):334-6.
- Robertson NG, Lu L, Heller S, Merchant SN, Eavey RD, McKenna M, Nadol JB Jr, Miyamoto RT, Linthicum FH Jr, Lubianca Neto JF, Hudspeth AJ, Seidman CE, Morton CC & Seidman JG. Mutations in a novel cochlear gene cause DFNA9, a human nonsyndromic deafness with vestibular dysfunction. *Nat Genet.* 1998;20(3):299-303.
- Ruf RG, Xu PX, Silvius D, Otto EA, Beekmann F, Muerb UT, Kumar S, Neuhaus TJ, Kemper MJ, Raymond RM Jr, Brophy PD, Berkman J, Gattas M, Hyland V, Ruf EM, Schwartz C, Chang EH, Smith RJ, Stratakis CA, Weil D, Petit C & Hildebrandt F. SIX1 mutations cause branchio-oto-renal syndrome by disruption of EYA1-SIX1-DNA complexes. *Proc Natl Acad Sci U S A.* 2004;101(21):8090-5.
- Sahin-Calapoglu N, Calapoglu M & Karaguzel A. Non-syndromic recessive hearing loss linked to *TMPRSS3* gene in the Turkish population. *SDÜ Týp Fak Derg.* 2005;12(3):31-35.
- Schraders M, Lee K, Oostrik J, Huygen PL, Ali G, Hoefsloot LH, Veltman JA, Cremers FP, Basit S, Ansar M, Cremers CW, Kunst HP, Ahmad W, Admiraal RJ, Leal SM & Kremer H. Homozygosity mapping reveals mutations of *GRXCR1* as a cause of autosomal-recessive nonsyndromic hearing impairment. *Am J Hum Genet.* 2010;86(2):138-47.
- Schraders M, Oostrik J, Huygen PL, Strom TM, van Wijk E, Kunst HP, Hoefsloot LH, Cremers CW, Admiraal RJ & Kremer H. Mutations in *PTPRQ* are a cause of autosomal-recessive nonsyndromic hearing impairment DFNB84 and associated with vestibular dysfunction. *Am J Hum Genet.* 2010;86(4):604-10.
- Schultz JM, Khan SN, Ahmed ZM, Riazuddin S, Waryah AM, Chhatre D, Starost MF, Ploplis B, Buckley S, Velásquez D, Kabra M, Lee K, Hassan MJ, Ali G, Ansar M, Ghosh M, Wilcox ER, Ahmad W, Merlini G, Leal SM, Riazuddin S, Friedman TB & Morell RJ. Noncoding mutations of *HGF* are associated with nonsyndromic hearing loss, DFNB39. *Am J Hum Genet.* 2009;85(1):25-39.
- Scott HS, Kudoh J, Wattenhofer M, Shibuya K, Berry A, Chrast R, Guipponi M, Wang J, Kawasaki K, Asakawa S, Minoshima S, Younus F, Mehdi SQ, Radhakrishna U, Papasavvas MP, Gehrig C, Rossier C, Korostishevsky M, Gal A, Shimizu N, Bonne-Tamir B & Antonarakis SE. Insertion of beta-satellite repeats identifies a transmembrane protease causing both congenital and childhood onset autosomal recessive deafness. *Nat Genet.* 2001;27(1):59-63.
- Shabbir MI, Ahmed ZM, Khan SY, Riazuddin S, Waryah AM, Khan SN, Camps RD, Ghosh M, Kabra M, Belyantseva IA, Friedman TB & Riazuddin S. Mutations of human *TMHS* cause recessively inherited non-syndromic hearing loss. *J Med Genet.* 2006;43(8):634-40.
- Shah KN, Dalal SJ, Desai MP, Sheth PN, Joshi NC & Ambani LM. White forelock, pigmentary disorder of irides, and long segment Hirschsprung disease: possible variant of Waardenburg syndrome. *J Pediatr.* 1981;99(3):432-5.
- Sheffield VC, Kraiem Z, Beck JC, Nishimura D, Stone EM, Salameh M, Sadeh O & Glaser B. Pendred syndrome maps to chromosome 7q21-34 and is caused by an intrinsic defect in thyroid iodine organification. *Nat Genet.* 1996;12:424-426.

- Shin JJ, Keamy DG Jr & Steinberg EA. Medical and surgical interventions for hearing loss associated with congenital cytomegalovirus: a systematic review. *Otolaryngol Head Neck Surg.* 2011;144(5):662-75.
- Siemens J, Lillo C, Dumont RA, Reynolds A, Williams DS, Gillespie PG & Müller U. Cadherin 23 is a component of the tip link in hair-cell stereocilia. *Nature.* 2004;428(6986):950-5.
- Sirmaci A, Erbek S, Price J, Huang M, Duman D, Cengiz FB, Bademci G, Tokgöz-Yilmaz S, Hişmi B, Ozdağ H, Oztürk B, Kulaksizoğlu S, Yildirim E, Kokotas H, Grigoriadou M, Petersen MB, Shahin H, Kanaan M, King MC, Chen ZY, Blanton SH, Liu XZ, Zuchner S, Akar N & Tekin M. A truncating mutation in *SERPINB6* is associated with autosomal-recessive nonsyndromic sensorineural hearing loss. *Am J Hum Genet.* 2010;86(5):797-804.
- Smith RJ & Hone S. Genetic screening for deafness. *Pediatr Clin North Am.* 2003;50(2):315-29.
- Smith RJ, Berlin CI, Hejtmancik JF, Keats BJ, Kimberling WJ, Lewis RA, Möller CG, Pelias MZ & Tranebjaerg L. Clinical diagnosis of the Usher syndromes. Usher Syndrome Consortium. *Am J Med Genet.* 1994;50(1):32-8.
- Smith RJ, Lee EC, Kimberling WJ, Daiger SP, Pelias MZ, Keats BJ, Jay M, Bird A, Reardon W, Guest M, et al. Localization of two genes for Usher syndrome type I to chromosome 11. *Genomics.* 1992;14(4):995-1002.
- Stevenson VA, Ito M & Milunsky JM. Connexin-30 deletion analysis in connexin-26 heterozygotes. *Genet Test.* 2003;7:151-4.
- Szabo R, Wu Q, Dickson RB, Netzel-Arnett S, Antalis TM & Bugge TH. Type II transmembrane serine proteases. *Thromb Haemost.* 2003; 90(2):185-93.
- Tassabehji M, Newton VE & Read AP. Waardenburg syndrome type 2 caused by mutations in the human microphthalmia (*MITF*) gene. *Nat Genet.* 1994;8(3):251-5.
- Tekin M, Arnos KS & Pandya A. Advances in hereditary deafness. *Lancet* 2001;358(9287):1082-90.
- Unal M, Tamer L, Doğruer ZN, Yildirim H, Vayisoğlu Y & Camdeviren H. N-acetyltransferase 2 gene polymorphism and presbycusis. *Laryngoscope.* 2005;115(12):2238-41.
- Vahava O, Morell R, Lynch ED, Weiss S, Kagan ME, Ahituv N, Morrow JE, Lee MK, Skvorak AB, Morton CC, Blumenfeld A, Frydman M, Friedman TB, King MC & Avraham KB. Mutation in transcription factor *POU4F3* associated with inherited progressive hearing loss in humans. *Science.* 1998;279(5358):1950-4.
- Van Camp G, Snoeckx RL, Hilgert N, van den Ende J, Fukuoka H, Wagatsuma M, Suzuki H, Smets RM, Vanhoenacker F, Declau F, Van de Heyning P & Usami S. A new autosomal recessive form of Stickler syndrome is caused by a mutation in the *COL9A1* gene. *Am J Hum Genet.* 2006;79(3):449-57.
- van der Loop FT, Heidet L, Timmer ED, van den Bosch BJ, Leinonen A, Antignac C, Jefferson JA, Maxwell AP, Monnens LA, Schröder CH & Smeets HJ. Autosomal dominant Alport syndrome caused by a *COL4A3* splice site mutation. *Kidney Int.* 2000;58(5):1870-5.
- Van Eyken E, Van Camp G & Van Laer L. The complexity of age-related hearing impairment: contributing environmental and genetic factors. *Audiol Neurootol.* 2007b;12(6):345-58.

- Van Eyken E, Van Camp G, Fransen E, Topsakal V, Hendrickx JJ, Demeester K, Van de Heyning P, Mäki-Torkko E, Hannula S, Sorri M, Jensen M, Parving A, Bille M, Baur M, Pfister M, Bonaconsa A, Mazzoli M, Orzan E, Espeso A, Stephens D, Verbruggen K, Huyghe J, Dhooge I, Huygen P, Kremer H, Cremers CW, Kunst S, Manninen M, Pyykkö I, Lacava A, Steffens M, Wienker TF & Van Laer L. Contribution of the N-acetyltransferase 2 polymorphism NAT2*6A to age-related hearing impairment. *J Med Genet.* 2007a;44(9):570-8.
- Van Eyken E, Van Laer L, Fransen E, Topsakal V, Lemkens N, Laureys W, Nelissen N, Vandeveld A, Wienker T, Van De Heyning P & Van Camp G. *KCNQ4*: a gene for age-related hearing impairment? *Hum Mutat.* 2006;27(10):1007-16
- Van Laer L, Huizing EH, Verstreken M, van Zuijlen D, Wauters JG, Bossuyt PJ, Van de Heyning P, McGuirt WT, Smith RJ, Willems PJ, Legan PK, Richardson GP & Van Camp G. Nonsyndromic hearing impairment is associated with a mutation in *DFNA5*. *Nat Genet.* 1998;20(2):194-7.
- Van Laer L, Huyghe JR, Hannula S, Van Eyken E, Stephan DA, Mäki-Torkko E, Aikio P, Fransen E, Lysholm-Bernacchi A, Sorri M, Huentelman MJ & Van Camp G. A genome-wide association study for age-related hearing impairment in the Saami. *Eur J Hum Genet.* 2010;18(6):685-93.
- Van Laer L, Van Eyken E, Fransen E, Huyghe JR, Topsakal V, Hendrickx JJ, Hannula S, Mäki-Torkko E, Jensen M, Demeester K, Baur M, Bonaconsa A, Mazzoli M, Espeso A, Verbruggen K, Huyghe J, Huygen P, Kunst S, Manninen M, Konings A, Diaz-Lacava AN, Steffens M, Wienker TF, Pyykkö I, Cremers CW, Kremer H, Dhooge I, Stephens D, Orzan E, Pfister M, Bille M, Parving A, Sorri M, Van de Heyning PH & Van Camp G. The grainyhead like 2 gene (*GRHL2*), alias *TFCP2L3*, is associated with age-related hearing impairment. *Hum Mol Genet.* 2008;17(2):159-69.
- Varga R, Avenarius MR, Kelley PM, Keats BJ, Berlin CI, Hood LJ, Morlet TG, et al. *OTOF* mutations revealed by genetic analysis of hearing loss families including a potential temperature sensitive auditory neuropathy allele. *J Med Genet.* 2006; 43(7):576-81.
- Verhoeven K, Van Laer L, Kirschhofer K, Legan PK, Hughes DC, Schatteman I, Verstreken M, Van Hauwe P, Coucke P, Chen A, Smith RJ, Somers T, Offeciers FE, Van de Heyning P, Richardson GP, Wachtler F, Kimberling WJ, Willems PJ, Govaerts PJ & Van Camp G. Mutations in the human alpha-tectorin gene cause autosomal dominant non-syndromic hearing impairment. *Nat Genet.* 1998;19(1):60-2.
- Verpy E, Masmoudi S, Zwaenepoel I, Leibovici M, Hutchin TP, Del Castillo I, Nouaille S, Blanchard S, Lainé S, Popot JL, Moreno F, Mueller RF & Petit C. Mutations in a new gene encoding a protein of the hair bundle cause non-syndromic deafness at the *DFNB16* locus. *Nat Genet.* 2001;29(3):345-9.
- Veske A, Oehlmann R, Younus F, Mohyuddin A, Müller-Myhsok B, Mehdi SQ & Gal A. Autosomal recessive non-syndromic deafness locus (*DFNB8*) maps on chromosome 21q22 in a large consanguineous kindred from Pakistan. *Hum Mol Genet.* 1996;5(1):165-8.
- Vikkula M, Mariman EC, Lui VC, Zhidkova NI, Tiller GE, Goldring MB, van Beersum SE, de Waal Malefijt MC, van den Hoogen FH, Ropers HH, Mayne R, Cheah KE, Olsen BR, Warman ML & Brunner HG. Autosomal dominant and recessive osteochondrodysplasias associated with the *COL11A2* locus. *Cell* 1995;80(3):431-7.

- Wagatsuma M, Kitoh R, Suzuki H, Fukuoka H, Takumi Y, Usami S. Distribution and frequencies of *CDH23* mutations in Japanese patients with non-syndromic hearing loss. *Clin Genet*. 2007;72(4):339-44.
- Wallis C, Ballo R, Wallis G, Beighton P & Goldblatt J. X-linked mixed deafness with stapes fixation in a Mauritian kindred: linkage to Xq probe pDP34. *Genomics*. 1988;3(4):299-301.
- Walsh T, Pierce SB, Lenz DR, Brownstein Z, Dagan-Rosenfeld O, Shahin H, Roeb W, McCarthy S, Nord AS, Gordon CR, Ben-Neriah Z, Sebat J, Kanaan M, Lee MK, Frydman M, King MC & Avraham KB. Genomic duplication and overexpression of *TJP2/ZO-2* leads to altered expression of apoptosis genes in progressive nonsyndromic hearing loss DFNA51. *Am J Hum Genet*. 2010;87(1):101-9.
- Walsh T, Walsh V, Vreugde S, Hertzano R, Shahin H, Haika S, Lee MK, Kanaan M, King MC & Avraham KB. From flies' eyes to our ears: mutations in a human class III myosin cause progressive nonsyndromic hearing loss DFNB30. *Proc Natl Acad Sci U S A*. 2002;99(11):7518-23.
- Wang A, Liang Y, Fridell RA, Probst FJ, Wilcox ER, Touchman JW, Morton CC, Morell RJ, Noben-Trauth K, Camper SA & Friedman TB. Association of unconventional myosin *MYO15* mutations with human nonsyndromic deafness DFNB3. *Science*. 1998; 280 (5368):1447-51.
- Wang QJ, Lu CY, Li N, Rao SQ, Shi YB, Han DY, Li X, Cao JY, Yu LM, Li QZ, Guan MX, Yang WY & Shen Y. Y-linked inheritance of non-syndromic hearing impairment in a large Chinese family. *J Med Genet*. 2004;41(6):e80.
- Wattenhofer M, Sahin-Calapoglu N, Andreasen D, Kalay E, Caylan R, Braillard B, Fowler-Jaeger N, Reymond A, Rossier BC, Karaguzel A & Antonarakis SE. A novel *TMPRSS3* missense mutation in a DFNB8/10 family prevents proteolytic activation of the protein. *Hum Genet*. 2005;117(6):528-35.
- Wayne S, Robertson NG, DeClau F, Chen N, Verhoeven K, Prasad S, Tranebjärg L, Morton CC, Ryan AF, Van Camp G & Smith RJ. Mutations in the transcriptional activator *EYA4* cause late-onset deafness at the DFNA10 locus. *Hum Mol Genet*. 2001;10(3):195-200.
- Weil D, Blanchard S, Kaplan J, Guilford P, Gibson F, Walsh J, Mburu P, Varela A, Levilliers J, Weston MD, Kelley PM, Kimberling WJ, Wagenaar M, Levi-Acobas F, Larget-Piet D, Munnich A, Steel KP, Brown SM & Petit C. Defective myosin VIIA gene responsible for Usher syndrome type 1B. *Nature*. 1995;374(6517):60-1.
- Wester DC, Atkin CL, Gregory MC. Alport syndrome: clinical update. *J Am Acad Audiol*. 1995;6(1):73-9.
- Weston MD, Lujendijk MW, Humphrey KD, Möller C, Kimberling WJ. Mutations in the *VLGR1* gene implicate G-protein signaling in the pathogenesis of Usher syndrome type II. *Am J Hum Genet*. 2004;74(2):357-66.
- Wilch E, Azaiez H, Fisher RA, Elfenbein J, Murgia A, Birkenhäger R, Bolz H, Da Silva-Costa SM, Del Castillo I, Haaf T, Hoefsloot L, Kremer H, Kubisch C, Le Marechal C, Pandya A, Sartorato EL, Schneider E, Van Camp G, Wuyts W, Smith RJ, Friderici KH. A novel DFNB1 deletion allele supports the existence of a distant cis-regulatory region that controls *GJB2* and *GJB6* expression. *Clin Genet*. 2010;78(3):267-74.

- Wilcox ER, Burton QL, Naz S, Riazuddin S, Smith TN, Ploplis B, Belyantseva I, Ben-Yosef T, Liburd NA, Morell RJ, Kachar B, Wu DK, Griffith AJ, Riazuddin S & Friedman TB. Mutations in the gene encoding tight junction claudin-14 cause autosomal recessive deafness DFNB29. *Cell*. 2001;104(1):165-72.
- Willems PJ. Gene localization and isolation in nonsyndromic hearing loss. Willems PJ. In: *Genetic Hearing Loss*. Marcel Dekker Inc. New York, 2004;203.
- Xia JH, Liu CY, Tang BS, Pan Q, Huang L, Dai HP, Zhang BR, Xie W, Hu DX, Zheng D, Shi XL, Wang DA, Xia K, Yu KP, Liao XD, Feng Y, Yang YF, Xiao JY, Xie DH & Huang JZ. Mutations in the gene encoding gap junction protein beta-3 associated with autosomal dominant hearing impairment. *Nat Genet*. 1998;20(4):370-3.
- Xiao S, Yu C, Chou X, Yuan W, Wang Y, Bu L, Fu G, Qian M, Yang J, Shi Y, Hu L, Han B, Wang Z, Huang W, Liu J, Chen Z, Zhao G & Kong X. Dentinogenesis imperfecta 1 with or without progressive hearing loss is associated with distinct mutations in *DSPP*. *Nat Genet*. 2001;27(2):201-4.
- Yan D & Liu XZ. Genetics and pathological mechanisms of Usher syndrome. *J Hum Genet*. 2010;55(6):327-35.
- Yasunaga S, Grati M, Cohen-Salmon M, El-Amraoui A, Mustapha M, Salem N, El-Zir E, Loiselet J & Petit C. A mutation in *OTOE*, encoding *OTOE*ferlin, a FER-1-like protein, causes DFNB9, a nonsyndromic form of deafness. *Nat Genet*. 1999;21(4):363-9.
- Yoon JS, Park HJ, Yoo SY, Namkung W, Jo MJ, Koo SK, Park HY, Lee WS, Kim KH & Lee MG. Heterogeneity in the processing defect of *SLC26A4* mutants. *J Med Genet*. 2008;45(7):411-9.
- Zelante L, Gasparini P, Estivill X, Melchionda S, D'Agruma L, Govea N, Milá M, Monica MD, Lutfi J, Shohat M, Mansfield E, Delgrosso K, Rappaport E, Surrey S & Fortina P. Connexin 26 mutations associated with the most common form of nonsyndromic neurosensory autosomal recessive deafness (DFNB1) in Mediterraneans. *Hum Mol Genet* 1997;6:1605-1609.
- Zeviani M, Moraes CT, DiMauro S, Nakase H, Bonilla E, Schon EA, Rowland LP. Deletions of mitochondrial DNA in Kearns-Sayre syndrome. *Neurology*. 1998;51(6):1525-33.
- Zhao L, Young WY, Li R, Wang Q, Qian Y, Guan MX. Clinical evaluation and sequence analysis of the complete mitochondrial genome of three Chinese patients with hearing impairment associated with the 12S rRNA T1095C mutation. *Biochem Biophys Res Commun*. 2004;325(4):1503-8.
- Zhu M, Yang T, Wei S, DeWan AT, Morell RJ, Elfenbein JL, Fisher RA, Leal SM, Smith RJ & Friderici KH. Mutations in the gamma-actin gene (*ACTG1*) are associated with dominant progressive deafness (DFNA20/26). *Am J Hum Genet*. 2003;73(5):1082-91.
- Zwaenepoel I, Mustapha M, Leibovici M, Verpy E, Goodyear R, Liu XZ, Nouaille S, Nance WE, Kanaan M, Avraham KB, Tekaiia F, Loiselet J, Lathrop M, Richardson G & Petit C. *OTOAnchorin*, an inner ear protein restricted to the interface between the apical surface of sensory epithelia and their overlying acellular gels, is defective in autosomal recessive deafness DFNB22. *Proc Natl Acad Sci U S A*. 2002;99(9):6240-5.



Hearing Loss

Edited by Dr. Sadaf Naz

ISBN 978-953-51-0366-0

Hard cover, 406 pages

Publisher InTech

Published online 28, March, 2012

Published in print edition March, 2012

Authored by 17 international researchers and research teams, the book provides up-to-date insights on topics in five different research areas related to normal hearing and deafness. Techniques for assessment of hearing and the appropriateness of the Mongolian gerbil as a model for age-dependent hearing loss in humans are presented. Parental attitudes to childhood deafness and role of early intervention for better treatment of hearing loss are also discussed. Comprehensive details are provided on the role of different environmental insults including injuries in causing deafness. Additionally, many genes involved in hearing loss are reviewed and the genetics of recessively inherited moderate to severe and progressive deafness is covered for the first time. The book also details established and evolving therapies for treatment of deafness.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Nejat Mahdieh, Bahareh Rabbani and Ituro Inoue (2012). Genetics of Hearing Loss, Hearing Loss, Dr. Sadaf Naz (Ed.), ISBN: 978-953-51-0366-0, InTech, Available from: <http://www.intechopen.com/books/hearing-loss/genetics-of-hearing-loss>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen