

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Genetic Diversity and Transferability of Rubus Microsatellite Markers to South American *Rubus* Species

Marta L. Marulanda*, Ana M. López and Marcela Uribe
Plant Biotechnology Laboratory, Universidad Tecnológica de Pereira
Apartado Aéreo, Pereira
Colombia

1. Introduction

Rubus L. (Rosaceae) is grown extensively worldwide, in altitudes ranging from 0 to 4500 m above sea level. Found in six continents, this genus is reported to contain approximately 800 species due to biological processes such as hybridization, apomixis, and polyploidy that weaken species boundaries (Thompson, 1995). *Rubus* has been divided into 12 subgenera of which only a few species have been domesticated (Table 1).

Subgenus	Number of species
<i>Anoplobatus</i> (Focke) Focke	6
<i>Chamaebatus</i> (Focke) Focke	5
<i>Chamaemoras</i> (Hill) Focke	1
<i>Comaropsis</i> (Rich.) Focke	2
<i>Cylactis</i> (Raf.)Focke	14 (4 series)
<i>Dalibarda</i> (L.) Focke	5
<i>Dalibardastrum</i> Focke	4
<i>Idaeobatus</i> (Focke) Focke	117 (9 sections)
<i>Lampobatus</i> Focke	10
<i>Malachobatus</i> (Focke) Focke	115 (7 sections)
<i>Orobatus</i> (Focke)	19
<i>Rubus</i> L. (= <i>Eubatus</i> Focke)	132 (6 sections)

Table 1. Subgenera and number of species of *Rubus* (Focke, 1910, 1911, 1914).

The subgenus *Idaeobatus* contains the “raspberries” that are distributed in the Northern Hemisphere, mainly Asia, Africa, Europe, and North America. The subgenus *Rubus* includes species found in Europe, Asia, and North America, whereas the subgenus *Orobatus* is

* Corresponding Author

exclusive to South America. Species representative of the subgenera *Rubus*, *Orobatus*, and *Idaeobatus* are found in the Colombian and Ecuadorian Andes (Ballington et al., 1993). Vargas (2002) reported six species in Colombia's central Andes: *R. bogotensis*, *R. glaucus*, *R. macrocarpus*, *R. nubigenus*, *R. porphyromallus*, and *R. urticifolius*. Romoleroux (1992) also reported the existence of nine edible species in Colombia of a total of 44 species reported, and that due to natural crossing, up to 500 varieties can be identified. Besides its ecological relevance, for example as an invasive weed, this genus is an economically important fruit crop for small farmers in the northern Andes, mainly because of the production of *Rubus glaucus* L., commonly known as the Andean blackberry (Fig. 1).




Fig. 1. Plants of *Rubus glaucus* L. growing on the Andes.

Of flowering plant species, the genus *Rubus* is one of the most complexes because of its hybridization, polyploidy, agamospermy, and loss of the concept of species (Alice & Campbell, 1999). Polyploidy and hybridization prevails in the subgenus *Rubus*. Only subgenera *Idaeobatus*, *Dalibarda*, and *Anoplobatus* are predominantly diploid, whereas *Dalibardastrum*, *Malachobatus*, and *Orobatus* are exclusively polyploid (Thompson, 1995, 1997). Hybridization occurs in *Rubus* between closely related species (Kraft et al., 1996; Naruhashi, 1979, 1990; Steele & Hodgdon, 1963, 1970) and, in some cases, between subgenera (Alice et al., 1997; Gustafsson, 1942; Jennings, 1978; Weber, 2003).

This species presents traits of two subgenera – *Idaeobatus* and *Rubus* – possibly being a fertile amphidiploid or allotetraploid ($n=7$, $4x=28$) (Delgado et al., 2010; Thompson, 1997). Sympatrically with *R. glaucus*, some other wild *Rubus* species are found in the Andean cordillera along with the introduced and cultivated Eurasian *R. idaeus* L. Because hybridization is a common process that affects species of any genus (Randell et al., 2004), it is reasonable to believe that gene flow is currently taking place between species of this genus.

Studies on the genetic diversity of *Rubus* have been carried out in temperate species, such as *Rubus idaeus* (Graham & McNicol, 1995; Graham et al., 1997; Parent & Fortin, 1993) and *Rubus occidentalis* (Parent & Page, 1998), and Asian species (Amsellem et al., 2000). These

works used RAPD, RFLP, and SCAR markers as well as SSR (Antonius-Klemola, 1999). ITS are also been used to study hybrids of *R. idaeus* and *R. caesius* (Alice et al., 1997). These markers made it possible to confirm the genetic origin of the hybrids and further phylogenetic and evolutionary studies in *Rubus* (Alice, 2002). Recently, major advances have been achieved worldwide in the use of molecular markers in temperate species of *Rubus*, such as DNA fingerprinting to study and characterize genotypes, development of linkage maps, marker-assisted selection, and mapping of QTLs (Antonius-Klemola, 1999; Graham et al., 2002).

To date, molecular markers such as randomly amplified polymorphic DNA (RAPD), amplified fragment length polymorphisms (AFLP) (Marulanda et al., 2007), and microsatellites (simple sequence repeats, SSRs) (Marulanda et al., 2011) have been used to study the genetic diversity of the Andean blackberry. Previous work carried out by Marulanda et al. (2007) and Marulanda & López (2009) on the genetic diversity of Colombian blackberries identified high phenotypic and molecular plasticity in the *R. glaucus* species known as the “Castilla” blackberry in Colombia’s central Andean area. Other wild *Rubus* species present in the Andean region are found near farms where the “Castilla” blackberry is commercially grown. These plants were also submitted to morphological, agronomic, and molecular characterizations using AFLP and SSR molecular markers (Marulanda & López, 2009). A genomic library enriched for microsatellite sequences was recently developed for *R. glaucus*.

This chapter presents the results of the molecular characterization of wild and cultivated *Rubus* species collected in the central Andes of Colombia using SSR markers from other *Rubus* species available in Genbank, together with 11 microsatellite markers isolated from *R. glaucus* and characterized in 39 samples of *Rubus* (Table 2). Intra- and inter-specific differences between *R. glaucus* and its wild relatives were established, generating not only information on the current status of populations, their uses, and distribution, but also information considered crucial to launch a breeding program for *R. glaucus*.

2. Materials and methods

2.1 Plant material and DNA extraction

A total of 39 *Rubus* samples were collected at altitudes ranging from 1800 to 2455 m above sea level in the central Andes of Colombia (between 1° 42' 10.7" and 6° 99' 44" N and 72° 92' 80" and 76° 25' 35.9" W), and placed on silica gel (1:10, plant tissue: silica gel) (Table 2). DNA was isolated using the Plant DNAeasy Miniprep kit (QIAGEN®), following the manufacturer’s instructions. Several samples did not show any DNA after the isolation procedure so it was necessary to reprocess these samples following the Doyle & Doyle (1990) protocol. In all cases, samples were purified using the protocol described by Castillo (2006).

2.2 Analysis with SSR markers

A total of 36 microsatellite sequences from other *Rubus* species, *R. idaeus* (23 primer pairs) (Series RhM, RiM and Rubus) and *R. occidentalis* (2 primer pairs) (Series mRaCIRRI), and 11 microsatellites from *R. glaucus* were used (Table 3). The microsatellite named as “Rg” was

developed using a genomic library enriched for microsatellite sequences from a cultivated genotype of *R. glaucus*, following the protocol described by Billotte et al. (1999).

Code	Latitude (N)	Longitude (W)	Cultivated/Wild	Altitude (m.a.s.l.)	Characteristics
CVM 1	4°48'06''	74°24'49''	<i>R. glaucus</i> Cultivated	2455	Thorny
CVM 2	4°47'37''	74°25'27''	<i>R. glaucus</i> Cultivated	2337	Thorny
CVM 3	4°79'33''	74°42'68''	<i>R. glaucus</i> Cultivated	2288	Thorny
CVM 4	4°09'09''	74°23'27''	<i>R. glaucus</i> Cultivated	2007	Thorny
CVM 6	6°99'44''	72°98'80''	<i>R. glaucus</i> Cultivated	2157	Thorny
CVM 7	6°99'44''	72°98'80''	<i>R. glaucus</i> Cultivated	2157	Thornless
CVM 8	6°59'39.1''	72°59'13''	<i>R. glaucus</i> Cultivated	2176	Thorny
CVM 9	7°00'38''	72°58'39''	<i>R. glaucus</i> Cultivated	2357	Thorny
CVM 10	4°40'00.8''	76°16'06.6''	<i>R. glaucus</i> Cultivated	2049	Thorny
CVM 11	4°13'23.8''	76°25'35.9''	<i>R. idaeus</i> Raspberry	2100	Thornless
CVM 12	4°13'23.8''	76°25'35.9''	<i>R. glaucus</i> Cultivated	2380	Thorny
CVM 13	4°13'23.8''	76°25'35.9''	<i>R. glaucus</i> Cultivated	2380	Thorny
CVM 15	6°14'52.3''	75°24'08.5''	<i>R. glaucus</i> Cultivated	2150	Thorny
CVM 18	6°09'15.4''	75°23'00.1''	<i>R. glaucus</i> Cultivated	2000	Thornless
CVM 19	6°09'15.4''	75°23'00.1''	<i>R. glaucus</i> Cultivated	2000	Thorny
CVM 20	6°09'15.4''	75°23'00.1''	<i>R. glaucus</i> Cultivated	2000	Thorny
CVM 22	1°57'45.2''	76°14'17.4''	<i>R. glaucus</i> Cultivated	1840	Thorny
CVM 25	1°51'41.8''	76°21'21.9''	<i>R. glaucus</i> Cultivated	2140	Thorny
CVM 26	1°53'42.8''	76°18'45.5''	<i>R. glaucus</i> Cultivated	1850	Thorny
CVM 27	1°42'10.7''	76°12'08.3''	<i>R. glaucus</i> Cultivated	2000	Thorny
CVM WILD	_____	_____	<i>R. glaucus</i> Wild	_____	Wild
CVM A	4° 44'45.1''	75° 36'39.6''	<i>R. glaucus</i> Cultivated	2000	Thornless
CVM B	4° 39'7''	75° 35'26.3''	<i>R. glaucus</i> Cultivated	2014	Thorny
CVM C	4°38'36''	75°28'41.5''	<i>R. glaucus</i> Cultivated	2300	Thornless
CVM D	4°48'99.2''	75°41'86''	<i>R. glaucus</i> Cultivated	1950	Thorny
CVM E	5° 2'2.7''	75° 27'10.5''	<i>R. glaucus</i> Cultivated	1800	Thorny
95	4°21'22.5''	75°42'10.1''	<i>R. glaucus</i> Wild	1805	Wild
107	4° 33'51.9''	75° 39'14.7''	<i>R. urticifolius</i>	1800	Wild
106	4°39'7''	75° 35'26.3''	<i>R. urticifolius</i>	1574	Wild
97	4° 41'23.3''	75°37'33.8''	<i>R. glaucus</i> Wild	1997	Wild
86	4° 38'36''	75° 29'9.5''	<i>R. glaucus</i> Wild	2430	Wild
22	5° 4' 35.8''	75° 32' 31.1''	<i>R. glaucus</i> Cultivated	1947	Thorny
37	5° 0'50.3''	75° 32'11.7''	<i>R. urticifolius</i>	1777	Wild
44	4° 45' 00.2''	75° 36' 39''	<i>R. urticifolius</i>	1879	Wild
MSA 1	4°52'15.0''	75°37'32.4''	<i>R. glaucus</i> Cultivated		Thornless
MSA 2	4° 44'45.1''	75° 36'39.6''	<i>R. glaucus</i> Cultivated	1850	Thornless
MSA 3	4°38'36''	75°28'41.5''	<i>R. glaucus</i> Cultivated		Thornless
MSA 4	4°11'36.1''	75°48'14.6''	<i>R. glaucus</i> Cultivated	2000	Thornless
MSA 5	5°0'50.3''	75°32'11.7''	<i>R. glaucus</i> Cultivated	1825	Thornless

Table 2. Samples of *Rubus* species, accessions of *R. glaucus*, and collection sites.

Amplification reactions were performed in a final volume of 12.5 µl, with 5 ng genomic DNA, 0.3 µM of each primer, 1X reaction buffer (10 mM Tris-HCl, pH 8.3, 50 mM KCl), 200 µM of each dNTP, 2 mM MgCl₂ and 1.0 unit of Taq DNA polymerase. The PCR was performed according to the following parameters: 94 °C for 4 min, 10 cycles of 94 °C for 30 sec, 65 °C (–1 °C/cycle) for 30 sec and 72 °C for 1 min; 35 cycles of 94 °C for 15 sec, annealing temperature (°C) for 30 sec and 72 °C for 1 min; and 72 °C for 5 min.

2.3 Statistical analysis

The allelic diversity of the SSR was evaluated by determining polymorphism information content (PIC) value, as described by Bonstein et al. (1980) and cited and modified by Anderson et al. (1993), as described in Equation 1:

$${}_n\text{PIC}_i = 1 - \sum_{j=1} P_{ij}^2 \quad (1)$$

Where P_{ij} is the frequency of the j th pattern, i is the sum, and n are the patterns. To measure the utility of the marker systems, average heterozygosity and expected and observed heterozygosity were calculated. The partitioning of genetic variation within and among the groups by the SSR marker system was achieved by analysis of molecular variance (AMOVA) using the same software. Analyses were performed in GenAlex (Peakall & Smouse, 2006) and Arlequin v.3.5 (Excoffier & Lischer, 2010).

3. Results and discussion

3.1 Genetic diversity and variability

A total of 41 loci and 133 alleles were detected. The number of alleles observed for each locus ranged from 2 to 6, with an average of 4.6 alleles per locus. The PIC value varied between 0.07 and 0.61 (average 0.48), and the discriminating power (D) ranged from 0.05 to 0.52 (average 0.27). Observed heterozygosities (H_o) were 0.078–1.0 (average 0.84) and expected heterozygosities (H_e) were 0.07–0.582 (average 0.473). The highest PIC value (0.61) was found in the Rg-D7 locus, which presented a high number of alleles (5). To compare the efficiency of markers to identify varieties, the D value of each primer was estimated. The highest D value (0.52) was also found in the Rg-D7 locus (Table 3).

Similar results were reported by Castillo (2006), who used 12 SSRs to analyze an extensive collection of North American *Rubus* (raspberry) germplasm. Results indicated from 3 to 16 alleles per locus, with an average of eight alleles per locus and a total number of alleles of 96. More recently, Flores et al. (2010) isolated 12 microsatellites from SSR-enriched genomic libraries of *R. idaeus*.

Another measure of genetic variability is the presence of exclusive alleles per loci and genotype (Tables 4 and 5). The genotypes presenting the highest number of exclusive alleles are listed in Table 4, with *R. idaeus* genotype CVM11 ranking highest, which could be attributed to the fact that most of the SSRs used are derived from *R. idaeus*. Of the evaluated microsatellites, those of the series “Rubus” were one of the most polymorphic groups and detected the highest number of alleles in the study population. It should be mentioned that exclusive alleles also appear in wild genotypes and in genotypes 106 and 107, which belong to the species *R. urticifolius*. The loci in which the private alleles were detected are very important for genotype differentiation, particularly in the case of the thornless genotypes.

Fig. 2 presents the results of the principal coordinates analysis. There is no clear differentiation of genotypes based on collection sites; however, the genotypes belonging to the species *R. urticifolius* (37, 44, 106, and 107) are separate from the species *R. glaucus*, both cultivated and wild. Likewise, the species *R. idaeus* is separate from the species *R. urticifolius*

	Locus	Motif	GenBank accession no.	Primer sequences (5'-3')	Size (bp)	T (°C)	Number of loci	PIC*	A*	D*	He*	Ho*
1	RhM018 ³	(CTT) ₆	FJ194447	F-ACCAATTGTACACCCAACAAC R-GATTGTGAGCTGGTTACCAA	379	52	1	0.3510	2	0.5094	0.279	0.435
2	RhM023 ³	(CAT) ₅	FJ194449	F-CGACAAACGACAAATCTCACATT R-GTTATCAAGCGATCTCGAGTT	196	52	1	0.4864	2	0.3388	0.463	0.835
3	RhM011 ³	(TC) ₁₈	FJ194446	F-AAAGACAAGGCTCCACAAC R-GTTATGCTTTGATTAGGCTGG	280	58	2	0.5419	3	0.1448	0.530	0.969
4	RhM001 ³	(CA) ₇	FJ194444	F-GGTTGGGATAGTTAATCTCCC R-CCAACTGTTGTAATGCAGGAA	232	50	1	0.1328	3	0.2148	0.117	0.078
5	RhM021 ³	(TC) ₆	FJ194448	F-CAGTCCCTTATAGGATCCAACG R-GAACTCCACCATCTCCTCTAG	282	50	1	0.5461	3	0.219	0.536	1.000
6	RhM003 ³	(TG) ₁₀	FJ194445	F-CCATCTCCAATTCAGTCTTCC R-AGCAGATCGGTTCTACAAGC	200	48	1	0.5151	3	0.047	0.512	1.000
7	RiM017 ³	(TC) ₆	FJ194453	F-GAAAACAGGTGGAAGAAACCTTG R-CATTGTGCTTATGATGTTTCG	194	58	1	0.5869	3	0.2148	0.556	0.897
8	RiM015 ³	(ATC) ₅	FJ194452	F-CGACACCGATCAGAGCTAATTC R-ATAGTGCATTGGCAGGCTTAT	350	58	1	0.4462	2	0.4451	0.367	0.610
9	Rubus 76b ²	(CT) ₅ (CT) ₄	—	F-CTCACCCGAAATGTCAACC R-GGCTAGGCCGAATGACTACA	190-210	55	1	0.6004	4	0.2463	0.567	0.897
10	Rubus 16a ²	(AT) ₈ (GT) ₁₁	—	F-TGTTGACGTGTGGGCTTT R-GGGTGTGTGCCAGTTTCAGT	169	55	3	0.3461	3	0.0588	0.533	1.000
11	Rubus 116a ²	(CT) ₁₂ (T) ₁₀	—	F-CCAAACCCAAAAACCTTCAAC R-GTTGTGGCATGGCCTTTTAT	299	55	2	0.5297	3	0.053	0.520	0.969
12	Rubus 105b ²	(AG) ₈	—	F-GAAAATGCAAGGCGAATTGT R-TCCATCACCAACACCACCTA	165 -173	55	2	0.5924	6	0.2171	0.571	0.948
13	Rubus 137a ²	(TG) ₈ (TA) ₄	—	F-TGTGACGAGATGAAGGAGCTA R-AGCAATTATTCGCGCAGTTT	198	55	1	0.4995	2	0.0465	0.496	0.969
14	Rubus 259f ²	(CT) ₄ (AG) ₈	—	F-TGGCAACAAGAAGCCTGTAAC R-TCCCATATCCCTCAGCATTC	265	55	1	0.5717	4	0.1727	0.552	1.000
15	Rubus 98d ²	(GAA) ₅ (GA) ₁₀	—	F-GGCTTCTCAATTGTCTGTGC R-TGATTTGAATCGTCCGGTTA	198	55	2	0.5880	4	0.2558	0.544	0.861
16	mRaCIRRI1G3 ¹	(GA) ₂₈	AF205116	F-CTCTACAAAAGGATCTGCATGA R-CAGCAAAAAGTGAATGGTTCA	195-265	55	2	0.5473	3	0.2566	0.524	0.881
17	mRaCIRRIV2A8 ¹	(CA) ₁₂ (CT) ₁₁	AF261693	F-TAAAAAGGCGCAACAGTCG R-AGACACAGAAACAGGCATCG	191-237	55	2	0.5604	3	0.23	0.548	0.959
18	Rubus 285a ²	(TC) ₉	—	F-TCCAGAAAGCTTGCTATGCTG R-GGATACCTCAATGGCTTCTTG	138-167	55	1	0.5497	4	0.2462	0.543	1.000

19	Rubus 145a ²	(GT) ₇	—	F-TGTCCCAGCTTTCTGGTTTC R-GGCATCTGTGCGGTAAAAAT	131	55	1
20	Rubus 262b ²	(AG) ₁₅	—	F-TGCATGAAGGCGATATAAAGG R-TCCGCAAGGGTTGTATCCTA	217-225	55	1
21	Rubus 107a ²	(AG) ₈	—	F-GCCAGCACCAAAAACCTACA R-TTTCACCGTCAAGAAGAAAGC	179	55	1
22	Rubus 251a ²	(GA) ₁₀	—	F-GCATCAGCCATTGAATTTC R-CCCACCTCCATTACCAACTC	157-283	55	1
23	Rubus 270a ²	(GA) ₁₀	—	F-GCATCAGCCATTGAATTTC R-CCCACCTCCATTACCAACTC	167-185	55	1
24	Rubus 2a ²	(GT) ₁₂ -G-(GT) ₈	—	F-TGAGGGAAGAAGAGGCAAGA R-CACGTGTGACCCCAATGATA	175-180	55	1
25	Rubus 194h ²	(GA) ₁₂	—	F-TGTGTTGTTCTCTGCAACCA R-AGCCCTTACTTTTCCTGCAA	100-110	55	1
26	Rg-A6 ⁴	(GAA) ₄	—	F-AGCGCAAGGACTTCTACC R-TTCAGCTCCGGTAGTAGC	282-293	55	1
27	Rg-A12-1 ⁴	(GA) ₂₃	HQ637525	F-TCATGTTTCATGTTTCGTGTG R-CTAACACGCGATAGAATAGG	387-432	48	1
28	Rg-A12-2 ⁴	(AC) ₈	HQ637525	F-GCGGGCATTCTCTTGCTTAC R-GCGGTTTCGTGACTCAGACAG	179-194	58	1
29	Rg-B7 ⁴	(CA) ₁₁	HQ637519	F-TACGCGTGGCACTAACAC R-TCAGAGCACTCCAAGAAGG	150-171	58	2
30	Rg-B8 ⁴	(AG) ₂₃	HQ637493	F-CTGATGAGTAGCTGGTTTCTC R-GTCTGAAATAGCACATGGAC	185-230	52	1
31	Rg-B9 ⁴	(CA) ₉	HQ637500	F-TTCAGAACCTCATGCAGAGCC R-TTCACTCAGGAGGAGGAGGTG	348-367	57	1
32	Rg-D2 ⁴	(GA) ₁₆	HQ637534	F-AAACCCTAATTGTGTTGAC R-CCCAAGCTAGATTTCCTC	418-449	48	1
33	Rg-D7 ⁴	(GA) ₂₅	HQ637494	F-AACCATCGGTGTCGACCTC R-GGCTCTTGACCGTAACTTAG	61-110	55	1
34	Rg-F3 ⁴	(GA) ₁₇ (GT) ₉	HQ637516	F-GCCAATGAAACGGAAAGAC R-GCCTTCACTCATATCATTCTCC	210-265	60	1
35	Rg-F7 ⁴	(CT) ₂₁	HQ637527	F-GCAGGTGTGTCTATGTGATATG R-ATTCCCAACGACTACATAATTC	290-331	55	1
36	Rg-F8 ⁴	(GA) ₁₄	HQ637499	F-CATGCTGGCTTTGCTGCTC R-TGCTTACGCGTGACTAACAG	267-294	60	1

¹ *R. alceifolius* (Amsellem et al., 2001); ² *R. idaeus* (Graham et al., 2002, 2004) ; ³ *R. idaeus* and blackberry “Marion” (Marulanda et al., 2010 (data not published).
* PIC = polymorphic information content; A = allele number; D = discrimination power; He = expected heterozygosity.

Table 3. Microsatellite sequences and characteristics of each SSR used to evaluate *Rubus* materials.

and from cultivated and wild genotypes of *R. glaucus*. Genotype CVM Wild of the species *R. glaucus* is notably separate from the genotypes of the same species.

Locus	Private alleles	Frequency
RhM003	2	0.125
RhM001	2	0.250
Rubus 76b	1	0.125
Rubus 98d	1	0.333
Rubus 98d	1	0.333
mRaCIRRI1G3	2	0.125
mRaCIRRIV2A8	5	0.125
Rubus 128a	3	0.125
Rubus 262b	3	0.125
Rubus 107a	5	0.250
RgA12-1	4	0.125
RgB7	2	0.125
RgB7	1	0.125
RgB7	2	0.125
RgF3	1	0.250
RgF7	1	0.250
Rubus 116a	1	0.250
Rubus 105b	2	0.100
Rubus 105b	5	0.100
Rubus 105b	4	0.200
RgA12-1	2	0.100
Rubus 98d	3	0.222

Table 4. Summary of private alleles.

Sample	No. of loci with private alleles	Loci with private alleles
<i>R. glaucus</i> cultivated (CVM 10)	1	RgB7
<i>R. idaeus</i> (CVM 11)	14	RhM003, RhM001, Rubus 76b, Rubus 98d, Rubus 98d, mRaCIRRI1G3, mRaCIRRIV2A8, Rubus 128a, Rubus 262b, Rubus 107a, RgA12-1, Rg B7, RgF3, RgF7
<i>R. glaucus</i> cultivated (CVM 13)	1	RgB7
<i>R. glaucus</i> wild (CVM WILD)	3	Rubus 116a, Rubus 105b, Rubus 105b
<i>R. glaucus</i> cultivated (22)	1	RgA12-1
<i>R. urticifolius</i> (107)	1	Rubus 98d
<i>R. urticifolius</i> (106)	1	Rubus 98d

Table 5. List of samples with one or more private alleles.

3.2 Transferability of microsatellite sequences

Microsatellites RhM and RiM developed from *R. idaeus* showed cross-species amplification in *R. glaucus* genotypes and in other two wild species of *Rubus*, with 2 and 3 alleles per locus. High D values were found for loci RhM018 (0.5094) and RiM015 (0.4451). The “*Rubus*” microsatellite markers, also from *R. idaeus*, were the most polymorphic and presented the highest number of alleles per locus: *Rubus* 105b with 6 alleles and *Rubus* 107a with 5. The microsatellite markers from *R. alceifolius* showed 3 alleles and D values ranging from 0.23 to 0.2566. The *R. glaucus* microsatellites (Rg series) amplified for other wild species, with the number of alleles ranging from 2 to 5 per locus.

The cross-species amplification data suggested that microsatellites developed for *R. idaeus*, *R. alceifolius*, and *R. glaucus* can be potentially useful for genetic diversity studies of different *Rubus* species. In the case of conservation programs, they should prove useful for characterizing natural populations and germplasm collections, as well as for determining the degree of relatedness between individuals or groups of accessions.

The microsatellites developed by Amsellem et al. (2001) to study *R. alceifolius*, subgenus *Malachobatus*, that grows in Southeast Asia were used for the characterization of Andean *Rubus*, the transferability and applicability of microsatellites of *R. alceifolius* to study and evaluate the diversity of *Rubus* species in the American Andes were demonstrated; results were similar to those obtained in Asian species.

Amsellem et al. (2001) also observed amplification from 3 to 4 alleles per individual in the species *R. alceifolius*, confirming the suspicion that this is a tetraploid species. Based on the analysis carried out by Amsellem et al. (2001), the present study produced between 3 and 5 alleles for *R. glaucus* and between 2 and 4 alleles for *R. urticifolius*., suggesting that both cultivated and wild materials of *R. glaucus* have ploidy levels greater than those of *R. urticifolius*. This polyploidy was also described by Hall (1990), who explained that *Rubus* species used in plant breeding programs have produced euploid and aneuploid hybrids and that diploid, triploid, tetraploid, hexaploid, septaploid, octaploid, and nonaploid cultivars have been selected, most of them tetraploids.

4. Conclusion

Wild forms are also usually found at sites where *Rubus* species are cultivated, particularly in forest clearings, along roadsides, and on hillsides. Both cultivated and wild forms have the potential for interacting in different ways with cultivated materials. Cultivars can influence the genetic diversity of natural populations through gene transfer by pollen and wild populations are a potential source of genetic material for improvement programs.

This evaluation of the status of genetic resources of the species *R. glaucus* and related wild species serves to provide guidelines for conservation and breeding efforts aiming to promote the development of cultivated species important for the rural economies of South America's Andean region.

Using microsatellites from other *Rubus* species has proven to be a very useful strategy to differentiate between wild and cultivated *R. glaucus* genotypes, as well as between thorny and thornless cultivars.

The development of a genomic library enriched with microsatellites and the design of microsatellite sequences for the Andean specie *Rubus glucus*, is allowing a deeper comprehension of the genetic variability existing among cultivated and wild genotypes as well as the relationships between the cultivated specie and the wild relatives.

The Analysis of molecular variation (AMOVA) showed a higher variability distributed between genotypes than between populations, which agrees with the results obtained in the principal coordinates analysis.

5. Acknowledgment

The authors would like to thank the blackberry farmers of Colombia, the Governor's Office of the Department of Risaralda, and the Colombian Ministry of Agriculture and Rural Development for financing and supporting the present research. Our sincere thanks also go to Juliana Arias Villegas for planning and managing the project, all laboratory and nursery staff of the Biodiversity and Biotechnology Research Group, and the Universidad Tecnológica de Pereira.

6. References

- Alice, L.A. (2002). Evolutionary relationships in *Rubus* (Rosaceae) based on molecular data. *Acta Horticulturae*, Vol.585, pp. 79–83, ISSN 0567-7572
- Alice, L.A. & Campbell, C.S (1999) Phylogeny of *Rubus* (Rosaceae) based on nuclear ribosomal DNA internal transcribed spacer region sequences. *American Journal of Botany*. 86:1:81-97
- Alice, L.A.; Eriksson, T.; Eriksen, B. & Campbell, C.S. (1997). Intersubgeneric hybridization between a diploid raspberry, *Rubus idaeus*, and a tetraploid blackberry, *R. caesius* (Rosaceae). *American Journal of Botany*, Vol.84, pp. 171, ISSN 0002-9122
- Amsellem L., Dutech C., Billotte N. (2001). Isolation and characterization of polymorphic microsatellite loci in *Rubus alceifolius* Poir. (Rosaceae), an invasive weed in La Reunion Island. *Molecular Ecology Notes*, Vol.1, pp. 33–35, ISSN 1471-8278
- Amsellem, L.; Noyer, J.L.; Le Bourgeois, T.; Hossaert-Mckey, M. (2000). Comparison of genetic diversity of the invasive weed *Rubus alceifolius* Poir. (Rosaceae) in its native range and in areas of introduction, using amplified fragment length polymorphism (AFLP) markers. *Molecular Ecology*, Vol.9, pp. 443–455, ISSN 0962-1083
- Anderson, J.A.; Churchill, G.A.; Autrique, J.E.; Tanksley, S.D. & Sorrells, M.E. (1993). Optimizing parental selection for genetic linkage maps. *Genome*, Vol.36, pp. 181–186, ISSN 0831-2796
- Antonius-Klemola. K. (1999). Molecular markers in *Rubus* (Rosaceae) research and breeding. *The Journal of Horticultural Science & Biotechnology*, Vol.74, No.2, pp. 149–160, ISSN 1462-0316
- Ballington, J. R.; Luteyn, M.M.; Thompson, K.; Romoleroux, K. & Castillo, R. (1993). *Rubus* and *Vacciniaceous* germplasm resources in the Andes of Ecuador. *Plant Genetic Resources Newsletter*, Vol.93, pp. 9–15, ISSN 1020-3362
- Billotte, N.; Lagoda, P.J.L.; Risterucci, A.M. & Baurens, F.C. (1999). Microsatellite-enriched libraries: applied methodology for the development of SSR markers in tropical crops. *Fruits*, Vol.54, No.4, pp. 277–288, ISSN 0248-1294


- Bonstein, D.; White, R. L.; Skolnick, M.E. & Davis, R.W. (1980). Construction of a genetic linkage map in man using restriction fragment length polymorphism. *American Journal of Human Genetics*, Vol.32, pp. 314–331, ISSN 0002-9297
- Castillo, N.R. (2006). Fingerprinting and genetic stability of *Rubus* using molecular markers. M.Sc. Thesis. Oregon State University, USA
- Delgado, L.M.; Uribe, M. & Marulanda, M.L. (2010). Estandarización de una técnica citogenética para el conteo de cromosomas mitóticos en *Rubus glaucus* Benth. *Scientia et Technica*, Vol.46, pp. 74–80, ISSN 0122-1701
- Doyle, J. J. & Doyle, J.L. (1990). Isolation of plant DNA from fresh tissue. *Focus*, Vol.12, pp. 13–15
- Excoffier, L. & Lischer, H.E.L. (2010). Arlequin suite ver 3.5: A new series of programs to perform population genetics analyses under Linux and Windows. *Molecular Ecology Resources*, Vol.10, pp. 564–567, ISSN 1755-098X
- Flores, N. R.; Reed, B.M.; Graham, J.; Fernandez-Fernandez, F. & Bassil, N.V. (2010). Microsatellite markers for raspberries and blackberries. *Journal of the American Society for Horticultural Science*, Vol.135, No.3, pp. 271–278, ISSN 0003-1062
- Focke, W.O. (1910-1914) Species *Ruborum*. *Bibliotheca Botanica*, Vol.72, pp. 1-274, ISSN 0067-7892
- Graham, J. & Mcnicol, R.J. (1995). An examination of the ability of RAPD markers to determine the relationships within and between *Rubus* species. *Theor. Appl Genet* 90: 1128-1132.
- Graham, J., Squire, G.R. Marshall, B. & Harrison, R.E. (1997). Spatially dependent genetic diversity within and between colonies of wild raspberry *Rubus idaeus* detected using RAPD markers. *Molecular Ecology* 6(11): 1001 – 1008.
- Graham, J.; Smith, K.; Mackenzie, K.; Jorgenson, L.; Hackett, C.A. & Powell, W. (2004). The construction of a genetic linkage map of red raspberry (*Rubus idaeus* subsp. *idaeus*) based on AFLPs, genomic SSR and EST-SSR markers. *Theoretical and Applied Genetics*, Vol.109, pp. 740–749, ISSN 0040-5752
- Graham, J.; Smith, K.; Woodhead, M. & Russell, J. (2002). Development and use of SSR markers in *Rubus* species. *Molecular Ecology*, Vol.2, pp. 250–252, ISSN 0962-1083
- Graham, J.; Squire, G.R.; Marshall, B.Y. & Harrison, R.E. (1997). Spatially dependent genetic diversity within and between colonies of wild raspberry *Rubus idaeus* detected using RAPD markers. *Molecular Ecology*, Vol.6, No.11, pp. 1001–1008, ISSN 0962-1083
- Gustafsson, A. (1942). The origin and properties of the European blackberry flora. *Hereditas*, Vol.28, pp. 249–277, ISSN 0018-0661
- Hall, H.K. (1990). Blackberry breeding, In: *Plant Breeding Reviews* Vol. 8, J. Janick (Ed.), 249–312, Timber Press, ISBN 9780881921731, Portland, OR
- Jennings, D.L. (1978). The blackberries of South America—an unexplored reservoir of germplasm. *Fruit Varieties Journal*, Vol.32, pp. 61–63, ISSN 0091-3642
- Jennings, D.L. (1988). *Raspberries and Blackberries: Their Breeding, Diseases and Growth*, Academic Press, ISBN 0123842409, New York
- Kollmann, J.; Steinger, T. & Roy, B.A. (2000). Evidence of sexuality in European *Rubus* (Rosaceae) species based on AFLP and allozyme analysis. *American Journal of Botany*, Vol.87, No.11, pp. 1592–1598, ISSN 0002-9122

- Kraft, T. & Nybom, H. (1995) DNA fingerprint and biometry can solve some taxonomic problems in apomictic blackberries (*Rubus* sub *Rubus*) *Watsonia* 20:329-43
- Kraft, T.; Nybom, H. & Werlemark, G. (1996). DNA fingerprint variation in some blackberry species *Rubus* sug. *Rubus*, Rosaceae. *Plant Systematics and Evolution*, Vol.199, pp. 93–108, ISSN 0378-2697
- Marulanda, M.L. & López, A.M. (2009). Characterization of thornless *Rubus glaucus* in Colombia. *Canadian Journal of Pure & Applied Sciences*, Vol.3, No.2, pp. 927–937, ISSN 1715-9997
- Marulanda, M.L.; López, A.M. & Aguilar, S.B. (2007). Genetic diversity of wild and cultivated *Rubus* species in Colombia using AFLP and SSR markers. *Crop Breeding and Applied Biotechnology*, Vol.7, pp. 243–253, ISSN 1518-7853
- Marulanda M.L.; López, A.M. & Uribe, M. (2011). Molecular characterization of *Rubus* spp using SSR markers. *Genetic Molecular Research (GMR) Brasil*. (In press)
- Michalakis, Y. & Excoffier, L. (1996). A generic estimation of population subdivision using distances between alleles with special reference for microsatellite loci. *Genetics*, 142, 1061-1064.
- Naruhashi, N. (1979). Notes on Japanese *Rubus* (3). *Journal of Phytogeography and Taxonomy*, Vol.27, pp. 38–40, ISSN 0388-6212
- Naruhashi, N. (1990). *Rubus* 3 semi-nepalensis, a new natural hybrid from Nepal Himalaya. *Journal of Japanese Botany*, Vol.65, pp. 186–191, ISSN 0022-2062
- Parent J. G. & M. G. Fortin. (1993). Identification of raspberry cultivars by random amplified polymorphic DNA (RAPD) analysis. *Canadian Journal of Plant Science* 73(4): 1115 – 1122.
- Parent, J.G. & Page, D. (1998). Identification of raspberry cultivars by sequence characterized amplified region DNA analysis. *HortScience*, Vol.33, No.1, pp. 140–142, ISSN 0018-5345
- Peakall, R. & Smouse, P. (2006). GENALEX 6: Genetic analysis in Excel. Population genetics software for teaching and research. *Molecular Ecology Notes*, Vol.6, pp. 288–295, ISSN 1471-8278
- Randell, R.A.; Howarth, D.G. & Morden, C.W. (2004). Genetic analysis of natural hybrids between endemic and alien *Rubus* (Rosaceae) species in Hawaii. *Conservation Genetics*, Vol.5, No.2, pp. 217–230, ISSN 1566-0621
- Romoleroux, K. (1992). Rosaceae in the paramos of Ecuador, In: *Paramo: An Andean Ecosystem under Human Influence*, H. Balslev & J.L. Luteyn (Eds.), 85-94, Academic Press, ISBN 0124604420, London
- Steele, F.L. & Hodgdon, A.R. (1963). Hybridization of *Rubus hispidus* and *R. setosus*. *Rhodora*, Vol.65, pp. 262–270, ISSN 0035-4902
- Steele, F.L. & Hodgdon, A.R. (1970). Hybrids in *Rubus* subgenus *Eubatus* in New England. *Rhodora*, Vol.72, pp. 240–249, ISSN 0035-4902
- Thompson, M.M. (1995). Chromosome numbers of *Rubus* cultivars at the National Clonal Germplasm Repository. *HortScience*, Vol. 30, pp. 1453–1456, ISSN 0018-5345
- Thompson, M.M. (1997). Survey of chromosome numbers in *Rubus* (Rosaceae: Rosoideae). *Annals of the Missouri Botanical Garden*, Vol. 84, pp. 128–164, ISSN 0026-6493
- Vargas, W. (2002). *Guía Ilustrada de las Plantas de las Montañas del Quindío y los Andes Centrales*. Universidad de Caldas and Corporación Autónoma Regional del Quindío, ISBN 958-8041-38-4, Manizales, Colombia

Weber, C.A. (2003). Genetic diversity in black raspberry detected by RAPD markers.
HortScience, Vol.38, No.2, pp. 269–272, ISSN 0018-5345

IntechOpen

IntechOpen



The Molecular Basis of Plant Genetic Diversity

Edited by Prof. Mahmut Caliskan

ISBN 978-953-51-0157-4

Hard cover, 374 pages

Publisher InTech

Published online 30, March, 2012

Published in print edition March, 2012

The Molecular Basis of Plant Genetic Diversity presents chapters revealing the magnitude of genetic variations existing in plant populations. Natural populations contain a considerable genetic variability which provides a genomic flexibility that can be used as a raw material for adaptation to changing environmental conditions. The analysis of genetic diversity provides information about allelic variation at a given locus. The increasing availability of PCR-based molecular markers allows the detailed analyses and evaluation of genetic diversity in plants and also, the detection of genes influencing economically important traits. The purpose of the book is to provide a glimpse into the dynamic process of genetic variation by presenting the thoughts of scientists who are engaged in the generation of new ideas and techniques employed for the assessment of genetic diversity, often from very different perspectives. The book should prove useful to students, researchers, and experts in the area of conservation biology, genetic diversity, and molecular biology.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Marta L. Marulanda, Ana M. López and Marcela Uribe (2012). Genetic Diversity and Transferability of Rubus Microsatellite Markers to South American Rubus Species, The Molecular Basis of Plant Genetic Diversity, Prof. Mahmut Caliskan (Ed.), ISBN: 978-953-51-0157-4, InTech, Available from:
<http://www.intechopen.com/books/the-molecular-basis-of-plant-genetic-diversity/genetic-diversity-and-transferability-of-rubus-microsatellite-markers-to-south-american-rubus-specie>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen