

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

185,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



## General Description of *Rhizoctonia* Species Complex

Genhua Yang and Chengyun Li

Key Laboratory of Agro-Biodiversity and Pest Management of Education  
Ministry of China, Yunnan Agricultural University, Kunming, Yunnan  
China

### 1. Introduction

The genus concept of *Rhizoctonia* spp. was established by de Candolle (1815) (Sneh *et al.*, 1998). However, the lack of specific characters led to the classification of a mixture of unrelated fungi as *Rhizoctonia* spp. (Parmeter and Whitney, 1970; Moore, 1987). Ogoshi (1975) enhanced the specificity of the genus concept for *Rhizoctonia* by elevating the following characteristics of *R. solani* to the genus level. Based on this revised genus concept, species of *Rhizoctonia* can be differentiated by mycelia color, number of nuclei per young vegetative hyphal cell and the morphology of their teleomorph. The teleomorph of *Rhizoctonia* spp. belongs to the sub-division Basidiomycota, class Hymenomycetes.

The anamorphs of *Rhizoctonia* are heterogeneous. Moore (1987) placed the anamorphs of *Thanatephorus* spp. in *Moniliopsis*. She reserved the genus *Rhizoctonia* for anamorph of ustomycetous fungi which have septa with simple pores. *Moniliopsis* species have smooth, broad hyphae with brown walls, multinucleate cells, dolipore septa with perforate parenthesomes and teleomorphs in the genera *Thanatephorus* and *Waitea*. Of the binucleate *Rhizoctonia* spp., the anamorphs of the *R. repens* group (teleomorph *Tulasnella*) were assigned to the new genus *Epulorhiza*. Anamorph of *Ceratobasidium* was assigned to the new genus *Ceratorhiza* (Moore, 1987). Moore's system is taxonomically correct and justified. At present, the concept of genus *Rhizoctonia* has become clear from these taxonomical studies at the molecular level (Gonzalez *et al.*, 2001). However, many researchers (Sneh *et al.*, 1998) in the world still retain the name *Rhizoctonia* for Moore's *Moniliopsis* spp., *Ceratorhiza* spp. and *Epulorhiza* spp.. Hence, I used the name of *Rhizoctonia* in this study.

Affinity for hyphal fusion (anastomosis) (Parmeter *et al.*, 1969; Parmeter and Whitney, 1970; Ogoshi *et al.*, 1983a; Burpee *et al.*, 1980a) has been used to characterize isolates among *R. solani*, *R. zeae*, *R. oryzae*, *R. repens* and binucleate *Rhizoctonia* spp. with *Ceratobasidium* teleomorphs. To date, isolates of *R. solani* have been assigned to 13 anastomosis groups (AG) and those of *R. zeae* and *R. oryzae* have each been assigned to their own one group (Sneh *et al.*, 1998; Carling *et al.*, 1999, 2002c).

Anastomosis reactions between hyphae of paired isolates of *R. solani* consist of several types; such as perfect fusion, imperfect fusion, contact fusion and no reaction (Matsumoto *et al.*, 1932). At present, four categories of anastomosis (C3 to C0) defined by Carling *et al.* (1996)

have been accepted by many researchers. These are useful for a better understanding of the genetic diversity of *R. solani* populations, because of the background genetically supported by vegetative or somatic compatibility (VC or SC) of confronted isolates (MacNish *et al.*, 1997). Each of categories is as follows:

**C3:** walls fuse; membranes fuse, accompanied with protoplasm connection; anastomosis point frequently is not obvious; diameter of anastomosis point is equal or nearly equal hyphal diameter; anastomosing cells and adjacent cells may die, but generally do not. This category occurs for the same anastomosis group, same vegetative compatibility population (VCP) and the same isolate.

**C2:** wall connection is obvious, but membrane contact is uncertain; anastomosing and adjacent cells always die. This category occurs in same AG, but not between different VCPs.

**C1:** wall contact between hyphae is apparent, but both wall penetration and membrane-membrane contact do not occur; occasionally one or both anastomosing cells and adjacent cells die. This category occurs between different AGs or in the same AG.

**C0:** no reaction. This category occurs between different AGs.

In general, hyphal fusion occurs at a high frequency (50% $\geq$ ) within members of the same AG, with the exception of non-self-anastomosing isolates (Hyakumachi and Ui, 1988). On the other hand, hyphal fusion among members of different AGs occurs at either a low frequency ( $\leq$ 30%) or no fusion occurs. *Rhizoctonia* isolates giving C3 to C1 reactions in anastomosing test have been taken to be the same AG.

To date, isolates of multinucleate *R. solani* have been assigned to 13 anastomosis groups (AG-1 to AG-13), some of which include several subgroups and isolates of *R. zeae* and *R. oryzae* have been assigned to WAG-Z and WAG-O, respectively (Sneh *et al.*, 1998; Carling *et al.*, 1999, 2002c). Isolates of binucleate *Rhizoctonia* spp. with *Ceratobasidium* teleomorphs have been reported. A system developed in Japan (Ogoshi *et al.*, 1979, 1983 a,b; Sneh *et al.*, 1998; Hyakumachi *et al.*, 2005) includes 21 anastomosis groups designated AG-A to AG-U, in which at present AG-J and AG-M still are in question as members of binucleate *Rhizoctonia*. Another system developed in the USA (Burpee *et al.*, 1980a) includes 7 anastomosis groups designed as CAG-1 to CAG -7. CAG-1 corresponds to AG-D, CAG-2 to AG-A, CAG-3 and CAG-6 to AG-E, CAG-4 to AG-F, CAG-5 to AG-R, and CAG-7 to AG-S (Sneh *et al.*, 1998; Ogoshi *et al.*, 1983a). At present, the anastomosis system based on AG-A through AG-U used in this review paper is widely accepted by many researchers.

Some homogenous groups of isolates of *R. solani* are well known as bridging isolates (AG-BI) that anastomose with members of different AGs (Carling, 1998). In general, there is no contradiction in the conventional anastomosis grouping system by taking anastomosis frequency into consideration. However, two exceptional cases where anastomosis frequency mismatched with morphological, physiological and pathogenic characteristics have been reported from tobacco (Nicoletti *et al.*, 1999) and soybean (Naito and Kanematsu, 1994). These demonstrate the limitations of using hyphal anastomosis as the sole criteria for characterization and identification of closely related fungi. In addition, it is not easy to determine the subgroup of isolates within the same AG because no differences occur in their anastomosis reaction. Thus, in order to determine AGs or subgroups in *R. solani*, genetic analysis using molecular approaches that employ multiple genetic loci is needed.

Isolates of *R. solani* that exhibits DNA base sequence homology and affinities for hyphal anastomosis may represent a diverging evolutionary unit (Kuninaga and Yokosawa, 1980). This hypothesis is supported by analysis of restriction fragment length polymorphisms (RFLPs) and the sequences with in ribosomal RNA genes (rDNA) among different anastomosis groups of *R. solani* (Vilgalys and Gonzalez, 1990; Gonzalez, *et al.*, 2001; Carling *et al.*, 2002b).

As mentioned above, many AGs and subgroups of *R. solani* and binucleate *Rhizoctonia* spp. have been reported as causal of agents Rhizoctonia diseases on a wide range of host species. However, little is known about the Rhizoctonia diseases and the anastomosis groups and subgroups of their causal fungi on vegetables, ornamentals and food crops in the Asian tropics especially the southern parts of China.

## 2. Characteristics of anastomosis groups and subgroups of *Rhizoctonia solani* and binucleate *Rhizoctonia* spp.

Disease symptoms and host range of each AG and its subgroups are summarized as follows. In this review, the book by Sneh *et al.*, 1998 entitled "Identification of *Rhizoctonia* Species" provided a substitute for the reference before 1998.

### 2.1 Multinucleate *Rhizoctonia* spp.

#### 1. AG-1: IA, IB, IC, ID

**AG-1 IA** (Li and Yan, 1990; Sneh *et al.*, 1998; Fenille *et al.*, 2002; Naito, 2004).

Symptoms: sheath blight, foliar blight, leaf blight, web-blight, head rot, bottom rot, and brown patch.

Host: rice (*Oryza sativa* L.), corn (*Zea mays* L.), barley (*Hordeum vulgare* L.), sorghum (*Sorghum vulgare* Pes.), potato (*Solanum tuberosum* L.), barnyard millet, common millet, soybean, peanut (*Arachis hypogaea* L.), lima bean, cabbage, leaf lettuce, Stevia, orchard grass, crimson clover, tall fescue (*Festuca arundinacea* Schreb), turfgrass, creeping bentgrass, perennial ryegrass, gentian (*Gentiana scabra*), and camphor.

Note: This group has a tendency to attack aerial parts of the plants. Basidiospore infection of rice has been reported, but sclerotia are more important as an infection source. The optimum growth temperature is higher than those of AG-1 IB.

**AG-1 IB** (Sneh *et al.*, 1998; Naito, 2004; Yang *et al.*, 2005b).

Symptoms: sheath blight, leaf blight, foliar blight, web-blight, root rot, damping-off, head rot, and bottom rot.

Host: corn, sugar beet, gay feather (*Liatris* spp.), common bean, fig (*Ficus* L.), adzuki bean, soybean, cabbage, leaf lettuce, redtop, bentgrass, orchard grass, leaf lettuce, apple (*Malus pumila* Mill), Japanese pear, European pear, lion'ear (*Leonotis leonurus*), hortensia (*Hydrangea* spp.), *Larix* spp., gazania (*Gazania* spp.) *Cotoneaster* spp., Egyptian atar-cluster (*Pentas lanceolata*), Chinese lantern plant (*Physalis alkekeng* var. *franchetii*), *Hypericum patulum*, marigold, *Acacia* spp., rosemary, *Eucalyptus* spp., pine (*Pinus* L.), *Larix* spp., cypress (*Cupressus* spp.), and elephant foot (*Amorphophallus Konjac*).

**AG-1 IC** (Sneh *et al.*, 1998; Naito, 2004).

Symptoms: damping-off, summer blight, foot rot, crown rot canker, and root rot.

Host: sugar beet, carrot (*Daucus carota* L.), buckwheat (*Eriogonum* Michx), flax (*Linum usitatissimum* L.), soybean, bean (*Phaseolus* L.), cabbage, pineapple (*Ananas comosus* (Linn.) Merr.), panicum (*Panicum* spp.), spinach (*Spinacia oleracea* L.), and radish (*Raphanus sativus* Linn.).

**AG-1 ID** (Priyatmojo *et al.*, 2001).

Symptom: leaf spot.

Host: coffee (*Coffea* Linn.).

Note: this subgroup was recently reported in the Philippines (Priyatmojo *et al.*, 2001)

Undetermined subgroup: buckwheat, flax, spinach, radish, and durian (*Durio zibethinus* Murr.).

## 2. **AG-2: 2-1, 2-2 IIIB, 2-2 IV, 2-2 Lp, 2-3, 2-4, 2-BI.**

**AG-2-1** (Satoh *et al.*, 1997; Camporota and Perrin, 1998; Sneh *et al.*, 1998; Rollins *et al.*, 1999; Khan and Kolte, 2000; Naito, 2004)

Symptoms: damping-off, leaf rot, leaf blight, root rot, foot rot, bottom rot, and bud rot.

Host: sugar beet, wheat (*Triticum aestivum* Linn.), potato, cowpea (*Vigna unguiculata* (Linn.) Walp), canola, rape (*Brassica napus* Linn.), cauliflower (*Brassica oleracea* var. *botrytis* Linn.), mustard (*Sinapis* Linn.), turnip (*Brassica rapa* Linn.), pepper (*Piper* Linn.), *Silene armeria*, spinach, leaf lettuce, strawberry (*Fragaria ananassa* Duchesne), tulip (*Tulipa gesneriana* Linn.), tobacco (*Nicotiana* Linn.), clover (*Medicago* Linn.), and table beet.

Note: This group includes the AG-2-1 tulip strain (former AG-2t) and the AG-2-1 tobacco strain (former homogenous Nt-isolates) (Kuninaga *et al.*, 2000).

**AG-2-2 III B** (Sneh *et al.*, 1998; Priyatmojo *et al.*, 2001; Naito, 2004).

Symptoms: brown sheath blight, dry root rot, root rot, brown patch, large patch, black scurf, stem rot, stem blight, *Rhizoctonia* rot, damping-off, stem rot, collar rot, and crown brace rot.

Host: rice, soybean, corn, sugar beet, edible burdock (*Arctium lappa*), taro (*Colocasia esculenta*), *Dryopteris* spp., elephant foot, crocus, saffron (*Crocus sativus* Linn.), redtop, bentgrass, St. Augustine grass, turf, balloon flower (*Platycodon grandiflorum*), Christmas-bells (*Sandersonia aurantiaca*), *Hedera rhombea*, mat rash, *gladiolus*, ginger, and *Iris* Linn..

**AG-2-2 IV:** (Sneh *et al.*, 1998; Naito, 2004).

Symptoms: leaf blight, foliage rot, root rot, and stem rot.

Host: sugar beet, carrot, eggplant (*Solanum* Linn), pepper, spinach, stevenia (*Stevenia* Adams et Fisch), and turfgrass.

**AG-2-2 LP:** (Aoyagi *et al.*, 1998).

Symptoms: large patch.



Host: Zoysia grass.

**AG 2-3:** (Naito and Kanematsu, 1994; Sumner *et al.*, 2003).

Symptoms: leaf blight and root rot.

Host: soybean.

Note: basidiospores cause leaf spot of soybean.

**AG-2-4:** (Sumner, 1985).

Symptoms: crown rot, brace rot, and damping-off.

Host: corn and carrot.

**AG-2-BI:** (Carling *et al.*, 2002b).

Symptoms: nonpathogenic.

Host: isolates, obtained only from soils and plants in forests.

Note: former name is AG-BI.

Undetermined subgroup: sesame (*Sesamum* Linn.), white mustard (*Sinapsis alba*), primrose (*Primula* spp.), white lace flower (*Ammi majus*), carnation, baby's-breath (*Gypsophila paniculata*), russel prairie gentian (*Eustoma grandiflorum*), snap bean, lima bean, and Chinese radish.

3. **AG 3: PT, TB** (Sneh *et al.*, 1998; Kuninaga *et al.*, 2000).

Symptoms: black scurf, leaf spot, target leaf spot, and damping-off.

PT: potato with black scurf symptoms.

TB: tobacco with target leaf spot symptoms.

Note: Undetermined subgroup: eggplant, sugar beet, tomato, and wheat. Their pathological and ecological information is less.

4. **AG-4: HG-I, HG-II, HG-III** (Baird, 1996; Holtz *et al.*, 1996; Sneh *et al.*, 1998; Fenille *et al.*, 2002; Ravanlou and Banihashemi, 2002; El Hussieni, 2003; Kuramae *et al.*, 2002, 2003; Naito, 2004; Yang *et al.*, 2005c).

Symptoms: damping-off, root rot, stem canker, fruit rot, and stem rot.

Host: pea, sugar beet, melon, soybean, adzuki bean, common bean, snap bean, lima bean, carrot, spinach, taro, tomato (*Lycopersicon esculentum* Mill.), potato, alfalfa (*Medicago sativa* Linn.), elephant foot, arrowleaf clover, beans, barley, buckwheat, cabbage, canola, turnip, carnation, cauliflower, Chinese chive, chrysanthemum, corn, cotton (*Gossypium* Linn.), table beet, tobacco, turfgrass, wheat, white lupine, parsley (*Petroselinum* Hill), *Cineraria* Linn., stock, poinsettia, primrose, hybrid bouvardia, *Citrus* Linn., cauliflower, *Euphorbia* spp., geranium (*Pelargonium* spp.), Russel prairie gentian, statice (*Limonium* spp.), baby's-breath, and *Astragalus membranaceus*.

5. **AG-5** (Li, *et al.*, 1998; Demirci, 1998; Sneh *et al.*, 1998; Ravanlou and Banihashemi, 2002; Eken and Demirci, 2004; Naito, 2004).

Symptoms: root rot, damping-off, black scurf, brown patch, and symbiosis (orchids).

Host: soybean, adzuki bean, apple, barley, chickpea, common bean, lima bean, potato, strawberry, sugar beet, table beet, tobacco, turfgrass, wheat, and white lupine.

6. **AG-6: HG-I, GV** (Mazzola, 1997; Meyer *et al.*, 1998; Sneh *et al.*, 1998; Carling *et al.*, 1999; Pope and Carter, 2001; Naito, 2004)

Symptom: root rot, crater rot, and symbiosis (orchids).

Host: apple, wheat, carrot, and carnation.

Note: all isolates from forests are nonpathogenic.

7. **AG-7:** (Naito, *et al.*, 1993; Baird and Carling, 1995; Carling, 1997, 2000; Carling *et al.*, 1998)

Symptoms: damping-off, root rot, and black scurf.

Host: carnation, cotton, soybean, watermelon (*Citrullus lanatus* (Thunb.) Mansfeld), *Raphanus* Linn., and potato.

8. **AG-8:** (Sneh *et al.*, 1998; Naito, 2004).

Symptoms: bare patch.

Host: barley, cereals, green pepper, potato, and wheat.

9. **AG-9:** (Sneh *et al.*, 1998; Naito, 2004).

Symptoms: black scurf.

Host: potato, crucifers, wheat, and barley.

10. **AG-10:** (Sneh *et al.*, 1998.)

Symptoms: weak pathogenic.

Host: barley and wheat.

11. **AG-11:** (Kumar *et al.*, 2002).

Symptoms: damping-off and hypocotyls rot.

Host: barley, lupine, soybean, and wheat.

Note: this group is considered as bridging isolates (anastomose with each members of AG-2-1, AG-2 BI, AG-8) (Carling *et al.*, 1996).

12. **AG-12:** (Kumar *et al.*, 2002).

Symptoms: symbiosis (orchids).

Host: *Dactylorhiza aristata* (Orchidaceae).

13. **AG-13:** (Carling *et al.*, 2002a).

Symptoms: none.

Host: cotton.

## 2.2 Binucleate *Rhizoctonia* spp.

### 1. AG-A: (Mazzola, 1997; Sneh *et al.*, 1998).

Symptoms: root rot, damping-off, browning, and tortoise shell.

Host: strawberry, sugar beet, bean, pea, sunflower (*Helianthus annuus* Linn.), tomato, melon, cucumbeare (*Cucumis sativas* Linn.), leaf lettuce, spinach, peanut, potato, *Solanum tuberosum*, and apple.

Note: Some isolates in this group form mycorrhizal associations with orchids.

### 2. AG-B: a and b.

**AG-Ba** (Sneh *et al.*, 1998).

Symptoms: grey sclerotium disease, sclerotium disease, gray southern blight.

Host: rice, *Echinochloa crugalli* subsp. *submitica* var. *typica*, and foxtail millet.

**AG-Bb** (Sneh *et al.*, 1998).

Symptoms: brown sclerotium disease, grey sclerotium disease, and sheath spot.

Host: fox tail, millet, and rice.

### 3. AG-C (Sneh *et al.*, 1998; Hayakawa *et al.*, 1999).

Symptoms: symbiosis (orchids).

Host: orchids, sugar beet seedlings, subterranean clover, and wheat.

Note: No important pathogens have been reported.

### 4. AG-D: I, II (Sneh *et al.*, 1998; Toda *et al.*, 1999).

Symptoms: sharp eye spot, yellow patch, foot rot, Sclerotium disease, snow mold, root rot, damping-off, lesions on stems, and winter stem rot.

Host: cereals, turf grass, wheat, barley, sugar beet, clove, pea, onions (*Allium cepa* Linn.), potato, cotton, bean, soybean, mat rush, foxtail millet, and subterranean clover.

Note: Recently this group is classified into subgroup AG-D (I) that causes Rhizoctonia patch and winter patch diseases. AG-D (II) causes elephant footprint disease.

### 5. AG-E (Sneh *et al.*, 1998).

Symptoms: web-blight, damping-off, seedlings, and symbiosis (orchids).

Host: bean, pea, radish, onion, leaf lettuce, tomato lima bean, snap bean, soybean, peanut, cowpea (*Vigna Savi*), flax, sugar beet, *Rhododendron* Linn., long leaf pine (*Pinus palustris* Mill.), slash, lobolly pine (*Pinus taeda* Linn.), and rye (*Secale cereale* Linn.).

### 6. AG-F (Sneh *et al.*, 1998; Eken and Demirci, 2004).

Symptoms: none.

Host: bean, pea, radish, onion, peanut, leaf lettuce, tomato, subterranean clover radish, tomato, cotton, taro, strawberry (source: DDJB), and *Fragaria x ananassa*.



7. **AG-G** (Mazzola, 1997; Sneh *et al.*, 1998; Leclerc *et al.*, 1999; Martin, 2000; Botha *et al.*, 2003; Fenille *et al.*, 2005).

Symptoms: damping-off, root rot, and browning.

Host: strawberry, sugar beet, bean, pea, tomato, melon, sunflower, peanut, yacoon, apple, *Rhododendron* Linn., and *Fragaria x ananassa*.

Note: Non-pathogenic binucleate *Rhizoctonia* spp. provide effective protection to young bean seedlings against root rot caused by *R. solani* AG-4 (Leclerc *et al.*, 1999).

8. **AG-H** (Hayakawa *et al.*, 1999).

Symptoms: symbiosis (orchids).

Host: *Dactylorhiza aristata* (Orchidaceae).

9. **AG-I** (Mazzola, 1997; Sneh *et al.*, 1998; Ravanlou and Banihashemi, 2002)

Symptoms: root rot and symbiosis (orchids).

Host: strawberry, sugar beet, wheat, apple, orchids, and *Fragaria x ananassa*.

10. **AG-J**: (Sneh *et al.*, 1998).

Symptoms: none.

Host: apple.

11. **AG-K** (Demirci, 1998; Li *et al.*, 1998; Sneh *et al.*, 1998; Ravanlou and Banihashemi, 2002).

Symptoms: none.

Host: sugar beet, radish, tomato, carrot, onion, wheat, maize, *Allium cepa* (source: DDJB),

*Pyrus communis* (pear) (source: DDJB), and *Fragaria x ananassa*.

12. **AG-L**: No special diseases have been reported (Sneh *et al.*, 1991).

13. **AG-N**: No special diseases have been reported (Sneh *et al.*, 1991).

14. **AG-O**: No special diseases have been reported (Mazzola, 1997; Sneh *et al.*, 1998).

Host: apple.

15. **AG-P**: (Sneh *et al.*, 1998; Yang *et al.*, 2006).

Symptoms: black rot and wirestem.

Host: tea (*Camellia* Linn.), red birch.

16. **AG-Q**: (Sneh *et al.*, 1998).

Symptoms: none.

Host: (Bentgrass).

17. **AG-R**: (Sneh *et al.*, 1998; Yang *et al.*, 2006).

Symptoms: wirestem

Host: bean, pea, radish, onion, leaf lettuce, tomato, lima bean, snap bean, soybean, cowpea, peanuts, red birch, and azalea.

18. **AG-S** (Demirci, 1998; Sneh *et al.*, 1998).

Symptoms: no specific diseases.

Host: azalea, wheat, barley, and azalea.

19. **AG-T**: (Hyakumachi *et al.*, 2005).

Symptoms: stem rot and root rot.

Host: miniature roses.

20. **AG-U**: (Hyakumachi *et al.*, 2005).

Symptoms: stem rot and root rot.

Host: miniature roses (*Rosa rugosa* Thunb.).

### 3. Summary

In this chapter, we described the classification of *Rhizoctonia* spp. complex. Mutinucleate *Rhizoctonia* spp. included 13 anastomosis, of which AG 1-4 were strong pathogenic on many plants and AG 6-10 were orchid mycorrhizae. Binucleate *Rhizoctonia* spp. included 18 anastomosis groups, but AG-U belonged to AG-P and AG-T belonged to AG-A (Sharon *et al.*, 2008), which were weak or nonpathogenic to plants and some AGs were orchid mycorrhizae.

### 4. Acknowledgments

This work was supported by the National Basic Research Program (No. 2011CB100400) from The Ministry of Science and Technology of China and the National Natural Science Funds, China (30660006, and 31160352), respectively.

### 5. References


- Aoyagi, T., Kageyama, K. & Hyakumachi, M. (1998) Characterization and survival of *Rhizoctonia solani* AG2-2 LP associated with large patch disease of zoysia grass. *Plant Dis.* 82, 857-863.
- Baird, R. E. (1996) First report of *Rhizoctonia solani* AG-4 on canola in Georgia. *Plant Dis.* 80(1), 104.
- Baird, R. E. & Carling, D. E. (1995) First report of *Rhizoctonia solani* AG-7 in Indiana. *Plant Dis.* 79(3), 321.
- Botha, A., Denman, S., Lamprecht, S. C., Mazzola, M. & Crous, P. W. (2003) Characterization and pathogenicity of *Rhizoctonia* isolates associated with black root rot of strawberries in the Western Cape Province, South Africa. *Australasian Plant Pathol.* 32(2), 195-201.
- Burpee, L. L., Sanders, P. L., Cole, H. Jr. & Sherwood, R. T. (1980a) Anastomosis groups among isolates of *Ceratobasidium cornigerum* and related fungi. *Mycologia* 72, 689-701.
- Camporota, P. & Perrin, R. (1998) Characterization of *Rhizoctonia* species involved in tree seedling damping-off in French forest nurseries. *Appl. Soil Ecol.* 10(1/2), 65-71.
- Carling, D. E. (1997) First report of *Rhizoctonia solani* AG-7 in Georgia. *Plant Dis.* 82(1), 127.

- Carling, D. E. (2000) Anastomosis groups and subsets of anastomosis groups of *Rhizoctonia solani*. Abstract in Proceedings of the Third International Symposium on *Rhizoctonia*. National Chung Hsing University, Taichung, Taiwan .14 pp.
- Carling, D. E., Baird, R. E., Gitaitis, R. D., Brainard, K. A. & Kuninaga, S. (2002a) Characterization of AG-13, a newly reported anastomosis group of *Rhizoctonia solani*. *Phytopathology* 92(8), 893-899.
- Carling, D. E., Brainard, K.A., Virgen-Calleros, G. & Olalde-Portugal, V.F. (1998) First report of *Rhizoctonia solani* AG-7 on potato in Mexico. *Plant Dis.* 82(1), 127.
- Carling, D. E., Kuninaga, S. & Brainard, K. A. (2002b) Hyphal anastomosis reactions, rDNA-internal transcribed spacer sequences, and virulence levels among subsets of *Rhizoctonia solani* anastomosis group-2 (AG-2) and AG-BI. *Phytopathology* 92, 43-50.
- Carling, D. E., Pope, E. J., Brainard, K. A. & Carter, D. A. (1999) Characterization of mycorrhizal isolates of *Rhizoctonia solani* from an orchid, including AG-12, a new anastomosis group. *Phytopathology* 89(10), 942-946.
- Carling, D. E. (1996) Grouping in *Rhizoctonia solani* by hyphal anastomosis reaction. In: Sneh, B., Jabaji-Hare, S., Neat, S. and Dijst, G. *et al.*, (eds). *Rhizoctonia* species: Taxonomy, Molecular Biology, Ecology, Pathology and Disease Control. Kluwer Academic Publishers, Dordrecht, The Netherlands, pp 37-47.
- Carling, D. E., Baird, R. E., Gitaitis, R. D., Brainard, K. A. & Kuninaga, S. (2002c) Characterization of AG-13, a newly reported anastomosis group of *Rhizoctonia solani*. *Phytopathology* 92,893-899.
- de Candolle, A. 1815. Uredo rouille des cereals In Forafran caise, famille des champignons p.83.
- Demirci, E. (1998) *Rhizoctonia* species and anastomosis groups isolated from barley and wheat in Erzurum, Turkey. *Plant Pathol.* 47(1), 10-15.
- Eken, C. & Demirci, E. (2004) Anastomosis groups and pathogenicity of *Rhizoctonia solani* and binucleate *Rhizoctonia* isolates from bean in Erzurum, Turkey J. *Plant Pathol.* 86(1), 49-52.
- Fenille, R.C., Ciampi, M.B., Souza, N.L., Nakataniand, A.K. and Kuramae, E.E.(2005) Binucleate *Rhizoctonia* sp. AG-G causing rot rot in yacon (*Smallanthus sonchifolius*) in Brazil. *Plant patho.* 54,325-330.
- Fenille, R. C., Luizde S. N. & Kuramae, E. E. (2002) Characterization of *Rhizoctonia solani* associated with soybean in Brazil. *Eur. J. of Plant Pathol.* 108(8), 783-792.
- Gonzalez, D., Carling, D. E., Kuninaga, S., Vilgalys, R. & Cubeta, M. A. (2001) Ribosomal DNA systematics of *Ceratobasidium* and *Thanatephorus* with *Rhizoctonia* anamorphs. *Mycologia* 93 (6), 1138-1150.
- Hayakawa, S., Uetake, Y. & Ogoshi, A.(1999) Identification of symbiotic rhizoctonia from naturally occurring protocorms and roots of *Dactylorhiza aristata* (Orchidaceae). *J. of the Faculty of Agriculture Hokkaido University* 69(2), 129-141.
- Holtz, B. A., Michailides, T. J., Feguson, L., Hancock, J. G. & Weinhold, A. R. (1996) First report of *Rhizoctonia solani* (AG-4) on pistachio rootstock seedlings. *Plant Dis.* 80(11), 1303.
- Hyakumachi, M., Priyatmojo, A., Kubota, M. & Fukui, H. (2005) New anastomosis groups, AG-Tand AG-U, of binucleate *Rhizoctonia* spp. causing root and stem rot of cut-flower and miniature roses. *Phytopathology* 95,784-792.
- Hyyakumachi, M. and Ui,T. (1988)Development of the teleomorph of non-self-anastomosing isolates of *Rhizoctonia solani* by a buried-slide method *plant pathol.* 37(3):438-440
- Khan, R. U. & Kolte, S. J. (2000) Some seedling diseases of rapeseed-mustard and their control. *Indian Phytopathol.* 55(1), 102-103.

- Kumar, S., Sivasithamparam, K. & Sweetingham, M. W. (2002) Prolific production of sclerotia in soil by *Rhizoctonia solani* anastomosis group (AG) 11 pathogenic on lupin. *Annal. Appl. Biol.* 141(1), 11-18.
- Kuninaga, S. & Yokosawa, R. (1980) A comparison of DNA compositions among anastomosis groups in *Rhizoctonia solani* Kühn. *Ann. Phytopathol. Soc. Japan* 46,: 150-158.
- Kuninaga, S., Nicoletti, R., Lahoz, E. & Naito, S. (2000). Ascription of Nt-isolates of *Rhizoctonia solani* to anastomosis group 2-1 (AG-2-1) on account of rDNA-ITS sequence similarity. *J. Plant Pathol.* 82, 61-64.
- Leclerc, P. C., Balmas, V., Charest, P. M. & Jabaji, H. S. (1999) Development of reliable molecular markers to detect non-pathogenic binucleate *Rhizoctonia* isolates (AG-G) using PCR. *Mycol. Res.* 103(9), 1165-1172.
- Li, H.R. & Yan, S.Q. (1990) Studies on the strains of pathogens of sheath blight of rice in the east and south of Sichuan Province. *Acta Mycologica Sinica* 9: 41-9. (Chinese with English abstract).
- Li, H.R., Wu, B.C. & Yan, S. Q. (1998) Aetiology of *Rhizoctonia* in sheath blight of maize in Sichuan. *Plant Pathol.* 47 (1), 16-21.
- MacNish, G. C., Carling, D. E. & Brainard, K. A. (1997) Relationship of microscopic vegetative reactions in *Rhizoctonia solani* and the occurrence of vegetatively compatible populations (VCP) in AG-8. *Mycol. Res.* 100, 61-68.
- Martin, F. N. (2000) *Rhizoctonia* spp. recovered from strawberry roots in central coastal California. *Phytopathology* 90, 345-353.
- Mazzola, M. (1997) Identification and pathogenicity of *Rhizoctonia* spp. isolated from apple roots and orchard soils. *Phytopathology* 87, 582-587.
- Meyer, L., Wehner, F. C., Nel, L. H. & Carling, D. E. (1998) Characterization of the crater disease strain of *Rhizoctonia solani*. *Phytopathology* 88, 366-371.
- Moore, R. T. (1987) The genera of *Rhizoctonia*-like fungi: *Asorhizoctonia*, *Ceratorhiza* gen. nov., *Epulorhiza* gen. nov., *Moniliopsis* and *Rhizoctonia*. *Mycotaxon* 29, 91-99.
- Naito, S. (2004) *Rhizoctonia* diseases: Taxonomy and population biology. *Proceeding of the International Seminar on Biological Control of Soilborne Plant Diseases, Japan-Argentina Joint Study, Buenos Aires, Argentina*, p.18-31.
- Naito, S. & Kanematsu, S. (1994) Characterization and pathogenicity of a new anastomosis subgroup AG-2-3 of *Rhizoctonia solani* Kühn isolated from leaves of soybean. *Ann. Phytopath. Soc. Japan* 60(6), 681-690.
- Naito, S., Mohamad, D., Nasution, A. & Purwanti, H. (1993) Soilborne diseases and ecology of pathogens on soybean roots in Indonesia. *JARQ* 26, 247-253.
- Nicoletti, R., Lahoz, E., Kanematsu, S., Naito, S. & Contillo, R. (1999) Characterization of *Rhizoctonia solani* isolates from tobacco fields related to anastomosis groups 2-1 and BI (AG 2-1 and AG BI). *J. Phytopathology* 147 (2), 71-77.
- Ogoshi, A. (1975) Grouping of *Rhizoctonia solani* Kühn and their perfect stages. *Rev. Plant. Protect. Res.* 8, 93-103.
- Ogoshi, A. & Ui, T. (1979) Specificity in vitamin requirement among anastomosis groups of *Rhizoctonia solani* Kühn. *Ann. Phytopathol. Soc. Jpn.* 45, 47-53.
- Ogoshi, A., Oniki, M., Araki, T. & Ui, T. (1983a) Anastomosis groups of binucleate *Rhizoctonia* in Japan and North America and their perfect states. *Mycol. Soc. Japan* 24, 79-87.

- Ogoshi, A., Oniki, M., Araki, T. & Ui, T. (1983b) Studies on the anastomosis groups of binucleate *Rhizoctonia* and their perfect states. J. Fac. Agr. Hokkaido Univ. 61, 244-260.
- Parmeter, J. R. J., Sherwood, R. T. & Platt, W. D. (1969) Anastomosis grouping among isolates of *Thanatephorus cucumeris*. Phytopathology 59, 1270-1278.
- Parmeter, J. R. Jr. & Whitney, H. S. (1970) Taxonomy and nomenclature of the imperfect state. Pages 7-19 in: J. R. Parmeter Jr., (ed.) Biology and Pathology of *Rhizoctonia solani*. University of California Press, Berkeley. 255 pp.
- Pope, E. J. & Carter, D. A. (2001) Phylogenetic placement and host specificity of mycorrhizal isolates belonging to AG-6 and AG-12 in the *Rhizoctonia solani* species complex. Mycologia 93(4), 712-719.
- Priyatmojo, A., Escopalao, V. E., Tangonan, N. G., Pascual, C. B., Suga, H., Kageyama, K. & Hyakumachi, M. (2001) Characterization of a new subgroup of *Rhizoctonia solani* anastomosis group 1 (AG-1 ID), causal agent of a necrotic leaf spot on coffee. Phytopathology 91, 1054-1061.
- Ravanlou, A. & Banihashemi, Z. (2002) Isolation of some anastomosis groups of *Rhizoctonia* associated with wheat root and crown in Fars province. Iranian J. Plant Pathol. 38(3-4), 67-69.
- Rollins, P. A., Keinath, A. P. & Farnham, M. W. (1999) Effect of inoculum type and anastomosis group of *Rhizoctonia solani* causing wirestem of cabbage seedlings in a controlled environment. Can. J. Plant Pathol. 21(2), 119-124.
- Satoh, Y., Kanehira, T. & Shinohara, M. (1997) Occurrence of seedling damping-off of Jew's mallow, *Corchorus olitorius* caused by *Rhizoctonia solani* AG-2-1. Nippon Kingakukai Kaiho 38(2): 87-91 (Japanese with English abstract).
- Sharon, M., Kuninaga, S., Hyakumachi, M., Naito, S., & Sneh B. (2008) Classification of *Rhizoctonia* spp. using rDNA-ITS sequence analysis supports the genetic basis of the classical anastomosis grouping. Mycoscience 49, 93-114.
- Sneh, B., Burpee, L. and Ogoshi, A. (1998) Identification of *Rhizoctonia* species. The APS, St. Paul, Minnesota.
- Sumner, D. R. (1985) First report of *Rhizoctonia solani* AG-2-4 on carrot in Georgia. Plant Dis. 69, 25-27.
- Sumner, D. R., Phatak, S. C. & Carling, D. E. (2003) Characterization and pathogenicity of a new anastomosis subgroup AG-2-3 of *Rhizoctonia solani* Kuün isolated from leaves of soybean. Plant Dis. 87(10), 1264.
- Toda, T., Hyakumachi, M., Suga, H., Kageyama, K., Tanaka, A. & Tani, T. (1999) Differentiation of *Rhizoctonia* AG-D isolates from turfgrass into subgroups I and II based on rDNA and RAPD analyses. Eur. J. Plant Pathol. 105, 835-846.
- Vilgalys, R. & Gonzalez, D. (1990) Ribosomal DNA restriction fragment length polymorphism in *Rhizoctonia solani*. Phytopathology 80, 151-158.
- Yang G. H., Chen H. R., Naito, S., Wu, J. Y., He, X. H. & Duan, C. F. (2005b) Occurrence of foliar rot of pak choy and Chinese mustard caused by *Rhizoctonia solani* AG-1 IB in China. J. Gen. Plant Pathol. 71, 377-379.
- Yang, G. H., Naito, S., Ogoshi, A. & Dong, W. H. (2006) Identification, isolation frequency and pathogenicity of *Rhizoctonia* spp. causing the wirestem of red birch in China. J. Phytopathology 154(2), 80-83.





## **Plant Pathology**

Edited by Dr. Christian Joseph Cumagun

ISBN 978-953-51-0489-6

Hard cover, 362 pages

**Publisher** InTech

**Published online** 04, April, 2012

**Published in print edition** April, 2012

Plant pathology is an applied science that deals with the nature, causes and control of plant diseases in agriculture and forestry. The vital role of plant pathology in attaining food security and food safety for the world cannot be overemphasized.

### **How to reference**

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Genhua Yang and Chengyun Li (2012). General Description of Rhizoctonia Species Complex, Plant Pathology, Dr. Christian Joseph Cumagun (Ed.), ISBN: 978-953-51-0489-6, InTech, Available from:  
<http://www.intechopen.com/books/plant-pathology/general-description-of-rhizocotonia-species-complex>

**INTECH**  
open science | open minds

### **InTech Europe**

University Campus STeP Ri  
Slavka Krautzeka 83/A  
51000 Rijeka, Croatia  
Phone: +385 (51) 770 447  
Fax: +385 (51) 686 166  
[www.intechopen.com](http://www.intechopen.com)

### **InTech China**

Unit 405, Office Block, Hotel Equatorial Shanghai  
No.65, Yan An Road (West), Shanghai, 200040, China  
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元  
Phone: +86-21-62489820  
Fax: +86-21-62489821



© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen