

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Novel Elicitors Induce Defense Responses in Cut Flowers

Anastasios I. Darras

*Department of Greenhouse Cultivation and Floriculture,
Technological Educational Institute of Kalamata
Greece*

1. Introduction

Cut flower production and trade in the E.U. and the rest of the world holds the main share within the ornamental horticulture industry. Despite the global economic crisis started in 2008, changes in cut flower trade, such as the merge of the 2 major auctions in Holland (i.e. VBA and FloraHolland), resulted in stabilization or even small increases in stem number sales for the years 2008-2010 (Evans & Van der Ploeg, 2008; Anonymous, 2011). In other words, the importance of cut flower industry in global economy is undisputed, but also reflects the human need for ornamental plant consumption as part of a better life.

Product quality of horticultural crops has been the main area of research the past decades. Growers and sellers have been seeking for best possible product quality and highest possible profits. However, problems in quality after pathogen infections at some point of production, or during storage or transportation eventually result in economic losses (van Meeteren, 2009). *Botrytis cinerea* is the single-most important pathogen infecting ornamental plants and cut flowers postharvest and substantially reduce growers' and sellers' income by increasing product rejections.

B. cinerea Pers. is a common fungal pathogen that infects glasshouse-grown ornamental crops under cool and humid conditions with latent symptoms, which develop during storage or transportation (Elad, 1988). Growers and sellers around the world are deeply concerned by such infection problems. In Europe, for instance, large quantities of *B. cinerea*-infected cut freesias from The Netherlands are rejected in the UK by wholesalers and retailers at certain times of the year (Darras et al., 2004). These rejections result in immediate economic losses and make cooperation between growers and importers problematical. The problem is equally substantial for roses (Elad, 1988; Elad et al., 1993), gerberas (Salinas & Verhoeff, 1995) and Geraldton waxflowers (Joyce, 1993), although species such as chrysanthemum (Dirkse, 1982), narcissus (O'Neill et al., 2004), lisianthus (Wegulo & Vilchez, 2007), dianthus, ranunculus and cyclamen (Seglie et al., 2009) eventually suffer infections by *B. cinerea*, but to a lesser extent.

Infections by *B. cinerea* are usually managed by conventional fungicides applied protectively at certain times of the year and especially during autumn and spring when most infections occur. However, extensive use of fungicides such as dicarboximides, has led to the

appearance of resistant *B. cinerea* strains (Pappas, 1997). Alternative methods to control *B. cinerea* disease (i.e. grey mold) within the concept of integrated disease management (IDM) programs are sought by growers and help overcome resistance by the pathogen.

Elicitor-based disease management constitutes an attractive socio-environmentally sound strategy (Joyce & Johnson, 1999). Known activators of plant defence reactions, such as 2,6-dichloroisonicotinic acid (INA), salicylic acid (SA), 3-aminobutyric acid (BABA), Acibenzolar-S-methyl [ASM; benzo(1,2,3)thiadiazole-7-carbothioic acid S-methyl ester; benzothiadiazole or BTH; CGA 245704] and methyl jasmonate (MeJA), have been shown to enhance natural defence mechanisms or induce systemic defence responses such as SAR or ISR in plants, thereby providing prospects for IDM (Terry & Joyce, 2004a).

1.1 *Botrytis cinerea* infecting cut flowers and ornamental pot plants

Botrytis cinerea Pers. belongs to the Class Deuteromycetes and the Phylum Ascomycota. The disease caused by *B. cinerea* is called grey mold. The fungus is pathogenic to most of the cultivated ornamental pot plants and cut flowers. For example, infection of gerbera (*Gerbera jamesonii*) flowers occurs inside the glasshouse during crop cultivation, but symptoms develop after a latent period at storage or transportation following fluctuations in temperature (Salinas & Verhoeff, 1995). Favourable temperature and relative humidity (RH) for the pathogen after harvest results in rapid disease development (Salinas et al., 1989). Grey mold on gerbera and freesia flowers is observed as small necrotic, dark-brown fleck lesions 'spots'. Similar symptoms developed in the laboratory under controlled conditions following artificial inoculation of gerbera or freesia inflorescences at temperatures ranging from 4 to 25°C (Salinas & Verhoeff, 1995; Darras et al., 2006a). Infection of freesia (*Freesia hybrida*) inflorescences after artificial inoculation occurred in less than 24 h at 12°C and 80-90% RH. Even at the low temperature of 5°C, disease symptoms were evident in a saturated atmosphere (ca. 100% RH) within the first 24 h of incubation.

B. cinerea is also pathogenic to Geraldton waxflower (*Chamelaucium uncinatum*), the Australian native plant which holds a high ornamental and commercial value (Joyce, 1993; Tomas et al., 1995). Geraldton waxflower sprigs artificially inoculated with *B. cinerea* showed increased abscission of flowers from their pedicels.

B. cinerea infects rose (*Rosa hybrida*) flowers and produces necrotic spots or blister-like patches on petal surfaces (Pie & De Leeuw, 1991; Williamson et al., 1995). Infection has been described by Elad (1988) as restricted, brown, volcano-like shaped lesions. *B. cinerea* damages phylloclades of ruscus (*Ruscus aculeatus*) by causing small, dark water soaked necrotic lesions encircled by a faint halo. These lesions later become brown without growing in size (Elad et al., 1993).

Infection of lisianthus (*Eustoma russellianum*) flowers has been recently reported by Wegulo & Vilchez (2007). Significant ($P \leq 0.03$) positive correlations between stem lesion length of naturally infected plants in the glasshouse ($R = 0.74$) and stem lesion length of artificially inoculated ones ($R = 0.62$) with the disease incidence score, and with the percent of necrosis ($R = 0.71$) of detached leaves were reported (Wegulo & Vilchez, 2007). From all the 12 lisianthus cultivars tested, 'Magic Champagne' was suggested as the most resistant and proposed as ideal for commercial cultivation.

In regards to pot plants, *B. cinerea* disease symptoms on geranium (*Pelargonium zonale*) flowers has been described by Strider (1985) as flower blight, leaf blight and stem rot. Martinez et al. (2008) published a detailed report on infection of *Pelargonium x hortorum*, *Euphorbia pulcherrima*, *Lantana camara*, *Lonicera japonica*, *Hydrangea macrophylla*, and *Cyclamen persicum* by *B. cinerea*. They reported that growth of *B. cinerea* isolates in-vitro from the above mentioned ornamentals varied significantly. *B. cinerea* showed a high degree of phenotypical variability among the isolates, not only as regards to visual aspects of the colonies but also to some morphological structures such as conidium length, conidiophores, sclerotia production, and hyphae (Martinez et al., 2008). Increased susceptibility to grey mold from 10% to 80% in stems and from 3% to 14% in leaves was observed after using elevated levels of N supply (i.e. from 7.15 to 57.1 mM) for begonia plant (*Begonia x tuberhybrida* Voss) cultivation (Pitchay et al., 2007).

1.2 Review on host-pathogen interactions and on defence responses

Host-pathogen specificity involves factors that determine the virulence of the pathogen and also factors that confer resistance on the host (Lucas, 1997). Many theories have been proposed concerning mechanisms by which pathogens either achieve or fail to infect host tissue. A model concerning specific gene-for-gene interactions determining the host range of pathogens in wild plant species was first proposed by Flor (1971). In a gene-for-gene system, recognition of the pathogen by the host occurs when a resistance (R) gene of the host interacts with an avirulence (avr) gene of the pathogen (Table 1).

Virulent or avirulent Pathogen genes	Resistant or susceptible genes in the plant	
	R (resistant) dominant	r (susceptible) recessive
A (avirulent) dominant	AR (-) ^a	Ar (+)
a (virulent) recessive	aR (+)	ar (+)

^a (-) indicate incompatible interaction and, therefore, no infection. (+) indicate compatible interaction and, therefore, infection.

Table 1. Quadratic check of gene combinations and disease reaction types in host-pathogen systems in which the gene-for-gene concept for one gene operates (Lucas, 1997).

According to this model, avr gene products secreted by hyphae or located on the surface of the pathogen bind to a receptor located on the cell membranes of host's epidermal cells. Binding triggers a cascade of defence responses by the host. Every other possible match in the system could lead to infection (Table 1). Thus, a combination of a resistant host gene and a virulent pathogen leads to a compatible host-pathogen interaction. In both cases, when an avr race of the pathogen matches with a susceptible host and a virulent pathogen matches with a susceptible host, the host fails to recognize the pathogen and infection occurs (Flor, 1971).

Culture filtrates or extracts from microbial cells can act as potent inducers of plant defence responses (Chappell & Hahlbrock, 1984; Kombrink & Hahlbrock, 1986; Fritzscheier et al.,

1987; Keller et al., 1999). For instance, extracts from fungal cell walls when applied to plant tissue induced the synthesis and accumulation of phytoalexins (Yoshikawa et al., 1993). Active components in such chemical, biological and physical extracts are referred to as elicitors. This term is now generally used to denote agents, which induce plant defence responses, including accumulation of PR-proteins, cell wall structural (strengthening) changes, and hypersensitive cell death (Kombrink & Hahlbrock, 1986).

1.2.1 Rapid defence responses

The first step in the rapid defence responses by plants is recognition of the infection attempt by the pathogen. Pathogen recognition results in a signalling cascade to neighbouring cells and in initial molecular defence responses (Kombrink & Somssich, 1995). Examples of elicitor-active components produced by pathogenic fungi include the β -glucan elicitor and the 42 kDa glycoprotein derived from the fungus *Phytophthora megasperma*, the oligo-1,4- α -galacturonides from *Cladosporium fulvum* and *Rhynchosporium secalis*, and the harpin protein from *Erwinia amylovora*. These compounds activate defence responses when they bind to host receptors during incompatible host-pathogen interactions (Ebel & Cosio, 1994; Kombrink & Somssich, 1995).

In parsley cells, the existence of a receptor was proposed by Ebel & Cosio (1994). The intracellular changes were subsequent signals activated by the receptor and transported to host plasma membrane. Changes in H^+ , K^+ , Cl^- and Ca^{2+} fluxes across the plasma membrane and H_2O_2 increase within 2-5 min can occur (Nurnberger et al., 1994; Nurnberger & Scheel, 2001).

The activity of active oxygen species (e.g. O^- , H_2O_2) and the rapidity of their production after invasion characterize the rapid defence response of the host (Dixon et al., 1994; Ebel & Cosio, 1994; Bolwell, 1999). These toxic active oxygen species cause host cell death at the infection site (Kombrink & Somssich, 1995). It has been suggested that reactive oxygen species (ROS) could have a dual function in disease resistance (Kombrink & Schmelzer, 2001). Firstly, ROS participate directly in cell death during HR and, thereby, results in direct pathogen inhibition. Secondly, ROS have a role in signal diffusion for cellular protectant induction and associated defence responses in neighbouring cells (Kombrink & Schmelzer, 2001).

The HR is part of the initial plant defence responses and involves localized cell death at the infection site (Kombrink & Schmelzer, 2001). Thus, the HR is a result of host recognition of infection attempts made by a pathogenic bacterium or fungus. Specific elicitor-molecules comprise signals, which induce these initial defence responses. When pathogenic bacteria are injected inside a non-host plant under artificial conditions they are killed by the HR as a result of being surrounded by dead cells. The HR may occur when either virulent strains of bacteria are injected inside a resistant host or avirulent strains of bacteria are injected inside a susceptible host (Agrios, 1997). HR associated isolation of the pathogen inside necrotic cells causes the pathogen loses its ability to take-up nutrients and grow into adjacent healthy cells (Kombrink & Schmelzer, 2001).

Elicitors which do not cause an HR can also activate defence-related compounds (Schroder et al., 1992; Atkinson, 1993; Kuć 1995; Kombrink & Schmelzer, 2001). Activation of these compounds can be similar for both compatible and incompatible host-pathogen interactions (Schroder et al., 1992; Kombrink and Schmelzer, 2001). However, only with compatible

interactions does the pathogen infect and colonize the host. Accumulation of phytoalexins can occur as part of the HR (Dixon et al., 1994). However, it is not clear whether the HR triggers the production of phytoalexins and other antimicrobial compounds or if their accumulation is a direct result of elicitation (Kombrink & Somssich, 1995).

1.2.2 Local acquired resistance

Phytoalexins are low molecular weight antimicrobial compounds produced de-novo by some plants. They accumulate during infection by pathogens or after injury or stress (Ebel, 1986; Isaac, 1992; Kuć, 1995). Accumulation of phytoalexins is mainly observed when fungi, rather than bacteria, viruses or nematodes, try to infect the plant. Accumulation is a result of specific elicitors released either by the fungal cell walls or by the plant cell walls (Ebel, 1986). Elicitors of phytoalexins include a large number of compounds including inorganic salts (Perrin & Cruickshank, 1965), oligoglucans (Sharp et al., 1984), ethylene (Chalutz & Stahmann, 1969), fatty acids (Bostock et al., 1981), and chitosan oligomers (Kendra & Hadwiger, 1984). Over 200 compounds, microorganisms and physiological stresses have been reported to elicit pisatin in pea, phaseollin and kievitone in green bean and glyceollin in soybean (Kuć, 1991).

Most phytoalexins have been isolated from dicot plants, but they are also present in monocots including rice, corn, sorghum, wheat, barley and onions (Kuć, 1995). There is no published work on phytoalexins in cut flower species. Phytoalexins have been found in almost every part of the plant including roots, stems, leaves and fruits (Kuć, 1995). Such plant species produces a characteristic set of phytoalexins derived from secondary metabolism, in most cases via the phenylpropanoid pathway (Ebel, 1986; Kombrink & Somssich, 1995; Kuć, 1995). Phytoalexins belong to a number of key chemical groups including phenolics (e.g. flavonoids and coumarins), polyacetylenes, isoprenes, terpenoids and steroids (Ebel, 1986). They are produced by both resistant and susceptible tissues and resistance appears to be related with the total phytoalexin concentration (Kuć, 1995). Phytoalexins can affect fungal growth by inhibiting germ tube elongation and colony growth (Elad, 1997). The main effect of phytoalexins on fungi is via their cell membranes. Direct contact of phytoalexins with fungal cell walls resulted in fungal plasma membrane disruption and loss of the ultrastructural integrity (Elad, 1997). In compatible interactions, the pathogen apparently tolerates accumulated phytoalexins, detoxifies them, suppresses phytoalexin accumulation and/or avoids eliciting phytoalexin production (Kuć, 1995). Overcoming phytoalexin accumulation is attributed to either suppressor molecules released by the pathogen (i.e. low molecular weight polysaccharides or glycopeptides) or suppression of the intensity and timing of signal genes that could trigger phytoalexin accumulation (Kuć, 1995).

Pathogenesis related proteins (PR-proteins) accumulate either in extracellular space or the vacuole after various types of plant stress, including pathogen infection (Stermer, 1995; Sticher et al., 1997). PR-proteins accumulate at the site of infection as well as in uninfected tissues (Van Loon & Gerritsen, 1989; Ryals et al., 1996). Although healthy plants may contain traces of PR-proteins, the transcription of genes encoding PR-proteins is up-regulated following pathogen attack, elicitor treatment, wounding or stress (Stermer, 1995; Sticher et al., 1997; Van Loon, 1997). Signal compounds responsible for initiating PR-protein production include salicylic acid, ethylene, the enzyme xylanase, the polypeptide systemin

and jasmonic acid (Agrios, 1997). The importance of PR-proteins lies in their range of defence activities (Van Loon et al., 1994). A number of PR-proteins release molecules that may act as elicitors (Keen & Yoshikawa, 1983). PR-proteins accumulation has been observed in monocots as well as in dicots (Redolfi, 1983). However, there is no published work on PR-proteins induced in flower species. Eleven families of PR-proteins have been recognized so far (Van Loon et al., 1994). Some inhibit pathogen development during microbial infection by inhibiting fungal spore production and germination. Others are associated with strengthening of the host cell wall via its outgrowths and papillae (Agrios, 1997). Both β -1,3-glucanases and chitinases, PR-2 and PR-3, respectively, are known to have direct antifungal activity (Mauch et al., 1988; Van Loon, 1997). However, many pathogens have evolved mechanisms to reduce the antifungal impact of PR-proteins (Van Loon, 1997). For example, many chitin-containing fungi are not inhibited by host-produced chitinases.

Plant secondary metabolites are divided into the three main categories of terpenes, phenolic compounds and nitrogen containing secondary metabolites (i.e. alkaloids) (Taiz & Zeiger, 1998). All secondary metabolites are produced through one of the major mevalonic, malonic or shikimic pathways (Taiz & Zeiger, 1998). Phenylalanine is a common amino acid produced via the shikimic pathway (Hahlbrock & Scheel, 1989). The most abundant classes of secondary phenolic compounds in plants are derived from phenylalanine via elimination of an ammonia molecule to form cinnamic acid. This reaction is catalyzed by phenylalanine ammonia lyase (PAL), the key enzyme of phenylpropanoid metabolism (Hahlbrock & Scheel, 1989). Derivatives of phenylpropanoid pathway include low-molecular-weight flower pigments, antibiotic phytoalexins, UV-protectants, insect repellents, and signal molecules in plant-microbe interactions (Hahlbrock & Scheel, 1989; Kombrink & Somssich, 1995).

The main phenylpropanoid pathway branches leading to formation of flavonoids, isoflavonoids, coumarins, soluble esters, wall bound phenolics, lignin and suberin. This diverse spectrum of compounds has a wide range of properties (Hahlbrock & Scheel, 1989). For example, the lignin pathway is an important phenylpropanoid pathway branch that produces precursors for lignin deposition (Grisebach, 1981). Various enzymes implicated in the biosynthesis of lignin appeared to be induced in plants in response to infection or elicitor treatment (Matern & Kneusel, 1988). However, not all studies show a role of lignin and cell lignification in disease inhibition (Garrod et al., 1982). Furanocoumarins derived from the furanocoumarin pathway in parsley are considered potent phytoalexins (Beier & Oertli, 1983). Flavonoid and furanocoumarin production as a response to UV light or fungal elicitor treatment respectively was associated with up-regulation of PAL, 4-coumarate: CoA-ligase (4CL) and chalcone synthase (CHS). Up-regulation was based on rapid changes in amounts and activities of the corresponding mRNAs (Chappell & Hahlbrock, 1984).

After pathogen recognition by the host, a cascade of early responses is induced including ion fluxes, phosphorylation events, and generation of active oxygen species (Kombrink & Somssich, 1995). SA acts as a secondary signal molecule and its levels increase during the defence induction process. Thus, SA is required for initiation of synthesis of various defence-related proteins such as the PR-proteins (Van Loon, 1997; Metraux, 2001). SA accumulation endogenously in tobacco and cucumber plants lead to the HR and the SAR responses. However, SA is not necessarily the translocated signal (elicitor) for the onset of SAR. Rather, SA exerts an effect locally (Vernooij et al., 1994; Ryals et al., 1996). Nonetheless,

SA is still required for SAR expression (Van Loon, 1997). The importance of SA in the onset of SAR was determined using transgenic tobacco and Arabidopsis plants engineered to over-express SA-hydroxylase. Transformed plants with the naphthalene hydroxylase G (NahG) gene produced low levels of SA and SAR expression was blocked.

SA is produced from phenylalanine via coumaric and benzoic acid (Mauch-Mani and Slusarenko, 1996; Ryals et al., 1996; Sticher et al., 1997). Biosynthesis of SA starts with the conversion of phenylalanine to trans-cinnamic acid (Sticher et al., 1997). From trans-cinnamic acid, either benzoic acid (BA) or ortho-coumaric acid are produced. Both compounds are SA precursors (Sticher et al., 1997). Pallas et al. (1996) showed that tobacco plants epigenetically suppressed in PAL expression produced a much lower concentration of SA and other phenylpropanoid derivatives when artificially inoculated with tobacco mosaic virus (TMV). This was seen, firstly, due to the lack of resistance to TMV upon secondary infection, and, secondly, to the absence of PR protein induction in systemic leaves (Pallas et al., 1996).

Jasmonic acid (JA) and its methyl ester (MeJA) are derived from linolenic acid. They are cyclopentanone-based compounds that occur naturally in many plant species (Sembdner & Parthier, 1993; Creelman & Mullet, 1997). Linolenic acid levels or its availability could determine JA biosynthetic rate (Farmer & Ryan, 1992; Conconi et al., 1996). The level of JA in plants varies as a function of tissue and cell type, developmental stage, and in response to various environmental stimuli (Creelman & Mullet, 1997). For example, in soybean seedlings, JA levels are higher in the hypocotyls hook (a zone of cell division) and young plumules as compared to the zone of cell elongation and more mature regions of the stem, older leaves and roots (Creelman & Mullet, 1997). High JA levels are also found in flowers and pericarp tissues of developing reproductive structures (Creelman & Mullet, 1997). Jasmonates are wide spread in Angiosperms, Gymnosperms and algae (Parthier, 1991). They can mediate gene expression in response to various environmental and developmental processes (Wasternack & Parthier, 1997). These processes include wounding (Schaller & Ryan, 1995), pathogen attack (Epple et al., 1997), fungal elicitation (Nojiri et al., 1996), touch (Sharkey, 1996), nitrogen storage (Staswick, 1990), and cell wall strengthening (Creelman et al., 1992). Wounding of tomato leaves produced an 18-amino acid polypeptide called systemin, the first polypeptide hormone discovered in plants so far (Pearce et al., 1991). Systemin was released from damaged cells into the apoplast and transported out of the wounded leaf via the phloem (Schaller & Ryan, 1995) (Fig. 1).

Upon herbivore wounding, a systemic signal is delivered from systemin and results in an ABA-dependent rise of linoleic acid. Systemin was believed to bind to the plasma membrane of target cells and thereby initiate JA biosynthesis (Schaller & Ryan, 1995). JA accumulation can also be induced by oligosaccharides derived from plant cell walls and by elicitors, such as chitosans derived from fungal cell walls (Gundlach et al., 1992; Doares et al., 1995; Nojiri et al., 1996). JA also activates gene expression encoding proteinase inhibitors (Creelman & Mullet, 1997). Proteinase inhibitors are known antidigestive proteins that block the action of herbivore proteolytic enzymes (Creelman & Mullet, 1997). Thereby, proteinase inhibitors help the host to avoid consumption by herbivores. Proteinase inhibitors were accumulated in tomato plants after wounding (O'Donnell et al., 1996) and after irradiation with UV-C (Conconi et al. 1996). In response to wounding, ethylene and JA act together to regulate gene expression of proteinase inhibitors (O'Donnell et al., 1996). Exposing tomato

leaves to increasing doses of 254 nm UV-C resulted in increased proteinase inhibitors gene expression. Expression of proteinase inhibitors in wounded (Doares et al., 1995; O'Donnell et al., 1996) or UV-C treated (Conconi et al., 1996) tomato leaves was markedly reduced upon treatment with SA. From linoleic acid, jasmonic acid is produced. Ethylene is required in the jasmonic-signalling cascade (O'Donnell et al., 1995).




Fig. 1. The octadecanoid-signalling pathway for defence gene expression in tomato (Schaller and Ryan, 1995).

1.2.3 Systemic defence responses (i.e. SAR, ISR) and signalling pathways

SAR is activated following induction of local acquired resistance (LAR). SAR is potentially induced after the HR and after challenge with virulent strains of a pathogen or elicitor treatment. It develops systemically in distant parts of the infected plant (Lawton et al., 1996; Ryals et al., 1996; Mettraux, 2001). SAR protects plants from a broad range of potential pathogens (Kessmann et al., 1994).

Specific genes induced in different plant species during SAR have been called SAR-genes (Kessmann et al., 1994; Stermer, 1995; Ryals et al., 1996; Sticher et al., 1997). Most of SAR-genes encode PR-proteins such as those accumulated after inoculation of tobacco with TMV (Ward et al., 1991). These include PR-1 (PR-1a, PR-1b, PR-1c), β -1,3-glucanases (PR-2a, PR-2b, PR-2c), chitinases (PR-3a, PR-3b), hevein-like proteins (PR-4a, PR-4b), thaumatin like proteins (PR-5a, PR-5b), acidic and basic isoforms of class III chitinase, an extracellular β -1,3-glucanase and the basic isoform of PR-1 (Ward et al., 1991). SAR and SAR-gene activation has been observed in various dicots (Kessmann et al., 1994; Ryals et al., 1996). SAR activation involves species specificity (Ryals et al., 1992). For example, acidic PR-1 is only weakly expressed in cucumber. In contrast, acidic PR-1 is the main protein accumulating in tobacco and Arabidopsis. A number of homologous SAR-genes have been identified in monocots. Homologs of the PR-1 family were found in maize and barley and other PR-proteins in maize (Nasser et al., 1988). Gorlach et al. (1996) isolated a group of wheat genes (WCI or wheat chemically induced) induced after chemical treatment with potent SAR inducers. WCI genes seemed to act in a similar manner to SAR-genes in dicots after chemical treatment with plant activators (Gorlach et al., 1996).

Recent research has revealed that JA and ethylene play key roles in signal transduction pathways associated with plant defence responses (Pieterse and van Loon, 1999; Thomma et al., 2000). Inoculation with a necrotizing pathogen resulted predominantly in activation of the SA-dependent SAR response. This response leads to the accumulation of salicylic acid inducible PR-proteins and the expression of SAR (Ryals et al., 1996; Pieterse & van Loon, 1999) (Fig. 2, pathway 2). JA and ethylene inducible defence responses are induced by non-necrotizing rhizobacteria and lead to the ISR phenomenon (Pieterse et al., 1996; Pieterse et al., 1998) (Fig. 2, pathway 1). Both pathways 1 and 2 are regulated in Arabidopsis plants carrying the NPR1 gene.

Depending on the invading pathogen, the composition of defence compounds produced after infection can vary between SA- and JA/ethylene-inducible pathways (Fig. 2, pathways 2 and 3) (Ryals et al., 1996; Epple et al., 1997; Dong, 1998).

Wounding can also result in JA and ethylene inducible defence response activation (Fig. 2, pathway 4) (O'Donnell et al., 1996; Wasternack & Parthier, 1997). However, resultant products of the wounding pathway differ from those induced upon pathogen infection (O'Donnell et al., 1996; Rojo et al., 1999). A second distinct wound-signalling pathway leading to wound responsive (WR) gene expression has been found in Arabidopsis plants (Titarenko et al., 1997; Rojo et al., 1998) (Fig. 2, pathway 6). Upon wounding, Arabidopsis plants carrying the *coi1* (JA-insensitive) mutant gene expressed the wound responsive genes choline kinase (CK) and wound responsive (WR3) indicating that the induced pathway was totally independent of JA. UV irradiation of tomato leaves also resulted in induction of the same defensive genes normally activated through the octadecanoid pathway after wounding (Conconi et al., 1996). This response is blocked after SA treatment, confirming the

antagonistic regulation of the two distinct pathways (Pena-Cortes et al., 1993; Lawton et al., 1995; Xu et al., 1994; Doares et al., 1995; O'Donnell et al., 1996; Niki et al., 1998; Gupta et al., 2000; Rao et al., 2000).

In the rhizobacteria-mediated induced systemic resistance (ISR) pathway, components from the JA/ethylene response acted in sequence in activating a systemic resistance response that, like pathogen induced SAR, was dependent on the regulatory protein NPR1 (Pieterse & van Loon, 1999). The ISR pathway shares signalling events with pathways initiated upon pathogen infection, but is not associated with the activation of genes encoding plant defensins, thionins or PR-proteins (Pieterse & van Loon, 1999) (Fig. 2, pathway 3). This observation indicates that ISR inducing rhizobacteria, such as *P. fluorescens* strain WCS417r, trigger a novel signalling pathway leading to the production of so far unidentified defense compounds (Pieterse et al., 1996; Pieterse et al., 1998). Protection of NahG Arabidopsis plants by gaseous MeJA suggested that induction of a SA non-dependent systemic pathway was regulated by JA (Thomma et al., 2000) (Fig. 2, pathway 3). Protection was provided against two necrotrophic fungi, *A. brassicicola* and *B. cinerea*.




Fig. 2. Model showing systemic signalling pathways that can be induced in plants by non-pathogenic rhizobacteria, pathogen infection and wounding, such as caused by foraging insects. 1: ISR is induced in NPR1 Arabidopsis plants as a result of JA and ethylene responses. 2: SAR is induced in NPR1 Arabidopsis plants after necrosis by pathogenic fungi, bacteria or virus. 3: JA/ethylene pathway is up-regulated after fungal infection. JA/ethylene expression leads to genes encoding plant defensins, thionins, proteinase inhibitors and SA-independent PR-proteins. 4 and 6: A number of genes are regulated after mechanical wounding. JA and ethylene levels rise after mechanical wounding. 5: Cross-talk between SA- and JA- dependent pathways exist. Adopted from Pieterse & van Loon (1999).

1.3 Elicitation of defence responses with chemical activators

Disease management in the past has been achieved by various methods including resistant cultivars, biological control, crop rotation, tillage, and chemical pesticides (Kessmann et al., 1994). Recently, the use of abiotic and/or biotic agents, as well as, synthetic compounds that induce host immune systems have offered a new prospect for disease management.

A chemical is generally characterized as a plant activator when it induces natural and/or systemic defence responses, activate gene expression and provide protection on the same spectrum of diseases exerted by a wild type host (Kessmann et al., 1996; Ruess et al., 1996). Plant activators, normally, do not exert direct antimicrobial activity against pathogens when used for disease control, but rather work through their mutagenic elicitation effect and help eliminate the risk of the development of resistant strains by the pathogen.

1.3.1 Acibenzolar-S-methyl (ASM; benzo[1,2,3]thiadiazole-7-carbothioic acid S-methyl ester; CGA 245704; benzothiadiazole or BTH)

The efficacy of ASM has been tested in field, glasshouse and pot trials (Table 2).

Host	Pathogen	Induced response	ASM concentration	Reference
Apple seedlings cv. Golden Delicious	<i>Erwinia amylovora</i>	β -1,3-glucanases, peroxidases	0.1-0.2 g AI L ⁻¹	Brisset et al., 2000
Cauliflower (<i>Brassica oleracea</i>)	<i>Peronospora parasitica</i>	ns	0.0015-0.075 g AI L ⁻¹	Godard et al., 1999
Cereals, tobacco	<i>Erysiphe graminis</i> , <i>Septoria</i> spp., <i>Puccinia</i> spp., <i>Peronospora hyoscyami</i> f. sp. <i>tabacina</i>	ns	12-30 g AI ha ⁻¹	Ruess et al., 1996
Cucumber (<i>Cucumis sativus</i> L.)	<i>Cladosporium cucumerinum</i>	Acidic peroxidase, class III chitinase and β -1,3-glucanase	32.4 g AI L ⁻¹	Narusaka et al., 1999
Cucumber (<i>Cucumis sativus</i> L.) and Japanese pear (<i>Pyrus pyrifolia</i> Nakai var. <i>culta</i> Nakai)	Many pathogens	ns	0.05-0.1 g AI L ⁻¹	Ishii et al., 1999
Cauliflower (<i>Brassica oleracea</i>)	<i>P. parasitica</i>	β -1,3-glucanase. PR-1 and PR-5	0.05 g AI L ⁻¹	Ziadi et al., 2001
Grapevine cv Merlot	<i>B. cinerea</i>	na	0.3 mM	Iriti et al., 2004
Melons cv. Early Yellow Hami	<i>Fusarium</i> spp., <i>Alternaria</i> spp., <i>Rhizopus</i> spp., <i>Trichothecium</i> sp.	ns	0.025 or 0.05 g AI L ⁻¹	Huang et al., 2000

Host	Pathogen	Induced response	ASM concentration	Reference
Melon fruit	<i>Fusarium pallidoroseum</i>	ascorbate peroxidase, guaiacol peroxidase, PAL, β -1,3-glucanase		Gondim et al., 2008
Parsley cells (<i>Petroselinum crispum</i> L.)	With or without elicitor (Pmg)	PAL, coumarins	0.32-6.48 g AI L ⁻¹	Katz et al., 1998
Pepper (<i>Capsicum annuum</i> L.)	<i>Xanthomonas campestris</i> pv. <i>vesicatoria</i>	ns	1.25-5 g AI L ⁻¹	Buonaurio et al., 2002
Soybean seedlings	<i>Sclerotinia sclerotiorum</i>	ns	0.035-0.375 g AI L ⁻¹	Dann et al., 1998
Strawberry plants cvs. Elsanta and Andana	<i>B. cinerea</i>	ns	0.25-2 g AI L ⁻¹	Terry and Joyce, 2000
Strawberry	Microbial populations	chitinase and β -1,3-glucanase	0.05-0.5 g.L ⁻¹	Cao et al., 2010
Tobacco plants cv. Kutsaga Mammoth 10	<i>Pseudomonas syringae</i> pv. <i>tabaci</i> , <i>Thanatephorus cucumeris</i> , <i>Cercospora nicotianae</i>	ns	0.05-30 g AI L ⁻¹	Cole, 1999
Tomato plants (<i>Lycopersicon esculentum</i>)	<i>Fusarium oxysporum</i> f.sp. <i>radicis-lycopersici</i>	Callose enriched wall appositions phenolic compounds	97.2 g AI L ⁻¹	Benhamou & Belanger, 1998
Tomato plants cv. Vollendung	<i>Cucumber mosaic virus</i> (CMV)	ns	0.1 mM	Anfoka, 2000
Wheat	<i>Erysiphe graminis</i> f.sp. <i>tritici</i>	WCI genes (1-5)	0.3 mM	Gorlach et al., 1996

Table 2. Effects of ASM on different host-pathogen interactions (ns: not shown, na: not applicable).

Although, most of ASM application were carried out preharvest, there is number of published research on ASM postharvest applications (i.e. Cao et al., 2010). Additionally, a considerable work on postharvest application of ASM on ornamentals has been published the recent years (i.e. Darras et al., 2007). ASM was introduced as a potent inducer of SAR and treated plants were resistant to the same spectrum of diseases as plants activated naturally (Kessmann et al., 1996; Friedrich et al., 1996). Although ASM and its metabolites exhibited no direct antimicrobial activity towards plant pathogens tested, they induced the same biochemical processes in the plant as those observed after natural activation of SAR (Friedrich et al., 1996; Lawton et al., 1996). The compound, which was inactive in plants that do not express the SAR-signaling pathway, required a lag time of approximately 30 days between application and protection (Lawton et al., 1996).

1.3.2 Jasmonates (plant hormones produced through the octadecanoid pathway)

The efficacy of jasmonates has been tested in field, glasshouse and pot trials (Table 3).

Host	Pathogen	Induced response	MeJA concentration	Reference
Arabidopsis (<i>Arabidopsis thaliana</i>)	<i>B. cinerea</i> , <i>A. brassicicola</i> , <i>Plectosphaerella cucumerina</i>	ns	0.5-50 μM and 0.001-1 $\mu\text{L L}^{-1}$ air	Thomma et al., 2000
Arabidopsis (<i>Arabidopsis thaliana</i>)	<i>A. brassicicola</i>	PDF1.2	45 μM	Penninckx et al., 1996
Grapefruit (<i>Citrus paradisi</i>) var. 'Marsh Seedless	<i>Penicillium digitatum</i>	ns	1-50 μM	Droby et al., 1999
Large number of species	na	PPO	na	Constabel and Ryan, 1998
Loquat fruit	<i>Colletotrichum acutatum</i>	chitinase and β -1,3-glucanase	10 $\mu\text{mol L}^{-1}$	Cao et al., 2008
Potato plants (<i>Solanum tuberosum</i>)	<i>Phytophthora infestans</i>	phytoalexins	1-10 μM	Il'inskaya et al., 1996
Sweet cherry	<i>Monilinia fructicola</i>	PAL, β -1,3-glucanase β -glucuronidase (GUS), osmotin protein	0.2 mM	Yao & Tian, 2005
Tobacco cell cultures	na	β -glucuronidase (GUS), osmotin protein	0.045-4550 μM	Xu et al., 1994
Tobacco cv. Xanthi-nc	<i>Phytophthora parasitica</i> var. <i>nicotianae</i> , <i>Cercospora nicotianae</i> , TMV	β -glucuronidase (GUS)	45 μM	Mitter et al., 1998
Tomato plants (<i>Lycopersicon esculentum</i>)	<i>Helicoverpa zea</i> , <i>Spodoptera exigua</i>	PPO, POD, LOX and PIs	0.1-10 mM	Thaler et al., 1996
Tomato plants (<i>Lycopersicon esculentum</i>)	<i>Spodoptera exigua</i> , <i>Pseudomonas syringae</i> pv. <i>tomato</i>	PPO	1 mM	Thaler et al., 1999

Table 3. Effects of MeJA on different host-pathogen interactions. (ns: not shown, na: not applicable).

Although, firstly tested preharvest, JA or MeJA has been extensively used postharvest at different hosts (i.e. fruits, vegetables, cut flowers), application modes (i.e. spray, pulse, gas) and incubation environments (i.e. storage or ambient temperatures). For example, JA and MeJA were tested on grapefruit for suppressing postharvest green mold decay [*Penicillium digitatum* (Pers.:Fr.) Sacc.] (Droby et al., 1999). Studies showed that 50 μM and 1 μM MeJA concentrations were effective against the disease and that the reduction in the decay was the same at incubation temperatures of 2 or 20°C. Moreover, as the in-vitro tests showed no direct antifungal activity of JA and MeJA, it was suggested that the disease suppression was achieved via natural resistance induction (Droby et al., 1999). Treatment of Arabidopsis plants with MeJA reduced *A. brassicicola*, *B. cinerea* and *Plectosphaerella cucumerina* disease development (Thomma et al., 2000). Application of gaseous MeJA to plants resulted in a greater disease reduction compared to that on plants sprayed with MeJA or treated with INA. Gaseous MeJA protected SA-degrading transformant NahG plants, suggesting that gaseous MeJA induced a non-SA dependent systemic response (Thomma et al., 2000). Combination of ASM and JA was tested against bacterial and insect attack on field grown tomato plants (Thaler et al., 1999). Two signaling pathways, one involving SA and another involving JA were proposed to provide resistance against pathogens and insect herbivores, respectively (Thaler et al., 1999).

1.4 Elicitation of defence responses in floriculture

The efficacy of ASM and MeJA on ornamental pot plants and on cut flowers has been tested pre- and postharvest, respectively (Table 4). Most of such tests were carried out in the very recent years and still increasing. For example, pre- and postharvest treatments with MeJA or ASM on cut flowers conferred a variable measure of protection against postharvest infections by *B. cinerea* (Dinh et al., 2007).

JA and MeJA provided systemic protection to various rose cultivars (e.g. Mercedes, Europa, Lambada, Frisco, Sacha and Eskimo) against *B. cinerea* (Meir et al., 1998). MeJA applied as postharvest pulse, significantly reduced *B. cinerea* lesion size on detached rose petals. In the same study, MeJA at concentrations of 100-400 μM showed in-vitro antifungal activity on *B. cinerea* spore germination and germ-tube elongation. Similarly, a postharvest pulse, spray, or vapour treatment with MeJA 200 μM , 600 μM or 1 $\mu\text{L L}^{-1}$, respectively, significantly reduced petal specking by *B. cinerea* on cut inflorescences of *Freesia hybrida* 'Cote d'Azur' (Darras et al., 2005; 2007). Moreover, 1-100 $\mu\text{L L}^{-1}$ MeJA postharvest vapour treatment reduced *B. cinerea* development on cut Geraldton waxflower 'Purple Pride' and 'Mullering Brook' sprigs (Eyre et al., 2006). Application of gaseous MeJA to fresh cut peonies resulted in the lowest disease severity and in an improvement of vase life compared to the untreated controls (Gast, 2001).

MeJA and ASM, applied preharvest had variable responses against postharvest infection by *B. cinerea*. ASM was not as effective as MeJA in suppressing the development of postharvest *B. cinerea* disease for glasshouse grown freesias (Darras et al., 2006b). Dinh et al. (2007) reported that multiple sprays of $\leq 1000 \mu\text{M}$ MeJA to field grown plants significantly reduced *B. cinerea* on Geraldton waxflower 'My Sweet Sixteen' cut sprigs, that were un-inoculated or artificially inoculated with *B. cinerea* (Dinh et al., 2007).

Host	Elicitor	Target pathogen	Application method and timing	Reference
a. Cut flowers				
Rose (<i>Rosa hybrida</i>)	ASM	<i>Diplocarpon rosae</i>	Spray - preharvest	Suo & Leung, 2002
	MeJA	<i>B. cinerea</i>	Pulse, spray - postharvest	Meir et al., 1998; 2005
Gerbera (<i>Gerbera jamesonii</i>)	UV-C	<i>B. cinerea</i>	Postharvest	Darras et al., 2012
Freesia (<i>Freesia hybrida</i>)	MeJA & ASM	<i>B. cinerea</i>	Spray - preharvest	Darras et al., 2006b
	MeJA	<i>B. cinerea</i>	Pulse, spray, gas - postharvest	Darras et al., 2005; 2007
Sunflower plants (<i>Helianthus annuus</i>)	ASM, MeJA, SA, INA	<i>B. cinerea</i>	Spray - preharvest	Dimitriev et al., 2003
Sunflower plants (<i>Helianthus annuus</i>)	ASM	<i>Plasmopara helianthi</i>	Spray - preharvest	Tosi et al., 1999
Geraldton waxflower (<i>Chamelaucium uncinatum</i>)	SA	<i>Alternaria</i> sp., <i>Epicoccum</i> sp.	Spray - preharvest	Beasley, 2001
	MeJA	<i>B. cinerea</i>	Gas - postharvest	Eyre et al., 2006
	MeJA & ASM	<i>B. cinerea</i>	Spray - pre- and postharvest	Dinh et al., 2007; 2008
Peonies (<i>Paeonia lactiflora</i>)	MeJA	<i>B. cinerea</i>	Gas - postharvest	Gast, 2001
b. Pot plants				
Cyclamen (<i>Cyclamen persicum</i>)	ASM	<i>Fusarium oxysporum</i> f. sp. <i>cyclaminis</i>	Spray - preharvest	Elmer, 2006a
Petunia (<i>Petunia hybrida</i>)		<i>Phytophthora infestans</i>	Spray - preharvest	Beckett et al., 2005
c. Landscape architecture plants				
Date palm	ASM	<i>Fusarium oxysporum</i> f. sp. <i>albedinis</i>	Injection in the trunk	Jaiti et al., 2009
d. Propagation material				
Gladiolus corms (<i>Gladiolus x hortulanus</i>)	ASM	<i>Fusarium oxysporum</i> f. sp. <i>gladioli</i>	Dip	Elmer, 2006b

Table 4. Chemical and biological elicitors tested on cut flowers and ornamental pot plants against various pathogens infecting either pre- or postharvest.

Chemical elicitors such as ASM have been applied in pot ornamentals such as petunia (Beckett et al., 2005), cyclamen (Elmer, 2006a) and in gladiolus corms (Elmer, 2006b), but effectiveness varied within the different experimental designs and conditions. In cyclamen, infection by *Fusarium oxysporum* f.sp. *cyclaminis* was reduced with increasing ASM doses (Elmer, 2006a). Additionally, the dry mass of ASM treated cyclamen plants increased with increasing ASM rates. However, as no further assays were carried out to assess possible induction of defence responses, it was not clear whether ASM reduced *F. oxysporum* f.sp. *cyclaminis* via induction of defence mechanisms or via a profound fungitoxic effect. It has been demonstrated in other research that ASM may exert direct toxic activity against *B. cinerea* (Darras et al., 2006b). In addition, ASM did not confer a significant level of protection on gladiolus corms against *F. oxysporum* f. sp. *gladioli*, and compared to conventional fungicides, although, the number of emerging flower spikes increased significantly compared to the ASM-untreated corms (Elmer, 2006b).

2. Elicitation of defence responses in cut *Freesia hybrida* flowers – A typical example

2.1 Background

Infection problems by *Botrytis cinerea* are typical to most geographical areas around the world and concern cut flower industry. Infection of cut flowers by the fungus results in visible lesions on flower petals (petal spotting or petal specking). According to Darras et al. (2004) freesia flower rejections at certain periods of the year (viz. April, May, October) lead in severe economic losses to growers, importers and sellers. Infection by *B. cinerea* of most cut flowers occurs in the glasshouse when a single conidium germinates and penetrates petal epidermal cells. A necrotic lesion appears postharvest after a brief incubation period under favourable environmental conditions (Darras et al., 2006a). Infection is difficult to control as it appears later in handling chain under various conditions during transport or storage.

In most cases, *B. cinerea* disease is controlled by conventional fungicides. However, extensive use of fungicides such as dicarboximides in the glasshouse has led to appearance of fungicide resistance (Pappas 1997). Alternative management methods within the concept of IDM can help overcome such problems.

For this reason, plant defence inducers (i.e. elicitors) such as ASM and MeJA have been tested with applications at various intervals, pre- or postharvest to activate systemic defence responses of the host (Kessmann et al., 1994; Meir et al., 1998; Thomma et al., 2000). For cut freesia flowers postharvest pulse, spray, or gaseous MeJA treatment at 200 μM , 600 μM , or 1 $\mu\text{L L}^{-1}$, respectively, significantly reduced petal specking by *B. cinerea* on cv. 'Cote d'Azur' inflorescences (Darras et al., 2005; 2007). An apparent induced defence response was recorded by both ASM and MeJA treatment. However, only MeJA conferred constant and significant disease reductions. MeJA vapour at 1 $\mu\text{L L}^{-1}$ significantly reduced lesion numbers and diameters on freesia petals by up to 56% and 50%, respectively (Darras et al., 2005).

2.2 Overview of published research and further discussion

Freesia inflorescences cv 'Cote d'Azur' gassed with 0.1 $\mu\text{L L}^{-1}$ MeJA showed significantly smaller lesions after artificial inoculation with *B. cinerea* (Fig. 3). Gaseous MeJA might have

induced a range of defence mechanisms to halt infection development. MeJA applied post-harvest as vapour at 1-100 $\mu\text{L L}^{-1}$ significantly reduced the development of *B. cinerea* on cut Geraldton waxflower 'Purple Pride' and 'Mullering Brook' sprigs (Eyre et al., 2006). In a very recent study, Darras et al. (2011) demonstrated that gaseous MeJA at 0.1 $\mu\text{L L}^{-1}$ significantly increased polyphenol oxidase (PPO) activities 24 and 36 h post-treatment. This observation suggests that MeJA-induced defence mechanisms might be associated with the production of quinones (Constabel and Ryan, 1998), which probably helped in *B. cinerea* disease reduction. The effects of PPO in *B. cinerea* disease control have been confirmed for gerbera flowers (Darras et al., 2012). A low dose of UV-C irradiation increased PPO activity and was positively correlated with the reduction of *B. cinerea* disease symptoms on the florets (Darras et al., 2012). This indicates that PPO might play an important role in *B. cinerea* disease control on cut flowers.



Fig. 3. *B. cinerea* necrotic lesions on artificially inoculated freesia cv. 'Cote d'Azur' flowers treated with 0.1 $\mu\text{L L}^{-1}$ gaseous MeJA (left) or left un-treated (control) (right) and incubated for 48 h at 20°C (Darras, 2003).

Lesion diameters on the detached freesia petals were significantly reduced with increasing MeJA spray, pulse or gaseous concentrations (Darras et al., 2007). The first published evidence of postharvest MeJA spray treatments enhancing protection of cut flowers against *B. cinerea* was the work by Meir et al. (2005) on cut roses. According to Meir et al. (2005), simultaneous MeJA pulsing and spraying under handling conditions resulted in suppression of gray mold in seven rose cultivars ('Eskimo', 'Profita', 'Tamara', 'Sun Beam', 'Pink Tango', 'Carmen', 'Golden Gate'). In an earlier study MeJA applied as a pulse variably reduced *B. cinerea* lesion numbers and diameters (Meir et al., 1998). Our findings are in agreement with those by Meir et al. (1998) that disease severity in both artificially inoculated and naturally infected rose flowers was reduced by a MeJA pulse at 0.2 mM at 20°C. On cut Geraldton waxflower 'Purple Pride' and 'Mullering Brook' sprigs, 1-100 µL L⁻¹ MeJA postharvest vapour treatment significantly reduced the development of *B. cinerea* (Eyre et al., 2006). However, it also induced flower fall incidence, which was correlated with a systemic resistance-associated up-regulation of ethylene biosynthesis.

Irrespective to the concentration tested, ASM provided no protection to artificially inoculated freesia flowers (Darras et al., 2007). However, natural infection was significantly ($P < 0.05$) reduced after ASM treatment during storage at 5 and at 12°C. On the contrary, postharvest treatments of strawberry cv. Camarosa fruit with ASM failed to reduce natural infection by *B. cinerea* at 5°C (Terry & Joyce, 2004b). Generally, ASM tended to provide protection on freesia flowers at lower incubation temperatures (Darras et al., 2007). However, it was not clear whether such disease reductions were the result of the induction of host's defence responses or a direct fungitoxic activity measured in the same study. Likewise, Terry & Joyce (2000) showed that ASM reduced in-vitro *B. cinerea* mycelial growth on ASM-amended agar. It is possible that the limited disease control on freesia flowers at 5°C was due to direct toxic effect of ASM rather than via SAR induction.

Elicitation of defence responses in cut flowers is an interesting prospect for *B. cinerea* disease control especially as it may offer alternatives to fungicide application. In series of postharvest experiments with freesia inflorescences the potential to induce natural defence mechanisms or directly controlling *B. cinerea* disease by application of biological and chemical elicitors was investigated. Postharvest treatments with ASM, MeJA or UV-C irradiation markedly suppressed *B. cinerea* specking on freesia petals by reducing disease severity, lesion numbers and lesion diameters. However, attempts to further minimise disease damage caused by *B. cinerea* using combined treatments with different plant activators (i.e. both ASM and MeJA), were not successful (Darras et al., 2011).

In summary, ASM was the least effective in reducing *B. cinerea* specking on cut freesia flowers (Fig. 4). In addition, it remained unclear as to whether or not SAR was induced. In contrast, gaseous MeJA reduced disease severity most probably by inducing JA-dependant biochemical responses. These contrasting results tend to concur with observations by Pieterse & van Loon (1999) and Thomma et al. (2001) that SA- versus JA-dependent pathways are effective against different pathogens. The results of a most recent paper (i.e. Darras et al., 2011) suggested that, SA-dependant pathway and consequently the SAR response was not effective in freesia flowers against *B. cinerea* infection. In contrast, the JA-dependant pathway was apparently induced and suppressive of *B. cinerea* infection (Darras et al., 2011).




Fig. 4. Ranking, in terms of relative efficacy, of postharvest biological (i.e. UV-C) and chemical (i.e. ASM, MeJA) elicitors tested on cut freesia inflorescences to control *B. cinerea* infection starting with the most effective (Darras, 2003).

3. Conclusions and recommendations for future research

Management of postharvest infection of cut freesia flowers by *B. cinerea*, was, in most cases, successful. ASM was somewhat effective compared to untreated controls mostly when applied preharvest at 1.43 μM . In-vitro studies showed direct antifungal activity of ASM against *B. cinerea* colony growth and conidial germination. Inconsistency of ASM applied pre- or postharvest may be explained by: 1) variability of environmental conditions in the glasshouse, which may affected defence enhancement (Herms & Mattson, 1992; Terry, 2002); and, 2) infection by *B. cinerea* might not necessarily be sensitive to induced SAR responses, and thus ASM treatments may not correspond to *B. cinerea* disease suppression (Thomma et al., 1998; Govrin & Levine, 2002). Friedrich et al. (1996) reported that ASM failed to control *B. cinerea* in tobacco, but was effective against other pathogens. The apparent inability of ASM to control *B. cinerea* was seemingly supported by the observation that PAL activity in ASM treated freesia inflorescences was not higher compared to the untreated controls. Therefore, ASM did not induce biochemical defence processes, such as the production of antifungal secondary metabolites like phytoalexins through the phenylpropanoid pathway (Kombrink & Somssich, 1995; Kuć, 1995).

In contrast to inconsistent effects of ASM, MeJA was markedly effective in suppressing *B. cinerea* specking on cut freesia flowers when applied either pre- or postharvest. MeJA effectiveness was application method and concentration dependent. MeJA applied as gas was more effective compared to pulsing or spraying. It is possible that MeJA may function as an airborne signal which activated disease resistance and the expression of defence related genes in plant tissue (Shulaev et al., 1997). This finding agrees with earlier findings in *Arabidopsis* presented by Thomma et al. (2000). In *Arabidopsis*, this effect was mediated via the JA-dependent defence responses (Thomma et al., 2000). MeJA did not exert any direct antifungal activity in-vitro except at the concentration of 600 μM and therefore it is possible that MeJA reduced *B. cinerea* disease on freesia flowers by inducing responses correlated with the JA-dependent pathway (Darras et al., 2005). PPO levels in freesia flowers after MeJA gaseous treatment increased by 47 and 57% compared to the untreated controls (Darras et al., 2011). However, PAL activity decreased markedly compared 12 h post MeJA application and maintained at minimum level (i.e. ≈ 0). These findings suggest that MeJA might suppress the action of PAL in the phenylpropanoid pathway and consequently reduce or block SA production. Antagonistic regulation of JA- and SA-dependent pathways has been documented in the past by Pena-Cortes et al. (1993), Conconi et al. (1996), Niki et al. (1998), Gupta et al. (2000), and Rao et al. (2000). The apparent suppression of PAL in freesia flowers by MeJA constitute additional evidence of a JA- and SA- antagonistic response.

MeJA applied to freesia plants 28 days before harvest suppressed postharvest flower specking caused by *B. cinerea* in both a temperature and variety dependent fashion. MeJA was highly effective when flowers were incubated at 20°C compared to incubation at 5 or 12°C. It is likely that low incubation temperatures slow down plant's metabolism and also the production of defence related compounds (Jarvis, 1980). Overall, MeJA provided a considerable level of protection against *B. cinerea* when applied preharvest and, thus, could be considered a promising tool in an IDM context. Further study at the molecular level is

warranted to help interpret the MeJA mode of action in cut flowers. Also, additional in-planta trials on extra freesia varieties and a wider range of MeJA concentrations may help in better understanding MeJA efficacy.

In view to the promising results using MeJA, it is likely that elicitor based strategies within IDM could be used for the control of *Botrytis* or other pathogens on freesias and ornamental pot plants, as well as on various cut flowers. In turn, IDM would minimise the risk of pathogens developing resistance to fungicides and also reduce public concerns over extensive fungicide use (Jacobsen & Backman, 1993).

More research could be undertaken into potential synergistic effects of combined pre- and postharvest treatments with plant activators and/or abiotic biological agents (i.e. UV-C irradiation). In due course, pre and/or postharvest use of plant activators could have commercial potential for postharvest disease suppression (Kessmann et al., 1994; Kessmann et al., 1996; Thaler et al., 1996; Meir et al., 1998; Huang et al., 2000; (Darras et al., 2011).

4. References

- Agrios, G. N. 1997. *Plant Pathology* (4th edition), Academic Press, London UK
- Anfoka, G.H. (2000). Benzo-(1,2,3)-thiadiazole-7-carbothioic acid S-methyl ester induces systemic resistance in tomato (*Lycopersicon esculentum*. Mill cv. Vollendung) to Cucumber mosaic virus. *Crop Protection*, Vol.19, pp. 401-405
- Anonymous, (2011). FloraHolland reposts 7% increase in turnover. In *Floraculture International*, 30/08/2011, Available from:
<http://www.floracultureinternational.com/index.php?option=com_content&view=article&id=2196:floraholland-reports-7-increase-inturnover&catid=52:bussiness&Itemid=376>
- Atkinson, M.M. (1993). Molecular mechanisms of pathogen recognition by plants. *Advances in Plant Pathology*, Vol.10, pp. 35-64
- Beasley, D.R. (2001). Strategies for control of *Botrytis cinerea* on Geraldton waxflower flowers. PhD thesis. The University of Queensland, Australia
- Becktell, M.C.; Daughtrey, M.L. & Fry, W.E. (2005). Epidemiology and management of petunia and tomato late blight in the greenhouse. *Plant Disease*, Vol.89(9), pp. 1000-1008.
- Beier, R.C. & Oertli, E.H. (1983). Psoralen and other linear furocoumarins as phytoalexins in celery. *Phytochemistry*, Vol.22, pp. 2595-2597
- Benhamou, N. & Bélanger, R.R. (1998). Benzothiadiazole-mediated induced resistance to *Fusarium oxysporum* f.sp. *radicis-lycopersici* in tomato. *Plant Physiology*, Vol.118, pp. 1203-1212
- Bostock, R.; Kuć, J., & Laine, R. (1981). Eicosapentaenoic and arachidonic acids from *Phytophthora infestans* elicit fungitoxic sesquiterpenes in potato. *Science*, Vol.212, pp. 67-69
- Bolwell, G.P. (1999). Role of active oxygen species and NO in plant defence responses. *Current Opinion in Plant Biology*, Vol.2, pp. 287-294

- Brisset, M-N.; Cesbron, S.; Thomson S.V. & Paulin, J-P. (2000). Acibenzolar-S-methyl induces the accumulation of defense-related enzymes in apple and protects from fire blight. *European Journal of Plant Pathology*, Vol.106, pp. 529-536
- Buonaurio, R.; Scarponi, L.; Ferrara, M.; Sidoti, P. & Bertona, A. (2002). Induction of systemic acquired resistance in pepper plants by acibenzolar-S-methyl against bacterial spot disease. *European Journal of Plant Pathology*, Vol.108, pp. 41-49
- Cao, S.; Zheng, Y.; Yang Z.; Tang, S.; Jin, P.; Wang, K. & Wang, X. (2008). Effect of methyl jasmonate on the inhibition of *Colletotrichum acutatum* infection in loquat fruit and the possible mechanisms. *Postharvest Biology and Technology*, Vol.49, pp. 301-307.
- Cao, S.; Hu, Z.C.; Zheng, Y.H.; Li, X.W.; Wang, H.O. & Pang, B. (2010). Effect of post-harvest treatment with BTH on fruit decay, microbial populations, and the maintenance of quality in strawberry. *Journal of Horticultural Science & Biotechnology*, Vol.85(3), pp. 185-190
- Chalutz, E. & Stahmann, M. (1969). Induction of pisatin by ethylene. *Phytopathology*, Vol.59, pp. 1972-1973
- Chappell, J. & Hahlbrock, K. (1984). Transcription of plant defence genes in response to UV light or fungal elicitor. *Nature*, Vol.311, pp. 76-78
- Cole, D.L. (1999). The efficacy of acibenzolar-S-methyl, an inducer of systemic acquired resistance, against bacterial and fungal diseases of tobacco. *Crop Protection*, Vol.18, pp. 267-273
- Conconi, A.; Smerdon, M.J.; Howe, G.A. & Ryan, C.A. (1996). The octadecanoid signaling pathway in plants mediates a response to ultraviolet radiation. *Nature*, Vol.383, pp. 226-229
- Constabel, C.P. & Ryan C.A. (1998). A survey of wound and methyl jasmonate induced leaf polyphenol oxidase in crop plants. *Phytochemistry*, Vol.47, pp. 507-511
- Creelman, R.A.; Tierney, M.L. & Mullet, J.E. (1992). Jasmonic acid/methyl jasmonate accumulate in wounded soybean hypocotyls and modulated wound gene expression. *Proceedings of the National Academy for Science U.S.A.*, Vol.89, pp. 4938-4941
- Creelman R.A. & Mullet J.E. (1997). Biosynthesis and action of jasmonates in plants. *Annual Review of Plant Physiology and Plant Molecular Biology*, Vol.48, pp. 355-381
- Dann, E.; Diers, B.; Byrum, J. & Hammerschmidt, R. (1998). Effect of treating soybean with 2,6-dichloroisonicotinic acid (INA) and benzothiadiazole (BTH) on seed yields and the level of disease caused by *Sclerotinia sclerotiorum* in field and greenhouse studies. *European Journal of Plant Pathology*, Vol.104, pp. 271-278
- Darras, A.I. (2003). Biology and management of freesia flower specking caused by *Botrytis cinerea*. PhD Thesis, Cranfield University, UK
- Darras, A.I.; Joyce, D.C. & Terry, L.A. (2004). A survey of possible associations between pre-harvest environment conditions and postharvest rejections of cut freesia flowers. *Australian Journal of Experimental Agriculture*, Vol.44, pp. 103-108
- Darras, A.I.; Terry, L.A. & Joyce, D.C. (2005). Methyl jasmonate vapour treatment suppresses specking caused by *Botrytis cinerea* on cut *Freesia hybrida* L. flowers. *Postharvest Biology and Technology*, Vol.38, pp. 175-182

- Darras, A.I.; Joyce, D.C.; Terry, L.A. & Vloutoglou I. (2006a). Postharvest infections of *Freesia hybrida* L. flowers by *Botrytis cinerea*. *Australasian Plant Pathology*, Vol.35, pp. 55-63
- Darras, A.I.; Joyce, D.C. & Terry, L.A. (2006b). Acibenzolar-S-methyl and methyl jasmonate of glasshouse-grown freesias suppress postharvest petal specking caused by *Botrytis cinerea*. *Journal of Horticultural Science & Biotechnology*, Vol.81(6), pp. 1043-1051
- Darras, A.I.; Joyce, D.C.; Terry, L.A.; Pompodakis, N.E. & Dimitriadis, C.I. (2007). Efficacy of postharvest treatments with acibenzolar-S-methyl and methyl jasmonate against *Botrytis cinerea* infecting cut *Freesia hybrida* L. flowers. *Australasian Plant Pathology*, Vol.36, pp. 332-340
- Darras, A.I.; Joyce, D.C. & Terry, L.A. (2011). MeJA and ASM protect cut *Freesia hybrida* inflorescences against *Botrytis cinerea*, but do not act synergistically. *Journal of Horticultural Science & Biotechnology*, Vol.86(1), pp. 74-78
- Darras, A.I.; Demopoulos, V. & Tiniakou, C.A. (2012). UV-C irradiation induces defence responses and improves vase life of cut gerbera flowers. *Postharvest Biology and Technology* Vol.64, pp. 168-174
- Dinh, S.-Q.; Joyce, D.C.; Irving, D.E. & Wearing, A.H. (2007). Field applications of three different classes of known host plant defence elicitors did not suppress infection of Geraldton waxflower by *Botrytis cinerea*. *Australasian Plant Pathology*, Vol.36, pp. 142-148
- Dinh, S.-Q.; Joyce, D.C.; Irving, D.E. & Wearing, A.H. (2008). Effects of multiple applications of chemical elicitors on *Botrytis cinerea* infecting Geraldton waxflower. *Australasian Plant Pathology*, Vol.37, pp. 87-94
- Delaney, T.P.; Uknes, S.; Vernooij, B.; Friedrich, L.; Weymann, K.; Negretto, D.; Gaffney, T.; Gut-Rella, M.; Kessmann, H. & Ward, E. (1994). A central role of salicylic acid in plant disease resistance. *Science*, Vol.266, pp. 1247-1249
- Dimitriev, A.; Tena, M. & Jorrin, J. (2003). Systemic acquired resistance in sunflower (*Helianthus annuus* L.). *TSitologiia i Genetika*, Vol.37, pp. 9-15
- Dirkse, FB. (1982). Preharvest treatment of chrysanthemum against *Botrytis cinerea*. *Acta Horticulturae*, Vol.125, pp. 221-226.
- Dixon, R.A.; Lamb, C.J. & Harrison, M.J. (1994). Early events in the activation of plant defense responses. *Annual Review of Phytopathology*, Vol.32, pp. 479-501
- Doares, S.H.; Syrover, T.; Weiler, E. & Ryan C.A. (1995). Oligogalacturonides and chitosan activate plant defensive genes through the octadecanoid pathway. *Proceedings of the National Academy of Science U.S.A.*, Vol.92, pp. 4095-4098
- Dong, X. (1998). SA, JA, ethylene, and disease resistance in plants. *Current Opinions in Plant Biology*, Vol.1, pp. 316-323
- Droby, S.; Porat, R.; Cohen, L.; Weiss, B.; Shapiro, S.; Philisoph-Hadas, S. & Meir, S. (1999). Suppressing green mold decay in grapefruit with postharvest jasmonate application. *Journal of the American Society for Horticultural Science*, Vol.124(2), pp. 184-188
- Ebel, J. (1986). Phytoalexin synthesis: the biochemical analysis of the induction process. *Annual Review of Phytopathology*, Vol.24, pp. 235-264

- Ebel, J. & Cosio, E.G. (1994). Elicitors of plant defense responses. *International Review of Cytology*, Vol.148, pp. 1-36
- Elad, Y. (1988). Latent infection of *Botrytis cinerea* in rose flowers and combined chemical and physiological control of the disease. *Crop Protection*, Vol.7, pp. 361-366
- Elad, Y.; Kirshner, B. & Gotlib, Y. (1993). Attempts to control *Botrytis cinerea* on roses by pre- and postharvest treatments with biological and chemical agents. *Crop Protection* Vol.12, pp. 69-73
- Elad, Y. (1997). Responses of plants to infection by *Botrytis cinerea* and novel means involved in reducing their susceptibility to infection. *Biological Reviews*, Vol.72, pp. 381-342
- Elmer, W.H. (2006a). Efficacy of preplant treatments of gladiolus corms with combinations of acibenzolar-S-methyl and biological or chemical fungicides for suppression of fusarium corm rot [*Fusarium oxysporum* f. sp. *gladioli*]. *Canadian Journal of Plant Pathology*, Vol.28(4), pp. 609-624
- Elmer, W.H. (2006b). Effects of acibenzolar-S-methyl on the suppression of Fusarium wilt on cyclamen. *Crop Protection*, Vol.25, pp. 671-676
- Epple P.; Apel, K. & Bohlmann, H. (1997). Overexpression of an endogenous thionin enhances resistance of Arabidopsis against *Fusarium oxysporum*. *The Plant Cell*, Vol.9, pp. 509-520
- Evans, A. & van der Ploeg, R. (2008). Auctions around the world. *FloraCulture International*, Vol.5, pp. 8-9
- Eyre, J.X.; Faragher, J.; Joyce, D.C. & Franz, P.R. (2006). Effects of postharvest jasmonate treatments against *Botrytis cinerea* on Geraldton waxflower (*Chamelaucium uncinatum*). *Australasian Plant Pathology*, Vol.46, pp. 717-723
- Farmer, E.E. & Ryan, C.A. (1992). Octadecanoid precursors of jasmonic acid activate the synthesis of wound-inducible proteinase inhibitors. *Plant Cell*, Vol.4, pp. 129-134
- Flor, H.H. (1971). Current status of the gene-for-gene concept. *Annual Review of Phytopathology*, Vol.9, pp. 275-296
- Friedrich, L.; Lawton, K.; Ruess, W.; Masner, P.; Specker, N.; Gut Rella, M.; Meiers, B.; Dincher, S.; Staub, T.; Uknes, S.; Metraux, J-P.; Kessmann, H. & Ryals, J. (1996). A Benzothiadiazole derivative induces systemic acquired resistance in tobacco. *Plant Journal*, Vol.10, pp. 61-72
- Fritzemeier, K-H.; Cretin, C.; Kombrink, E.; Rohwer, F.; Taylor, J.; Scheel, D. & Hahlbrock, K. (1987). Transient induction of phenylalanine ammonia-lyase and 4-coumarate:CoA legase mRNAs in potato leaves infected with virulent or avirulent races of *Phytophthora infestans*. *Plant Physiology*, Vol.85, pp. 34-41
- Gast, K. (2001). Methyl jasmonate and long term storage of fresh cut peony flowers. *Acta Horticulturae*, Vol.543, pp. 327-330.
- Garrod, B.; Lewis, B.G.; Brittain, M.J. & Davies, W.P. (1982). Studies on the contribution of lignin and suberin to the impedance of wounded carrot root tissue to fungal invasion. *New Phytologist*, Vol.90, pp. 99-108
- Godard, J.-F.; Ziadi, S.; Monot, C.; Le Corre, D. & Silue, D. (1999). Benzothiadiazole (BTH) induces resistance in cauliflower (*Brassica oleracea* var *botrytis*) to downy

- mildew of crucifers caused by *Peronospora parasitica*. *Crop Protection*, Vol.18, pp. 397-405
- Gondim, D.M.F.; Terao, D.; Martins-Miranda, A.S.; Vansconcelos, I.M. & Oliveira, J.T.A. (2008). Benzo-thiadiazole-7-carbothioic acid S-methyl ester does not protect melon fruits against *Fusarium pallidoroseum* infection but induces defence responses in melon seedlings. *Journal of Phytopathology*, Vol.156(10), pp. 607-614
- Gorlach, J.; Volrath, S.; Knauf-Beiter, G.; Hengy, G.; Beckhove, U.; Kogel, K.-H.; Oostendorp, M.; Staub, T.; Ward, E.; Kessmann, H. & Ryals, J. (1996). Benzothiadiazole, a novel class of inducers of systemic acquired resistance, activates gene expression and disease resistance in wheat. *Plant Cell*, Vol.8, pp. 629-643
- Govrin, E.M. & Levine, A. (2002). Infection of Arabidopsis with a necrotrophic pathogen, *Botrytis cinerea*, elicits various defense responses but does not induce systemic acquired resistance (SAR). *Plant Molecular Biology*, Vol. 48, pp. 267-276
- Grisebach, H. (1981). *Lignins*. In 'The Biochemistry of Plants' (Ed. E.E. Conn). Chapter 7, pp. 457-478, Academic Press, New York, USA
- Gundlach, H.; Muller, M.J.; Kutchan, T.M. & Zenk, M.H. (1992). Jasmonic acid is a signal transducer in elicitor-induced plant cell cultures. *Proceedings of the National Academy of Science USA*, Vol.89, pp. 2389-2393
- Gupta, V.; Willits, M.G. & Glazebrook, J. (2000). *Arabidopsis thaliana* EDS4 contributes to salicylic acid (SA)-dependent expression of defense responses: evidence for inhibition of jasmonic acid signaling by SA. *Molecular Plant Microbe Interactions*, Vol.13, pp. 503-5011
- Hahlbrock, K. & Scheel, D. (1989). Physiology and molecular biology of phenylpropanoid pathway. *Annual Review of Plant Physiology and Plant Molecular Biology*, Vol.40, pp. 347- 369
- Hermes, D.A. & Mattson, W.J. (1992). The dilemma of plants: to grow or defend. *The Quarterly Review of Biology*, Vol.67, pp. 283-335
- Huang, Y.; Deverall, B.J.; Tang, W.H.; Wang, W. & Wu, F.W. (2000). Foliar application of acibenzolar-S-methyl and protection of postharvest rock melons and Hami melons from disease. *European Journal of Plant Pathology*, Vol.106, pp. 651-656
- Il'inskaya, L.I.; Goenburg, E.V.; Chalenko, G.I. & Ozeretskovskaya, O.L. (1996). Involvement of jasmonic acid in the induction of potato resistance to *Phytophthora* infection. *Russian Journal of Plant Physiology*, Vol.43(5), pp. 622-628
- Iriti, M.; Rossoni, M.; Borgo, M. & Faoro, F. (2004). Benzothiadiazole enhances resveratrol and anthocyanin biosynthesis in grapevine, meanwhile improving resistance to *Botrytis cinerea*. *Journal of Agricultural and Food Chemistry*, Vol.52, pp.4406-4413
- Isaac, S. (1992). *Fungal-Plant Interactions*. Chapman & Hall, London, UK
- Ishii, H.; Tomita, Y.; Horio, T.; Narusaka, Y.; Nakazawa, Y.; Nishimura, K. & Iwamoto, S. (1999). Induced resistance of acibenzolar-S-methyl (CGA 245704) to cucumber and Japanese pear diseases. *European Journal of Plant Pathology*, Vol.105, pp. 77-85
- Jacobsen, B.J. & Backman, P.A. (1993). Biological and cultural plant disease controls: alternative and supplements to chemicals in IPM systems. *Plant Disease*, Vol.77, pp. 311-315

- Jaiti, F.; Verdeil, J.L. & El Hadrami, I. (2009). Effect of jasmonic acid on the induction of polyphenoloxidase and peroxidase activities in relation to date palm resistance against *Fusarium oxysporum* f. sp. *albedinis*. *Physiological and Molecular Plant Pathology*, Vol.74(1), pp. 84-90
- Jarvis, W.R. (1977). *Botryotinia and Botrytis species: taxonomy, physiology, and pathogenicity. A guide to the literature*. Research Branch Canada Department of Agriculture, Monograph No 15
- Jarvis, W.R. (1980). *Epidemiology*. In 'The Biology of Botrytis' Eds. Coley-Smith, J.R., Verhoeff, K., and Jarvis, W.R. pp. 219-245, Academic Press, York, London
- Joyce, D.C. (1993). Postharvest floral organ fall in Geraldton waxflowers (*Chamelaucium uncinatum* Schauer). *Australian Journal of Experimental Agriculture*, Vol.33, pp. 481-487
- Joyce, D.C. & Johnson, G.I. (1999). Prospects for exploitation of natural disease resistance in harvested horticultural crops. *Postharvest News and Information*, Vol.10, pp. 45-48
- Katz, V.A.; Thulke, O.U. & Conrath, U. (1998). A Benzothiadiazole primes parsley cells for augmented elicitation of defense responses. *Plant Physiology*, Vol.117, pp. 1333-1339
- Keen, N.T. & Yoshikawa, M. (1983). β -1,3-endoglucanase from soybean releases elicitor-active carbohydrates from fungus cell walls. *Plant Physiology*, Vol.71, pp. 460-465
- Keller, H.; Pamboukdjian, N.; Ponchet, M.; Poupet, A.; Delon, R.; Verrier, J-L.; Roby, D. & Ricci, P. (1999). Pathogen-induced elicitor production in transgenic tobacco generates a hypersensitive response and nonspecific disease resistance. *The Plant Cell*, Vol.11, pp. 223-235
- Kendra, D.F. & Hadwiger, L.A. (1984). Characterization of the smallest chitosan oligomer that is maximally antifungal to *Fusarium solani* and elicits pisatin formation in *Pisum sativum*. *Experimental Mycology*, Vol.8, pp. 276-282
- Kessmann, H.; Staub, T.; Hofmann, C.; Maetzke, T. & Herzog, J. (1994). Induction of systemic acquired disease resistance in plants by chemicals. *Annual Review of Phytopathology*, Vol.32, pp. 439-459
- Kessmann, H.; Oostendorp, M.; Ruess, W.; Staub, T.; Kunz, W. & Ryals, J. (1996). Systemic activated resistance - a new technology for plant disease control. *Pesticide Outlook* pp. 10-13
- Kombrink, E. & Hahlbrock, K. (1986). Responses of cultured parsley cells to elicitors from phytopathogenic fungi. Timing and dose dependency of elicitor-induced reactions. *Plant Physiology*, Vol.81, pp. 216-221
- Kombrink, E. & Somssich, I. (1995). Defense responses of plants to pathogens. *Advances in Botanical Research*, Vol.21, pp. 2-34
- Kombrink, E. & Schmelzer, E. (2001). The hypersensitive response and its role in local and systemic disease resistance. *European Journal of Plant Pathology*, Vol.107, pp. 69-78
- Kuč, J. (1991). *Phytoalexins: perspectives and prospects*. In 'Mycotoxins and phytoalexins' (Eds. Sharma, R., and Salunkhe, D.), . pp. 595-603, Boca Raton: CRC, USA

- Kuč, J. (1995). Phytoalexins, stress metabolism and disease resistance in plants. *Annual Review of Phytopathology*, Vol.33, pp. 275-297
- Lawton, K.A.; Weymann, K.; Friedrich, L.; Vernooij, B.; Uknes, S. & Ryals, J. (1995). Systemic acquired resistance in Arabidopsis requires salicylic acid but not ethylene. *Molecular Plant-Microbe Interactions*, Vol.8, pp. 863-870
- Lawton, K.; Friedrich, L.; Hunt, M.; Weymann, K.; Delaney, T.; Kessmann, H.; Staub, T. & Ryals, J. (1996). Benzothiadiazole induces disease resistance in Arabidopsis by activation of the systemic acquired resistance signal transduction pathway. *Plant Journal*, Vol.10, pp. 71-82
- Lucas, J.A. (1997). *Plant Pathology and Plant Pathogens*. 3rd edition, Blackwell Science, UK
- Martinez, J.A.; Valdes, R.; Vicente, M.J.; & Banon, S. (2008). Phenotypical differences among *Botrytis cinerea* isolates from ornamental plants. *Communications in Agricultural and Applied Biological Sciences*, Vol.73, pp. 121-129
- Matern, U. & Kneusel, R. (1988). Phenolic compounds in plant disease resistance. *Phytoparasitica*, Vol.16, pp. 153-170
- Mauch, F.; Hadwiger, L.A. & Boller, T. (1988). Antifungal hydrolases in pea tissue. 2. Inhibition of fungal growth by combinations of chitinase and β -1,3-glucanase in pea pods by pathogens and elicitors. *Plant Physiology*, Vol.76, pp. 607-611
- Mauch-Mani, B. & Slusarenko, A.J. (1996). Production of salicylic acid precursors is a major function of phenylalanine ammonia lyase in the resistance of Arabidopsis to *Peronospora parasitica*. *The Plant Cell*, Vol.8, pp. 203-212
- Meir, S.; Droby, S.; Davidson, H.; Alsvia, S.; Cohen, L.; Horev, B. & Philosoph-Hadas, S. (1998). Suppression of Botrytis rot in cut rose flowers by postharvest application of methyl jasmonate. *Postharvest Biology and Technology*, Vol.13, pp. 235-243
- Meir, S.; Droby, S.; Kochanek, S.; Salim, S. & Philosoph-Hadas, S. (2005) Use of methyl jasmonate for suppression of Botrytis rot in various cultivars of cut rose flowers. *Acta Horticulturae*, Vol.669, pp. 91-98
- Métraux, J-P. (2001). Systemic acquired resistance and salicylic acid: current state of knowledge. *European Journal of Plant Pathology*, Vol.107, pp. 13-18
- Mitter, N.; Kazan, K.; Way, H.M.; Broekaert, W.F. & Manners, J.M. (1998). Systemic induction of an Arabidopsis plant defensin gene promoter by tobacco mosaic virus and jasmonic acid in transgenic tobacco. *Plant Science*, Vol.136, pp. 169-180
- Narusaka, Y.; Narusaka, M.; Horio, T. & Ishii, H. (1999). Induction of disease resistance in cucumber by acibenzolar-S-methyl and expression of resistance-related genes. *Annals of the Phytopathological Society of Japan*, Vol.65, pp. 116-122
- Nasser, W.; De Tapia, M.; Kauffmann, S.; Montasser-Kouhsari, S. & Burkard, G. (1988). Identification and characterization of maize pathogenesis related proteins. Four maize PR proteins are chitinases. *Plant Molecular Biology*, Vol.11, pp. 529-538
- Niki, T.; Mitsuhara, I.; Seo, S.; Ohtsubo, N. & Ohashi, Y. (1998). Antagonistic effect of salicylic acid and jasmonic acid on the expression of pathogenesis-related (PR) protein genes in wounded mature tobacco leaves. *Plant Cell Physiology*, Vol.39(5), pp. 500-507
- Nojiri, H.; Sugimori, M.; Yamane, H.; Nishimura, Y.; Yamada, A.; Shibuya, N.; Kodama, O.; Murofushi, N. & Omori, T. (1996). Involvement of jasmonic acid in elicitor-induced

- phytoalexin production in suspension-cultured rice cells. *Plant Physiology*, Vol.110, pp. 387-392
- Nurnberger, T.; Nennstiel, D.; Jabs, T.; Sacks, W.R.; Hahlbrock, K. & Scheel, D. (1994). High affinity binding of a fungal oligopeptide elicitor to parsley plasma membranes triggers multiple defenses responses. *The Plant Cell*, Vol.78, pp. 449-460
- Nurnberger, T. & Scheel, D. (2001). Signal transmission in the plant immune response. *Trends in Plant Science*, Vol.6(8), pp. 372-379
- O'Donnell, P.J.; Calvert, C.; Atzorn, R.; Wasternack, C.; Leyser, H.M.O. & Bowles, D.J. (1996). Ethylene as a signal mediating the wound response of tomato plants. *Science*, Vol.274, pp. 1914-1917
- O'Neil, T.M.; Hanks, G.R. & Wilson, T.W. (2004). Control of smoulder (*Botrytis narcissicola*) in narcissus with fungicides. *Annals of Applied Biology*, Vol.145, pp. 129-137
- Pallas, J.A.; Paiva, N.L.; Lamb, C. & Dixon, R.A. (1996). Tobacco plants epigenetically suppressed in phenylalanine ammonia lyase expression do not develop systemic acquired resistance in response to infection by tobacco mosaic virus. *The Plant Journal*, Vol.10, pp. 281-293
- Parthier, B. (1991). Jasmonates, new regulators of plant growth and development: many facts and few hypotheses on their actions. *Botanica Acta* pp. 446-454
- Pappas, A.C. (1997). Evolution of fungicide resistance in *Botrytis cinerea* in protected crops in Greece. *Crop Protection*, Vol.16, pp. 257-263
- Pearce, G.; Strydom, D.; Johnson, S. & Ryan, C.A. (1991). A polypeptide from tomato leaves induces wound-inducible proteinase inhibitor proteins. *Science*, Vol.253, pp. 895-898
- Pena-Cortes, H.; Albrecht, T.; Prat, S.; Weiler, E.W. & Willmitzer, L. (1993). Aspirin prevents wound-induced gene expression in tomato leaves by blocking jasmonic acid biosynthesis. *Planta*, Vol.191, pp. 123-128
- Penninckx, I.A.M.A.; Eggermont, K.; Terras, F.R.G.; Thomma, B.P.H.J.; De Samblanx, G.W.; Buchala, A.; Mettraux, J-P.; Manners J.M. & Broekaert, W.F. (1996). Pathogen-induced systemic activation of a plant defensin gene in Arabidopsis follows a salicylic acid-independed pathway. *The Plant Cell*, Vol.8, pp. 2309-2323
- Perrin, D.R. & Cruickshank, I.A.M. (1965). Studies on phytoalexins VII. The chemical stimulation of pisatin formation in *Pisum sativum* L. *Austrian Journal of Biological Science*, Vol.18, pp. 803-816
- Pichay, D.S.; Frantz, J.M.; Locke, J.C.; Krause, C.R. & Fernandez, G.C.J. (2007). Impact of applied nitrogen concentration on growth of elatior Begonia and new guinea impatiens and susceptibility of Begonia to *Botrytis cinerea*. *Journal of the American Society for Horticultural Science*, Vol.132, pp. 193-201
- Pie, K. & de Leeuw, G.T.N. (1991). Histopathology of the initial stages of interaction between rose flowers and *Botrytis cinerea*. *Netherlands Journal of Plant Pathology*, Vol.97, pp. 335-344
- Pieterse, C.M.J.; van Wees, S.; Hoffland, E.; van Pelt, J.A. & van Loon, L.C. (1996). Systemic resistance in Arabidopsis induced by biocontrol bacteria is independent of salicylic acid accumulation and pathogenesis related gene expression. *The Plant Cell*, Vol.8, pp. 1225-1237

- Pieterse, C.M.J.; van Wees, S.; van Pelt, J.A.; Knoester, M.; Laan, R.; Gerrits, H.; Weisbeek, P.J. & van Loon L.C. (1998). A novel signalling pathway controlling induced systemic resistance in Arabidopsis. *The Plant Cell*, Vol.10, pp. 1571-1580
- Pieterse, C.M.J. & van Loon, L.C. (1999). Salicylic acid-independent plant defence pathways. *Trends in Plant Science*, Vol.4(2), pp. 52-58
- Rao, M.V.; Lee, H-I.; Creelman, R.A.; Mullet, J.E. & Davis, K.R. (2000). Jasmonic acid signaling modulates ozone-induced hypersensitive cell death. *The Plant Cell*, Vol.12, pp. 1633-1646
- Redolfi, P. (1983). Occurrence of pathogenesis related and similar proteins in different plant species. *Netherlands Journal of Plant Pathology*, Vol.89, pp. 245-254
- Rojo, E.; Titarenko, E.; Leon, J.; Berger, S.; Vancanneyt, G. & Sanchez-Serrano, J.J. (1998). Reversible protein phosphorylation regulates jasmonic acid-dependent and independent wound signal transduction pathways in *Arabidopsis thaliana*. *The Plant Journal*, Vol.13, pp. 153-165
- Rojo, E.; Leon, J. & Sanchez-Serrano, J.J. (1999). Cross-talk between wound signaling pathways determines local versus systemic gene expression in *Arabidopsis thaliana*. *The Plant Journal*, Vol.20, pp. 136-142
- Ross, A.F. (1961). Systemic acquired resistance induced by localized virus infections in plants. *Virology*, Vol.14, pp. 341-358
- Ruess, W.; Mueller, K.; Knauf-Beiter, G.; Kenz, W. & Staub, T. (1996). Plant activator CGA 245704: an innovative approach for disease control in cereals and tobacco. *Brighton Crop Protection Conference-Pests and Diseases*, Vol.2A-6, pp. 53-60, 18-21 November 1996
- Ryals, J.A.; Ward, E. & Metraux J.P. (1992). *Systemic acquired resistance: An inducible defense mechanism in plants*. In 'Inducible Plant Proteins: Their Biochemistry and Molecular Biology', (Ed. J.L. Wray), pp. 205-229, Cambridge University Press, Cambridge UK
- Ryals, J.A.; Neuenschwander, U.H.; Willits, M.G.; Molina, A.; Steiner, H-Y. & Hunt, M.D. 1996. Systemic Acquired Resistance. *The Plant Cell*, Vol.8, pp. 1809-1819
- Salinas, J.; Glandorf, D.C.M.; Picavet, F.D. & Verhoeff, K. (1989). Effects of temperature, relative humidity and age of conidia on the incidence of specking on gerbera flowers caused by *Botrytis cinerea*. *Netherlands Journal of Plant Pathology*, Vol.95, pp. 51-64
- Salinas, J. & Verhoeff, K. (1995). Microscopic studies of the infection of gerbera flowers by *Botrytis cinerea*. *European Journal of Plant Pathology*, Vol.101, pp. 377-386
- Seglie, L.; Spadano, D.; Devecchi, M.; Larcher, F.; Gullino, M.L. (2009). Use of 1-methylcyclopropene for the control of *Botrytis cinerea* on cut flowers. *Phytopathologia Mediterranea*, Vol.48, pp. 253-261
- Schaller, A. & Ryan, C.A. (1995). Systemin - a polypeptide defense signal in plants. *BioEssays*, Vol.18(1), pp. 27-33
- Schroder, M.; Hahlbrock, K. & Kombrink, E. (1992). Temporal and spatial patterns of 1,3- β -glucanase and chitinase induction in potato leaves infected by *Phytophthora infestans*. *Plant Journal*, Vol.2, pp. 161-172
- Sembdner, G. & Parthier, B. (1993). The biochemistry and the physiological and molecular actions of jasmonates. *Annual Review of Plant Physiology and Plant Molecular Biology*, Vol.44, pp. 569-589


- Sharkey, T.D. (1996). Emission of low molecular mass hydrocarbons from plants. *Trends in Plant Science*, Vol.3, pp. 78-82
- Sharp, J.; Albersheim, P.; Ossowski, O.; Pilotti, A.; Garegg, P. & Lindberg, B. (1984). Comparison of the structure of elicitor activities of a synthetic and mycelial-wall-derived hexa-(β -D-glucopyranosyl)-D-glucitol. *Journal of Biological Chemistry*, Vol.259, pp. 11341-11345
- Shulaev, V.; Silverman, P. & Raskin, I. (1997). Airborne signaling by methyl salicylate in plant pathogen resistance. *Nature*, Vol.385, pp. 718-721
- Staswick, P.E. (1990). Novel regulation of vegetative storage protein genes. *The Plant Cell*, Vol.2, pp. 1-6
- Stermer, B.A. (1995). *Molecular regulation of systemic induced resistance*. In "Induced Resistance to Disease in Plants". Eds. R. Hammerschmidt and Kuć, J. pp. 111-140, Kluwer Academic Publishers, USA
- Sticher, L.; Mauch-Mani, B. & Metraux, J.P. (1997). Systemic Acquired Resistance. *Annual Review of Phytopathology*, Vol.35, pp. 235-270
- Strider, D.L. (1985). *Diseases of Floral Crops*. Vol. 2, Praeger Publishers, New York, USA
- Suo, Y.; & Leung, D.W.M. (2002). BTH-induced accumulation of extracellular proteins and blackspot disease in rose. *Biologia Plantarum*, Vol.45, pp. 273-279
- Taiz, L. & Zeiger, E. (1998). *Plant Physiology*. 2nd ed. pp. 61-80, Sinauer Associates, Inc., Publishers, USA
- Terry, L.A. & Joyce, D.C. (2000). *Suppression of grey mould on strawberry fruit with the chemical plant activator acibenzolar*. *Pest Management Science*, Vol.56, pp. 989-992
- Terry, L.A. (2002). Natural disease resistance in strawberry fruit and Geraldton waxflower flowers. PhD Thesis, Cranfield University, UK
- Terry, L.A. & Joyce, D.C. (2004a) Elicitors of induced disease resistance in postharvest horticultural crops: a brief review. *Postharvest Biology and Technology*, Vol.32, pp. 1-13
- Terry, L.A. & Joyce, D.C. (2004b) Influence of growing conditions on efficacy of acibenzolar and botryticides in suppression of *Botrytis cinerea* on strawberry fruit. *Advances in Strawberry Research*, Vol.23, pp. 11-19
- Thaler, J.S.; Stout, M.J.; Karban, R. & Duffey, S.S. (1996). Exogenous jasmonates simulate insect wounding in tomato plants (*Lycopersicon esculentum*) in the laboratory and field. *Journal of Chemical Ecology*, Vol.22, pp. 1767-1781
- Thaler, J.S.; Fidantsef, A.L.; Duffey, S.S. & Bostock R.M. (1999). Trade-offs in plant defense against pathogens and herbivores: a field demonstration of chemical elicitors of induced resistance. *Journal of Chemical Ecology*, Vol.25, pp. 1597-1609
- Thomma, B. P.H.J.; Eggermont, K.; Penninckx, I.A.M.A.; Mauch-Mani, B.; Vogelsang, R.; Cammue, B.P.A. & Broekaert, W. (1998). Separate jasmonate-dependent and salicylate-dependent defense response pathways in Arabidopsis are essential for resistance to distinct microbial pathogens. *Proceedings of the National Academy of Science USA*, Vol.95, pp. 15107-15111
- Thomma, B. P.H.J.; Eggermont, K.; Broekaert, W. & Cammue, B.P.A. (2000). Disease development of several fungi on Arabidopsis can be reduced by treatment with methyl jasmonate. *Plant Physiology and Biochemistry*, Vol.38, pp. 421-427.

- Thomma, B. P.H.J.; Penninckx, I.A.M.A.; Broekaert, W.F. & Cammue, B.P.A. (2001). The complexity of disease signaling in *Arabidopsis*. *Current Opinion in Immunology*, Vol.13, pp. 63-68
- Titarenko, E.; Rojo, E.; Leon, J. & Sanchez-Serrano, J.J. (1997). Jasmonic acid-dependent and independent signaling pathways control wound-induced gene activation in *Arabidopsis thaliana*. *Plant Physiology*, Vol.115, pp. 817-826
- Tomas, A.; Wearing, A.H. & Joyce, D.C. (1995). *Botrytis cinerea*: a causal agent of premature flower drop in packaged Geraldton waxflower. *Australasian Plant Pathology*, Vol.24, pp. 26-28
- Tosi, L. & Zizzerini, A. (1999). Benzothiadiazole induces resistance to *Plasmopara helianthi* in sunflower plants. *Journal of Phytopathology*, Vol.147, pp. 365-370
- Van Loon, L.C. & Gerritsen, Y.A.M. (1989). Localization of pathogenesis-related proteins in infected and non-infected leaves of Samsun NN tobacco during the hypersensitive reaction to tobacco mosaic virus. *Plant Science*, Vol.63, pp. 131-140
- Van Loon, L.C.; Pierpoint, W.S.; Boller, T. & Conejero, V. (1994). Recommendations for naming plant pathogenesis-related proteins. *Plant Molecular Biology Reporter*, Vol.12, pp. 245-264
- Van Loon, L.C. (1997). Induced resistance in plants and the role of pathogenesis-related proteins. *European Journal of Plant Pathology*, Vol.103, pp. 753-765
- Van Meeteren, U. (2009). Causes of quality loss of cut flowers - A critical analysis of postharvest treatments. *Acta Horticulturae*, Vol.847, pp. 27-36
- Vernooij, B.; Friedrich, L.; Morse, A.; Reist, R., Kolditz-Jawhar, R.; Ward, E.; Uknes, S.; Kessmann, H. & Ryals, J. (1994). Salicylic acid is not the translocated signal responsible for inducing systemic acquired resistance but is required in signal transduction. *The Plant Cell*, Vol.6, pp. 959-965
- Ward, E.R.; Uknes, S.J.; Williams, S.C.; Dincher, S.S.; Wiederhold, D.L.; Alexander, D.C.; Ahi-Goy, P.; Mettraux, J.P. & Ryals, J.A. (1991). Coordinate gene activity in response to agents that induce systemic acquired resistance. *The Plant Cell*, Vol.3, pp. 1085-1094
- Wegulo, S.N. & Vilchez, M. (2007). Evaluation of lisianthus cultivars for resistance to *Botrytis cinerea*. *Plant Disease*, Vol.91, pp. 997-1001
- Wasternack, C. & Parthier, B. (1997). Jasmonate-signalled plant gene expression. *Trends in Plant Science*, Vol.2, pp. 302-307
- Williamson, B.; Duncan, G.H.; Harrison, J.G.; Harding, L.A.; Elad, Y. & Zimand, G. (1995). Effect of humidity on infection of rose petals by dry-inoculated conidia of *Botrytis cinerea*. *Mycological Research*, Vol.99, pp. 1303-1310
- Xu, Y.; Linda Chang, P-F.; Liu, D.; Narasimhan, M.L.; Raghothama, K.G.; Hasegawa, P.M. & Bressan, R.A. (1994). Plant defense genes are synergistically induced by ethylene and methyl jasmonate. *The Plant Cell*, Vol.6, pp. 1077-1085
- Yao, H. & Tian, S. (2005). Effects of pre- and post-harvest application of salicylic acid or methyl jasmonate on inducing disease resistance of sweet cherry fruit in storage. *Postharvest Biology and Technology*, Vol.35, pp. 253-262
- Yoshikawa, M.; Yamaoka, N. & Takeuchi, Y. (1993). Elicitors: their significance and primary modes of action in the induction of plant defense reactions. *Plant Cell Physiology*, Vol.34, pp. 1163-1173

Ziadi, S.; Barbedette, S.; Godard, J.F.; Monot, C.; Le Corre, D. & Silue, D. (2001). Production of pathogenesis-related proteins in the cauliflower (*Brassica oleracea* var. *botrytis*)-downy mildew (*Peronospora parasitica*) pathosystem treated with acibenzolar-S-methyl. *Plant Pathology*, Vol.50, pp. 579-586

IntechOpen

IntechOpen



Plant Pathology

Edited by Dr. Christian Joseph Cumagun

ISBN 978-953-51-0489-6

Hard cover, 362 pages

Publisher InTech

Published online 04, April, 2012

Published in print edition April, 2012

Plant pathology is an applied science that deals with the nature, causes and control of plant diseases in agriculture and forestry. The vital role of plant pathology in attaining food security and food safety for the world cannot be overemphasized.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Anastasios I. Darras (2012). Novel Elicitors Induce Defense Responses in Cut Flowers, Plant Pathology, Dr. Christian Joseph Cumagun (Ed.), ISBN: 978-953-51-0489-6, InTech, Available from:
<http://www.intechopen.com/books/plant-pathology/novel-elicitors-induce-defense-responses-in-cut-flowers>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen