

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



# Antioxidant Adaptive Response of Malignant Glioma Related to Resistance to Antitumor Treatment

Tomohiro Sawa and Takaaki Akaike

*Department of Microbiology, Graduate School of Medical Sciences, Kumamoto University  
Japan*

## 1. Introduction

Glioblastomas are the most frequent and most malignant nervous system tumors [Stewart & Kleihues, 2003]. Despite technological advances in surgical treatment and new regimens of radiotherapy combined with chemotherapy, the median survival of patients with these tumors is approximately 1 year, and only 3% of patients survive more than 3 years [Stupp et al., 2005]. Glioblastomas have been traditionally defined as two clinically and cytogenetically distinct diseases: the primary or de novo glioblastomas and the secondary glioblastomas. The latter commonly appear in younger people (median age at onset ~45 years) as low-grade gliomas and possess aberrations in genes encoding platelet-derived growth factor receptor (*PDGFR*) and *TP53* [Stewart & Kleihues, 2003]. Primary glioblastomas occur more frequently (>80% of cases) and develop rapidly in older people (median age at onset ~60 years); survival of patients with such tumors is short, less than 3 months [Stewart & Kleihues, 2003]. The genetic profile of primary glioblastomas includes amplification and overexpression of the gene encoding epidermal growth factor receptor (*EGFR*), mutations of the phosphatase and tensin homolog (*PTEN*) gene, *p16<sup>INK4A</sup>* deletions, and loss of chromosome 10 [Stewart & Kleihues, 2003]. Several inhibitors that target *EGFR* or its downstream signaling cascade including Akt and mTOR have been evaluated for potential application in glioblastoma treatment [Krakstad & Chekenya, 2010]. Recent clinical trials of *EGFR* inhibitors, however, showed no therapeutic benefit [Prados et al., 2006; Rich et al., 2004].

Chemotherapy plays an important role in combined treatment of gliomas, whereas it appears to fail in a significant clinical outcome. This may be largely due to drug resistance of malignant gliomas developed [Lu & Shervington, 2008; Sarkaria, et al, 2008; Frosina, 2009; Bleau et al., 2009]. Multiple mechanisms may be involved in the development of drug resistance in gliomas, including DNA repair enzyme activities, particularly O<sup>6</sup>-methylguanine methyltransferase for alkylating agents, overexpression of antiapoptotic proteins such as Bcl-2 or Bcl-X<sub>L</sub> and ABC transporters that efflux anticancer drugs. Clarification of mechanisms for glioma drug resistance may provide us an important insight for the development of promising strategies for the treatment of gliomas.

A number of studies have suggested that heme oxygenase-1 (HO-1) expression occurs in various types of tumors, both animal models and human cancers including glioblastomas

[Fang et al., 2004a; Hara et al., 1996]. HO is a rate-limiting enzyme that catalyzes the initial step of heme degradation in which oxidative cleavage of the porphyrin ring results in generation of biliverdin, carbon monoxide (CO), and free iron [Maines, 1988; Schacter, 1988]. Cytosolic biliverdin reductase then reduces the biliverdin to form the potent antioxidant bilirubin. Three mammalian HO isoforms have been identified: HO-1, HO-2, and HO-3. Among them, HO-1 is a member (HSP-32) of the heat shock protein family, and HO-1 expression is triggered by various stress-inducing stimuli including hypoxia [Motterlini et al., 2000], heavy metals [Keyse & Tyrrell, 1989; Mitani et al., 1993], UV irradiation [Keyse & Tyrrell, 1989], and reactive oxygen species (ROS) and nitric oxide (NO) [Doi et al., 1999; Foresti et al., 1997]. Because of the antioxidant and cytoprotective nature of HO products, elevated HO-1 expression in glioblastomas may be a key component of cellular adaptation to oxidative stress and toxic insults induced by chemotherapeutic agents [Fang et al., 2004b]. The fact that HO-1 expression is related to resistance of glioma cells to oxidative stress and anticancer agents indicates that the pathways involved in regulating HO-1 expression in gliomas may be molecular targets for treatment. This chapter describes the mechanisms and regulation of HO-1 expression and the adaptive response of glioma cells. The chapter pays particular attention to the adaptive response signaling that is mediated by 8-nitroguanosine 3',5'-cyclic monophosphate (8-nitro-cGMP), a newly discovered nitrated nucleotide that functions as an endogenous inducer of the response [Akaike et al., 2010; Fujii et al., 2010; Sawa et al., 2007; Zaki et al., 2009]. Also discussed is the potential application of HO inhibitors as a new class of chemotherapeutic and chemosensitizing agents.

## **2. Antioxidant adaptive response mediated by a nitrated cyclic nucleotide**

The antioxidant adaptive response to oxidative stress is a critically important mechanism that allows aerobic organisms to maintain redox homeostasis. The Keap1 (Kelch-like ECH-associated protein 1)/Nrf2 (nuclear factor-erythroid 2-related factor 2) system plays a key role in adaptation to oxidative stress and toxic insults of electrophilic compounds (including many chemotherapeutic agents) by inducing antioxidant and cytoprotective enzymes such as HO-1.

### **2.1 The Keap1/Nrf2 system as a sensor of oxidants and electrophiles**

Keap1 contains reactive cysteine residues that can serve as sensing moieties for oxidants and electrophiles [Dinkova-Kostova et al., 2005; Itoh et al., 2004; Sekhar et al., 2010]. Various electrophiles called Nrf2 inducers, both naturally occurring and synthetic, have reportedly reacted with Keap1 cysteine residues.

Keap1, which exists as dimers inside cells, comprises five distinct domains: (i) the N-terminal region (NTR, amino acids 1-60); (ii) the BTB (Bric-a-brac, Tramtrack, Broad-complex) domain (amino acids 61-178), which is an evolutionarily conserved protein-protein interaction motif that occurs in actin-binding proteins and zinc finger transcription factors and participates in binding to Rbx1-bound Cullin 3 (Cul3) [Dinkova-Kostova et al., 2005] and formation of homodimers [Zipper & Mulcahy, 2002]; (iii) the intervening region (IVR, amino acids 179-321), which is a central linker domain, especially rich in cysteine, that also takes part in binding to Rbx1-bound Cul3 [A. Kobayashi et al., 2004]; (iv) the Kelch repeat domain (amino acids 322-608), which mediates binding to the Nrf2 domain of Nrf2 [M.

Kobayashi et al., 2002]; and (v) the C-terminal region (CTR, amino acids 609-625). Site-directed mutagenesis assays have revealed that three cysteine residues, Cys151, Cys273, and Cys288, play critical roles in the Keap1-Nrf2 complex interaction [D.D. Zhang & Hannink, 2003]. Cys151, located in the BTB domain, is probably the major site that is directly alkylated by chemopreventive agents that work by activating the antioxidant response element (ARE) through Nrf2 [Eggler et al., 2007]. This modification of Cys151 leads to conformational changes in the BTB domain via perturbation of the homodimerization site and results in switching ubiquitination from Nrf2 to Keap1, thereby facilitating translocation of Nrf2 to the nucleus and its accumulation there [Eggler et al., 2005]. Cys273 and Cys288, which are located in the IVR domain of Keap1, are essential for stability of the Nrf2-Keap1 complex and maintenance of Nrf2 homeostasis [Wakabayashi et al., 2004]. ARE inducers, via conformational changes in Keap1—i.e., covalent modification or oxidation of cysteine thiol groups in the IVR region of Keap1—make Keap1 incapable of binding to Nrf2 and thereby facilitate nuclear translocation of Nrf2.

Nrf2, a member of the cap'n'collar family of basic region leucine zipper transcription factors, has a crucial role in regulating several stress-responsive genes, including HO-1 [Dinkova-Kostova et al., 2005; Bloom & Jaiswal, 2003; Itoh et al., 2004; Zhu et al., 2005]. Under basal conditions, transcriptional activity of Nrf2 is negatively regulated by the Nrf2-binding partner Keap1, a cysteine-rich cytoplasmic protein [Dinkova-Kostova et al., 2005; Sekhar et al., 2010; Surh et al., 2009; Zhao et al., 2010]. Keap1, serving as a substrate adaptor protein for a Cul3-dependent ubiquitin ligase complex, targets Nrf2 for ubiquitination and proteasomal degradation [Dinkova-Kostova et al., 2005; Sekhar et al., 2010; Surh et al., 2009; Zhao et al., 2010]. During oxidative stress, chemical modification of one or more cysteine residues of Keap1 facilitates Nrf2 dissociation, which leads to nuclear translocation of Nrf2. After migration to the nucleus, Nrf2 forms heterodimers with small musculoaponeurotic fibrosarcoma (sMaf) proteins and then binds to the cis-acting ARE [Motohashi & Yamamoto, 2007]. The result is transcriptional activation of a battery of genes that encode various phase II detoxifying or antioxidant enzymes, such as glutathione *S*-transferase (GST), NAD(P)H:quinone oxidoreductase (NQO1), and HO-1, as well as other cytoprotective proteins [Nguyen et al., 2003].

## **2.2 Nitrated guanine nucleotide as an endogenous electrophile that activates the Keap1/Nrf2 system**

Both NO and ROS are critically involved, via Nrf2-dependent mechanisms, as endogenous inducers of HO-1 expression in gliomas. 8-Nitro-cGMP is a nitrated nucleotide that was identified as a second messenger in NO and ROS signaling in induction of an adaptive response mediated by the Keap1/Nrf2 system [Akaike et al., 2010; Fujii et al., 2010; Ihara et al., 2011b; Sawa et al., 2007, 2011; Zaki et al., 2009]. 8-Nitro-cGMP also functions in extracellular signal-regulated kinase pathways in neurons [Kurauchi et al., 2011]. Formation of 8-nitro-cGMP can be triggered in rat C6 glioma cells by treatment with a chemical NO donor (*S*-nitroso-*N*-acetylpenicillamine or treatment with inflammatory stimuli (lipopolysaccharide plus pro-inflammatory cytokines) that activate endogenous NO production via expression of inducible NO synthase [Fujii et al., 2010]. Under these conditions, ROS production is also increased and depends on both NADPH oxidase and mitochondria [Ahmed et al., 2011a]. Pharmacological studies have suggested that 8-nitro-cGMP is formed from its precursor 8-nitroguanosine 5'-triphosphate by means of catalytic action of soluble guanylate cyclase [Fujii

et al., 2010; Ihara et al., 2011a]. So far, guanine nitration has been identified in various biological systems including lungs of mice suffering from viral pneumonia [Akaike et al., 2003], lungs of patients with idiopathic pulmonary fibrosis and lung cancer [Terasaki et al., 2006], and urine from cigarette smokers [Sawa et al., 2006].

Activation of the Keap1/Nrf2-mediated adaptive response by 8-nitro-cGMP involves a unique post-translational modification of the cysteine residues of Keap1 [Akaike et al., 2010; Fujii et al., 2010; Sawa et al., 2007; Zaki et al., 2009]. 8-Nitro-cGMP is the first known endogenously formed electrophilic nucleotide [Akaike et al., 2010; Fujii et al., 2010; Saito et al., 2008; Sawa et al., 2007; Zaki et al., 2009]. Because of its electrophilic characteristics, 8-nitro-cGMP reacts with cysteine residues in proteins, particularly residues with low  $pK_a$  values that facilitate deprotonation of those residues to form a cysteine thiolate anion [Ahmed et al., 2011b; Akaike et al., 2010; Sawa et al., 2010]. Reaction of 8-nitro-cGMP with the thiolate anion results in cGMP adduction to the cysteine residue, a process called "protein S-guanylation" (Fig. 1) [Sawa et al., 2007]. Keap1 was identified as a possible target during 8-nitro-cGMP-induced protein S-guanylation in cells [Fujii et al., 2010; Sawa et al., 2007; Zaki et al., 2009]. In fact, Keap1 S-guanylation was clearly demonstrated when rat C6 glioma cells were treated with lipopolysaccharide plus pro-inflammatory cytokines [Fujii et al., 2010]. Mass spectrometry revealed that Keap1 S-guanylation in rat C6 cells occurred predominantly at Cys434 [Fujii et al., 2010]. X-ray crystallographic analysis suggested that the Cys434 residue is located at blade 3 and is exposed to the outer surface of the  $\beta$ -propeller structure [Lo et al., 2006; Padmanabhan et al., 2006].

With regard to the mechanism of how S-guanylation of Cys434 causes Nrf2 activation, two possibilities present themselves. One is that S-guanylation of Cys434 may weaken Keap1 binding to the ETGE and DLG motifs of Nrf2, because Cys434 is located close to the Nrf2-binding region of the DC domain. The alternative possibility is that Cys434 modification may affect the integrity of the entire Keap1-Nrf2 complex. Fitting the atomic DC domain model into the overall Keap1 homodimer structure obtained via single-particle electron microscopy showed that the globular part of the Keap1 cherry-bob structure was bulkier than the DC domain, which suggests that the external DC domain surface is wrapped with the other part of Keap1, perhaps the IVR between the BTB and DC domains [Ogura et al., 2010]. Thus, S-guanylation may cause disruption of the globular structure incorporating the DC domain, which would disturb the entire structure and result in reduced ubiquitin ligase activity of the Cul3-Keap1 complex.

Among genes regulated by Nrf2, HO-1 has been regarded as conferring important protection against oxidative stress [T.J. Chen et al., 2006; Y.C. Chen et al., 2006; Konorev et al., 2002]. HO-1 may induce cytoprotective responses through various mechanisms including (i) reducing prooxidant levels (heme) [Jeney et al., 2002]; (ii) raising antioxidant levels (bilirubin) [Baranano et al., 2002]; (iii) generating the antiapoptotic CO [Brouard et al., 2000]; (iv) inducing ferritin, which detoxifies and removes free ferric ions [Balla et al., 1992]; and (v) blocking overstimulation of an immune response [Lee & Chau, 2002]. We found that 8-nitro-cGMP increased Nrf2 accumulation in the nucleus and HO-1 expression in rat C6 glioma cells (Fig. 2) [Fujii et al., 2010]. Treatment with 8-nitro-cGMP caused C6 cells to become resistant to cell death induced by oxidative stress related to hydrogen peroxide exposure [Fujii et al., 2010]. Therefore, 8-nitro-cGMP conceivably participates in an antioxidant signaling pathway involved in cytoprotection or adaptive responses to ROS and oxidative stress.







Fig. 2. 8-Nitro-cGMP-mediated cytoprotection as a result of S-guanylation of Keap1. S-Guanylation of Keap1 facilitates dissociation of Nrf2 from Keap1 and leads to translocation of Nrf2 to the nucleus. In the nucleus, Nrf2 forms a heterodimer with sMaf proteins, thereby inducing expression of various cytoprotective enzymes.

3. Genetic regulation of the Keap1/Nrf2 system in cancer cells

Hypermethylation of the promoter CpG island is an epigenetic mechanism that leads to gene silencing [Warnecke & Bestor, 2000]. Quite recently, promoter methylation of the *KEAP1* gene was found in malignant gliomas [Muscarella et al., 2011]. A strong inverse correlation was discovered between methylation levels and *KEAP1* mRNA transcript in tumor tissue [Muscarella et al., 2011], which suggests the occurrence of *KEAP1* gene

silencing in gliomas. The reduction in *KEAP1* expression caused by promoter methylation may contribute to constitutive Nrf2 stabilization and subsequent activation of an antioxidant response in gliomas, as seen in certain cancer cells such as those in the lung [Wang et al., 2008] and prostate [P. Zhang et al., 2010]. Inactivation of Keap1 functions by somatic mutations causing amino acid substitutions has been found in cancer cells in the breast, lung, gallbladder, and liver [Shibata et al., 2008a; Singh et al., 2006; Taguchi et al., 2011]. Also, mutations in the *NRF2* gene that are related to the reduction in Keap1-Nrf2 interaction were identified in human cancers of the lung, head and neck, and esophagus [Shibata et al., 2008b; Taguchi et al., 2011]. Such mutations in the *KEAP1* and *NRF2* genes, however, have not yet been found in gliomas and hence warrant further investigation.

#### **4. HO-1 as a target for chemosensitization: The therapeutic potential of HO inhibitors**

Many antitumor agents, such as cisplatin, doxorubicin, mitomycin C, vinblastine, and arsenic trioxide, produce cytotoxic effects via generation of ROS and/or electrophilic actions, which lead to oxidative stress [Dilda & Hogg, 2007; Fang et al., 2007; Simizu et al., 1998]. Hence, those antitumor agents activate Nrf2-dependent HO-1 induction, either by direct actions or by indirect means, i.e., by augmenting the intrinsic signaling cascade mediated by endogenous inducers including 8-nitro-cGMP, as mentioned above. Tumor cells subsequently become resistant to treatment with those antitumor agents. That the HO-1-dependent antioxidant system would be a potential target for chemosensitization is therefore conceivable. The following discussion will cover the effects of disrupting the HO-1-dependent antioxidant system on chemosensitivity, with particular attention to HO inhibitors as new chemosensitizers.

Knockdown of HO-1 or its master regulator Nrf2 is a straightforward approach to disrupting the HO-1 antioxidant system. We found that knockdown of HO-1 by means of small interfering RNA (siRNA) resulted in the induction of apoptosis in SW480 human colon cancer cells in culture [Fang et al., 2003]. Liu et al. [2011] reported that Nrf2 knockdown effectively led to suppressed HO-1 expression and enhanced sensitization to arsenic trioxide-induced apoptosis in cultured human glioma cells U251MG and A172. Quite recently, HO-1 expression was found to be regulated post-transcriptionally by microRNA (miRNA). Beckman et al. [2011], by using an HO-1 reporter assay, found that a combination of two miRNAs (miR-377 and miR-217) markedly reduced HO-1 expression. Mature miRNAs are ~21-22 nucleotides long and affect post-translational expression of genes by interacting with complementary target sites in the 3'-untranslated region of the mRNA [Jackson & Standart, 2007]. The exact molecular mechanisms used by miRNAs to mediate translational repression are still being studied intensively; however, modulation of HO-1 activity by miRNAs may be an alternative strategy for sensitizing tumor cells to chemotherapy.

Direct inhibition of HO-1 enzyme activity may be an alternative approach for disruption of the HO-1-mediated antioxidant system in tumor cells. Metalloporphyrins are a class of compounds in which various metals such as cobalt, zinc, manganese, chromium, or tin replace the central iron in heme [Drummond, 1987]. These metalloporphyrins can act as competitive inhibitors of the HO reaction because they bind inefficiently to molecular oxygen, so HO is prevented from degrading the metalloporphyrins [Drummond, 1987].

Antitumor and chemosensitizing effects of metalloporphyrin-type HO inhibitors such as zinc protoporphyrin (ZnPP) and cobalt protoporphyrin have been demonstrated in cultured tumor cells and in animal models [Doi et al., 1999; Fang et al., 2003, 2004b; Liu et al., 2011; Tanaka et al., 2003]. Combining chemotherapeutic agents such as camptothecin, doxorubicin, arsenic trioxide, gemcitabine, and imatinib, as well as irradiation, with ZnPP or related inhibitors may lead to improved antitumor treatment [Fang et al., 2004b; Gleixner et al., 2009; Liu et al., 2011; Mayerhofer et al., 2008; Miyake et al., 2010].

Polyethylene glycol-conjugated ZnPP (PEG-ZnPP) is a water-soluble derivative of ZnPP (Fig. 3) [Sahoo et al., 2002]. PEG-ZnPP accumulates in solid tumor tissues after intravenous injection because of its unique ability to form micelles in aqueous media [Fang et al., 2003; Sahoo et al., 2002]. As evidenced by accumulation of PEG-ZnPP in tumors, tumor-targeted inhibition of HO activity may be achieved by using PEG-ZnPP, which induces apoptosis in solid tumors via induction of oxidative stress [Fang et al., 2003]. Potent synergistic activity was also observed when PEG-ZnPP was combined with antitumor agents [Fang et al., 2004b]. This combined treatment was tolerated well, with no evident signs of severe toxicity and no dose-limiting toxicity so far. These observations point to new applications of PEG-ZnPP in combination therapy with various anticancer agents as well as irradiation.





Fig. 3. Chemical structures of HO inhibitors used in combination with anticancer agents to treat tumors.

In addition to the metalloporphyrin-type HO inhibitors, non-porphyrin-based HO inhibitors have been developed (Fig. 3). Azalanstat, an imidazole-dioxolane compound, was produced as such an HO inhibitor [Vlahakis et al., 2005]. With azalanstat as the lead compound, so far



more than 100 imidazole-based compounds have been synthesized and their HO inhibitory activities evaluated [Kinobe et al., 2006; Roman et al., 2007; Vlahakis et al., 2005]. Some of these compounds are quite selective for HO-1, with selectivity indices greater than 300 relative to HO-2, a constitutive isoform of HO [Kinobe et al., 2008]. One recent study showed that the imidazole-based compound OB-24 exhibited potent antitumor activity in a hormone-refractory prostate cancer model, especially in combination with paclitaxel [Alaoui-Jamali et al., 2009], which suggests that additional investigations of this type of HO inhibitor and possible uses for tumor chemosensitization are warranted.

### Chapter summary

- Drug resistance is a major cause of treatment failure in patients with malignant gliomas.
- Multiple mechanisms may be involved in the development of drug resistance in gliomas, including augmented DNA repair enzyme activities, overexpression of antiapoptotic proteins and ABC transporters, whereas detailed mechanisms remain unknown.
- Heme oxygenase-1 (HO-1), a rate-limiting enzyme for heme degradation, may contribute to the development of chemoresistance in gliomas, via inducing gene expression of antioxidant, antiapoptotic, cytoprotective enzymes and drug transporters.
- Nitric oxide, reactive oxygen species, and their down stream electrophilic second messengers such as nitrated cyclic nucleotide play crucial roles in HO-1 expression via activation of redox-sensitive transcriptional factor (Nrf2).
- HO inhibitors exhibit potential to sensitize tumor cells against chemotherapy.

### 5. Conclusion

This chapter describes induction mechanisms of the antioxidant adaptive response in gliomas. Nrf2 stabilization may be induced by post-translational cysteine modification of Keap1, which is a repressor of Nrf2, under conditions of excess production of NO and ROS. 8-Nitro-cGMP is a newly discovered second messenger for NO and ROS signaling that can induce S-guanylation of Keap1 in glioma cells. *KEAP1* gene silencing via promoter methylation is an alternative mechanism that may contribute to constitutive Nrf2 stabilization. Activation of Nrf2 signaling leads to the expression of genes encoding phase II detoxifying or antioxidant enzymes as well as other cytoprotective proteins. Among these enzymes, HO-1 is known as a strong antioxidant and antiapoptotic enzyme that protects cells from oxidative stress-related injury. Many studies have shown that inhibition of HO-1 activity, by using either siRNA or HO-1 inhibitors, is a promising strategy for chemosensitization of cancer cells. In addition to direct antioxidant actions of HO-1, which support cancer cell growth as discussed in this chapter, HO-1 expression may contribute to the growth of gliomas by facilitating neovascularization via a vascular endothelial growth factor-dependent mechanism [Morita et al., 2009; Nishie et al., 1999]. HO-1 expression is also correlated with progression of gliomas, possibly through FoxP3-mediated T cell immune suppression [Deininger et al., 2000; El Andaloussi & Lesniak, 2007]. Improved understanding of the mechanisms mediating the Keap1/Nrf2-HO-1 antioxidant response and regulation by 8-nitro-cGMP and its downstream signaling cascade is necessary for development of chemotherapeutic drugs that will target malignant gliomas.

## 6. Acknowledgment

We thank J.B. Gandy for excellent editing of the manuscript.

## 7. References

- Ahmed, K.A., Sawa, T., Ihara, H., Kasamatsu, S., Yoshitake, J., Rahaman, M.M., Okamoto, T., Fujii, S. & Akaike, T. (2011a). Regulation by mitochondrial superoxide and NADPH oxidase of cellular formation of nitrated cyclic GMP: potential implications for ROS signaling. *Biochem. J.*, in press.
- Ahmed, K.A., Sawa, T. & Akaike, T. (2011b). Protein cysteine S-guanylation and electrophilic signal transduction by endogenous nitro-nucleotides. *Amino Acids*, Vol. 162, No. 8, pp. 123-130.
- Akaike, T., Fujii, S., Sawa, T. & Ihara, H. (2010). Cell signaling mediated by nitrated cyclic guanine nucleotide. *Nitric Oxide*, Vol. 23, No. 3 pp. 166-174.
- Akaike, T., Okamoto, S., Sawa, T., Yoshitake, J., Tamura, F., Ichimori, K., Miyazaki, K., Sasamoto, K. & Maeda, H. (2003). 8-Nitroguanosine formation in viral pneumonia and its implication for pathogenesis. *Proc. Natl. Acad. Sci. USA*, Vol. 100, No. 2, pp. 685-690.
- Alaoui-Jamali, M.A., Bismar, T.A., Gupta, A., Szarek, W.A., Su, J., Song, W., Xu, Y., Xu, B., Liu, G., Vlahakis, J.Z., Roman, G., Jiao, J. & Schipper, H.M. (2009). A novel experimental heme oxygenase-1-targeted therapy for hormone-refractory prostate cancer. *Cancer Res.*, Vol. 69, No. 20, pp. 8017-8024.
- Balla, G., Jacob, H.S., Balla, J., Rosenberg, M., Nath, K., Apple, F., Eaton, J.W. & Vercellotti, G.M. (1992). Ferritin: a cytoprotective antioxidant strategem of endothelium. *J. Biol. Chem.*, Vol. 267, No. 25, pp. 18148-18153.
- Baranano, D.E., Rao, M., Ferris, C.D. & Snyder, S.H. (2002). Biliverdin reductase: a major physiologic cytoprotectant. *Proc. Natl. Acad. Sci. USA*, Vol. 99, No. 25, pp. 16093-16098.
- Beckman, J.D., Chen, C., Nguyen, J., Thayanithy, V., Subramanian, S., Steer, C.J. & Vercellotti, G.M. (2011). Regulation of heme oxygenase-1 protein expression by miR-377 in combination with miR-217. *J. Biol. Chem.*, Vol. 286, No. 5, pp. 3194-3202.
- Bleau, A.M., Huse, J.T. & Holland, E.C. (2009). The ABCG2 resistance network of glioblastoma. *Cell Cycle* 8; 2936-44.
- Bloom, D.A. & Jaiswal, A.K. (2003). Phosphorylation of Nrf2 at Ser<sup>40</sup> by protein kinase C in response to antioxidants leads to the release of Nrf2 from INrf2, but is not required for Nrf2 stabilization/accumulation in the nucleus and transcriptional activation of antioxidant response element-mediated NAD(P)H:quinone oxidoreductase-1 gene expression. *J. Biol. Chem.*, Vol. 278, No. 45, pp. 44675-44682.
- Brouard, S., Otterbein, L.E., Anrather, J., Tobiasch, E., Bach, F.H., Choi, A.M. & Soares, M.P. (2000). Carbon monoxide generated by heme oxygenase 1 suppresses endothelial cell apoptosis. *J. Exp. Med.*, Vol. 192, No. 7, pp. 1015-1026.
- Chen, T.J., Jeng, J.Y., Lin, C.W., Wu, C.Y. & Chen, Y.C. (2006). Quercetin inhibition of ROS-dependent and -independent apoptosis in rat glioma C6 cells. *Toxicology*, Vol. 223, No. 1-2, pp. 113-126.

- Chen, Y.C., Chow, J.M., Lin, C.W., Wu, C.Y. & Shen, S.C. (2006). Baicalein inhibition of oxidative-stress-induced apoptosis via modulation of ERKs activation and induction of HO-1 gene expression in rat glioma cells C6. *Toxicol. Appl. Pharmacol.*, Vol. 216, No. 2, pp. 263-273.
- Deininger, M.H., Meyermann, R., Trautmann, K., Duffner, F., Grote, E.H., Wickboldt, J. & Schluesener, H.J. (2000). Heme oxygenase (HO)-1 expressing macrophages/microglial cells accumulate during oligodendroglioma progression. *Brain Res.*, Vol. 882, No. 1-2, pp. 1-8.
- Dilda, P.J. & Hogg, P.J. (2007). Arsenical-based cancer drugs. *Cancer Treat. Rev.*, Vol. 33, No. 6, pp. 542-564.
- Dinkova-Kostova, A.T., Holtzclaw, W.D. & Kensler, T.W. (2005). The role of Keap1 in cellular protective responses. *Chem. Res. Toxicol.*, Vol. 18, No. 12, pp. 1779-1791.
- Doi, K., Akaike, T., Fujii, S., Tanaka, S., Ikebe, N., Beppu, T., Shibahara, S., Ogawa, M. & Maeda, H. (1999). Induction of haem oxygenase-1 nitric oxide and ischaemia in experimental solid tumours and implications for tumour growth. *Br. J. Cancer.*, Vol. 80, No. 12, pp. 1945-1954.
- Drummond, G.S. (1987). Control of heme metabolism by synthetic metalloporphyrins. *Ann. NY Acad. Sci.*, Vol. 514, pp. 87-95.
- Eggler, A.L., Liu, G., Pezzuto, J.M., van Breemen, R.B. & Mesecar, A.D. (2005). Modifying specific cysteines of the electrophile-sensing human Keap1 protein is insufficient to disrupt binding to the Nrf2 domain Neh2. *Proc. Natl. Acad. Sci. USA*, Vol. 102, No. 29, pp. 10070-10075.
- Eggler, A.L., Luo, Y., van Breemen, R.B. & Mesecar, A.D. (2007). Identification of the highly reactive cysteine 151 in the chemopreventive agent-sensor Keap1 protein is method-dependent. *Chem. Res. Toxicol.*, Vol. 20, No. 12, pp. 1878-1884.
- El Andaloussi, A. & Lesniak, M.S. (2007). CD4<sup>+</sup>CD25<sup>+</sup>FoxP3<sup>+</sup> T-cell infiltration and heme oxygenase-1 expression correlate with tumor grade in human gliomas. *J. Neurooncol.*, Vol. 83, No. 2, pp. 145-152.
- Fang, J., Akaike, T. & Maeda, H. (2004a). Antiapoptotic role of heme oxygenase (HO) and the potential of HO as a target in anticancer treatment. *Apoptosis*, Vol. 9, No. 1, pp. 27-35.
- Fang, J., Nakamura, H. & Iyer, A.K. (2007). Tumor-targeted induction of oxystress for cancer therapy. *J. Drug Target.*, Vol. 15, No. 7-8, pp. 475-486.
- Fang, J., Sawa, T., Akaike, T., Akuta, T., Sahoo, S.K., Khaled, G., Hamada, A. & Maeda, H. (2003). In vivo antitumor activity of pegylated zinc protoporphyrin: targeted inhibition of heme oxygenase in solid tumor. *Cancer Res.*, Vol. 63, No. 13, pp. 3567-3574.
- Fang, J., Sawa, T., Akaike, T., Greish, K. & Maeda, H. (2004b). Enhancement of chemotherapeutic response of tumor cells by a heme oxygenase inhibitor, pegylated zinc protoporphyrin. *Int. J. Cancer.*, Vol. 109, No. 1, pp. 1-8.
- Foresti, R., Clark, J.E., Green, C.J. & Motterlini, R. (1997). Thiol compounds interact with nitric oxide in regulating heme oxygenase-1 induction in endothelial cells. Involvement of superoxide and peroxynitrite anions. *J. Biol. Chem.*, Vol. 272, No. 29, pp. 18411-18417.

- Frosina, G. (2009). DNA repair and resistance of gliomas to chemotherapy and radiotherapy. *Mol Cancer Res* 7; 989-99.
- Fujii, S., Sawa, T., Ihara, H., Tong, K.I., Ida, T., Okamoto, T., Ahtesham, A.K., Ishima, Y., Motohashi, H., Yamamoto, M. & Akaike, T. (2010). The critical role of nitric oxide signaling, via protein S-guanylation and nitrated cyclic GMP, in the antioxidant adaptive response. *J. Biol. Chem.*, Vol. 285, No. 31, pp. 23970-23984.
- Gleixner, K.V., Mayerhofer, M., Vales, A., Gruze, A., Hormann, G., Cerny-Reiterer, S., Lackner, E., Hadzijusufovic, E., Herrmann, H., Iyer, A.K., Krauth, M.T., Pickl, W.F., Marian, B., Panzer-Grumayer, R., Sillaber, C., Maeda, H., Zielinski, C. & Valent, P. (2009). Targeting of Hsp32 in solid tumors and leukemias: a novel approach to optimize anticancer therapy. *Curr. Cancer Drug Targets*, Vol. 9, No. 5, pp. 675-689.
- Hara, E., Takahashi, K., Tominaga, T., Kumabe, T., Kayama, T., Suzuki, H., Fujita, H., Yoshimoto, T., Shirato, K. & Shibahara, S. (1996). Expression of heme oxygenase and inducible nitric oxide synthase mRNA in human brain tumors. *Biochem. Biophys. Res. Commun.*, Vol. 224, No. 1, pp. 153-158.
- Ihara, H., Ahmed, K.A., Ida, T., Kasamatsu, S., Kunieda, K., Okamoto, T., Sawa, T. & Akaike, T. (2011a). Methodological proof of immunocytochemistry for specific identification of 8-nitroguanosine 3',5'-cyclic monophosphate formed in glia cells. *Nitric Oxide* 25; 169-75.
- Ihara, H., Sawa, T., Nakabeppu, Y. & Akaike, T. (2011b). Nucleotides function as endogenous chemical sensors for oxidative stress signaling. *J. Clin. Biochem. Nutr.*, Vol. 48, No. 1, pp. 33-39.
- Itoh, K., Tong, K.I. & Yamamoto, M. (2004). Molecular mechanism activating Nrf2-Keap1 pathway in regulation of adaptive response to electrophiles. *Free Radic. Biol. Med.*, Vol. 36, No. 10, pp. 1208-1213.
- Jackson, R.J. & Standart, N. (2007). How do microRNAs regulate gene expression? *Sci. STKE*, Vol. 2007, No. 367, p. re1.
- Jeney, V., Balla, J., Yachie, A., Varga, Z., Vercellotti, G.M., Eaton, J.W. & Balla, G. (2002). Pro-oxidant and cytotoxic effects of circulating heme. *Blood*, Vol. 100, No. 3, pp. 879-887.
- Keyse, S.M. & Tyrrell, R.M. (1989). Heme oxygenase is the major 32-kDa stress protein induced in human skin fibroblasts by UVA radiation, hydrogen peroxide, and sodium arsenite. *Proc. Natl. Acad. Sci. USA*, Vol. 86, No. 1, pp. 99-103.
- Kinobe, R.T., Dercho, R.A. & Nakatsu, K. (2008). Inhibitors of the heme oxygenase-carbon monoxide system: on the doorstep of the clinic? *Can. J. Physiol. Pharmacol.*, Vol. 86, No. 9, pp. 577-599.
- Kinobe, R.T., Vlahakis, J.Z., Vreman, H.J., Stevenson, D.K., Brien, J.F., Szarek, W.A. & Nakatsu, K. (2006). Selectivity of imidazole-dioxolane compounds for in vitro inhibition of microsomal haem oxygenase isoforms. *Br. J. Pharmacol.*, Vol. 147, No. 3, pp. 307-315.
- Kobayashi, A., Kang, M.I., Okawa, H., Ohtsuji, M., Zenke, Y., Chiba, T., Igarashi, K. & Yamamoto, M. (2004). Oxidative stress sensor Keap1 functions as an adaptor for



- Cul3-based E3 ligase to regulate proteasomal degradation of Nrf2. *Mol. Cell. Biol.*, Vol. 24, No. 16, pp. 7130-7139.
- Kobayashi, M., Itoh, K., Suzuki, T., Osanai, H., Nishikawa, K., Katoh, Y., Takagi, Y. & Yamamoto, M. (2002). Identification of the interactive interface and phylogenetic conservation of the Nrf2-Keap1 system. *Genes Cells*, Vol. 7, No. 8, pp. 807-820.
- Konorev, E.A., Kotamraju, S., Zhao, H., Kalivendi, S., Joseph, J. & Kalyanaraman, B. (2002). Paradoxical effects of metalloporphyrins on doxorubicin-induced apoptosis: scavenging of reactive oxygen species versus induction of heme oxygenase-1. *Free Radic. Biol. Med.*, Vol. 33, No. 7, pp. 988-997.
- Krakstad, C. & Chekenya, M. (2010). Survival signalling and apoptosis resistance in glioblastomas: opportunities for targeted therapeutics. *Mol. Cancer*, Vol. 9, p. 135.
- Kurauchi, Y., Hisatsune, A., Isohama, Y., Sawa, T., Akaike, T., Shudo, K. & Katsuki, H. (2011). Midbrain dopaminergic neurons utilize nitric oxide/cyclic GMP signaling to recruit ERK that links retinoic acid receptor stimulation to up-regulation of BDNF. *J. Neurochem.*, Vol. 116, No. 3, pp. 323-333.
- Lee, T.S. & Chau, L.Y. (2002). Heme oxygenase-1 mediates the anti-inflammatory effect of interleukin-10 in mice. *Nat. Med.*, Vol. 8, No. 3, pp. 240-246.
- Liu, Y., Liang, Y., Zheng, T., Yang, G., Zhang, X., Sun, Z., Shi, C. & Zhao, S. (2011). Inhibition of heme oxygenase-1 enhances anti-cancer effects of arsenic trioxide on glioma cells. *J. Neurooncol*, Vol. 104, No. 2, pp. 449-458.
- Lo, S.C., Li, X., Henzl, M.T., Beamer, L.J. & Hannink, M. (2006). Structure of the Keap1:Nrf2 interface provides mechanistic insight into Nrf2 signaling. *EMBO J.*, Vol. 25, No. 15, pp. 3605-3617.
- Lu, C. & Shervington, A. (2008). Chemoresistance in gliomas. *Mol Cell Biochem* 312; 71-80.
- Maines, M.D. (1988). Heme oxygenase: function, multiplicity, regulatory mechanisms, and clinical applications. *FASEB J.*, Vol. 2, No. 10, pp. 2557-2568.
- Mayerhofer, M., Gleixner, K.V., Mayerhofer, J., Hoermann, G., Jaeger, E., Aichberger, K.J., Ott, R.G., Greish, K., Nakamura, H., Derdak, S., Samorapoompichit, P., Pickl, W.F., Sexl, V., Esterbauer, H., Schwarzing, I., Sillaber, C., Maeda, H. & Valent, P. (2008). Targeting of heat shock protein 32 (Hsp32)/heme oxygenase-1 (HO-1) in leukemic cells in chronic myeloid leukemia: a novel approach to overcome resistance against imatinib. *Blood*, Vol. 111, No. 4, pp. 2200-2210.
- Mitani, K., Fujita, H., Fukuda, Y., Kappas, A. & Sassa, S. (1993). The role of inorganic metals and metalloporphyrins in the induction of haem oxygenase and heat-shock protein 70 in human hepatoma cells. *Biochem. J.*, Vol. 290 (Pt 3), pp. 819-825.
- Miyake, M., Fujimoto, K., Anai, S., Ohnishi, S., Nakai, Y., Inoue, T., Matsumura, Y., Tomioka, A., Ikeda, T., Okajima, E., Tanaka, N. & Hirao, Y. (2010). Inhibition of heme oxygenase-1 enhances the cytotoxic effect of gemcitabine in urothelial cancer cells. *Anticancer Res.*, Vol. 30, No. 6, pp. 2145-2152.
- Morita, K., Lee, M.S. & Her, S. (2009). Possible relation of hemin-induced HO-1 expression to the upregulation of VEGF and BDNF mRNA levels in rat C6 glioma cells. *J. Mol. Neurosci.*, Vol. 38, No. 1, pp. 31-40.




- Motohashi, H. & Yamamoto, M. (2007). Carcinogenesis and transcriptional regulation through Maf recognition elements. *Cancer Sci.*, Vol. 98, No. 2, pp. 135-139.
- Motterlini, R., Foresti, R., Bassi, R., Calabrese, V., Clark, J.E. & Green, C.J. (2000). Endothelial heme oxygenase-1 induction by hypoxia. Modulation by inducible nitric-oxide synthase and S-nitrosothiols. *J. Biol. Chem.*, Vol. 275, No. 18, pp. 13613-13620.
- Muscarella, L.A., Barbano, R., D'Angelo, V., Copetti, M., Coco, M., Balsamo, T., la Torre, A., Notarangelo, A., Troiano, M., Parisi, S., Icolaro, N., Catapano, D., Valori, V.M., Pellegrini, F., Merla, G., Carella, M., Fazio, V.M. & Parrella, P. (2011). Regulation of KEAP1 expression by promoter methylation in malignant gliomas and association with patient's outcome. *Epigenetics*, Vol. 6, No. 3, pp. 317-325.
- Nguyen, T., Sherratt, P.J. & Pickett, C.B. (2003). Regulatory mechanisms controlling gene expression mediated by the antioxidant response element. *Annu. Rev. Pharmacol. Toxicol.*, Vol. 43, pp. 233-260.
- Nishie, A., Ono, M., Shono, T., Fukushima, J., Otsubo, M., Onoue, H., Ito, Y., Inamura, T., Ikezaki, K., Fukui, M., Iwaki, T. & Kuwano, M. (1999). Macrophage infiltration and heme oxygenase-1 expression correlate with angiogenesis in human gliomas. *Clin. Cancer Res.*, Vol. 5, No. 5, pp. 1107-1113.
- Ogura, T., Tong, K.I., Mio, K., Maruyama, Y., Kurokawa, H., Sato, C. & Yamamoto, M. (2010). Keap1 is a forked-stem dimer structure with two large spheres enclosing the intervening, double glycine repeat, and C-terminal domains. *Proc. Natl. Acad. Sci. USA*, Vol. 107, No. 7, pp. 2842-2847.
- Padmanabhan, B., Tong, K.I., Ohta, T., Nakamura, Y., Scharlock, M., Ohtsuji, M., Kang, M.I., Kobayashi, A., Yokoyama, S. & Yamamoto, M. (2006). Structural basis for defects of Keap1 activity provoked by its point mutations in lung cancer. *Mol. Cell*, Vol. 21, No. 5, pp. 689-700.
- Prados, M.D., Lamborn, K.R., Chang, S., Burton, E., Butowski, N., Malec, M., Kapadia, A., Rabbitt, J., Page, M.S., Fedoroff, A., Xie, D. & Kelley, S.K. (2006). Phase 1 study of erlotinib HCl alone and combined with temozolomide in patients with stable or recurrent malignant glioma. *Neuro-oncol.*, Vol. 8, No. 1, pp. 67-78.
- Rich, J.N., Reardon, D.A., Peery, T., Dowell, J.M., Quinn, J.A., Penne, K.L., Wikstrand, C.J., Van Duyn, L.B., Dancey, J.E., McLendon, R.E., Kao, J.C., Stenzel, T.T., Ahmed Rasheed, B.K., Tourt-Uhlig, S.E., Herndon, J.E., 2nd, Vredenburgh, J.J., Sampson, J.H., Friedman, A.H., Bigner, D.D. & Friedman, H.S. (2004). Phase II trial of gefitinib in recurrent glioblastoma. *J. Clin. Oncol.*, Vol. 22, No. 1, pp. 133-142.
- Roman, G., Riley, J.G., Vlahakis, J.Z., Kinobe, R.T., Brien, J.F., Nakatsu, K. & Szarek, W.A. (2007). Heme oxygenase inhibition by 2-oxy-substituted 1-(1H-imidazol-1-yl)-4-phenylbutanes: effect of halogen substitution in the phenyl ring. *Bioorg. Med. Chem.*, Vol. 15, No. 9, pp. 3225-3234.
- Sahoo, S.K., Sawa, T., Fang, J., Tanaka, S., Miyamoto, Y., Akaike, T. & Maeda, H. (2002). Pegylated zinc protoporphyrin: a water-soluble heme oxygenase inhibitor with tumor-targeting capacity. *Bioconjug. Chem.*, Vol. 13, No. 5, pp. 1031-1038.

- Saito, Y., Taguchi, H., Fujii, S., Sawa, T., Kida, E., Kabuto, C., Akaike, T. & Arimoto, H. (2008). 8-Nitroguanosines as chemical probes of the protein S-guanylation. *Chem. Commun. (Camb.)*, No. 45, pp. 5984-5986.
- Sarkaria, J.N., Kitange, G.J., James, C.D., Plummer, R., Calvert, H., Weller, M. & Wick, W. (2008). Mechanisms of chemoresistance to alkylating agents in malignant glioma. *Clin Cancer Res* 14; 2900-8.
- Sawa, T., Arimoto, H. & Akaike, T. (2010). Regulation of redox signaling involving chemical conjugation of protein thiols by nitric oxide and electrophiles. *Bioconjug. Chem.*, Vol. 21, No. 7, pp. 1121-1129.
- Sawa, T., Ihara, H. & Akaike, T. (2011). Antioxidant effect of a nitrated cyclic nucleotide functioning as an endogenous electrophile. *Curr. Top. Med. Chem.*, in press.
- Sawa, T., Tatemichi, M., Akaike, T., Barbin, A. & Ohshima, H. (2006). Analysis of urinary 8-nitroguanine, a marker of nitrative nucleic acid damage, by high-performance liquid chromatography-electrochemical detection coupled with immunoaffinity purification: association with cigarette smoking. *Free Radic. Biol. Med.*, Vol. 40, No. 4, pp. 711-720.
- Sawa, T., Zaki, M.H., Okamoto, T., Akuta, T., Tokutomi, Y., Kim-Mitsuyama, S., Ihara, H., Kobayashi, A., Yamamoto, M., Fujii, S., Arimoto, H. & Akaike, T. (2007). Protein S-guanylation by the biological signal 8-nitroguanosine 3',5'-cyclic monophosphate. *Nat. Chem. Biol.*, Vol. 3, No. 11, pp. 727-735.
- Schacter, B.A. (1988). Heme catabolism by heme oxygenase: physiology, regulation, and mechanism of action. *Semin. Hematol.*, Vol. 25, No. 4, pp. 349-369.
- Sekhar, K.R., Rachakonda, G. & Freeman, M.L. (2010). Cysteine-based regulation of the CUL3 adaptor protein Keap1. *Toxicol. Appl. Pharmacol.*, Vol. 244, No. 1, pp. 21-26.
- Shibata, T., Kokubu, A., Gotoh, M., Ojima, H., Ohta, T., Yamamoto, M. & Hirohashi, S. (2008a). Genetic alteration of Keap1 confers constitutive Nrf2 activation and resistance to chemotherapy in gallbladder cancer. *Gastroenterology*, Vol. 135, No. 4, pp. 1358-1368.e1-4.
- Shibata, T., Ohta, T., Tong, K.I., Kokubu, A., Odogawa, R., Tsuta, K., Asamura, H., Yamamoto, M. & Hirohashi, S. (2008b). Cancer related mutations in *NRF2* impair its recognition by Keap1-Cul3 E3 ligase and promote malignancy. *Proc. Natl. Acad. Sci. USA*, Vol. 105, No. 36, pp. 13568-13573.
- Simizu, S., Takada, M., Umezawa, K. & Imoto, M. (1998). Requirement of caspase-3(-like) protease-mediated hydrogen peroxide production for apoptosis induced by various anticancer drugs. *J. Biol. Chem.*, Vol. 273, No. 41, pp. 26900-26907.
- Singh, A., Misra, V., Thimmulappa, R.K., Lee, H., Ames, S., Hoque, M.O., Herman, J.G., Baylin, S.B., Sidransky, D., Gabrielson, E., Brock, M.V. & Biswal, S. (2006). Dysfunctional KEAP1-NRF2 interaction in non-small-cell lung cancer. *PLoS Med.*, Vol. 3, No. 10, p. e420.
- Stewart, B.W. & Kleihues, P. (Eds.). (2003). *World Cancer Report*, IARC Press, ISBN, Lyon.
- Stupp, R., Mason, W.P., van den Bent, M.J., Weller, M., Fisher, B., Taphoorn, M.J., Belanger, K., Brandes, A.A., Marosi, C., Bogdahn, U., Curschmann, J., Janzer,

- R.C., Ludwin, S.K., Gorlia, T., Allgeier, A., Lacombe, D., Cairncross, J.G., Eisenhauer, E. & Mirimanoff, R.O. (2005). Radiotherapy plus concomitant and adjuvant temozolomide for glioblastoma. *N. Engl. J. Med.*, Vol. 352, No. 10, pp. 987-996.
- Surh, Y.J., Kundu, J.K., Li, M.H., Na, H.K. & Cha, Y.N. (2009). Role of Nrf2-mediated heme oxygenase-1 upregulation in adaptive survival response to nitrosative stress. *Arch. Pharm. Res.*, Vol. 32, No. 8, pp. 1163-1176.
- Taguchi, K., Motohashi, H. & Yamamoto, M. (2011). Molecular mechanisms of the Keap1-Nrf2 pathway in stress response and cancer evolution. *Genes Cells*, Vol. 16, No. 2, pp. 123-140.
- Tanaka, S., Akaike, T., Fang, J., Beppu, T., Ogawa, M., Tamura, F., Miyamoto, Y. & Maeda, H. (2003). Antiapoptotic effect of haem oxygenase-1 induced by nitric oxide in experimental solid tumour. *Br. J. Cancer*, Vol. 88, No. 6, pp. 902-909.
- Terasaki, Y., Akuta, T., Terasaki, M., Sawa, T., Mori, T., Okamoto, T., Ozaki, M., Takeya, M. & Akaike, T. (2006). Guanine nitration in idiopathic pulmonary fibrosis and its implication for carcinogenesis. *Am. J. Respir. Crit. Care Med.*, Vol. 174, No. 6, pp. 665-673.
- Vlahakis, J.Z., Kinobe, R.T., Bowers, R.J., Brien, J.F., Nakatsu, K. & Szarek, W.A. (2005). Synthesis and evaluation of azalanstat analogues as heme oxygenase inhibitors. *Bioorg. Med. Chem. Lett.*, Vol. 15, No. 5, pp. 1457-1461.
- Wakabayashi, N., Dinkova-Kostova, A.T., Holtzclaw, W.D., Kang, M.I., Kobayashi, A., Yamamoto, M., Kensler, T.W. & Talalay, P. (2004). Protection against electrophile and oxidant stress by induction of the phase 2 response: fate of cysteines of the Keap1 sensor modified by inducers. *Proc. Natl. Acad. Sci. USA*, Vol. 101, No. 7, pp. 2040-2045.
- Wang, R., An, J., Ji, F., Jiao, H., Sun, H. & Zhou, D. (2008). Hypermethylation of the Keap1 gene in human lung cancer cell lines and lung cancer tissues. *Biochem. Biophys. Res. Commun.*, Vol. 373, No. 1, pp. 151-154.
- Warnecke, P.M. & Bestor, T.H. (2000). Cytosine methylation and human cancer. *Curr. Opin. Oncol.*, Vol. 12, No. 1, pp. 68-73.
- Zaki, M.H., Fujii, S., Okamoto, T., Islam, S., Khan, S., Ahmed, K.A., Sawa, T. & Akaike, T. (2009). Cytoprotective function of heme oxygenase 1 induced by a nitrated cyclic nucleotide formed during murine salmonellosis. *J. Immunol.*, Vol. 182, No. 6, pp. 3746-3756.
- Zhang, D.D. & Hannink, M. (2003). Distinct cysteine residues in Keap1 are required for Keap1-dependent ubiquitination of Nrf2 and for stabilization of Nrf2 by chemopreventive agents and oxidative stress. *Mol. Cell. Biol.*, Vol. 23, No. 22, pp. 8137-8151.
- Zhang, P., Singh, A., Yegnasubramanian, S., Esopi, D., Kombairaju, P., Bodas, M., Wu, H., Bova, S.G. & Biswal, S. (2010). Loss of Kelch-like ECH-associated protein 1 function in prostate cancer cells causes chemoresistance and radioresistance and promotes tumor growth. *Mol. Cancer Ther.*, Vol. 9, No. 2, pp. 336-346.
- Zhao, C.R., Gao, Z.H. & Qu, X.J. (2010). Nrf2-ARE signaling pathway and natural products for cancer chemoprevention. *Cancer Epidemiol.*, Vol. 34, No. 5, pp. 523-533.

- Zhu, H., Itoh, K., Yamamoto, M., Zweier, J.L. & Li, Y. (2005). Role of Nrf2 signaling in regulation of antioxidants and phase 2 enzymes in cardiac fibroblasts: protection against reactive oxygen and nitrogen species-induced cell injury. *FEBS Lett.*, Vol. 579, No. 14, pp. 3029-3036.
- Zipper, L.M. & Mulcahy, R.T. (2002). The Keap1 BTB/POZ dimerization function is required to sequester Nrf2 in cytoplasm. *J. Biol. Chem.*, Vol. 277, No. 39, pp. 36544-36552.



## **Novel Therapeutic Concepts in Targeting Glioma**

Edited by Prof. Faris Farassati

ISBN 978-953-51-0491-9

Hard cover, 306 pages

**Publisher** InTech

**Published online** 04, April, 2012

**Published in print edition** April, 2012

Novel Therapeutic Concepts for Targeting Glioma offers a comprehensive collection of current information and the upcoming possibilities for designing new therapies for Glioma by an array of experts ranging from Cell Biologists to Oncologists and Neurosurgeons. A variety of topics cover therapeutic strategies based on Cell Signaling, Gene Therapy, Drug Therapy and Surgical methods providing the reader with a unique opportunity to expand and advance his knowledge of the field.

### **How to reference**

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Tomohiro Sawa and Takaaki Akaike (2012). Antioxidant Adaptive Response of Malignant Glioma Related to Resistance to Antitumor Treatment, Novel Therapeutic Concepts in Targeting Glioma, Prof. Faris Farassati (Ed.), ISBN: 978-953-51-0491-9, InTech, Available from: <http://www.intechopen.com/books/novel-therapeutic-concepts-in-targeting-glioma/antioxidant-adaptive-response-of-malignant-glioma-related-to-resistance-to-antitumor-treatment>

**INTECH**  
open science | open minds

### **InTech Europe**

University Campus STeP Ri  
Slavka Krautzeka 83/A  
51000 Rijeka, Croatia  
Phone: +385 (51) 770 447  
Fax: +385 (51) 686 166  
[www.intechopen.com](http://www.intechopen.com)

### **InTech China**

Unit 405, Office Block, Hotel Equatorial Shanghai  
No.65, Yan An Road (West), Shanghai, 200040, China  
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元  
Phone: +86-21-62489820  
Fax: +86-21-62489821



© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen