

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

185,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Wildlife Forestry

Daniel J. Twedt
 U.S. Geological Survey
 Patuxent Wildlife Research Center Vicksburg, Mississippi
 USA

1. Introduction

Wildlife forestry is management of forest resources, within sites and across landscapes, to provide sustainable, desirable habitat conditions for all forest-dependent (silvicolous) fauna while concurrently yielding economically viable, quality timber products. In practice, however, management decisions associated with wildlife forestry often reflect a desire to provide suitable habitat for rare species, species with declining populations, and exploitable (i.e., game) species. Collectively, these species are deemed priority species and they are assumed to benefit from habitat conditions that result from prescribed silvicultural management actions.

Early wildlife conservation efforts largely focused on controlling indiscriminate slaughter of wildlife by restricting the season or sex of harvested species (Graham, 1947). Subsequent conservation efforts targeted increased protection of populations through creation of sanctuaries or reserves (Knight, 1999), such as national parks and wildlife refuges, within which harvest of wildlife was prohibited or greatly restricted. Typically parks and wildlife reserves were located in areas with abundant wildlife populations, and therefore, little emphasis was placed on active management to improve or maintain suitable habitat.

In his description of a “land ethic” in which people are members of the natural community, Leopold (1949) indicated the appropriateness of habitat alteration and management to aid the continued existence of wildlife species. His philosophical principles were succinctly stated as “*A thing is right when it tends to preserve the integrity, stability, and beauty of the biotic community. It is wrong when it tends otherwise.*”

Anthropogenic alteration of forest habitat for the benefit wildlife species has long been practiced, particularly with regard to intentional use of fire (Bonnicksen, 2000, Ford et al., 2002). Even so, active manipulation of forest structure to purposefully enhance wildlife habitat has been slow to evolve. Most early efforts regarding habitat alteration focused on providing suitable escape cover, foraging habitat, or distinct habitat features (e.g., nest sites) for game species. Within forested landscapes, improving wildlife habitat often meant increasing local heterogeneity and providing more forest edge habitat. However, increased knowledge of the possible negative effects associated with forest edges, such as greater predation and nest parasitism rates, has called into question the practice of providing

increased heterogeneity within forested landscapes. At the same time, a desire to provide habitat for specific species that may require specialized habitat conditions, as is the case for some endangered and threatened species as well as other charismatic megafauna with widespread public support, has prompted development of habitat management prescriptions designed to improve forest habitat conditions for specific species (Wilson et al., 2007).

Partly in response to what has been perceived as management for single-species, but also to advocate a desire to prevent future population declines of common species, recent emphases in wildlife conservation have been placed on the development of comprehensive, eco-regional, conservation plans whose scope encompasses multiple species or the entirety of a species-group. This approach gained momentum with the North American Waterfowl Management Plan (1986) which identified conservation actions intended to return waterfowl populations to their former abundance and distribution. Subsequently, this basic concept has been expanded to other species groups such as songbirds (Rich et al., 2004), reptiles and amphibians (Bailey et al., 2006), and shorebirds (Brown et al., 2001), as well as other species groups. Although these conservation plans differ markedly with regard to their recommendations for habitat management, many of these conservation plans advocate use of prophylactic management prescriptions to enhance or maintain suitable habitat conditions for priority wildlife species. Indeed, most conservation plans recognize the need for alteration of habitat and landscapes via management actions so as to attain the desired distribution, abundance, and viability of priority wildlife species.

Before managers can provide appropriate forest habitat for priority wildlife, the landscape and site characteristics that contribute to viable populations of these species must be identified. Preferably, those forest habitat characteristics identified can be quantified via standard measurement protocols. If possible, the desired or acceptable range of values should be determined, and either quantitatively or qualitatively stated, for each of these characteristics. In addition, a threshold value for each characteristic should be determined, that if exceeded would justify prescription of management actions to return site characteristics within the desired range (Fig. 1).

Because wildlife forestry silviculture does not target optimal production of forest products, many forest management practitioners (i.e., foresters) have been reluctant to adopt these alternative silvicultural practices. Indeed, these concerns may be justified, as desired stand conditions may include less than fully stocked stand densities, retention of some economically mature trees, maintaining species diversity which includes less merchantable tree species, and preservation of less vigorous (decadent) trees. However, at present we lack credible, long-term economic data to suggest that development and maintenance of quality forest habitat for priority wildlife species will result in a sacrifice in timber production or quality. Regardless of these overarching financial concerns, silvicultural management actions prescribed to enhance desirable forest habitat conditions for priority wildlife species are expected to be commercially viable. Maintaining the long-term commercial viability of prescribed silviculture is important, because without sufficient financial returns, needed management actions will not be undertaken.




Fig. 1. Characterization of wildlife forestry silvicultural methodology.

2. Forest management

Silviculture is the manipulation of the establishment, growth, composition, and quality of forest vegetation to achieve management objectives. Historically a primary objective on managed forest lands has been to maximize financial returns from production of forest

products such as timber and wood pulp. Production-oriented silviculture has focused on reducing the length of production cycles between stand regeneration and harvest while concurrently maximizing wood product yields. Reduction of production cycles (i.e., rotation length) tends to lessen the time regenerating forests are in early seral habitats, and consequently the size and age of shrubs and there their likelihood of persistence within subsequent seral stages is reduced (Hagar, 2004). Often production-forestry objectives favored a limited number of tree species due to their high economic value or growth potential. Concomitantly, to increase growth and yield of desired trees, other species with lesser economic value, as well as congeneric competitors, were culled. These actions result in increased spatial homogeneity within managed stands. The apogee of this trend towards increased spatial regularity is expressed within planted plantation forests. Climate, soils, species, and other factors affect the optimal length of production cycles within production-oriented forests (Lindenmayer et al., 1999). Thus, rotation periods within plantations vary markedly from rotations of up to 150 years to as few as 10 years. Notably, recent developments and increased advocacy of short-rotation woody coppice systems for use in biomass production have further compressed the length of production cycles to <4 years (Tubby and Armstrong, 2002).

Even in non-plantation forest management systems, historical forest management practices, especially those focused on timber or pulpwood production, have often resulted in relatively homogeneous mature forest conditions. This homogeneity has been exacerbated by environmental changes that limit natural disturbances within forest stands (e.g., fire suppression). When these homogenous forests are managed using even-aged silviculture, during which all or most trees are harvested and the entire stand regenerated, successional 'boom and bust' conditions are dispersed throughout landscapes. The forest conditions resulting from even-aged management (e.g., clear cuts) are temporarily favorable for some species, specifically benefiting species that exploit early-successional forest habitat (Askins, 2001, Hunter et al., 2001). Even so, these early-successional habitats are transitory and are followed by an extended period of stand development (stem exclusion and understory reinitiation; Oliver and Larson, 1996) during which the homogeneous, closed-canopy forest structure with sparse understory vegetation offers attraction (forage or cover) primarily for common (non-priority) forest wildlife species. Indeed, the abundance of homogeneous, closed canopy forests with sparse understory cover that have resulted from historical silvicultural practices may have contributed to why species for which these habitat conditions are suitable are considered common and are not priority species in need of management actions to increase their abundance. In addition, the altered (often depauperate) species compositions that have resulted from selective harvest and culling during past management practices may amplify the potential for catastrophic damage from outbreaks of insects or diseases or limit the seasonality of mast crops that are exploited by wildlife.

In contrast to the relatively short periodicity of production-oriented silviculture and the periodic boom-bust cycles associated with even-aged silviculture, natural successional processes within forests may require hundreds, or potentially a thousand, years to achieve complete replacement via small-scale disturbance or as the duration between catastrophic stand-replacement events. Owing to differences in their successional development, structural and ecological characteristics within production-oriented managed forests and in forests managed using even-aged silviculture are manifestly different from those same characteristics within natural forests (Seymour and Hunter, 1999). Ecological differences

between managed and natural forest stands are exacerbated over time, as natural stands approach climax or 'old-growth' conditions while intensively managed forests are repeatedly regenerated.

To mitigate the detrimental effects of forest management on priority wildlife species, recommendations have been made to increase the length of production cycles, thereby allowing increased time for development of heterogeneous forest structure (Kerr, 1999). Unfortunately, small increases in rotation lengths provide little opportunity for increased complexity of forest structure within these stands – and periodic, complete harvest and stand regeneration thwart retention of forest structure over time. Similarly, forest managers have argued that temporal and spatial dispersion of clear-cuts throughout landscapes will provide continual availability of wildlife habitat – albeit at different locations. Although early-successional habitat may be continuously afforded via such dispersion, few of the other benefits conferred by increased forest structural heterogeneity, such as emergent or senescent legacy trees, are realized.

Increased recognition of the structural and ecological deficiencies within production-oriented forests has spurred interest in and development of alternative silviculture (Franklin, 1989). Alternative silviculture has been especially valuable for stands where management objectives are not solely financial but where stewardship objectives include preservation of regional biodiversity, enhancement of wildlife habitat, maintaining landscape aesthetics, or providing increased recreational opportunities.

Alternative silvicultural methods developed for use in many forest types have been based on identification and quantification of natural disturbance regimes, with subsequent implementation of silvicultural methods intended to emulate these disturbances (Mitchell et al., 2002, 2006; Palik et al., 2002). For many forest types, emulation of natural disturbance regimes includes small-scale disturbances implemented via single-tree and patch-cut harvests (Franklin et al., 2007, North and Keeton, 2008). If successfully implemented, emulation of natural disturbance regimes should promote natural stand development and succession. Unfortunately, natural stand development and concurrent successional changes may conflict with landowner objectives if succession results in changed species composition or reduced financial returns. Moreover, strict adherence to silvicultural regimes that mimic natural disturbance may not be necessary to maintain biodiversity or to provide desired wildlife habitat (Palik et al., 2002).

Another common alternative silviculture has been inspired by the desire to achieve forest structure reminiscent of old-growth forest (Bauhus et al., 2009). Although the structure of old-growth forest varies among forest types (Hayward, 1991), common attributes generally include increased vertical heterogeneity via a multi-layered canopy, increased horizontal heterogeneity associated with canopy gaps and different regeneration cohorts, the presence of large trees that are often in older-age classes, abundant snags (standing dead trees) or senescent trees, and large diameter downed woody debris (Bauhus et al., 2009, Keeton, 2006). Myriad ecological benefits, including increased biodiversity, have been ascribed to old-growth forests (Lindenmayer and Franklin, 2002). As such, forest reserve lands have been designated (e.g., wilderness areas and parks) which afford unfettered, natural development of these characteristics. However, on lands with timber production as an objective, even if not the sole objective, implementation of production cycles that span hundreds of years is not likely to occur. In addition, merely increasing the length of

production cycles may provide little enhancement of biodiversity if the forest structure favored by production-oriented silviculture is maintained (Carey, 2006). Furthermore, where forest reserves are small or located within inhospitable landscapes, full ecological benefits of old-growth forests may not be attainable or sustainable (Kneeshaw and Gauthier, 2003).

Fortunately, it is likely that many of the benefits conferred by old-growth forests, or at least specific species associated with old-growth forests, are positively related to the heterogeneous structure (vertical and horizontal) and other structural attributes found within these forests, not necessarily with the prolonged existence of these forests (Beggs, 2004, Carey, 2003b). As such, alternative silvicultural methods have been developed which retain or enhance structural heterogeneity within managed forests, thereby promoting structural attributes of old-growth forests and encouraging greater biodiversity (Bauhus et al., 2009, Garman et al., 2003; McClellan, 2004). Most often, increased heterogeneity and other desired attributes have been achieved via retention of living or dead trees (or other forest elements). These retained structures have been referred to as legacy elements (Franklin et al., 1997).

Based on the concept of retained legacy elements, various alternative silvicultural methods have been advocated to enhance structural complexity within managed forests (Whitman and Hagan, 2003). Some of these methods have been designated as green-tree retention harvests (Zenner, 2000), variable-retention harvests (Aubry et al., 2004, Maguire et al., 2007), dispersed retention harvests (Heithecker and Halpern, 2006), variable density thinning (Aukema and Carey, 2008), and active intentional management (Carey et al., 1999, Carey, 2006). The legacy elements retained under these different silvicultural regimes may differ markedly. Retained elements may differ in quantity (e.g., % canopy cover or % basal area retained; Fig. 2a,b) but they may also differ in their composition, as different canopy species are preferentially harvested or specific regenerating species favored for retention. In addition, the quality of retained elements may differ via selection of retained trees with differing perceived fates – from robust, canopy-emergent trees through decadent, dead or dying trees. Moreover, retained elements may differ in their dispersion (e.g., evenly dispersed or aggregated; Fig. 2c,d) and the spatial juxtaposition of retention (e.g., clumped or linear).

Wildlife forestry silviculture exploits the versatility of legacy retention by combining specific retention and removal of canopy to achieve habitat conditions that are deemed suitable for priority wildlife species. The resultant forest stand may exhibit a mixture of tree species, size classes, and decadence within a heterogeneous distribution of aggregated retention and canopy gaps (Fig. 3).

2.1 Landscape considerations

Although specific silvicultural methods have most often been developed for and are implemented within forest management units (a.k.a. forest stands), a similar ecologically based approach has evolved with regard to maintaining or restoring landscapes surrounding managed forest stands (Lindenmayer et al., 2002, Lindenmayer & Hobbs, 2007). Intact forested landscapes are conducive to maintaining floristic and wildlife diversity, but also provide tangible social and economic benefits (Poudyal et al., 2010).




Fig. 2. Potential distribution of legacy elements after application of alternative silviculture, differing in quantity of retained canopy cover (a, b) or dispersion of retained elements (c, d).

Loss of diversity within landscapes is often a reflection of the increased homogeneity within forest stands. Although forest structure within a landscape ranges from regenerating stands to mature merchantable stands, these stands may be monotypic in species composition and physical structure. In addition, ensuring appropriate size of habitat patches and maintaining connectivity among these patches is critical to sustaining viable wildlife populations within landscapes (Franklin and Forman, 1987). Although the appropriate patch size is dependent upon the species under consideration, increased fragmentation tends to disrupt dispersion of fauna and flora and when habitat area is reduced without appropriate connectivity, extirpation of some species can result (MacArthur and Wilson, 1967). Finally, maintaining landscapes capable of sustaining viable wildlife populations requires that evaluation, planning, and management decisions reflect the temporal processes inherent in the ecosystem. Four landscape conditions have been recognized as contributing to ecosystems that support sustainable wildlife populations: 1) diversity within the landscape, 2) connectivity among landscape elements, 3) appropriate habitat patch sizes, and 4) sufficient time to achieve ecological functions (Silva, 1992). The impact of the degradation of forest landscapes was succinctly summarized by Franklin (1989) as: *"In general, we have tended to forget that what is good for wood production is not necessarily good for other organisms or processes in a forest ecosystem. Fully stocked young forests,... are the most simplified stage of forest development in terms of structure and function, and the most impoverished in terms of biological*

diversity.. . . Simplification--genetic, structural, landscape and temporal – reduces ecosystem resilience...[Thus] the key to retaining resilience must be in maintaining ecological complexity or diversity”.




Fig. 3. Canopy cover accounted for by individual tree crowns within bottomland hardwood forests on an untreated control stand (left) and on a stand subjected to wildlife forestry variable-retention silvicultural treatment (right), 6 years after treatment (Twedt & Wilson, 2007).

As when considering disturbance within the context of forest stand development, landscape conditions have often been influenced by large-scale disturbance events, particularly fires (Kaufmann et al., 2003, Van Willgenburg and Hobson, 2008). Forest landscapes that result from management based on historical fire regimes, when compared with landscapes with extensive reserves and traditional silvicultural management, had increased area of late-successional habitat, more overstory structure in young forests, and larger forest patches (Cissel et al., 1999). Similarly, when compared with human altered landscapes, natural landscapes were composed of larger habitat patches which formed the landscape matrix, had greater connectivity among patches, the range of variation in patch size and complexity was greater, as was the likelihood of encountering ‘rare’ habitat types (Mladenoff et al., 1993).

Despite some understanding of desirable characteristics of forested landscapes, the appropriate landscape area remains nebulous. In part this is because an appropriate landscape area is dependent upon the species being considered, landscape characteristics,

desired response, and the management context (Karau and Keane, 2007; Mayer and Cameron, 2003). As an example, the landscape appropriate for a grizzly bear is vastly different from the appropriate landscape for a shrew.

Even with knowledge of the appropriate landscape area and characteristics, manipulation of landscape conditions is usually beyond the purview of individual landowners. This is true even for large industrial or governmental landowners. Nevertheless, landscape conditions influence the ecological benefits conferred by forest stands, including the distribution, abundance, and viability of wildlife species within forest stands. In boreal forests, stand structure accounted for 54% of variability in bird and small mammal communities but landscape characteristics accounted for 29% (St Laurent et al., 2008). Similarly, bird species abundance within coniferous dominated landscapes was positively related to the proportion of broadleaf forest, with landscape thresholds for individual species ranging from 1% to 25% canopy cover. A strong negative relationship was noted between landscape broadleaf forest and avian population trends, such that the species most associated with broadleaf forests had greatest population declines. Therefore, maintaining or restoring broadleaf vegetation is important within coniferous dominated landscapes (Betts et al., 2010).

Because of landscape influences, providing desired forest conditions within forest stands may be insufficient for some priority species if these stands are located within inhospitable landscapes. Moreover, for long-term sustainable populations, most sites, even those with suitable habitat conditions, must have spatial and temporal linkages that facilitate dispersal, colonization, and interchange within metapopulations (Hanski, 1998). Towards that end, desired landscape conditions for priority wildlife species are usually composed of large forest patches as the preponderance of the landscape matrix, have a wide range of variation in patch sizes and complexities, and provide extensive connectivity among forest patches.

2.2 Why wildlife forestry?

The application of wildlife forestry silviculture is in many respects the logical extension of the various alternative silviculture methods whose objectives are to maintain or increase forest biodiversity (Table 1). Although forest management targeting increased biodiversity is a laudable objective in and of itself, managers of public conservation lands (e.g., National Wildlife Refuges and State Wildlife Management Areas) or private lands under conservation easements may have legal, or self-appointed, mandates to manage and conserve priority wildlife species rather than the more general goal of improving biodiversity. On other private lands, the economic and recreational values associated with silvicolous wildlife, including fee hunting, hunt-club memberships, wildlife viewing and other eco-tourism activities, have income potential that may equal or exceed that of extractive timber harvest. Hence, ensuring sustainable populations of wildlife may be essential for maintaining financial revenues from these forest lands.

In addition, land ownership patterns of private forest lands are changing. Changing ownership demographics and associated alteration of management objectives of these forest land owners have resulted in smaller landownership holdings. For owners of small landholdings, maintaining continuous forest cover and the aesthetic considerations associated with forests may be of paramount importance (Kaetzel et al., 2011).

Legacy characteristic	Wildfire	Wind	Insect / disease	Flood	Clearcut	Wildlife forestry
Live trees	Sparse	Variable	Variable	Sparse	Sparse	Variable
Snags	Abundant	Variable	Abundant	Abundant	Sparse	Variable
Downed woody debris	Variable	Abundant	Variable	Variable	Sparse	Variable
Understory development	Abundant	Abundant	Variable	Variable	Abundant	Variable
Spatial heterogeneity	Variable	Variable	High	Variable	Low	High
Duration of altered state	Variable	Variable	Variable	Variable	Long	Variable

Table 1. State of habitat features following different disturbances and prescribed wildlife forestry silviculture (expanded from Swanson et al., 2011).

Even on large land ownerships, such as industrial forest lands or lands managed by timber consortiums, there are increased public expectations to provide sustainable, ‘green’ forest products (Barneycastle, 2001). Concurrently, corporate efforts to maintain positive environmental images have prompted greater reliance on alternative silviculture, such as wildlife forestry. Finally, even on lands where implementation of traditional silviculture may be the preferred management option, exurban encroachment may constrain traditional forest management practices (Egan and Luloff, 2000).

2.3 Priority species

What are ‘priority’ species? For some groups of wildlife species, conservation planners have established priority rankings based on objective criteria such as current population abundance, distributions, availability of suitable habitats, perceived threats to habitat, etc. (Mehlman et al., 2004). For example, for North American landbirds, Partners in Flight has established ecoregional priority rankings (Panjabi et al., 2005). For other wildlife groups, priority status may be conferred based on species being listed as threatened or endangered by national or regional governmental agencies (Greenwald et al., 2006) or listed as a species of concern by private conservation organizations (Butcher, 2007). Regardless of the underlying rationale for their selection, priority wildlife species within managed forests differ among regions and forest types. Moreover, the priority species targeted for habitat improvement via prescribed management actions ultimately are determined by the landowner’s objectives.

Why favor these species over others? All forest management, including no active management (a.k.a., passive management or benign neglect), influences the abundance and composition of wildlife species. For many priority species, insufficient habitat, or an overabundance of habitat with unsuitable conditions, likely contributed to their designation as a priority species. Therefore, where modification of forest habitat will promote habitat conditions conducive to supporting sustainable populations of these species, it behooves managers to prescribe silvicultural actions that will result in increased habitat for these species.

3. Desired habitat conditions for priority species

Because priority species differ among regions and among forest types, quantification of specific habitat characteristics suitable for all forests and all species is not practical. Even so,

empirical evaluations of wildlife species responses to different silvicultural treatments provide insight into the general habitat conditions that benefit different species groups.

Amphibians – Within landscapes, amphibians benefit from protection of existing forest, avoiding further forest fragmentation, and disturbances that mimic natural processes (Kingsburg & Gibson, 2002). Within stands, seasonally available water, especially vernal pools, should be protected from disturbance (e.g., siltation) by retaining forest canopy ($\geq 75\%$) immediately surrounding pools and maintaining $>50\%$ canopy for up to 120 m from seasonal waters (Calhoun & deMaynadier, 2004). Amphibians tend to benefit from retention of sufficient canopy throughout the forest stand to provide a partially shaded forest floor. Their abundance also tends to be positively associated with a deep litter layer and abundant woody debris on the forest floor (Maguire et al., 2005). Within southern pine forests (e.g., longleaf pine [*Pinus palustris*]), thinning and burning are positively related to increased abundance of priority (open-pine adapted) amphibian species (Steen et al., 2010).

Reptiles – The effects of silvicultural treatments on reptiles have been mixed, often exhibiting no discernable effect. Despite inconclusive studies, development of heterogeneous canopies, with numerous gaps enabling light to reach the forest floor, enhances thermoregulation opportunities for reptiles. Herpetofauna (reptiles and amphibians) may exhibit different regional responses to silviculture. Similarly, species-specific response to silviculture may result in species responding positively, negatively, or not at all (Russell et al., 2004). Thus, more research is needed on the effects of thinning and retained structural elements on reptiles and amphibians (Jones et al., 2009; Verschuyt et al., 2011).

Birds: The intensity of harvest, otherwise stated as the magnitude of retention, influences species composition of bird communities. More canopy retention favors retention of mature-forest species, whereas more intense harvest tends to favor early-successional species and forest generalist species (Schieck et al., 2000). In northern hardwood forests, retention of $20 \text{ m}^2 \text{ ha}^{-1}$ basal area maintained habitat for mature-forest birds yet afforded habitat for birds using early-successional habitat (Homes and Pitt, 2007). Similarly, in South American hardwood forests, combined dispersed and aggregated retention permitted colonization by early successional birds yet retained species characteristic of old-growth forests within retained aggregates (Lencinas et al., 2009). In Tasmanian eucalypt forest, long-term, aggregated retention harvests better sustained mature forest bird communities, post-harvest, compared with clearcut or dispersed retention harvests (Lefort and Grove, 2009). In young, conifer-dominated stands, thinning promotes diversity of breeding birds – a variety of thinning intensities and patterns, ranging from no thinning to widely spaced residual trees, will maximize avian diversity at the landscape scale and provide structural diversity within and among stands (Hagar et al., 2004). In southern pine forests, bird abundance and richness were positively related to volume of coarse woody debris and density of snags (Jones et al., 2009). In Midwestern oak-hickory forests, thinning harvest and prescribed fire increased avian nest survival (Streby and Miles, 2010). However, responses to variable retention harvests are species specific and temporally dynamic: Response of some species is immediate but short-lived, whereas other species exhibit a deferred (maybe several years post-treatment) but long-lasting response (Hagar and Friesen, 2009; Twedt and Somershoe, 2009).

Bats – Many bat species avoid spatial clutter: Thus bats tend to be more abundant in forests with open structure (harvested or old-growth) compared to closed-structured mid-successional forest (Loeb and O'keefe, 2006; Patriquin and Barclay, 2003; Menzel et al., 2005).

Forest structure appears more important than forest composition for foraging bats (Loeb and O'Keefe, 2006). Increased species and structural diversity of woody species, including providing broadleaf trees and shrubs in corridors and surrounding water features, tend to enhance foraging opportunities for bats. Conversely, forest type (or tree species) may be an important determinant of roosting location or behavior (Barclay and Kurta, 2007; Perry et al., 2007). Specifically, roost trees of bats are generally tall with large diameters and are located in stands with open canopies and high snag densities (Kalcounis-Rüppell et al., 2005). Compared to trees used by foliage-roosting bats, cavity-roosting bats use trees within stands that have more open canopies and trees that are closer to water than are random trees (Kalcounis-Rüppell et al., 2005).

Deer – Within landscapes, decreased road density and lessening of associated traffic reduces the negative impacts of disturbance. Silviculture should promote structurally complex, uneven-aged stands, that include forest openings (1 – 20 ha), and provide structural heterogeneity within and among stands (Nelson et al., 2008). Resulting habitat should provide understory forage and cover while retaining species diversity of trees.

Bear – Habitat selection is largely influenced by food abundance (Costello and Sage, 1994). A diversity of tree and shrub species producing hard mast (acorns and nuts) as well as soft mast (berries and fruits) is beneficial. Thinning, group selection, or patch cuts should be used to promote regeneration and recruitment of mast producing species, as well to increase understory food production, escape cover, and bedding areas. Care should be exercised to retain some large trees within each stand and to protect potential den trees during harvest.

Small Mammals – Silviculture that retains legacy structures, variable-density thinning, and management for increased forest decadence combined to support 3 species of squirrel in Douglas-fir forests (Carey, 2000). In southwestern conifer forests, thinning and burning treatments had positive effects on most small mammals. Even so, effects of silviculture on small mammals are species-specific, such that a positive response in abundance of one species may be offset by negative response in abundance of another species.

Effective strategies to achieve habitat for different species across spatial and temporal scales should include developing: 1) measurable objectives, 2) integrated conservation goals and silvicultural prescriptions, 3) clear and practical guidelines, effective training, and communication programs, and 4) a monitoring and an adaptive management process to evaluate and improve results (Munks et al., 2009).

3.1 Desired landscape conditions

Landscape conditions deemed suitable for priority species are highly dependent upon which species have been designated as priority species within the landscape under consideration. Furthermore, even after priority species have been designated, determining the nature and extent of desired landscape conditions have relatively little empirical justification (Moilanen, 1999). Even so, various methods have been employed to evaluate the appropriate landscape area and context for priority species.

A relatively simple, 6-step approach was put forward to identify the appropriate landscapes for forest bird conservation in the Mississippi Alluvial Valley: 1) establish priority species, 2) establish habitat priorities, 3) identify habitat requirements for priority species, 4) determine

the extent and location of extant habitat, 5) set site-specific habitat goals, and 6) establish metapopulations goals (Mueller et al., 2000). A variation of this method was used to characterize priority of lands for forest restoration within the Mississippi Alluvial Valley and thereby delineate 107 discrete landscape units that were deemed capable of supporting viable populations (assumed to be ≥ 500 pairs) of priority silvicolous birds (Twedt et al., 2006; Fig. 4a).

Building upon this model, the Lower Mississippi Valley Joint Venture Forest Resource Conservation Working Group (Wilson et al., 2007) quantified desired forest conditions within these landscapes as:

1. $>70\%$ forest habitat, with large contiguous forest areas preferred;
2. 70-95% of forest area actively managed via silviculture intended to encourage desired stand conditions;
3. 35-50% of forest should meet desired forest conditions at any point in time, (Recognizing that habitat conditions are dynamic, silvicultural treatments should target conditions that yield extended temporal duration of desired stand conditions);
4. $\leq 10\%$ of forest should be in regeneration harvests (where $>80\%$ of overstory is removed) that are >3 ha in area;
5. $\sim 5\%$ of area should be in shrub-scrub (thamnic woody vegetation) habitat, but this may including early seral forest stages; and
6. 5-30% of forest area should be passively managed as wilderness, natural areas, or set-aside areas that are not subjected to silvicultural manipulation.

A further refinement of this method has been used within the West Gulf Coastal Plain and Ouachita Mountains Ecoregion to characterize landscape requirements of birds reliant on open-canopy, mature pine forests (Keister et al., 2011). The first steps were again to identify priority species and define suitable habitats for priority species. However, area requirements for each of these priority species were determined by assessing the minimum territory size or home range of breeding pairs. Connectivity among territories of appropriate size was assumed to be related to the likely dispersal distance of each priority species. The minimum viable population size was estimated for each species (Hanski et al., 1996). These species-specific characteristics were used to evaluate existing landscape suitability for priority species. The sum of the total area of all habitats that were capable of supporting a breeding pair (assuming silvicultural management would yield suitable stand conditions) and that were joined by virtue of being within dispersal distance of other suitable habitat was used to evaluate landscapes. Landscapes were deemed suitable when their total areas were sufficient to support a minimum viable population of the species: Non-suitable landscapes were characterized with regard to their suitability based on the proportion of a minimum viable population that could be supported (Fig. 4b-d). Desired landscape conditions for multiple species were then jointly considered (Fig. 4, right).

3.2 Desired site conditions

Desired site conditions are dependent upon which species have been designated as priority species. Even so, as identified above, there are commonalities among stand characteristics that have been found to benefit many species and species groups. Prescribed wildlife

forestry silvicultural practices can induce disturbance within forests and thereby stimulate development of desired structural conditions. Specific needs to be addressed when prescribing management actions include: 1) development and maintenance of structural (vertical and horizontal) heterogeneity, 2) achieving and retaining site appropriate tree species diversity, 3) maintaining an appropriate [sufficient yet acceptable] level of decadence, and 4) providing adequate reproduction to ensure sustainable habitat conditions and maintain future management options (Fig. 5).




Fig. 4. Landscapes identified for priority birds based on forest restoration (a) in the Mississippi Alluvial Valley (Twedt et al. 2006) and on proportion of minimum viable population supported for Bachman's sparrow (b), brown-headed nuthatch (c), and red-cockaded woodpecker (d) within the West Gulf Coastal Plain (Keister et al. 2011).

Maintaining or enhancing diversity of species – A diversity of flora, in the canopy, mid-story, and understory, buffers fluctuations (annual, seasonal, and temporal) in phenology, productivity, susceptibility, and merchantability of forests, although this is not universally applicable to canopy species (e.g., longleaf pine stand canopies may be predominately one species). Because diversity of fruiting species affords temporal and spatial availability of fruits, and because fruit use by wildlife in temperate forests is substantial, with fruits being consumed primarily during winter (McCarty et al., 2002), maintaining and enhancing diversity of fruit bearing species should be encouraged. Moreover, a variety of fruits provide alternative foods in the event of mast crop failure of other species. Similarly, different plants host different caterpillar (larval Lepidoptera) species, some of which have species-specific relationships. Because

butterfly and moth larvae are consumed by many wildlife species, especially birds, maintaining a diversity of trees and other flora within forest stands extends the temporal availability and increases the diversity of these insects (Hagar, 2004, Twedt and Best, 2004). In addition, forests with diverse species compositions are less vulnerable to catastrophic effects of disease and insects. Indeed, recent detrimental oak mortality in the Ozark Mountains Ecoregion has been attributed, in part, to the preponderance of oaks within these forests (Riggins et al. 2009). Similarly, species-specificity of many diseases and insects (e.g. chestnut blight [*Cryphonectria parasitica*] impact on American chestnut [*Castanea dentate*] or emerald ash borer [*Agrilus planipennis*] infestation of ash species [*Fraxinus* spp.]) provides argument for the maintenance of forest diversity to ameliorate the potentially catastrophic effects of regional outbreaks of disease or insect pests.



Fig. 5. Forest structure observed during summer (left) and winter (right) within bottomland hardwood forests circa 4 years after variable-retention, wildlife-forestry based silvicultural treatments (bottom) and on untreated control stands (top).

Finally, our naïve perception of the value of “undesirable species” is likely insufficient to comprehend the value of these species within the ecosystem. Therefore, preferential retention or diminution of native species may have unintended detrimental consequences (Lockhart 2004). For example, although the presence of piñon pine (*Pinus edulis*) is among the most important habitat features for many birds in piñon-juniper forests, the favored (86%) location of nests in juniper (*Juniperus osteosperma*) suggests that managers should avoid preferential thinning of juniper within these forests (Francis et al., 2011).

Increased structural heterogeneity (horizontal and vertical) – More structure generally provides a greater number of habitat niches for occupancy by different wildlife species, thereby increasing species richness, which may include priority species. Managers must, however,

be cautious of excessive application of silviculture intended to promote heterogeneity as these excesses may promote what could be viewed as 'homogeneity' of this habitat structure at a more dispersed scale. As different intensity of silvicultural treatment result in different stand complexity, silvicultural decisions made at each stage of stand development affect forest structure (Table 2).

Lower Basal Area - Thinning canopies has a positive or neutral effect on diversity and abundance of most taxa, albeit intensity and biophysical setting influence wildlife response (Verschuyl et al., 2011). In general, lower basal areas are associated with increased density of understory vegetation which adds to structural diversity, provides cover for escape, thermoregulation, and nesting or bedding. Moreover patches of dense understory enhance forage opportunities for species that browse grasses, herbs, or seedlings, species that glean insects from leaves, and frugivores.

Large, old, decadent trees - These features tend to be especially characteristic of old-growth-like forest conditions. As such, these characteristics are difficult to achieve given historical forest management where short rotation lengthen and harvest of economically mature trees limit availability of these features. Large cavity trees are used as den sites by bears, as well as other meso-mammalian (e.g., martens, fishers, raccoons, porcupines, weasels) and larger avian species (e.g., woodpeckers, owls, mergansers, and ducks). Similarly, dominant emergent trees provide perches and nesting platforms for large raptors (e.g., eagles, hawks, and kites). Although avian excavators (e.g., woodpeckers) are the primary cavity producers in North America, producing 77% of nesting cavities, this is not the case globally ($\leq 26\%$) where most cavity formation results from damage and decay (Cockle et al., 2011). Thus, cavity-using communities are highly dependent upon maintaining decadent trees within forest stands. As such, the presence of large, decadent trees may be essential for occupancy of stands by many of these large fauna.

Abundant standing snags and downed woody debris - Diversity and abundance of birds and biomass of invertebrates are positively associated with the volume of downed coarse woody debris and snags (Riffell et al., 2011). In unmanaged oak and oak-beech forests, 25% of dead wood was standing, whereas 75% was downed (Vanderkerkhove et al., 2009). Snags (and other decadent trees, such as those with heart rot), are characteristically exploited by primary-cavity excavators (e.g., woodpeckers) as well as those species that forage on saprophytic insects. Excavated cavities are subsequently used by myriad secondary-cavity using species. Lack of snags or other decadent trees may restrict this entire suite of cavity using species (Cockle et al. 2011).

Providing sufficient and suitable regeneration - As successional changes in forest species composition are often beyond human life-spans, the need for sufficient and suitable regeneration of canopy trees is often neglected. However, within the concept of wildlife forestry, regeneration of canopy species is ongoing (generally not a single event - as in even-aged forest management). Thus, ensuring sufficient regeneration of marketable trees, and thereby enhancing the long-term merchantability of forest stands, is imperative if desired habitat conditions for priority wildlife species are to be sustained through repeated, prescribed, silvicultural treatments. Notably, where maintaining shade-intolerant species within future generations is deemed desirable for anticipated management options, silviculturally induced gaps (patch cuts or group selections) should be of sufficient area to allow regeneration and development of these shade-intolerant species.

Treatment	Increasing forest diversity, structure, and complexity →		
Regeneration harvest	Clearcut	Legacy retention; dispersed or aggregated	Wildlife forestry; variable retention with canopy gaps
Restoration planting	Monoculture	Monoculture with supplemental natural reproduction	Mixture of species; hard and soft mast; shade tolerant and intolerant
Stand modification (TSI)	Herbicide removal of understory and midstory	Partial removal of competitors	Heterogeneous retention and removal of understory competitors
Thinning	Systematic single species retention	Multi-species retention	Heterogeneous retention and thinning; underrepresented species preferably retained

Table 2. Effects of wildlife forestry and other silviculture on stand structure (expanded from Carey, 2003b).

Although not specially referring to wildlife forestry, Carey (2006) succinctly encapsulated the holistic approach of wildlife forestry as - “*Management ... of forest development and landscape dynamics is more likely to be successful in maintaining ecosystem and landscape function than just providing select structural elements in stands and select structural stages in landscapes, as is often suggested for conservation.*”

3.3 Regional examples of desired stand conditions

Bottomland hardwood forests - A preponderance of closed canopy, second-growth forests with relatively homogeneous structure prompted recommendations for increased area of open canopied forests with greater structural diversity and more understory vegetation development (Wilson et al., 2007). Basal area should be reduced to 14-16 m² ha⁻¹ with ≥25% of the basal area approaching biological maturity (i.e., in older age classes). Forest canopy cover should be 60%-70% with at least 5 stems ha⁻¹ being canopy dominant trees - emergent canopies preferred. Understory vegetation should be robust yet patchily distributed with 25%-40% cover. Density of snags ≥25 cm diameter at breast height (dbh) should be >15 snags ha⁻¹ or have a basal area >0.9 m² ha⁻¹. Volume of coarse woody debris should exceed 14 m³ ha⁻¹ and advanced reproduction should be present on 30%-40% of the stand (Table 3).

Northwestern U.S. coniferous forests - Bird species richness and abundance are positively related to proportion of broadleaf forest, from 1% to 25% canopy cover, within coniferous dominated landscapes. Thus, it is import to maintain or restore broadleaf vegetation in these coniferous dominated stands (Betts et al., 2010). Desired stand conditions reflect a structure similar to that of old-growth forest conditions (Cissel et al., 2006). Multiple tree species, representing multiple age cohorts, should occupy stands and this mix should include at least 1 shade tolerant species. Large diameter (>125 cm dbh) conifers should be present at densities of 5-15 stems ha⁻¹. At least half of the 20-30 snags ha⁻¹ should be >60 cm dbh and a third of coarse woody debris should be of large (>60 cm) diameter (Table 3).

Southeastern U.S. pines - Historically open-canopied pine forests were found throughout the southeastern United States. These open forests were maintained by frequent fires. Prescribed burning, complimented with thinning of pine canopies and herbicide treatments

of encroaching hardwoods can restore these historic conditions. Open canopy conditions with an abundant and floristically diverse understory increase species richness of priority open pine associated amphibians (Guyer, and Bailey, 1993, Steen et al., 2010). Similarly, abundance of endangered gopher tortoises (*Gopherus polyphemus*) and red-cockaded woodpecker (*Picoides borealis*) respond positively to prescribed fire within open pine stands. Desired stand conditions include open (40-60%) canopies that are dominated by large (≥ 35 cm dbh) pines. These large pines should account for ≥ 4.5 m² ha⁻¹ of a total basal area of 11 – 16 m² ha⁻¹. Mid-story should be sparse (<15%) with very limited hardwoods (<5%). Ground cover and understory should be lush (>80% cover) and comprised predominately of grasses and forbs. At least 7 snags ha⁻¹ should be present. Relatively frequent fire (≤ 3 year interval) is likely required to maintain these desired conditions (Table 3).

4. Economic considerations

Wildlife forestry's primary intent is to provide suitable habitat conditions for priority wildlife. Even so, silvicultural practices that require long-term commitments or repeated expenditures may not be undertaken and thus, may not provide desired results. Indeed, silvicultural prescriptions intended to enhance wildlife habitat must be cognizant of their expense and their long-term effect on future merchantability of the stand. With that caveat, wildlife forestry silviculture may have both positive and negative economic effects.

A positive impact of most wildlife forestry practices is a reduction in wildfire hazards. Indeed, managing tree density and species composition with prophylactic silvicultural treatments that include thinning, surface fuel treatments, and prescribed fire reduce the risk of wildfire (Graham et al., 1999).

Negative economic factors include wounding of potentially merchantable trees that remain within stands after silvicultural treatments, as these wounds may result in future economic loss due to mortality or reduction in quality of timber (Hennon and DeMars, 1997). Thus, harvest prescriptions should account for this possibility and provide recommendations to reduce unintended damage to residual trees. On the other hand, from a wildlife use perspective, basal wounds to large retained trees have the potential to increase the number of decant large trees within a stand.

Another negative economic impact of wildlife silviculture may be successional changes in species composition. As noted for ensuring adequate tree regeneration within stands, managers should be aware of unintended shifts in species composition as a consequence of prescribed wildlife silviculture, usually due to an increased presence of shade-tolerant species. Therefore, if shade-intolerant species are intended to be a part of future forest canopies, silvicultural treatments must ensure their regeneration and development (Twedt and Somershoe, in press).

All forests are subject to windthrow (toppling of live standing trees due to wind), although its severity is influenced by wind speed, climate, topography, hydrology, and soils (Ruel, 1995): Toppled trees are present in mature unharvested forests as well as along the edges of clearcuts. Even so, windthrow may increase after partial harvest silvicultural treatments. More retention within a stand, however, tends to lessen the severity of windthrow (Franklin et al., 1997). Windthrow can also be mitigated through judicious silviculture, as some species are more susceptible to windthrow, possibly due to differences in form. Trees with

more extensive crown structure or shallow, poorly developed roots are more likely to be toppled by wind (Beese, 2001). Retained, sound, dominant trees are less subject to windthrow than decadent or subdominant trees. The dispersion, orientation, and shape of retention also influences susceptibility to windthrow, with tear-drop shaped, aggregated clumps, oriented with prevailing winds better able to withstand wind damage (Franklin et al., 1997). Even so, small (1.5 ha) patch cuts may be least vulnerable to wind (Beese, 2001).

Stand feature	Bottomland hardwood (Wilson et al., 2007)	Longleaf pine (Bragg, 2004; East Gulf Coastal Plain Joint Venture, unpublished)	Pacific northwest conifers (Cissel et al. 2006)	Young conifers in northern pacific rainforest Altman and Hagar, 2007)
Basal area	14 – 16 m ² ha ⁻¹	11 – 16 m ² ha ⁻¹	9 – 27 m ² ha ⁻¹	
Size class	≥25% of BA approaching biological maturity	>4 m ² ha ⁻¹ , ≥35 cm dbh	several tree species of varying size and age; large emergent trees 5 – 15 ha ⁻¹ , >125 cm dbh conifers;	large conifer trees
Density	>5 dominant ha ⁻¹	12 – 30 pines ha ⁻¹ , >76 cm dbh	60 – 85 ha ⁻¹ , 38-76 cm dbh; 250 – 500 ha ⁻¹ <38 cm dbh	<500 ha ⁻¹ ; thin to promote growth
Canopy cover	60 – 70%	40 – 60% (≤5% hardwood) Patchy	multi-layered canopies	variation in overstory achieved via silviculture, low % canopy cover to encourage understory
Midstory	25 – 40%	≤15% (≤5% hardwood) Open	open space among the lower branches of canopy	retain and protect old shrubs and shrub patches
Ground cover	25 – 40%	>80%		variation in understory associated with variable canopy
Snags & stressed trees	>15 ha ⁻¹ , ≥25 cm dbh; or ≥5 ha ⁻¹ , ≥ 51 cm dbh; or >0.9 m ² ha ⁻¹ , >25 cm dbh	≥7 (12-25) ha ⁻¹ , red heart; 10 – 50% cull in retained trees	20 – 30 ha ⁻¹ >25 cm dbh; 50% > 60 cm dbh; cavities in standing trees	large
Coarse woody debris	≥14 m ³ ha ⁻¹	21 – 54 m ³ ha ⁻¹ (includes snags)	275 linear m >25 cm diameter; 33% > 60 cm dbh	retain, protect, and recruit large diameter CWD
Reproduction	30 – 40% of area with advanced reproduction	10% of area with advanced reproduction		encourage regeneration and retention of a diversity of species
Other	>10 visible cavities ha ⁻¹ ; 1 den tree (4 ha) ⁻¹	fire return ≤3 year	≥2 species (≥1 shade-tolerant); ≥2 age cohorts; >150 years old	retain and promote deciduous trees and shrubs

Table 3. Described and quantified desired site (stand) conditions that are deemed to provide enhanced habitat for priority wildlife species in different forest types and seral stages.

There is little basis for economic comparisons of wildlife forestry and other, traditional, silvicultural systems. Wildlife forestry may provide sustained revenues from exploitive use of wildlife, as well as from periodic harvest of forest products. But even without income from wildlife, little economic loss was reported from variable retention silviculture compared to even-aged, shelterwood cuts in *Nothofagus pumilio* forests (Pastur et al., 2009). Similarly, simulation models spanning 300 years that compared economic returns and ecological benefits within Pacific northwest forests being managed for biodiversity with forests being managed for timber production, and incorporating narrow riparian reserves (i.e., set asides), found significantly greater ecological benefits associated with biodiversity management (Carey et al., 1999). Concurrently, the economic cost of achieving those benefits was relatively modest, with 18% loss in net present value (NPV), only 4% loss if riparian protection was similar to that afforded within the biodiversity management model, and fully a 13% gain in NPV when newly enacted state laws for riparian protection were enforced (Carey, 2003a). More notably, under management for biodiversity, tree quality improved, decadal revenues increased by 150%, forest-based employment quadrupled, and manufacturing diversified and became more reliant on high quality products and value added manufacturing (Carey, 2003a, Carey et al., 1999, Lippke et al., 1996).

5. Restoration

If successfully implemented, wildlife forestry is self-sustaining, such that reforestation (e.g., artificial regeneration or stand regeneration) is rare. Even so, appropriate practices are increasingly being sought in conjunction with forest restoration that better enable attainment of desired forest conditions. For example, decades of bottomland afforestation, wherein agricultural lands are being restored to forested wetlands, has provided valuable insight regarding appropriate species and planting densities (Wilson et al., 2007). Management practices similar to those employed for wildlife forestry have also been used to restore longleaf pine forests on sites converted to forests dominated by other species. Similarly, within production-oriented plantation forests management alternatives are sought that will enhance biodiversity and improve wildlife habitat, yet minimally compromise product output.

Recommendations for restoration and mid-rotation management have included planting or maintaining species mixtures to ensure stand diversity and improve forest product quality (Twedt and Best, 2004, Lockhart et al. 2008). Mixed species planting of native species are favored over monocultures of exotic species. Some legacy trees (live or snags) and understory vegetation should be retained at harvest through a second rotation. Site-preparation should reflect natural disturbances and conserve coarse woody debris. Thinning earlier in the rotation and increasing rotation length also increase structural diversity (Hartley, 2002; Kerr, 1999). Finally, for some forest types, maintaining herbaceous or early succession habitat for a longer period provides benefits to priority bird species (Altman and Hagar, 2007).

6. Evaluating results of management actions

Quantitatively defined landscape conditions and desired stand conditions have been identified for only a few ecoregional landscapes and only a few forest types within these landscapes. Wildlife biologists, foresters, and conservation planners must develop a shared

vision regarding what each perceives as the desired habitat condition within forests. In the landscape context, these conditions need not be the same for all forests, nor will they be identical for all priority species. Research and monitoring will be required to evaluate species responses to different ranges of habitat conditions perceived as desirable. Judgments may be needed when priority species exhibit conflicting responses to what are perceived as desired conditions.

At present we have little empirical data regarding the range of economic returns from wildlife silviculture relative to traditional production-oriented silviculture. Appropriate measures of economic returns, including an evaluation of the long-term sustainability of different management options, are needed. In addition, we have assessed the faunal responses of relatively few species to wildlife forestry silviculture. Future efforts must assess comparable responses of less-studied priority species and species groups as well as evaluate appropriate demographic responses of all species.

To determine if prescribed management actions elicit intended responses from priority species, monitoring is required. Monitoring should target specific management actions and evaluate spatial and temporal responses. Silvicultural prescriptions should consider incorporation of the principles of adaptive management during their development, implementation, and evaluation (Fig. 1). Feedback from such efforts can provide an assessment of the adequacy and sustainability of wildlife forestry silviculture. Science based knowledge of the results of these management practices is crucial to their long-term success, as *“Conservation is paved with good intentions which prove to be futile, or even dangerous, because they are devoid of critical understanding either of the land, or of economic land use”* (Leopold, 1953).

7. Acknowledgement

I am grateful to the members of the Lower Mississippi Valley Joint Venture Forest Resource Conservation Working Group for thoughtful discussions and arguments regarding the merits of wildlife forestry. Similarly, members of the West Gulf Coastal Plain and Ouachita Mountains Landbird working group provided insight and understanding regarding application of wildlife forestry principles within southern pine systems. Finally, I am grateful to the many pioneers and advocates of alternative silviculture from which most of our understanding of wildlife forestry was derived.

8. References

- Altman, B. & Hagar, J. (2007). *Rainforest birds- A land manager's guide to breeding bird habitat in young conifer forests in the Pacific Northwest*, U.S. Geological Survey Scientific Investigations Report 2006-5304, Reston, Virginia
- Askins, R. (2001). Sustaining biological diversity in early succession communities: the challenge of managing unpopular habitats. *Wildlife Society Bulletin*, Vol.29, No.2, pp. 407-412
- Aubry, K.; Halpern, C. & Maguire, D. (2004). Ecological effects of variable-retention harvests in the northwestern United States: the DEMO study. *Forest Science and Landscape Research*, Vol.78, No.1-2, pp. 119-134

- Aukema, J. & Carey, A. (2008). *Effects of variable-density thinning on understory diversity and heterogeneity in young Douglas-fir forests*. Research Paper PNW-RP-575. U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station, Portland, Oregon
- Bailey, M.; Holmes, J.; Buhlmann, K. & Mitchell, J. (2006). *Habitat Management Guidelines for Amphibians and Reptiles of the Southeastern United States*. Partners in Amphibian and Reptile Conservation Technical publication HMG-2, Montgomery, Alabama
- Barclay, R. & Kurta, A. (2007). Ecology and behavior of bats roosting in tree cavities and under bark, In: *Bats in forests: conservation and management*, M.J. Lacki; J.P. Hayes & A. Kurta, (Eds.), pp. 17-59, Johns Hopkins University Press, Baltimore, Maryland
- Barneycastle, C. (2001). The Sustainable Forestry Initiative of the American Forest & Paper Association, In: *Proceedings of the symposium on Arkansas forests: a conference on the results of the recent forest survey of Arkansas*, J.M. Guldin, (Ed.), 110-112, U.S. Department of Agriculture, Forest Service, Southern Research Station, General Technical Report SRS 41, Asheville, North Carolina
- Bauhus, J.; Puettmann, K. & Messier, C. (2009). Silviculture for old-growth attributes. *Forest Ecology and Management* Vol.258, No.4, pp. 525-537
- Beese, W. (2001). Windthrow monitoring of alternative silvicultural systems in montane coastal forests, In: *Proceedings of the Windthrow Researchers Workshop: Windthrow Assessment and Management in British Columbia*, S.J. Mitchell & J. Rodney, (Compilers), pp. 2-11, Richmond, British Columbia
- Beggs, L. (2004). *Vegetation response following thinning in young Douglas-fir forests of western Oregon: Can thinning accelerate development of late-successional structure and composition?* M.S. thesis, Oregon State University, Corvallis, Oregon
- Betts, M.; Hagar, J.; Rivers, J.; Alexander, J.; McGarigal, K. & McComb, B. (2010). Thresholds in forest bird occurrence as a function of the amount of early-seral broadleaf forest at landscape scales. *Ecological Applications* Vol.20, No.8, pp. 2116-2130
- Bonnicksen, T.M. (2000). *America's ancient forests: from the ice age to the age of discovery*. John Wiley, New York.
- Bragg, D. (2004). A prescription for old-growth-like characteristics in southern pines, In: *Silviculture in special places: Proceedings of the National Silviculture Workshop*, W.D. Sheppard & L.G. Eskew, (Compilers), pp. 80-92, U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Proceedings RMRS-P-34, Fort Collins, Colorado
- Brown, S.; Hickey, C.; Harrington, B. & Gill, R., (Eds.). (2001). *United States Shorebird Conservation Plan*. Manomet Center for Conservation Sciences, Manomet, Massachusetts
- Butcher, G.S. (2007). Common birds in decline, a state of the birds report. *Audubon*, Vol.109, No.4, pp. 58-62
- Calhoun, A. & deMaynadier, P. (2004). *Forestry habitat management guidelines for vernal pool wildlife*. MCA Technical Paper No. 6, Metropolitan Conservation Alliance, Wildlife Conservation Society, Bronx, New York
- Carey, A. (2000). Effects of new forest management strategies on squirrel populations. *Ecological Applications*, Vol.10, No.1, pp. 248-257

- Carey, A. (2003a). Restoration of landscape function: reserve or active management? *Forestry*, Vol.76, No.2, pp. 221-230.
- Carey, A. (2003b). Managing for wildlife: A key component for social acceptance of compatible forest management, In: *Compatible Forest Management*, A. Monserud, R.W. Haynes, & A. C. Johnson, (Eds.), Kluwer Academic Publishers, Dordrecht, The Netherlands
- Carey, A. (2006). Active and passive forest management for multiple values. *Northwestern Naturalist*, Vol.87, No.1, pp. 18-30
- Carey, A., Lippke, B. & Sessions, J. (1999). Intentional systems management: managing forests for biodiversity. *Journal of Sustainable Forestry*, Vol.9, No.3-4, pp. 83-125
- Cissel, J.; Anderson, P.; Olson, D.; Puettmann, K.; Berryman, S.; Chan, S. & Thompson, C. (2006), *BLM Density Management and Riparian Buffer Study: Establishment Report and Study Plan*, U.S. Geological Survey, Scientific Investigations Report 2006-5087, Reston, Virginia, USA
- Cissel, J.; Swanson, F. & Weisberg, P. 1999. Landscape management using historical fire regimes: Blue River, Oregon. *Ecological Applications*, Vol.9, No.4, pp. 1217-1231
- Cockle, K.; Martin, K. & Wesołowski, T. (2011). Woodpeckers, decay, and the future of cavity-nesting vertebrate communities worldwide. *Frontiers in Ecology and the Environment*, Vol.9, No.7, pp. 377-382
- Costello, C. & Sage Jr., R. (1994). Predicting black bear habitat selection from food abundance under 3 forest management systems, *Bears: Their Biology and Management: A Selection of Papers from the Ninth International Conference on Bear Research and Management*, pp. 375-387, , Missoula, Montana, USA, February 23-28, 1992
- Egan, A. & Luloff, A. (2000). The exurbanization of America's forests: Research in rural social science. *Journal of Forestry*, Vol.98, No.3, pp. 26-30
- Ford, W.; Russell, K.; Moorman, C. (Eds.). (2002). *Proceedings: the role of fire for nongame wildlife management and community restoration: traditional uses and new directions*, Nashville, Tennessee, September 15, 2000. U.S. Department of Agriculture, Forest Service, Northeastern Research Station, General Technical Report NE-288, Newtown Square, Pennsylvania, USA
- Francis, C.; Ortega, C. & Hansen, J. (2011). Importance of juniper to birds nesting in piñon-juniper woodlands in northwest New Mexico. *Journal of Wildlife Management*, Vol.75, No.7, pp. 1574-1580
- Franklin, J. (1989). Toward a new forestry. *American Forests*, (November/December 1989), pp. 37-44
- Franklin, J.; Berg, D.; Thornburgh, D. & Tappeiner, J. (1997). Alternative silvicultural approaches to timber harvesting: Variable retention harvest systems, In: *Creating a Forestry for the 21st Century: The Science of Ecosystem Management*, K.A. Kohn & J.F. Franklin (Eds.), 111-139, Island Press, Washington, DC, USA
- Franklin, J.; Forman, R. (1987). Creating landscape patterns by forest cutting: Ecological consequences and principles. *Landscape Ecology*, Vol.1, No.1, pp. 5-18
- Franklin, J.; Mitchell, R. & Palik, B. (2007). *Natural disturbance and stand development principles for ecological forestry*, General Technical Report NRS-19, U.S. Department of

- Agriculture, Forest Service, Northern Research Station, Newtown Square, Pennsylvania, USA
- Garman, S.; Cissel, J. & Mayo, J. (2003). *Accelerating development of late-successional conditions in young managed Douglas-fir stands: a simulation study*. General Technical Report PNW-GTR-557, U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. Portland, Oregon, USA
- Graham, E. (1947). *The land and wildlife*, Oxford Press, New York, New York, USA
- Graham, R.; Harvey, A.; Jain, T. & Tonn, J. (1999). *The effects of thinning and similar stand treatments on fire behavior in Western forests*, General Technical Report PNW-GTR-463, U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station, Portland, Oregon, USA
- Greenwald, N.; Suckling, K. & Taylor, M. (2006). Factors affecting the rate and taxonomy of species listings under the U.S. Endangered Species Act, In: *The Endangered Species Act at 30: Renewing the Conservation Promise*, D.D. Goble, J.M. Scott & F.W. Davis (Eds.), 50-67, Island Press, Washington, DC, USA
- Guyer, C. & Bailey, M. (1993). Amphibians and reptiles of longleaf pine communities, In: *Proceedings of the Tall Timbers Fire Ecology Conference Number 18, The Longleaf Pine Ecosystem; Ecology, Restoration and Management*, 139-158, Tallahassee, Florida, USA
- Hagar, J.; Howlin, S. & Ganio, L. (2004). Short-term response of songbirds to experimental thinning of young Douglas-fir forests in the Oregon Cascades. *Forest Ecology and Management*, Vol.199, No.2-3, pp. 333-347
- Hagar, J. (2004). *Functional relationships among songbirds, arthropods, and understory vegetation in Douglas-fir forests, western Oregon*, Ph.D. dissertation, Oregon State University, Corvallis, Oregon, USA
- Hagar, J. & Friesen, C. (2009). *Young stand thinning and diversity study- response of songbird community one decade post-treatment*, U.S. Geological Survey Open-File Report 2009-1253, Reston, Virginia, USA
- Hanski, I. (1998). Metapopulation dynamics. *Nature* Vol.396, pp. 41-49
- Hanski, I.; Moilanen, A. & Gyllenberg, M. (1996). Minimum viable metapopulation size. *American Naturalist* Vol.147, No.4, pp. 527-541
- Hartley, M. (2002). Rationale and methods for conserving biodiversity in plantation forests. *Forest Ecology and Management*, Vol.155, No.1-3, pp. 81-95
- Hayward, G. (1991). Using population biology to define old-growth forests. *Wildlife Society Bulletin*, Vol.19, No.1, pp. 111-116.
- Heithecker, T. & Halpern, C. (2006). Variation in microclimate associated with dispersed-retention harvests in coniferous forests of western Washington. *Forest Ecology and Management*, Vol.226, No.1-3, pp. 60-71
- Hennon, P. & Demars, D. 1997. Development of wood decay in wounded western hemlock and Sitka spruce in southeast Alaska. *Canadian Journal of Forest Research*, Vol.27, pp. 1971-1978
- Holmes, S. & Pitt, D. (2007). Response of bird communities to selection harvesting in a northern tolerant hardwood forest. *Forest Ecology and Management*, Vol.238, No.1-3, pp. 280-292

- Hunter, W.; Buehler, D.; Canterbury, R.; Confer, J. & Hamel, P. (2001). Conservation of disturbance-dependent birds in eastern North America. *Wildlife Society Bulletin*, Vol.29, No.2, pp. 440-455
- Jones, P.; Hanberry, B. & Demarais, S. (2009). Stand-level wildlife habitat features and biodiversity in southern pine forests: A review. *Journal of Forestry*, Vol.107, No.8, pp. 398-404
- Kaetzel, B.; Hodges, D. & Fly, J. (2011). Landowner motivations for owning woodland on the Tennessee Northern Cumberland Plateau. *Southern Journal of Applied Forestry*, Vol.35, No.1, pp. 39-43
- Kalcounis-Rüppell, M.; Psyllakis, J. & Brigham, R. (2005). Tree roost selection by bats: An empirical synthesis using meta-analysis. *Wildlife Society Bulletin*, Vol.33, No.3, pp.1123-1132
- Karau, E. & Keane, R. (2007). Determining landscape extent for succession and disturbance simulation modeling. *Landscape Ecology* Vol.22, No.7, pp. 993-1006
- Kaufmann, M.; Huckaby, L.; Fornwalt, P.; Stoker, J. & Romme, W. (2003). Using tree recruitment patterns and fire history to guide restoration of an unlogged ponderosa pine/Douglas-fir landscape in the southern Rocky Mountains after a century of fire suppression. *Forestry*, Vol.76, No.2, pp. 231-241
- Keeton, W. (2006). Managing for late-successional/old-growth characteristics in northern hardwood-conifer forests. *Forest Ecology and Management* Vol.235, No.1-3, pp. 129-142
- Keister, A.; Twedt, D.; McKnight, S. & Tirpak, J. (2011). Prioritization of open pine management for landbirds of the West Gulf Coastal Plain and Ouachita Mountains, In: *Big Thicket Science Conference*, Stephen F. Austin State University, Nacogdoches, Texas, USA, April 8-10, 2011, Available from [http://atcofa.sfasu.edu/images/stories/pdf/\(2011\)_WCGPBTSC_Program_Book_3_30_2011_Final.pdf](http://atcofa.sfasu.edu/images/stories/pdf/(2011)_WCGPBTSC_Program_Book_3_30_2011_Final.pdf)
- Kerr, G. (1999). The use of silvicultural systems to enhance the biological diversity of plantation forests in Britain. *Forestry*, Vol.72, No.3, pp. 191-205
- Kingsbury, B. & Gibson, J. (2002). *Habitat management guidelines for amphibians and reptiles of the Midwest*. Partners in Amphibian and Reptile Conservation Technical Publication HMG-1, Fort Wayne, Indiana, USA, Available from <http://herpcenter.ipfw.edu/downloadables/MWHabitatGuide/Index.htm>.
- Kneeshaw, D. & Gauthier, S. (2003). Old growth in the boreal forest: A dynamic perspective at the stand and landscape level. *Environmental Reviews*, Vol.11, No.1 (supplement), pp. 99-114
- Knight, R. 1999. Private lands: The neglected geography. *Conservation Biology*, Vol.13, No.2, pp. 223-224
- Lefort, P. & Grove, S. (2009). Early responses of birds to clearfelling and its alternatives in lowland wet eucalypt forest in Tasmania, Australia. *Forest Ecology and Management*, Vol.258, No.4, pp. 460-471
- Lencinas, M.; Pastur, G.; Gallo, E. & Cellini, J. (2009). Alternative silvicultural practices with variable retention improve bird conservation in managed South Patagonian forests. *Forest Ecology and Management*, Vol.258, No.4, pp. 472-480
- Leopold, A. (1949). *A Sand County Almanac and Sketches Here and There*, Oxford University Press, New York, New York, USA.

- Leopold, A. (1953). *The Land Ethic Round River*, Oxford University Press, New York, New York, USA.
- Lindenmayer, D. & Franklin, J. (2002). *Conserving Forest Biodiversity: A Comprehensive Multiscaled Approach*, Island Press, Washington, DC, USA
- Lindenmayer, D. & Hobbs, R. (2007). *Managing and designing landscapes for conservation: Moving from perspectives to principles*, Blackwell Publishing, Malden, Massachusetts, USA
- Lindenmayer, D.; MacCarthy, M. & Donnelly, G. (1999). Factors affecting stand structure in forest - are there climatic and topographic determinants. *Forest Ecology and Management* Vol.123, No.1, pp. 55-63
- Lindenmayer, D.; Manning, A.; Smith, P.; Possingham, H.; Fischer, J.; Oliver, I. & McCarthy, M. (2002). The focal-species approach and landscape restoration: a critique. *Conservation Biology*, Vol.16, No.2, pp. 338-345
- Lippke, B.; Sessions, J. & Carey, A. (1996). *Economic analysis of forest landscape management alternatives*. CINTRAFOR Special Paper 21, College of Forest Resources, University of Washington, Seattle, Washington, USA
- Lockhart, B. (2004). All species have value. *Journal of Forestry*, Vol.102, No.1, (January/February 2004), pp. 60
- Lockhart, B.; Gardiner, E.; Leininger, T. & Stanturf, J. (2008). A stand-development approach to oak afforestation in the lower Mississippi Alluvial Valley. *Southern Journal of Applied Forestry*, Vol.32, No.3, pp. 120-129
- Loeb, S. & O'Keefe, J. (2006). Habitat use by forest bats in South Carolina in relation to local, stand, and landscape characteristics. *Journal of Wildlife Management*, Vol.70, No.5, pp. 1210-1218
- MacArthur, R. & Wilson, E. (1967). *The Theory of Island Biogeography*, Princeton University Press, Princeton, New Jersey, USA
- Maguire, D.; Halpern, C. & Phillips, D. (2007). Changes in forest structure following variable-retention harvests in Douglas-fir dominated forests. *Forest Ecology and Management*, Vol.242, No.2-3, pp. 708-726
- Maguire, C.; Manning, T; West, S & Gitzen, R. (2005). Green-tree retention in managed forests: post-harvest responses of salamanders, In: *Balancing ecosystem values: innovative experiments for sustainable forestry – proceedings of a conference*, C.E. Peterson & D.A. Maguire (Eds.), 265-270, General Technical Report PNW-GTR-635. U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station, Portland, Oregon, USA
- Mayer, A. & Cameron, G. (2003). Consideration of grain and extent in landscape studies of terrestrial vertebrate ecology. *Landscape and Urban Planning*, Vol.65, No.4, pp. 201-217
- McCarty, J.; Levey, D.; Greenberg, C. & Sargent, S. (2002). Spatial and temporal variation in fruit use by wildlife in a forested landscape. *Forest Ecology and Management*, Vol.164, No.1-3, pp. 277-291
- McClellan, M. (2004). Development of silvicultural systems for maintaining old-growth conditions in the temperate rainforest of southeast Alaska. *Forest Snow and Landscape Research*, Vol.78, No.1-2, pp. 173-190

- Mehlman, D.; Rosenberg, K; Wells, J. & Robertson, B. (2004). A comparison of North American avian conservation priority ranking systems. *Biological Conservation*, Vol.120, No.3, pp. 383-390
- Menzel, J.; Menzel Jr, M.; Kilgo, J.; Ford, W.; Edwards, J. & McCracken, G. (2005). Effect of habitat and foraging height on bat activity in the coastal plain of South Carolina. *Journal of Wildlife Management*, Vol.69, No.1, pp. 235-245
- Mitchell, R.; Hiers, J.; O'Brien, J.; Jack, S. & Engstrom, R. (2006). Silviculture that sustains: the nexus between silviculture, frequent prescribed fire, and conservation of biodiversity in longleaf pine forests of the southeastern United States. *Canadian Journal of Forest Research*, Vol.36, No.11, pp. 2724-2736
- Mitchell, R.; Palik, B. & Hunter, M. (2002). Natural disturbance as a guide to silviculture. *Forest Ecology and Management*, Vol.155, No.1-3, pp. 315-317
- Mladenoff, D.; White, M; Pastor, J. & Crow, T. (1993). Comparing spatial pattern in unaltered old-growth and disturbed forest landscapes. *Ecological Applications*, Vol.3, No.2, pp. 294-306
- Moilanen, A. (1999). Patch occupancy models of metapopulation dynamics: efficient parameter estimation using implicit statistical inference. *Ecology*, Vol.80, No.3, pp. 1031-1043
- Mueller, A.; Twedt, D. & Loesch, C. (2000). Development of management objectives for breeding birds in the Mississippi Alluvial Valley, In: *Strategies for bird conservation: the partners in flight planning process. Proceedings of the third Partners in Flight workshop*, 1-5 October 1995, Cape May, New Jersey, R. Bonney, D.N. Pashley, R. Cooper, & L. Niles, (Eds.), 12-17, Proceedings RMRS-P-16, U. S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Ogden, Utah, USA
- Munks, S.; Koch, A. & Wapstra, M. (2009). From guiding principles for the conservation of forest biodiversity to on-ground practice: Lessons from tree hollow management in Tasmania. *Forest Ecology and Management*, Vol.258, No.4, pp. 516-524
- Nelson, J.; Cottam, D.; Holman, E.; Lancaster, D.; McCorquodale, S. & Person, D. (2008). *Habitat guidelines for black-tailed deer: Coastal Rainforest Ecoregion*. Mule Deer Working Group, Western Association of Fish and Wildlife Agencies. Available from http://www.muledeerworkinggroup.com/Docs/Coastal_Rainforest_BT_D_Habitat_Guidelines.pdf
- North American Wildlife Management Plan. (1986). *North American Wildlife Management Plan: A Strategy for Cooperation*. Available from <http://www.fws.gov/birdhabitat/NAWMP/files/NAWMP.pdf>
- North, M. & Keeton, W. (2008). Emulating natural disturbance regimes: an emerging approach for sustainable forest management, In: *Patterns and processes in forest landscapes – multiple use and sustainable management*, R. Laforzezza, J. Chen, G. Sanesi, & T.R. Crow, (Eds.), 341-372, Springer, The Netherlands
- Oliver, C. & Larson, B. (1996). *Forest Stand Dynamics*. John Wiley and Sons, New York, New York, USA
- Palik, B.; Mitchell, R. & Hiers, J. (2002). Modeling silviculture after natural disturbance to sustain biodiversity in the longleaf pine (*Pinus palustris*) ecosystem: balancing

- complexity and implementation. *Forest Ecology and Management*, Vol.155, No.1-3, pp. 347-356
- Panjabi, A.; Dunn, E.; Blancher, P.; Hunter, W.; Altman, B.; Bart, J.; Beardmore, C.; Berlanga, H.; Butcher, G.; Davis, S.; Demarest, D.; Dettmers, R.; Easton, W.; de Silva Garza, H.; Iñigo-Elias, E.; Pashley, D.; Ralph, C.; Rich, T.; Rosenberg, K.; Rustay, C.; Ruth, J.; Wendt, J. & Will, T. (2005). *The Partners in Flight handbook on species assessment. Version 2005. Partners in Flight Technical Series Number 3*. Available from <http://www.rmbo.org/pubs/downloads/Handbook2005.pdf>
- Pastur, G.; Lencinasa, M.; Cellini, J.; Peric, P. & Estebana, R. (2009). Timber management with variable retention in *Nothofagus pumilio* forests of Southern Patagonia. *Forest Ecology and Management*, Vol.258, No.4, pp. 436-443
- Patriquin, K. & Barclay, R. (2003). Foraging by bats in cleared, thinned and unharvested boreal forest. *Journal of Applied Ecology*, Vol.40, No.4, pp. 646-657
- Perry, R.; Thill, R. & Leslie, D. (2007). Selection of roosting habitat by forest bats in a diverse forested landscape. *Forest Ecology and Management*, Vol.238, No.1-3, pp. 156-166
- Poudyal, N.; Hodges, D.; Fenderson, J. & Tarkington, W. (2010). Realizing the economic value of a forested landscape in a viewshed: an assessment of the Scottsboro-Bells Bend forest area near Nashville, Tennessee. *Southern Journal of Applied Forestry*, Vol.34, No.2, pp. 72-28
- Rich, T.; Beardmore, C.; Berlanga, H.; Blancher, P.; Bradstreet, M.; Butcher, G.; Demarest, D.; Dunn, E.; Hunter, W.; Iñigo-Elias, E.; Kennedy, J.; Martell, A.; Panjabi, A.; Pashley, D.; Rosenberg, K.; Rustay, C.; Wendt, J. & Will, T. (2004). *Partners in Flight North American Landbird Conservation Plan*, Cornell Lab of Ornithology, Ithaca, New York, USA, Available from http://www.partnersinflight.org/cont_plan/
- Riffell, S.; Verschuyt, J.; Miller, D. & Wigley, T. (2011). Biofuel harvests, coarse woody debris, and biodiversity - A meta-analysis. *Forest Ecology and Management*, Vol.261, No.4, pp. 878-887
- Riggins, J.; Galligan, L. & Stephen, F. (2009). Rise and fall of red oak borer in the Ozark Mountains of Arkansas, USA. *The Florida Entomologist*, Vol.92, No.3 (September 2009), pp. 426-433
- Ruel, J. (1995). Understanding windthrow: Silvicultural implications. *Forestry Chronicle*, Vol.71, No.4, pp. 434-444
- Russell, K.; Wigley, T.; Baughman, W.; Hanlin, H. & Ford, W. (2004). Responses of Southeastern amphibians and reptiles to forest management: A review, In: *Southern forest science: past, present, and future*, H.M. Rauscher & K. Johnsen, (Eds.), 319-334, General Technical Report SRS-75, U.S. Department of Agriculture, Forest Service, Southern Research Station, Asheville, North Carolina, USA
- Schieck, J.; Stuart-Smith, K. & Norton, M. (2000). Bird communities are affected by amount and dispersion of vegetation retained in mixedwood boreal forest harvest areas. *Forest Ecology and Management*, Vol.126, No.2, pp. 239-254
- Seymour, R. & Hunter, M. (1999). Principles of ecological forestry, In: *Maintaining Biodiversity in Forest Ecosystems*, M.L. Hunter, (Ed.), 22-62, Cambridge University Press, Cambridge, Massachusetts, USA

- Silva Ecosystem Consultants. (1992). *Landscape ecology literature review*. Available from <http://www.silvafor.org/assets/silva/PDF/Literature/LandscapeEcologyOverview.pdf>
- Steen, D.; Rall McGee, A; Hermann, S.; Stiles, J.; Stiles, S & Guyer, C. (2010). Effects of forest management on amphibians and reptiles: generalist species obscure trends among native forest associates. *Open Environmental Sciences*, Vol.4, pp. 24-30
- Streby, H. & Miles, D. (2010). Assessing ecosystem restoration alternatives in eastern deciduous hardwood forests using avian nest survival. *Open Environmental Sciences*, Vol.4, pp. 39-48
- St Laurent, M.; Ferron, J.; Haché, S. & Gagnon, R. (2008). Planning timber harvest of residual forest stands without compromising bird and small mammal communities in boreal landscapes. *Forest Ecology and Management*, Vol.254, No.2, pp. 261-275
- Swanson, M.; Franklin, J.; Beschta, R.; Crisafulli, C.; DellaSala, D.; Hutto, R.; Lindenmayer, D. & Swanson, F. (2011). The forgotten stage of forest succession: early-successional ecosystems on forest sites. *Frontiers in Ecology and the Environment*, Vol.9, No.2, pp. 117-125
- Tubby, I. & Armstrong, A. (2002). *The establishment and management of short rotation coppice – a practitioner's guide*. Forestry Commission Practice Note. Forestry Commission, Edinburgh. Available from [http://www.forestry.gov.uk/pdf/fcpn7.pdf/\\$FILE/fcpn7.pdf](http://www.forestry.gov.uk/pdf/fcpn7.pdf/$FILE/fcpn7.pdf)
- Twedt, D. & Best, C. (2004). Restoration of floodplain forests for the conservation of migratory landbirds. *Ecological Restoration*, Vol.22, No.3, pp. 194-203
- Twedt, D. & Somershoe, S. (2009). Bird response to prescribed silvicultural treatments in bottomland hardwood forests. *Journal of Wildlife Management*, Vol.73, No.7, pp. 1140-1150
- Twedt, D. & Somershoe, S. (In press). Regeneration in bottomland forest canopy gaps six years after variable retention harvests to enhance wildlife habitat, *Proceedings of the Fifteenth Biennial Southern Silvicultural Research Conference*, J. Guldin, (Ed.), Hot Springs, Arkansas, USA, November 17-20, 2008
- Twedt, D.; Uihlein III, W. & Elliott, A. (2006). Habitat restoration for forest bird conservation: A spatially explicit decision support model. *Conservation Biology*, Vol.20, No.1, pp. 100-110
- Twedt, D. & Wilson, R. (2007). Management of Bottomland Forests for Birds, In: *Proceeding of Louisiana Natural Resources Symposium*, T.F. Shupe, (Ed.), 49-64, Louisiana State University AgCenter, Baton Rouge, Louisiana, USA, August 13-14, 2007
- Vandekerckhove, K.; De Keersmaeker, L.; Menke, N.; Meyer, P. & Verschelde, P. (2009). When nature takes over from man: Dead wood accumulation in previously managed oak and beech woodlands in north-western and central Europe. *Forest Ecology and Management*, Vol.258, No.4, pp. 425-435
- Van Wilgenburg, S. & Hobson, K. (2008). Landscape-scale disturbance and boreal forest birds: Can large single-pass harvest approximate fires? *Forest Ecology and Management*, Vol.256, No.1-2, pp. 136-146
- Verschuyt, J.; Riffell, S.; Miller, D. & Wigley, T. (2011) Biodiversity response to intensive biomass production from forest thinning in North American forests - A meta-analysis. *Forest Ecology and Management*, Vol.261, No.2, pp. 221-232

- Whitman, A. & Hagan, J. (2003). *Legacy Retention: A tool for retaining biodiversity in managed forests*. Mosaic Science Notes #2003-1, Manomet Center for Conservation Sciences, Brunswick, Maine, USA
- Wilson, R.; Ribbeck, K.; King, S. & Twedt, D., (Eds.). (2007). *Restoration, Management, and Monitoring of Forest Resources in the Mississippi Alluvial Valley: Recommendations for Enhancing Wildlife Habitat*. LMJV Forest Resource Conservation Working Group. Lower Mississippi Valley Joint Venture, Vicksburg, Mississippi, USA
- Zenner, E. (2000). Do residual trees increase structural complexity in Pacific Northwest coniferous forests? *Ecological Applications*, Vol.10, No.3, pp. 800–810



Global Perspectives on Sustainable Forest Management

Edited by Dr. Dr. Clement A. Okia

ISBN 978-953-51-0569-5

Hard cover, 300 pages

Publisher InTech

Published online 25, April, 2012

Published in print edition April, 2012

This book is dedicated to global perspectives on sustainable forest management. It focuses on a need to move away from purely protective management of forests to innovative approaches for multiple use and management of forest resources. The book is divided into two sections; the first section, with thirteen chapters deals with the forest management aspects while the second section, with five chapters is dedicated to forest utilization. This book will fill the existing gaps in the knowledge about emerging perspectives on sustainable forest management. It will be an interesting and helpful resource to managers, specialists and students in the field of forestry and natural resources management.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Daniel J. Twedt (2012). Wildlife Forestry, Global Perspectives on Sustainable Forest Management, Dr. Dr. Clement A. Okia (Ed.), ISBN: 978-953-51-0569-5, InTech, Available from:

<http://www.intechopen.com/books/global-perspectives-on-sustainable-forest-management/wildlife-forestry>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen