We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900
Open access books available

185,000
International authors and editors

200M
Downloads

Our authors are among the

154
Countries delivered to

TOP 1%
most cited scientists

12.2%
Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Chapter 9

Soil Physical Properties and Nitrous Oxide Emission from Agricultural Soils

Natalya P. Buchkina, Elena Y. Rizhiya, Sergey V. Pavlik and Eugene V. Balashov

Additional information is available at the end of the chapter

http://dx.doi.org/10.5772/53061

1. Introduction

1.1. N_2O in agriculture and microbial processes of nitrification and denitrification

N_2O is an important greenhouse gas as it contributes to global warming and the depletion of the stratospheric ozone layer (Bouwman, 1990; Crutzen, 1979; Houghton et al., 1990). At the present time the atmospheric concentration of N_2O is rising linearly at a rate of 0.3% per year (IPCC, 2007). Its current concentration in the atmosphere is equal to 319 ppbv (IPCC, 2007). A warming potential of 1 kg of N_2O is 310 times greater than 1 kg of CO_2 over a 100-year period (IPCC, 2007).

It is now widely accepted that agriculture is the main source of anthropogenic N_2O. The agriculture contributes to 60% of the global N_2O emissions (OECD. 2000). Agricultural soils are recognized as the major source of atmospheric N_2O, globally contributing 1.7-4.8 Tg N yr^{-1} (IPCC, 2007; Mosier, 1998).

N_2O is formed in two microbiological processes – nitrification and denitrification (Bremner, 1997; Firestone and Davidson, 1989) (Figure 1).

$$\begin{array}{cc} N_2O & N_2O \\ \uparrow & \uparrow \end{array}$$

$$NH_3 \rightarrow NH_2OH \rightarrow NO_2^- \rightarrow NO_3^- \text{ (a)}$$

Figure 1. Production of N_2O in soils during (a) nitrification and (b) denitrification processes. (Natalya P. Buchkina, Elena Y. Rizhiya, Sergey V. Pavlik, Eugene V. Balashov)

INTECH
open science | open minds

© 2013 P. Buchkina et al.; licensee InTech. This is an open access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The contribution of these processes to N_2O emission will vary with climate, soil management and properties (Granli, Bockman, 1994; Skiba, Smith, 2000). During nitrification N_2O is formed as a by-product in the reactions of oxidation of NH_3 to NH_2OH and NH_2OH to NO_2^- under strictly aerobic conditions. These two reactions are catalysed by such enzymes as ammonia monooxigenase and hydroxylamine oxidoreductase (McCarty, 1999; Wood, 1986). The reaction of NO_3^- production from NO_2^- is catalysed by nitrite oxidoreductase (Bock et al., 1986).

In contrast to nitrification, N_2O is an obligatory intermediate product of denitrification. Denitrification is a stepwise reduction of NO_3^- to N_2 by facultative anaerobic microorganisms. Enzymes catalysing these above-mentioned four reactions are nitrate reductase, nitrite reductase, nitric oxide reductase and nitrous oxide reductase (Hochstein, Tomlinson, 1988).

Despite denitrification is considered as a strictly anaerobic process, it is now accepted that N_2O can be produced during nitrifier denitrification (as a pathway of nitrification) under aerobic conditions. During a first part of nitrifier denitrification an oxidation of NH_3 to NO_2^- (nitrification) occurred, whereas a reduction of NO_2^- to N_2O and N_2 is a second part of the microbial process (Poth, Focht, 1985; Ritchie, Nicholas, 1972).

The rates of these processes and of N_2O production are controlled by several soil factors: soil water-filled pore space (WFPS), oxygen availability, pH, mineral nitrogen (NO_3^-, NH_4^+), available soil organic carbon (SOC), and temperature (Dobbie et al., 1999; Dobbie, Smith, 2003; Khalil et al., 2004., Wlodarczyk et al., 2003). Nitrification is the main process of N_2O emission at 35-60% WFPS. Denitrification becomes more important at soil water contents greater than 60% WFPS due to a decreased oxygen supply (Dobbie, Smith, 2003; Drury et al., 2003).

N_2O flux from agricultural soils depends on a complex interaction between climatic factors, soil properties and soil management. Management practices such as tillage systems and fertilizer applications can significantly affect the production and consumption of N_2O because of alteration in soil physical, chemical, and biochemical properties. The only way to reduce N_2O emissions from agricultural soils is to affect the soil properties through: 1. application of appropriate soil tillage system, 2. improving efficiency of N fertilizers with better placement, timing and rates of required N for growing relevant crops with high N uptake efficiency (Chatskikh, Olesen, 2007).

1.2. Role of soil tillage and soil physical properties in N_2O emission from soils

Tillage systems change soil structural, air, and water status (Beare et al., 1994; Green et al., 2007; Simansky et al., 2008). Nevertheless, continuous use of conventional moulboard ploughing in traditional farming systems can lead to degradation of soil structure and to losses in soil organic matter through erosion and oxidation (Lal, 2004; Lopez-Garrido et al., 2011; Norton et al., 2006; Reicosky, 2002). H. Riley et al. (2008) showed that a 15-year conventional ploughing of loamy soil caused a significant increase in bulk density from 1.29 to 1.43 Mg m^{-3} and a significant decrease in total porosity from 49.1 to 44.8%. E. Balashov and N. Buchkina (2011) showed that a 75-year use of conventional mouldboard ploughing re-

http://dx.doi.org/10.5772/53061

sulted in a significant decline in a concentration of SOC (from 43.1 to 30.2 g C kg^{-1} soil) as well as in an amount of water-stable aggregates (from 90.1 to 70.2%) and clay content (from 30.5 to 26.9%), compared to the fallow land in a clayey loam Haplic Chernozem.

There is a trend for the adoption of no-tillage and reduced tillage around the world, because these systems have been shown to improve soil organic matter status, structural state and water regime (Green et al., 2007; Simansky et al., 2008; Angers et al., 1997; Fernandez et al., 2010; Nyakatawa et al., 2000; Six et al., 2002; West, Post, 2002). V. Simansky (2008) reported that reduced tillage to a depth of 10-12 cm, compared to conventional tillage to the depth of 22-25 cm, led to a significant increase in the average amount of water-stable macro-aggregates of Orthic Luvisol. Long-term studies also suggested that the reduced tillage resulted in an essential change in soil structure that increased soil porosity and decreased bulk density (Zhang et al., 2007).

The conversion from mouldboard ploughing to no-tillage can result in a reduced N_2O emission (Almaraz et al., 2009) as well as in an increased N_2O emission from soils (Lemke et al., 1999). This unfavorable increase in N_2O emission can be caused by higher WFPS and bulk density of soils recently converted to no-tillage system (Liu et al., 2007).

Direct measurements of N_2O flux from soils with different tillage systems have shown that their effects on the greenhouse gas emission are governed by climatic conditions. According to the results reported in (Six et al., 2002), a conversion of conventional tillage to no-tillage resulted to an increase in soil organic matter level in temperate and tropical soils in a wide range of agroecosystems. In the temperate soils an increase in N_2O emissions and CH_4 uptake was observed under no-tillage compared with conventional tillage. In general, light-textured, well-aerated soils under drier climate conditions do not produce more N_2O when under no-tillage or reduced tillage system compared to conventional tillage. However, poorly aerated soils in wet climate can produce more N_2O when conventional tillage is converted for no-tillage or reduced tillage system (Chatskikh, Olesen, 2007; Ball et al., 1999; Choudhary et al., 2002).

Changes in the total porosity and the amount of water-stable aggregates may result in a formation of favorable conditions for denitrification as it can take place in soils with high availability of organic matter and low availability of oxygen. Oxygen is regarded as a key determinant of denitrification rates, while the importance of available SOC and NO_3^- (or other nitrogen oxides) will vary in the ecosystems (Robertson, Groffman, 2007). A proportion of the total gaseous products of denitrification that is actually emitted to the atmosphere as N_2O depends on the soil aggregation and water content (Smith et al., 2003). A destruction of soil aggregates by tillage can lead to an increase in availability of SOC for soil denitrifying microorganisms (Gregorich et al., 1989; Petersen et al., 2008).

Despite positive effects of no-tillage and reduced tillage systems on soil aggregation and porosity, denitrification can occur even in the soil aggregates under soil aerobic conditions (Smith, 1980). K. Khalil (2004) reported that significant denitrification rates (i.e. greater than 0.2 mg N kg^{-1}) from 2-3-mm aggregates were observed in the 0 and 0.35 kPa O_2 pressure treatments, whereas nitrification (both NH_4^+ and NO_2^- oxidation rates) was reduced by a fac-

tor of 4–10 when O_2 concentration decreased from 20.4 to 0.35 kPa. According to the results reported by Uchida et al. (2008), the highest N_2O-N fluxes of 9920 μg m^{-2} h^{-1} occurred in the 0–1.0 mm aggregates, while the respective values in the 1.0–2,0; 2.0–4.0 and 4.0–5.6 mm aggregates were 3985, 2135 and 2750 μg m^{-2} h^{-1}.

An absence of tillage can lead to soil compaction and, as a result, to a reduced air-filled porosity and availability of oxygen (Rasmussen, 1999). At high soil water content, the effects of increasing soil bulk density may contribute to a formation of anaerobic conditions (Douglas, Crawford, 1998). R. Ruser et al. (1998) reported that an increase in bulk density of fine silty soil only by 20% led to increasing N_2O emission. According to the results reported by M. Beare et al. (2009), there was a strong exponential increase in N_2O production above 0.60 cm^3 cm^{-3} WFPS of a compacted fine loamy soil where denitrification was a dominant pathway of N_2O production.

Therefore, tillage systems can have contradictory effects on soil physical status and on N_2O emissions from soils. If it is necessary to obtain valid information on N_2O emissions from soils differing in soil physical status and tillage systems the site-specific measurements need to be carried out in various agroecosystems with different agricultural uses.

1.3. Role of increasing nitrogen rates in N_2O emission from soils

Application of mineral N-fertilizers into agricultural soils usually results in increasing N_2O emissions (Smith et al., 2003; Jones et al., 2007; MacKenzie et al., 1998; Rizhiya et al., 2011). However, there is contradictory information on linearity between applied N rates and N_2O emissions from soils. According to results reported by Gregorich et al. (2005), N_2O emission from agricultural soils increased linearly with the applied amount of mineral N fertilizer. At N rates not exceeding or equal to those required for maximum yields, N rates tended to create a linear response in N_2O emissions, with approximately 1% of applied mineral N lost as N_2O (Bouwman, 1996; Halvorson et al., 2008).

However, there are threshold N rates, which can exceed the crop and ecosystem uptake capacity (Grant et al., 2006). At N rates exceeding crop requirements, N_2O emissions are more variable and often rise exponentially with increasing N fertilization (Hobet et al., 2011; Snyder et al., 2009). C.P. McSwiney and G.P. Robertson (2005) reported on nonlinear increase between N_2O emission and N rates. N_2O emission was moderately low (20 g N_2O-N ha^{-1} day^{-1}) at N rate of 101 kg N ha^{-1}, but at N rate above 134 kg N ha^{-1}, N_2O emissions sharply increased to about 450 g N ha^{-1} day^{-1}. The greatest percentage of fertilizer N lost as N_2O (7%) occurred at the N rate of 134 kg N ha^{-1}. The authors also (1) found that the proportion of applied N lost as N_2O decreased to 2–4% (as emission factor) with N rates exceeding 134 kg N ha^{-1} and (2) stated that this threshold of N_2O response to N fertilization could be reduced with no or little yield penalty by reducing N fertilizer inputs to levels that just satisfy crop needs.

According to J.W. Van Groenigen et al. (2010), agricultural management practices to reduce N_2O emissions should focus on optimizing fertilizer-N use efficiency under median rates of

N input, rather than on minimizing N application rates. The authors proposed to assess the N_2O emissions as a function of crop N uptake and crop yield.

1.4. Role of crop roots in N_2O emission from soils

Living roots of crops can play an integral role in soil N_2O production by providing their labile organic exudates and mineral N to microbial community involved into nitrificarion and denitrification. J.A. Bird (2011) reported that living roots of *Avena barbata* increased the turnover and loss of belowground ^{13}C compared with unplanted soils. The rhizosphere of *Avena barbata* shifted the microbial community composition, resulting in greater abundances of gram-negative markers and lower abundances of gram- positive, actinobacteria and cyclopropyl PLFA markers compared to unplanted soil. According to the results of Khalid et al. in (Khalid et al., 2007), the presence of grass species significantly increased the size of a number of dissolved nutrient pools in comparison to the unplanted soil (e.g. dissolved organic carbon, total phenolics in solution) but had a little affect on other pools (e.g. free amino acids).

The plant roots change such soil physical properties as air porosity, pore distribution, penetrometer resistance, permeability and bulk density (Williams, Weil, 2004; Mitchel et al., 1995; Balashov, Bazzoffi, 2003). Balashov and Bazzoffi (2003) showed that the resilience capacity of wheat root system at the boot stage was sufficient for the formation and stabilization of water-stable aggregates in a sandy loam Typic Udorthents and a clayey loam Vertic Xerorthent compacted with ground contact pressures of 51 and 103 kPa. After the compaction with ground contact pressure of 154 kPa, the root system was unable to maintain the water-stable aggregation of soils even at its initial level. Therefore, roots of plants can improve soil physical status of compacted soils and can reduce a risk of formation of favorable conditions for denitrification, which are observed at WFPS exceeding 60%.

A rather high attention has been given to the dynamics of N_2O profile concentration and its subsequent diffusion to the soil surface. The concentrations and fluxes of N_2O are dependent on WFPS, mineral N content, microbial activity, temperature, and macro-porosity in the soil profiles (Burton, Beauchamp, 1994; Clough et al., 2005; Jacinthe, Lal, 2004; Kusa et al., 2010; Li et al., 2002; Muller et al., 2004; Phillips et al., 2012; Van Groenigen et al., 2005; Velthof et al., 1996; Yoh et al., 1997).

J.W. Van Groenigen et al. (2005) found that maximum N_2O concentrations were observed at depths of 48 and 90 cm at WFPS > 70% in a sandy soil. The authors reported that during the summer the maximum N_2O fluxes from the soil surface were coinciding with high subsoil N_2O concentrations suggesting that denitrification in the subsoil was a main determinant of N_2O formation. In winter with low air and soil temperatures as well as with low soil microbial activity high N_2O concentrations in the subsoil did not lead to corresponding high fluxes of N_2O from the soil surface.

K. Kusa et al. (2010) showed that the contribution ratios of the N_2O produced in the top soil (0–0.3 m depth) to the total N_2O emitted from the soil to the atmosphere in the treatments with the Gray Lowland soil and the Andosol were 0.86 and 1.00, respectively, indicating that

the N_2O emitted from the soils to the atmosphere was mainly produced in the top soil. Variations in the profile N_2O concentrations between two soils were caused by those in the soil structure, in particular, because of the presence of macro-pores and cracks in the soil structure, which resulted in different production and movement of N_2O in the soils. The higher N_2O production in the subsoil of the Gray Lowland soil could have been activated by NO_3^- leaching through macro-pores and cracks, and subsequently the N_2O produced in the subsoil could have been rapidly emitted to the atmosphere through the macro-pores and cracks.

The aim of this study was to quantify direct N_2O emissions from a light-textured arable soil under several crops grown on ridges, in North-Western Russia, that was typical of the region, during four growing seasons, with special attention to soil physical properties.

2. Materials and methods

The measurements of N_2O emission were conducted at the Menkovo Experimental Station of the Agrophysical Research Institute in the St. Petersburg region of Russia (59°34'N, 30°08'E) in 2004–2007. Average annual rainfall in the area for the previous 10 years was 1,109 mm, with 50–60% falling during the growing season (May–September). Average annual air temperature was +4.5°C, and average air temperature in the growing season was +13.6°C. The studied soil was the most typical soil of the area: a sandy loam Spodosol containing 7–8% clay, 17–23 g C kg^{-1} soil, and with bulk density ranging from 0.95 to 1.5 g cm^{-3}. Three 0.5-ha fields containing this soil were used to conduct the experiment. The first of them received 160 t ha^{-1} (656 kg of total N ha^{-1} or 219 kg of available N ha^{-1}), the second – 80 t ha^{-1} (328 kg of total N ha^{-1} or 108 kg of available N ha^{-1}) of farmyard manure (FYM) in spring of 2003 and early summer of 2004, and the third received no FYM.

Part of the field with no FYM was allocated for potato (*Solanum tuberosum* L.) in 2004 and 2005. Part of the field with the highest application of FYM was allocated for potato in 2005. All the three fields were used to grow cabbage (*Brassica oleracea capitat* L.) in 2006 and carrot (*Daucus carota* L.) in 2007. By 2005 the SOC content in the soils of the plots receiving 160 t ha^{-1} of FYM, 80 t ha^{-1} of FYM and no FYM at the beginning of the experimental studies was equal to 21, 19 and 17 g C kg^{-1} soil, respectively.

In 2004-2005 large single plots were used for the experiment while smaller but randomized replicate plots were used in 2006 and 2007. The use of single plots earlier in the experiment was a necessary consequence of the cropping arrangements adopted by the manager of the experimental station. The experimental study described here was therefore the only available way in which to conduct this project. Mineral N fertilizer (ammonium nitrate, AN) was applied to the soil at different rates (Table 1). The mineral N was applied (broadcast) to the soil in granular form mechanically: before the furrows and ridges were formed (1st application) and the second and third time (for potato crop only) the fertilizer was applied to furrows. The depth of mouldboard ploughing for all the plots was 22–24 cm. Most of the plots were ploughed in spring between 13 and 20 May. Crops were planted within a day or two after mouldboard ploughing and harvested at the end of August/early September.

http://dx.doi.org/10.5772/53061

The closed chamber technique was used to measure direct N_2O fluxes from the soil. Gas samples were collected two-three times a week (between noon and 2 pm) throughout the growing seasons (May-September) of 2004–2007. Chambers were made of inverted cylindrical plastic buckets, 18.9 cm in diameter and 11 cm high, and put into the field every time before gas samples were collected (Figure 2). Eight replicate chambers were used at all the plots — four in the furrows and four on the ridges.

Year	Crop	SOC, g C kg^{-1} soil	Ammonium nitrate fertilizer application					
			1st	Date	2nd	Date	3rd	Date
2004	Potato	17	72	14/05	48	17/07	0	
2005	Potato	17	72	13/05	38	10/06	10	27/07
		21	72	13/05	38	10/06	10	27/07
2006	Cabbage	17	0		0		0	
		17	70	15/05	0		0	
		19	0		0		0	
		19	90	15/05	0		0	
		22	0		0		0	
		22	110	15/05	0		0	
2007	Carrot	17	40	20/05	0		0	
		17	110	20/05	0		0	
		19	40	20/05	0		0	
		19	120	20/05	0		0	
		23	60	20/05	0		0	
		23	130	20/05	0		0	

SOC – soil organic carbon

Table 1. Crops grown and rates of mineral fertilizers used during the growing seasons of 2003-2007 (Natalya P. Buchkina, Elena Y. Rizhiya, Sergey V. Pavlik, Eugene V. Balashov).




Figure 2. Chamber placement on the ridges and in the furrows of the potato (*Solanum tuberosum* L.), cabbage (*Brassica oleracea capitat* L.) and carrot (*Daucus carota* L.) plots. (Natalya P. Buchkina, Elena Y. Rizhiya, Sergey V. Pavlik, Eugene V. Balashov)

Chambers were pressed into the soil to a depth about 2 cm and the soil outside the chambers was compacted against the chamber walls to make them gas-tight. A three-way tap on the top of each chamber was closed only after the chamber was fixed into the topsoil, to avoid extra air pressure inside the chamber. After 60 min gas samples were collected via the three-way tap. Similar sampling at the end of the closure period has been employed elsewhere (Ball et al., 2007; Hergoualc'h, 2008).

Gas samples were taken with a 60-ml syringe; 50 ml were flushed through a 10 ml vial that had been previously flushed with air, then the remaining 10 ml were forced into the vial to over-pressurise it. 3-ml subsamples from the vials were analysed for N_2O in a Carlo Erba 4130 gas chromatograph (GC) fitted with an electron capture detector. The GC was calibrated with standard gas mixtures. The detection limit for the sampling/analytical system was 0.05 ppm. Daily fluxes were calculated using the arithmetic mean of the four replicate chambers. Cumulative N_2O fluxes for different months as well as for whole growing seasons of 2004–2007 were calculated by plotting daily fluxes through time, interpolating linearly between them, and integrating the area under the curve. Cumulative standard errors were also determined by the same way (Dobbie, Smith, 2003; Buchkina et al., 2010).

The climatic factors most likely to affect the amount of N_2O produced in the soil were monitored during the experiment. Data on daily rainfall and air temperatures (maximum, minimum, and average) were collected from the Institute's meteorological station, situated c. 100 m from the experimental plots. Soil temperature at a 10-cm depth was measured with a digital thermometer on each sampling occasion. Soil bulk density and water content were meas-

http://dx.doi.org/10.5772/53061

ured (in three replicates) 1–2 times a month by standard methods (Rastvorova, 1988; Soil Survey, 1996) and the results were used to calculate water-filled pore space (WFPS). Soil samples for mineral nitrogen (NH_4^+ and NO_3^-), were also collected 1–2 times a month from the 0–10 cm layer. At least 10 subsamples were collected from each plot and combined. The composite samples were dried at room temperature and then amounts of NH_4^+ and NO_3^- were measured in water extractions using ion-selective electrodes in three replicates (Bankin et al., 2005). SOC content was measured by a wet combustion method using potassium dichromate ($K_2Cr_2O_7$) solution in sulfuric acid (Soil Survey, 1996).

All the measurements of the soil properties were done in three replicates. Means and standard deviations were calculated for each parameter within each treatment. Significance of differences between treatments was estimated by analysis of variance (one-way ANOVA) at $p \leq 0.05$. Relationships between the soil parameters were assessed with a linear regression analysis using computer statistical package.

3. Results and discussion

3.1. Weather conditions

The amount of precipitation and its distribution over the growing season (1 May–30 September) varied between the studied years significantly (Figure 3). The wettest growing season was 2004 and the driest 2006, with 845 and 456 mm of rain falling, respectively, compared to 575 mm in 2005, and 555 mm in 2007. During the growing season there were 75 days with rain in 2004, 55 in 2005, 41 in 2006 and 52 in 2007. Almost 75% of the rain fell on days when precipitation was greater than 10 mm: there were 25 such occasions in 2004, 15 in 2005, 13 in 2006 and 20 in 2007. The longest dry spells were observed in 2006: 27 days in late April-May, 15 days in July and 11 days in August. Even the wettest growing seasons of 2004 had a 10-day dry spell in May. The growing seasons of 2005 and 2007 had their longest dry spells in July (9 and 11 days, respectively) and in August-September (12 and 16 days, respectively). Amount of rainfall exceeded 100 mm in June, July, August and September of 2004, in May, June, and July of 2005, in May of 2006, in May and July of 2007.




Figure 3. Monthly rainfall during the growing seasons of 2004-2007. (Natalya P. Buchkina, Elena Y. Rizhiya, Sergey V. Pavlik, Eugene V. Balashov)

Average air temperature for the studied growing seasons was more stable than the precipitation. July and August were always the warmest months, (16–20°C). In May, June and September the average was always between 10 and 15°C. The warmest growing season overall was 2006, averaging 15°C, compared to 14.4°C in 2005 and 2007, and to 13.9°C in 2004. In the growing season of 2005 the minimum air temperature never fell below zero but in all the other growing season it did: to -0.5°C at the end of May of 2004 and to -2.1°C in early May of 2006 and 2007. Maximum air temperatures were 27.2°C in 2004, 29.2°C in 2005, 33.8°C in 2006 and 31.4°C in 2007.

3.2. Soil properties

Soil bulk density varied during the growing seasons of 2004-2007 from 0.9 to 1.4 g cm^{-3} being the highest in potato furrows during the wettest growing season of 2004 and the lowest on cabbage ridges during the driest growing season of 2006 (Figure 4). The differences in soil bulk density between ridges and furrows were higher in the wetter growing seasons of 2004, 2005 and 2007 than in dryer growing season of 2006. That must be a result of soil compaction in the furrows during crop management operations. It was shown that the studied soils were much more easily compacted when wet than when they were relatively dry. In wetter growing seasons the differences in soil bulk density between ridges and furrows were greater in spring and early summer time when the ridges were formed for crop planting and mechanical cultivation of soil was regularly done (as a weed control operation). It also coincided with higher monthly rainfalls at the beginning of the wetter growing seasons. Later in the season the differences in soil bulk density between ridges and furrows were smaller. During the driest growing season of the four – 2006 – the difference in soil bulk density between ridges and furrows was less pronounced and did not change much during the season. The difference in soil bulk density between soils with the low (19 g C kg^{-1} soil) and high (22 g C kg^{-1} soil) SOC content was also often significant with more fertile soils having lower bulk density than the less fertile soils. Still, the differences in bulk density between soil ridges and furrows were almost always more significant than the differences in that for ridges (or furrows) between soils with different SOC content. The significantly higher bulk density of soil furrows, compared to that of soil ridges, could result in a formation of soil physical conditions more favorable for microbial process of denitrification in the studied soil (Beare et al., 2009).

The soil water content during the four studied growing seasons changed from 8 to 36% (of weight) and was higher than 20% for most of the wettest growing season of 2004, varied from 8 and 36% and from 8 and 31% during the growing seasons of 2005 and 2007, respectively, and never exceeded 24% during the driest growing season of 2006 (Figure 5). The differences in soil water content between ridges and furrows were greater in wetter growing seasons of 2004, 2005 and 2007 than in the driest of the four growing seasons of 2006. A combination of greater water content and higher bulk density in the soil of furrows as compared to the soil of ridges could lead to decreasing oxygen availability and to increasing N_2O fluxes as the intermediate of accelerated process of denitrification under such soil physical conditions (Drury et al., 2003; Beare et al., 2009).

http://dx.doi.org/10.5772/53061




Figure 4. Soil bulk density (BD) in furrows and on ridges at the plots with potato (2004, 2005), cabbage (2006) and carrot (2007) at different soil organic matter (SOC) content (17, 19 and 21 g C kg^{-1} soil). (Natalya P. Buchkina, Elena Y. Rizhiya, Sergey V. Pavlik, Eugene V. Balashov)




Figure 5. Soil water content (W) in furrows and on ridges at the plots with potato (2004, 2005), cabbage (2006) and carrot (2007) at different soil organic matter (SOC) content (17, 19 and 21 g C kg^{-1} soil) (Natalya P. Buchkina, Elena Y. Rizhiya, Sergey V. Pavlik, Eugene V. Balashov).

The relationship between soil water content (average for a month) and monthly precipitation is shown on Figure 6 and 7. A linear correlation ($p < 0.05$) between the two parameters was higher for the soil with the lower SOC content than for the soil with the higher SOC (especially for furrows). The water-holding capacity of the studied soils was not very high as they were light- textured and should mainly depend on rainfall and SOC content. The higher SOC of a light-textured soil the more water it would be able to hold and would be less dependent upon rainfall if the dry spells were not very long.




Figure 6. Relationships between monthly precipitation and average (for a month) soil water content (W, % of weight) for (a) soil ridges and (b) soil furrows with the highest (22 g C kg^{-1} soil) soil organic carbon content during the growing seasons of 2004-2007 (Natalya P. Buchkina, Elena Y. Rizhiya, Sergey V. Pavlik, Eugene V. Balashov)

During the studied growing seasons the soil WFPS varied widely within and between years (Figure 8). In general, the soils of all plots were wetter in 2004 than in 2005, 2006, and 2007 with WFPS for most of the growing seasons varying from 35 to 70%, compared to 15–50% in 2005, 15–40% in 2006, and to 20–45% in 2007. In 2004, in contrast to all the other studied seasons, maximum values of WFPS were observed in June and July (40–85%). During the wettest part of the growing season soil in furrows was 20–40% wetter than soil on ridges (60–85% versus to 40–45%, respectively). In May and August of 2004 the soil WFPS was lower than in the middle of the summer, changing from 20 to 40% and from 38 to 58%, respectively. The difference in soil water content between ridges and furrows during the drier part of the season was 10–20%. In 2005, the lowest values of soil WFPS were observed at the end of July, August and September, when WFPS on potato ridges went down to 10–22%, being the lowest of the year. During this dry spell the soil WFPS in potato furrows varied between 19 and 38%. The highest values of soil WFPS for the whole growing season of 2005 were observed in May and June but even then they were varying between 38 and 62%, and the difference between furrows and ridges was 11-13%. In 2006, the driest growing season of all, soil WFPS was very low in July (15-20%). The highest values of soil WFPS, but still quite low compared to the highest values of the other growing seasons, were observed in May and September (33-43%). The difference in soil WFPS between furrows and ridges during the growing season of 2006 was never higher than 8%. In 2007, the lowest values of the soil WFPS were found at the end of May-early June and at the end of June – early August (15-25%). Early May and August of 2007 were wetter but the highest values of the soil WFPS were not exceeding 53%. The difference in soil WFPS on ridges and furrows in 2007 was 7-10% during dry spells and 10-15% during wetter spells of the season. As was indicated

http://dx.doi.org/10.5772/53061

above, nitrification is the main process of N_2O emission at 35-60% WFPS, but denitrification is more important at soil water content greater than 60% WFPS due to a decreased oxygen supply (Drury et al., 2003; Dobbie, Smith, 2003). Therefore, the observed soil moisture conditions were often more favorable for nitrification rather than denitrification during the growing seasons of 2004-2007.




Figure 7. Relationships between monthly precipitation and average (for a month) soil water content (W, % of weight) for (a) soil ridges and (b) soil furrows with the lowest (19 g C kg^{-1} soil) soil organic carbon content during the growing seasons of 2004-2007 (Natalya P. Buchkina, Elena Y. Rizhiya, Sergey V. Pavlik, Eugene V. Balashov).




Figure 8. Soil water-filled pore space (WFPS) for furrows and ridges for potato (2004 and 2005), cabbage (2006) and carrot (2007) at different soil organic matter (SOC) content (17, 19 and 21 g C kg^{-1} soil) (Natalya P. Buchkina, Elena Y. Rizhiya, Sergey V. Pavlik, Eugene V. Balashov).

The linear correlation between monthly rainfall and soil WFPS (average for a month) was strong for soil furrows as well as for soil ridges only during the driest growing season of 2006 (R^2=0.80-0.88, p<0.05). For the growing season of 2007 this relationship was strong only for soil ridges (R^2=0.92-0.94, p < 0.05) but not for furrows (R^2 = 0.65-0.81, p = 0.1-0.19) while in 2004 it was strong for soil furrows (R^2=0.95, p < 0.01) but not for ridges (R^2=0.57, p = 0.14). The relationship between monthly rainfall and soil WFPS (average for a month) for all 5 growing seasons is shown on the Figure 9. In average, the soil WFPS in furrows is more sensitive to monthly rainfall (R^2=0.64, p < 0,001) than that on ridges (R^2=0.37, p < 0,001). Our results corresponded with the data reported by L. Meng et al. (2005) where the strong correlations between WFPS and precipitation were also observed.

According to T. Granli and O.C. Bockman (1994) optimum soil temperatures for N_2O emissions vary between 25-40°C. During the growing seasons soil temperatures at 10 cm depth varied between 3 and 26°C, and were highest (15–26°C) always at the end of July and in early August. Average soil temperature was almost 1°C lower during the growing season of 2004 (14.3°C) and 2007 (14.4°C) than of 2003 (15.1°C), 2005 (15.2°C) and 2006 (15.0°C). Over the 4 years, higher soil temperatures were always found in those plots where soil was more exposed to the sun. The soil temperature was often higher in the ridges than in the furrows.

The experimental soil was very low with mineral N content when N-fertilizers were not used and contained 2–9 mg N kg^{-1} soil at the beginning of all studied growing seasons. Soil mineral N concentrations always increased after N-fertilizer was applied to the soil to 20-100 mg N kg^{-1} soil but were never high for longer than 3 weeks and were always back to below 10 mg N kg^{-1} soil at the end of the growing seasons.




Figure 9. Relationships between monthly rainfall and soil water filled pore space (WFPS, average for a month) for (a) soil ridges and (b) soil furrows during the growing seasons of 2004-2007 (Natalya P. Buchkina, Elena Y. Rizhiya, Sergey V. Pavlik, Eugene V. Balashov).

K. Dobbie and K.A. Smith (2003) reported that high N_2O emission could be even observed at WFPS (65%), soil temperature (4.5°C) and NO_3^--N content (5 mg kg^{-1} soil). The results of our studies showed that these threshold limits had been observed or exceeded during the growing seasons of 2004-2007.

http://dx.doi.org/10.5772/53061

3.3. N_2O emissions

3.3.1. Daily fluxes

During the growing seasons 0f 2004-2007 daily N_2O fluxes varied highly between plots differed in fertilization, soil physical conditions and crops. Agricultural soils without any fertilization can produce N_2O during the growing season because of mineralization of soil organic matter and N rich plant residues. In our study the daily N_2O fluxes for the four studied growing seasons were never higher than 10 g N_2O–N ha^{-1} on the unfertilized plots (both in furrows and on ridges). Conversely, daily N_2O fluxes from the soil could be very low even on fertilized soils (especially on ridges), indicating that other parameters, for example, soil mineral N content, temperature, soil WFPS and oxygen availability were main determinants controlling production of N_2O as the by-product of nitrification (Khalil et al., 2004; Meng et al., 2005; Conen et al., 2000).

Our data corresponded with the results published by H. Flessa and P. Dorsch (1995) and also by B. Ball et al. (2002) who found that application of N-fertilizers by itself did not always enhance N_2O emission and that weather conditions at the time of fertilizer application influenced N_2O emissions substantially. Nevertheless, C. Henault et al. (1998) reported that N_2O emissions were higher 20 g N ha^{-1} day^{-1} from soils with low water content as a result of domination of nitrification.

Possible differences in N_2O emissions from soils with different crop types depend on the management practices and climatic conditions. Tillage, sowing and other cultivations for agricultural plants can disturb soil aggregates, increase soil organic matter mineralization and deteriorate soil physical status and, as a result, can lead to an increased N_2O emissions (Elder, Lal, 2008; Ruser et al., 1998; Beare et al., 2009; Six et al., 1998). Crops can (1) demonstrate different rates of soil water uptake, (2) prevent the accumulated soil water against evaporation under different weather conditions, and (3) remove more N from the soil that will result in different availability of mineral N for nitrifying and denitrifying microorganisms.

During the growing seasons of 2004-2007 daily N_2O fluxes from the soil with crops grown on ridges (potato, carrot, cabbage) were often higher than those from the soil with cereals, grasses, grass-clover mixture and grass with oats (the latter crops grown in the same experiment are not being discussed in this paper but were described earlier in our paper - Buchkina et al., 2010).

For the latter crops daily N_2O fluxes were never exceeding 20 g N_2O-N ha^{-1} while for the crops grown on ridges N_2O fluxes from the soil of furrows could be as high as 115 g N_2O-N ha^{-1} day^{-1} (Figure 10). Higher N_2O emissions from ridges can be explained by a greater mineralisation of soil organic matter that resulted to an increased N_2O emission from a bare soil subjected to a higher warming (Maljanen et al., 2002). According to the results reported by M. Maljanen et al. (2002), higher N_2O fluxes were observed from soils kept bare by tillage or cutting presumably from lack of competition for nitrate (NO_3^-) between microbes and plants.

Daily N_2O fluxes during the growing seasons of 2004-2007 were never affected by WFPS if no N had been applied with the mineral fertilizers as the original soil was very low in available N. However, in the N-fertilized plots the greatest N_2O fluxes were found when WFPS

exceeded 40% (Figure 11) and that happened more often in the soil of furrows with the higher soil bulk density rather than in the soil of ridges with the lower soil bulk density. The highest N_2O emission in the cultivated soils generally are observed at 70–90% WFPS due to denitrification which rapidly increased when WFPS exceeded 60%, whereas at 30–70% WFPS nitrification was the main source of N_2O (Granli, Bockman, 1994; Smith et al., 2003; Maljanen et al., 2002).

It was shown for field experiments in Scotland (Dobbie et al., 1999; Conen et al., 2000) that high N_2O fluxes actually occurred more often when WFPS was higher than 60%. In our experiment the soil WFPS quite rarely exceeded 60% (the longest period for the four growing seasons was about 20 days in 2004 and only for potato furrows) as the rainfall of the area was less than it was in Scotland and also the soil of the experimental plot was light-textured with relatively easy drainage down the soil profile. The work of E.A. Davidson (1991) suggests that denitrification played its role in N_2O emissions from our soil only occasionally, when there were anaerobic conditions in some parts of the soil, while the main source of N_2O emission most of the time was nitrification. The same results were reported in our earlier work (Buchkina et al., 2010) for other crops (cereals, grasses, and grass-legume mixtures) at our experimental station.

Soils in the ridges and furrows could differ in bulk density, air and total porosity, available SOC, mineral N and oxygen availability. For instance, the average bulk density of the soil on ridges and in furrows was equal to 0.9 and 1.4 g cm^{-3}, respectively. Therefore, the soil furrows could show more favorable conditions for denitrification before and after a rainfall. Our results corresponded with the data of Beare et al. (2009) who showed that most (88%) of the total N_2O production from compacted soil occurred after soil rewetting, at a time when there was ample NO_3^--N and dissolved organic carbon available and the air-filled porosity was low (0.22 cm^3 cm^{-3}), rather than during the drying phase when compacted soil was accumulating NO_3^-N and air-filled porosity was high (0.62 cm^3 cm^{-3}). P. Merino et al. (2012) also reported that the interaction of WFPS and soil NO_3^- content was statistically significant ($p < 0.001$), indicating that the response of N_2O emission from a loamy clay soil to changes in NO_3^- content was very dependent on WFPS.




Figure 10. Daily N_2O fluxes from the soil under (a) cereals, grasses, grass-legume mixtures, grass-cereal mixtures during the growing seasons of 2003-2005 (Buchkina et al., 2010) and (b) crops grown on ridges (potato, cabbage, carrot) during the growing seasons of 2004-2007 (Natalya P. Buchkina, Elena Y. Rizhiya, Sergey V. Pavlik, Eugene V. Balashov).

http://dx.doi.org/10.5772/53061




Figure 11. Soil water-filled pore space (WFPS) and daily N_2O fluxes (both from furrows and ridges) from the soil with potato (2004, 2005), cabbage (2006) and carrot (2007): (a) unfertilized plots, (b) fertilized plots (Natalya P. Buchkina, Elena Y. Rizhiya, Sergey V. Pavlik, Eugene V. Balashov).

3.3.2. Cumulative fluxes and emission factors

Cumulative N_2O fluxes for the growing seasons of 2004-2007 from the studied soil for different crops varied from 0.34±0.09 to 1.83±0.45 kg N_2O-N ha^{-1} (Table 2). Within the same growing season N_2O cumulative fluxes were higher from the soils of furrows than from the soils of ridges (especially, on the plots where mineral nitrogen was applied with fertilizers) but the difference was not significant when all the four growing seasons were taken into account: the cumulative N_2O fluxes from soils in furrows varied from 0.37±0.10 to 1.83±0.45 kg N_2O-N ha^{-1} while from soils on ridges – from 0.36±0.011 to 1.25±0.29 kg N_2O-N ha^{-1}.

The lowest cumulative N_2O fluxes for both positions were measured for the growing season of 2006 – the driest of the four. During this growing season cumulative N_2O fluxes from soils of furrows were often even lower than those from soils of ridges in wetter growing seasons (Table 2, Figure 12). During the dry growing season of 2006 cumulative N_2O fluxes were not affected by N-fertilizer rates on the soils having SOC content of 17 and 19 g C kg^{-1}. On the plot containing 23 g C kg^{-1} only the soil in furrows emitted more N_2O for this growing season when higher amount of N-fertilizer was applied to it. During the dry growing season the compacted soil in the furrows demonstrated greater soil water content than the uncompacted soil on the ridges. Hence, the first one had a higher amount of smaller pores than the latter one with a higher amount of larger pores. Therefore, during this growing season the compacted furrow soil with smaller pores could (1) hold more water, (2) have more favorable conditions for nitrification and denitrification, and (3) response in the higher values of N_2O emissions to the mineral N fertilization than the uncompacted ridge soil.

The cumulative N_2O fluxes from the soils showed the higher values during the wetter growing season of 2007 compared to the driest growing season of 2006 when there was no relationship between fertilizer rates and cumulative N_2O fluxes. In 2007, an increase in a rate on mineral N-fertilizer from 40 kg N ha^{-1} to 110 kg N ha^{-1} did not result in an increase of cumulative N_2O fluxes from the soil on ridges under carrot (Table 2). Only the rates of 120 and 130 kg N ha^{-1} caused a significant increase of cumulative N_2O fluxes from this soil. Thus, the re-

lationship of the cumulative N_2O fluxes with the mineral N rates was nonlinear for soil on ridges for the growing season of 2007. The cumulative N_2O fluxes from the soil in furrows under carrot were significantly higher than those from the ridges, and demonstrated a non-linear and unstable increase with increasing N rates from 40 kg N ha^{-1} to 130 kg N ha^{-1}. However, the highest cumulative N_2O fluxes from the soil both on ridges and in furrows for the growing season of 2007 were observed at the N rate of 130 kg N ha^{-1}.

Year	Crop	Crop yield, kg ha^{-1}	SOC, g kg^{-1} soil	N applied, kg N ha^{-1}	Position	Cumulative N_2O flux, kg N_2O-N ha^{-1}	Emission factor
2004	Potato	17000	17	120	Ridges	1.13±0.39	0.94
					Furrows	1.51±0.69	1.26
2005	Potato	18000	17	120	Ridges	0.60±0.21	0.50
					Furrows	0.91±0.35	0.76
	Potato	21000	21	120	Ridges	1.25±0.29	1.04
					Furrows	1.83±0.45	1.52
2006	Cabbage	41400	17	0	Ridges	0.37±0.08	-
					Furrows	0.45±0.11	-
	Cabbage	59900	17	70	Ridges	0.36±0.05	0.51
					Furrows	0.42±0.08	0.60
	Cabbage	45040	19	0	Ridges	0.43±0.06	-
					Furrows	0.53±0.07	-
	Cabbage	75900	19	90	Ridges	0.45±0.07	0.50
					Furrows	0.56±0.08	0.62
	Cabbage	81100	22	0	Ridges	0.34±0.09	-
					Furrows	0.37±0.10	-
	Cabbage	103320	22	110	Ridges	0.36±0.11	0.33
					Furrows	0.60±0.12	0.54
2007	Carrot	54830	17	40	Ridges	0.62±0.02	1.55
					Furrows	0.65±0.07	1.63
	Carrot	61220	17	110	Ridges	0.64±0.03	0.58
					Furrows	0.93±0.07	0.85
	Carrot	68110	19	40	Ridges	0.71±0.01	1.78
					Furrows	0.92±0.10	2.3
	Carrot	70170	19	120	Ridges	0.74±0.01	0.62
					Furrows	0.99±0.11	0.83
	Carrot	74570	23	60	Ridges	0.65±0.03	1.08
					Furrows	1.09±0.05	1.82
	Carrot	70040	23	130	Ridges	0.80±0.04	0.62
					Furrows	1.45±0.14	1.12

SOC – soil organic carbon

Table 2. Crops, amounts of mineral N-fertiliser applied, cumulative N_2O fluxes and emission factors.

http://dx.doi.org/10.5772/53061

According to several authors, there is the threshold of N rates which can exceed the N requirements of crops (Granli, Bockman, 1994) and if the N rates exceed crop requirements, N_2O emissions can become more variable and increase exponentially with increasing N fertilization (Hoben et al., 2011; Snyder et al., 2009). Our data supported the results received by these authors. In our opinion, the N rates of 120 and 130 kg N ha^{-1} exceeded the threshold of N requirements of carrot during the growing season of 2007. Moreover, there was only a weak correlation (r=0.16) between the yields of carrot and the mineral N-fertilizer rates for the growing season of 2007.




Figure 12. Differences in cumulative N_2O fluxes between soil in ridges and on furrows (F) in relatively dry (2006) and wet (2007) growing seasons at different N-fertilizer rates (0, 60, 70, 130 kg N ha^{-1}) (Natalya P. Buchkina, Elena Y. Rizhiya, Sergey V. Pavlik, Eugene V. Balashov)

It was recommended by J.W. Van Groenigen et al. (2010) that to obtain more valid information on N_2O emissions from agricultural soils it is better to assess N_2O emissions as a function of crop yield. According to our results, the ratios of N_2O cumulative fluxes (average for furrows and ridges) to carrot yields were equal to $1.16*10^{-5}$, $1.17*10^{-5}$, $1.27*10^{-5}$, $1.20*10^{-5}$ and $1.61*10^{-5}$ kg N_2O-N ha^{-1} kg^{-1} yield at the N-fertilizer rates of 40, 60, 110, 120 and 130 kg N ha^{-1}, respectively. These results also demonstrated that the N rate of 130 kg ha^{-1} exceeded the N requirements of carrot during the growing season of 2007.

When all the data for all the growing seasons were taken into account we observed that the significant positive correlation between amount of N applied into the soil with mineral fertilizers and cumulative N_2O flux for a growing season was higher in the soils of furrows (r=0.66, p =0.01) than in the soil of ridges (r=0.56, p=0.01) (Figure 13). The soil of furrows contained the same amount of mineral N and carbon as the soil of ridges but often had higher water content, bulk density and WFPS and, as a result, demonstrated more favorable soil physical conditions for microbial process of denitrification.

The emission factor of 1% is recommended by (IPCC, 2006) for evaluating the efficiency of direct N_2O emissions from agricultural soils. The emission factor (calculated for 5 months' cumulative fluxes) for the different crops varied during the four growing seasons from 0.33 to 2.30%. The emission factors were expectedly higher in wetter growing seasons of 2004, 2005 and 2007 (0.50-2.30%) than in drier growing season of 2006 (0.50-0.62%) The emission

factors were often higher than 1% for the soil of both furrows and ridges only in wetter growing seasons of 2004, 2005 and 2007 but not in dry growing season of 2006.




Figure 13. Relationship between the rates of mineral nitrogen fertilizer and cumulative N_2O flux from a sandy loam Spodosol in furrows (black dots and line) and on ridges (empty dots and dotted line) for the growing seasons of 2004-2007 (Natalya P. Buchkina, Elena Y. Rizhiya, Sergey V. Pavlik, Eugene V. Balashov)

4. Conclusions

Cumulative N_2O fluxes from a sandy loam Spodosol for the growing seasons of 2004–2007 varied between 0.34±0.09 and 1.83±0.45 kg N_2O-N ha^{-1} for different crops studied in this experiment. Seasonal N_2O cumulative fluxes were higher from the soils of furrows than from those of ridges (especially on the plots where mineral nitrogen was applied with fertilizers) only in wetter growing seasons of 2004, 2005 and 2007. This difference was not significant for the drier growing season of 2006 or when all the four growing seasons were taken into consideration as climatic conditions affected the relationship and made it much weaker.

The N application contributed to a higher N_2O emission from soil in furrows, where the soil was more compacted with higher water-filled pore space, than from soil on ridges but only in wetter growing seasons. During the dry growing season of 2006 there was no significant difference between N_2O emissions from soils on ridges and in furrows. There was a nonlinear N_2O response to increasing N fertilizer rates from the soil on ridges and in furrows for carrot during the wet growing season of 2007. The mineral N-fertilizer rate of 130 kg N ha^{-1} could exceed the N requirements of carrot during the growing season of 2007.

Soil water-filled pore space affected N_2O emission from the soil only if mineral N-fertilizer was applied into the soil. Plots receiving no extra N never emitted much N_2O whatever the soil water-filled pore space.

Acknowledgements

The authors are very grateful to Prof. V.A. Semenov, the founder, and to Dr. E.A. Olenchenko, the main research manager, of the field experiment where all the described data were collected. Data on the crop yields in 2006 and 2007 were provided by Dr. E.A. Olenchenko (not published). The Royal Society of London supported the project (ref no 15336) in 2003-2005. The project would have been impossible without the donation of the gas chromatograph by the University of Edinburgh to the Agrophysical Research Institute. We thank the staff of the Menkovo Experimental Station of the Agrophysical Research Institute for providing the information on daily weather conditions.

Author details

Natalya P. Buchkina*, Elena Y. Rizhiya, Sergey V. Pavlik and Eugene V. Balashov

*Address all correspondence to: buchkina_natalya@mail.ru

Department of Soil Physics, Physical Chemistry and Biophysics, Agrophysical Research Institute, St. Petersburg, Russia

References

[1] Almaraz JJ, Mabood F, Zhou X, Madramootoo C, Rochette P, Ma B, Smith DL. Carbon dioxide and nitrous oxide fluxes in corn grown under two tillage systems in Southwestern Quebec. Soil Science Society of America Journal 2009;73 113–119.

[2] Angers DA, Bolinder MA, Carter MR, Gregorich EG, Drury CF, Liang BC, Voroney RP, Simard RR, Donald RG, Beyaert RP, Martel J. Impact of tillage practices on organic carbon and nitrogen storage in cool, humid soils of eastern Canada. Soil and Tillage Research 1997;41 191–201.

[3] Balashov E, Bazzoffi P. Aggregate water stability of differently compacted sandy and clayey loam soils with and without wheat plants. International Agrophysics 2003;17 151-155.

[4] Balashov E, Buchkina N. Effect of short- and long-term agricultural use of chernozem on its quality indicators. International Agrophysics 2011;25 1-5.

[5] Ball BC, McTaggart IP, Watson CA. Influence of organic ley-arable management and afforestation in sandy loam to clay loam soils on fluxes of N2O and CH4 in Scotland. Agriculture, Ecosystems and Environment 2002;90 305–317.

[6] Ball BC, Scott A, Parker JP. Field. N2O, CO2 and CH4 fluxes in relation to tillage, compaction and soil quality in Scotland. Soil and Tillage Research 1999;53 29–39.

[7] Ball BC, Watson CA, Crichton I. Nitrous oxide emissions, cereal growth, N recovery and soil nitrogen status after ploughing organically managed grass/clover swards. Soil Use and Management 2007;23 145–155

[8] Bankin MP, Bankina TA, Korobeinikova LP. Physicochimicheskiye metody v agrochimii i biologii pochv (Physico-chemical methods in agrochemistry and soil biology). Izdatelstvo Sankt-Peterburgskogo Universiteta, Sankt-Peterburg; 2005.

[9] Beare MH, Hendrix PF, Coleman DC. Water-stable aggregates and organic matter fractions in conventional- and no-tillage. Soil Science Society of America Journal 1994;58 777–786.

[10] Beare MH, Gregorich. EG, St-Georges P. Compaction effects on CO2 and N2O production during drying and rewetting of soil. Soil Biology and Biochemistry 2009; 41 611-621.

[11] Bird JA, Herman DJ, Firestone MK. Rhizosphere priming of soil organic matter by bacterial groups in a grassland soil. Soil Biology and Biochemistry 2011;43 718-725.

[12] Bock E, Koops HP, Harms H. Cell biology of nitrifying bacteria. In: Prosser JI. (ed) Nitrification. Special Publications of the Society for General Microbiology; 1986. 20 p17–38.

[13] Bouwman AF. Exchange of greenhouse gas between terrestrial ecosystems and atmosphere. In: Bouwman AF. (ed) Soil and the greenhouse effects. Wiley, Chichester, 1990. p.61–127.

[14] Bouwman AF. Direct emission of nitrous oxide from agricultural soils. Nutrient cycling in agroecosystems 1996;46 53-70.

[15] Bremner, JM. Sources of nitrous oxide in soils. Nutrient Cycling in Agroecosystems 1997;49 7-16.

[16] Buchkina NP, Balashov EV, Rizhiya EY, Smith KA. Nitrous oxide emissions from a light-textured arable soil of North-Western Russia: effects of crops, fertilizers, manures and climate parameters. Nutrient Cycling in Agroecosystems 2010;87 429-442.

[17] Burton DL, Beauchamp EG. Profile nitrous oxide and carbon dioxide concentrations in a soil subject to freezing. Soil Science Society of America Journal 1994;58 115–122.

[18] Chatskikh D, Olesen JE. Soil tillage enhanced CO2 and N2O emissions from loamy sand soil under spring barley. Soil and Tillage Research 2007;97 5–18.

[19] Choudhary MA, Akramkhanov A, Saggar S. Nitrous oxide emission from a New Zealand cropped soil: tillage effects, spatial and seasonal variability. Agriculture, Ecosystems and Environment 2002;93 33–43.

[20] Clough TJ, Sherlock RR, Rolston DE. A review of the movement and fate of N2O in the subsoil. Nutrient Cycling in Agroecosystems 2005;72 3–11.

[21] Conen F, Dobbie KE, Smith KA. Predicting N2O emission from agricultural land through related soil parameters. Global Change Biology 2000;6 417–426.

[22] Crutzen PJ. The role of NO and N2O in the chemistry of the troposphere and stratosphere. Annual Review of Earth and Planetary Sciences 1979;7 443-472.

[23] Davidson EA. Fluxes of nitrous oxide and nitric oxide from terrestrial ecosystems. In: Rogers JE, Whitman WB (eds) Microbial production and consumption of greenhouse gases: methane, nitrogen oxides and halo-methanes. Washington, DC: American Society of Microbiology; 1991; p219–235.

[24] Dobbie KE, Smith KA. Nitrous oxide emission factors for agricultural soils in Great Britain: The impact of soil water-filled pore space and other controlling variables. Global Change Biology 2003;9 204–218.

[25] Dobbie KE, Smith KA. Impact of different forms of N fertilizer on N2O emissions from intensive grassland. Nutrient Cycling in Agroecosystems 2003;67 37–46.

[26] Dobbie KE, McTaggart IP, Smith KA. Nitrous oxide emissions from intensive agricultural systems: variations between crops and seasons; key driving variables; and mean emission factors. Journal of Geophysical Research 1999;104 26891- 26899.

[27] Douglas JT, Crawford CE. Soil compaction effects on utilization of nitrogen from livestock slurry applied to grassland. Grass and Forage Science 1998;53 31-40.

[28] Drury CF, Zhang TQ, Kay BD. The non-limiting and least limiting water ranges for soil nitrogen mineralization. Soil Science Society of America Journal 2003;67 1388–1404.

[29] Elder JW, Lal R. Tillage effects on gaseous emissions from an intensively farmed organic soil in North Central Ohio. Soil and Tillage Research 2008; 98 45–55.

[30] Fernandez R, Quiroga A, Zorati C, Noellmeyer E. Carbon contents and respiration rates of aggregate size fractions under no-till and conventional tillage. Soil and Tillage Research 2010;109 103-109.

[31] Firestone MK, Davidson EA. Microbiological basis of NO and nitrous oxide production and consumption in soil, In Andreae MO, Schimel DS. (eds.) Exchange of Trace Gases between Terrestrial Ecosystems and the Atmosphere. New York: John Wiley and Sons; 1989. p7-21.

[32] Flessa H, Dorsch P. Seasonal variation on N2O and CH4 fluxes in differently managed arable soils in southern Germany. Journal of Geophysical Research 1995;100(D11):23115–23124.

[33] Granli T, Bøckman OC. Nitrous oxide from agriculture. Norwegian Journal of Agricultural Science 1994;12 128 pp.

[34] Grant RF, Pattey E, Goddard TW, Kryzanowski LM, Puurveen H. Modeling the effects of fertilizer application rate on nitrous oxide emissions. Soil Science Society of America Journal 2006;70 235–248.

[35] Green VS, Stott DE, Cruz JC, Curi N. Tillage impacts on soil biological activity and aggregation in a Brazilian Cerrado Oxisol. Soil and Tillage Research 2007;92 114–121.

[36] Gregorich EG, Kachanoski RG, Voroney RP. Carbon mineralization in soil size fractions after various amounts of aggregate disruption. Journal of Soil Science 1989;40 649:659.

[37] Gregorich EG, Rochette P, VandenBygaart AJ, Angers DA. Greenhouse gas contributions of agricultural soils and potential mitigations practices in Eastern Canada. Soil and Tillage Research 2005;83 53-72.

[38] Halvorson AD, Del Grosso SJ, Reule CA. Nitrogen, tillage, and crop rotation effects on nitrous oxide emissions from irrigated cropping systems. Journal of Environmental Quality 2008;37 1337–1344.

[39] Henault C, Devis X, Page S, Justes E, Reau R, Germon JC. Nitrous oxide emissions from different soil and land management conditions. Biology and Fertility of Soils 1998;26 199–207.

[40] Hergoualc'h K, Skiba U, Harmand J-M, Henault C. Fluxes of greenhouse gases from Andosols under coffee in monoculture or shaded by Inga densiflora in Costa Rica. Biogeochemistry 2008;89 329–345.

[41] Hoben JR, Gehl RJ, Millar N, Grace PR, Robertson GP. Nonlinear nitrous oxide (N2O) response to nitrogen fertilizer in on-farm corn crops of the US Midwest. Global Change Biology 2011;17 1140-1152.

[42] Hochstein LI, Tomlinson GA. The enzymes associated with denitrification. Annual Review of Microbiology 1988;42 231-261.

[43] Houghton JT, Jenkins GJ, Ephraums JJ., editors. Climate Change: The IPCC Scientific Assessment. Cambridge: Cambridge University Press; 1990.

[44] IPCC. Changes in atmospheric constituents and in radiative forcing. Cambridge: Cambridge University Press; 2007.

[45] Intergovernmental Panel on Climate Change, IPCC Guidelines for National Greenhouse Gas Inventories, prepared by the National Greenhouse Gas Inventories Programme. In: Eggleston, HS, Buendia L, Miwa K, Ngara T, Tanabe K. (eds.) N2O Emissions from Managed Soils, and CO2 Emissions from Lime and Urea Application, vol. 4. Hayama: IGES (Chapter 11); 2006.

[46] Jacinthe PA, Lal R. Effects of soil cover and land-use on the relations flux-concentration of trace gases. Soil Science 2004;169 243–259.

[47] Jones SK, Rees RM, Skiba UM, Ball BC. Influence of organic and mineral N fertiliser on N2O fluxes from a temperate grassland. Agriculture Ecosystems and Environment 2007;121 74–83.

[48] Kaiser EA, Ruser R. Nitrous oxide emissions from arable soils in Germany – an evaluation of six long-term field experiments. Journal of Plant Nutrition and Soil Science 2000;163 249–259.

[49] Kavdir Y, Hellebrand HJ, Kern J. Seasonal variations of nitrous oxide emission in relation to nitrogen fertilization and energy crop types in sandy soil. Soil and Tillage Research 2008;98 175–186.

[50] Khalid M, Soleman N, Jones DL. Grassland plants affect dissolved organic matter and nitrogen dynamics in soil. Soil Biology and Biochemistry 2007;39 378-381.

[51] Khalil K, Mary B, Renault P. Nitrous oxide production by nitrification and denitrification in soil aggregates as affected by O2 concentration. Soil Biology and Biochemistry 2004;36 687–699.

[52] Kusa K, Sawamoto T, Hu R, Hatano R. Comparison of N2O and CO2 concentrations and fluxes in the soil profile between a Gray Lowland soil and an Andosol. Soil Science and Plant Nutrition 2010;56 186–199.

[53] Lal R. Soil carbon sequestration to mitigate climate change. Geoderma 2004;123 1–22.

[54] Lemke RL, Izaurralde RC, Nyborg M, Solberg ED. Tillage and N source influence soil-emitted nitrous oxide in the Alberta Parkland region. Canadian Journal of Soil Science 1999;79:15–24.

[55] Li X, Inubushi K, Sakamoto K. Nitrous oxide concentrations in an andisol profile and emissions to the atmosphere as influenced by the application of nitrogen fertilizers and manure. Biology and Fertility of Soils 2002;35 108–113.

[56] Liu Hui, Zhao Ping, Lu Ping, Wang Yue-Si, Lin Yong-Biao, Rao Xing-Quan. Greenhouse gas fluxes from soils of different land-use types in a hilly area of South China. Agriculture, Ecosystems and Environment 2008;124 125–135.

[57] Liu XJ, Mosier AR, Halvorson AD, Reule CA, Zhang FS. Dinitrogen and N2O emissions in arable soils: Effect of tillage, N source and soil moisture Soil Biology and Biochemistry 2007;39 2362–2370.

[58] Lopez-Garrido R, Madejon E, Murillo JM, Moreno F. Soil quality alteration by mouldboard ploughing in a commercial farm devoted to no-tillage under Mediterranean conditions. Agriculture, Ecosystems and Environment 2011;140 182–190.

[59] MacKenzie AF, Fan MX, Cadrin F. Nitrous oxide emission in three years as affected by tillage, corn-soybean-alfalfa rotations, and nitrogen fertilization. Journal of Environmental Quality 1998;27 698–703.

[60] Maljanen M, Martikainen PJ, Aaltonen H, Silvola J. Short-term variation in fluxes of carbon dioxide, nitrous oxide and methane in cultivated and forested organic boreal soils. Soil Biology and Biochemistry 2002;34 577–584.

[61] McCarty GW. Modes of action of nitrification inhibitors. Biology and Fertility of Soils 1999;29 1-9.

[62] McSwiney CP, Robertson GP. Nonlinear response of N2O flux to incremental fertilizer addition in a continuous maize (Zea mays L.) cropping system. Global Change Biology 2005;11 1712–1719.

[63] Meng L, Ding W, Cai Z. Long-term application of organic manure and nitrogen fertilizer on N2O emissions, soil quality and crop production in a sandy loam soil. Soil Biology and Biochemistry 2005;37 2037–2045.

[64] Merino P, Artetxe A, Castellon A, Menendez S, Aizpurua A, Estavillo JM. Warming potential of N2O emissions from rapeseed crop in Northern Spain. Soil and Tillage Research 2012;123 29–34.

[65] Mitchell AR, Ellsworth TR, Meek BD. Effect of root systems on preferential flow in swelling soil. Communications in Soil Science and Plant Analysis 1995;26 2655-2666.

[66] Mosier A, Kroeze C, Nevison C, Oenema O, Seitsinger S, Van Cleemput O. Closing the global N2O budget: nitrous oxide emissions through the agricultural nitrogen cycle. Nutrient Cycling in Agroecosystems 1998;52 225–248.

[67] Muller C, Stevens R, Laughlin R, Jager HJ, Microbial processes and the site of N2O production in a temperate grassland soil. Soil Biology and Biochemistry 2004;36 453–461.

[68] Norton LD, Mamedov AI, Huang C. Soil aggregate stability as affected long-term tillage and clay mineralogy. In: Horn R, Fleige H, Peth S, Peng X. (eds.) Soil Management for Sustainability: Advances in Geoecology. Catena Verlag, Reiskirchen; 2006.

[69] Nyakatawa EZ, Reddy KC, Lemunyon JL. Predicting soil erosion in conservation tillage cotton production systems using the revised universal soil loss equation (RUSLE). Soil and Tillage Research 2000;57 213–224.

[70] OECD. 2000. Environmental indicators for agriculture, methods, and results, executive summary. Paris, France (www.oecd.org/dataoecd/0/9/19166629.pdf)

[71] Petersen SO, Schjonning P, Thomsen IK, Christensen BT. Nitrous oxide evolution from structurally intact soil as influenced by tillage and soil water content. Soil Biology and Biochemistry 2008;40 967–977.

[72] Phillips RL, Wick AF, Liebig MA, West MS, Lee DW. Biogenic emissions of CO2 and N2O at multiple depths increase exponentially during a simulated soil thaw for a northern prairie Mollisol. Soil Biology and Biochemistry 2012;45 14-22.

[73] Poth M, Focht DD. 15N kinetic analysis of N2O production by Nitrosomonas europaea: an examination of nitrifier denitrification. Applied and Environmental Microbiology 1985;49 1134–1141.

[74] Rasmussen KJ, Impact of ploughless soil tillage on yield and soil quality: a Scandinavian review. Soil and Tillage Research 1999;53 3–14.

[75] Rastvorova OG. Soil physics: practical use. Leningrad: Leningrad University Press (in Russian). 1988.

[76] Reicosky DC. Long-term effect of mouldboard ploughing on tillage-induced CO2 loss. In: Kimble JM, Lal R, Follet RF. (eds.) Agricultural Practices and Policies for Carbon Sequestration in Soil. Florida: CRC/Lewis, Boca Raton; 2002. p.87–97.

[77] Riley H, Pommeresche R, Eltun R, Hansen S, Korsaeth A. Soil structure, organic matter and earthworm activity in a comparison of cropping systems with contrasting tillage, rotations, fertilizer levels and manure use. Agriculture, Ecosystems and Environment 2008;124 275–284.

[78] Ritchie GAF., Nicholas DJD. Identification of the sources of nitrous oxide produced by oxidative and reductive processes in Nitrosomonas europaea. Biochemistry Journal 1972;126 1181–1191.

[79] Rizhiya EY, Boitsova LV, Buchkina NP, Panova GG. The influence of plant recidues with different C/N ratio on nitrous oxide emission from derno-podzolic loamy-sand soil. Eurasian Journal of Soil Science 2011;10 1251-1259.

[80] Robertson GP, Groffman PM. Nitrogen Transformation. In: Paul EA. (ed.) Soil Microbiology, Biochemistry, and Ecology. New York: Springer; 2007. p341-364.

[81] Ruser R, Flessa H, Schilling R, Steindl H, Beese F. Soil compaction and fertilization effects on nitroius oxide and methane fluxes in potato fields. Soil Science Society of America Journal 1998;62 1587-1595.

[82] Simansky V, Tobiasova E, Chlpik J. Soil tillage and fertilization of Orthic Luvisol and their influence on chemical properties, soil structure stability and carbon distribution in water-stable macro-aggregates. Soil and Tillage Research 2008;100 125–132.

[83] Six J, Elliott ET, Paustian K. Aggregate and SOM dynamics under conventional and no-tillage systems. Soil Science Society of America Journal 1998;63 1350–1358.

[84] Six J, Feller Ch, Denef C, Ogle SM, de Morales Sa JC, Albrecht A. Soil organic matter, biota and aggregation in temperate and tropical soils – effects of no-tillage. Agronomie 2002;22 755–775.

[85] Skiba U, Smith KA. The control of nitrous oxide emissions from agricultural and natural soils. Chemosphere 2000;2 379–386.

[86] Smith KA. A model of the extent of anaerobic zones in aggregated soils, and its potential application to estimates of denitrification. Journal of Soil Science 1980;31 263–277.

[87] Smith KA, Ball T, Conen F, Dobbie KE, Massheder J, Rey A. Exchange of greenhouse gases between soil and atmosphere: interactions of soil physical factors and biological processes. European Journal of Soil Science 2003;54 779–791.

[88] Snyder CS, Bruulsema TW, Jensen TL, Fixen PE. Review of greenhouse gas emissions from crop production systems and fertilizer management effects. Agriculture, Ecosystems and Environment 2009;133 247-266.

[89] Soil Survey. Laboratory methods manual, soil survey investigations report N42, version 3.0, January 1996. Washington: US Department of Agriculture, Natural Resources Conservation Service, National Soil Survey Center; 1996.

[90] Uchida Y, Clough TJ, Kelliher FM, Sherlock RR. Effects of aggregate size, soil compaction, and bovine urine on N2O emissions from a pasture soil. Soil Biology and Biochemistry 2008;40 924–931.

[91] Van Groenigen JW, Kuikman PJ, de Groot WJM., Velthof GL. Nitrous oxide emission from urine-treated soil as influenced by urine composition and soil physical conditions. Soil Biology and Biochemistry 2005;37 268–274.

[92] Van Groenigen JW, Velthof GL, Oenema O, van Groenigen KJ, van Kessel C. Towards an agronomic assessment of N2O emissions: a case study for arable crops. European Journal of Soil Science 2010;61 903–913.

[93] Velthof GL, Brader AB, Oenema O. Seasonal variations in nitrous oxide losses from managed grasslands in the Netherlands. Plant and Soil 1996;181 263–274.

[94] West TO, Post WM. Soil organic carbon sequestration rates by tillage and crop rotation: a global data analysis. Soil Science Society of America Journal 2002;66 1930–1946.

[95] Williams SM, Weil RR. Crop cover root channels may alleviate soil compaction effects on soybean crop. Soil Science Society of America Journal 2004;68 1403-1409.

[96] Wlodarczyk T, Stepniewska Z, Brzezinska M. Denitrification, organic matter, and redox potential transformations in Cambisols. International Agrophysics 2003;17 219-227.

[97] Wood PM. Notrification as a bacterial energy source. In: Prosser J.I. (ed.) Nitrification. Special Publications of the Society for General Microbiology 1986;20 39-62.

[98] Yoh M, Toda H, Kanda K, Tsuruta H. Diffusion analysis of N2O cycling in a fertilized soil. Nutrient Cycling in Agroecosystems 1997;49 29–33.

[99] Zhang GS, Chan KY, Oates A, Heenan DP, Huang GB. Relationship between soil structure and runoff/soil loss after 24 years of conservation tillage. Soil and Tillage Research 2007;92 122–128