

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

185,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Hermansky-Pudlak Syndrome

Naoki Oiso and Akira Kawada

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/53573>

1. Introduction

Oculocutaneous albinism is classified into non-syndromic oculocutaneous albinism (OCA) and syndromic OCA including Hermansky-Pudlak syndrome (HPS), Chediak-Higashi syndrome (CHS) and Griscelli syndrome (GS). Both non-syndromic and syndromic OCAs are autosomal recessive disorders. Human HPS is genetically divided into nine forms, HPS type 1 (HPS-1) to HPS-9. Human HPS can be sub-classified into four subgroups which are associated with protein complexes encoded by the causative genes. In this session, we summarize (1) the clinical features of HPS, (2) the mice and rat models of HPS, and (3) the molecular functions.

2. The clinical features of HPS

In 1959, Hermansky and Pudlak described two cases of OCA associated with hemorrhagic diathesis.¹ Currently, the condition is known as HPS. HPS is a rare heterogeneous autosomal recessive syndrome which is typically characterized by OCA, bleeding diathesis, and lysosomal ceroid storage resulting from defects of multiple cytoplasmic organelles: melanosomes, platelet dense core granules, and lysosomes.² The storage of ceroid-like material in lysosomes induces restrictive lung disease, ulcerative colitis, kidney failure, and cardiomyopathy.

Accumulation of mice models, identification of causative genes and functional analysis indicated that HPS could be sub-classified into four groups according to four protein complexes, biogenesis of lysosome-related organelles complex-3 (BLOC-3) (HPS-1 and HPS-4), adaptor protein-3 (AP-3) (HPS-2), BLOC-2 (HPS-3, HPS-5 and HPS-6) and BLOC-1 (HPS-7, HPS-8 and HPS-9).³⁻⁵ Currently, more than 16 mice strains and more than 2 rat strains are known as models of human HPS (**Table 1**). HPS-1 is caused by mutation in *HPS1*,⁶ HPS-2 is caused by mutation in *AP3B1*,⁷ HPS-3 is caused by mutation in *HPS3*,⁸ HPS-4 is caused by mutation in *HPS4*,⁹ HPS-5 is caused by mutation in *HPS5*,¹⁰ HPS-6 is caused by mutation in

HPS6,¹⁰ HPS-7 is caused by mutation in *DTNBP1*,¹¹ HPS-8 is caused by mutation in *BLOC1S3*,¹² and HPS-9 is caused by mutation in *PLDN*.¹³ Functional analyses identify that most of all HPS proteins construct complexes, BLOC-1, BLOC-2, BLOC-3, AP3, class C vacuolar protein sorting (VPS), and Rab geranylgeranyl transferase (RABGGT).

Mouse models	Human type	Genes	Protein complexes
<i>pale ear</i>	HPS-1	<i>HPS1</i>	BLOC-3
<i>pearl</i>	HPS-2	<i>AP3B1</i>	AP3
<i>cocoa</i>	HPS-3	<i>HPS3</i>	BLOC-2
<i>light ear</i>	HPS-4	<i>HPS4</i>	BLOC-3
<i>ruby-eye-2</i>	HPS-5	<i>HPS5</i>	BLOC-2
<i>ruby-eye</i>	HPS-6	<i>HPS6</i>	BLOC-2
<i>sandy</i>	HPS-7	<i>DTNBP1</i>	BLOC-1
<i>reduced pigmentation</i>	HPS-8	<i>BLOC1S3</i>	BLOC-1
<i>pallid</i>	HPS-9	<i>PLDN</i>	BLOC-1
<i>buff</i>	?	<i>VPS33A</i>	class C VPS
<i>cappuccino</i>	?	<i>CNO</i>	BLOC-1
<i>gunmetal</i>	?	<i>RABGGTA</i>	RABGGT
<i>misty</i>	?	<i>DOCK7</i>	
<i>mocha</i>	?	<i>AP3D1</i>	AP3
<i>muted</i>	?	<i>MUTED</i>	BLOC-1
<i>subtle gray</i>	?	<i>SLC7A11</i>	
Rat models			
Fawn-Hooded rat	?	<i>RAB38</i>	
Tester-Moriyama rat	?	<i>RAB38</i>	

Table 1. Animal models, human types, causative genes and their protein complexes in HPS.

HPS-1 and HPS-4, the group of BLOC-3, are the most dominant and typical subtypes. The founder effect in HPS-1 is present in the region of northwest Puerto Rico.⁸ HPS-1 and HPS-4 are characterized by OCA by deficiency of melanosomes, bleeding by loss of platelet dense core granules, and systemic organ involvement (restrictive lung disease, granulomatous colitis, kidney failure, and cardiomyopathy) by the storage of lysosomal ceroid-like substances due to impaired lysosomes. The clinical features of OCA and bleeding diathesis are present in infancy. Bleeding tendency is important to diagnose.

HPS-2, the group of AP3, is the most severe and rare subtype, with 15 cases reported in the literature.¹⁴ Clinical manifestations include OCA, a platelet storage pool defect, interstitial lung disease, and recurrent bacterial and viral infections due to immunodeficiency.¹⁴ Patients with HPS-2 exhibit neutropenia that is responsive to granulocyte colony-stimulation factor, deficiency of natural killer and natural killer T-cells, T-lymphocyte dysfunction, and in one case hemophagocytic lymphohistiocytosis.¹⁴

HPS-3, HPS-5, and HPS-6, the group of BLOC-2, are the milder and relatively rare subtypes. The founder effect in HPS-3 is present in the area of central Puerto Rico.⁸ HPS-3, HPS-5 and HPS-6 are relatively milder forms of the disease in that both OCA and bleeding diathesis are mild and pulmonary fibrosis and granulomatous colitis generally does not develop.^{10, 15-17}

HPS-7, HPS-8, and HPS-9, the group of BLOC-1, are extremely rare subtypes. HPS-7 is only found in a 48-year-old Portuguese woman with OCA, a bleeding tendency, mild shortness of breath on exertion and reduced lung compliance but otherwise normal pulmonary function.¹¹ HPS-8 is found in a Pakistani family¹² and an Iranian patient.¹⁸ HPS-8 is characterized by typical OCA and a bleeding diathesis. Pulmonary fibrosis, granulomatous colitis, or neutropenia are not detected in the cases.^{12, 18} HPS-9 is only found in a 9-month-old male of Indian ancestry.¹³ The patient showed OCA with generalized hypopigmentation, nystagmus, iris transillumination, and retinal hypopigmentation; respiratory distress requiring a 3 week admission to a neonatal intensive-care unit for respiratory support; and platelet electron microscopy showing absent platelet delta granules.¹³

Recombinant factor VIIa (rFVIIa) is useful for dangerous bleeding such as refractory menorrhagia.¹⁹ Progressive HPS-1 pulmonary fibrosis is effectively treated by pirfenidone, a small molecule that inhibits TGF-beta-mediated fibroblast proliferation and collagen synthesis *in vitro*.²⁰ Infliximab is effective for granulomatous colitis in HPS patients.²¹ The efficacy of infliximab suggests that TNF- α plays a pivotal role in the pathogenesis.²¹

3. The mice and rat models of HPS

Mice models of HPS can be grouped into BLOC-3 (*pale ear*^{22, 23} and *light ear*⁹), BLOC-2 (*cocoa*²⁴, *ruby-eye-2*¹⁰ and *ruby-eye*¹⁰), BLOC-1 (*sandy*¹¹, *reduced pigmentation*²⁵, *pallid*²⁶, *cappuccino*²⁷ and *muted*²⁸), AP-3 (*pearl*²⁹ and *mocha*³⁰), class C VPS (*buff*³¹), RABGGT (*gunmetal*³²), and others (*misty*³³ and *subtle gray*³⁴) (**Table 1**). Two rat models of HPS, Fawn-Hooded Rat and Tester-Moriyama Rat, are genetically identical with no expression of *RAB38*.³⁵

Gautam *et al.* contacted mutant mice doubly or triply deficient in protein subunits of the various BLOC complexes and/or the AP-3 adaptor complex and tested for viability and for abnormalities of lysosome-related organelles (LROs) including melanosomes, lamellar bodies of lung type II cells and platelet dense granules.³⁶ They showed that double and triple mutant HPS mice provide unique and practical experimental advantages in the study of LROs.³⁶ Long-Evans Cinnamon rats with a point mutation in the initiation codon of Rab38 small GTPase are investigated for the pathogenesis of interstitial pneumonia via aberrant lung surfactant secretion.³⁷ Thus, mice and rat models are indispensable for recognizing the molecular function in LROs and the pathogenesis of HPS.

4. The molecular functions

The complexes involving in the pathogenesis of HPS are BLOC-3, BLOC-2, BLOC-1, AP-3 adaptor complex, and class C VPS. BLOC-3 is composed of HPS1/pale ear and HPS4/light

ear.³⁸⁻⁴⁰ BLOC-2 comprises HPS3/cocoa, HPS5/ruby-eye-2 and HPS6/ruby-eye.^{10, 41} BLOC-1 is constructed by proteins of HPS7/DTNBP1/sandy, HPS8/BLOC1S3/reduced pigmentation/BLOS3, cappuccino, muted, pallid, BLOS1, BLOS2 and snapin.^{42, 43} AP-3 (subunits δ /mocha, β 3/HPS2/AP3B1/pearl, μ 3, σ 3) is one of the family of heterotetrameric clathrin adaptors.⁴⁴ The class C VPS is composed of VPS11, VPS16, VPS18, and VPS33/buff.⁴⁵




Figure 1. BLOC-1, -2, -3, AP-3 and class C VPS complexes involve in membrane trafficking from endosomes to lysosomes and lysosome-related organelles, melanosomes and dense core granules in platelets.

Adaptor protein complexes are composed of heterotetramers (two large subunits, a medium-sized subunit and a small subunit) and sort cargo into vesicles for transport from one membrane compartment of the cell to another.⁴⁶ AP-3 traffics cargo from tubular endosomes to late endosomes, lysosomes, and related organelles via the bound to BLOC-1, vimentin, clathrin and others.^{46, 47}

The class C VPS core complex (VPS33A/B, VPS11, VPS16 and VPS18) is essential for late endosome and lysosome assembly and for numerous endolysosomal trafficking pathways.⁴⁸ Two class C VPC complexes, homotypic fusion and protein sorting (HOPS) and class C core vacuole/endosome tethering (CORVET), incorporate diverse biochemical functions: they tether membranes, stimulate Rab nucleotide exchange, guide SNARE assembly to drive membrane fusion, and possibly act as ubiquitin ligases.⁴⁸

BLOC-1 functions in selective cargo exit from early endosomes toward lysosomes and lysosome-related organelles such as melanosomes and BLOC-2 act sequentially in the same pathway.⁴⁹ Melanosome maturation requires at least two cargo transport pathways directly from early endosomes to melanosomes, one pathway mediated by AP-3 and one pathway mediated by BLOC-1 and BLOC-2.⁴⁹ BLOC-3 is constructed by HPS1 and HPS4 heterodimers.⁵⁰ BLOC-3 interacts with the GTP-bound form of the endosomal GTPase, Rab9. BLOC-3 might function as a Rab9 effector in the biogenesis of lysosome-related organelles.⁵⁰

5. Conclusion

Now, HPS is a representative disorder of aberrant membrane trafficking. HPS genes have been identified with mice models. The function of encoded proteins has been accompanied with cell biology in yeast, worm, fly and animal models. Membrane trafficking is crucial for cells to survive and play their active functions. Further emerging investigation will reveal more precise pathogenesis in HPS.

Aberrations

biogenesis of lysosome related organelle complex (BLOC)-1, -2, -3

adaptor protein complex 3 (AP3)

vacuolar protein sorting (VPS)

Rab geranylgeranyl transferase (RABGGT)

homotypic fusion and protein sorting (HOPS)

class C core vacuole/endosome tethering (CORVET)

Author details

Naoki Oiso and Akira Kawada

Departments of Dermatology, Kinki University Faculty of Medicine, Osaka-Sayama, Osaka, Japan

6. References

- [1] Hermansky F, Pudlak P. Albinism associated with hemorrhagic diathesis and unusual pigmented reticular cells in the bone marrow: report of two cases with histochemical studies. *Blood* 1959; 14: 162-9.
- [2] Oh J, Ho L, Ala-Mello S, *et al.* Mutation analysis of patients with Hermansky-Pudlak syndrome: a frameshift hot spot in the HPS gene and apparent locus heterogeneity. *Am J Hum Genet* 1998; 62: 593-598.
- [3] Li W, Rusiniak ME, Chintala S, *et al.* Murine Hermansky-Pudlak syndrome genes: regulators of lysosome-related organelles. *Bioessays* 2004; 26: 616-28.

- [4] Di Pietro SM, Dell'Angelica EC. The cell biology of Hermansky-Pudlak syndrome: recent advances. *Traffic* 2005; 6: 525-33.
- [5] Huizing M, Helip-Wooley A, Westbroek W, *et al.* Disorders of lysosome-related organelle biogenesis: clinical and molecular genetics. *Annu Rev Genomics Hum Genet* 2008; 9: 359-86.
- [6] Oh J*, Bailin T*, Fukai K*, Feng GH*, *et al* (*co-first author). Positional cloning of a gene for Hermansky-Pudlak syndrome, a disorder of cytoplasmic organelles. *Nat Genet* 1996; 14: 300-6.
- [7] Dell'Angelica EC, Shotelersuk V, Aguilar RC, *et al.* Altered trafficking of lysosomal proteins in Hermansky-Pudlak syndrome due to mutations in the beta 3A subunit of the AP-3 adaptor. *Mol Cell* 1999; 3: 11-21.
- [8] Anikster Y, Huizing M, White J, *et al.* Mutation of a new gene causes a unique form of Hermansky-Pudlak syndrome in a genetic isolate of central Puerto Rico. *Nat Genet* 2001; 28: 376-80.
- [9] Suzuki T, Li W, Zhang Q, *et al.* Hermansky-Pudlak syndrome is caused by mutations in HPS4, the human homolog of the mouse light-ear gene. *Nat Genet* 2002; 30: 321-4.
- [10] Zhang Q*, Zhao B*, Li W*, Oiso N*, *et al* (*co-first author). Ru2 and Ru encode mouse orthologs of the genes mutated in human Hermansky-Pudlak syndrome types 5 and 6. *Nat Genet* 2003; 33: 145-53.
- [11] Li W*, Zhang Q*, Oiso N*, *et al* (*co-first author). Hermansky-Pudlak syndrome type 7 (HPS-7) results from mutant dysbindin, a member of the biogenesis of lysosome-related organelles complex 1 (BLOC-1). *Nat Genet* 2003; 35: 84-9.
- [12] Morgan NV, Pasha S, Johnson CA, *et al.* A germline mutation in BLOC1S3/reduced pigmentation causes a novel variant of Hermansky-Pudlak syndrome (HPS8). *Am J Hum Genet* 2006; 78: 160-6.
- [13] Cullinane AR, Curry JA, Carmona-Rivera C, *et al.* A BLOC-1 mutation screen reveals that PLDN is mutated in Hermansky-Pudlak Syndrome type 9. *Am J Hum Genet* 2011; 88: 778-87.
- [14] Gochuico BR, Huizing M, Golas GA, *et al.* Interstitial lung disease and pulmonary fibrosis in Hermansky-Pudlak syndrome type 2, an adaptor protein-3 complex disease. *Mol Med* 2012; 18: 56-64.
- [15] Huizing M, Anikster Y, Fitzpatrick DL, *et al.* Hermansky-Pudlak syndrome type 3 in Ashkenazi Jews and other non-Puerto Rican patients with hypopigmentation and platelet storage-pool deficiency. *Am J Hum Genet* 2001; 69: 1022-32.
- [16] Huizing M, Hess R, Dorward H, *et al.* Cellular, molecular and clinical characterization of patients with Hermansky-Pudlak syndrome type 5. *Traffic* 2004; 5: 711-22.
- [17] Huizing M, Pederson B, Hess RA, *et al.* Clinical and cellular characterisation of Hermansky-Pudlak syndrome type 6. *J Med Genet* 2009; 46: 803-10.
- [18] Cullinane AR, Curry JA, Golas G, *et al.* A BLOC-1 mutation screen reveals a novel BLOC1S3 mutation in Hermansky-Pudlak Syndrome type 8. *Pigment Cell Melanoma Res* 2012; 25: 584-91.
- [19] Lohse J, Gehrish S, Tauer JT, *et al.* Therapy refractory menorrhagia as first manifestation of Hermansky-Pudlak syndrome. *Hamostaseologie* 2011; 31 Suppl 1: S61-3.

- [20] O'Brien K, Troendle J, Gochuico BR, *et al.* Pirfenidone for the treatment of Hermansky-Pudlak syndrome pulmonary fibrosis. *Mol Genet Metab* 2011; 103: 128-34.
- [21] Grucela AL, Patel P, Goldstein E, *et al.* Granulomatous enterocolitis associated with Hermansky-Pudlak syndrome. *Am J Gastroenterol* 2006; 101: 2090-5.
- [22] Gardner JM, Wildenberg SC, Keiper NM, *et al.* The mouse pale ear (ep) mutation is the homologue of human Hermansky-Pudlak syndrome. *Proc Natl Acad Sci U S A* 1997; 94: 9238-43.
- [23] Feng GH, Bailin T, Oh J, *et al.* Mouse pale ear (ep) is homologous to human Hermansky-Pudlak syndrome and contains a rare 'AT-AC' intron. *Hum Mol Genet* 1997; 6: 793-7.
- [24] Suzuki T*, Li W*, Zhang Q, *et al* (*co-first author). The gene mutated in cocoa mice, carrying a defect of organelle biogenesis, is a homologue of the human Hermansky-Pudlak syndrome-3 gene. *Genomics* 2001; 78: 30-7.
- [25] Gwynn B, Martina JA, Bonifacino JS, *et al.* Reduced pigmentation (rp), a mouse model of Hermansky-Pudlak syndrome, encodes a novel component of the BLOC-1 complex. *Blood* 2004; 104: 3181-9.
- [26] Huang L, Kuo YM, Gitschier J, *et al.* The pallid gene encodes a novel, syntaxin 13-interacting protein involved in platelet storage pool deficiency. *Nat Genet* 1999; 23: 329-32.
- [27] Cicciotte SC, Gwynn B, Moriyaka K, *et al.* Cappuccino, a mouse model of Hermansky-Pudlak syndrome, encodes a novel protein that is part of the pallidin-mutated complex (BLOC-1). *Blood* 2003; 101: 4402-7.
- [28] Zhang Q*, Li W*, Novak EK, *et al* (*co-first author). The gene for the muted (mu) mouse, a model for Hermansky-Pudlak syndrome, defines a novel protein which regulates vesicle trafficking. *Hum Mol Genet* 2002; 11: 697-706.
- [29] Feng L*, Seymour AB*, Jiang S, *et al* (*co-first author). The beta3A subunit gene (Ap3b1) of the AP-3 adaptor complex is altered in the mouse hypopigmentation mutant pearl, a model for Hermansky-Pudlak syndrome and night blindness. *Hum Mol Genet* 1999; 8: 323-30.
- [30] Kantheti P, Qiao X, Diaz ME, *et al.* Mutation in AP-3 delta in the mocha mouse links endosomal transport to storage deficiency in platelets, melanosomes, and synaptic vesicles. *Neuron* 1998; 21: 111-22.
- [31] Suzuki T*, Oiso N*, Gautam R*, *et al* (co-first author). The mouse organellar biogenesis mutant buff results from a mutation in Vps33a, a homologue of yeast vps33 and *Drosophila* carnation. *Proc Natl Acad Sci U S A* 2003; 100: 1146-50.
- [32] Detter JC, Zhang Q, Mules EH, *et al.* Rab geranylgeranyl transferase alpha mutation in the gunmetal mouse reduces Rab prenylation and platelet synthesis. *Proc Natl Acad Sci U S A* 2000; 97: 4144-9.
- [33] Blasius AL, Brandl K, Crozat K, *et al.* Mice with mutations of Dock7 have generalized hypopigmentation and white-spotting but show normal neurological function. *Proc Natl Acad Sci U S A* 2009; 106: 2706-11.
- [34] Chintala S, Li W, Lamoreux ML, *et al.* Slc7a11 gene controls production of pheomelanin pigment and proliferation of cultured cells. *Proc Natl Acad Sci USA* 2005; 102: 10964-9.

- [35] Oiso N, Riddle SR, Serikawa T, *et al.* The rat Ruby (R) locus is Rab38: identical mutations in Fawn-hooded and Tester-Moriyama rats derived from an ancestral Long Evans rat sub-strain. *Mamm Genome* 2004; 15: 307-14.
- [36] Gautam R, Novak EK, Tan J, *et al.* Interaction of Hermansky-Pudlak Syndrome genes in the regulation of lysosome-related organelles. *Traffic* 2006; 7: 779-92.
- [37] Osanai K, Higuchi J, Oikawa R, *et al.* Altered lung surfactant system in a Rab38-deficient rat model of Hermansky-Pudlak syndrome. *Am J Physiol Lung Cell Mol Physiol* 2010; 298: L243-51.
- [38] Chiang PW, Oiso N, Gautam R, *et al.* The Hermansky-Pudlak syndrome 1 (HPS1) and HPS4 proteins are components of two complexes, BLOC-3 and BLOC-4, involved in the biogenesis of lysosome-related organelles. *J Biol Chem* 2003; 278: 20332-7.
- [39] Martina JA, Moriyama K, Bonifacino JS. BLOC-3, a protein complex containing the Hermansky-Pudlak syndrome gene products HPS1 and HPS4. *J Biol Chem* 2003; 278: 29376-84.
- [40] Nazarian R, Falcon-Perez JM, Dell'angelica EC. Biogenesis of lysosome-related organelles complex 3 (BLOC-3): a complex containing the Hermansky-Pudlak syndrome (HPS) proteins HPS1 and HPS4. *Proc Natl Acad Sci U S A* 2003; 100: 8770-5.
- [41] Gautam R, Chintala S, Li W, *et al.* The Hermansky-Pudlak syndrome 3 (cocoa) protein is a component of the biogenesis of lysosome-related organelles complex-2 (BLOC-2). *J Biol Chem* 2004; 279: 12935-42.
- [42] Falcon-Perez JM, Starcevic M, Gautam R, *et al.* BLOC-1, a novel complex containing the pallidin and muted proteins involved in the biogenesis of melanosomes and platelet-dense granules. *J Biol Chem* 2002; 277: 28191-9.
- [43] Starcevic M, Dell'angelica EC. Identification of snapin and three novel proteins (BLOS1, BLOS2, and BLOS3/reduced pigmentation) as subunits of biogenesis of lysosome-related organelles complex-1 (BLOC-1). *J Biol Chem* 2004; 279: 28393-401.
- [44] Simpson F, Bright NA, West MA, *et al.* A novel adaptor-related protein complex. *J Cell Biol* 1996; 133: 749-60.
- [45] Sato TK, Rehling P, Peterson MR, *et al.* Class C Vps protein complex regulates vacuolar SNARE pairing and is required for vesicle docking/fusion. *Mol Cell* 2000; 6: 661-71.
- [46] Hirst J, Barlow LD, Francisco GC, *et al.* The fifth adaptor protein complex. *PLoS Biol* 2011; 9: e1001170.
- [47] Newell-Litwa K, Seong E, Burmeister M, *et al.* Neuronal and non-neuronal functions of the AP-3 sorting machinery. *J Cell Sci* 2007; 120: 531-41.
- [48] Nickerson DP, Brett CL, Merz AJ. Vps-C complexes: gatekeepers of endolysosomal traffic. *Curr Opin Cell Biol* 2009; 21: 543-51.
- [49] Setty SR, Tenza D, Truschel ST, *et al.* BLOC-1 is required for cargo-specific sorting from vacuolar early endosomes toward lysosome-related organelles. *Mol Biol Cell* 2007; 18: 768-80.
- [50] Kloer DP, Rojas R, Ivan V, *et al.* Assembly of the biogenesis of lysosome-related organelles complex-3 (BLOC-3) and its interaction with Rab9. *J Biol Chem* 2010; 285: 7794-804.