

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Tuberculosis Pharmacogenetics: State of The Art

Raquel Lima de Figueiredo Teixeira,
Márcia Quinhones Pires Lopes,
Philip Noel Suffys and Adalberto Rezende Santos

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/54984>

1. Introduction

The interindividual variability in the metabolism of xenobiotics and drug response is extensive and many factors are involved with this variation including genetic composition, gender, age, co-administration of medication, individual physiology, pathophysiology and presence of other environmental factors (alcohol consumption, smoking, eating habits).

To produce their therapeutic effects, the drug must be present in appropriate concentrations at its site of action. Although the therapeutic concentrations are dependent on the given dose, they will also depend on the magnitude and rate of absorption, distribution, biotransformation, and excretion. Pharmacokinetics studies the course and distribution of drug and its metabolites in different tissues, covering the mechanisms of absorption, transport, metabolism and excretion. In addition, pharmacodynamics concentrates on the biochemical and physiological effects of drugs and their mechanism of action. Proteins involved in drug effects are defined as target molecules and include not only (direct) receptors, but also proteins associated with mechanism of action such as e.g. signal transducer proteins [1].

After its administration, a drug is absorbed and then distributed throughout the body, requiring the coordinated functioning of various proteins, including metabolic enzymes, trafficking proteins, receptor proteins, and others. Medication can enter the body as either active drugs or as inactive prodrugs. Most drugs are metabolized in the liver to make them more soluble for subsequent elimination through the kidneys or intestines. Prodrugs require metabolic conversion, also called biotransformation, to liberate the active compound. Complete biotransformation of any one drug typically requires several different enzymes. [2]. Genetic variability has been described to have effect on drug absorption and metabolism and its interactions with the receptors. This forms the basis for

slow and rapid drug absorption, poor, efficient or ultrarapid drug metabolism and poor or efficient receptor interactions [3]. The consequences of such variations can lead to adverse drug reaction and/or therapeutic failure.

In this context, pharmacogenetics is the study of genetic variations associated with individual variability in drug response, including differences in efficacy, drug-drug interactions, and the relative risk of an adverse response to drugs. It includes the study of genetic polymorphisms that could affect the expression or activity of drug transporters, drug metabolizing enzymes and drug receptors [2-4].

It's estimated that 99.9% of the human genome sequence between individuals is identical and genetic differences in populations are called mutations if they are present in less than 1% and polymorphisms when present in at least 1% of a population. A single-nucleotide polymorphism (SNP) involves a replacement of one nucleotide base with any one of the other three and occurring at approximately one out of every 1,000 bases in the human genome [5].

A mutation or polymorphism in genes that encode metabolic enzymes, carriers or receptors can affect the drug pharmacokinetics and pharmacodynamics leading to undesired therapeutic effects. The identification of these genetic markers which predicted if a person responds well or not to a specific drug could help to select the right medication in right dosage, maximizing the efficacy and preventing or reducing the adverse drug reactions.

2. Problem statement

TB is an important global public health problem but has cure in almost 100% of the new cases if correct chemotherapy is applied. The American Thoracic Society (ATS) treatment guidelines recommend an initial phase for TB treatment which consists of rifampicin 10 mg/kg (maximum 600 mg), isoniazid 5 mg/kg (maximum 300 mg), pyrazinamide 15–30 mg/kg (maximum 2 g), and ethambutol 15–20 mg/kg (maximum 1.6 g) given daily for 8 weeks, followed by a continuous phase of isoniazid 15 mg/kg (maximum 900 mg) and rifampicin 10 mg/kg (maximum 600 mg) administered 2–3 times/week for 18 weeks [6]. The use of fixed-dose combination (FDC) tablets containing anti-TB drugs has been recommended by the World Health Organization (WHO) as an additional measure to improve treatment adherence by reducing the number of tablets to be taken. The principal disadvantages of combining three or more drugs in one tablet include (a) the possibility of overdosage or underdosage resulting from a prescription error, (b) changes in the bioavailability of rifampicin and (c) difficulties in determining which drug is responsible for adverse effects [7].

Isoniazid (INH) is an important drug in the TB treatment and was introduced in chemotherapeutic scheme since 1952. It is the hydrazine of isonicotinic acid and shows cytotoxic activity for *Mycobacterium tuberculosis* both in rest (during latency) and proliferation phases. This drug enters easily in macrophage cells to kill bacilli in multiplication and is specific for mycobacteria [1].

INH-induced adverse reactions include fever, nausea, vomiting, hepatotoxicity, skin reactions, gastrointestinal and neurological disorders. Only in the early 1970s, the occurrence of severe liver injury as a side effect of this drug was recognized, resulting in the death of some patients [8]. Among the first-line anti-TB drugs, INH is the main associated with drug-induced hepatotoxicity with a frequency ranging from 1 to 30% in different populations [9]. Other drugs causing liver injury are mainly reported in combination with INH [10, 11]. Drug-induced hepatotoxicity is defined as a serum alanine aminotransferase (ALT) level three times greater than the upper limit of normal (ULN) with clinical symptoms or five times the ULN without symptoms. In both cases treatment should be interrupted and, generally, a modified or alternative regimen is introduced [9]. Because these adverse reactions do not only affect morbidity and mortality rate but also lead to treatment interruptions, failure and relapse, adverse reactions contribute to the spread of the disease and the emergence of multidrug resistance (MDR).

Adverse Drug Reactions (ADRs) are common causes of hospitalization and lead to large costs to society. There are two main financial burdens due to illnesses caused by ADRs: that of treating and that of avoiding them [12]. The occurrence of serious and fatal ADRs has been extensively studied in hospitalized patients and a meta-analysis of prospective studies in approximately forty hospitals in the United States of America (USA) suggests that 6-7% of hospitalized patients suffer from serious ADRs and 0.32% of patients develop fatal ADRs [13]. This results in approximately 100,000 deaths annually in the U.S. and an annual cost of over a hundred billion dollars to the society due to prolonged hospitalization and reduced productivity [3, 13]. Furthermore, it has been estimated that ADRs are responsible for up to 7% of all admissions in hospitals in the United Kingdom (UK) and 13% in medical clinics in Sweden [3], which shows the magnitude of this problem in the context of chemotherapy and drug development. Additionally, in France, a 10-year study in the Liver Unit of Hôpital Beaujon in Paris showed that among all patients hospitalized with acute hepatitis, 10% were due to adverse reaction to drugs and the prevalence of drug hepatotoxicity in patients older than fifty years exceeded 40%. In Japan and other Eastern countries, drugs are responsible for about 10-20% of cases of fulminant hepatitis [14].

Liver injury is the most common ADR and the main complication during chemotherapy since liver is the central organ for the biotransformation and excretion of most drugs and xenobiotics [14-17]. There are basically six mechanisms involving primarily the hepatocyte injury. The reactions of mono-oxygenase cytochrome P450 (CYP450) with certain drugs generate toxic metabolites that bind to intracellular proteins, leading to calcium homeostasis pump dysfunction with consequent disruption of actin fibers and cell lysis. Some drugs affect transport proteins in the cell membrane interrupting the flow of bile and then causing cholestasis. Several reactions involving CYP P450 can promote binding of the drug to the enzyme, with consequent exposure of this complex on the cell surface for recognition by T cells and antibody production as part of the autoimmune response. Finally, certain drugs may promote hepatic injury mediated by programmed cell death (apoptosis) or being capable of inhibiting respiration and/or mitochondrial beta-oxidation [17].

Xenobiotics are usually lipophilic and this facilitates their transport in association with lipoproteins in the blood stream and their penetration of lipid membranes and entrance into organs. However, physicochemical properties of drug molecules difficult their removal from the organism by biliary or renal excretion and therefore, these substances require enzymatic conversion to water soluble compounds [1]. The xenobiotics metabolism, often through multiple pathways, can generate metabolites that are more toxic than the substrate and through their interaction with target macromolecules such as DNA, RNA, proteins and receptors, generate the toxic effects. The organ affected is generally that responsible for drug metabolism or excretion of metabolites [1].

The enzyme systems responsible for the biotransformation of many drugs are located in the endoplasmic reticulum of the liver (microsomal fraction). Such enzymes are also present in the kidneys, lungs and gastrointestinal epithelium, although at a lower concentration [1]. The metabolic modification in biotransformation usually takes place in two consecutive steps and results in the loss of biological activity. Phase I reactions convert the xenobiotic into a metabolite with higher polarity by oxidation, reduction or hydrolysis and generates a pharmacologically inactive or less active, or in the case of a pro-drug, more active molecule. This metabolite is then either eliminated or go through Phase II reactions (so-called synthesis or conjugation reactions), involving binding to a primary metabolite or endogenous substrate such as glucuronate, sulfate, acetate, amino acids or glutathione (tripeptide). Such enzymatic reactions include glucuronidation, methylation, sulfation, acetylation, conjugation with glutathione and conjugation with glycine [1].

The risk for developing hepatotoxicity is associated both with genetic and acquired factors. The acquired factors include: age, gender, nutritional habits, drug abuse, pregnancy and extrahepatic disease. Genetic variations in isoenzymes involved in drug biotransformation can result in abnormal reactions leading to toxic effects [14,17]. In the case of INH in particular, advanced age is a risk factor for hepatotoxicity whereas deficiency in the ability of N-acetylation represent a genetic risk factor for liver injury.

INH is administered orally and rapidly absorbed through the gastrointestinal tract passing through the liver by the portal venous system before reaching the general circulation where is metabolized by a process known as the first pass effect with reduction of its bioavailability. About 75% to 95% of the INH is excreted by the kidneys during the first 24 hours, mainly as the metabolic forms acetyl-isoniazid and isonicotinic acid [1].

In the liver, INH is metabolized to acetylisoniazid by N-acetyltransferase 2 (NAT2), followed by hydrolysis to acetylhydrazine and then oxidized by cytochrome P4502E1 (CYP2E1) to hepatotoxic intermediates [18, 19]. These metabolites can destroy hepatocytes either by interfering with cell homeostasis or by triggering immunologic reactions in which reactive metabolites that are bound to hepatocyte plasma proteins may act as haptens [17]. The other metabolic pathway to generate toxic metabolites is direct hydrolysis of INH to hydrazine, a potent hepatotoxin. NAT2 is also responsible for converting acetylhydrazine to diacetylhydrazine, a nontoxic component [18, 20, 21] (Figure 1). Glutathione S-transferase (GST), an important phase II detoxification enzyme, is thought to play a protective role as an intracellular free radical scavenger, which conjugates glutathione with toxic metabolites that are generated

from CYP2E1 [22]. Sulphydryl conjugation facilitates the elimination of metabolites from the body and reduces the toxic effect [23] (Figure 1).




Figure 1. Schematic representation of the INH metabolism. The major enzymes involved in this pathway are indicated in boxes [20, 24].

In the last few years, an increasing number of studies have suggested that genetic polymorphisms in *NAT2*, *CYP2E1* and *GST* genes would be associated with susceptibility to drug-induced hepatotoxicity during TB treatment. The present work focused in an overview of the role of such polymorphisms in occurrence of liver injury induced by anti-TB drugs, and by INH in particular.

3. State of the art

3.1. N-acetyltransferase 2

NAT2, the main enzyme responsible for the metabolism and inactivation of INH in humans, is a Phase II enzyme that catalyzes the transfer of the acetyl group from the cofactor acetyl coenzyme A (acetyl-CoA) to the nitrogen terminal of the drug. Variations in activity of *NAT2* were discovered over 50 years ago when observing interindividual differences in the metabolism of INH and the level of drug-induced toxicity in TB patients. *NAT2* is encoded by the *NAT2* gene and according family genetic studies, variability of *NAT2* was directly related to the emergence of different phenotypes of acetylation [25].

The molecular study of human N-acetyltransferases revealed the presence of three genetic loci, two very homologous encoding the enzymes *NAT1* and *NAT2*, and a third including the

pseudogene *pNAT* (Figure 2). These loci are located on chromosome 8 between 170-360Kb at 8p22 [26]. The *pNAT* is a pseudogene containing a premature stop codon, and is not transcribed. *NAT1* and *NAT2* genes consist of 873 bp, are intronless, and encode proteins of 34 kDa. Protein sequence homology between both enzymes is 81% while that between their respective genes is 87%. Both enzymes have N-acetylation, O-acetylation and NO-transfer in different xenobiotics and carcinogens but differ considerably in their tissue distribution and expression levels during embryonic development [26-28].

Both *NAT1* and *NAT2* are polymorphic genes and SNPs in their coding region can alter the enzymatic activity [29, 30] and are the basis of the three major genetically determined phenotypes, being rapid, intermediate and slow acetylators, which are inherited as a codominant trait [31, 32]. The reference *NAT2**4 allele (without mutations / wild-type) and 66 variants were identified and classified in human populations depending on the combination of up to four SNPs present throughout the *NAT2* coding region [33]. So far, over 30 SNPs have been identified in this region, including several rare mutations described in different populations [34]. Among these, the seven most frequent are the 191 G>A (R64Q), 282 C>T (silent), 341 T>C (I114T), 481 C>T (silent), 590 G>A (R197Q), 803 A>G (K268R) and 857 G>A (G286T) SNPs identified in different human populations [35]. *NAT2* alleles containing the 191G>A, 341T>C, 590G>A or 857G>A SNPs are associated with slow acetylator *NAT2* alleles [33].




Figure 2. Schematic representation of *NAT* genes on human chromosome 8p22. Distribution of the seven most common SNPs in *NAT2*. *D8S21* represents a polymorphic marker situated in the *NAT2* locus [26, 36].

Presence of different SNPs in *NAT2* can be easily determined by genotyping procedures such as PCR-RFLP [37], allele specific PCR [38] or direct sequencing [39]. To achieve the *NAT2* genotype of each individual and predict the phenotype, the haplotype of both chromosomes is usually reconstructed using the statistic software (PHASEv2.1.1[40, 41]). Using haplotype data, many studies have reported the frequencies of the different acetylation profiles among ethnically different populations showing the high diversity around the world. In Asians and Ameridians, the fast acetylator phenotype is more frequent [42-44] whereas in Euro-descendants slow acetylators account for 50% of the study population [37,45]. The molecular basis for such discrepancy is that the most common *NAT2* allele in Euro-descendants is very rare in Asians and may represent a different selective advantage within the gene pools of these separate populations. Description of new alleles of *NAT2* is still occurring in recent studies [34].

In an attempt to establish an association between acetylation profiles and development of disease, cohort or case-control studies have been performed using of genotyping and phenotyping tools. Evidence was found for an association between the slow acetylator predicted phenotype and developing urinary bladder cancer, while rapid acetylators seem more susceptible to development of colon cancer. For a review, see [27, 46].

For many years, INH has been considered the main cause of hepatotoxicity during TB treatment and association studies between the acetylation phenotypes and susceptibility to liver-related ADRs have been performed. Two early studies conducted in oriental populations investigated the association of the acetylator phenotype with INH induced hepatotoxicity and observed an increased risk of developing hepatotoxicity by INH among the slow acetilators [47, 48]. This observation was confirmed in several other studies performed in different populations [49-52].

Several studies reported the absence of a relationship between acetylation status and hepatotoxicity during TB treatment [53-55] but some, suggested the rapid acetylators as more susceptible to side effects [55, 56]. Reasons for these different findings range from genotyping methods to ethnicity. In some studies, *NAT2* acetylation phenotypes were determined by an enzymatic method leading to possible misclassification of the acetylation status [53, 56, 57]. Indeed, it is difficult to compare the accuracy of different NAT phenotyping methods or different cut-off points using the same phenotyping method. In addition, for genotyping, investigators sometimes select a small number of SNPs to define the acetylation status [54, 55]. Since the frequencies of *NAT2* alleles are different among worldwide populations and new alleles are been identified in some countries, investigators need to characterize such alleles in their own study population in order to choose appropriate SNPs for genotyping and classify the acetylation status of individuals, otherwise overestimation of slow acetylators may be obtained, contributing to a spurious results in the association study.

Recently, a study with an admixed population showed that *NAT2* is a genetic factor for predisposition to anti-TB drug-induced hepatitis. In this case, *NAT2* genes were well characterized by direct sequencing and their genotypes achieved by haplotype reconstruction using the PHASE software. In addition, functional unknown genotypes were disconsidered and others confounding variables for hepatotocixity were taken into account. The incidence of elevated levels of serum transaminases was significantly higher in slow acetylators than those

of the rapid/intermediate type. These results corroborate with the current hypothesis that the acetylator status may be a risk factor for the hepatic side effects of isoniazid [58].

Finally, a meta-analysis was conducted to solve the problem of inadequate statistical power and controversial results based on accumulated data with small sample size [59]. Data from 14 studies performed between 2000 and 2011 were pooled and showed that TB patients with a slow acetylator genotype had a higher risk of anti-tuberculosis drug induced hepatotoxicity than patients with rapid or intermediate acetylation ($p < 0.001$). Moreover, subgroup analyses indicate that both Asians and non-Asians slow acetylators develop anti-tuberculosis drug induced hepatotoxicity more frequently. Additionally, there were statistically significant associations between NAT2*5/*7, NAT2*6/*6, NAT2*6/*7 and NAT2*7/*7 and the risk of anti-TB drug induced hepatotoxicity [59].

As a final consideration, NAT acetylates more slowly not only isoniazid but also acetylhydrazine, the immediate precursor of toxic intermediates, to the harmless diacetylhydrazine [60, 61]. This protective acetylation is further suppressed by INH competition. Therefore, slow acetylators may be prone to higher accumulation rates of INH toxic metabolites. Another important route to generate toxic intermediates is the direct hydrolysis of unacetylated INH [62], producing hydrazine that also induces hepatic injury [62, 63]. Pharmacokinetic studies showed that the serum concentration of hydrazine was significantly higher in slow acetylators than in rapid acetylators, probably due to the high INH concentration. The high amount of INH disposed of through this pathway is likely to lead to enhanced hydrolysis to hydrazine, since the rate of metabolic conversion of INH to acetylisoniazid is lower in slow than in rapid acetylators [64, 65]. All of these drug-disposal processes may support the finding that slow acetylators are prone to INH-induced hepatitis. We therefore conclude that screening of patients for the NAT2 genetic polymorphisms can prove clinically useful for the prediction and prevention of anti-tuberculosis drug induced hepatotoxicity.

3.2. CYP450

Cytochromes P450 (CYP450) are hemoproteins and form the most important enzymatic group for Phase I biotransformation. The main activity of isozymes of CYP450 system is oxidation and they are located in the smooth endoplasmic reticulum, mainly in liver cells. However, these mono-oxygenases are also localized in the intestine, pancreas, brain, lung, kidney, bone marrow, skin, ovary and testicles [66]. The CYP450 proteins are clustered into families and subfamilies according to the similarity between the amino acid sequences: where family members have $\geq 40\%$ identity in amino acid sequence, members of the same subfamily share $\geq 55\%$ identity [67].

The CYP450s are responsible for the metabolism of several endogenous substrates and the synthesis of hydrophobic lipids such as cholesterol, steroid hormones, bile acids and fatty acids. Moreover, some enzymes of P450 complex metabolize exogenous substances including drugs, environmental chemicals and pollutants as well as products derived from plants. The metabolism of exogenous substances by CYP450 usually results in detoxification of the xenobiotic; however, the reactions triggered by such enzymes can

lead to generation of toxic metabolites that contribute to the increased risk of developing cancers and other toxic effects [68].

The complete sequencing of the human genome revealed the presence of about 115 genes of CYP450, including 57 active genes and 58 pseudogenes [67]. They belong to families 1-3 and are responsible for 70-80% of Phase I-dependent metabolism of clinically used drugs. Other families of CYPs are involved in metabolism of endogenous components [66]. The CYP2 constitutes the largest family of isoenzymes and comprises one third of all human CYPs. Genes encoding these enzymes are polymorphic and the frequency distribution of allelic variants in different ethnic groups differs. Overall, four phenotypes based on genotypes can be identified: (i) poor metabolizers who present low enzymatic activity, (ii) intermediate metabolizers, usually heterozygous for a defective allele, (iii) rapid metabolizers, who have two normal alleles and (iv) ultrarapid metabolizers, who have several gene copies [69].

The enzyme CYP2E1 is expressed mainly in the liver but can be found in other organs such as kidney, gastrointestinal tract and brain and involved in oxidation of substrates such as ethanol and the metabolism of many drugs and pre-carcinogens. Besides ethanol, CYP2E1 can be induced by various drugs such as INH but also by hydrocarbons, benzene, chloroform and various organic solvents [70].

The activity of CYP2E1 is also modulated by polymorphisms in several locations of its gene and more activity of this enzyme may increase the synthesis of hepatotoxins. Two polymorphisms upstream of the *CYP2E1* transcriptional start site are characterized by *Pst* I and *Rsa* I digestion and appear to be in complete linkage disequilibrium (Figure 3). These two polymorphisms are located in a putative HNF-q binding site and thus may play a role in the regulation of *CYP2E1* transcription and subsequent protein expression [71]. Genotypes of *CYP2E1* are classified as being *1A/*1A, *1A/*5 or *5/*5 by *Rsa* I based restriction analysis. The polymorphism detectable by *Dra* I (7632 T>A) is located in intron 6 and characterizes the allelic variant *CYP2E1**6. The other polymorphism is an insertion/deletion of 96 bp (*CYP2E1**1D and *1C alleles) that regulates the expression of the gene [72]. Some studies have shown that allelic variants *CYP2E1* *5, *6 and *1D would increase enzyme activity [71, 73]. However, other authors did not confirm any relationship with these polymorphisms with CYP2E1 activity [74].




Figure 3. Polymorphic and corresponding restriction enzyme cutting sites at *CYP2E1* [24].

Several studies have described the involvement of polymorphisms in *CYP2E1* in cancer development but results are controversial. The studies showed that the frequency of SNP -1053 C>T in the promoter region varies significantly in different ethnic groups. The mutant allele is present with a frequency of 2-8% in Euro-descendants but varies in Asia from 25 to 36% [75].

In 2003, Huang and coworkers showed an association of the wild-type genotype *1A/*1A with risk of developing liver damage induced by isoniazid in adult TB patients, regardless of their profile of acetylation (OR 2.52; 95% CI 1.26 to 5.05) [76]. Later, Vuilleumier and colleagues showed association between this CYP and isoniazid-induced hepatotoxicity, without hepatitis, during chemoprophylaxis for TB (OR 3.4; 95% CI 1.1 to 12; $p = 0.02$). The risk of having high levels of liver enzymes was 3.4-fold higher when compared with all other *CYP2E1* genotypes [55]. Another study on Indian children with TB showed association between risk of hepatotoxicity and polymorphisms in *CYP2E1*, despite of low sample size [77]. However, a study with on a Korean population found no relationship between hepatic adverse effects with genotype *1A/*1A of *CYP2E1* during anti-TB treatment [51]. Lack of association between this CYP and antituberculosis drug-induced liver injury was also observed in Brazil [58]. The discrepancy of these results may be due to differences in the frequencies of *CYP2E1**1A and *CYP2E1**5 alleles among the populations and the different criteria to define hepatotoxicity used.

Finally, *CYP2E1* converts acetyl hydrazine into hepatotoxins like acetyldiazene, ketene and acetylonium ion. The reaction of acetyl hydrazine (at high levels) with *CYP2E1* leads to covalent binding of these secondary metabolites with intracellular proteins (Figure 1). As a consequence, intracellular changes occur resulting in loss of ionic gradients and decrease of ATP levels and consequent disruption of actin followed by cell lysis. Further studies in different populations and with a larger sample size are needed to determine the true influence of *CYP2E1* gene polymorphisms on the occurrence of liver injury during treatment for TB.

3.3. Glutathione S-transferases

Glutathione S-transferases constitute a superfamily of multifunctional ubiquitous enzymes that play an important role in cellular detoxification by protecting macromolecules against reactive electrophilic attack. The GSTs are Phase II enzymes that catalyze the nucleophilic attack of glutathione (GSH) into components that contain an electrophilic carbon, nitrogen or sulfur atom. The combination of the GSH with these compounds often leads to formation of less reactive and more water soluble products, more easily excreted by the body [23, 78].

Glutathione transferases are of great interest to pharmacologists and toxicologists, since they are drug targets for the treatment of asthma and cancer, in addition to metabolize drugs, insecticides, herbicides, carcinogens and products of oxidative stress. Polymorphisms in *GST* genes are often correlated with susceptibility to various cancers, as well as alcoholic liver disease [23, 78-81].

In humans, eight gene families of soluble (or cytosolic) GSTs have been described: alpha (α) located on chromosome 6, mu (μ) on chromosome 1, theta (θ) on chromosome 22, pi (π) on chromosome 11; zeta (ζ) on chromosome 14, sigma (σ) on chromosome 4; kappa (κ) (chromo-

somal location not given) and omega (Ω) on chromosome 10 [80]. This classification is based on amino acid sequences, substrate specificity, chemical affinity, protein structure and enzyme kinetics. These enzymes are highly expressed in the liver and constitute up to 4% of total soluble proteins but can be seen in several other tissues [82]. GSTs have an overlap of specific substrates and the deficiency in one isoform can be compensated by other isoforms. Glutathione S-transferase mu (GSTM), glutathione S-transferase theta (GSTT) and glutathione S-transferase Pi (GSTP) have been the most studied isoform [83-88].

The subfamily GST mu is encoded by five genes arranged in tandem (5'-*GSTM4*-*GSTM2*-*GSTM1*-*GSTM5*-*GSTM3*-3'), forming a 100 kb gene cluster on chromosome 1p13.3 (Figure 4). Polymorphisms have been identified and clinical consequences of genotypes resulting from combinations of alleles *GSTM1**0, *GSTM1**A, and *GSTM1**B have been widely investigated [78, 81, 89, 90]. Individuals who possess the homozygous null for *GSTM1* (*GSTM1**0/*GSTM1**0) do not express this protein. Thus, the absence of this gene can cause an increased accumulation of reactive metabolites in the body, increasing the interaction with cellular macromolecules and tumor initiation process. *GSTM1**A and *GSTM1**B differ in only one base in exon 7 and encode monomers that form active dimers. The catalytic activity of these enzymes are very similar [91].

The *GSTM1* gene is flanked by two almost identical 4.2-kb regions. *GSTM1**0 originates from homologous recombination between the two repeat regions which results in a 16 Kb deletion containing the entire gene *GSTM1* (Figure 4). *GSTM1* is precisely excised leaving the adjacent *GSTM2* and *GSTM5* genes intact [78]. In a study of liver specimens of 168 autopsied Japanese subjects, observed was that the *GSTM1**0 null allele was more frequent in livers with hepatitis and hepatocellular carcinoma compared to control livers [92].




Figure 4. Structural localization of 100 kb gene cluster encoding the GST mu subfamily (chromosome 1p13.3). The figure indicates the homologous recombination event that can happen causing the null allele (*GSTM1**0 - no *GSTM1*). Figure adapted from [78].

The subfamily GST theta consists of two genes, *GSTT1* and *GSTT2*, located on chromosome 22q11.2 and separated by approximately 50 Kb (Figure 5). Analysis of the 119 Kb portion

containing these genes revealed two regions flanking *GSTT1*, HA3 and HA5, with more than 90% homology. HA3 and HA5 contain two identical 403-bp repeats and the occurrence of *GSTT1*0* allele is probably caused by homologous recombination between the two regions [78]. In humans, *GSTT1* is also expressed in erythrocytes and probably plays a global role in early detoxification of xenobiotics and carcinogens.




Figure 5. Structural localization of gene cluster encoding the GST subfamily theta (chromosome 22q11.2). The *GSTT1* null allele (*GSTT1*0*) arises by homologous recombination of the left and right 403-bp repeats, which results in a 54-kb deletion containing the entire *GSTT1* gene. Figure adapted from [78].

Deficiencies in the GST activity due to the null genotypes of *GSTM1* and *GSTT1* may modulate susceptibility to the development of hepatotoxicity induced by drugs and xenobiotics. Furthermore, it was observed that the frequencies of *GSTT1*0* and *GSTM1*0* alleles vary within different ethnic groups [78, 82]. Liver injury induced by INH has been associated with the depletion of glutathione content and reduction of GST activity in an animal model for hepatotoxicity by anti-TB drugs [22].

In 2001, Roy and colleagues demonstrated that individuals, homozygous for the null *GSTM1*, had a relative risk of 2.12 for developing hepatotoxicity induced by anti-TB drugs. However, these authors found no association of the *GSTT1* null genotype with this side effects [54]. Similarly, another study in the Thai population found that only the *GSTM1* null genotype increases the risk of liver injury (OR 2.23, 95% CI 1.07 to 4.67) [93]. The opposite was observed by Leiro and colleagues: individuals with the *GSTT1* null genotype had an increased risk of developing hepatotoxicity induced by anti-TB drugs and no significant association was observed between *GSTM1*0/*0* genotype and liver injury [94]. These studies suggest a protective effect of glutathione S-transferases to the hepatotoxic effects of isoniazid.

On the other hand, recent studies in different population showed no relationship between *GSTM1*0/*0* or *GSTT1*0/*0* genotypes and liver injury during anti-TB treatment [58, 95, 96]. In a population-based prospective antituberculosis treatment cohort in China, a more robust case-control study was conducted and there was no statistically significant association between null genotypes and hepatotoxicity induced by anti-TB drugs [97].

These controversial results may be due to the small sample size in many studies and the different frequencies of the null genotypes. New populations should be evaluated with large sample size to see which of these polymorphisms can be used as genetic markers for the risk of side effects during anti-TB treatment.

4. Conclusion

The concept of personalized medicine is not really new, but it has been receiving increasing attention in recent years for improvement of drug regulation and medical guidelines. There is considerable interindividual variability in metabolism, partly due to human differences on a genetic level. Genetic polymorphisms in drug-metabolizing enzymes can affect enzyme activity and may cause differences in treatment response or drug toxicity, for example, due to an increased formation of reactive metabolites. Such polymorphisms may explain differences in incidence of anti-TB drugs induced hepatotoxicity between different populations.

Genotyping cannot completely predict the phenotype on an individual level because of the additional contribution of epigenetic, endogenous and environmental factors. However, pharmacogenetics is able to add important information in many cases where therapeutic drug scheme is inappropriate or not sufficient. Nowadays, we can cite three examples of personalized medicine application in clinical practice, (i) AIDS treatment (abavir / skin hypersensitivity / *HLA-B*5701*), (ii) anticoagulation (warfarin / bleeding / *CYP2C9*) and (iii) treatment of acute lymphoblastic leukemia (azathioprine / treatment resistance / *TPMT*) [98].

Although limited information exists regarding isoniazid concentrations that cause toxic reactions, it has been proposed to adjust isoniazid dosage depending on individuals acetylator status: a lower dosage for slow acetylators to reduce the risk of liver injury and a higher isoniazid dosage for fast acetylators to increase the early bactericidal activity and thereby lower the probability of treatment failure [50]. However, more robust clinical prospective studies are needed to evaluate the real contribution of these different polymorphisms in the occurrence of liver side effects during anti-TB treatment. Future studies should include larger sample size, different ethnic population, simultaneous analysis of different genetic markers, different degrees of liver injury and consideration of possible confounding factors.

Author details

Raquel Lima de Figueiredo Teixeira*, Márcia Quinhones Pires Lopes, Philip Noel Suffys and Adalberto Rezende Santos

*Address all correspondence to: raquelft@ioc.fiocruz.br

Laboratory of Molecular Biology Applied to Mycobacteria – Oswaldo Cruz Institute – Fiocruz, Rio de Janeiro, Brazil

References

- [1] Hardman, JG., Limbird LE, Molinoff PB, Ruddon RW, Goodman AG. Goodman and Gilman's The Pharmacological Basis of Therapeutics. 9th ed. New York, NY: McGraw-Hill, 1996.
- [2] Prows CA, Prows DR. Medication Selection by genotype: How genetics is changing drug prescribing and efficacy. *American Journal of Nursing* 2004; 104 60-70.
- [3] Ingelman-Sundberg M. Pharmacogenetics: an opportunity for a safer and more efficient pharmacotherapy. *Journal of Internal Medicine* 2001; 250 186-200.
- [4] Roses AD. Pharmacogenetics place in modern medical science and practice. *Life Sciences* 2002; 70 1471-1480.
- [5] Sachidanandam R, Weissman D, Schmidt SC, Kakol JM, Stein LD, Marth G, Sherry S, Mullikin JC, Mortimore BJ, Willey DL, Hunt SE, Cole CG, Coggill PC, Rice CM, Ning Z, Rogers J, Bentley DR, Kwok PY, Mardis ER, Yeh RT, Schultz B, Cook L, Davenport R, Dante M, Fulton L, Hillier L, Waterston RH, McPherson JD, Gilman B, Schaffner S, Van Etten WJ, Reich D, Higgins J, Daly MJ, Blumenstiel B, Baldwin J, Stange-Thomann N, Zody MC, Linton L, Lander ES, Altshuler D. International SNP Map Working Group. A map of human genome sequence variation containing 1.42 million single nucleotide polymorphisms. *Nature* 2001; 409(6822) 928-33.
- [6] Hall II RG, Leff RD, Gumbo T. Treatment of Active Pulmonary Tuberculosis in Adults: Current Standards and Recent Advances: Insights from the Society of Infectious Diseases Pharmacists. *Pharmacotherapy* 2009; 29(12) 1468-1481.
- [7] Blomberg B, Fourie B. Fixed-dose combination drugs for tuberculosis: application in standardised treatment regimens. *Drugs* 2003;63(6) 535-53.
- [8] Garibaldi RA, Drusin RE, Ferebee SH, Gregg MB. Isoniazid-associated hepatitis. Report of an outbreak. *American Review of Respiratory Disease*. 1972; 106 357-365.
- [9] Saukkonen JJ, Cohn DL, Jasmer RM, Schenker S, Jereb JA, Nolan CM, Peloquin CA, Gordin FM, Nunes D, Strader DB, Bernardo J, Venkataramanan R, Sterling TR. An Official ATS Statement: Hepatotoxicity of Antituberculosis Therapy. *American Journal of Respiratory and Critical Care Medicine* 2006; 174 935-952.
- [10] Xue HY, Hou YN, Liu HC. The general investigation of the increased hepatotoxicity caused by isoniazid in combination with rifampin. *Chinese Journal Modern Applied Pharmacy* 2002; 19 463-465.
- [11] Tostmann A, Boeree MJ, Aarnoutse RE, Lange WC, van der Ven AJ, Dekhuijzen R. Antituberculosis drug-induced hepatotoxicity: concise up-to-date review. *Journal of Gastroenterology and Hepatology* 2007; 6 1440-1446.

- [12] Lundkvist J, Jonsson B. Pharmacoeconomics of adverse drug reactions. *Fundamental and Clinical Pharmacology* 2004; 18 275-280.
- [13] Hug H, Bagatto D, Dannecker R, Schindler R, Horlancher O, Gut J. ADRIS – The adverse drug reactions information scheme. *Pharmacogenetics*. 2003; 13:767-772.
- [14] Larrey D. Epidemiology and individual susceptibility to adverse drug reactions affecting the liver. *Seminars in Liver Disease* 2002; 22 145-155.
- [15] Bagheri H, Michel F, Lapeyre-Mestre M, Lagier E, Cambus JP, Valdiguié P, Montastruc JL. Detection and incidence of drug-induced injuries in hospital: a prospective analysis from laboratory signal. *British Journal of Clinical Pharmacology* 2000; 50 479-484.
- [16] Bissell DM, Gores GJ, Laskin DL, Hoofnagle JH. Drug-induced liver injury: mechanisms and test systems. *Hepatology* 2001; 33 1009-1013.
- [17] Lee WM. Drug-induced hepatotoxicity. *New England Journal of Medicine* 2003; 349 474-485.
- [18] Nelson SD, Mitchell JR, Timbrell JA, Snodgrass WR, Corcoran GB. Isoniazid and iproniazid: activation of metabolites to toxic intermediates in man and rat. *Science* 1976; 193(4256) 901-903.
- [19] Timbrell JA, Mitchell JR, Snodgrass WR, Nelson SD. Isoniazid hepatotoxicity: the relationship between covalent binding and metabolism in vivo. *Journal of Pharmacology and Experimental Therapeutics* 1980; 213 364-369.
- [20] Mitchell JR, Snodgrass WR, Gillette JR. The Role of Biotransformation in Chemical-Induced Liver Injury. *Environmental Health Perspectives* 1976; 15 27-38.
- [21] Woodward KN, Timbrell JA. Acetylhydrazine hepatotoxicity: the role of covalent binding. *Toxicology* 1984; 30 65-74.
- [22] Sodhi CP, Rana SV, Mehta SK, Vaiphei K, Attri S, Thakur S, Mehta S. Study of oxidative stress in isoniazid-induced hepatic injury in young rats with and without protein-energy malnutrition. *Journal of Biochemical and Molecular Toxicology* 1996; 11 139-146.
- [23] Hayes JD, Flanagan JU, Jowsey IR. Glutathione transferases. *Annual Review of Pharmacology and Toxicology* 2005; 45 51-88.
- [24] Roy PD, Majumder M, Roy B. Pharmacogenomics of anti-TB drugs-related hepatotoxicity. *Pharmacogenomics* 2008; 9 311-321.
- [25] Evans DAP. N-acetyltransferase. *Pharmacology & Therapeutics* 1989; 42:157-234
- [26] Sim E, Payton M, Noble M, Minchin R. An update on genetic, structural and functional studies of arylamine N-acetyltransferases in eucaryotes and procaryotes. *Human Molecular Genetics* 2000; 9 2435-41.

- [27] Hein DW, Doll MA, Fletland AJ, Leff MA, Webb SJ, Xiao GH, Devanaboyina US, Nangju NA, Feng Y. Molecular genetics and epidemiology of the NAT1 and NAT2 acetylation polymorphisms. *Cancer Epidemiology, Biomarkers and Prevention* 2000; 9 29-42.
- [28] Hein DW. Molecular genetics and function of NAT1 and NAT2: role in aromatic amine metabolism and carcinogenesis. *Mutation Research* 2002; 506-507 65-77.
- [29] Fretland AJ, Leff MA, Doll MA, Hein DW. Functional characterization of human N-acetyltransferase 2 (NAT2) single nucleotide polymorphisms. *Pharmacogenetics* 2001; 11 207-215.
- [30] Zang Y, Doll MA, Zhao S, States JC, Hein DW. Functional characterization of single-nucleotide polymorphisms and haplotypes of human N-acetyltransferase 2. *Carcinogenesis* 2007; 28 1665-1671.
- [31] Parkin DP, Vandenplas S, Botha FJ, Vandenplas ML, Seifart HI, van Helden PD, van der Walt BJ, Donald PR, van Jaarsveld PP. Trimodality of isoniazid elimination: phenotype and genotype in patients with tuberculosis. *American Journal of Respiratory and Critical Care Medicine* 1997; 155 1717-1722.
- [32] Chen B, Zhang WX, Cai WM. The influence of various genotypes on the metabolic activity of NAT2 in Chinese population. *European Journal of Clinical Pharmacology* 2006; 62 355-359.
- [33] Consensus Human Arylamine N-Acetyltransferase Gene Nomenclature www.louisville.edu/medschool/pharmacology/NAT.html (accessed 15 August 2012).
- [34] Teixeira RL, Silva Jr FP, Silveira AR, Cabello PH, Mendonça-Lima L, Rabahi MF, Kritski AL, Mello FC, Suffys PN, de Miranda AB, Santos AR. Sequence analysis of NAT2 gene in Brazilians: identification of undescribed single nucleotide polymorphisms and molecular modeling of the N-acetyltransferase 2 protein structure. *Mutation Research* 2010; 683 43-49.
- [35] García-Martín E. Interethnic and intraethnic variability of NAT2 single nucleotide polymorphisms. *Current Drug Metabolism* 2008; 9 (6) 487-497.
- [36] Boukouvala S, Sim E. Structural Analysis of the Genes for Human Arylamine N-Acetyltransferases and Characterisation of Alternative Transcripts. *Basic & Clinical Pharmacology & Toxicology* 2005; 96 343-351.
- [37] Cascorbi I, Drakoulis N, Brockmoller J, Maurer A, Sperling K, Roots I. Arylamine N-acetyltransferase (NAT2) mutations and their allelic linkage in unrelated Caucasian individuals: Correlation with phenotypic activity. *American Journal of Human Genetics* 1995; 57 581-592.
- [38] Lin HJ, Han CY, Lin BK, Hardy S. Slow acetylators mutations in the human polymorphic N-acetyltransferase gene in 786 Asians, blacks, Hispanics, and whites: applica-

- tion to metabolic epidemiology. *American Journal of Human Genetics* 1993; 52(4) 827-34.
- [39] Teixeira RL, Miranda AB., Pacheco AG, Lopes MQ, Fonseca-Costa J, Rabahi MF, Melo HM, Kritski AL, Mello FC, Suffys PN, Santos AR. Genetic profile of the arylamine N-Acetyltransferase 2 coding gene among individuals from two different regions of Brazil. *Mutation Research* 2007; 624 31-40.
 - [40] Stephens M, Smith NJ, Donnelly P. A new statistical method for haplotype reconstruction from population data. *American Journal of Human Genetics* 2001; 68 978-989.
 - [41] Stephens M, Donnelly P. A comparison of Bayesian methods for haplotype reconstruction. *American Journal of Human Genetics* 2003; 73 1162-1169.
 - [42] Xie HG, Xu ZH, Ou-Yang DS, Shu Y, Yang DL, Wang JS, Yan XD, Huang SL, Wang W, Zhou HH. Meta-analysis of phenotype and genotype of NAT2 deficiency in Chinese populations. *Pharmacogenetics* 1997; 7(6) 503-14.
 - [43] Sekine A, Saito S, Iida A, Mitsunobu Y, Higuchi S, Harigae S, Nakamura Y. Identification of single-nucleotide polymorphisms (SNPs) of human N-acetyltransferase genes NAT1, NAT2, AANAT, ARD1 and L1CAM in the Japanese population. *Journal of Human Genetics* 2001; 46 314-319.
 - [44] Jorge-Nebert LF, Eichelbaum M, Griese EU, Inaba T, Arias TD. Analysis of six SNPs of NAT2 in Ngawbe and Embera Amerindians of Panama and determination of the Embera acetylation phenotype using caffeine. *Pharmacogenetics* 2002; 12 39-48.
 - [45] Cascorbi I, Brockmoller J, Mrozikiewicz PM, Muller A, Roots I. Arylamine N-acetyltransferase activity in man. *Drug Metab Rev* 1999; 31 489-502.
 - [46] Hein, D. N-acetyltransferase 2 genetic polymorphism: Effects of carcinogen and haplotype on urinary bladder cancer risk. *Oncogene* 2006; 25(11) 1649-1658.
 - [47] Ohno M, Yamaguchi I, Yamamoto I, Fukuda T, Yolota S, Maekura R, Ito M, Yamamoto Y, Ogura T, Maeda K., Komuta K, Igarashi T, Azuma J. Slow N-acetyltransferase 2 genotype affects the incidence of isoniazid and rifampicin-induced hepatotoxicity. *International Journal of Tuberculosis Lung Disease*. 2000; 4 256-261.
 - [48] Huang YS, Chern HD, Su WJ, Wu JC, Lai SL, Yang SY, Chang FY, Lee SD. Polymorphism of the N-acetyltransferase 2 gene as a susceptibility risk factor for antituberculosis drug-induced hepatitis. *Hepatology* 2002; 35 883-889.
 - [49] Kinzig-Schippers M, Tomalik-Scharte D, Jetter A, Scheidel B, Jakob V, Rodamer M, Cascorbi I, Doroshenko O, Sörgel F, Fuhr U. Should we use N-acetyltransferase type 2 genotyping to personalize isoniazid doses? *Antimicrobial Agents and Chemotherapy* 2005; 49 1733-1738.
 - [50] Shimizu Y, Dobashi K, Mita Y, Endou K, Moriya S, Osano K, Koike Y, Higuchi S, Yabe S, Utsugi M, Ishizuka T, Hisada T, Nakazawa T, Mori M. DNA microarray gen-

otyping of N-acetyltransferase 2 polymorphism using carbodiimide as the linker for assessment of isoniazid hepatotoxicity. *Tuberculosis* 2006; 86(5) 374-81.

- [51] Cho HJ, Koh WJ, Ryu YJ, Ki CS, Nam MH, Kim JW, Lee SY. Genetic polymorphisms of NAT2 and CYP2E1 associated with antituberculosis drug-induced hepatotoxicity in Korean patients with pulmonary tuberculosis. *Tuberculosis* 2007; 87:551-556.
- [52] Higuchi N, Tahara N, Yanagihara K, Fukushima H, Suyama N, Inoue Y, Miyazaki Y, Kobayashi T, Yoshiura K, Niikawa N, Wen CY, Isomoto H, Shikuwa S, Omagari K, Mizuta Y, Kohno S, Tsukamoto K. NAT2*6A, a haplotype of the N-acetyltransferase 2 gene, is an important biomarker for risk of anti-tuberculosis drug-induced hepatotoxicity in Japanese patients with tuberculosis. *World J Gastroenterology* 2007; 13 6003-6008.
- [53] Singh J, Arora A, Garg PK, Thakur VS, Pande JN, Tandon RK. Antituberculosis treatment-induced hepatotoxicity: role of predictive factors. *Postgraduate Medical Journal* 1995; 71 359-362.
- [54] Roy B, Chowdhury A, Kundu S, Santra A, Dey B, Chakraborty M, Majumder PP. Increased risk of antituberculosis drug-induced hepatotoxicity in individuals with glutathione S-transferase M1 "null" mutation. *Journal of Gastroenterology and Hepatology* 2001; 16 1033-1037.
- [55] Vuilleumier N, Rossier MF, Chiappe A, Degoumois F, Dayer P, Mermillod B, Nicod L, Desmeules J, Hochstrasser D. CYP2E1 genotype and isoniazid-induced hepatotoxicity in patients treated for latent tuberculosis. *European Journal of Clinical Pharmacology* 2006; 62 423-429.
- [56] Mitchell JR, Thorgeisson UP, Black M, Timbrell JA, Snodgrass WR, Potter WZ, Jollow HR, Keiser HR. Increased incidence of isoniazid hepatitis in rapid acetylators: possible relation to hydralazine metabolites. *Clinical Pharmacology & Therapeutics* 1975; 18 70-79.
- [57] Yamamoto T, Suou T, Hirayama C. Elevated serum aminotransferase induced by isoniazid in relation to isoniazid acetylator phenotype. *Hepatology* 1986; 6 295-298.
- [58] Teixeira RL, Morato RG, Cabello PH, Muniz LM, Moreira Ada S, Kritski AL, Mello FC, Suffys PN, Miranda AB, Santos AR. Genetic polymorphisms of NAT2, CYP2E1, GST enzymes and the occurrence of antituberculosis drug-induced hepatitis in Brazilian TB patients. *Memórias do Instituto Oswaldo Cruz* 2011; 106(6) 716-24.
- [59] Wang PY, Xie SY, Hao Q, Zhang C, Jiang BF. NAT2 polymorphisms and susceptibility to anti-tuberculosis drug-induced liver injury: a meta-analysis. *The International Journal of Tuberculosis and Lung Disease* 2012; 16(5) 589-95.
- [60] Ellard GA, Gammon PT. Pharmacokinetics of isoniazid metabolism in man. *Journal of Pharmacokinetics and Biopharmaceutics* 1976; 4 83-113.

- [61] Lauterburg BH, Smith CV, Todd EL, Mitchell JR. Pharmacokinetics of the toxic hydrazine metabolites formed from isoniazid in humans. *J Pharmacol Exp Ther* 1985; 235 566-570.
- [62] Timbrell JA, Wright JM, Baillie TA. Monoacetylhydrazine as a metabolite of isoniazid in man. *Clinical Pharmacology & Therapeutics* 1977; 22 602-608.
- [63] Timbrell JA, Mitchell JR, Snodgrass WR, Nelson SD. Isoniazid hepatotoxicity: the relationship between covalent binding and metabolism in vivo. *Journal of Pharmacology and Experimental Therapeutics* 1980; 213 364-369.
- [64] Sarma GR, Immanuel C, Kailasam S, Narayana ASL, Venkatesan P. Rifampin-induced release of hydrazine from isoniazid: a possible cause of hepatitis during treatment of tuberculosis with regimens containing isoniazid and rifampin. *American Review of Respiratory Disease* 1986; 133 1072-1075.
- [65] Fukino K, Sasaki Y, Hirai S, Nakamura T, Hashimoto M, Yamagishi F, Ueno K. Effects of NAT2, CYP2E1 and GST genotypes on the serum concentrations of isoniazid and metabolites in tuberculosis patients. *The Journal of Toxicological Sciences* 2008; 33 187-95.
- [66] Wijnen PAHM, Op Den Buijsch RAM, Drent M, Kuipers PMJC, Neef C, Bast A, Bekers O, Koek GH. Review article: the prevalence and clinical relevance of cytochrome P450 polymorphisms. *Alimentary Pharmacology & Therapeutics* 2007; 26 Supl: 211-219.
- [67] The CYP 450 nomenclature. <http://drnelson.uthsc.edu/CytochromeP450.html> (accessed 15 August 2012).
- [68] Nebert DW, Russell DW. Clinical importance of the cytochromes P450. *Lancet* 2002; 360 1155-1162.
- [69] Ingelman-Sundberg M. Pharmacogenetics of cytochrome P450 and its applications in drug therapy: the past, present and future. *Trends in Pharmacological Sciences* 2004; 25 193-200.
- [70] Caro AA, Cederbaum AI. Oxidative stress, toxicology, and pharmacology of CYP2E1. *Annual Review of Pharmacology and Toxicology* 2004; 44 27-42.
- [71] Watanabe J, Hayashi S, Kawajiri K. Different regulations and expression of the human CYP2E1 gene due to the Rsa I polymorphism in the 5' flanking region. *The Journal of Biochemistry* 1994; 116 321-326.
- [72] CYP2E1 allele nomenclature. www.cypalleles.ki.se/cyp2e1.htm (accessed 15 August 2012).
- [73] Carriere V, Berthou F, Baird S, Belloc C, Beaune P, Waziers ID. Human cytochrome P4502E1 (CYP2E1): from genotype to phenotype. *Pharmacogenetics* 1996; 6 203-211.

- [74] Powell H, Kitteringham NR, Pirmohamed M, Smith DA, Perik BK. Expression of cytochrome P-4502E1 in human liver: assessment by mRNA, genotype and phenotype. *Pharmacogenetics* 1998; 8 411-421.
- [75] Neuhaus T, Ko YD, Lorenzen K, Fronhoffs S, Harth V, Bröde P, Vetter H, Bolt HM, Pesch B, Brüning T. Association of cytochrome P450 2E1 polymorphisms and head and neck squamous cell cancer. *Toxicology* 2004; 151 273-282.
- [76] Huang YS, Chern HD, Su WJ, Wu JC, Chang SC, Chiang CH, Chang FY, Lee SD. Cytochrome P450 2E1 genotype and the susceptibility to antituberculosis drug-induced hepatitis. *Hepatology* 2003; 37 924-930.
- [77] Roy B, Ghosh SK, Sutraghar D, Sikdar N, Mazumder S, Barman S. Predisposition of antituberculosis drug induced hepatotoxicity by cytochrome P450 2E1 genotype and haplotype in pediatric patients. *Journal of Gastroenterology and Hepatology* 2006; 21 781-786.
- [78] Parl FF. Glutathione S-transferase genotypes and cancer risk. *Cancer Letters* 2005; 221 123-129.
- [79] Strange RC, Jones PW, Fryer AA. Glutathione S-transferase: genetics and role in toxicology. *Toxicology Letters* 2000; 112-113 357-363.
- [80] Strange RC, Spiteri MA, Ramachandran S, Fryer AA. Glutathione-S-transferase family of enzymes. *Mutation Research* 2001; 482 21-26.
- [81] Coughlin SS, Hall IJ. Glutathione S-transferase polymorphisms and risk of ovarian cancer: A HuGE review. *Genetics Medicine* 2002; 4 250-257.
- [82] Landi S. Mammalian class theta GST and differential susceptibility to carcinogens: a review. *Mutation Research* 2000; 463 247-283.
- [83] Pemble S, Schroeder KR, Spencer SR, Meyer DJ, Hallier E, Bolt HM, Brian Ketterer, Taylor JB. Human glutathione S-transferase Theta (GSTT1): cDNA cloning and the characterization of a genetic polymorphism. *Biochemical Journal* 1994; 300 271-276.
- [84] Geisler SA, Olshan AF. GSTM1, GSTT1, and the Risk of Squamous Cell Carcinoma of the Head and Neck: A Mini-HuGE Review. *American Journal of Epidemiology* 2001; 154 95-105.
- [85] Henrion-Caude A, Roussey C, Housset M, Flahault C, Fryer A, Chadelat AA, Strange RC, Clement A. Liver disease in pediatric patients with cystic fibrosis is associated with glutathione S-transferase P1 polymorphism. *Hepatology* 2002; 36 913-917.
- [86] Vineis P, Veglia F, Anttila S, Benhamou S, Clapper ML, Dolzan V, Ryberg D, Hirvonen A, Kremers P, Le Marchand L, Pastorelli R, Rannug A, Romkes M, Schoket B, Strange RC, Garte S, Taioli E. CYP1A1, GSTM1 and GSTT1 polymorphisms and lung cancer: a pooled analysis of gene-gene interactions. *Biomarkers* 2004; 9 298-305.
- [87] Raimondi S, Paracchini V, Autrup H, Barros-Dios JM, Benhamou S, Boffetta P, Cote ML, Dialyna IA, Dolzan V, Filiberti R, Garte S, Hirvonen A, Husgafvel-Pursiainen K,

- Imyanitov EN, Kalina I, Kang D, Kiyohara C, Kohno T, Kremers P, Lan Q, London S, Povey AC, Rannug A, Reszka E, Risch A, Romkes M, Schneider J, Seow A, Shields PG, Sobti RC, Sørensen M, Spinola M, Spitz MR, Strange RC, Stücker I, Sugimura H, To-Figueras J, Tokudome S, Yang P, Yuan JM, Warholm M, Taioli E. Meta- and Pooled Analysis of GSTT1 and Lung Cancer: A HuGE-GSEC Review. *American Journal of Epidemiology* 2006; 164 1027–1042
- [88] Holley SL, Fryer AA, Haycock JW, Grubb SEW, Strange RC, Hoban PR. Differential effects of glutathione S-transferase pi (GSTP1) haplotypes on cell proliferation and apoptosis. *Carcinogenesis* 2007; 28 2268-2273.
- [89] Brockmöller J, Kerb R, Drakoulis N, Staffeldt B, Roots I. Glutathione S-transferase M1 and its variants A and B as host factors of bladder cancer susceptibility: a case-control study. *Cancer Research* 1994; 54 4103-4111.
- [90] Cotton SC, Sharp L, Little J, Brockton N. Glutathione S-transferase polymorphisms and colorectal cancer: a HuGE review. *American Journal of Epidemiology* 2000; 151 7-32.
- [91] Widersten M, Pearson WR, Engström A, Mannervik B. Heterologous expression of the allelic variant mu-class glutathione transferases mu and psi. *Biochemical Journal* 1991; 276 519-524.
- [92] Harada S, Abei M, Tanaka N, Agarwal DP, Goedde HW. Liver glutathione S-transferase polymorphism in Japanese and its pharmacogenetic importance. *Human Genetics* 1987; 75 322-325.
- [93] Huang YS. Genetic polymorphisms of drug-metabolizing enzymes and the susceptibility to antituberculosis drug-induced liver injury. *Expert Opinion on Drug Metabolism & Toxicology* 2007; 3 1-8.
- [94] Leiro V, Fernandez-Villar A, Valverde D, Constenla L, Vazquez R, Pineiro L, González-Quintela A. Influence of glutathione S-transferase M1 and T1 homozygous null mutations on the risk of antituberculosis drug-induced hepatotoxicity in Caucasian population. *Liver International* 2008; 28 835-839.
- [95] Chatterjee S, Lyle N, Mandal A, Kundu S. GSTT1 and GSTM1 gene deletions are not associated with hepatotoxicity caused by antitubercular drugs. *Journal of Clinical Pharmacy and Therapeutics* 2010; 35(4) 465-70.
- [96] Sotsuka T, Sasaki Y, Hirai S, Yamagishi F, Ueno K. Association of isoniazid-metabolizing enzyme genotypes and isoniazid-induced hepatotoxicity in tuberculosis patients. *In Vivo* 2011; 25(5) 803-12.
- [97] Tang SW, Lv XZ, Zhang Y, Wu SS, Yang ZR, Xia YY, Tu DH, Deng PY, Ma Y, Chen DF, Zhan SY. CYP2E1, GSTM1 and GSTT1 genetic polymorphisms and susceptibility to antituberculosis drug-induced hepatotoxicity: a nested case-control study. *Journal of Clinical Pharmacy and Therapeutics* 2012 doi: 10.1111/j.1365-2710.2012.01334.x.

- [98] Cascorbi I. The promises of personalized medicine. *European Journal of Clinical Pharmacology* 2010; 66 749–754.

IntechOpen

IntechOpen