

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Current Advances on Genetic Resistance to Rice Blast Disease

Xueyan Wang, Seonghee Lee, Jichun Wang,
Jianbing Ma, Tracy Bianco and Yulin Jia

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/56824>

1. Introduction

1.1. The historical and contemporary aspects of rice blast disease

Rice (*Oryza sativa* L.) is one of the most important staple foods that feed more than half of the world's population, with Asia and Africa as the largest consuming regions [1]. Blast disease caused by *Magnaporthe oryzae* (Hebert) Barr is one of the most damaging diseases of rice. This disease was first known as rice fever disease in China as early as 1637 [2]. Blast disease was first reported in the United States in 1876, and has been identified in 85 rice-producing countries or regions worldwide (Figure 1).

Blast severely affects lowland rice in temperate and subtropical areas of Asia, and is highly destructive to upland rice in tropical areas of Asia, Latin America, and Africa [3]. Although blast is considered the most destructive rice disease due to the favorable environmental conditions for disease occurrence and worldwide distribution, little information about annual yield losses are available. Table 1 summarizes reported blast outbreaks with annual yield losses from five countries. In China, 40-50% yield losses were observed under severe rice blast infection; in some cases, 100% yield losses were found in severely infected fields [4]. Yield losses of 5-10%, 8%, and 14% were reported in India from 1960 to 1961, Korea from the mid-1970s, and China from 1980 to 1981, respectively [3]. The highest yield losses were recorded in the Philippines; ranging from 50% to 85% in 1963 [3]. It was estimated that 1.6 billion dollars were lost from 1975-1990 due to blast disease worldwide [5]. The estimated annual loss of rice was enough to feed 60 million people for one year [6] (Table 1).



Figure 1. Worldwide distribution of rice blast disease. Red dots show the countries or regions where blast disease has been reported.

Yield loss (%)	Country	Year
5-10	India	1960-61
50-60	Philippines	1963
70-85	Philippines	1969-70
8	Korea	mid-70s
14	China	1980-81
60	Thailand	1982

Source: <http://www.knowledgebank.irri.org/ipm/rice-blast.html>

Table 1. Yield losses due to blast.

1.2. The biology of *M. oryzae*

The most common symptoms in commercial rice fields induced by *M. oryzae* can be found on all the above ground parts of the rice plant at all growth stages. Seeds display brown spots after infection, which may have resulted from the infection of the florets as they mature into seeds. Infected roots have also been observed; however, lesions on the sheaths were relatively rare. Infections on young seedlings are initiated when the conidia are deposited on the surface of the leaves. Water is essential for spores to germinate and attach to the leaf surface [7, 8]. Under optimal conditions, spore germination occurs rapidly and the polarized germ tubes are

formed within hours after landing on the leaf [9]. The secondary cycles are initiated by the spores produced by the lesions on the young seedlings, which can be repeated many times through the growing season. Thousands of spores can be produced from a single lesion in 15 days after infection. Typically blast lesions are diamond shaped (Figure 2A). Initial lesions appear dark green or grey with brown borders; while, older lesions are light tan with necrotic borders. Under favorable conditions, lesions can merge together and rapidly enlarge to several centimeters in length, eventually killing the leaf, and ultimately resulting in plant death. On resistant cultivars, lesions induced by *M. oryzae* usually remain small in size (1-2mm) and appear brown to dark brown in color. Disease severity of rice blast and the amount of spores produced on single lesion depends on temperature, field conditions, relative humidity, fertilization levels, and genotype of rice cultivars. In general, moderate temperatures (~24°C), high relative humidity (90-92%), and high moisture with at least a 12-hour period are advantageous for rice blast. The disease severity of the vegetative phase during the growing season highly influences the amount of disease during the reproductive phase. Spores produced at the end of the growing season may result in collar blast and neck blast; neck blast often causes direct crop loss (Figure 2B).

Existence of physiological races of *M. oryzae* complicates the identification of resistance (*R*) genes. Physiological races of *M. oryzae* were first reported by Sasaki in Japan as early as 1922 [10]. From 1950s to 60s differential rice lines resistant to races of *M. oryzae* were identified in Japan, the United States, India, the Philippines, and South Korea. In 1961, 18 physiological races of *M. oryzae* were identified with 12 differential rice varieties in Japan. During that time, an international differential system using 8 rice varieties was established [11]. In China, the identification of *M. oryzae* races was initiated in late 1970s. Seven rice varieties, Tebo, Zhenlong13, Sifeng43, Dongnong363, Kanto51, Hejiang18, and Lijiangxintuanheigu (LTH), and 43 isolates of *M. oryzae* were used. In 1976, Yamada and his colleagues identified 23 races of *M. oryzae* from 2245 isolates with 9 differential rice varieties. Duan et al [12] used Yuyun 1 (with *Pia*), Gaoliangdao (with *Pii*), Kanto51 (with *Pik*), Chugeng1 (with *Pikm*), Dianyu1 × Fook Kam (with *Piz*), Dali782 (with *Pita*), Dan83-3 (with *Pita2*), and Chengbao1 (with *Pizt*) as differential varieties to characterize races of *M. oryzae* in China. These blast *R* genes are described in more details in the part II of this chapter. Near-isogenic lines (NILs) were chosen to better identify races of *M. oryzae* in a gene-for-gene specific manner. The NILs with *indica* high-susceptible variety CO39 background was developed at the International Rice Research Institute (IRRI), the Philippines [13]. In the United States, Marchetti [14] reported that the races IB-54, ID-13, IG-1, and IH-1 of *M. oryzae* were the most common. Most recently, monogenic lines with 24 major blast *R* genes in BC1 of LTH were developed by scientists at IRRI and Japan [15].

Extensive analysis of rice germplasm with physiological races in the past century reveals that complete genetic resistance (vertical resistance) is conferred by major blast *R* genes named as *Piricularia* genes or *Pi*-genes. These genes are often specific in preventing infections by strains of *M. oryzae* that contain the corresponding avirulence genes; whereas, incomplete resistance (slow-blasting components or horizontal resistance, field resistance, or dilatory resistance) is often conditioned by more than one gene on different chromosomal regions. These genes are




Figure 2. Symptoms of leaf (A) and neck blast diseases (B) in commercial rice fields.

referred to as quantitative resistant loci (QTLs). Resistant germplasm carrying both major and minor *R* genes are extremely important genetic resources that rice breeders can use to improve blast resistance in elite rice varieties.

2. Mapped blast *R* genes

Blast *R* genes are predicted to play important roles in the frontier of rice defense responses. During interactions between rice and blast pathogens, products of the *R* gene can specifically recognize the corresponding elicitors of *M. oryzae*. Since the *Pia* gene, identified in 1967 by Kiyosawa as the first blast *R* gene from the *japonica* variety Aichi Asahi [16], 99 blast *R* genes have been identified; in which 45% were found in *japonica* cultivars, 51% in *indica* cultivars, and the rest 4% in wild rice species (Table 2 to 5). Most deployed *R* genes have often been identified in Asian cultivated rice, specially rice cultivars from Japan and China, with the exception of *Pi9*, *Pi54rh*, *Pi40(t)*, and *Pirf2-1(t)*, which were domesticated from *O. minuta*, *O.*

rhizomatis, *O. australiensis*, and *O. rufipogon*, respectively. All *R* genes have been mapped on all rice chromosomes except for chromosome 3 (Tables 2 to 4; Fig 3). Host genotypes, chromosomal loci, and molecular markers that are tightly linked to blast *R* genes are summarized in Figure 2 and Table 2 (60 major *R* gene) and Table 3 (17 minor *R* gene). Among them, three major *R* gene clusters have been well characterized: the *Piz* locus on Chromosome 6, the *Pik* locus on Chromosome 11, and the *Pita* locus on Chromosome 12 (Figure 3). More detailed information of mapped blast *R* genes can be found at <http://www.ricedata.cn/gene/>, <http://www.shigen.nig.ac.jp/rice/oryzabaseV4/>, and <http://www.gramene.com>.

Chromosome	Major R genes mapped*	Minor R genes mapped*	R gene cloned*	Total*
1	2	2	3	7
2	7	3	1	11
3	0	0	0	0
4	1	2	1	4
5	2	1	0	3
6	12	1	6	19
7	1	0	0	1
8	4	2	1	7
9	3	0	1	3
10	1	1	0	2
11	12	3	8	23
12	15	2	1	18
Total	60	17	22	99

* refers to number of the genes on the chromosome.

Table 2. Summary of blast *R* (major and minor, mapped and cloned) genes on rice chromosomes.

Chr.	Name of R gene	Name of germplasm	Map position (cM) >#	Markers	Name of pathogenic Strains	Ref.
1	<i>Pi27(t)*</i>	Q14	28.4-38.8	RM151, RM259	CHL0335, CHL0888, CHL0918	[17]
1	<i>Pitp(t)</i>	Tetep	114.1	RM246	IC9	[18]
2	<i>Pi14(t)</i>	Maowangu		Amp-1		[19]
2	<i>Pi16(t)</i>	Aus373		Amp-1	Hoko1, Ina72, TH67-22, Ai75-61	[20]
2	<i>PiDa(t)</i>	Dacca6	10.8-14.4	RM211, RM5529		[21]

Chr.	Name of R gene	Name of germplasm	Map position (cM) >#	Markers	Name of pathogenic Strains	Ref.
2	<i>Pid1(t)</i>	Digu	87.5-89.9	RM262	ZB13	[22]
2	<i>Pitq-5</i>	Teqing	150.5-157.5	RG520, RZ446b	IC-17, IB-49, IE-1, IG-1	[23]
2	<i>Pig(t)</i>	Guangchangzhan	142.0-154.1	RM166, RM208	Ken53-33	[24]
2	<i>Piy(t)</i>	Yanxian No.1	153.2-154.1	RM3284, RM208	97-27-2, Zhong10-8-14	[25]
4	<i>Pi39(t)*</i>	Chubu 111	107.4-108.2	RM3743, RM5473		[26]
5	<i>Pi10(t)</i>	Tongil	88.5-102.8	RG13	IB46	[27]
5	<i>Pi23(t)</i>	Suweon 365	59.3-99.5			[28]
6	<i>Pi2-1</i>	Tianjingyeshengdao	Allilic to Pi2/9	AP4791, AP4007	CHL477, CHL473, P06-6, IC-17, 87-4	[29]
6	<i>Pi2-2</i>	Jefferson	58.7	RM19817, AP5659-5	HN318-2, CHL438, KJ201, ROR1, PO6-6	[30]
6	<i>Pi8(t)</i>	Kasalath	74.6-78.2	Amp-3	Race 447.1	[31]
6	<i>Pi13(t)*</i>	Maowangu	74.6-78.2	Amp-3		[19]
6	<i>Pi13(t)*</i>	Kasalath	67.7-68.5	RM2123, RM20155	Ken54-04, 95Mu-29, Ina86-137	[32]
6	<i>Pi22(t)</i>	Suweon 365	38.4-41.9		KJ-201	[28]
6	<i>Pi26(t)*</i>	Gumei 2	51.0-61.6	B10, R674	Ca89	[33]
6	<i>Pi40(t)</i>	IR65482-4-136-2-2 O. <i>australiensis</i>	54.1-61.6	RM527, RM3330	KJ105, Ca89, PO6-6, M101-1-29-1, M64-1-3-9	[34]
6	<i>Pi50(t)</i>	Er-Ba-Zhan	46.8	GDAP51, GDAP16	09-3041a, SC0602, SCRB14, HN0102, W06-18a	[35]
6	<i>Pigm(t)</i>	Gumei 4	65.8	C5483, C0428	CH109 (ZC13), CH147 (ZB25), CH131 (ZA1)	[36]
6	<i>Piz</i>	Zenith	58.7	z4792, z60510, z5765		[37]
6	<i>Pitq-1</i>	Teqing	103.0-124.4	C236, RG653	IC-17, IB-49, IE-1	[23]
7	<i>Pi17(t)</i>	DJ123	94.0-104.0	Est9		[38]
8	<i>Pi42(t)</i>	Zhe733	58.5	RM72	IE1K	[39]
8	<i>Pi33(t)</i>	IR64	45.4	RM72, C483	Guy11	[40]
8	<i>Pi55(t)</i>	Yuejingsimiao 2	99.1-102.1	RM1345, RM3452	CHL688	[41]
8	<i>PiGD-1(t)</i>	Sanhuangzhan 2	53.7	RG1034	GD RFDW-I	[42]
9	<i>Pi3(t)</i>	C104PKT,	31.3-33.0	40N23r	PO6-6	[43]

Chr.	Name of R gene	Name of germplasm	Map position (cM) >#	Markers	Name of pathogenic Strains	Ref.
9	<i>Pi15(t)</i>	GA25	31.3-34.9	CRG3, CRG4	CHL0416 , Hoku 1	[44]
9	<i>Pi56(t)</i>	Sanhuangzhan 2	31.3	RM24022	PO6-6	[42]
10	<i>PiGD-2(t)</i>	Sanhuangzhan 2		R16, R14B	PO6-6	[42]
11	<i>Pi18(t)</i>	Suweon 365	117.9	RZ536	KI-313	[45]
11	<i>Pi38(t)</i>	Tadukan	79.1-88.7	RM206, RM21	B157	[46]
11	<i>Pi44(t)</i>	Moroberekan	91.4-117.9	AF349	C9240-1	[47]
11	<i>PiCO39(t)</i>	CO39	49.1	S2712	6082	[48]
11	<i>Pilm-2</i>	Lemont	56.2-117.9	R4, RZ536	IB54, IG1	[23]
11	<i>Pi7(t)</i>	Moroberekan	71.4-84.3	RG103, RG16	PO6-6	[49]
11	<i>Pi47(t)</i>	Xiangzi 3150	104.2-120.1	RM206, RM224		[50]
11	<i>Pi43(t)</i>	Zhe733	109.5	RM1233	IE1K	[39]
11	<i>Piks</i>	Bengal, M201	115.1-117.3	RM224, RM1233		[51]
11	<i>Pikg(t)</i>	GA20	119.9-120.3			[19]
11	<i>Piy(t)</i>	Yunyin	54	RM202	Sichuang-43	[52]
11	<i>Pizy(t)</i>	Ziyu44	102.9	RM206	ZB13, ZE1	[53]
12	<i>Pi19(t)</i>	Aichi Asahi	50.4-51.5	RM27937, RM1337	CHNO58-3-1, IRBL19-A	[54]
12	<i>Pita-2</i>	Tetep, Pi No.4	50.4			‡
12	<i>Pi6(t)</i>	Apura	12.2-47.9	RG457, RG869		[55]
12	<i>Pi62(t)</i>	Yashiro-mochi	12.2-26.0	RG9, RZ816	4360-R-62	[56]
12	<i>Pi24(t)*</i>	Zhong 156	51.5	RG241A	92-183 (ZC15)	[57]
12	<i>Pi12(t)</i>	Moroberekan	47.9	RG869		[58]
12	<i>Pi20(t)</i>	IR24	51.5-51.8	RM1337, RM5364, RM7102	BN111	[58]
12	<i>PiGD-3</i>	Sanhuangzhan 2	55.8	RM179	GD RFDW-IV	[42]
12	<i>Pi51(t)</i>	Tianjingyeshengdao		RM5364, RM27990	CHL447, RB5, CHL473, PO6-6	[29]
12	<i>Pi39(t)*</i>	Q15	50.4	RM27933, RM27940	CHL724	[59]
12	<i>Pi41(t)</i>	93-11		RM28130	CHL1789, CHL347, CHL688	[60]
12	<i>Pi157(t)</i>	Moroberekan	49.5-62.2	RG341, RG9		[61]

Chr.	Name of R gene	Name of germplasm	Map position (cM) >#	Markers	Name of pathogenic Strains	Ref.
12	<i>Pi48(t)</i>	Xiangzi 3150		RM5364, RM7102		[50]
12	<i>Pitq-6</i>	Teqing	47.9	RG869, RZ397	IC-17, IB-49, IE-1, IB-54	[23]
12	<i>Pih1(t)</i>	Hongjiaozhan	47.9	RG869, RG81	ZB1	[62]

* This R gene shares the same name with another R gene.

The map positions were integrated into IRGSP map according to marker information. Detail information can be found on <http://rgp.dna.affrc.go.jp/E/IRGSP/index.html>.

‡ Information is known, but has not been published.

Table 3. Summary of major blast R genes including their resistance specificity, chromosomal location, map position, and tightly linked DNA markers.

Chr.	Name of R gene	Donor	Map position (cM) >#	DNA Markers	Avirulent race/isolate	Ref.
1	<i>Pi24(t)*</i>	Azucena	64.4	K5	CL6	[63]
1	<i>Pi35(t)</i>	Hokkai 188	132.0-136.6	RM1216, RM1003		[64]
2	<i>Pir2-3</i>	IR64	141.7	RM263, RM250	Race 173	[65]
2	<i>Pi25(t)*</i>	IR64	157.9	RG520	BR26, CH66, CH72	[63]
2	<i>Pirf2-1(t)</i>	<i>O. rufipogon</i>	172.3	RM206, RM266	Race 001	[65]
4	<i>Pikur1</i>	Kuroka	86.0			[66]
4	<i>Pikahei1(t)</i>	Kahei	108.2	RM17496, RM6629		[67]
5	<i>Pi26(t)*</i>	Azucena	22.5-24.7	RG313	PH68	[63]
6	<i>Pi27(t)*</i>	IR64	51.9	Est-2	CH66	[63]
8	<i>Pi11(t)</i>	Zhaiyeqing8	53.2-84.8	BP127A, RZ617	18-2, ZH7-2, Zhong10-2-4,	[68]
8	<i>Pi29(t)</i>	IR64	69	RZ617, RGA-IR86	CL6	[63]
10	<i>Pi28(t)</i>	Azucena	114.7	RZ500	PH68	[63]
11	<i>Pi30(t)</i>	IR64	59.4-60.4	OpZ11-f, RGA-IR14	CH66, CH72	[63]
11	<i>Pi34(t)</i>	Chubu32	79.1-91.4	Z77, Z150-5		[69]
11	<i>Pif</i>	St No. 1	119-120			[70]
12	<i>Pi31(t)</i>	IR64	44.3	O10-800	PH68, CD69	[63]
12	<i>Pi32(t)</i>	IR64	47.5	AF6	BR26	[63]

* This R gene shares the same name with another R gene.

The map positions were integrated into IRGSP map according to marker information. Detail information can be found on <http://rgp.dna.affrc.go.jp/E/IRGSP/index.html>.

Table 4. Summary of minor blast R genes, donors, map position, tightly linked DNA markers, and associated blast races.




Figure 3. Location of cloned and mapped *R* genes on rice chromosomes. The locations of *R* genes have been integrated into IRGSP map according to marker information, then the map was built using Mapmaker software. Centimorgan was used to measure the map positions showing in the right column of the chromosome. The underlined words indicate either SSR or RFLP markers (see additional resources: <http://www.shigen.nig.ac.jp/rice/oryzabaseV4/insd/detail/3554>).

Chr.	Donor and R gene cultivar or line cloned carrying the gene	Map position (cM) ^{>#}	Markers	Locus structure	Protein type	Subcellular localization	FNPs	Expression	Ref.	
1	<i>Pit</i>	K59, Tjahaja	12.2	T256	Multiple	CC-NBS-LRR		Repressible	[71]	
1	<i>Pi37</i>	St. No1	136.1	RM302, RM212	Multiple	NBS-LRR	Cytoplasm V 239 A, I 247 M	Constitutive	[72]	
1	<i>Pish</i>	Shin 2, Norin 22	148.7-154.8		Multiple	CC-NBS-LRR		Constitutive	[73]	
2	<i>Pib</i>	Engkatek, Tohoku IL9, Teqing, Tjinam, BL1	154.1	RM208	Multiple	NBS-LRR		Inducible	[74]	
4	<i>pi21</i>	Owarihatamochi	58.6	P702D03	Multiple	NBS-LRR	Cytoplasm		[75]	
6	<i>Pid2</i>	Digu (I)	65.8		Single	Receptor kinase	Membrane	I 441 M	Constitutive	[76]
6	<i>Pi9</i>	<i>O.minuta</i> , 75-1-127	58.7		Multiple	NBS-LRR		Constitutive	[77]	
6	<i>Pi2</i>	C101A51	58.7		Multiple	NBS-LRR	R 838 S	Constitutive	[78]	
6	<i>Piz-t</i>	TKM, Toride 1	58.7	zt56591	Multiple	NBS-LRR	S 839 R	Constitutive	[79]	
6	<i>Pid3</i>	Digu	65.2-65.8		Single	NBS-LRR	Q 737 Stop		[79]	
6	<i>Pi25*</i>	Gumei 2	63.2-64.6		Multiple	CC-NBS-LRR			[80]	
8	<i>Pi36</i>	Q61, Kasalath	21.6-25.2	CRG3	Single	NBS-LRR	S 590 D	Constitutive	[81]	
9	<i>Pi5</i>	Tetep, RIL 260	31.3-33.0	76B14r, 40N23r	Multiple	CC-NBS-LRR	Cytoplasm	Pi5-1 is inducible, Pi5-2 is constitutive	[82]	
11	<i>Pi1</i>	LAC23, C101LAC	112.1-117.9		Multiple	NBS-LRR			[83]	
11	<i>Pik</i>	To-To, Kusabue, Kanto 51, K60, Chugoku 31, Shin 2-1, K2, K3, , Minehikari, GA 20	119.9-120.3	k8823, k8824, k3951, k39512	Multiple	CC-NBS-LRR		Constitutive	[84], [85]	
11	<i>Pikm</i>	Hokushi,Tsuyuake , IRBLkm-Ts	115.1-117.0	k2167, k6441	Multiple	NBS-LRR		Constitutive	[86]	
11	<i>Pikp</i>	Tetep K60	119.9-120.3	k3957	Multiple	CC-NBS-LRR		Constitutive	[87]	
11	<i>Pikh</i>	Tetep, K3, Kaybonnet, Lemont, Lebonnet	101.9	RM224	Multiple	NBS-LRR		Inducible	[88]	
11	<i>Pi54rh</i>	<i>O. rhizomatis</i>	119.9-120.3		Multiple	CC-NBS-LRR	Extracellular	Inducible	[89]	
11	<i>Pia</i>	Aichi Asahi	36.0	Yca72	Multiple	NBS-LRR			[90]	
11	<i>Pb1</i>	Modan, Tsukinohikari, St NO. 1	85.7-91.4		Single	CC-NBS-LRR		Age- dependent	[91]	
12	<i>Pita</i>	Tetep, Katy, Teqing	50.4		Single	NBS-LRR	Cytoplasm	A 918 S	Constitutive	[92]

^a This R gene shares the same name with another R gene. # The map positions were integrated into IRGSP map according to marker information. Detail information can be found on <http://rgp.dna.affrc.go.jp/E/IR>.

Table 5. Summary of cloned *R* genes, map position, closely linked DNA markers, and their expression.

3. Structure and function of blast *R* genes

Among the mapped *R* genes (Table 3 and 4), 22 genes including 20 major and 2 minor *R* genes (*Pb1* and *pi21*) have been molecularly characterized (Table 5). Noticeably, *Pid2*, *Pid3*, *Pi36*, *Pb1*, and *Pita* are single copy genes; while others are members of small gene families. A total of eight *R* genes have been identified on chromosome 11, with six at the *Pik* locus; six *R* genes on chromosome 6, four of which are at the *Piz* locus. Most cloned blast *R* genes are adequate in providing complete resistance to strains of *M. oryzae* that contain the corresponding avirulence genes. Interestingly, two different members of each of *Pi5*, *Pik*, *Pikp*, *Pikm*, and *Pia* are required for complete resistance to some avirulent races.

Similar to other plant *R* genes, all cloned blast *R* genes to date encode predicted proteins with centrally located nucleotide binding sites (NBS) and leucine rich repeat (LRR) at the carboxyl terminus (Figure 4), with the exception of *Pid2* and *pi21* encoding a B-lectin kinase protein and a proline containing protein, respectively. Plant NBS-LRR proteins can be divided into two subgroups based on whether they contain a Toll-interleukin receptor (TIR)-like domain (TIR-




Figure 4. Structure of all cloned *R* genes. The light green bar represents the length of the *R* genes. The highlighted bars represent the different domains of the *R* genes.

NBS-LRR) or a putative coiled-coil (CC) structure (CC-NBS-LRR) in their amino-terminal region. The rice genome has 500 NBS-LRR gene families, and most of them belong to the CC-NBS-LRR family. The NBS domain contains kinase 1a (p-loop), kinase 2 and 3a (RNBS-B) motif, which presumably bind to ATP and trigger downstream signal transduction; whereas, the LRR is predicted to recognize pathogen effectors, either directly or indirectly. Other noticeable protein domains of plant *R* proteins were also summarized in Figure 4.

The observed structural similarities of blast *R* proteins might imply that their predicted conserved regions are associated with functional roles in triggering resistance to *M. oryzae*. Cloned blast *R* genes can be separated into two clades, I and II (Figure 5). Clade I consists of all NBS-LRR genes and clade II contains both NBS-LRR and non-NBS-LRR gene, such as *Pid2* and *pi21*. Among them, *Pi1-5*, *Pik-1*, *Pikp-1*, and *Pikm1-TS* on chromosome 11 share substantial homology to the *Pi9* locus on Chromosome 6; whereas, *Pi1-6*, *Pik-2*, *Pikp-2*, and *Pikm2-T* are more similar to *Pid2* and *pi21*, which are not NBS-LRR *R* genes. Homologous sequences of blast *R* genes can be found in the diverse germplasm of cultivated species including domesticated landrace varieties and wild relatives of rice. These observations suggest that genetics of rice immunity is ancient and may have been evolved during speciation and domestication.




Figure 5. Phylogenetic tree of all cloned blast *R* genes. The tree was constructed using protein sequences by software Mega 5.0 NJ method.

4. *R* gene-mediated signaling transduction pathways

It is now commonly accepted that products of *R* genes in plants can specifically recognize avirulence genes from the pathogen directly or indirectly to initiate innate immunity system responses. Direct recognition of the putative product of the avirulence gene, *AVR-Pita1* by the *Pita* protein was first reported in 2000 [93]. Over a decade later, in 2012, another blast *R* protein, *Pik*, with similar structure to *Pita*, was found to directly recognize the corresponding avirulence gene *AvrPik* [94]. Direct interactions between other blast *R* and avirulence genes have not been reported; suggesting that indirect interactions may be responsible in triggering effective signal transduction pathways. Other plant genes involved in signal transduction have been investigated by the use of *R* proteins as bait in the yeast two-hybrid system (Y2H). While Y2H is a highly effective tool, it is limited in indentifying immediate plant components of *R* proteins. Molecular basis of blast *R* gene-mediated signaling has been a subject of intensive investigation worldwide. Abundant genes that may be involved in *R* gene-mediated signaling have been identified with DNA microarray [95, 96], and most of them were pathogenicity related genes. Genetic analysis using mutagenesis has been another commonly used alternative to identify downstream components. However, most mutants identified, thus far, are lesion mimic mutants [97]. A major effort to identify *Pita* mediated signaling was accomplished by treating 20,000 Katy seeds with *Pita/Pita²/Pik^s* using fast neutrons, ethyl methyl sulfate (EMS), and gamma irradiation [98]. A total of 142 rice seedlings, with altered disease reactions, were identified from independent M2. The susceptibility of M2 individuals was verified in subsequent generations, and 20 of them were confirmed to be derived from Katy using 20 diagnostic single sequence repeat (SSR) markers. Consequently, the *Ptr(t)* gene in rice was identified to be essential for *Pita* mediated signal transudation [99]. Molecular cloning of *Ptr(t)* will shed light on the interaction mechanism of *Pita* and *Ptr(t)*, and subsequent plant genes involved in defense responses.

5. The management of blast disease-marker assisted selection

Blast disease has been effectively managed by a combination of fungicides and *R* genes integrated into diverse cultural practices. These include seed treatment with fungicide; preventive application of fungicide before heading; crop rotation; balanced application of fertilizers with nitrogen, potassium, and phosphate; and maintaining a sufficient water level during tillering and flowering stages. However, the most effective way to manage rice blast is by the utilization of resistant cultivars due to its environmental and economic sustainability. Incorporating major blast *R* genes have been traditionally accomplished by classical breeding methods and can be accelerated by the use of marker assisted selection (MAS) [100]. MAS has become a practical tool in cultivar improvement by selecting important traits at the early growth stages based on DNA markers, thus breeders can screen for resistance without having to maintain pathogen culture. MAS is efficient and consistent in the field and greenhouse [101]. MAS is also reliable in dealing with traits whose phenotype is affected by the environment. To date, 99 blast *R* genes have been mapped with closely linked DNA markers; and some of

them can be used for MAS. DNA markers were also developed from portions of cloned *R* genes, such as *Pi-ta* and *Pi-b*, for their introduction into elite rice cultivars using MAS. Markers for *Pita*, one of the most important *R* genes for blast in the United States, were developed [102]; while, linked markers for 4 blast *R* genes (*Pik*, *Pib*, *Pita2*, and *Pii*) are effective against eight to ten races of *M. oryzae* were identified [103]. Using MAS, *R* genes like *Pi1*, *Pi5*, *Piz-5*, and *Pita* have been established in different rice genotypes [82, 100, 104, 105]

6. Future prospects

Blast disease is a moving target where the fungus can rapidly adapt to the host. The major difficulty in controlling rice blast is the durability of genetic resistance. Rice cultivars containing only a single *R* gene to a specific pathogen race often become susceptible over time due to the emergence of new virulent races. In theory, *R* genes can be found in rice germplasm in different rice production areas. Stacking *R* genes with overlapped resistance spectra can lead to long lasting resistance. Knowledge of genetic identity of contemporary *M. oryzae* is crucial for precise deployment of rice cultivars with different *R* genes [104]. Effective blast management also requires unprecedented international cooperation. IRRI and research institutions worldwide have been coordinating their resources for both genotyping using next generation of DNA sequencing and phenotyping at different geographic locations. The knowledge gained by this massive collaborative effort ought to lead to more effective methods to reduce crop loss due to blast disease worldwide.

Acknowledgements

We thank the Arkansas Rice Research and Promotion Board, and the US National Science Foundation (Plant Research Program no. 0701745), Natural Science Foundation of China (Program no. 31000847), Zhejiang Natural Science Foundation (Program no. Y3100577), and Qianjiang Talents Project supported by Science Technology Department of Zhejiang Province (Program no. 2011R10038) for their partial financial supports. USDA is an equal opportunity provider and employer.

Author details

Xueyan Wang¹, Seonghee Lee², Jichun Wang³, Jianbing Ma⁴, Tracy Bianco⁵ and Yulin Jia^{5*}

*Address all correspondence to: yulin.jia@ars.usda.gov

¹ China Jiliang University, Hangzhou, China

² Noble Foundation, Oklahoma, USA

3 Jilin Academy of Agricultural Sciences, Changchun, China

4 University of Arkansas Rice Research and Extension Center, Stuttgart, Arkansas, USA

5 USDA ARS Dale Bumpers National Rice Research Center, Stuttgart, Arkansas, USA

References

- [1] United States Department of Agriculture-Economic Research Service. USDA-ERS: Topics, Crops: rice. <http://www.ers.usda.gov/topics/crops/rice/background> (Accessed 20 September 2012)
- [2] Song Y. Tian Gong Kai Wu. China: 1637.
- [3] International Rice Research Institute-Rice Knowledge Bank. IRRI: Crop Health, Disease: Rice Blast. <http://www.knowledgebank.irri.org/ipm/rice-blast/symptoms.html>
- [4] Zhang L, Zhang Q, Cheng Z, Yang H, Zhou Y, Lu C. Resistances of four sets of rice varieties to 19 strains of *Magnaporthe grisea* in Jiangsu area. Jiangsu Journal of Agricultural Science 2010; 26(5) 948-953.
- [5] Baker B, Zambryski P, Staskawicz B, Dinesh-Kumar SP. Signaling in plant-microbe interactions. Science 1997; 276(5313) 726-733.
- [6] Pennisi E. Armed and dangerous. Science 2010; 327 (5867) 804-805.
- [7] Hamer JE, Howard RJ, Chumley FG, Valent B. A mechanism for surface attachment in spores of a plant pathogenic fungus. Science 1988; 239(4837) 288-290.
- [8] Bourett TM, Howard RJ. *In vitro* development of penetration structures in the rice blast fungus *Magnaporthe grisea*. Canadian Journal of Botany 1990; 68(2) 329-432.
- [9] Talbot NJ. On the Trail of a Cereal Killer: Exploring the biology of *Magnaporthe grisea*. Annual Review of Microbiology 2003; 57(1) 177-202.
- [10] Sasaki R. Extensive of strains in rice blast fungus. Journal of plant protection 1992; 9 634-644
- [11] Ling KC, Ou SH. Standardization of the international race numbers of *Pyricularia oryzae*. Phytopathology 1969; 59 339-342.
- [12] Duan Y, Zhu Y, Chen H, Na X. The blast resistance mechanisms and genetic regularity. Journal of Yunnan Agricultural University 1987; 2(2) 11-16.
- [13] Mackill DJ, Bonman JM. Inheritance of blast resistance in near isogenic lines of rice. Phytopathology 1992; 82 (7) 746-749.

- [14] Marchetti MA. Race-specific and rate-reducing resistance to rice blast in US rice cultivars. In: Zeigler RS, Leong SA, Teng PS (eds.) Rice blast disease. Wallingford: Cab International; 1994. pp 231-244.
- [15] Tsunematsu H, Yanoria MJT, Ebron LA, Hayashi N, Ando I, Kato H, Imbe T, Khush GS. Development of Monogenic Lines of Rice for Rice Blast Resistance. *Breeding Science* 2000; 50(3) 229-234.
- [16] Kiyosawa, S.. Genetic studies on host-pathogen relationship in the rice blast disease. In: Ogura T. (ed.) 1967: rice disease and their control by growing resistant varieties and other measures: proceedings of the Symposium of Agriculture, Forestry and Fisheries Research Council, 1967, Tokyo, Japan.
- [17] Zhu M, Wang L, Pan Q. Identification and characterization of a new blast resistance gene located on rice chromosome 1 through linkage and differential analyses. *Phytopathology* 2004; 94(5) 515-519.
- [18] Barman SR, Gowda M, Venu RC, Chattoo BB. Identification of a major blast resistance gene in the rice cultivar 'Tetep'. *Plant Breeding* 2004; 123(3) 300-302.
- [19] Pan Q, Wang L, Ikehashi H, Yamagata H, Tanisaka T. Identification of two new genes conferring resistance to rice blast in the Chinese native cultivar 'Maowangu'. *Plant breeding* 1998; 117(1) 27-31.
- [20] Pan Q, Wang L, Tanisaka T. A new blast resistance gene identified in the Indian native rice cultivar Aus373 through allelism and linkage tests. *Plant Pathology Journal* 1999; 48(2) 288-293.
- [21] Shi B, Zhang J, Zheng Y, Liu Y, Vera Cruz C, Zheng T, Zhao M. Identification of a new resistance gene Pi-Da(t) from Dacca6 against rice blast fungus (*Magnaporthe oryzae*) in Jin23B background. *Molecular Breeding* 2012; 30(2)1089-1096.
- [22] Chen X, Li S, Xu J, Zhai W, Ling Z, Ma B, Wang Y, Wang W, Cao G, Ma Y, Shang J, Zhao X, Zhou K, Zhu L. Identification of two blast resistance genes in a rice variety, Digu. *Journal of Phytopathology* 2004; 152(2) 77-85.
- [23] Tabien RE, Li Z, Paterson AH, Marchetti MA, Stansel JW, Pinson SRM. Mapping of four major rice blast resistance genes from 'Lemont' and 'Teqing' and evaluation of their combinatorial effect for field resistance. *Theoretical and Applied Genetics* 2000; 101(8)1215-1225.
- [24] Zhou J, Wang J, Xu J, Lei C Ling Z. Identification and mapping of a rice blast resistance gene Pi-g(t) in the cultivar Guangchangzhan. *Plant pathology* 2004; 53(2) 191-196.
- [25] Lei C, Huang D, Li W, Wang J, Liu Z, Wang X, Shi K, Cheng Z, Zhang X, Ling Z, Wan J. Molecular mapping of a blast resistance gene in an indica rice cultivar Yanxian No.1. *Rice Genetics Newsletter* 2005; 22 76-77.

- [26] Terashima T, Fukuoka S, Saka N, Kudo S. Mapping of a blast field resistance gene *Pi39(t)* of elite rice strain Chubu 111. *Plant Breeding* 2008; 127(5) 485-489.
- [27] Naqvi NI, Bonman JM, Makill DJ, Nelson RJ, Chattoo BB. Identification of RAPD markers linked to major blast resistance gene in rice. *Molecular Breeding* 1995;1(4) 341-348.
- [28] Ahn SN, Kim YK, Hong HC, Han SS, Choi HC, McCouch SR, Moon HP. Mapping of genes conferring resistance to Korean isolates of rice blast fungus using DNA markers. *Korean Journal of Breeding* 1997; 29(4) 416-423.
- [29] Wang Y, Wang D, Deng X, Liu J, Sun P, Liu Y, Huang H, Jiang N, Kang H, Ning Y, Wang Z, Xiao Y, Liu X, Liu E, Dai L, Wang G. Molecular mapping of the blast resistance genes *Pi2-1* and *Pi51(t)* in the durably resistant rice 'Tianjingyeshengdao'. *Phytopathology* 2012; 102(8) 779-786.
- [30] Jiang N, Li Z, Wu J, Wang Y, Wu L, Wang S, Wang D, Wen T, Liang Y, Sun P, Liu J, Dai L, Wang Z, Wang C, Luo M, Liu X, Wang G. Molecular mapping of the *Pi2/9* allelic gene *Pi2-2* conferring broad-spectrum resistance to *Magnaporthe oryzae* in the rice cultivar Jefferson. *Rice* 2012; 5(29) 1-7.
- [31] Pan Q, Wang L, Ikehashi H, Taniska T. Identification of a new blast resistance gene in the *indica* rice cultivar Kasalath using Japanese differential cultivars and isozyme markers. *Phytopathology* 1996; 86(10) 1071-1076.
- [32] Ebitani T, Hayashi N, Omoteno M, Ozaki H, Yano M, Morikawa M, Fukuta Y. Characterization of *Pi13*, a blast resistance gene that maps to chromosome 6 in *indica* rice (*Oryza sativa* L. variety, Kasalath). *Breeding Science* 2011; 61(3) 251-259.
- [33] Wu J, Fan Y, Li D, Zheng K, Leung H, Zhuang J. Genetic control of rice blast resistance in the durably resistant cultivar Gumei 2 against multiple isolates. *Theoretical and Applied Genetics* 2005; 111(1) 50-56.
- [34] Jeung JU, Kim BR, Cho YC, Han SS, Moon HP, Lee YT, Jena KK. A novel gene, *Pi40(t)*, linked to the DNA markers derived from NBS-LRR motifs confers broad spectrum of blast resistance in rice. *Theoretical and Applied Genetics* 2007; 115 (8) 1163-1177.
- [35] Zhu X, Chen S, Yang J, Zhou S, Zeng L, Han J, Su J, Wang L, Pan Q. The identification of *Pi50(t)*, a new member of the rice blast resistance *Pi2/Pi9* multigene family. *Theoretical and Applied Genetics* 2012; 124(7) 1295-1304.
- [36] Deng Y, Zhu X, Shen Y, He Z. Genetic characterization and fine mapping of the blast resistance locus *Pigm(t)* tightly linked to *Pi2* and *Pi9* in a broad-spectrum resistant Chinese variety. *Theoretical and Applied Genetics* 2006; 113(4) 705-713.
- [37] Hayashi K, Hashimoto N, Daigen M, Ashikawa I. Development of PCR-based SNP markers for rice blast resistance genes at the *Piz* locus. *Theoretical and Applied Genetics* 2004; 108(7) 1212-1220.

- [38] Pan Q, Tanisaka T, Ikehashi H. Studies on the genetics and breeding of blast resistance in rice VI. Gene analysis for the blast resistance of two Yunnan native cultivars GA20 and GA25. *Breeding Science* 1996; 46(Suppl 2) 70.
- [39] Lee S, Wamishe Y, Jia Y, Liu G, Jia M. Identification of two major resistance genes against race IE-1k of *Magnaporthe oryzae* in the *indica* rice cultivar Zhe733. *Molecular Breeding* 2009; 24(2) 127-134.
- [40] Berruyer R, Adreit H, Milazzo J, Gaillard S, Berger A, Dioh W, Lebrun MH, Tharreau D. Identification and fine mapping of Pi33, the rice resistance gene corresponding to the *Magnaporthe grisea* avirulence gene ACE1. *Theoretical and Applied Genetics* 2003; 107(6) 1139-1147.
- [41] He X, Liu X, Wang L, Wang L, Lin F, Cheng Y, Chen Z, Pan Q. Identification of the novel recessive gene *pi55(t)* conferring resistance to *Magnaporthe oryzae*. *Science China Life Sciences* 2012; 55(2) 141-150.
- [42] Liu B, Zhang S, Zhu X, Yang Q, Wu S, Mei M, Mauleon R, Leach J, Mew T, Leung H. Candidate defense genes as predictors of quantitative blast resistance in rice. *Molecular Plant-Microbe Interactions* 2004; 17(10) 1146-1152.
- [43] Jeon J S, Chen D, Yi G, Wang G, Ronald PC. Genetic and physical mapping of *Pi5(t)*, a locus associated with broad-spectrum resistance to rice blast. *Molecular Genetics and Genomics* 2003; 269(2) 280-289.
- [44] Lin F, Liu Y, Wang L, Liu X Pan Q. A high-resolution map of the rice blast resistance gene *Pi15* constructed by sequence-ready markers. *Plant Breeding* 2007; 126(3) 287-290.
- [45] Ahn SH, Kim YK, Hong HC, Han SS, Kwon SJ, Choi HC, Moon HP, McCouch SR. Molecular mapping of a new gene for resistance to rice blast (*Pyricularia grisea* Sacc.). *Euphytica* 2000; 116(1) 17-22.
- [46] Gowda M, Roy-Barman S, Chattoo BB. Molecular mapping of a novel blast resistance gene *Pi38* in rice using SSLP and AFLP markers. *Plant Breeding* 2006; 125(6) 596-599.
- [47] Chen D, dela Viña M, Inukai T, Mackill DJ, Ronald PC, Nelson RJ. Molecular mapping of the blast resistance gene, *Pi44(t)*, in a line derived from a durably resistant rice cultivar. *Theoretical and Applied Genetics* 1999; 98 1046-1053.
- [48] Chauhan RS, Farman ML, Zhang HB, Leong SA. Genetic and physical mapping of a rice blast resistance locus, *Pi-CO39(t)*, that corresponds to the avirulence gene AVR1-CO39 of *Magnaporthe grisea*. *Molecular Genetics and Genomics* 2002; 267(5) 603-612.
- [49] Campbell MA, Chen D, Ronald PC. Development of co-dominant amplified polymorphic sequence markers in rice that flank the *Magnaporthe grisea* resistance gene *Pi7(t)* in recombinant inbred line 29. *Phytopathology* 2004; 94(3) 302-307
- [50] Huang H, Huang L, Feng G, Wang S, Wang Y, Liu J, Jiang N, Yan W, Xu L, Sun P, Li Z, Pan S, Liu X, Xiao Y, Liu E, Dai L, Wang G. Molecular mapping of the new blast

resistance genes *Pi47* and *Pi48* in the durably resistant local rice cultivar Xiangzi 3150. *Phytopathology* 2011; 101(5) 620-626.

- [51] Fjellstrom R, Conaway-Bormans CA, McClung AC, Marchetti MA, Shank A, Park WD. Development of DNA markers suitable for marker assisted selection of three *Pi*-genes conferring resistance to multiple *Pyricularia grisea* pathotypes. *Crop Science* 2004; 44(5) 1790-1798.
- [52] Zhang J, Wang G, Xie H, Ling Z. Genetic analysis and mapping of blast resistance genes in *japonica* rice Yunyin from Yunnan province. *Journal of Agricultural Biotechnology* 2003; 11(3) 241-244.
- [53] Zhang J, Tan Y, Hong R, Fan J, Luo Q, Zeng Q. Genetic analysis and gene mapping of rice blast resistance in *japonica* variety Ziyu 44. *Chinese journal of rice science* 2009; 23(1) 31-35.
- [54] Koide Y, Telebanco-Yanoria MJ, dela Peña FD, Fukuta Y, Kobayashi N. Characterization of rice blast isolates by the differential system and their application for mapping a resistance gene, *Pi19(t)*. *Journal of Phytopathology* 2011; 159(2) 85-93.
- [55] Causse MA, Fulton TM, Cho YG, Ahn SN, Chunwongse J, Wu K, Xiao J, Yu Z, Ronald PC, Harrington SE, Second G, McCouch SR, Tanksley SD. Saturated molecular map of the rice genome based on an interspecific backcross population. *Genetics* 1994; 138(4) 1251-1274.
- [56] Wu KS, Martinez C, Lentini Z, Tohme J, Chumley FG, Scolnik PA, Valent B. Cloning a blast resistance gene by chromosome walking. In: Khush GS. (ed.) *IRRI1996: map-based gene cloning: proceedings of the Third International Rice Genetic Symposium, IRRI, 1996, Manila, Philippines.*
- [57] Zhuang J, Ma W, Wu J, Chai R, Lu J, Fan Y, Jin M, Leung H, Zheng K. Mapping of leaf and neck blast resistance genes with resistance gene analog, RAPD and RFLP in rice. *Euphytica* 2002; 128(3) 363-370
- [58] Inukai T, Zeigler RS, Sarkarung S, Bronson M, Dung LV, Kinoshita T, Nelson RJ. Development of pre-isogenic lines for rice blast-resistance by marker-aided selection from a recombinant inbred population. *Theoretical and Applied Genetics* 1996; 93(4) 560-567.
- [59] Liu X, Yang Q, Lin F, Hua L, Wang C, Wang L, Pan Q. Identification and fine mapping of *Pi39(t)*, a major gene conferring the broad-spectrum resistance to *Magnaporthe oryzae*. *Molecular Genetics and Genomics* 2007; 278(4) 403-410.
- [60] Yang Q, Lin F, Wang L, Pan Q. Identification and mapping of *Pi41*, a major gene conferring resistance to rice blast in the *Oryza sativa* subsp. *indica* reference cultivar, 93-11. *Theoretical and Applied Genetics* 2009; 118(6) 1027-1034.

- [61] Naqvi NI, Chattoo BB. Development of a sequence characterized amplified region (SCAR) based indirect selection method for a dominant blast-resistance gene in rice. *Genome* 1996; 39(1) 26-30.
- [62] Zheng K, Qiang H, Zhuang J, Lu J, Lin H. Mapping of rice resistance genes to blast using DNA marker. *Acta Phytopathologica Sinica* 1995; 25(4) 307-313.
- [63] Sallaud C, Lorieux M, Roumen E, Tharreau D, Berruyer R, Svestasrani P, Garsmeur O, Ghesquiere A, Notteghem JL. Identification of five new blast resistance genes in the highly blast-resistant rice variety IR64 using a QTL mapping strategy. *Theoretical and Applied Genetics* 2003; 106(5) 794-803.
- [64] Nguyen TT, Koizumi S, La TN, Zenbayashi KS, Ashizawa T, Yasuda N, Imazaki I, Miyasaka A. *Pi35(t)*, a new gene conferring partial resistance to leaf blast in the rice cultivar Hokkai 188. *Theoretical and Applied Genetics* 2006; 113(4) 697-704.
- [65] Utami DW, Moeljopawiro S, Aswidinnoor H, Setiawan A, Hanarida I. Blast resistance genes in wild rice *Oryza rufipogon* and rice cultivar IR64. *Indonesian Journal of Agriculture* 2008; 1(2) 71-76
- [66] Goto I. Genetic studies on resistance of rice plant to blast fungus (VII). Blast resistance genes of Kuroka. *Annals of Phytopathological Society of Japan* 1988; 54 460-465.
- [67] Xu X, Chen H, Fujimura T, Kawasaki S. Fine mapping of a strong QTL of field resistance against rice blast, Pikahei-1(t), from upland rice Kahei, utilizing a novel resistance evaluation system in the greenhouse. *Theoretical and Applied Genetics* 2008; 117(6) 997-1008.
- [68] Wang G, Mackill D, Bonman J, McCouch SR, Champoux MC, Nelson RJ. RFLP mapping of genes conferring complete and partial resistance to blast in a durably resistance rice cultivar. *Genetics* 1994; 136(4) 1421-1434.
- [69] Zenbayashi-Sawata K, Fukuoka S, Katagiri S, Fujisawa M, Matsumoto T, Ashizawa T, Koizumi S. Genetic and physical mapping of the partial resistance gene, *pi34*, to blast in rice. *Phytopathology* 2007; 97(5) 598-602.
- [70] Toriyama K. Breeding for resistance to major rice disease in Japan. In: IRRI (ed.) *Rice Breeding*. Manila: IRRI; 1972. p253-281.
- [71] Hayashi K, Yoshida H. Refunctionalization of the ancient rice blast disease resistance gene *Pit* by the recruitment of a retrotransposon as a promoter. *The Plant Journal* 2009; 57(3) 413-425.
- [72] Lin F, Chen S, Que Z, Wang L, Liu X, Pan Q. The blast resistance gene *Pi37* encodes a nucleotide binding site leucine-rich repeat protein and is a member of a resistance gene cluster on rice chromosome 1. *Genetics* 2007; 177(3) 1871-1880.

- [73] Takahashi A, Hayashi N, Miyao A, Hirochika H. Unique features of the rice blast resistance *Pish* locus revealed by large scale retrotransposon-tagging. *BMC Plant Biology* 2010; 10175.
- [74] Wang Z, Yano M, Yamanouchi U, Iwamoto M, Monna L, Hayasaka H, Katayose Y, Sasaki T. The *Pib* gene for rice blast resistance belongs to the nucleotide binding and leucine-rich repeat class of plant disease resistance genes. *The Plant Journal* 1999; 19(1) 55-64.
- [75] Fukuoka S, Saka N, Koga H, Ono K, Shimizu T, Ebana K, Hayashi N, Takahashi A, Hirochika H, Okuno K, Yano M. Loss of function of a proline-containing protein confers durable disease resistance in rice. *Science* 2009; 325(5943) 998-1001.
- [76] Chen X, Shang J, Chen D, Lei C, Zou Y, Zhai W, Liu G, Xu J, Ling Z, Cao G, Ma B, Wang Y, Zhao X, Li S, Zhu L. A B-lectin receptor kinase gene conferring rice blast resistance. *The Plant Journal* 2006; 46(5) 794-804.
- [77] Qu S, Liu G, Zhou B, Bellizzi M, Zeng L, Dai L, Han B, Wang G. The broad-spectrum blast resistance gene *Pi9* encodes a nucleotide-binding site-leucine-rich repeat protein and is a member of a multigene family in rice. *Genetics* 2006; 172(3) 1901-1914.
- [78] Zhou B, Qu S, Liu G, Dolan M, Sakai H, Lu G, Bellizzi M, Wang G. The eight amino-acid differences within three leucine-rich repeats between *Pi2* and *Piz-t* resistance proteins determine the resistance specificity to *Magnaporthe grisea*. *Molecular Plant-Microbe Interactions* 2006; 19(11) 1216-1228.
- [79] Shang J, Tao Y, Chen X, Zou Y, Lei C, Wang J, Li X, Zhao X, Zhang M, Lu Z, Xu J, Cheng Z, Wan J, Zhu L. Identification of a new rice blast resistance gene, *Pid3*, by genome-wide comparison of paired nucleotide-binding site-leucine-rich repeat genes and their pseudogene alleles between the two sequenced rice genomes. *Genetics* 2009; 182(4) 1303-1311.
- [80] Chen J, Shi Y, Liu W, Chai R, Fu Y, Zhuang J, Wu J. A *Pid3* allele from rice cultivar Gumei2 confers resistance to *Magnaporthe oryzae*. *Journal of Genetics and Genomics* 2011; 38(5) 209-216.
- [81] Liu X, Lin F, Wang L, Pan Q. The in silico map-based cloning of *Pi36*, a rice coiled-coil nucleotide-binding site leucine-rich repeat gene that confers race-specific resistance to the blast fungus. *Genetics* 2007; 176(4) 2541-2549.
- [82] Lee SK, Song MY, Seo YS, Kim HK, Ko S, Cao PJ, Suh JP, Yi G, Roh JH, Lee S, An G, Hahn TR, Wang GL, Ronald P, Jeon JS. Rice *Pi5*-mediated resistance to *Magnaporthe oryzae* requires the presence of two coiled-coil-nucleotide-binding-leucine-rich repeat genes. *Genetics* 2009; 181(4) 1627-1638.
- [83] Hua L, Wu J, Chen C, Wu W, He X, Lin F, Wang L, Ashikawa I, Matsumoto T, Wang L, Pan Q. The isolation of *Pi1*, an allele at the *Pik* locus which confers broad spectrum resistance to rice blast. *Theoretical and Applied Genetics* 2012; 125(5) 1047-1055.

- [84] Zhai C, Lin F, Dong Z, He X, Yuan B, Zeng X, Wang L, Pan Q. The isolation and characterization of *Pik*, a rice blast resistance gene which emerged after rice domestication. *New Phytologist* 2011;189(1) 321-334.
- [85] Ashikawa I, Hayashi N, Abe F, Wu J, Matsumoto T. Characterization of the rice blast resistance gene *Pik* cloned from Kanto51. *Molecular Breeding* 2012; 30(1) 485-494.
- [86] Ashikawa I, Hayashi N, Yamane H, Kanamori H, Wu J, Matsumoto T, Ono K, Yano M. Two adjacent nucleotide-binding site-leucine-rich repeat class genes are required to confer *Pikm*-specific rice blast resistance. *Genetics* 2008; 180(4) 2267-2276.
- [87] Yuan B, Zhai C, Wang W, Zeng X, Xu X, Hu H, Lin F, Wang L, Pan Q. The *Pik-p* resistance to *Magnaporthe oryzae* in rice is mediated by a pair of closely linked CC-NBS-LRR genes. *Theoretical and Applied Genetics* 2011; 122(5) 1017-1028.
- [88] Sharma TR, Madhav MS, Singh BK, Shanker P, Jana TK, Dalal V, Pandit A, Singh A, Gaikwad K, Upreti HC, Singh NK. High-resolution mapping, cloning and molecular characterization of the *Pi-k(h)* gene of rice, which confers resistance to *Magnaporthe grisea*. *Molecular Genetics and Genomics* 2005; 274(6) 569-578.
- [89] Das A, Soubam D, Singh PK, Thakur S, Singh NK, Sharma TR. A novel blast resistance gene, *Pi54rh* cloned from wild species of rice, *Oryza rhizomatis* confers broad spectrum resistance to *Magnaporthe oryzae*. *Functional and Integrative Genomics* 2012; 12(2) 215-228.
- [90] Okuyama Y, Kanzaki H, Abe A, Yoshida K, Tamiru M, Saitoh H, Fujibe T, Matsuura H, Shenton M, Galam DC, Undan J, Ito A, Sone T, Terauchi R. A multifaceted genomics approach allows the isolation of the rice *Pia*-blast resistance gene consisting of two adjacent NBS-LRR protein genes. *The Plant Journal* 2011; 66(3) 467-479.
- [91] Hayashi N, Inoue H, Kato T, Funao T, Shirota M, Shimizu T, Kanamori H, Yamane H, Hayano-Saito Y, Matsumoto T, Yano M, Takatsuji H. Durable panicle blast-resistance gene *Pb1* encodes an atypical CC-NBS-LRR protein and was generated by acquiring a promoter through local genome duplication. *The Plant Journal* 2010; 64(3) 498-510.
- [92] Bryan GT, Wu KS, Farrall L, Jia Y, Hershey HP, McAdams SA, Faulk KN, Donaldson GK, Tarchini R, Valent B. tA single amino acid difference distinguishes resistant and susceptible alleles of the rice blast resistance gene *Pi-ta*. *The Plant Cell* 2000; 12(11) 2033-2046.
- [93] Jia Y, McAdams SA, Bryan GT, Hershey HP, Valent B. Direct interaction of resistance gene and avirulence gene products confers rice blast resistance. *EMBO Journal* 2000; 19(15): 4004-4014.
- [94] Kanzaki H, Yoshida K, Saitoh H, Fujisaki K, Hirabuchi A, Alaux L, Fournier E, Tharreau D, Terauchi R. Arms race co-evolution of *Magnaporthe oryzae* AVR-*Pik* and rice *Pik* genes driven by their physical interactions. *The Plant Journal* 2012; 72(6) 894-907.

- [95] Vergne E, Ballini E, Marques S, Sidi Mammar B, Droc G, Gaillard S, Bourot S, DeRose R, Tharreau D, Notteghem JL, Lebrun MH, Morel JB. Early and specific gene expression triggered by rice resistance gene *Pi33* in response to infection by *ACE1* avirulent blast fungus. *New Phytologist* 2007; 174(1) 159-171.
- [96] Gupta SK, Rai AK, Kanwar SS, Chand D, Singh NK, Sharma TR. The single functional blast resistance gene *Pi54* activates a complex defence mechanism in rice. *Journal of Experimental Botany* 2012; 63(2) 757-772.
- [97] Jia Y. Registration of lesion mimic mutant of Katy rice. *Crop Science* 2005; 45(4) 1675.
- [98] Jia Y, Xie J, Rutger JN. Development and characterization of Katy deletion mutant populations. *Plant Mutation Reports* 2006; 1(1) 43-47.
- [99] Jia Y, Martin R. Identification of a new locus, *Ptr(t)*, required for rice blast resistance gene *Pi-ta*-mediated resistance. *Molecular Plant-Microbe Interactions* 2008; 21(4) 396-403.
- [100] Hittalmani S, Parco A, Mew TV, Zeigler RS, Huang N. Fine mapping and DNA marker-assisted pyramiding of the three major genes for blast resistance in rice. *Theoretical and Applied Genetics* 2000; 100(7) 1121-1128.
- [101] Moose SP, Mumm RH. Molecular plant breeding as the foundation for 21st century crop improvement. *Plant Physiology* 2008; 147(3) 969-977.
- [102] Jia, Y, Wang, Z, Singh P. Development of dominant rice blast resistance *Pi-ta* gene markers. *Crop Science* 2002; 42(3) 2145-2149.
- [103] Jia, Y. Understanding the molecular mechanisms of rice blast resistance using rice mutants. In: Shu Q. (ed.) *Induced Plant Mutations in the Genomics Era*. Rome: FAO; 2009. p 375-378.
- [104] Liu S, Li X, Wang C, Li X, He Y. Improvement of resistance to rice blast in Zhenshan 97 by molecular marker-aided selection. *Acta Botanica Sinica* 2003; 45(11) 1346-1350.
- [105] Narayanan NN, Baisakh N, Vera Cruz CM, Gnanamanickam SS, Datta K, Datta SK. Molecular breeding for the development of blast and bacterial blight resistance in rice cv. IR50. *Crop Science* 2002; 42(6) 2072-2079.
- [106] Xing J, Jia Y, Correll JC, Lee FN, Cartwright R, Cao M, Yuan L. Analysis of genetic and molecular identity among field isolates of the rice blast fungus with an international differential system, Rep-PCR, and DNA sequencing. *Plant disease* 2013; 97(4) 491-495.

