

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



---

# **Persistence of *Helicobacter pylori* Infection: Genetic and Epigenetic Diversity**

---

Mohammed Benghezal, Jonathan C. Gauntlett,  
Aleksandra W. Debowski, Alma Fulurija,  
Hans-Olof Nilsson and Barry James Marshall

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/57428>

---

## **1. Introduction**

*Helicobacter pylori* is a Gram negative bacterium found on the luminal surface of the gastric epithelium. Infection is generally acquired during childhood and persists life-long in the absence of antibiotic treatment. *H. pylori* has a long period of co-evolution with humans, going back at least since human migration out of Africa about 60, 000 years ago [1, 2]. This co-evolution is reflected in DNA sequence signatures observed in *H. pylori* strains of different geographic origin and has enabled the mapping of human migration out of Africa. This prolonged and intimate relationship is likely to have shaped the large and diverse repertoire of strategies which *H. pylori* employs to establish robust colonization and persist in the gastric niche. Key challenges that *H. pylori* encounters are fluctuation of acidic pH of the gastric lumen, peristalsis of the mucus layer leading to washout in the lower intestine, nutrient scarcity, and the innate and adaptive immune responses promoting local inflammation or gastritis [3-8]. These challenges, particularly host immune responses, are likely to represent the selective pressure driving *H. pylori* micro-evolution during transmission leading to persistence in the human host.

Host defences against *H. pylori* have been extensively studied including mechanisms which *H. pylori* uses to avoid or inhibit an effective host immune response and review of these related studies is beyond the scope of this chapter (see reviews [9-24]). Instead, key strategies of *H. pylori* immune escape with emphasis on regulation of inflammation are succinctly presented in the context of *H. pylori* persistence. *H. pylori* has evolved to avoid detection by pattern recognition receptors of the innate immune system, such as toll-like receptors and C-type

lectins. Indeed, the TLR4 determinant of *H. pylori* lipopolysaccharide is a very weak stimulus as a result of its altered and highly conserved lipid A structure [25, 26]. In addition, the lipopolysaccharide O-antigen mimics Lewis antigen expressed on host cells and has been shown to regulate dendritic cell function through its binding of DC-SIGN [27-32]. Mutation of the TLR5 recognition site in the flagellin and the sheath protecting the flagella prevent strong activation of the TLR5 signalling pathway [33-35]. *H. pylori* inhibits the adaptive immune response by blocking T-cell proliferation at different levels via at least three different factors, the gamma-glutamyltranspeptidase [36], the cytotoxin VacA [37] and its unique glucosyl cholesterol derivatives [38] (produced from the cholesterol *H. pylori* extracts from host cells). A recent study on the role of the inflammasome during *H. pylori* infection unveiled the pro-inflammatory and regulatory properties of caspase-1 mediated by its substrates IL-1 $\beta$  and IL-18, respectively [39]. In light of the acid-suppressive properties of IL-1 $\beta$  [40], the latter observation exemplifies how seamlessly adapted *H. pylori* is to its human host in its ability to balance gastric pH, inflammation and avoid overt gastric pathology to maintain the physiology of its niche and persist for decades. It would therefore be interesting to note the higher risk for atrophic gastritis in patients with IL-1 $\beta$  polymorphisms that leads to increased expression of IL-1 $\beta$  [41-43] as elevated IL-1 $\beta$  levels might interfere with the dual role of caspase-1 and promote overt inflammation during *H. pylori* chronic infection. Further studies on the activation/regulation of the inflammasome are warranted to gain new insights into gastric cancer caused by *H. pylori* infection.

The scope of this chapter is to review *H. pylori* genetic and epigenetic plasticity and discuss the hypothesis that this plasticity promotes *H. pylori* adaptation to individual human hosts by generating phenotypically diverse populations. Emphasis has been put on mathematical modelling of *H. pylori* chronic infection [44], its micro-evolution and related mechanisms for the generation of diversity including genetic [45-49] and epigenetic diversity [50, 51]. Mechanisms of horizontal gene transfer and the generation of intra-strain genetic diversity are reviewed and the implication of phase-variation-mediated epigenetic diversity is discussed in the context of bacterial population and adaptation.

Examples of experimental strategies to study and decipher *H. pylori* persistence are presented and include bacterial genetics combined with the use of animal models as well as *H. pylori* comparative genomics during chronic and acute infection in humans. The chapter summarises the mechanism of *H. pylori* micro-evolution, in particular the tension between generation of genetic diversity to adapt and genome integrity. Finally, alternatives to antibiotic treatment by targeting *H. pylori* persistence are discussed based on the urease enzyme.

## 2. *H. pylori* persistence: Mathematical modelling

*H. pylori* survive in the gastric niche in a dynamic equilibrium of replication and death by manipulating the host immune system to keep a favourable balance that allows for persistence and transmission. Blaser and Kirschner developed an elegant mathematical model of *H. pylori* persistence based on the Nash equilibrium, specifically that *H. pylori* uses the evolution-

ary stable strategy based on cross-signalling and feedback loop regulations between the host and the bacteria [44]. In this model, a set of interactions between bacteria and the host is defined as well as their corresponding rate parameters. Two populations are considered, the non-replicating free swimming bacteria in the mucus and the adherent bacteria replicating in a nutrient-rich site. This model predicts clearance of the bacteria in the presence of a strong host immunological response and persistence if the host response is weaker. However, this model does not take into account random fluctuations for stochastic phenotype transitions. *H. pylori* is likely to exhibit phenotypic and genetic plasticity to adapt to changing gastric environments but it has relatively few sensors of gastric environment change (e.g. pH, immunological responses, receptor availability, and nutrients). *H. pylori*'s apparently limited gene regulation and its small genome suggest alternative adaptive mechanisms, different from exclusive maintenance of active sensory machinery that is costly. Possibilities include small RNA regulation [52], automatic random genetic switches for generating diverse adaptive phenotypes [53], exemplified by the frameshift-prone repetitive sequences at the beginning of certain phase variable genes [47, 50, 51], and the numerous duplicate and divergent outer membrane genes, which could be part of a more general gene regulation network, so far unidentified. Thus further refinement of this model is required to understand the mechanisms involved in establishing the optimal balance between sensing changes and random phenotype switching. Introducing random fluctuations for stochastic phenotype transitions in this model is highly relevant to phase variation and phasevarion, two mechanisms *H. pylori* uses to generate phenotypic changes and adapt.

### 3. Genetic diversity

The above mentioned mathematical model based on cross-signalling and feedback loop regulation between the host and the bacteria predicts a unique *H. pylori* population in every human host. In other words, *H. pylori* transmission results in adaptation to a specific host during the acute phase of transmission as well as in the chronic phase. The Nash equilibrium model for *H. pylori* colonization is in line with the genetic diversity of *H. pylori* populations as the result of human migration out of Africa and with vertical transmission. Indeed, *H. pylori* strains transmitted within families are genetically less diverse than strains from unrelated infected persons. This highlights the isolation of *H. pylori* strains within a host and genetic adaptation to human subpopulations. Multi-locus sequence typing analysis has identified 6 ancestral populations of *H. pylori* named ancestral European 1, ancestral European 2, ancestral East Asia, ancestral Africa1, ancestral Africa2 [2], and ancestral Sahul [1].

#### 3.1. Intra-strain generation of genetic diversity

Adaptive evolution of species relies on a balance between genetic diversity and genome stability promoted by genome maintenance mechanisms and DNA repair preventing mutations and ensuring cell viability. Intra-strain or intracellular genetic changes have several origins including spontaneous chemical instability of DNA such as depurination and deamination, errors during DNA replication and the action of DNA damaging metabolites, either

endogenous or exogenous. The DNA repair machinery is essential to all living organisms and has been best studied for the model organism *Escherichia coli*. The advances in DNA sequencing technologies and comparative genomics provided a unique opportunity to better understand genome maintenance beyond *E. coli* model organism by comparing the DNA repair gene content in different bacterial species. This is of particular interest for bacterial pathogens that have to overcome immune responses and associated DNA damaging oxidative stress [54]. Comparative genomics of nine human pathogens (*Helicobacter pylori*, *Campylobacter jejuni*, *Haemophilus influenza*, *Mycobacterium tuberculosis*, *Neisseria gonorrhoea*, *Neisseria meningitidis*, *Staphylococcus aureus*, *Streptococcus pneumoniae* and *Streptococcus pyogenes*) revealed a reduced number of genes in DNA repair, recombination and replication compared to *E. coli* [54].

During replication DNA polymerase encountering DNA damage could either be blocked or continue and introduce a mutation into the daughter strand. Maintenance of the template for DNA replication before the replication fork reaches the DNA lesion is therefore an effective DNA repair strategy employed by the cell to avoid mutation or replication arrest. A blocked replication fork requires the homologous recombination machinery to repair the damaged DNA and to resume replication. DNA template maintenance is achieved through several mechanisms pre- and post-replication:

- Direct repair that reverses base damage.
- Excision repair that removes the lesion from the DNA duplex. There are three types of excision repair:
  - Base excision repair (BER) – PolI dependent [54].
  - Nucleotide excision repair (NER) – PolI dependent [54].
  - Alternative excision repair (AER) has been described in a limited number of organisms such as *Schizosaccharomyces pombe* and *Deinococcus radiodurans* – Endonuclease and DNA ligase dependent [55].
- Mismatch repair (MMR) is a post-replication mechanism which contributes to the DNA polymerase fidelity by identifying mismatched bases and removing them from the daughter strand [54].
- Recombinational repair that exchanges the isologous strands between the sister DNA molecules.

Table 1 shows that nucleotide excision repair is the only fully conserved repair pathway amongst the nine pathogens mentioned above and that the SOS response related genes are completely missing from *H. pylori* [46, 54]. Direct repair and mismatch repair are often completely absent whereas base excision repair, recombinational repair and replication (*dnaA*, *dnaB*, *dnaG*, *gyrA*, *gyrB*, *parC*, *parE*, *priA*, *rep*, *topA* and *polA*) are often missing one or several genes. This absence of DNA repair and replication genes suggests either that functional homologs remain to be discovered or that specific genome dynamics and genome integrity maintenance strategies are at play in different microbial pathogens to adapt to their niche.

Pathway	Protein	<i>H. pylori</i> gene	Protein function	Bacterial species	
				Ec	Hp
<b>Direct repair</b>					
	Ada		Methyltransferase	+	-
	AlkB		Oxidative demethylase	+	-
	Ogt	HP0676	Methyltransferase	+	+
	Phr/Spl		Photolyase	+	-
<b>Base excision repair</b>					
	MutY	HP0142	Glycosylase (adenine)	+	+
	MutM		Glycosylase (8-oxoG)	+	-
	Nei		Endonuclease VIII	+	-
	Nth	HP0585	Endonuclease III	+	+
	Tag		Glycosylase I (adenine)	+	-
	AlkA		Glycosylase II (adenine)	+	-
	Ung	HP1347	Glycosylase (uracil)	+	+
	Xth	HP1526	Exonuclease III	+	+
	Mpg		Glycosylase (purine)	+	-
	YgjF		Glycosylase (thymine)	+	-
	Nfo		Endonuclease IV	+	-
	MagIII	HP0602	Glycosylase ( adenine)	-	+
<b>Nucleotide excision repair</b>					
	UvrA	HP0705	DNA damage recognition	+	+
	UvrB	HP1114	Exinuclease	+	+
	UvrC	HP0821	Exinuclease	+	+
	UvrD	HP1478	Helicase II	+	+
	Mfd	HP1541	Transcription-repair coupling factor	+	+
<b>Mismatch excision repair</b>					
Mismatch recognition	MutS1		Mismatch recognition	+	-
	MutS2	HP0621	Repair of oxidative DNA damage	-	+
	MutL		Recruitment of MutS1	+	-
	MutH		Endonuclease	+	-



Pathway	Protein	<i>H. pylori</i> gene	Protein function	Bacterial species	
				Ec	Hp
<b>Recombinational repair</b>					
	RecA	HP0153	DNA strand exchange and recombination	+	+
RecBCD pathway	RecB (AddA)	HP1553	Exonuclease V, β subunit	+	+
	RecC (AddB)	HP0275	Exonuclease V, γ subunit	+	+
	RecD		Exonuclease V, α subunit	+	-
RecFOR pathway	RecF		Gap repair protein	+	-
	RecJ	HP0348	5'→3' ssDNA exonuclease	+	+
	RecO	HP0951	Gap repair protein	+	+
	RecR	HP0925	Gap repair protein	+	+
	RecN	HP1393	ATP binding	+	+
	RecQ		3'-5'DNA helicase	+	-
	Branch migration	RuvA	HP0883	Binds junctions; helicase (with RuvB)	+
RuvB		HP1059	5'-3'junction helicase (with RuvA)	+	+
RecG		HP1523	Resolvase, 3'-5'junction helicase	+	+
Resolvase	RuvC	HP0877	Junction endonuclease	+	+
<b>Chromosome dimer resolution</b>					
	XerC		Recombinase	+	-
	XerD		Recombinase	+	-
	XerH	HP0675	Recombinase	-	+

Adapted from References Kang and Blaser, Nat Rev Microbiol. 2006; 4(11):826-36 and Ambur et al., FEMS Microbiol.Rev 2009; 33:453-470. AP, apurinic/aprymidinic; ds, double stranded; ss, single stranded.

Adapted from References Kang and Blaser, Nat Rev Microbiol. 2006; 4(11):826-36 and Ambur et al., FEMS Microbiol.Rev. 2009; 33:453-470. AP, apurinic/aprimidinic; ds, double stranded; ss, single stranded.

**Table 1.** Comparative analysis of DNA repair and recombination pathways in *E. coli* and *H. pylori*

*H. pylori* specific DNA repair and replication pathways and their potential role in colonization, virulence and persistence are discussed below based on experimental evidence.

### 3.1.1. DNA repair and mutagenesis

The most striking feature of *H. pylori* DNA repair gene content is the absence of the mismatch repair. A distant homolog of *mutS* was identified [56, 57] and phylogenetic analysis revealed

that MutS belongs to the MutS2 subfamily of proteins [58] that are not associated with MMR. Functional analysis of *H. pylori* MutS2 identified a role of this protein in repair of oxidative DNA damage and Muts2 is required for robust colonization in the mouse model of *H. pylori* infection [59]. Deficiency in MMR activity leads to an increase in mutation rate and is known as the mutator phenotype in Enterobacteriaceae and *Pseudomonas aeruginosa* [60, 61]. The apparent lack of MMR is in line with *H. pylori* mutation rate that is about 2 orders of magnitude higher than in *E. coli* [45]. *H. pylori* mutator phenotype could confer genetic diversity and a selective advantage to adapt and persist in the changing gastric niche. Alternatively, the mutator phenotype of *H. pylori* might promote transmission as postulated for *Neisseria meningitidis* based on the observation of high prevalence of mutations in MMR genes in a *N. meningitidis* epidemic [62].

Numerous reports have confirmed *H. pylori* dependence on DNA repair to establish robust colonization and to persist, suggesting that the human gastric niche induces bacterial DNA lesions [63].

Four of the base excision repair proteins only (MutY, Nth, Ung and Xth) are present in *H. pylori* [54, 64-67] in addition to a novel 3-methyladenine DNA glycosylase (MagIII) that defines a new class within the endonuclease III family of base excision repair glycosylases resembling the Tag protein [68, 69]. *magIII* and *xth* mutants were identified in a signature-tagged mutagenesis screen based on the mouse model of *H. pylori* infection suggesting a role during colonization [70]. Deletion mutants *mutY*, *ung* and *xth* exhibited higher spontaneous mutation frequencies compared to wild-type, with a *mutY* mutant displaying the highest frequency of spontaneous mutation. *mutY* mutants colonized the stomach of mice less robustly compared to wild-type, demonstrating a role for MutY in base excision repair *in vivo* to correct oxidative DNA damage [64]. The presence of an adenine homopolymeric tract in *mutY* suggests that MutY phase varies. This raises an interesting question whether *H. pylori* can vary its mutation rate to adapt to its gastric niche, and highlights the tension between mutation and repair. Deletion of the *nth* gene also led to hypersensitivity to oxidative stress, reduced survival in macrophages and an increased mutation rate compared to wild-type [71]. The *nth* mutant also colonized the mouse stomach poorly 15 days post challenge and was almost cleared after 60 days [71].

Mutants in nucleotide excision repair genes *uvrA*, *uvrB*, *uvrC* and *uvrD* have been constructed in *H. pylori* [49, 72, 73] and their UV sensitivity phenotype confirmed their role in DNA repair. Although surprisingly *uvrA* and *uvrB* mutants had lower mutation rate and recombination frequencies [49]. This phenomenon can be explained by nucleotide exchange of undamaged DNA and was hypothesized to be another mechanism *H. pylori* uses to generate genetic diversity [49]. Furthermore, *uvrC* mutation led to an increase in the length of DNA import, suggesting that NER influences homologous recombination. UvrD limited homologous recombination between strains [49, 73]. A mutant deficient in Mfd, the transcription repair coupling factor, was found to be more sensitive to DNA damaging agents [74], suggesting that *H. pylori* may also detect blocked RNA polymerase as a damage recognition signal in addition to the DNA distortion recognition properties of UvrA and UvrB. In summary, NER has opposite dual functions; maintenance of genome integrity by excision repair versus generation



of genetic diversity by increasing the spontaneous mutation rate and controlling the rate of homologous recombination and corresponding import length of DNA. Full conservation of the NER pathway in *H. pylori* contrasts with other lacunar DNA repair pathways and highlights the importance of the dual role of NER for *H. pylori* during its replication cycle to balance genetic diversity and genome integrity. To date, the role of NER in genetic diversification has not been tested *in vivo*. Only the *mfd* mutant was identified in a signature-tagged mutagenesis screen based on the mouse model of *H. pylori* infection, suggesting a role of NER during colonization [70].


Finally, recombinant *H. pylori* overexpressing DNA polymerase I displays a mutator phenotype suggesting a role of replication in generating genetic diversity. Bacterial DNA polymerase I participates in both DNA replication and DNA repair. *H. pylori* DNA PolI lacks a proofreading activity, elongates mismatched primers and performs mutagenic translesion synthesis. Conversely, the DNA polymerase I deficient mutant exhibited lower mutation frequency compared to wild-type.

### 3.1.2. DNA recombination

#### 3.1.2.1. Homologous recombination

Recombination between similar sequences is called homologous recombination (HR). HR participates in DNA repair of double strand breaks and stalled replication forks. It is dependent on RecA, a protein that binds and exchanges single stranded DNA. As depicted in Figure 1, HR is a three-step process involving presynapsis, synapsis and postsynapsis. The presynapsis pathway is dictated by the nature of the DNA substrate. Two categories of proteins prepare the single stranded DNA for binding by RecA. A linear DNA duplex with a double-strand end (that could arise during partial replication of incoming single stranded DNA during conjugation, transduction, or DNA damage) is processed by RecBCD. Gapped DNA (that may form during replication) is processed to single-stranded DNA by RecQ and RecJ, whereas RecFOR inhibits RecQ and RecJ activities to allow RecA binding. The result is a nucleoprotein filament that is ready for the search of homologous sequence in the DNA duplex and RecA-mediated strand exchange once that homologous sequence is found. This synapsis step leads to the formation of a structure termed the D-loop. Postsynapsis involves D-loop branch migration and Holliday junction formation catalysed by RuvAB prior to resolution by RuvC or RusA. RecG has also been shown to be involved in recombination and to catalyse branch migration, in addition to its role in replication fork reversal. Interestingly, RuvC-mediated Holliday junction resolution is biased towards non-crossover, avoiding the formation of a chromosome dimer that requires the *Xer/dif* machinery for resolution.

*H. pylori* expresses most of the HR proteins of *E. coli* including; RecA, AddAB instead of RecBCD, RecOR (lacking RecF and RecQ), RuvABC (lacking RusA), RecG, and *XerH/dif<sub>H</sub>* for chromosome dimer resolution. The presence of most HR genes in *H. pylori* suggests that HR plays an important role in *H. pylori* gastric colonization. Intragenomic recombination in families of genes encoding outer membrane proteins leads to *H. pylori* cell surface remodelling to adapt to the human host by adjusting bacterial adhesive properties, antigen



**Figure 1.** Homologous recombination. Two DNA substrates can be processed by the HR machinery **a)** double strand break **b)** gapped DNA. Three stages of HR are presented starting with presynapsis (DNA processing to ssDNA for RecA loading), synapsis (search of the homologous sequence in the DNA duplex and RecA-mediated strand exchange leading to the formation of a structure termed D-loop) and postsynapsis (D-loop branch migration and Holliday junction formation catalysed by RuvAB before resolution by RuvC or RusA). Proteins involved at the different steps are indicated in black for the model organism *E. coli* and in blue for *H. pylori*.

mimicry [75, 76] and modulation of the immune system [76]. HR was suggested to be the underlying recombination mechanism for *homa/homB* and *galT/Jhp0562* allelic diversity [77, 78], whereas gene conversion (non-reciprocal recombination) is responsible for *sabA* diversity [79]. Mutants in *recA*, *addA* or *recG* had lower rates of *sabA* adhesin gene conversion suggesting that RecA-independent gene conversion exists and that this recombination may be initiated by a double-strand break [79].

The RecA deficient mutants are sensitive to DNA damaging agents such as UV light, methyl methanesulfonate, ciprofloxacin, and metronidazole [80, 81]. RecA was the first HR protein to be characterized in *H. pylori* and it was found not to complement an *E. coli* RecA deficient mutant [80]. Lack of cross-species complementation was first attributed to the putative post-translational modification of RecA [80], however, studies showed that the lack of complementation was due to species specific interaction of RecA with proteins involved in presynapsis such as RecA loading on the single stranded DNA by AddAB [82]. RecA's role *in vivo* is supported by poor colonization of a RecA deficient mutant [82]. Interestingly, RecA was shown to integrate the transcriptional up-regulation of DNA damage responsive genes (upon DNA uptake) and natural competence genes (upon DNA damage) in a positive feedback loop. The

interconnection of natural competence and DNA damage through RecA highlights the role of HR in persistence and in generating genetic diversity. Alternatively, and not exclusive to a role in generation of genetic diversity, RecA-mediated genetic exchange might represent a mechanism for genome integrity maintenance in an extreme DNA-damaging environment.

Several gene deletion studies have shown that *H. pylori* has two separate and non-overlapping presynaptic pathways, AddAB and RecOR, contrasting with the redundancy of RecBCD and RecFOR in *E. coli* [83, 84]. The single *addA* mutant and double mutant *addA recO* exhibit similar sensitivity to double strand break inducing agents, suggesting that AddAB is involved in the double strand break repair pathway, and RecOR in gap repair. Finally, RecOR is involved in intragenomic recombination and AddAB in intergenomic recombination. Both pathways are required *in vivo* for robust colonization and persistence based on the lower colonization loads of single *addA*, *recO*, and *recR* mutants in the mouse model of *H. pylori* infection with the double *addA recO* mutant displaying the lowest bacterial load. As expected for recombinational repair proteins, RecN mediated DNA double strand break recognition and initiation of DNA recombination is also required *in vivo* for robust colonization [85].

Resolution of the Holliday junction formed by the action of RecA is performed by RuvABC in *H. pylori* and *recA*, *ruvB*, or *ruvC* mutants exhibited similar UV sensitivities [86, 87]. Colonization of the recombinational repair mutant, *ruvC*, was affected and 35 days post-infection the *ruvC* mutant was cleared by mice. Thus, although dispensable for the initial colonization step, recombinational DNA repair and HR are essential to *H. pylori* persistent infection. Furthermore, the *ruvC* deletion mutant elicited a Th1 biased immune response compared to a Th2 biased response observed for wild-type, highlighting the role of homologous recombination in *H. pylori* immune modulation and persistence [88].

Unexpectedly, the RecG homolog in *H. pylori* limits recombinational repair [86] by competing with the helicase RuvB. Mutation of RecG increased recombination frequencies in line with a role of RecG in generating genetic diversity. The term 'DNA antirepair' was coined to highlight the tension between the generation of diversity and genome integrity maintenance for *H. pylori* adaptation to its niche. Further regulation of homologous recombination is mediated by the MutS2 protein that displays high affinity for DNA structures such as recombination intermediates thus inhibiting DNA strand exchange and consequent recombination [89]. MutS2 deficient cells have a 5-fold increase in recombination rate [90].

### 3.1.2.2. Non-homologous recombination

XerH/*dif<sub>H</sub>* machinery for chromosome dimer resolution was found to be essential for *H. pylori* colonization [87]. Deletion of *xerH* in *H. pylori* caused: (i) a slight growth defect in liquid culture, as is typical of *xer* mutants of *E. coli* [91], (ii) markedly increased sensitivity to DNA breakage inducing and homologous recombination stimulating UV irradiation and ciprofloxacin, (iii) increased UV sensitivity of a *recG* mutant [86], and (iv) a defect in chromosome segregation. The inability of the *xerH* mutant to survive in the gastric niche contrasts with *ruvC* mutant colonization and further supports the idea that XerH is not involved in DNA repair but in chromosome maintenance such as chromosome dimer resolution, regulation of replication and possibly in chromosome unlinking. This, in turn, suggests that slow growing *H. pylori* depends

on unique chromosome replication and maintenance machinery to thrive in their special gastric niche.

Rearrangement of the middle region of the *cagY* gene, independent of RecA [92], leads to in frame insertion or deletion of CagY and gain or loss of function of the CagA type IV secretion system. Recombination of *cagY* was proposed to be a mechanism to regulate the inflammatory response to adapt and persist in the gastric niche [93]. To date the exact recombination mechanism involving direct repeats in the middle region of *cagY* remains unknown.


### 3.1.3. Phase variation

Host adapted human pathogens, such as *H. influenzae*, *Neisseria* species and *H. pylori*, have evolved genetic strategies to generate extensive phenotypic variation by regulating the expression of surface bound (or secreted) protein antigens that directly (or indirectly) interact with host cells. Phenotypic variation of the bacterial external composition will alter the appearance of the bacterium as sensed by the host immune system. One common regulatory mechanism to achieve antigen diversity within a bacterial population is known as phase variation [94, 95].

In pathogens, simple sequence repeats (SSRs) are tandem iterations of a single nucleotide or short oligonucleotides that, with respect to their length, are hypermutable (Figure 2). Reversible slipped strand mutation/mispairing of SSRs within protein coding regions cause frame shifts, resulting in the translation of proteins that vary between being in-frame (on), producing functional full-length proteins, and out-of-frame (off), where a truncated or non-sense polypeptide is produced [96]. Additionally, SSRs may occur in the promoter region of genes where variation in their length may affect promoter strength by mechanisms such as alteration of the distance between -10 and -35 elements.

In *H. pylori*, phase variation regulates the expression of genes that are likely to be important for adaptation in response to environmental changes and for immune evasion in order to establish persistent colonization of the host. Analysis of DNA sequence motifs based on annotated genomes of *H. pylori* strains 26695 [57] and J99 [97] revealed substantial occurrence, intra- and intergenic, of homopolymeric tracts and dinucleotide repeats. Certain categories of genes (or their promoter region) were particularly prone to contain SSRs, such as those coding for LPS biosynthesis enzymes, outer membrane proteins and DNA restriction/modification systems, and thus have been identified as possible candidates regulated by phase variation [98].

Further genome analysis of *H. pylori* strains 26695 and J99 demonstrated an expanded repertoire of candidate phase-variable genes. In addition to previous sequence motif analyses of the annotated genomes by Tomb and Alm, 13 novel putative phase-variable antigens were identified *in silico* [99]. Poly-A and poly-T repeats were almost exclusively found in intergenic regions whereas poly-C and poly-G repeats were mostly intragenic. Five classes of gene function were described; i) LPS biosynthesis (seven genes), ii) cell surface associated proteins (22 genes), iii) DNA restriction/modification systems (nine genes), iv) metabolic or other proteins (three genes), and v) hypothetical ORFs with unidentified homology (five genes). This



**Figure 2.** Phase variation and generation of epigenetic diversity by phasevarions. Slipped strand mispairing of short sequence repeats during DNA replication results in alteration of repeat length in descendants. Figure adapted from Thomson *et al.* [316]. Repeat sequences, represented by a white box, may be present in gene promoters or within the coding sequence of a gene where variation in repeat length alters gene expression, or changes the reading frame leading to the generation of a premature stop codon, respectively. This leads to phenotypic variation due to the presence or absence of the encoded protein. In instances where the phase varying gene encodes a methyltransferase, phase variable expression of the methylase results in either the methylation, or absence of methylation, of target DNA sequences of the enzyme. In instances where methylation affects gene promoter activity this results in varied gene expression. A set of genes, referred to as a phasevarion, may be regulated in this manner resulting in rapid, reversible epigenetic generation of phenotypic diversity.



analysis highlights the importance for a bacterial species such as *H. pylori* to be able to regulate cell surface antigen expression that is responsible for direct interaction with a changing environment.

Multi locus sequence typing (MLST) analysis was then applied for analysis of sequence motif variation in 23 *H. pylori* strains selected on the basis of ethnicity and country of origin (Table 2). Four strain types were investigated, i) hpEastAsia, ii) hpLadakh, iii) hpEurope, and iv) hpAfrica1 and hpAfrica2. In conclusion, approximately 30 genes have been identified as likely phase-variable and it has been postulated that a much higher degree of recombination occurs for genes under constant selective pressure as opposed to more neutral genes such as those encoding 'housekeeping' functions [99]. DNA sequence analysis of *H. pylori* strains indicated that recombination of LPS biosynthesis genes may reflect genetic exchange within the population lineage and that phase variable gene evolution occurs at a high rate [100].

Gene function	CDS		Repeat	Strain variation <sup>1</sup>		
	26695 (HP)	J99 (jhp)		+ <sup>2</sup>	- <sup>3</sup>	Abs <sup>4</sup>
<b>LPS biosynthesis</b>						
$\alpha$ -1, 3-fucosyltransferase*	0651	0596	A	7	16	0
			C	10	13	0
$\alpha$ -1, 3-fucosyltransferase*	0379	1002	A	12	11	0
			C	14	9	0
FutC ( $\alpha$ -1, 2-fucosyltransferase)*	0093-4	0086	C	23	0	0
			A	19	4	0
Lex2B	0619	0563	C	20	0	3
RfaJ ( $\alpha$ -1, 2-glycosyltransferase)	0208	0194	GA	20	0	3
RfaJ ( $\alpha$ -1, 2-glycosyltransferase)	NH <sup>5</sup>	0820	C	8	11	4
$\beta$ -1, 4- <i>N</i> -acetylgalactoamyl transferase*	0217	0203	G	21	2	0
			G	3	20	0
RfaJ homologue	0159	-	-	Putative <sup>6</sup>		
RfaJ homologue	1416	-	-	Putative		
<b>OMP<sup>7</sup></b>						
FliP (flagellar protein)	0684-5	0625	C	6	16	1
OM adherence protein	1417	1312	GA	14	0	9
Streptococcal M protein	0058	0050	C	12	2	9
HopH (OipA)	0638	0581	CT	16	7	0
TlpB (chemotaxis protein)	0103	0095	G	20	0	3
BabA (Leb binding protein)	1243	0833	CT	Putative		



Gene function	CDS		Repeat	Strain variation <sup>1</sup>		
	26695 (HP)	J99 (jhp)		+ <sup>2</sup>	- <sup>3</sup>	Abs <sup>4</sup>
BabB (Leb binding protein homologue)	0896	1164	TC	22	0	1
SabA (sialic acid binding adhesion)	0725	0662	TC	17	2	4
SabB (sialic acid binding adhesion)	0722	0659	TC	7	16	0
HopZ (adhesion)	0009	0007	TC	22	0	1
Oxaluglutarate	0143	0131	A	23	0	0
PldA (phospholipase A)	0499	0451	G	20	3	0
HcpA (cysteine rich protein)	0211	0197	-	Putative		
HcpB (cysteine rich protein)	0335	0318	G	13	0	10
<b>DNA Restriction/modification</b>						
Methyltransferase*	1353-4	1272	G	15	0	8
			G	15	0	8
Type III restriction enzyme	1369-70	1284	G	18	3	2
Type II R/M enzyme	1471	1364	G	14	1	8
Type III R/M enzyme	1522	1411	G	6	0	17
HsdR (type I restriction enzyme)*	0464	0416	C	23	0	0
			A	2	21	0
MbolIR (type II restriction enzyme)	1366	1422	A	6	12	5
<b>Metabolic proteins</b>						
FrxA (NADPH flavin oxidoreductase)	0642	0586	G	6	15	2
<b>Hypothetical ORFs</b>						
Hypothetical protein	0744	0681	AG	18	0	5
Hypothetical protein	1433	1326	C	18	2	3
Hypothetical protein	0767	NA <sup>8</sup>	G	7	0	16
Hypothetical protein	NH	0540	A	20	0	3
<sup>1</sup> Phase variation investigated for 30 genes in 23 <i>H. pylori</i> strains. Adapted from Salaün et al [99, 122]						
<sup>2</sup> Gene present with repeat						
<sup>3</sup> Gene present with repeat absent or stabilized						
<sup>4</sup> Gene absent						
<sup>5</sup> No homolog in genome						
<sup>6</sup> Putative phase variable gene						
<sup>7</sup> Outer membrane protein						
<sup>8</sup> Not annotated						
*‘on’-‘off’ gene status directed by more than one repeat						

**Table 2.** Phase variation in *H. pylori* strains from different geographic regions

### 3.1.3.1. Lipopolysaccharide (LPS) biosynthesis

All Gram-negative bacterial outer membranes contain a structurally important component called LPS (or endotoxin). *H. pylori* LPS consists of three major moieties; a lipid A membrane anchor, a core- and an O-polysaccharide antigen. Although structurally similar to many other Gram-negative bacteria, *H. pylori* LPS has low immunological activity [101]. The O-polysaccharide chain of the LPS of most *H. pylori* strains contains carbohydrates that are structurally related to human blood group antigens, such as Lewis a, b, x and y. The structural oligosaccharide pattern of the LPS of some pathogenic bacteria, including *H. pylori*, is regulated by phase variable fucosyl- and glycosyltransferases; enzymes that transfer sugar residues to its acceptor.

*H. pylori* strain NCTC 11637 (ATCC 43504, CCUG 17874) expresses the human blood group antigen Lewis x ( $\text{Le}^x$ ) in a polymeric form ( $\text{Le}^x$ )<sub>n</sub> on its core antigen. However,  $\text{Le}^x$  expression is not stable and can lead to different LPS variants in single cell populations. Loss of  $\alpha$ 1, 3-linked fucose resulted in a non-fucosylated (lactosamine)<sub>n</sub> core antigen, known as the i antigen, that was reversible. Other LPS variants lost the ( $\text{Le}^x$ )<sub>n</sub> main chain resulting in the expression of monomeric ( $\text{Le}^y$ )-core-lipid A or had acquired  $\alpha$ 1, 2-linked fucose expressing polymeric  $\text{Le}^x$  and  $\text{Le}^y$  simultaneously. Most *H. pylori* isolates have been shown to be able to switch back to the parental phenotype but with varying frequency [102].

Moreover, poly-C tract length variation causes frame shifts in *H. pylori*  $\alpha$ 3-glycosyltransferases that can inactivate gene products in a reversible manner. Serological data suggested that LPS structural diversification arises from phase variable regulation of glycosyltransferase genes, provisionally named *futA* and *futB* [103]. Phase variation of *futA* and *futB* genes independently has been confirmed and genetic exchange between these loci was shown to occur in single colonies from the same patient and also during *in vitro* passage [104].

*H. pylori* strain NCTC 11637 also has been shown to express blood group antigen H type I. This epitope demonstrated high frequency phase variation that was reversible. Insertional mutagenesis of gene *jhp563* (a poly-C tract sequence containing an ORF homologous to glycosyltransferases) in NCTC 11637 showed that LPS then lacked the H type I epitope. DNA sequence analysis confirmed gene-on and gene-off variation. In *H. pylori* strain G27 mutagenesis of *jhp563* yielded a mutant expressing  $\text{Le}^x$  and  $\text{Le}^y$  as opposed to wild-type [H type 1,  $\text{Le}^a$ ,  $\text{Le}^x$  and  $\text{Le}^y$ ]. *Jhp563* may encode a  $\beta$ 3-galactosyltransferase involved in H type I synthesis that phase varies due to poly-C tract changes [105].

*H. pylori* ORF HP0208, and its homologues HP0159 and HP1416, show homology to the *waaJ* gene that encodes a  $\alpha$ 1, 2-glycosyltransferase required for core LPS biosynthesis in *Salmonella typhimurium*. HP0208 contains multiple repeats of the dinucleotide 5'GA at its 5' end and transcription of its gene product has been predicted to be controlled by phase variation. Most strains examined, including strains 26695, J99 and NCTC 11637, had repeat numbers inconsistent with expression of the gene; i.e. placing the translational initiation codon out-of-frame with the full length ORF. A 'phase-on' HP0208 was constructed in the genome of strain 26695. Tricine gel and Western blot analysis demonstrated a role for HP0208 as well as HP0159 and HP1416 in the biosynthesis of core LPS [106]. It is likely that the biosynthesis machinery of not only the *H. pylori* LPS O-antigen side chain but also the core oligosaccharide of *H. pylori* LPS

is subject to phase variation. These complex processes possibly give rise to the diversification of LPS observed in clonal populations of *H. pylori*.

#### 3.1.3.2. *Lewis expression in vitro*

The  $\alpha 1$ , 2-fucosyltransferase (*futC*) of *H. pylori* catalyses the conversion of Le<sup>x</sup> to Le<sup>y</sup>, the repeating units of the LPS O-antigen. *futC* is subjected to phase variation through slipped strand mispairing involving a poly-C tract. Single colonies ( $n=379$ ) from *in vitro* cultures have been examined for Lewis expression and demonstrated equal distribution of Le<sup>x</sup> and Le<sup>y</sup> expression and the phenotypes correlated with *futC* frame status. The founding population remained, since phenotypes did not change significantly over additional hundreds of generations *in vitro* [107].

Two single colonies of the same isolate of *H. pylori* that expressed Le<sup>y</sup> of different molecular weights demonstrated wild-type Lewis phenotype after 50 *in vitro* passages after expansion of a larger cell mass; however after 50 *in vitro* passages of single colonies, ~5% of the analysed strains also expressed considerable levels of Le<sup>x</sup> in addition to low levels of Le<sup>y</sup>, suggesting reduced expression of *futC*. Successive *in vitro* passaging of single colonies introduced a much more frequent phenotypic diversification in terms of O-antigen size and Le<sup>x</sup> expression [104].

#### 3.1.3.3. *Lewis expression in vivo*

With a limited number of passages of strains in the laboratory, analysis of the phenotypic diversity of Lewis antigen expression from 180 clonal *H. pylori* populations from primary cultures of 20 gastric biopsies indicated a substantial difference in Lewis expression in 75% of the patients. The variation of Lewis expression was unrelated to the overall genetic diversity. In experimentally infected rodents however, Lewis expression was highly uniform [108]. Intra population diversity of Lewis expression has since been confirmed. *H. pylori* isolates with identical DNA signatures (arbitrary primed PCR) from the same chronically infected patient demonstrated variations in the amount and size (length) of the O-antigen and immunoassays detected exclusively the presence of Le<sup>y</sup>, suggesting simultaneous expression of both  $\alpha 1$ , 2- and  $\alpha 1$ , 3-fucosyltransferases. LPS diversification has also been investigated in transgenic mice expressing Le<sup>b</sup> on gastric epithelial cells. The challenging strain expressed a high molecular weight O-antigen and showed a strong antibody response against Le<sup>y</sup>. More than 90% of the mouse output isolates produced glycolipids of low molecular weight compared with the input strain. Subsequent immunoblot analysis demonstrated decreased or no Le<sup>y</sup> expression [104].

#### 3.1.3.4. *Adhesins and cell surface proteins*

Expression of bacterial outer membrane proteins can be regulated by environmental changes through signal transduction as well as the generation of genetic changes controlling protein function. Cell surface associated proteins are the most abundant group of *H. pylori* proteins that is subject to phase variation. Such proteins include so called adhesins, flagellar and flagellar hook proteins, pro-inflammatory proteins, cysteine-rich proteins as well as some other categories. With exception of adhesins, most proteins in this group remain uncharacterized.

The outer membrane of *H. pylori* partially comprises adhesins, which bind to host gastric epithelial cell surface receptors. Gene functions of *H. pylori* adhesins, many of which belong to the so called *hop* gene family, are regulated through phase variation.

The sialic acid binding adhesin (SabA) of *H. pylori* adhere to glycosphingolipids that display sialyl Lewis x antigens. Such antigens have been shown to be upregulated on human epithelial cell surfaces as a result of gastric inflammation [109]. DNA sequence analysis demonstrated that locus HP0725 (*sabA*) of *H. pylori* strain 26695 contained repetitive CT dinucleotides at the 5' end of the ORF [57]. Translational modification may encounter premature termination (non-functional protein) or a full length functional adhesion protein. *sabA* has promoter poly-T as well as ORF 5' end CT tract repeats. Multiple length alleles have been shown to occur in single colonies isolated from the same individual and genome sequence analyses of isolates of *H. pylori* strains demonstrated genetic and phenotypic variation of SabA [110].

The *H. pylori* protein HopZ is a candidate to be involved in the adherence to host gastric epithelial cells and *hopZ* is likely phase variable due to a CT dinucleotide repeat in the signal sequence of the gene [111]. Human volunteers have recently been challenged with a *H. pylori* strain with *hopZ* 'off' status. Out of 56 re-isolated strains (from 32 volunteers at 3 month post inoculation) 68% had switched to a *hopZ* 'on' status. After 4 years, paired isolates had 54% *hopZ* 'on' status. Sequence analysis of *hopZ* 'on' and 'off' status in 54 *H. pylori* strains representing seven different phylogeographic populations (hpAsia2, hpEurope, hpNEAfrica, hpAfrica1, hpAfrica2, hpEastAsia and hpSahul) and 11 subpopulations, demonstrated variability between and within most populations; only two subpopulations (hspAfrica2SA and hspAmerind) were exclusively *hopZ* 'off' [112].

Many *H. pylori* strains bind to the human blood group antigen Le<sup>b</sup>. This adherence is mediated by the blood group antigen binding adhesin BabA. Some strains contain two alleles; *babA1* which is 'silent' and *babA2* that expresses the adhesin [113]. Although BabA expression has not been identified to be phase variable by DNA sequence analysis [99], experimental infection of rhesus macaques showed that, in some animals, compared with its parent challenge strain, output strains lost BabA expression due to an alteration in dinucleotide CT repeats in the 5' coding region. Output strains from other macaques that also had lost Le<sup>b</sup> binding, had *babA* exchanged for *babB*. Duplication of *babB* has also been observed in human clinical *H. pylori* isolates [75]. *babB* and *babA* have very similar 5' and 3' end sequences but *babB* lacks the mid region that codes for the Le<sup>b</sup> binding epitope. BabB is an uncharacterized outer membrane protein and *babB* contains repetitive sequence motifs and is likely subject to frameshift-based phase variation [57, 97].

Gene function of *hopH* (*oipA*) depends on slipped strand mispairing in a CT dinucleotide repeat in the signal sequence of the gene. HopH has been associated with increased interleukin-8 (IL-8) production in epithelial cells and gastric epithelial adherence *in vitro* [114, 115]. *H. pylori* isolates with a *hopH* 'in-frame' status were more common than 'out-of-frame' strains in patients with chronic gastritis, a feature that correlated with the virulence factor status of the strains, particularly *cagA* [114]. In a patient setting, *hopH* (*oipA*) 'in-frame' status was associated with a higher colonization density of *H. pylori* and clinical presentation regarding gastric inflammation and mucosal IL-8 production [116].

### 3.1.3.5. Acid adaptation

*H. pylori* can reversibly change its membrane phospholipid composition, producing variants with differing concentrations of lysophospholipids. Lysophospholipid-rich cells are more adherent, secrete more VacA and are more haemolytic. As opposed to neutral culture conditions, growth at low pH (3.5) renders an accumulation of membrane lysophospholipids. This variation in lipid composition is mediated by phase variation in the phospholipase A (*pldA*) gene. A change in the C-tract length of the ORF results either in a functional full-length or a truncated non-functional gene product [117].

The structure and composition of LPS adapts to an acid environment. Under acidic growth conditions (pH 5) *in vitro*, the LPS core and lipid A moieties seem not to be altered. However, the O-side chain backbone is partially fucosylated forming Le<sup>x</sup>, whereas the terminal sugar residues on the O-side chain are modified differently and terminate with Le<sup>y</sup> instead of Le<sup>x</sup> [118].

### 3.1.3.6. Immune modulation

*H. pylori* genomes contain a family of genes coding for proteins designated Helicobacter cysteine rich proteins (Hcp). HcpA, a secreted protein, has partially been characterized. Recombinant HcpA, as opposed to HcpC, induced maturation of non-adherent human Thp1 monocytes into macrophages (star-like morphology with filopodia) with phagocytic ability and surface adherent properties [119].

Surfactant protein D (SP-D), a component of innate immunity, is expressed in human gastric mucosa. SP-D has an affinity for simple sugars and likely functions as a mucosal receptor that recognizes pathogen associated molecular patterns (PAMPs). SP-D induces aggregation of microorganisms facilitating pathogen clearance by neutrophil and macrophage phagocytosis. Some *H. pylori* strains lack a ligand for SP-D and this 'escape' mechanism is associated with phase variation of the LPS structure. Fucosylation of the O-side chain, determined by slipped strand mispairing in a fucosyltransferase gene leading to terminal Le<sup>x</sup> (SP-D binding) or Le<sup>y</sup> (escape), controls the *H. pylori* ligand recognized by SP-D [120].

DC-SIGN, a C-type lectin, is a surface receptor expressed on dendritic cells that captures and aids the internalization of microbial antigens. *H. pylori* strains that express Lewis antigens on their LPS bind DC-SIGN and thereby block T helper cell (Th) 1 development, whereas some strains that do not express Lewis antigens escape binding to dendritic cells and promote a Th 1 response. Phase variation of LPS in terms of Lewis expression may influence host immunity through the dendritic cell pathway. Clonal populations of *H. pylori*, with high frequency of subclone phase variation in LPS biosynthesis genes, are proposed to manipulate the Th response for optimal persistence that prevents severe atrophy and destruction of the ecological niche [29, 121].

### 3.1.3.7. Experimental infections

Phase variable OMP gene switch status of *H. pylori* have been investigated in an experimental mouse model of infection. The gene switch status of several OMPs (*hopH*, *hopZ*, *hopO* and *hopP*) influenced *H. pylori* density in gastric tissue and its ability to colonise mice. If two or



more of the genes were in the switch 'off' mode, the colonization ability was markedly reduced. These results correlated well with observations in humans; i.e. patients with strains whose *hop* genes were in the 'off' mode had lower bacterial load [115].

In a murine model of infection, repeat length and function of 31 phase variable genes of several *H. pylori* strains were followed for up to one year. At endpoint, 15 genes had a change in repeat length. However, a third of these did not lead to an alteration in protein expression. Ten genes demonstrated a frame shift to an 'on' mode of the encoded protein. At early time points (3 and 21 days), mixed *pldA* phenotypes rapidly and exclusively changed to 'on' status, followed by LPS biosynthesis genes modifying terminal sugars. Glycosyltransferases modifying LPS core structures remained in 'off' configuration throughout the study. From 21 days onward some OMPs (*babB* and *hopZ*) switched from 'on' to 'off'. Restriction/modification systems did not show a particular pattern over time [122].

BabA expression has been shown to be lost during experimental infection in rhesus macaques, either by allele replacement with *babB*, or phase variation. A follow up study investigated this phenomenon in other animal hosts using additional *H. pylori* strains. Murine and gerbil experimental infection models further demonstrated loss of Le<sup>b</sup> binding (as a result of *babA* recombination) of output strains by varying mechanisms. In the mouse, BabA expression was lost due to phase variation in a 5' CT repeat region of *H. pylori* strain J166 [123].

To conclude, *H. pylori* randomly exhibits phase variation in sets of genes that directly interact with the environment. These include LPS biosynthesis genes, adhesins and genes with an impact on the structural composition of the bacterial outer membrane. Phase variation also influences the expression of some genes affecting host immune responses. Taken together, these traits are likely to aid a continuous adaptation to the ecological niche and persistence of the bacterium.

Selection pressure in terms of host niche physiology and maturation of host immune response likely contributes to the genetic regulation and diversification of bacterial adherence properties as well as the composition of the outer membrane. For bacteria such as *H. pylori* that may cause life-long colonization, surface antigen diversity likely requires parallel evolution with host cell subsets over time for continuous adaptation to a dynamic host environment.

#### 3.1.4. Epigenetic diversity: phasevarion

As previously described, phase variation via the high-frequency reversible ON/OFF switching of gene expression is beneficial to pathogenic bacteria, including *H. pylori*, as a means of rapidly generating the genotypic and phenotypic diversity required for adaptation to the host environment and evasion from the immune system [124]. However, certain evolutionary advantages arise when expression of a repertoire of genes is brought under the influence of a single phase-variable gene as is the case for a phase-variable regulon or "phasevarion" (Figure 2).

The phasevarion is an epigenetic regulatory system whereby the expression of a set of genes is randomly switched as coordinated by the activity of the modification (Mod) component of a restriction-modification (R-M) system [51]. R-M systems share two components; the restric-



tion (Res) component that specifically cleaves unmethylated DNA at a recognition sequence, and the Mod component that methylates the same recognition sequence to prevent cleavage by the Res component [125]. Most R-M systems fall within one of three major families (Type I, II, III). Type II and type III Mod proteins recognise specific DNA sequences whereas type I Mod proteins require an additional specificity subunit [126]. A role for R-M systems in host-pathogen interaction was unexpected, as these systems generally function to protect the genome integrity of bacteria from invasion by foreign DNA by restriction of DNA that does not share the same modifications as that of the host. However, where the Res component is absent or not functional, phase variation of the Mod component results in the random switching of methylation of DNA sequences recognized by the *mod* encoded methyltransferase. For genes where DNA methylation by the phase-variable Mod affects promoter activity, either by altering the DNA binding affinity of regulatory proteins for promoters [127], or by other mechanisms, differential gene expression results.

Although phase-variable R-M systems had previously been identified in a number of bacteria [128], the first experimental evidence for the phasevarion was from *H. influenzae*, a pathogen of the upper respiratory tract [51]. The *mod* gene of the sole type-III R-M system of *H. influenzae* contains tetranucleotide repeats consistent with an ability to phase vary. Microarray analysis of a mutant in this gene revealed differential expression of 16 genes, including genes implicated in pathogenicity, in the absence of *mod*. The differential expression of these genes was shown by reporter gene fusions to be dependent on the phase variation of *mod*. Hence the phase variation of a single gene, *mod*, was shown to influence the expression of multiple genes, suggesting the presence of a phase-variable regulon, or “phasevarion.”

Phasevarion mediated gene switching has now been confirmed in *H. influenzae* [51], *N. gonorrhoeae* and *N. meningitidis* [129], and *H. pylori* [50]. This wide distribution, and the allelic diversity of phase-variable methyltransferases, indicates that there is strong selective pressure on phasevarions. It has been postulated that it may be simpler for a gene to evolve to join a phasevarion than become phase variable. The evolution of phase-variation requires the generation of repeat sequences without destroying either promoter or gene function, whereas joining a phasevarion requires only a few key point mutations to generate the methyltransferase recognition site in a region where methylation would affect gene expression [51]. A further evolutionary advantage of the phasevarion may be that it represents an extension of the regulation achieved by phase variation. Rather than randomly reversibly switching a single gene, the phasevarion switches a whole set of genes, thereby differentiating a bacterial population into two different cell types based on many phenotypic characteristics [50, 130]. This switching between different physiological states, rather than merely individual proteins, may assist the bacterium in taking advantage of microenvironments within the host.

Genome sequencing of *H. pylori* 26695 [57] and J99 [97] revealed a surprisingly large number of R-M systems (22 in 26695) when compared to other sequenced bacterial and archaeal genomes. Each strain of *H. pylori* contains its own complement of R-M systems, some of which may have the potential to phase vary due to the presence of repeat regions [47, 97, 131, 132]. Typically multiple *mod* genes may be present in any given strain and there may be multiple different *mod* alleles for each *mod* gene within a given species [128]. Diversity of *mod* genes can

be driven by recombination of DNA recognition domains between non-orthologous genes and horizontal gene transfer [133, 134]. Phylogenetic analysis of clinical isolates of *H. pylori* revealed a diverse set of 17 alleles of the *modH* gene that differed in the DNA recognition domain and phase-varying repeat region. This diversity in *mod* genes indicates corresponding diversity in the set of genes regulated by Mod and also indicates that there may be many phasevarions present in *H. pylori* [50]. Any R-M system present within the genome with an inactive Res may represent a phasevarion. Phasevarions may therefore represent common epigenetic mechanisms for generating rapid reversible phenotypic diversity in bacterial host-adapted pathogens such as *H. pylori*.

Functional analysis of the Type-II R-M systems in J99 and 26695 revealed that less than 30% were fully functional (with both Res and Mod functional) and that there were many functional Mod enzymes with no apparent functional Res partner [132, 135], indicative of these *mod* genes regulating a phasevarion. Repeat sequences and homopolymeric tracts, indicative of phase variation, were identified in a number of R-M systems (both type-II and type-III) in *H. pylori* [47, 57, 97, 98, 130]. Direct experimental evidence of phase variation within an R-M system was first given by *lacZ* reporter fusions to a type-III R-M system [136]. An R-M system has also been shown to phase vary in the mouse model, suggesting a role for these systems in host-adaptation [122].

Given that *H. pylori* contains a diverse repertoire of R-M systems that phase vary and R-M systems with orphaned Mod proteins, it is likely that many of these systems regulate a phasevarion. The first evidence that the activity of a phase-variable Mod influenced the transcription of other genes in *H. pylori* came from the analysis of the type-II methyltransferase M.HpyAIV in 26695. M.HpyAIV was found to phase vary due to the presence of homopolymeric tract of adenine residues. Analysis of the genomes of 26695 and J99 revealed 60 genes common to both strains where the M.HpyAIV DNA-methylation sites occurred in the intergenic region upstream of ORFs, indicating a possible role for these sites in regulating gene expression. Differential expression of these genes was studied by qPCR and the expression of catalase encoding *katA* was shown to be significantly decreased in a 26695 mutant of M.HpyAIV [137].

Further evidence of the presence of phasevarions in *H. pylori* came from deletion of the *modH5* type-III methyltransferase from *H. pylori* P12. Microarray analysis of the mutant strain revealed six genes, including outer membrane protein HopG and flagellar associated proteins, that were differentially expressed compared to wild-type [50].

A phasevarion regulated by *hpyAVIBM* may also be present in *H. pylori*. The presence of AG repeats in this type-II methyltransferase indicates that this gene may have the potential to phase vary, although this has not been experimentally demonstrated. Deletion of *hpyAVIBM* from *H. pylori* SS1 and clinical isolate AM5 resulted in differential regulation in a diverse set of genes, including outer membrane proteins, genes involved in motility, pathogenicity, LPS biosynthesis, and R-M systems, when compared to WT by microarray analysis. Further analysis revealed corresponding alterations in phenotype of the *hpyAVIBM* mutant such as altered motility, increased expression of CagA as determined by Western blot, altered LPS profile, improved ability to induce IL-8 production in human AGS cells, and a decrease in

transformation efficiency. Differences observed in gene expression and phenotype due to deletion of *hpyAVIBM* varied between the two strains investigated potentially due to different distribution of the methylation recognition site of HpyAVIBM across the genome of the two strains. The occurrence of the *hpyAVIBM* allele in *H. pylori* strains isolated from individuals with duodenal ulcer and healthy individuals revealed that the methyltransferase was present in most strains isolated from symptomatic patients but absent in most strains isolated from healthy individuals, indicating that *hpyAVIBM* expression may be clinically relevant [138]. Even if *hpyAVIBM* does not phase vary, this study demonstrates the wide ranging regulatory capabilities of DNA methyltransferases.

### 3.2. Inter-strain generation of diversity

#### 3.2.1. Natural transformation

*H. pylori* is naturally competent, it is able to be transformed by the uptake and incorporation of foreign DNA into its genome [139]. Potential reasons for natural competence in bacteria are a matter of discussion. It is postulated that bacteria may utilize the uptake of foreign DNA for nutrition, for DNA repair, for evolution via horizontal gene transfer, or that DNA-uptake is an evolutionary spandrel of adhesion and twitching motility [140, 141]. As we will discuss, it is becoming increasingly clear that natural transformation plays an important role in *H. pylori* genome evolution and host adaptation.

##### 3.2.1.1. Quick overview of players in uptake and recombination

A complete picture of the process of DNA uptake followed by integration into the genome in *H. pylori* is beginning to emerge, although many details remain to be clarified. Uptake of foreign DNA into the bacterial cell is achieved by a two-step mechanism as shown in Figure 3. Firstly, dsDNA is taken up through the outer membrane by the ComB type-IV secretion system [142]. The ComEC system is then responsible for DNA uptake through the inner membrane [143, 144]. Transport of DNA by ComEC likely results in the entry of single-stranded DNA into the cytoplasm based on the function of ComFA in *Bacillus subtilis*, although this has not been directly experimentally demonstrated in *H. pylori*. Incoming DNA is at some point subjected to the activity of restriction endonucleases [145]. Once in the cytoplasm DprA and RecA cooperatively bind the incoming ssDNA forming a heterodimer [146]. During recombination RecA mediates strand invasion of incoming DNA with chromosomal DNA and this process is subject to interference from UvrD and MutS2 [49, 73, 90, 147]. RecA mediated synapsis with chromosomal DNA results in the formation of four-way branched DNA intermediate structures referred to as Holliday junctions whose migration and resolution are mediated by DNA processing enzymes. Branch migration of Holliday junctions is mediated by either the competing RuvAB or RecG helicases [86]. Resolution of the Holliday junction is primarily by DprB in instances of natural transformation with homeologous DNA from other *H. pylori* strains, although RuvC, the DNA-repair resolvase, can partially compensate for a loss of DprB [148].

### 3.2.1.2. Structure of the uptake system

In contrast to other bacterial species where DNA-uptake occurs via systems related to type IV pili, the ComB system of *H. pylori* is related to type IV secretion systems and its components have been named for their homologues in the *Agrobacterium tumefaciens* VirB type IV secretion apparatus [142, 149]. The genes encoding ComB are organized in two separate loci with an operon consisting of *comB6 – 10* and a second operon consisting of *comB2 – 4* [142, 150-152]. All the *comB* and *comEC* genes are essential for competence with the exception of *comB7* which is postulated to play a stabilizing role for the comB complex [142, 143, 151]. Sequence homology with the VirB type IV secretion system and topological mapping of the ComB proteins has given some insight into the structure of the ComB apparatus [151, 153]. ComB2 is postulated to be located as a “stump structure” in the external membrane and have a role in initial DNA-uptake. It has also been associated with adhesion to human gastric tissue [152]. ComB7 is also associated with the outer membrane and may serve to stabilize ComB9, which is present in the periplasm with an anchor to the outer membrane via a disulphide bond. ComB8 contains a large periplasmic domain and spans the inner membrane and may interact with ComB9 and/or ComB10 which is postulated to be anchored in the inner membrane where it may be present as a homodimer. ComB4 is cytoplasmic and serves as the ATPase that energises the ComB machinery. A role for ComB3 is largely unknown although it is predicted to contain one transmembrane domain.

### 3.2.1.3. Process of DNA uptake

The process of DNA uptake by *H. pylori* has been studied using fluorescently labelled DNA and single molecule analysis with laser tweezers [143]. The initial step in transformation is the binding of extracellular DNA to the surface of the bacterium. The ComB machinery may play a role in DNA-binding to the cell as mutants lacking inner-membrane components of the ComB machinery showed impaired DNA-binding [151]. Once bound, DNA was found to be rapidly taken into the periplasm of the cell through ComB via an ATP-dependent mechanism likely driven by the ATPase ComB4. Multiple DNA-uptake complexes were found to be simultaneously active. Uptake of DNA through the outer membrane by ComB appears to be non-specific as uptake of DNA was not distinguished on the basis of DNA sequence. Following transport of DNA into the periplasm by ComB, ssDNA enters the cytoplasm via ComEC. Transport of DNA by ComB and ComEC appears to be spatially and temporally uncoupled. The identity of any motor driving uptake of DNA by comEC remains to be uncovered. Transport of DNA by ComEC appears to be more discriminating than ComB, as covalently labelled DNA transported by ComB could not enter the cytoplasm via ComEC. As DNA sequence does not seem to play a role in the initial uptake of DNA by *H. pylori*, discrimination of incoming DNA and protection from the potentially hazardous consequences of the incorporation of foreign DNA may come from the numerous restriction-modification systems of *H. pylori*.



#### 3.2.1.4. *The restriction barrier - frequency*

The fate of incoming DNA is either one of restriction or recombination. Restriction of foreign DNA forms the most significant barrier to natural transformation in *H. pylori* [145]. Like other bacteria, *H. pylori* discriminates the DNA of self from non-self by the modification of bases by methylation. Restriction modification systems (R-M) consist of a methyltransferase that methylates specific DNA sequences and a restriction endonuclease that cleaves non-methylated DNA at the same sequence. Incoming foreign DNA that does not share the same methylation pattern as that of the host is thus digested. In this manner, many R-M systems function to prevent transformation and protect the host from foreign DNA [125]. The number and diversity of R-M systems in the *H. pylori* genome is notable, with many being strain specific [57]. The diversity of R-M systems can be driven by deletion and acquisition of such systems by horizontal gene transfer [154]. These strain specific R-M systems may be responsible for the observation that competence of different *H. pylori* strains varies [155]. The barrier posed by R-M systems to competency has been experimentally demonstrated by assessing transformation frequency in the presence and absence of R-M systems. The removal of four active type-II restriction endonucleases from *H. pylori* 26695 lead to higher transformation efficiency both of donor DNA from *E. coli* and other *H. pylori* strains [156]. The removal of two active type-II restriction endonucleases from NSH57 greatly reduced the barrier to transformation resulting in greater transformation frequency with DNA from a J99 donor [145].

#### 3.2.1.5. *The restriction barrier – integration length*

In addition to transformation frequency, restriction of incoming DNA also influences the length of incoming DNA that is integrated into the host chromosome. It has been proposed that although *H. pylori* takes up long DNA fragments by natural transformation, only shorter fragments are integrated into the genome. Genomic sequencing of isolates revealed that sequences recombined with imported DNA varied in length from 261 to 629 bp and were clustered. This observation was suggested to be consistent with the uptake of long stretches of DNA, corresponding to the length of the region in which recombination sites were clustered, that has subsequently been broken up and partially integrated [157]. Mutants of *H. pylori* NSH57 lacking active type-II restriction endonucleases were found to integrate longer DNA fragments into their genome following natural transformation than the WT strain [145].

#### 3.2.1.6. *DrpA overcomes restriction*

Although restriction presents a barrier to transformation, *H. pylori* achieves a balance between restriction and recombination. It has been proposed that the concentration of restriction enzymes in the cell may be limited to produce only partial cleavage of incoming DNA in order to allow a basal level of transformation [145]. In addition to this proposal, the DNA processing protein A (DrpA), has been shown to lower the barrier to recombination in a number of bacterial species and is widely conserved [146]. Deletion of *drpA* in *H. pylori* has been reported to result in either a significant decrease [158, 159], or abrogation of transformation frequency [145]. In *H. pylori* DrpA has a polar localisation and interacts with incoming DNA, binding ssDNA and also dsDNA to a lesser extent. DrpA protects incoming DNA from restriction by

both preventing the access of type-II restriction endonucleases to DNA and enhancing methylation of incoming DNA by direct interaction with methyltransferases. It thus appears to play a key role in the balance between restriction and recombination [160]. However, the temporal and spatial aspects of restriction endonuclease cleavage and DprA activity are not yet clearly understood. Single-stranded DNA is thought to enter the cytoplasm following uptake by ComEC and DprA binds preferentially to ssDNA, yet restriction enzymes, which contribute the most significant barrier to transformation, find ssDNA a poor substrate in preference to dsDNA.

#### 3.2.1.7. DNA processing enzymes and competence


As depicted in Figure 3, following entry into the cytoplasm, incoming DNA is co-operatively bound by DrpA and RecA. RecA acts to mediate strand invasion of foreign DNA with the host chromosome and promotes homologous recombination. The nucleotide excision repair helicase, UvrD, likely disrupts this process by removal of RecA from DNA, preventing potential recombination events. Mutants in RecA therefore are unable to undergo recombination whereas mutants in UvrD display a hyper-recombination phenotype [49, 73]. MutS2 is also proposed to interrupt RecA-mediated strand invasion. Deletion of MutS2 from *H. pylori* results in an increase in transformation efficiency suggesting a role of MutS2 in regulation of homologous recombination. Analysis of MutS2 indicates that it is not a member of the mismatch repair pathway but rather has a distinct function in strand displacement of incoming DNA from RecA-mediated D-loop formation during strand invasion of incoming DNA with the host chromosome. This function is independent of the degree of homology of the two strands [90, 147]. Thus in addition to the restriction barrier, enzymes involved in determining the outcome between formation or dissolution of the D-loop also appear to play a role in determining the fate of foreign DNA. The influence on transformation of other DNA processing enzymes acting during the recombination event is less clear. Reports regarding transformation efficiency of a *recG* mutant vary [145, 161] and integration length in a *recG* mutant was reported to be decreased [145]. In contrast to an initial report where deletion of *ruvC* resulted in a decrease in transformation frequency [162], two reports have found no decrease in transformation frequency [145, 148] but an increase in integration length in a *ruvC* mutant was noted [145]. Reports regarding the phenotype of a mutant in *dprB*, the recombination specific Holliday junction resolvase, are consistent that deletion of *dprB* results in a decrease in transformation frequency but does not influence integration length [145, 148].

#### 3.2.1.8. Other factors

A further factor required for competence that appears to be unique to *H. pylori* is ComH. Mutants in *comH* could not be transformed with chromosomal or plasmid DNA [163, 164]. ComH is a surface-exposed outer membrane protein that binds ssDNA but whose function in competence remains to be further characterized [165].

The role of the nuclease NucT in *H. pylori* competence is also unclear. NucT is an outer membrane bound nuclease that preferentially cleaves ssDNA. The observation that transformation rates are reduced in a *nucT* mutant leads to the proposal that NucT functions either in





**Figure 3.** Natural transformation in *H. pylori*. Schematic representation of the uptake of foreign DNA and its integration into the host chromosome in *H. pylori*. Double stranded DNA is taken up through the outer membrane by the ComB machinery and through the inner membrane by ComEC. Following DNA uptake ssDNA is bound by RecA and DprA, which affects the activity of restriction endonucleases and methyltransferases. Strand invasion of the host chromosome by foreign DNA is mediated by RecA and this process is inhibited by MutS2 and UvrD. Branch migration and resolution are mediated by RecG and DprB, respectively. NucT may function as a nuclease for the acquisition of purines and the exact role of ComH in competence has yet to be determined.

initial DNA binding, or translocation into the cell [166]. In contrast, a later study found that deletion of *nucT* did not influence transformation efficiency, but did result in an increase in the length of DNA integrated into the chromosome [145]. The presence of NucT as a stable outer membrane nuclease and the requirement of *H. pylori* to scavenge purines for growth as a result of its inability to synthesise purines *de novo* led to an investigation of the role of NucT in purine acquisition [167]. NucT was found to be required for growth when exogenous DNA is the only purine source, indicating that the primary function of NucT is the digestion of human DNA in the gastric mucosa as a purine source. Digestion of dsDNA in the absence of its preferred ssDNA substrate *in vitro* could be responsible for the conflicting results obtained in assays of transformation efficiency.

#### 3.2.1.9. Regulation of competence is by DNA damage

Competence is usually a tightly regulated and transient feature in bacteria. *H. pylori* is unusual in that it displays very high levels of competence. Competence was found to vary across stages of growth with each strain displaying a different pattern of peaks in transformation efficiency in different growth phases. The pattern observed was independent of the type of donor DNA [168]. Competence is upregulated in response to DNA damage by the induction of the transcription and translation of competence machinery. A marginal increase in transformation frequency was observed after UV induced DNA damage [73]. Transcription of several genes involved in competence were found to be upregulated in both cDNA microarrays of cells where acute double strand break DNA damage had been induced by ciprofloxacin treatment and in cells where chronic DNA damage had accumulated in *addA* mutants deficient in double-strand break repair. These genes include components of the ComB system and a lysozyme proposed to function by lysing neighbouring cells to provide DNA for uptake. This transcriptional response was dependent on both *recA* and the ability to uptake DNA. The authors proposed a RecA-dependent positive feedback loop in the induction of DNA damage responsive genes that is initiated by DNA damage and amplified by DNA uptake. This system was found not to be important in initial mouse colonization but may be of relevance to persistence. These results demonstrate that unlike other bacteria, *H. pylori* does not mount an SOS response to DNA damage but relies on homologous recombination for maintenance of genome integrity [169]. It seems plausible that *H. pylori* responds to stress induced by the immune system during persistence in the human host by increasing competence to not only repair DNA damage but also to increase genetic diversity in order to continually adapt to a changing host-mediated niche [63].

#### 3.2.2. Competence *in vitro*

##### 3.2.2.1. Substrate requirements for transformation

The characteristics of natural transformation in *H. pylori in vitro* have been studied in detail. Investigations into the influence of the properties of substrate DNA on natural transformation of *H. pylori* 26695 revealed a number of interesting features [155]:

- Transformation efficiency decreases with shorter DNA fragments although transformants could be obtained with fragments as short as 50 bp.
- Although uptake of DNA occurs within minutes of exposure, transformation efficiency increases with increasing time prior to transformant selection to allow time for uptake, recombination, and expression of a new phenotype.
- Transformation frequency of selectable alleles decreases with decreasing length of flanking sequences, although transformants were obtained with as little as 5 bp flanking sequence.
- Transformation frequencies are higher with chromosomal DNA than PCR products.
- Transformants could be obtained with both single stranded and double stranded DNA but with 1000-fold greater efficiency for double stranded DNA.
- Transformation efficiency is greater with homologous DNA than homeologous DNA.
- Different *H. pylori* strains vary in their competence.
- DNA uptake could be saturated at high DNA concentrations.

#### 3.2.2.2. Length of insertions

Study of transformation *in vitro* has also revealed that *H. pylori* typically imports fragments of short length into the chromosome in comparison to other bacteria and that these imports are regularly interrupted by wild-type recipient sequences. Mean lengths of between 1294 and 3853 bp of integrated DNA were observed from transformations of rifampicin sensitive recipient strains with DNA from resistant donors with different import lengths observed with different donor/recipient combinations [170]. Transformation of streptomycin sensitive strains with DNA of streptomycin resistant strains obtained a mean length of integration of 1300 bp. Furthermore the effect of the restriction barrier may have been observed, as endpoints of integration were found to be clustered, consistent with restriction of incoming DNA at those sites [171]. Short length of imported fragments has also been recorded *in vivo* [157].

#### 3.2.2.3. Why are insertions interspersed?

The region of integrated DNA is commonly reported to be interrupted by short interspersed sequences of the recipient (ISR) and multiple explanations for this observation are present in the literature. Lin *et al.* have suggested ISR could result from two separate but neighboring strand invasion events that would be consistent with the restriction of an invading strand prior to recombination [171]. Conversely, Kulick *et al.* reported that overexpression of the base excision repair glycosylase MutY increased the occurrence of ISR within imported regions, implicating it in their formation. MutY was also found to influence integration length. It was proposed that MutY-mediated DNA repair at impaired bases following recombination results in the insertion of host sequence within recombined regions [170]. The allelic variation generated by ISR is a further mechanism increasing genotypic variation.

#### 3.2.2.4. Promiscuity of DNA uptake

*H. pylori* is very promiscuous as it does not require incoming DNA sequences for transformation to be as closely related as what other bacterium do. This may be due to the absence of a mismatch repair system, a lack of DNA sequence specificity in DNA uptake machinery, absence of DNA uptake sequences, and a relaxed restriction barrier. Recombination between unrelated strains in co-infected hosts has been observed. Analysis of synonymous distance between recombined alleles within the genomes of *H. pylori* isolates from two South African families revealed recombination with sequences from unrelated strains, in contrast to studies of *S. enterica*, *E. coli*, and *B. cereus* where recombination only occurred between members of the same lineage [172]. Recombination can also occur between different *Helicobacter* species. Transformation of *H. pylori* 26695 with both homologous and homeologous streptomycin resistance conferring *rpsL* DNA derived from 26695 and *H. cetorum* respectively, yielded transformants, although with 1.5 log<sub>10</sub> greater efficiency with homologous DNA. No transformants could be obtained with DNA from *C. jejuni*. Competition between DNA from different sources for transformation revealed that *H. pylori* DNA could compete with *H. pylori* DNA but DNA from unrelated sources could not compete with *H. pylori* DNA, indicating that *H. pylori* can distinguish DNA from different sources on the basis of DNA sequence [155, 173]. This discrimination does not occur at the level of DNA uptake by ComB/ComEC, as cytoplasmic uptake of  $\lambda$ -phage DNA was comparable to *H. pylori* DNA [143], but is likely a consequence of the restriction barrier. Thus although transformation of *H. pylori* is efficient with DNA from unrelated strains, and even to a lesser degree with DNA of other *Helicobacter* species, it has not been observed with DNA from other bacterial genera.

*H. pylori* displays considerable allelic diversity and genetic variability. The high degree of competence facilitates frequent horizontal transfer of genetic material to the extent that the genome has been found to be in linkage equilibrium [174, 175]. Although genetic variability typifies *H. pylori*, with populations being regarded as panmictic, clonality is observed in the natural transmission of strains within closely related and co-habiting individuals [176]. Also, *H. pylori* populations can be grouped according to geographic location as strains located within a region are more closely related to each other than to strains outside the region [2]. The uptake of foreign DNA by natural transformation and resulting recombination generates a considerable portion of the genetic diversity observed in *H. pylori*.

#### 3.2.3. Plasmids and mobilizable transposons conjugation

In many bacterial species plasmid transfer by conjugation is a significant contributor to the acquisition of genetic material by horizontal gene transfer and often mediates the dissemination of genes of particular phenotypic importance such as antibiotic resistance, virulence determinants, and the ability to utilize certain substrates. Bacterial conjugation can generate chromosomal rearrangements due to plasmid insertion and excision and can also transfer chromosomal genetic material when errors in excision occur. Although diversity exists in bacterial conjugative mechanisms, a generalized overview of the process can be formed. Initially a mating pair of cells must make contact and be brought into proximity. This is typically achieved by the expression of the sex pilus, an elongated tubular appendage, by the

donor cell. Once in contact a conjugative pore or some other mechanism for the transfer of DNA to the recipient must be established. This is commonly achieved by a type-IV secretion system. In order to transfer plasmid DNA, a relaxase and accessory proteins, the relaxosome, bind to the plasmid origin of transfer (*oriT*) where they cleave a single strand. The single strand and bound relaxase is recognized by a coupling protein and transferred into the recipient cell in tandem with the rolling circle replication of the intact strand of the plasmid. The relaxase then circularizes the single strand which is replicated to form an intact plasmid within the recipient [177].

*H. pylori* commonly carry plasmids of varying sizes [178], often of low copy number, which can be divided into two groups [179, 180]. The first are homologous to Gram positive vectors that replicate via the rolling circle mechanism. The second, more common group, are proposed to replicate via the theta mechanism predominately utilising the replication protein, RepA, which binds to short tandem repeats in the plasmid origin of replication. There are several reports characterising the properties of various plasmids in *H. pylori* [180-185].

Plasmid transfer has been demonstrated to occur between different strains of *H. pylori* [186, 187]. Mating experiments with *H. pylori* P8 and P12 revealed that transfer of two conjugative *H. pylori* plasmids could occur through three distinct routes. Firstly, natural transformation, which was dependent on the ComB type-IV secretion system and was DNaseI sensitive, secondly, DNaseI insensitive mobilisation which was dependent on both ComB and the plasmid encoded relaxase (*mobA*), and finally, an alternative DNaseI resistant (ADR) pathway that is independent of ComB. No evidence was found for the involvement of chromosomal relaxases in any pathway [187]. Backert *et al.* also demonstrated conjugative plasmid transfer in *H. pylori* but used two mobilizable vectors containing a broad host range *oriT* [186]. In this instance transfer was insensitive to DNaseI and dependent on viable cell contact, indicative of a conjugative process. No role was found for ComB, indicating that natural transformation was not taking place, but both the chromosomally encoded TraG coupling protein homolog and relaxase *rlx1* were required. The mechanism of transfer thus appears similar to the ADR pathway observed by Rohrer *et al.* It may be that the ADR pathway is a minor pathway for conjugative plasmids with an endogenous relaxase as utilized by Rohrer *et al.*, but may be the only pathway for transfer of mobilizable plasmids via the utilization of chromosomal mobilisation genes. Transfer of *H. pylori* conjugative plasmids was found to occur at a rate orders of magnitude higher than that for the introduced mobilizable plasmids ( $10^{-4}$  vs  $10^{-7}$ ). The observation that *H. pylori* is capable of transferring plasmids with a broad-host range *oriT* utilising a chromosomal mobilisation system indicates that *H. pylori* may be promiscuous in its ability to uptake foreign plasmids. In both studies the *cagPAI* and *tfs3* type-IV secretion systems were found to play no role in plasmid transfer. Given that no type-IV secretion apparatus has been implicated in the ADR pathway, the mode of DNA transfer by this route has yet to be determined.

The high rate of plasmid occurrence and the presence of three pathways for plasmid acquisition indicate that genes carried by plasmids may assist in host adaptation and be an important source of genetic diversity amongst *H. pylori* strains. The observation of chromosomal sequences in *H. pylori* plasmids indicates that they are capable of acquiring genetic material



from the chromosome [181, 188]. Additionally, the presence of genes, displaying homology to chromosomal genes of unknown function, flanked by repeat and IS sequences on pHel4 and pHel5 gave rise to the proposal that *H. pylori* plasmids have a modular structure where genes can be integrated from the chromosome or other plasmids at repeat sequences by recombination or IS sequences by transposition events. Such shuffling of genes could represent a mechanism generating inter-strain diversity in *H. pylori* [180].

A role for plasmid encoded genes in virulence or host adaptation has yet to be clearly demonstrated as most plasmids characterized to date carry predominately only elements required for replication and hypothetical genes of unknown function [179]. Genes with homology to the *E. coli* microcin toxin operon have been identified in plasmids pHel4 and pHPM8. These genes function in *E. coli* to block protein biosynthesis in closely related bacteria, but their function has yet to be demonstrated in *H. pylori*. Similarly, both plasmids also contain a gene with homology to the tetracycline resistance determinant, *tetA*, but which has been demonstrated not to confer tetracycline resistance to host cells and thus remains of unknown function [179, 180, 188].

Plasmids are not the only entities transferred by conjugation in *H. pylori*. Initial evidence that *H. pylori* may be capable of transferring chromosomal elements by conjugation came from mating experiments in the presence and absence of DNaseI [189]. Subsequently, the transfer of chromosomally encoded streptomycin resistance by conjugation was demonstrated from *H. pylori* into *C. jejuni*. The transfer required cell to cell contact and was independent of any known type-IV secretion system [190].

More recently, the horizontal transfer of *H. pylori* plasticity zones has been investigated. Comparison of the first two genome sequences of *H. pylori* revealed the presence of chromosomal regions, termed plasticity zones, which contained strain specific genes. These regions vary from the rest of the genome in their G+C content, indicating possible acquisition from a foreign source [56]. Plasticity zones have been found to contain genes of interest, including DNA processing enzymes, type IV secretion systems [191, 192], and genes implicated in disease outcome. Plasticity regions have been found to be diverse in their gene content and presence in *H. pylori* strains [191, 193-195]. A large number of studies have implicated plasticity region localized genes, particularly JHP947 and surrounding genes, as virulence determinants in *H. pylori*. The presence or absence of these genes in clinical isolates have variously been correlated with gastric carcinoma and duodenal ulcer disease status and have been associated with variations in inflammatory cytokines [194-202]. The plasticity zones of *H. pylori* have been found to be able to horizontally transfer as conjugative transposons [191, 193].

Conjugative transposons are a form of integrative and conjugative elements (ICEs). These elements reside within, and are replicated by, the host chromosome and contain the elements required for their excision and integration. Following excision they circularize, are replicated and transferred to a recipient cell by conjugation. Following transfer, the ICE integrates into the recipient chromosome and the copy remaining in the donor also reintegrates. Many plasticity zones and genomic islands present in many bacterial species may be functional ICEs or remnants of mobile elements [203]. Given the importance of genomic and pathogenicity islands in the physiology of many bacteria, in particular the *cag* pathogenicity island and genes

implicated in disease within the plasticity zones of *H. pylori*, ICEs may represent a rapid means of generating genetic diversity of clinical relevance between different *H. pylori* strains. Deletion of plasticity zones has been found to decrease the fitness of some strains *in vivo* and mediate the induction of inflammatory cytokines in human AGS cells [193].

An investigation into the nature of plasticity zones by sequencing *H. pylori* strains to identify their gene content and genomic location lead to the proposal that they were conjugative transposons termed 'transposon plasticity zone' (TnPZ) [193]. The plasticity zones in the sequenced strains could be classified according to their gene content and arrangement. Plasticity zones of a particular class could be identified at different genomic contexts in different strains, indicating their ability to move as a discrete unit. The plasticity zones were often found to be flanked by direct repeats and contained terminal inverted sequences, characteristic of transposable elements. There was also evidence of recombination occurring between different classes of TnPZs and many strains appeared to contain vestigial remnants TnPZs including J99 and 26695 whose plasticity regions were described as 'complex mosaics of TnPZ remnants'. Full length TnPZs were found to contain DNA processing enzymes, including a candidate transposase, *xerT*, which displays homology with the recombinases XerC and XerD of *E. coli* and the *H. pylori* *dif*-specific XerH recombinase. Transfer of TnPZ1 of *H. pylori* strain P12 was demonstrated by co-cultivation experiments performed with both WT and mutant donor and recipient strains [191]. Transfer of the plasticity zone was observed by a conjugative mechanism that could occur, although at reduced frequency, in the absence of the TnPZ-encoded type-IV secretion system and was dependent on the TNPZ-encoded *xerT*. Circular DNA intermediates from donor strains could be visualised by PCR. Excision of the TnPZ was found to leave one copy of the direct repeat in the donor chromosome while the circular intermediate contained the second.

The genes present on TnPZ may represent a pool of genes that can be quickly acquired as a set, as the conjugative method of transfer is not subjected to the same restriction barrier as transformation. It seems that plasticity zones have been present in *H. pylori* for a long time as similar zones are also found in other *Helicobacter* species including *H. acinonychis* and *H. cetorum* and many have been subject to gene deletion and recombination over that time. Plasticity zones are widely present in *H. pylori* strains with each strain having a unique combination of such zones and the strain specific genes found within, they thus contribute significantly to the *Helicobacter* pan-genome [191]. The transfer of TnPZs may represent a further method utilized by *H. pylori* to generate a diverse population that displays fitness in a diverse set of niches. The physiological function of genes in plasticity zones requires further investigation to clarify their potential roles in host adaptation and disease outcome. Although conjugative transfer of TnPZs, other chromosomal elements, and plasmids does occur in *H. pylori*, the precise mechanisms of such transfer and their regulation are not completely understood. The contribution of conjugative processes to genetic diversity in *H. pylori* may not be as significant as that contributed by natural transformation but is still of evolutionary significance.

### 3.2.4. Transduction of bacteriophage

Phages are often important drivers of genetic diversity and host adaptation. Phages represent a significant mode of horizontal gene transfer responsible for a large portion of strain-specific genetic diversity in many bacterial species. The action of temperate phages may play a significant role in the evolution of bacterial genomes by the disruption of host genes during insertion events, delivery of bacterial DNA for recombination by transduction, or the introduction of new genes (morons) that are carried, but not required, by an integrating phage. This has phenotypic consequences and the presence of prophages in the bacterial genome may alter bacterial fitness and influence host-bacterial interactions. This is particularly significant in many instances where morons encode virulence factors such as bacterial toxins [204]. Despite the significance of phages in the evolution of other bacterial species, the presence of phages in *Helicobacter* has drawn little attention until recently.

Bacteriophages are relevant to the physiology of *H. pylori* *in vivo*. Early observations of *H. pylori* noted the presence of bacteriophage-like particles within cells in electron microscope images of gastric biopsies [205]. Similarly, electron microscopy of clinical isolate SchReck 290 [206] revealed the presence of the bacteriophage HP1 that could be propagated in a lytic cycle *in vitro* [207]. Active *H. pylori* bacteriophages can be isolated from human faeces [208].

More recently, several bacteriophages have been characterised in *H. pylori*. Genome sequencing of clinical isolate, B45, revealed the presence of a 24.6 kb prophage within the genome [209]. The phage was found to be similar to a previously identified phage in *H. acinonychis* [210] and could be induced by UV irradiation, producing phage particles. Screening of a further 341 clinical strains from different geographical regions by PCR revealed that the phage was widespread, being detected in 21.4% of isolates. This high prevalence could indicate that bacteriophages play a role in *Helicobacter* evolution [209].

The bacteriophage 1961P was isolated from a clinical isolate of *H. pylori*. It has a genome of 27 kb containing 33 genes. Sequence comparison revealed that prophages or prophage remnants similar to 1961P are likely present in six other sequenced *H. pylori* strains. Interestingly, 1961P was demonstrated to be a transducing phage demonstrating that transduction of *H. pylori* by bacteriophages does occur [211].

The sequence of two phages isolated from East-Asian-type isolates from Japanese patients has been reported [212] and one, KHP30, has been characterized further [213]. The phage did not appear to be integrative and may represent a novel category of bacteriophage.

Although it is becoming clear that bacteriophages may be widely present in *H. pylori*, with some recently well characterized, a potential role for bacteriophages in genomic evolution and host-adaptation in *H. pylori* remains largely unexplored. Genomic sequencing revealed the presence of two prophages within the feline-associated *H. acinonychis* genome which account for 26% of genes present in the sequenced *H. acinonychis* but absent in *H. pylori* J99 and 26695. Although most of these genes appear to have been acquired after the divergence of the two species and largely encode hypothetical proteins, three genes from one of the two prophages were implicated by the authors as potentially having played a role in enabling the host jump of the *H. acinonychis* ancestor into felines [210].

Lehours *et al.* have raised the hypothesis that plasticity zones in *H. pylori* could have a phage origin much as phages mediate the transfer of pathogenicity islands in *Staphylococcus* [214]. Kersulyte *et al.* have proposed that plasticity zones transfer as conjugative transposons [193]. Further research will be required to determine if the bacteriophages in *Helicobacter* have played any significant role in the evolution of its genome.

#### 4. Small RNAs and regulation

To date only a limited number of transcriptional regulators have been discovered in the small 1.67 Mbp genome of *H. pylori*, including a mere three sigma factors, namely RpoD, FliA and RpoN [57]. Consequently *H. pylori* was thought not to have any extensive regulatory networks. However, a recent comprehensive analysis of the *H. pylori* primary transcriptome has revealed that the compact genome produces an abundant number of antisense and small RNA (sRNA) transcripts portending a great potential for the use of riboregulation by *H. pylori* in gene expression [215].

Small RNAs have emerged as essential regulators that allow organisms to cope with environmental changes and stresses [216, 217]. Like transcription factors, sRNA can modulate the expression of multiple target genes and thereby function as key regulators of metabolic pathways and stress responses. In bacteria, sRNAs have been discovered to regulate processes as varied as carbon metabolism, iron homeostasis, RNA polymerase function, virulence, quorum sensing, biofilm formation, as well as response to stresses such as oxidation and outer membrane perturbation [216, 218-220].

Bacterial sRNAs exhibit a diverse range of molecular mechanisms of action. One class of bacterial sRNA acts by binding directly to protein targets and modulating their activity [221, 222]. Another group of sRNAs known as riboswitches consist of RNA sequences located in the 5' untranslated regions (UTRs) of mRNAs. Riboswitches regulate gene expression through their capacity to adopt different conformations that are mediated by factors such as temperature or small molecule metabolites that specifically bind to the riboswitch [223, 224].

The most extensively studied class of sRNAs are those that act through antisense base pairing with target mRNAs. These sRNAs are classified as either *cis*-encoded, because they are transcribed from the strand of DNA opposite their mRNA targets and so have extensive complementarity to their target, or as *trans*-encoded sRNAs, which are transcribed from a genomic location different from those of their targets. The base pairing interactions of *trans*-encoded sRNAs tend to involve less complementarity with their targets, permitting the regulation of multiple genes [217]. Furthermore many *trans*-encoded sRNAs in enteric bacteria require binding to Hfq, a chaperone protein that aids in sRNA stability and base pairing with target mRNAs [225]. Antisense base pairing of sRNAs can exert either negative or positive regulatory effects on their mRNA targets including RNA degradation, termination of transcription, inhibition of translation and the relief of intrinsic translation-inhibitory structure formation in the mRNA's 5' UTR (For detailed reviews of basepairing sRNAs and their regulatory mechanisms, refer to [220, 226-228]).



The discovery of substantial antisense transcription as well as a high number (>60) of sRNAs which included potential regulators of *cis*- and *trans*-encoded mRNA targets has revealed a new class of molecules that should be considered for their role in mediating *H. pylori* persistence. Since sRNAs in *H. pylori* were discovered only recently, knowledge regarding their target genes and the mechanism by which they regulate them is still very limited. The current literature on riboregulation for *H. pylori* has been recently reviewed [229]. When corrected by genome size, the sRNA repertoire of *H. pylori* rivals that of *E. coli*, the model organism of bacterial RNA research [220], and thus the mechanisms of sRNA action in *H. pylori* are likely to be much more diverse than what is currently known.

With the exception of a handful of housekeeping RNAs, none of the enterobacterial sRNAs are conserved in *H. pylori* [215]. *H. pylori* has no Hfq homolog, an RNA chaperone shown to be important for mediating sRNA function in enteric bacteria, and *H. pylori* lacks homologs of endonucleolytic RNases E/G and other common processing factors of stable RNAs [57, 230]. It has been hypothesized that antisense-mediated processing by double-strand specific ribonuclease RNase III may compensate for this paucity and act as the major regulator as has been recently suggested for *S. aureus* [229, 231]. To date the only sRNA shown to be essential in *H. pylori* and required for stress response is tmRNA, which is involved in the rescue of stalled ribosomes [232].

One of the most abundant transcripts identified in *H. pylori* is a homolog of 6S RNA [215], a ubiquitous riboregulator which mimics an open promoter complex and thereby sequesters RNA polymerase [233, 234]. In *E. coli*, deletion of 6S RNA has no obvious phenotype during exponential growth, however altered growth phenotypes are observed during stationary phase and under extreme stress conditions [222]. Whether 6S RNA has a similar role during stress response, stationary growth, or if it impacts on virulence as seen for *Legionella* [235] and therefore has implications in *H. pylori* persistence in the host still needs to be investigated.

Antisense transcripts have been identified for one third of the putative phase-variable genes with functions in lipopolysaccharide biosynthesis, surface structure and DNA restriction/modification [215], raising the possibility of antisense regulation of surface structure and host interactions. Furthermore, acid-stress-induced antisense RNAs opposite to known acid-stress-repressed genes have also been detected, suggesting that antisense regulation would also play an important role in adapting to changing environmental pH. To date only a few physiological effects of sRNA regulation have been demonstrated in *H. pylori*. The *ureAB* operon has been shown to be negatively regulated by a *cis*-encoded antisense 5' *ureB*-sRNA which is induced by unphosphorylated ArsR under neutral conditions [236]. The chemotaxis receptor TlpB, which is involved in pH-taxis, quorum sensing, colonization and gastric mucosa inflammation, has been shown to be negatively regulated by the *trans*-encoded HPnc5490 sRNA [237-239].

It has become increasingly clear that sRNAs serve as diverse regulators that impact almost every aspect of bacterial physiology in response to changes in the environment. It is therefore highly likely that riboregulation may have a very important role in *H. pylori* persistence. The stomach is a highly dynamic environment undergoing constant changes in pH, due to long periods of fasting interspaced with the intake of various food material, while continual shedding of gastric mucus, where a majority of the bacteria reside [240], leads to washout into



the lower intestine where the bacteria are exposed to an anaerobic environment and greater exposure to the host immune system through Peyer's patches. Here riboregulation may be very important for the quick changes and modulation, such as initiation of peptidoglycan remodelling and alterations in the lipid content [241, 242], necessary to transition from the spiral to coccoid form that is phagocytosed by dendritic cells in Peyer's patches [243].

## 5. Experimental strategies to study *Helicobacter pylori* persistence

A multidisciplinary approach is required to study persistence *in vivo*, through the combination of microbiology, immunology, genetics and clinics. Although powerful techniques in cellular microbiology as well as the flurry of transgenic mouse strains have been instrumental in investigating *H. pylori* pathogenesis, *in vivo* studies are limited to loss of function mutants based on gene deletion. This prevents the distinction between colonization and/or persistence functions of the gene of interest. Furthermore, deletion mutants also suffer the drawback of potential rapid genetic adaption in case of a strong selective pressure (which might often be the case already *in vitro*) *in vivo*, and this is particularly relevant in view of *H. pylori*'s genome plasticity.

### 5.1. Gene deletion

Both bacterial and host genetics have led to a tremendous progress in our understanding of *H. pylori* pathogenesis. However, the mechanisms of persistence of *H. pylori* have not yet been rigorously tested in animal models due to the difficulty in assessing the temporal requirement of virulence factor expression for colonization and/or persistence. Only a few genes have been strictly shown to be required for persistence based on the inability of mutant strains to colonize in the long-term despite identical short-term colonization loads compared to wild-type. For example, the peptidoglycan deacetylase (*pgdA*) mutant displayed significant attenuation in its ability to colonize mouse stomachs at 9 weeks infection although it is dispensable for up to 3 weeks during the initial colonization [244]. Furthermore, compared to wild-type *H. pylori*-infected mice, elevated levels of MIP-2, IL-10 and TNF $\alpha$  were observed in mice infected with the *pgdA* mutant, indicating that peptidoglycan deacetylation modulates the host immune system [244]. Another study reported the DNA repair *ruvC* mutant to be spontaneously cleared from the gastric mucosa of mice between 36 to 67 days of the initial challenge of mice [88]. Interestingly, the *ruvC* mutant was less efficient in modulating the murine immune system, suggesting that DNA recombination is critical for immune modulation and persistence of *H. pylori* [88]. Catalase (KatA) and the catalase associated proteins (KapA) have also been implicated in persistence of *H. pylori* infection [245]. KapA and KatA were proposed to promote survival of *H. pylori* in the inflamed gastric mucosa where the concentration of reactive oxygen species, particularly hydrogen peroxide, is high [245]. Very recently, a comparative analysis of the colonization loads between wild-type *H. pylori* and the *lpxE* isogenic mutant, harbouring a partial TLR4 determinant restoration, was conducted in C57/BL6J and the corresponding TLR4 knockout transgenic mice [26]. This two-sided genetic approach demonstrated that although LpxE-mediated avoidance of TLR4 recognition had little effect on the early coloni-

zation phase 15 days) *lpxE* mutant was prevented from persisting for more than 45 days due to TLR4 activation [26] that likely led to sterilizing adaptive immunity.

The natural competence of *H. pylori* has also recently been demonstrated to promote chronic colonization. A competition experiment between wild-type and *comB10* or *drpA* mutant strains showed that mutation of either the Com apparatus or a cytosolic competence factor resulted in reduced persistence despite normal initial colonization [246]. This recent data strongly suggests that the exchange of DNA between a heterogeneous population and genome plasticity and integrity are important for *H. pylori* to maintain chronic infection.


## 5.2. Gene mutagenesis: The urease example

As an alternative to gene deletion, protein engineering was applied to the urease complex to investigate its role in the host-pathogen interaction without affecting the enzymes ureolytic activity which is essential for colonization [247]. Surface exposed loops that display high thermal mobility were targeted for in-frame insertion mutagenesis (Figure 4). *H. pylori* expressing urease with insertions at four different sites retained urease activity and were able to establish robust infections. Bacteria expressing one of these four mutant ureases, however, had reduced bacterial loads after longer term (3 to 6 months) colonization. These results indicate that a discrete surface region of the urease complex, distinct from the ureolytic activity *per se*, is important for *H. pylori* persistence during gastric colonization [247].

*H. pylori* urease consists of a ball-like supramolecular structure of about 1000 kDa. It is composed of twelve UreA-UreB subunits (26.5 and 61.7 kDa, respectively), assembled in a high molecular weight complex of four alpha/beta trimers [248, 249]. Urease is an abundantly expressed enzyme (10% of the cell weight) that decreases the acidity of *H. pylori*'s immediate environment by generating ammonia and carbonate from the urea we secrete as metabolic waste and is one of the first characterized factors identified as essential for colonization by *H. pylori* [250-252]. The majority of the urease is localized in the bacterial cytoplasm, however, urease is also present on the cell surface and in the extracellular medium. The secretion of urease is still a matter of debate as both autolysis and specific secretion [250, 253, 254] have been postulated to be responsible for the release of urease.

Uptake of urea into the cytoplasm is controlled by an acid gated urea channel, UreI, that does not open until the periplasmic pH decreases below pH 6.5 to avoid alkalization of the cytoplasm [255-257]. The recruitment and interaction of the urease complex with UreI at the inner membrane is hypothesized to enable coupling of urea transport and urease activity for efficient pH homeostasis of the periplasm [258]. It is tempting to postulate that the surface properties of urease plays a key role in the intrabacterial regulation of its enzymatic activity by docking of the urease complex to the urea channel (UreI) to maintain periplasmic pH.

Although such local control of gastric acidity is considered essential, urease-negative *H. pylori* strains were unable to colonize piglets whose acid secretion had been suppressed, suggesting an additional role for urease [259]. Possible explanations include the utilization of urease generated ammonia by *H. pylori* to synthesize essential metabolites, especially amino acids [260]; protection from peroxynitrite [261], enhanced survival in macrophages [262],



**Figure 4.** Recombinant regions of urease and selection for enzyme function (Figure adapted from [247]). **a)** Molecular structure of urease showing insertion sites on the surface of urease. Urease subunit A (green) and subunit B (blue) associate to form a dodecameric supramolecular molecule [248, 249]. Sites 1 to 8 correspond to residues 102, 231 and 238 from UreA and residues 1, 66, 326, 541 and 549 from UreB, respectively. Insertion sites 1, 3, 4, and 8 are indicated in red. Urease activity could not be retained when altered at sites 2, 5, 6, and 7 (pink). **b)** Selection of bacteria producing functional urease on acidified media supplemented with the urease substrate, urea. Left side: X47 wild type; the colour change observed on the left side indicated that bacterial colonies were producing functional urease and growing. Right side: X47  $\Delta$ ureA: there was no colour (X47 wild-type). Colour change did not occur on the right side, indicating that inoculated colonies were unable to grow or functional urease was not being produced (X47  $\Delta$ ureA). **c)** A schematic showing insertion sites at the urease locus of DNA coding epitopes and linkers. Insertions were made in DNA corresponding to insertion after amino positions 102 (site 1) and 238 (site 3) of UreA (GenBank AAD07144.1), and amino acid positions 1 (site 4) and 549 (site 8) of UreB (GenBank AAD07143.1). Insertions at sites 3 and 4 correspond to the C- and N-termini of UreA and UreB, respectively. DNA coded HA(T): hemagglutinin T cell epitope; HA(B) hemagglutinin B cell epitope; SR linker: semi-random linker; linker: GPSL linker; FLAG: FLAG epitope; STOP: STOP codon.

evasion of phagocytosis [263], complement mediated opsonisation [264], and decreased viscosity in gastric mucin at high pH facilitating *H. pylori* motility [265]. An alternative explanation invokes the existence of urease-host tissue interactions, that are independent of urease enzymatic activity, and is based on *in vitro* studies that detected urease mediated

activation of macrophages [266], monocytes [267] and blood platelets [266, 267], dysregulation of gastric epithelial tight junctions [268] and induction of cytokine production from gastric epithelial cells [269] through binding to CD74 (MHC class II invariant chain) [270].

The above examples highlight the importance of studying the kinetics of colonization so as to gain deeper insights into the *H. pylori* persistence mechanisms and revisit the role of well-known and studied virulence factors such as urease.

### 5.3. Conditional mutants

The use of conditional *H. pylori* mutants makes it possible to investigate genes for their role in persistence by turning off target gene expression *in vivo* once colonization has been established. Several *H. pylori* conditional knockouts based on the *lac* operon have been constructed to study essential genes *in vitro* [271, 272]. So far however, this system has not been tested in the mouse model of *H. pylori* infection. Due to certain limitations, including leakiness of transgene expression and limited bioavailability of the IPTG transgene-inducers, this *lac* operon-based system is not very well suited for *in vivo* studies. On the contrary, the tetracycline-mediated gene expression regulation has been extensively used to construct mouse conditional knockouts as well as to study host-pathogen interactions using tetracycline-regulated bacterial conditional knockouts [273, 274]. Tetracycline gene regulation has recently been developed in *H. pylori* [275-277] (Figure 5) and successfully applied *in vivo* to monitor changes in colonization load upon the down regulation of *ureB* gene expression [277].


This system will enable study of the temporal requirements of specific genes during *H. pylori* colonization as well as during persistence. More importantly, tetracycline-mediated *H. pylori* gene regulation will allow for subtle characterization, *in vivo* and *ex vivo*, of changes in the innate and adaptive immune responses by monitoring the activation of antigen presenting cells and T cell proliferation, respectively.

The tetracycline gene regulation combined with host genetics will facilitate the investigation of *H. pylori* persistence mechanisms and may allow for the discovery of new markers for cancer risk and for novel therapeutic interventions specifically targeting persistence.

## 6. Comparative genomics

The genome era created the possibility to study gene contents of individual *H. pylori* strains, allelic diversity and related genetic plasticity. The first *H. pylori* strain, 26695, was sequenced in 1997 and was isolated from an English patient with chronic gastritis. The chromosome of strain 26695 is circular and composed of 1 667 867 base pairs [57]. J99, isolated from an American patient with a duodenal ulcer, was sequenced in 1999 and for the first time, a comparison of two unrelated genomes was made [97]. Compared to 26695, J99 has a slightly smaller circular chromosome (1 643 831 base pairs). *H. pylori* is believed to possess a large degree of genomic diversity but the overall genomic organization, gene order and predicted proteomes of the two sequenced strains was found to be similar. The two genomes displayed





**Figure 5.** Using conditional mutants to study the role of genes in persistence. In the top panel the conditional *H. pylori* mutant constitutively expresses the *tet* repressor (orange circles), encoded by *tetR*, which binds to the tandem *tet* operators, *tetO*, preventing transcription of the target virulence gene. The host immune cells encounter bacteria lacking a virulence factor and produce a particular measurable response (immune response A). In the bottom panel, the molecular inducer of TetR, Tetracycline (Tc), indicated by filled yellow triangles, is administered to the system. Tc diffuses into the bacterial cell, binds to the repressor-operator complex and triggers a conformational change in TetR resulting in rapid dissociation of TetR from the DNA and a burst of target gene expression. The host immune cells encounter bacteria which express the target virulence factor and in turn produce a different response (immune response B).

a high degree of diversity in terms of insertions and deletions. There are 1406 genes shared by both strains, but 86 open reading frames are absent from strain 26695. Both strains contain a complete *cag* pathogenicity island that codes for a type IV secretion system to deliver the CagA cytotoxin protein. Comparison of the two genomes revealed between 6 to 7% of the genes were specific to each strain, with almost half of these genes being clustered in a single hyper-variable region or plasticity zone. The presence of a variable gene pool is indicative of horizontal gene transfer. Because strain-specific genes could be involved in gastric adaptation during co-evolution, this flexible gene pool was extensively investigated in many strains worldwide. *In silico* analysis of the two *H. pylori* genomes revealed that both housekeeping genes and virulence genes are transferred between *H. pylori* strains [278, 279].



Next generation sequencing has enabled the study of unrelated *H. pylori* strains isolated from unrelated individuals and further studies have focused on evolution of *H. pylori* within the same host. Commonly, isolates have been derived from chronically infected individuals that have been exposed to *H. pylori* for decades, such as 26695 and J99. More recently, analysis of the genetic relationships between strains of *H. pylori* that had been sequentially isolated from the same host at different times also showed high mutation rates and genetic diversity.

Inter-strain diversity is represented by variations in the number and contents of genes, chromosomal rearrangements and allelic diversities, and is not unique to *H. pylori* [280]. For *H. pylori*, each strain contains many strain-specific genes. It has been proposed that a particular bacterial species contains a core set of genes and auxiliary genes. The core genome contains genes that are present in all or nearly all of the strains. It determines the properties that are characteristic for the species. The auxiliary genes are present in some strains, and are determinants of the biological properties unique to each strain.

### 6.1. Chronic infection

The first studies of genetic diversity and evolution amongst *H. pylori* strains were performed using isolates derived from chronically infected individuals. Early studies of genetic change during *H. pylori* chronic infection were restricted to selected housekeeping genes within a bacterial population in single and mixed infected hosts.

Molecular fingerprinting studies using RAPD [281] and amplified fragment length polymorphisms (AFLP) [282] first showed that strains isolated from single patients were closely related, but were subtly different. Strains isolated at the same time or months and years apart also showed similar divergence [281-285]. Thus, while populations diverge within the host, divergence appears to occur slowly.

Newer technologies allowed for more accurate comparisons of *H. pylori* genomes. Salama *et al.* reported genetic diversity amongst 15 unrelated *H. pylori* strains from different geographical locations and identified strain-specific genes based on whole genome *H. pylori* DNA microarray. Analysis of the genomic content of *H. pylori* strains from chronically infected patients found that 22% of *H. pylori* genes were dispensable and defined a minimal core genome of 1,281 *H. pylori* genes [286]. Of these, more than 300 genes were not homogeneously distributed and many of the dispensable genes were located in plasticity zones and in the *cag* pathogenicity island. Core genes encoded mostly metabolic and cellular processes, while strain-specific genes included genes unique to *H. pylori*, restriction modification genes, transposases and genes encoding cell surface proteins.

Gressmann *et al.* used a collection of 56 globally representative *H. pylori* strains that included examples from all known populations and subpopulations in comparative genome hybridizations with a microarray representing the genes of the combined genomes of 26695 and J99. The study found 1,150 genes were conserved between the 56 strains and estimated that the core genome conserved in all *H. pylori* strains is ~1,111 genes. The remaining 400 genes that each *H. pylori* genome contains come from a pool of genes that are strain-specific and located within the *cag* pathogenicity island [287], consistent with the findings of the previous study.

Other studies also identified the *cag* pathogenicity island and plasticity zones as regions of genetic diversity and DNA exchange. Israel *et al.* examined the gene content of two *H. pylori* strains that had similar virulence genotypes (*cagA*<sup>+</sup> *vacA* s1a *iceA1*), but were phenotypically different and showed by *H. pylori* whole genome microarray analysis that the gene content within the *cag* island differed substantially [288]. Occhialini *et al.* examined the composition of the plasticity zone in a collection of 43 *H. pylori* strains from diverse clinical origins and showed that the plasticity zone is highly mosaic and represents a large fragment of foreign DNA integrated into the genome [194].

Genetic variation can be generated in a bacterial population by mutation and/or recombination between different strains. The studies described so far focus on unrelated clinical isolates from unrelated infected individuals. However, to address the question of mutation and recombination rates of *H. pylori* during chronic infection, a comparison of *H. pylori* isolates from the same host must be made.

One of the first such studies compared the genome of J99, originally isolated in 1994, with multiple strains that had been isolated from the same patient six years later [289]. All follow-up strains differed from the original isolate by one or multiple gene losses or gains. Randomly amplified polymorphic DNA PCR and DNA sequencing of four unlinked loci revealed that these isolates were closely related to the original strain. In contrast, microarray analysis revealed differences in genetic content among all of the isolates that were not detected by randomly amplified polymorphic DNA PCR or sequence analysis [289]. Whole-genome microarray revealed a difference in genomic composition in 3% of J99 loci as well as the relatedness of these isolates to each other when compared with *H. pylori* isolates from other individuals.

Subsequent studies observed sequentially isolated *H. pylori* strains from the same host. One of the early reports examined the sequences of ten gene fragments (encoding house-keeping enzymes and virulence associated proteins) of paired sequential isolates from 26 patients in two geographical areas at between 3 month and 48 month intervals (average 1.8 year interval) and found that point mutations occur in the stomach of a single host and that mostly small DNA segments (median size of 417 base pairs) are exchanged with other bacteria in the stomach [175]. A Bayesian model was used to calculate mutation rate, import size and the frequency of recombination. On average, pairs of bacteria differed by ~100 DNA imports, corresponding to three percent of the genome or 50 kb. A mutation rate of  $4.1 \times 10^{-5}$  per year and base pair was estimated. Recombination occurred at a rate of 60 imports spanning 25,000 base pair per genome per year. Authors concluded that recombination is so frequent that appreciable fractions of the entire genome are exchanged during the colonization of a single human, resulting in a highly flexible genome content and frequent shuffling of sequence polymorphisms throughout the local gene pool [175].

Kraft *et al.* examined paired strains of *H. pylori* with respect to their genomic contents using the DNA microarray method and also reported evolutionary changes in the *H. pylori* genome. Isolates were obtained from the same patients at intervals of 3 to 36 months. Of the 21 pairs of strains examined, 4 pairs showed differences in their genomic contents, suggesting the occurrence of evolutionary recombination events. These included a complete deletion and a

partial loss of the *cag* pathogenicity island, a replacement of an open reading frame of unknown function, an acquisition of 14 genes in the plasticity zone, a duplication of the *ceuE* genes (HP1561/HP1562) and a truncation of tandem arranged *ackA* and *pta* genes resulting in the formation of pseudo-genes [290].

A more recent study estimated the short-term mutation and recombination rates of *H. pylori* by sequencing an average of 39, 300 base pairs in 78 gene fragments from 97 isolates [291]. These isolates included 34 pairs of sequential samples at intervals of 3 months to 10.2 years. They also included single isolates from 29 individuals from 10 families. The accumulation of sequence diversity increased with time of separation in a clock-like manner in the sequential isolates. Approximate Bayesian Computation was used to estimate the mutation and recombination rates, mean length of recombination tracts, and average diversity in those tracts. The short-term mutation rate is estimated to be 1.461026 (serial isolates) to 4.561026 (family isolates) per nucleotide per year and that three times as many substitutions are introduced by recombination as by mutation. Comparisons with the recent literature show that short-term mutation rates vary dramatically and can span a range of several orders of magnitude.

The above studies analysed the genetic relationships between strains of *H. pylori* sequentially isolated from the same patient. The studies are based mainly on multi-locus sequence analysis of mostly house-keeping genes and have limitations. Bayesian inference was used to estimate recombination and mutation but the multi-locus approach did not allow conclusions about the chromosomal distribution of import events or about the relative frequencies of imports in different categories of genes.

More recently, mutation and recombination rates have been estimated by a genome-wide analysis using pyro-sequencing technology. Kennemann *et al.* analysed the genomes of five sets of sequential isolates of *H. pylori*, including four pairs of isolates from the earlier studies (with isolation intervals of 3 years) and recent follow-up isolates for two of the pairs that were obtained 16 years after the first isolates. The genome comparisons of the four sets of isolates confirmed previous estimates of the length of imported fragments and reported an average length of 394 base pairs, which is in agreement with the previous estimate of 417 base pairs. The 16-year isolates differed from the initial isolates by far more SNPs and CNPs than the 3-year isolates, indicating that diversity caused by mutation and recombination had accumulated over time. The average genome-wide mutation rate for the four 3-year pairs of sequential isolates from chronically infected individuals was found to be  $2.5 \times 10^{-5}$  (range =  $0.5\text{--}6.5 \times 10^{-5}$ ) per year per site. This rate is ~18-fold faster than the mutation rate previously calculated for serial *H. pylori* isolates based on analysis of housekeeping genes. The rate of recombination was  $5.5 \times 10^{-5}$  recombination events per initiation site and year, similar to what was previously estimated [92], but 122-fold higher than the rate of  $4.4 \times 10^{-7}$  calculated from housekeeping genes. These differences can be attributed to multiple factors. Faster rates can be expected for a genome-wide analysis, because housekeeping genes are likely to be under strong purifying selection, whereas the genome-wide analysis comprises noncoding DNA as well as genes under diversifying selection. In addition, the rates of mutation and recombination varied strongly between different infected individuals in both studies, which could be because of strain properties, the extent of mixed infections that determines the availability of exogenous

DNA, or varying selective forces in infected hosts. This study of *H. pylori* genomic evolution during human infection shows genome-wide recombination in *H. pylori* colonizing humans in a high-prevalence area with a high rate of mixed infections. Genome-wide analyses of the length of individual import events were in good agreement with earlier estimates, but two important findings have emerged: 1) imports were often clustered and 2) imports frequently affected genes coding for outer membrane proteins of the Hop family.

Taken together, these provide evidence of genetic variability and DNA exchange among *H. pylori* strains and demonstrate that gene contents of *H. pylori* isolates from the same and different individuals displays between 3% and 22% variability, respectively. Data indicate that within an apparently homogeneous population, remarkable genetic differences exist among single-colony isolates of *H. pylori* and provide direct evidence that this bacterium has the capacity to lose and possibly acquire exogenous DNA and is consistent with the theory of continuous microevolution within a cognate host.

*H. pylori* are naturally competent for transformation [151] and non-random distributed repetitive sequences are found in the genome, which leads to frequent recombination events [92]. Calculations of mutation and recombination frequencies with respect to insert sizes revealed that genetic diversity displayed by the panmictic population structure is a result of continuous DNA exchange between parental strains and daughter strains, which have accumulated mutations. This was supported by gene content analysis of isolates taken from single patients at different time points, which demonstrated that the great majority of genetic changes were caused by homologous recombination, indicating that adaptation of *H. pylori* to the host individual is more frequently mediated by sequence changes acquired by recombination events rather than loss or gain of genes [290].

The comparative studies of the *H. pylori* genome reveal the genomic changes during the cycle of invasion, colonization and transmission to a new host. Invasion into a new host seems to have little effect on the gene composition of *H. pylori*, suggesting that the current genome of *H. pylori* has sufficient capacities for permitting bacterial invasion into a human host. Once the infection is established, the bacterium has to cope with the dynamic changes of the physiological environment during the long-term coexistence with the host. Genomic diversifications, or gain and/or loss of genes, occur in response to these changes. The diversifications involve genes that are mainly those strain-specific genes observed from comparative studies of unrelated strains of *H. pylori*. Intra-host evolution of *H. pylori*, thus, results in the creation of a unique and strain-specific combination of genes enabling persistence in individual hosts.

## 6.2. Acute infection and human challenge studies

Most studies of *H. pylori* adaptation to the human host have relied on samples from adults who had most likely been infected since childhood thus showing changes only after a long adaption of the bacterium to the individual host. It is also of interest to understand the evolution of the *H. pylori* genome in the first weeks and months after infection to reflect the selective pressure occurring during adaptation to a new host.



A limited number of *H. pylori* challenge studies in humans have been performed over the past 2 decades, mainly to support the development of anti-*Helicobacter* vaccines. These studies have provided valuable samples for the study of intra-host adaptation of *H. pylori* during the acute phase of *H. pylori* infection.

The first human challenge study was documented by Graham *et al.* to develop a reliable challenge model to evaluate *H. pylori* vaccine candidates [292]. Healthy human volunteers were infected with the *H. pylori* strain BCS 100 and *H. pylori* clones were re-isolated from stomach biopsies 3 months after challenge. The Baylor challenge strain (BCS 100) is *cag* pathogenicity island negative, and positive for *vacA s1c-m1*.

Another challenge study using the same BCS 100 strain study was reported for the development of a vaccine against *H. pylori* [293]. Performed in Germany at the Paul Ehrlich Institute, a live vaccine against *H. pylori* was tested in human volunteers sero-negative for, and without evidence of, active *H. pylori* infection. Volunteers (n=58) were immunised orally with *Salmonella enterica* serovar Typhi Ty21a expressing *H. pylori* urease or HP0231, or solely with Ty21a, and then challenged with *H. pylori*. Gastric biopsies were taken before and after vaccination and pre- and post-challenge.

A third unpublished acute human challenge study using a CagA positive *H. pylori* strain was reported by Malfertheiner *et al.* at the AGA Annual Meeting 2011 (AGA abstract #432, S-86). In brief, 34 healthy subjects that tested negative for *H. pylori* infection received an experimental vaccine or placebo prior to subsequent challenge with *H. pylori* isolate at a dose of  $5 \times 10^6$  CFU. The isolate was susceptible to all current antibiotics used in *H. pylori* therapy. Subjects were evaluated at 2, 4 and 12 weeks after challenge and the published data revealed changes in pepsinogen I and II and gastrin-17 levels. Authors concluded that challenge with a CagA positive *H. pylori* strain induced moderate dyspeptic symptoms that resolve within a few days and profoundly affected gastric physiology with a distinct reactive pattern of serum pepsinogen I and II and gastrin-17.

With the challenge strain BCS 100 and the reisolate 8A3, the analysis of a pair of isolates for which the time elapsed between the infection and re-isolation was exactly known, allowed the rate of genetic changes occurring in the early phase of *H. pylori* infection to be addressed. The genomes of the challenge strain (BCS 100) of the vaccine trial and a reisolate (8A3) from a volunteer (non-vaccinated control group) who had been infected with BCS 100 for 3 months [293] were sequenced. Whole-genome sequence comparison revealed very few differences between the two isolates [157]. Three point mutations were identified and confirmed by Sanger sequencing. All three were nonsynonymous mutations, leading to predicted single amino acid changes in the proteins  $\delta$ -1-pyrroline-5-carboxylate dehydrogenase (PutA; HP0056), pyridoxal phosphate biosynthetic protein J (PdxJ; HP1582), and HP1181 (a predicted multidrug efflux transporter). In addition to point mutations, noted were repeat length differences (RLDs) in two different dinucleotide repeat sequences and one repeat consisting of multiple copies of an 8-basepair motif.

In striking contrast to the pairs from the earlier study of isolates from chronically infected individuals, no single recombination event was detected. This is most likely due to the lack of



co-infection, more commonly observed in chronic infection. *H. pylori* prevalence has fallen in most Western countries, including Germany where the study was conducted, reducing the risk of acquiring multiple strains. A recent study of two pairs of sequential *H. pylori* isolates from Sweden also did not detect any evidence of recombination during chronic infection [294]. Likewise, there was no evidence of recombination during 3 months of infection of a volunteer in Germany, providing evidence that *H. pylori* can establish chronic infection after infection with a single strain and that its genome can be stable in the absence of mixed infection.

In addition, *H. pylori* isolates from four healthy adults (patients 101, 103, 104, and 105) who participated in the BCS 100 human challenge study [292] collected 15 days or 90 days post-infection were examined. Adult volunteers were not related to each other and were not related to the patient from which the challenge strain was obtained. Therefore, if host-specific differences select for genetic changes in the bacteria, such conditions would be present during this human challenge experiment. Comparative genome array analysis with single colony isolates demonstrated that their genomic contents were identical to the challenge strain. No rapid changes in gene content or sequence divergence up to 3 months after transmission was observed [295].

To date, no data has been published on the infectivity and colonisation rate of vaccinated and non-vaccinated subjects challenged with the CagA positive strain, or the genetic sequence of isolates collected at post-challenge time-points compared to the original challenge strain. This type of genome sequence analysis would provide new information on the adaptation of *H. pylori* during the acute phase of infection in the host and would allow comparison of genetic events occurring in the presence or absence of virulence genes such as CagA.

A fourth *H. pylori* human challenge study has recently been performed by investigators at Ondek Pty. Ltd. for the development of mucosal delivery technology based on live *H. pylori* (Benghezal *et al.* unpublished data). Five genotypically different strains of *H. pylori*, including CagA positive strains, were used to challenge healthy human volunteers in a Phase I study (clinical trial reference: SCGH HREC #2009-062). Briefly, 36 *H. pylori* sero-negative subjects were screened and randomised into 6 groups to receive either one of the 5 *H. pylori* strains or placebo. Subjects were monitored for the duration of the 12 week study and stomach biopsies were collected at 2 and 12 weeks post challenge, time points that depict an acute and start of the chronic phase of infection, respectively. Single colony isolates collected at the 2 and 12 week time points represent a unique resource for the investigation of *H. pylori* adaptation in humans based on a multi-strain, multi-person experimental study. Indeed, humans differ in physiologic and immunologic traits, and these traits change with age and in response to *H. pylori* infection. Such host diversity should select for adaptive changes in *H. pylori* genes important for host interaction. Yet the types of adaptive changes that *H. pylori* undergo during colonisation, the underlying mechanisms and functional significance and the resulting changes in the host response remain largely unknown. Thus a sweeping characterization of *H. pylori* adaptation is now made possible by the unique opportunity to access strains from the Ondek-sponsored clinical trial in which 6 subjects per group were each challenged with one of several *H. pylori* strains. *H. pylori* recovered during the acute (2 week) and chronic (12 week) phases of infection will be compared with input strains by genome sequencing to gain new perspec-

tives on bacterial adaptation to different human traits. Of note, comparative genomic analysis of the input/output strains of *H. pylori* will provide insight into the genome plasticity and stability of a live bacterial vector after human challenge for further development of a novel mucosal delivery technology.

## 7. Perspectives

### 7.1. Insight into bacterial pathogenesis and microbial evolution

#### 7.1.1. Genomic diversity

The sequencing of *H. pylori* host isolates has revealed the nature and extent of recombination arising from inter-strain horizontal transfer *in vivo*. Generation of genetic diversity by recombination likely relies on the presence of co-infection with multiple strains. Individuals who are already colonised are able to be infected with new additional strains [296]. A study of 127 individuals from three regions in Venezuela found evidence of mixed colonization in 55% of subjects [297]. Genome sequencing of sequential *H. pylori* isolates from chronically infected individuals in Columbia revealed high rates of recombination attributable to uptake of DNA from unrelated strains during co-infection [157]. Thus co-infection with multiple strains is likely to be common, at least in areas with endemic *H. pylori* infection [172, 298], as was the case through most of human history. Where recombination occurs between strains it has been found to introduce 100-fold more genetic alterations than mutation [172] and occurs genome-wide [157].

However, co-infection may be less pervasive in most Western countries where *H. pylori* prevalence is low. In the absence of mixed infection recombination has little effect on genetic diversity [172]. No evidence of recombination was found in a German individual experimentally infected with a single strain [157] or in sequential *H. pylori* isolates from Sweden [294]. The genome sequencing of *H. pylori* isolates from 52 members of two South African families found both high and low rates of recombination which the authors attributed to the presence of co-infection with multiple strains or infection with only a single strain, respectively [172]. Thus it is hypothesized that in instances of co-infection recombination can introduce genomic diversity but that a single strain of *H. pylori* can still establish a robust colonization and stably maintain its genome in the absence of recombination of DNA between strains, suggesting that intra-strain genetic diversity is a sufficient driver to adapt to the changing host. However, lack of opportunity for horizontal transfer between strains in mixed infections in developed countries may be accelerating the disappearance of *H. pylori* in these regions [176]. Indeed transient and self-clearing infections do occur [299] suggesting, in these instances, an inability of the bacterium to adapt to the host or a role of multiple infections in *H. pylori* transmission.

The competence of *H. pylori* likely confers evolutionary advantages. Competence has been found to confer a fitness advantage over non-competent strains *in vitro* [164]. Wild-type G27 was able to adapt to laboratory conditions more rapidly than a competence null *comH* mutant. Although competence is not required for gastric colonization in the gerbil model

[300], competence may be important for host adaptation. Natural transformation generates diversity by the introduction of new alleles and mosaic alleles in recipient strains. Greater genetic diversity amongst strains acts to increase the pool of variants available from which selection can act to confer an advantage on those who are fitter in the changing environment of the host or in different individuals. In the presence of co-infection recombination occurs throughout the time-course of infection. The continued requirement for host-adaptation results in the *in vivo* selection for genes that have undergone recombination that favour persistence [157, 172, 284, 301]. Evidence for recombination is often found in genes related to virulence and persistence.

Analysis of a mixed infection from a Lithuanian patient found that recombination between strains resulted in the loss of the *cag* pathogenicity island and changes in alleles encoding VacA and outer membrane proteins [301]. Genome sequencing of *H. pylori* strains isolated sequentially from chronically infected individuals demonstrated that the number of clustered nucleotide polymorphisms (CNPs), attributable to genetic recombination with DNA from different strains during co-infection, was much greater in strains isolated later in infection (16 years) than in those isolated earlier (3 years), demonstrating that recombination continues through the time-course of infection. CNPs occurred more regularly in some gene families than others; most notably in genes encoding outer membrane proteins, particularly those of the *hop* family [157] whose proteins are known to play key roles in adhesion [24]. This indicates that diversifying selection acts *in vivo* on recombined genes [157]. Indeed, genetic evidence of low effective population sizes in isolates from South Africa may reflect population bottlenecks arising from selection pressures imposed by the host immune system [172].

Horizontal gene transfer by natural transformation in the presence of multiple strains is a key driver in the genetic diversity of *H. pylori*. Evidence has emerged that this generation of genetic diversity plays a role in the ability of *Helicobacter* to continually adapt to the host throughout prolonged infection. Recombination has the advantage of allowing multiple beneficial mutations to be combined within the same genetic background by horizontal transfer rather than by sequential mutation. This would allow more rapid fixation of beneficial alleles [164]. Indeed, evidence is also emerging that *H. pylori* experiences selective pressure for fitter variants by the host immune system.

### 7.1.2. Epigenetic diversity

Recently, SMRT sequencing has been utilized to sequence the methylomes of *H. pylori* 26695 and J99. Analysis of the methylome following genetic manipulation of candidate methylases has allowed characterisation of a number of methyltransferases in *H. pylori*. Interestingly, a methyltransferase (HP1353) has been identified that contains two phase-variable repeat sequences. One phase variable repeat appears to function canonically to regulate expression of the methyltransferase. Phase variation of the second repeat appears to switch the protein between two forms that vary in their methylation recognition sequence. The protein thus has three different phase-variable states; off, methylation of recognition sequence 1, and methylation of recognition sequence 2 [302]. It is tempting to speculate that this represents an additional layer of complexity to the phasevarion. Should this methyltransferase regulate the

transcription of other genes, it could, as a single methyltransferase, function to regulate two separate phasevarions.

Phase-variable methyltransferases have been identified in *H. pylori* and the investigation of their corresponding phasevarions has been initiated. The coordinated reversible switching of many genes within a phasevarion is postulated to play a role in host adaptation, immune evasion and pathogenicity. However, given the potential significance of the phasevarion, experimental evidence is lacking regarding the role of the phasevarion *in vivo* in host colonization, persistence, and disease pathology. In one study investigating the significance of methylation by the *iceA-hpyIM* R-M system, expression levels of *hpyIM* as determined from RT-PCR and/or RNA slot-blot analysis of RNA isolated from gastric biopsy specimens could not be correlated with *iceA* expression, disease sequelae, colonization density, or mucosal IL-8 levels. However, this methyltransferase has not been identified as having the potential to phase-vary and appears to be a stationary growth phase regulator [303].

Phase variation of individual genes and phasevarions impart to the bacterium an ability to rapidly and reversibly switch between a vast array of different phenotypes. It is postulated that for pathogenic bacteria, there is significant pressure to constantly avoid detection by the immune system. The ability to generate epigenetic and phenotypic variation in this way may enable a bacterial population to play a lottery against the human immune system in which at least the phenotype of one individual imparts sufficient fitness within a changing environment to allow persistence. Thus, both the continued characterisation of phasevarions and their *in vivo* relevance for host adaptation and virulence is of importance in our understanding of *H. pylori* persistence and pathogenesis.

### 7.1.3. *H. pylori* regulation of genetic diversity and evolution constraints?

The ability of *H. pylori* to generate genetic diversity is likely to be regulated so as to achieve a subtle balance between the generation of genetic diversity and maintaining the integrity of its small genome to establish persistent infection. The different mutation rates reported to date are in line with regulation of generation of genetic diversity in *H. pylori*. Phase variation of DNA glycosylase MutY, an enzyme of the base excision repair process, represents further evidence for regulation of genetic diversity [66, 67]. Alternatively, a constitutive high mutation rate (named the mutator phenotype in *E. coli*) could be counterbalanced by efficient mechanisms to maintain genome integrity, such as homologous recombination using DNA template from neighbouring cells by natural competence. This possibility has gained some support due to recent reports describing an increase of DNA uptake and RecA homologous recombination upon DNA damage [169] and poor persistence of the *dprA* mutant deficient in natural competence, although the initial colonization level was comparable to wild-type [246].

The ability of *H. pylori* to genetically adapt to the human host is remarkable based on adaptation to human subpopulations with 6 ancestral populations of *H. pylori* named ancestral European 1, ancestral European 2, ancestral East Asia, ancestral Africa1, ancestral Africa2 [2], and ancestral Sahul [1]. As described above, *H. pylori* genetic plasticity is the result of several mechanisms ranging from poor replication fidelity, lack of DNA repair genes and horizontal gene transfer promoted by natural competence, bacteriophage transduction, conjugation of



TnPZs and plasmid transfer. An extreme example of *H. pylori*'s adaptive ability is exemplified by the host jump from humans to large felines, predicted to have occurred some 200,000 years ago that resulted in a new species; *Helicobacter acinonychis*. This particular host jump was accompanied by relative conservation of the core genome, inactivation by different mechanisms of genes encoding surface proteins and acquisition of genes involved in sialylation of the bacterial surface to evade immune responses in the new host [210].

Altogether *H. pylori* micro-evolution within the human host and the above host jump conserved the core genome with changes affecting mainly hotspots and accessory genes. This suggests that constraints in the genome architecture and gene repertoire exist and these limit the evolutionary trajectories of *H. pylori*. Recent advances in evolutionary biology suggest a plurality of constraints on evolution, including the sequence type (coding sequence, structural RNAs, micro RNA and else) and genome architecture and gene repertoire underlying the sum of all phenotypic traits of the organism or phenome [304]. Thus to make sense of *H. pylori*'s high mutation and recombination rates and investigate the role of genetic diversity in phenotypic adaptation to the human host, it is important to investigate the constraints, their networks and interactions limiting *H. pylori* evolvability.

#### 7.1.4. Robustness and evolvability

Robustness of systems is the resistance to change under perturbation. Robustness, either mutational, or phenotypic, has been proposed to make biological systems more evolvable [305-308]. Studying *H. pylori* mutational and phenotypic robustness is a highly attractive approach to understand its adaptation to, and persistence in, the human host based on the latest findings and concepts of evolutionary biology.

The evolvability, or capacity of biological systems for adaptive evolution, was proposed to depend on their genotype-phenotype maps [309]. Evolutionary change takes place in a population with each population member having some genotype defining a collection of genotypes in the genotype space [308]. Through mutations, members of the population can change their location in the genotype space. In the population some individuals have a phenotype either superior, or inferior to the existing well-adapted phenotype. Natural selection, while eliminating poorly adapted individuals, preserves the well-adapted ones and selects superior ones. The genotype space reconciles the key problem of evolutionary adaptation of finding the rare superior genotype while preserving the population of well-adapted ones. A first characteristic of the genotype space is the existence of a set of genotypes with the same phenotype or connected genotype networks. The second characteristic of the genotype space is the number of genotypes that can be derived from any one genotype via mutation. The set of these genotypes is named neighbourhood genotype and its size is a simple measure of phenotypic variability of a genotype in response to mutation. In summary, the first feature of the genotype space allow individuals in a population to preserve their phenotype while changing their genotype and the neighbourhood genotype allows exploration of novel superior phenotypes in the population. Thus robustness is beneficial to both the individual and the population. As a consequence, robustness in the presence of genetic mutation/



recombination allows cryptic genetic diversity to accumulate (in the sense of a capacitor) and promotes evolutionary adaptation through greater phenotypic variability in the population.

Robustness as a variability principle in the context of the mutator phenotype of *H. pylori* (due to its high mutation and recombination rates) would reconcile the interest of the single cell and the cell population and may underlie *H. pylori* persistence in the gastric niche. A robust cell phenotype during persistent colonization of the gastric mucosa would not easily be disturbed by mutations, would be beneficial to single cells and would allow greater phenotypic variability to be achieved at the cell population level on a (micro-) evolutionary time scale by accumulation of cryptic genotypic diversity.

The presence of phasevarions in *H. pylori* raises the possibility to influence robustness by dramatically increasing phenotypic variability of the neighbourhood genotype by simple mutation. Greater robustness of *H. pylori* would enhance adaptation and persistence in the changing human gastric niche. Other interesting questions related to evolutionary dynamics are the size of the *H. pylori* population and its mutation rate and how these parameters influence robustness and adaptation to the host. For example, one could hypothesize that the lack of inter-strain recombination weakens the robustness of *H. pylori* leading to its disappearance in Western countries where multiple infections are rare.

To conclude, the investigation of *H. pylori* robustness to better understand the adaptation of this bacterium to the human host will require both computational modelling and experimental data on genotype-phenotype maps of *H. pylori* populations made accessible by the latest sequencing and high-content technologies.

#### 7.1.5. Refining *H. pylori* persistence mathematical model

*H. pylori* apparently lacks active and costly sensory machinery gene regulation and its small genome suggests alternative adaptive mechanisms including small RNA regulation [52], automatic random genetic switches for generating diverse adaptive phenotypes [47, 50, 51, 53], and the numerous duplicate and divergent outer membrane genes, which could be part of a more general gene regulation network so far unidentified. Introducing random fluctuations for stochastic phenotype transitions in the *H. pylori* mathematical model of persistence published by Blaser and Kirshner [44] is highly relevant to the robustness principle of biological systems and their evolvability. Thus further refinement of this model by introducing random fluctuations for stochastic phenotype transitions in the spirit of Kussell and Leibler [310], could help to further our understanding of how *H. pylori* establishes the optimal balance between sensing changes and random phenotype switching to adapt to its niche. Model predictions could be tested experimentally in animal models, using the power of molecular-bacterial genetics, including tetracycline-based gene regulation [275, 277], to validate or invalidate aspects of the model and gain insight into the dynamics of *H. pylori* populations.

## 7.2. Therapeutic potential of targeting persistence mechanisms

*H. pylori* chronic infection remains a significant health burden, a long regimen of triple or quadruple antibiotic therapy is currently the only available treatment and antibiotic resistance

is emerging [311]. Since no prophylactic or therapeutic vaccine has been successfully developed to date, new target ideas are needed for the development of innovative drug therapies to achieve *H. pylori* eradication. Targeting *H. pylori* persistence mechanisms is one such strategy that could encompass immune evasion, micro RNA regulation (of urease for example [52]), genetic and phenotypic diversity generation underlying host adaptation and robustness of biological systems. In this regard, phasevarions, recombination (XerH for example [312]), DNA repair, replication (low fidelity Polymerase I for example [48]) and competence would represent target pathways of choice. Also a drug targeting discrete sites on the urease complex surface shown to be required for infection [247] would have several advantages over eliciting site-specific urease antibodies to inhibit *H. pylori* persistence; e.g. better bioavailability in the mucus layer and a higher concentration that is more likely to achieve inhibition of the large amount of urease produced by *H. pylori*. Other potential targets are the UreI channel (whose structure is available for drug design [313]) that is required for buffering the periplasm during colonization [255-257], the gamma-glutamyltransferase required for immune modulation and bacterial metabolism to maintain colonization [36], *H. pylori* LPS biosynthetic proteins contributing to gastric adaptation, adhesion and modulation of the host immune response by changing antigen expression in the same human host over time [314]. Compounds targeting the biosynthesis of glycolipids that *H. pylori* uses for immune evasion showed promising inhibition *in vitro* and further work is required to assess their potential [315]. With the recent adaptation of the tetracycline-based gene regulation system in *H. pylori* and its functionality *in vivo* [275, 277], potential targets can now be validated in animal models using tetracycline-based conditional knockouts. Furthermore, conditional knockouts will be instrumental in testing the rate of emergence of resistance for target candidates to select targets less prone to genetic adaptation before embarking on a long and costly drug discovery and development path. *H. pylori*-specific drugs targeting persistence mechanisms would have the advantage of leaving the gastrointestinal microbiota intact, avoiding side effects such as diarrhoea and increased patient compliance. Based on the use of triple and quadruple antibiotic therapies and the genetic plasticity of *H. pylori*, it is likely that more than one *H. pylori*-specific drug targeting persistence will be required to avoid emergence of resistance and achieve high eradication rate.

## Acknowledgements

This work was supported by a NHMRC Sir McFarlane Burnett Fellowship grant (572723) to BJM, NHMRC Project Grant (634465) to BJM and MB and NHMRC Early Career Fellowship grant to AWD (1073250). We thank the team at Transitttranslations who proofread the manuscript.

## Author details

Mohammed Benghezal\*, Jonathan C. Gauntlett, Aleksandra W. Debowski, Alma Fulurija, Hans-Olof Nilsson and Barry James Marshall

\*Address all correspondence to: [mohammed.benghezal@uwa.edu.au](mailto:mohammed.benghezal@uwa.edu.au)

Ondek Pty Ltd. and *Helicobacter pylori* Research Laboratory, Marshall Centre for Infectious Disease Research and Training, The University of Western Australia, School of Pathology & Laboratory Medicine, UWA Discipline of Microbiology & Immunology, Crawley WA, Australia

## References

- [1] Moodley Y, Linz B, Yamaoka Y, Windsor HM, Breurec S, et al. (2009) The peopling of the Pacific from a bacterial perspective. *Science* 323: 527-530.
- [2] Falush D, Wirth T, Linz B, Pritchard JK, Stephens M, et al. (2003) Traces of human migrations in *Helicobacter pylori* populations. *Science* 299: 1582-1585.
- [3] Dunn BE, Vakil NB, Schneider BG, Miller MM, Zitzer JB, et al. (1997) Localization of *Helicobacter pylori* urease and heat shock protein in human gastric biopsies. *Infect Immun* 65: 1181-1188.
- [4] Suerbaum S, Michetti P (2002) *Helicobacter pylori* infection. *N Engl J Med* 347: 1175-1186.
- [5] Blaser MJ, Atherton JC (2004) *Helicobacter pylori* persistence: biology and disease. *J Clin Invest* 113: 321-333.
- [6] Monack DM, Mueller A, Falkow S (2004) Persistent bacterial infections: the interface of the pathogen and the host immune system. *Nat Rev Microbiol* 2: 747-765.
- [7] Algood HM, Cover TL (2006) *Helicobacter pylori* persistence: an overview of interactions between *H. pylori* and host immune defenses. *Clin Microbiol Rev* 19: 597-613.
- [8] Fischer W, Prassl S, Haas R (2009) Virulence mechanisms and persistence strategies of the human gastric pathogen *Helicobacter pylori*. *Curr Top Microbiol Immunol* 337: 129-171.
- [9] Muller A, Solnick JV (2011) Inflammation, immunity, and vaccine development for *Helicobacter pylori*. *Helicobacter* 16 Suppl 1: 26-32.
- [10] Ricci V, Romano M, Boquet P (2011) Molecular cross-talk between *Helicobacter pylori* and human gastric mucosa. *World J Gastroenterol* 17: 1383-1399.

- [11] Kabir S (2011) The role of interleukin-17 in the *Helicobacter pylori* induced infection and immunity. *Helicobacter* 16: 1-8.
- [12] Sundquist M, Quiding-Jarbrink M (2010) *Helicobacter pylori* and its effect on innate and adaptive immunity: new insights and vaccination strategies. *Expert Rev Gastroenterol Hepatol* 4: 733-744.
- [13] Molnar B, Galamb O, Sipos F, Leiszter K, Tulassay Z (2010) Molecular pathogenesis of *Helicobacter pylori* infection: the role of bacterial virulence factors. *Dig Dis* 28: 604-608.
- [14] Allison CC, Ferrero RL (2010) Role of virulence factors and host cell signaling in the recognition of *Helicobacter pylori* and the generation of immune responses. *Future Microbiol* 5: 1233-1255.
- [15] Peek RM, Jr., Fiske C, Wilson KT (2010) Role of innate immunity in *Helicobacter pylori*-induced gastric malignancy. *Physiol Rev* 90: 831-858.
- [16] Watanabe T, Asano N, Kitani A, Fuss IJ, Chiba T, et al. (2011) Activation of type I IFN signaling by NOD1 mediates mucosal host defense against *Helicobacter pylori* infection. *Gut Microbes* 2: 61-65.
- [17] Atherton JC, Blaser MJ (2009) Coadaptation of *Helicobacter pylori* and humans: ancient history, modern implications. *J Clin Invest* 119: 2475-2487.
- [18] Kobayashi M, Lee H, Nakayama J, Fukuda M (2009) Roles of gastric mucin-type O-glycans in the pathogenesis of *Helicobacter pylori* infection. *Glycobiology* 19: 453-461.
- [19] O'Keefe J, Moran AP (2008) Conventional, regulatory, and unconventional T cells in the immunologic response to *Helicobacter pylori*. *Helicobacter* 13: 1-19.
- [20] D'Elia MM, Andersen LP (2007) *Helicobacter pylori* inflammation, immunity, and vaccines. *Helicobacter* 12 Suppl 1: 15-19.
- [21] Robinson K, Argent RH, Atherton JC (2007) The inflammatory and immune response to *Helicobacter pylori* infection. *Best Pract Res Clin Gastroenterol* 21: 237-259.
- [22] Suarez G, Reyes VE, Beswick EJ (2006) Immune response to *H. pylori*. *World J Gastroenterol* 12: 5593-5598.
- [23] Velin D, Michetti P (2006) Immunology of *Helicobacter pylori* infection. *Digestion* 73: 116-123.
- [24] Salama NR, Hartung ML, Muller A (2013) Life in the human stomach: persistence strategies of the bacterial pathogen *Helicobacter pylori*. *Nat Rev Microbiol* 11: 385-399.
- [25] Yokota S, Okabayashi T, Rehli M, Fujii N, Amano K (2010) *Helicobacter pylori* lipopolysaccharides upregulate toll-like receptor 4 expression and proliferation of gastric



epithelial cells via the MEK1/2-ERK1/2 mitogen-activated protein kinase pathway. *Infect Immun* 78: 468-476.

- [26] Cullen TW, Giles DK, Wolf LN, Ecobichon C, Boneca IG, et al. (2011) *Helicobacter pylori* versus the host: remodeling of the bacterial outer membrane is required for survival in the gastric mucosa. *PLoS Pathog* 7: e1002454.
- [27] Gringhuis SI, den Dunnen J, Litjens M, van der Vlist M, Geijtenbeek TB (2009) Carbohydrate-specific signaling through the DC-SIGN signalosome tailors immunity to *Mycobacterium tuberculosis*, HIV-1 and *Helicobacter pylori*. *Nat Immunol* 10: 1081-1088.
- [28] Necchi V, Manca R, Ricci V, Solcia E (2009) Evidence for transepithelial dendritic cells in human *H. pylori* active gastritis. *Helicobacter* 14: 208-222.
- [29] Bergman MP, Engering A, Smits HH, van Vliet SJ, van Bodegraven AA, et al. (2004) *Helicobacter pylori* modulates the T helper cell 1/T helper cell 2 balance through phase-variable interaction between lipopolysaccharide and DC-SIGN. *J Exp Med* 200: 979-990.
- [30] Appelmek BJ, van Die I, van Vliet SJ, Vandenbroucke-Grauls CM, Geijtenbeek TB, et al. (2003) Cutting edge: carbohydrate profiling identifies new pathogens that interact with dendritic cell-specific ICAM-3-grabbing nonintegrin on dendritic cells. *J Immunol* 170: 1635-1639.
- [31] Moran AP (2008) Relevance of fucosylation and Lewis antigen expression in the bacterial gastroduodenal pathogen *Helicobacter pylori*. *Carbohydr Res* 343: 1952-1965.
- [32] Aspinall GO, Monteiro MA, Pang H, Walsh EJ, Moran AP (1994) O Antigen chains in the lipopolysaccharide of *H. pylori* NCTC 11637. O Antigen chains in the lipopolysaccharide of *H. pylori* NCTC 11637 1: 151-156.
- [33] Andersen-Nissen E, Smith KD, Strobe KL, Barrett SL, Cookson BT, et al. (2005) Evasion of Toll-like receptor 5 by flagellated bacteria. *Proc Natl Acad Sci U S A* 102: 9247-9252.
- [34] Gewirtz AT, Yu Y, Krishna US, Israel DA, Lyons SL, et al. (2004) *Helicobacter pylori* flagellin evades toll-like receptor 5-mediated innate immunity. *J Infect Dis* 189: 1914-1920.
- [35] Lee SK, Stack A, Katzowitsch E, Aizawa SI, Suerbaum S, et al. (2003) *Helicobacter pylori* flagellins have very low intrinsic activity to stimulate human gastric epithelial cells via TLR5. *Microbes Infect* 5: 1345-1356.
- [36] Schmees C, Prinz C, Treptau T, Rad R, Hengst L, et al. (2007) Inhibition of T-cell proliferation by *Helicobacter pylori* gamma-glutamyl transpeptidase. *Gastroenterology* 132: 1820-1833.

- [37] Gebert B, Fischer W, Weiss E, Hoffmann R, Haas R (2003) *Helicobacter pylori* vacuolating cytotoxin inhibits T lymphocyte activation. *Science* 301: 1099-1102.
- [38] Wunder C, Churin Y, Winau F, Warnecke D, Vieth M, et al. (2006) Cholesterol glucosylation promotes immune evasion by *Helicobacter pylori*. *Nat Med* 12: 1030-1038.
- [39] Hitzler I, Sayi A, Kohler E, Engler DB, Koch KN, et al. (2012) Caspase-1 Has Both Pro-Inflammatory and Regulatory Properties in *Helicobacter* Infections, Which Are Differentially Mediated by Its Substrates IL-1 $\beta$  and IL-18. *J Immunol*.
- [40] Takashima M, Furuta T, Hanai H, Sugimura H, Kaneko E (2001) Effects of *Helicobacter pylori* infection on gastric acid secretion and serum gastrin levels in Mongolian gerbils. *Gut* 48: 765-773.
- [41] Troost E, Hold GL, Smith MG, Chow WH, Rabkin CS, et al. (2003) The role of interleukin-1 $\beta$  and other potential genetic markers as indicators of gastric cancer risk. *Can J Gastroenterol* 17 Suppl B: 8B-12B.
- [42] El-Omar EM, Rabkin CS, Gammon MD, Vaughan TL, Risch HA, et al. (2003) Increased risk of noncardia gastric cancer associated with proinflammatory cytokine gene polymorphisms. *Gastroenterology* 124: 1193-1201.
- [43] El-Omar EM, Carrington M, Chow WH, McColl KE, Bream JH, et al. (2000) Interleukin-1 polymorphisms associated with increased risk of gastric cancer. *Nature* 404: 398-402.
- [44] Blaser MJ, Kirschner D (2007) The equilibria that allow bacterial persistence in human hosts. *Nature* 449: 843-849.
- [45] Bjorkholm B, Sjolund M, Falk PG, Berg OG, Engstrand L, et al. (2001) Mutation frequency and biological cost of antibiotic resistance in *Helicobacter pylori*. *Proc Natl Acad Sci U S A* 98: 14607-14612.
- [46] Kang J, Blaser MJ (2006) Bacterial populations as perfect gases: genomic integrity and diversification tensions in *Helicobacter pylori*. *Nat Rev Microbiol* 4: 826-836.
- [47] Salaun L, Linz B, Suerbaum S, Saunders NJ (2004) The diversity within an expanded and redefined repertoire of phase-variable genes in *Helicobacter pylori*. *Microbiology* 150: 817-830.
- [48] Garcia-Ortiz MV, Marsin S, Arana ME, Gasparutto D, Guerois R, et al. (2011) Unexpected role for *Helicobacter pylori* DNA polymerase I as a source of genetic variability. *PLoS Genet* 7: e1002152.
- [49] Moccia C, Krebs J, Kulick S, Didelot X, Kraft C, et al. (2012) The nucleotide excision repair (NER) system of *Helicobacter pylori*: role in mutation prevention and chromosomal import patterns after natural transformation. *BMC Microbiol* 12: 67.
- [50] Srikhanta YN, Gorrell RJ, Steen JA, Gawthorne JA, Kwok T, et al. (2011) Phasevarion mediated epigenetic gene regulation in *Helicobacter pylori*. *PLoS One* 6: e27569.

- [51] Srikhanta YN, Maguire TL, Stacey KJ, Grimmond SM, Jennings MP (2005) The phase-varion: a genetic system controlling coordinated, random switching of expression of multiple genes. *Proc Natl Acad Sci U S A* 102: 5547-5551.
- [52] Wen Y, Feng J, Sachs G (2013) *Helicobacter pylori* 5'ureB-sRNA, a cis-encoded anti-sense small RNA, negatively regulates ureAB expression by transcription termination. *J Bacteriol* 195: 444-452.
- [53] Palmer ME, Lipsitch M, Moxon ER, Bayliss CD (2013) Broad conditions favor the evolution of phase-variable loci. *MBio* 4: e00430-00412.
- [54] Ambur OH, Davidsen T, Frye SA, Balasingham SV, Lagesen K, et al. (2009) Genome dynamics in major bacterial pathogens. *FEMS Microbiol Rev* 33: 453-470.
- [55] McCready SJ, Osman F, Yasui A (2000) Repair of UV damage in the fission yeast *Schizosaccharomyces pombe*. *Mutat Res* 451: 197-210.
- [56] Alm RA, Trust TJ (1999) Analysis of the genetic diversity of *Helicobacter pylori*: the tale of two genomes. *J Mol Med (Berl)* 77: 834-846.
- [57] Tomb JF, White O, Kerlavage AR, Clayton RA, Sutton GG, et al. (1997) The complete genome sequence of the gastric pathogen *Helicobacter pylori*. *Nature* 388: 539-547.
- [58] Eisen JA (1998) A phylogenomic study of the MutS family of proteins. *Nucleic Acids Res* 26: 4291-4300.
- [59] Wang G, Alamuri P, Humayun MZ, Taylor DE, Maier RJ (2005) The *Helicobacter pylori* MutS protein confers protection from oxidative DNA damage. *Mol Microbiol* 58: 166-176.
- [60] LeClerc JE, Li B, Payne WL, Cebula TA (1996) High mutation frequencies among *Escherichia coli* and *Salmonella* pathogens. *Science* 274: 1208-1211.
- [61] Oliver A, Canton R, Campo P, Baquero F, Blazquez J (2000) High frequency of hypermutable *Pseudomonas aeruginosa* in cystic fibrosis lung infection. *Science* 288: 1251-1254.
- [62] Richardson AR, Yu Z, Popovic T, Stojiljkovic I (2002) Mutator clones of *Neisseria meningitidis* in epidemic serogroup A disease. *Proc Natl Acad Sci U S A* 99: 6103-6107.
- [63] Dorer MS, Sessler TH, Salama NR (2011) Recombination and DNA repair in *Helicobacter pylori*. *Annu Rev Microbiol* 65: 329-348.
- [64] Eutsey R, Wang G, Maier RJ (2007) Role of a MutY DNA glycosylase in combating oxidative DNA damage in *Helicobacter pylori*. *DNA Repair (Amst)* 6: 19-26.
- [65] Huang S, Kang J, Blaser MJ (2006) Antimutator role of the DNA glycosylase mutY gene in *Helicobacter pylori*. *J Bacteriol* 188: 6224-6234.

- [66] Kulick S, Moccia C, Kraft C, Suerbaum S (2008) The *Helicobacter pylori* mutY homologue HP0142 is an antimutator gene that prevents specific C to A transversions. *Arch Microbiol* 189: 263-270.
- [67] Mathieu A, O'Rourke EJ, Radicella JP (2006) *Helicobacter pylori* genes involved in avoidance of mutations induced by 8-oxoguanine. *J Bacteriol* 188: 7464-7469.
- [68] Eichman BF, O'Rourke EJ, Radicella JP, Ellenberger T (2003) Crystal structures of 3-methyladenine DNA glycosylase MagIII and the recognition of alkylated bases. *EMBO J* 22: 4898-4909.
- [69] O'Rourke EJ, Chevalier C, Boiteux S, Labigne A, Ielpi L, et al. (2000) A novel 3-methyladenine DNA glycosylase from *Helicobacter pylori* defines a new class within the endonuclease III family of base excision repair glycosylases. *J Biol Chem* 275: 20077-20083.
- [70] Baldwin DN, Shepherd B, Kraemer P, Hall MK, Sycuro LK, et al. (2007) Identification of *Helicobacter pylori* genes that contribute to stomach colonization. *Infect Immun* 75: 1005-1016.
- [71] O'Rourke EJ, Chevalier C, Pinto AV, Thiberge JM, Ielpi L, et al. (2003) Pathogen DNA as target for host-generated oxidative stress: role for repair of bacterial DNA damage in *Helicobacter pylori* colonization. *Proc Natl Acad Sci U S A* 100: 2789-2794.
- [72] Thompson SA, Latch RL, Blaser JM (1998) Molecular characterization of the *Helicobacter pylori* uvr B gene. *Gene* 209: 113-122.
- [73] Kang J, Blaser MJ (2006) UvrD helicase suppresses recombination and DNA damage-induced deletions. *J Bacteriol* 188: 5450-5459.
- [74] Lee GH, Jeong JY, Chung JW, Nam WH, Lee SM, et al. (2009) The *Helicobacter pylori* Mfd protein is important for antibiotic resistance and DNA repair. *Diagn Microbiol Infect Dis* 65: 454-456.
- [75] Solnick JV, Hansen LM, Salama NR, Boonjakuakul JK, Syvanen M (2004) Modification of *Helicobacter pylori* outer membrane protein expression during experimental infection of rhesus macaques. *Proc Natl Acad Sci U S A* 101: 2106-2111.
- [76] Amundsen SK, Fero J, Hansen LM, Cromie GA, Solnick JV, et al. (2008) *Helicobacter pylori* AddAB helicase-nuclease and RecA promote recombination-related DNA repair and survival during stomach colonization. *Mol Microbiol* 69: 994-1007.
- [77] Oleastro M, Cordeiro R, Menard A, Gomes JP (2010) Allelic diversity among *Helicobacter pylori* outer membrane protein genes homB and homA generated by recombination. *J Bacteriol* 192: 3961-3968.
- [78] Pohl MA, Kienesberger S, Blaser MJ (2012) Novel functions for glycosyltransferases Jhp0562 and GalT in Lewis antigen synthesis and variation in *Helicobacter pylori*. *Infect Immun* 80: 1593-1605.



- [79] Talarico S, Whitefield SE, Fero J, Haas R, Salama NR (2012) Regulation of *Helicobacter pylori* adherence by gene conversion. *Mol Microbiol* 84: 1050-1061.
- [80] Schmitt W, Odenbreit S, Heuermann D, Haas R (1995) Cloning of the *Helicobacter pylori* recA gene and functional characterization of its product. *Mol Gen Genet* 248: 563-572.
- [81] Thompson SA, Blaser MJ (1995) Isolation of the *Helicobacter pylori* recA gene and involvement of the recA region in resistance to low pH. *Infect Immun* 63: 2185-2193.
- [82] Amundsen SK, Fero J, Salama NR, Smith GR (2009) Dual nuclease and helicase activities of *Helicobacter pylori* AddAB are required for DNA repair, recombination, and mouse infectivity. *J Biol Chem* 284: 16759-16766.
- [83] Marsin S, Mathieu A, Kortulewski T, Guerois R, Radicella JP (2008) Unveiling novel RecO distant orthologues involved in homologous recombination. *PLoS Genet* 4: e1000146.
- [84] Marsin S, Lopes A, Mathieu A, Dizet E, Orillard E, et al. (2010) Genetic dissection of *Helicobacter pylori* AddAB role in homologous recombination. *FEMS Microbiol Lett* 311: 44-50.
- [85] Wang G, Maier RJ (2008) Critical role of RecN in recombinational DNA repair and survival of *Helicobacter pylori*. *Infect Immun* 76: 153-160.
- [86] Kang J, Blaser MJ (2008) Repair and antirepair DNA helicases in *Helicobacter pylori*. *J Bacteriol* 190: 4218-4224.
- [87] Debowski AW, Carnoy C, Verbrugghe P, Nilsson H, Gauntlett JC, et al. (2012) Xer Recombinase and Genome Integrity in *Helicobacter pylori*, a Pathogen without Topoisomerase IV. *PLoS One* in press.
- [88] Robinson K, Loughlin MF, Potter R, Jenks PJ (2005) Host adaptation and immune modulation are mediated by homologous recombination in *Helicobacter pylori*. *J Infect Dis* 191: 579-587.
- [89] Fukui K, Nakagawa N, Kitamura Y, Nishida Y, Masui R, et al. (2008) Crystal structure of MutS2 endonuclease domain and the mechanism of homologous recombination suppression. *J Biol Chem* 283: 33417-33427.
- [90] Pinto AV, Mathieu A, Marsin S, Veaute X, Ielpi L, et al. (2005) Suppression of homologous and homeologous recombination by the bacterial MutS2 protein. *Mol Cell* 17: 113-120.
- [91] Kuempel PL, Henson JM, Dircks L, Tecklenburg M, Lim DF (1991) dif, a recA-independent recombination site in the terminus region of the chromosome of *Escherichia coli*. *New Biol* 3: 799-811.

- [92] Aras RA, Fischer W, Perez-Perez GI, Crosatti M, Ando T, et al. (2003) Plasticity of repetitive DNA sequences within a bacterial (Type IV) secretion system component. *J Exp Med* 198: 1349-1360.
- [93] Barrozo RM, Cooke CL, Hansen LM, Lam AM, Gaddy JA, et al. (2013) Functional plasticity in the type IV secretion system of *Helicobacter pylori*. *PLoS Pathog* 9: e1003189.
- [94] van der Woude MW, Baumler AJ (2004) Phase and antigenic variation in bacteria. *Clin Microbiol Rev* 17: 581-611, table of contents.
- [95] Salaun L, Snyder LA, Saunders NJ (2003) Adaptation by phase variation in pathogenic bacteria. *Adv Appl Microbiol* 52: 263-301.
- [96] Guo X, Mrazek J (2008) Long simple sequence repeats in host-adapted pathogens localize near genes encoding antigens, housekeeping genes, and pseudogenes. *J Mol Evol* 67: 497-509.
- [97] Alm RA, Ling LS, Moir DT, King BL, Brown ED, et al. (1999) Genomic-sequence comparison of two unrelated isolates of the human gastric pathogen *Helicobacter pylori*. *Nature* 397: 176-180.
- [98] Saunders NJ, Peden JF, Hood DW, Moxon ER (1998) Simple sequence repeats in the *Helicobacter pylori* genome. *Mol Microbiol* 27: 1091-1098.
- [99] Salaun L, Linz B, Suerbaum S, Saunders NJ (2004) The diversity within an expanded and redefined repertoire of phase-variable genes in *Helicobacter pylori*. *Microbiology-Sgm* 150: 817-830.
- [100] Salaun L, Saunders NJ (2006) Population-associated differences between the phase variable LPS biosynthetic genes of *Helicobacter pylori*. *BMC Microbiol* 6: 79.
- [101] Moran AP (1996) *Helicobacter pylori* expresses Lewis X. *Helicobacter* 1: 190-191.
- [102] Appelmelk BJ, Shiberu B, Trinkis C, Tapsi N, Zheng PY, et al. (1998) Phase variation in *Helicobacter pylori* lipopolysaccharide. *Infect Immun* 66: 70-76.
- [103] Appelmelk BJ, Martin SL, Monteiro MA, Clayton CA, McColm AA, et al. (1999) Phase variation in *Helicobacter pylori* lipopolysaccharide due to changes in the lengths of poly(C) tracts in alpha3-fucosyltransferase genes. *Infect Immun* 67: 5361-5366.
- [104] Nilsson C, Skoglund A, Moran AP, Annuk H, Engstrand L, et al. (2008) Lipopolysaccharide diversity evolving in *Helicobacter pylori* communities through genetic modifications in fucosyltransferases. *PLoS One* 3: e3811.
- [105] Appelmelk BJ, Martino MC, Veenhof E, Monteiro MA, Maaskant JJ, et al. (2000) Phase variation in H type I and Lewis a epitopes of *Helicobacter pylori* lipopolysaccharide. *Infect Immun* 68: 5928-5932.

- [106] Langdon R, Craig JE, Goldrick M, Houldsworth R, High NJ (2005) Analysis of the role of HP0208, a phase-variable open reading frame, and its homologues HP1416 and HP0159 in the biosynthesis of *Helicobacter pylori* lipopolysaccharide. *J Med Microbiol* 54: 697-706.
- [107] Sanabria-Valentin E, Colbert MT, Blaser MJ (2007) Role of futC slipped strand mispairing in *Helicobacter pylori* Lewisy phase variation. *Microbes Infect* 9: 1553-1560.
- [108] Wirth HP, Yang M, Peek RM, Jr., Hook-Nikanne J, Fried M, et al. (1999) Phenotypic diversity in Lewis expression of *Helicobacter pylori* isolates from the same host. *J Lab Clin Med* 133: 488-500.
- [109] Linden S, Boren T, Dubois A, Carlstedt I (2004) Rhesus monkey gastric mucins: oligomeric structure, glycoforms and *Helicobacter pylori* binding. *Biochem J* 379: 765-775.
- [110] Goodwin AC, Weinberger DM, Ford CB, Nelson JC, Snider JD, et al. (2008) Expression of the *Helicobacter pylori* adhesin SabA is controlled via phase variation and the ArsRS signal transduction system. *Microbiology* 154: 2231-2240.
- [111] Peck B, Ortkamp M, Diehl KD, Hundt E, Knapp B (1999) Conservation, localization and expression of HopZ, a protein involved in adhesion of *Helicobacter pylori*. *Nucleic Acids Res* 27: 3325-3333.
- [112] Kennemann L, Brenneke B, Andres S, Engstrand L, Meyer TF, et al. (2012) In vivo sequence variation in HopZ, a phase-variable outer membrane protein of *Helicobacter pylori*. *Infect Immun* 80: 4364-4373.
- [113] Ilver D, Arnqvist A, Ogren J, Frick IM, Kersulyte D, et al. (1998) *Helicobacter pylori* adhesin binding fucosylated histo-blood group antigens revealed by retagging. *Science* 279: 373-377.
- [114] Dossumbekova A, Prinz C, Mages J, Lang R, Kusters JG, et al. (2006) *Helicobacter pylori* HopH (OipA) and bacterial pathogenicity: genetic and functional genomic analysis of hopH gene polymorphisms. *J Infect Dis* 194: 1346-1355.
- [115] Yamaoka Y, Kita M, Kodama T, Imamura S, Ohno T, et al. (2002) *Helicobacter pylori* infection in mice: Role of outer membrane proteins in colonization and inflammation. *Gastroenterology* 123: 1992-2004.
- [116] Yamaoka Y, Kikuchi S, el-Zimaity HM, Gutierrez O, Osato MS, et al. (2002) Importance of *Helicobacter pylori* oipA in clinical presentation, gastric inflammation, and mucosal interleukin 8 production. *Gastroenterology* 123: 414-424.
- [117] Tannaes T, Dekker N, Bukholm G, Bijlsma JJ, Appelmek BJ (2001) Phase variation in the *Helicobacter pylori* phospholipase A gene and its role in acid adaptation. *Infect Immun* 69: 7334-7340.
- [118] Moran AP, Knirel YA, Senchenkova SN, Widmalm G, Hynes SO, et al. (2002) Phenotypic variation in molecular mimicry between *Helicobacter pylori* lipopolysaccharides and human gastric epithelial cell surface glycoforms. Acid-induced phase

variation in Lewis(x) and Lewis(y) expression by *H. Pylori* lipopolysaccharides. *J Biol Chem* 277: 5785-5795.

- [119] Dumrese C, Slomianka L, Ziegler U, Choi SS, Kalia A, et al. (2009) The secreted *Helicobacter* cysteine-rich protein A causes adherence of human monocytes and differentiation into a macrophage-like phenotype. *FEBS Lett* 583: 1637-1643.
- [120] Khamri W, Moran AP, Worku ML, Karim QN, Walker MM, et al. (2005) Variations in *Helicobacter pylori* lipopolysaccharide to evade the innate immune component surfactant protein D. *Infect Immun* 73: 7677-7686.
- [121] Bergman M, Del Prete G, van Kooyk Y, Appelmelk B (2006) *Helicobacter pylori* phase variation, immune modulation and gastric autoimmunity. *Nat Rev Microbiol* 4: 151-159.
- [122] Salaun L, Ayraud S, Saunders NJ (2005) Phase variation mediated niche adaptation during prolonged experimental murine infection with *Helicobacter pylori*. *Microbiology* 151: 917-923.
- [123] Styer CM, Hansen LM, Cooke CL, Gundersen AM, Choi SS, et al. (2010) Expression of the BabA adhesin during experimental infection with *Helicobacter pylori*. *Infect Immun* 78: 1593-1600.
- [124] Weiser JN, Maskell DJ, Butler PD, Lindberg AA, Moxon ER (1990) Characterization of repetitive sequences controlling phase variation of *Haemophilus influenzae* lipopolysaccharide. *J Bacteriol* 172: 3304-3309.
- [125] Boyer HW (1971) DNA restriction and modification mechanisms in bacteria. *Annu Rev Microbiol* 25: 153-176.
- [126] Wilson GG, Murray NE (1991) Restriction and modification systems. *Annu Rev Genet* 25: 585-627.
- [127] Low DA, Weyand NJ, Mahan MJ (2001) Roles of DNA adenine methylation in regulating bacterial gene expression and virulence. *Infect Immun* 69: 7197-7204.
- [128] Srikhanta YN, Fox KL, Jennings MP (2010) The phasevarion: phase variation of type III DNA methyltransferases controls coordinated switching in multiple genes. *Nat Rev Microbiol* 8: 196-206.
- [129] Srikhanta YN, Dowideit SJ, Edwards JL, Falsetta ML, Wu HJ, et al. (2009) Phasevarions mediate random switching of gene expression in pathogenic *Neisseria*. *PLoS Pathog* 5: e1000400.
- [130] Fox KL, Srikhanta YN, Jennings MP (2007) Phase variable type III restriction-modification systems of host-adapted bacterial pathogens. *Mol Microbiol* 65: 1375-1379.
- [131] Akopyants NS, Fradkov A, Diatchenko L, Hill JE, Siebert PD, et al. (1998) PCR-based subtractive hybridization and differences in gene content among strains of *Helicobacter pylori*. *Proc Natl Acad Sci U S A* 95: 13108-13113.



- [132] Lin LF, Posfai J, Roberts RJ, Kong H (2001) Comparative genomics of the restriction-modification systems in *Helicobacter pylori*. *Proc Natl Acad Sci U S A* 98: 2740-2745.
- [133] Furuta Y, Kobayashi I (2012) Mobility of DNA sequence recognition domains in DNA methyltransferases suggests epigenetics-driven adaptive evolution. *Mob Genet Elements* 2: 292-296.
- [134] Gawthorne JA, Beatson SA, Srikhanta YN, Fox KL, Jennings MP (2012) Origin of the diversity in DNA recognition domains in phasevarion associated *modA* genes of pathogenic *Neisseria* and *Haemophilus influenzae*. *PLoS One* 7: e32337.
- [135] Kong H, Lin LF, Porter N, Stickel S, Byrd D, et al. (2000) Functional analysis of putative restriction-modification system genes in the *Helicobacter pylori* J99 genome. *Nucleic Acids Res* 28: 3216-3223.
- [136] de Vries N, Duinsbergen D, Kuipers EJ, Pot RG, Wiesenekker P, et al. (2002) Transcriptional phase variation of a type III restriction-modification system in *Helicobacter pylori*. *J Bacteriol* 184: 6615-6623.
- [137] Skoglund A, Bjorkholm B, Nilsson C, Andersson AF, Jernberg C, et al. (2007) Functional analysis of the M.HpyAIV DNA methyltransferase of *Helicobacter pylori*. *J Bacteriol* 189: 8914-8921.
- [138] Kumar R, Mukhopadhyay AK, Ghosh P, Rao DN (2012) Comparative transcriptomics of *H. pylori* strains AM5, SS1 and their hpyAVIBM deletion mutants: possible roles of cytosine methylation. *PLoS One* 7: e42303.
- [139] Nedenskov-Sørensen P, Bukholm G, Bøvre K (1990) Natural competence for genetic transformation in *Campylobacter pylori*. *J Infect Dis* 161: 365-366.
- [140] Bakkali M (2013) Could DNA uptake be a side effect of bacterial adhesion and twitching motility? *Arch Microbiol* 195: 279-289.
- [141] Seitz P, Blokesch M (2013) Cues and regulatory pathways involved in natural competence and transformation in pathogenic and environmental Gram-negative bacteria. *FEMS Microbiol Rev* 37: 336-363.
- [142] Hofreuter D, Odenbreit S, Haas R (2001) Natural transformation competence in *Helicobacter pylori* is mediated by the basic components of a type IV secretion system. *Mol Microbiol* 41: 379-391.
- [143] Stingl K, Muller S, Scheidgen-Kleyboldt G, Clausen M, Maier B (2010) Composite system mediates two-step DNA uptake into *Helicobacter pylori*. *Proc Natl Acad Sci U S A* 107: 1184-1189.
- [144] Yeh YC, Lin TL, Chang KC, Wang JT (2003) Characterization of a *ComE3* homologue essential for DNA transformation in *Helicobacter pylori*. *Infect Immun* 71: 5427-5431.

- [145] Humbert O, Dorer MS, Salama NR (2011) Characterization of *Helicobacter pylori* factors that control transformation frequency and integration length during inter-strain DNA recombination. *Mol Microbiol* 79: 387-401.
- [146] Mortier-Barrière I, Velten M, Dupaigne P, Mirouze N, Piètrement O, et al. (2007) A key presynaptic role in transformation for a widespread bacterial protein: DprA conveys incoming ssDNA to RecA. *Cell* 130: 824-836.
- [147] Kang J, Huang S, Blaser MJ (2005) Structural and functional divergence of MutS2 from bacterial MutS1 and eukaryotic MSH4-MSH5 homologs. *J Bacteriol* 187: 3528-3537.
- [148] Zhang XS, Blaser MJ (2012) DprB facilitates inter- and intragenomic recombination in *Helicobacter pylori*. *J Bacteriol* 194: 3891-3903.
- [149] Smeets LC, Kusters JG (2002) Natural transformation in *Helicobacter pylori*: DNA transport in an unexpected way. *Trends Microbiol* 10: 159-162; discussion 162.
- [150] Hofreuter D, Odenbreit S, Henke G, Haas R (1998) Natural competence for DNA transformation in *Helicobacter pylori*: identification and genetic characterization of the comB locus. *Mol Microbiol* 28: 1027-1038.
- [151] Karnholz A, Hoefler C, Odenbreit S, Fischer W, Hofreuter D, et al. (2006) Functional and topological characterization of novel components of the comB DNA transformation competence system in *Helicobacter pylori*. *J Bacteriol* 188: 882-893.
- [152] Lin TL, Shun CT, Chang KC, Wang JT (2006) Isolation and characterization of a competence operon associated with transformation and adhesion in *Helicobacter pylori*. *Microbes Infect* 8: 2756-2765.
- [153] Hofreuter D, Karnholz A, Haas R (2003) Topology and membrane interaction of *Helicobacter pylori* ComB proteins involved in natural transformation competence. *Int J Med Microbiol* 293: 153-165.
- [154] Aras RA, Takata T, Ando T, van der Ende A, Blaser MJ (2001) Regulation of the HpyII restriction-modification system of *Helicobacter pylori* by gene deletion and horizontal reconstitution. *Mol Microbiol* 42: 369-382.
- [155] Levine SM, Lin EA, Emara W, Kang J, DiBenedetto M, et al. (2007) Plastic cells and populations: DNA substrate characteristics in *Helicobacter pylori* transformation define a flexible but conservative system for genomic variation. *FASEB J* 21: 3458-3467.
- [156] Zhang XS, Blaser MJ (2012) Natural transformation of an engineered *Helicobacter pylori* strain deficient in type II restriction endonucleases. *J Bacteriol* 194: 3407-3416.
- [157] Kennemann L, Didelot X, Aebischer T, Kuhn S, Drescher B, et al. (2011) *Helicobacter pylori* genome evolution during human infection. *Proc Natl Acad Sci U S A* 108: 5033-5038.

- [158] Ando T, Israel DA, Kusugami K, Blaser MJ (1999) HP0333, a member of the dprA family, is involved in natural transformation in *Helicobacter pylori*. *J Bacteriol* 181: 5572-5580.
- [159] Smeets LC, Bijlsma JJ, Kuipers EJ, Vandenbroucke-Grauls CM, Kusters JG (2000) The dprA gene is required for natural transformation of *Helicobacter pylori*. *FEMS Immunol Med Microbiol* 27: 99-102.
- [160] Dwivedi GR, Sharma E, Rao DN (2013) *Helicobacter pylori* DprA alleviates restriction barrier for incoming DNA. *Nucleic Acids Res* 41: 3274-3288.
- [161] Kang J, Tavakoli D, Tschumi A, Aras RA, Blaser MJ (2004) Effect of host species on recG phenotypes in *Helicobacter pylori* and *Escherichia coli*. *J Bacteriol* 186: 7704-7713.
- [162] Loughlin MF, Barnard FM, Jenkins D, Sharples GJ, Jenks PJ (2003) *Helicobacter pylori* mutants defective in RuvC Holliday junction resolvase display reduced macrophage survival and spontaneous clearance from the murine gastric mucosa. *Infect Immun* 71: 2022-2031.
- [163] Smeets LC, Bijlsma JJ, Boomkens SY, Vandenbroucke-Grauls CM, Kusters JG (2000) comH, a novel gene essential for natural transformation of *Helicobacter pylori*. *J Bacteriol* 182: 3948-3954.
- [164] Baltrus DA, Guillemin K, Phillips PC (2008) Natural transformation increases the rate of adaptation in the human pathogen *Helicobacter pylori*. *Evolution* 62: 39-49.
- [165] Kostidis K, Thackray P, Bardhan KD, Potter CW, Sayers JR ComH: isolation, purification and characterisation of a unique *H. pylori* protein. *Gut* 56: A166.
- [166] O'Rourke EJ, Pinto AV, Petroni EA, Tolmasky ME, Ielpi L (2004) Evidence for the active role of a novel nuclease from *Helicobacter pylori* in the horizontal transfer of genetic information. *J Bacteriol* 186: 2586-2593.
- [167] Liechti GW, Goldberg JB (2013) *Helicobacter pylori* Salvages Purines from Extracellular Host Cell DNA Utilizing the Outer Membrane-Associated Nuclease NucT. *J Bacteriol* 195: 4387-4398.
- [168] Baltrus DA, Guillemin K (2006) Multiple phases of competence occur during the *Helicobacter pylori* growth cycle. *FEMS Microbiol Lett* 255: 148-155.
- [169] Dorer MS, Fero J, Salama NR (2010) DNA damage triggers genetic exchange in *Helicobacter pylori*. *PLoS Pathog* 6: e1001026.
- [170] Kulick S, Moccia C, Didelot X, Falush D, Kraft C, et al. (2008) Mosaic DNA imports with interspersions of recipient sequence after natural transformation of *Helicobacter pylori*. *PLoS One* 3: e3797.

- [171] Lin EA, Zhang XS, Levine SM, Gill SR, Falush D, et al. (2009) Natural transformation of *Helicobacter pylori* involves the integration of short DNA fragments interrupted by gaps of variable size. *PLoS Pathog* 5: e1000337.
- [172] Didelot X, Nell S, Yang I, Woltemate S, van der Merwe S, et al. (2013) Genomic evolution and transmission of *Helicobacter pylori* in two South African families. *Proc Natl Acad Sci U S A* 110: 13880-13885.
- [173] Israel DA, Lou AS, Blaser MJ (2000) Characteristics of *Helicobacter pylori* natural transformation. *FEMS Microbiol Lett* 186: 275-280.
- [174] Suerbaum S, Smith JM, Bapumia K, Morelli G, Smith NH, et al. (1998) Free recombination within *Helicobacter pylori*. *Proc Natl Acad Sci U S A* 95: 12619-12624.
- [175] Falush D, Kraft C, Taylor NS, Correa P, Fox JG, et al. (2001) Recombination and mutation during long-term gastric colonization by *Helicobacter pylori*: estimates of clock rates, recombination size, and minimal age. *Proc Natl Acad Sci U S A* 98: 15056-15061.
- [176] Suerbaum S, Josenhans C (2007) *Helicobacter pylori* evolution and phenotypic diversification in a changing host. *Nat Rev Microbiol* 5: 441-452.
- [177] Schröder G, Lanka E (2005) The mating pair formation system of conjugative plasmids-A versatile secretion machinery for transfer of proteins and DNA. *Plasmid* 54: 1-25.
- [178] Penfold SS, Lastovica AJ, Elisha BG (1988) Demonstration of plasmids in *Campylobacter pylori*. *J Infect Dis* 157: 850-851.
- [179] Höfler C, Fischer W, Hofreuter D, Haas R (2004) Cryptic plasmids in *Helicobacter pylori*: putative functions in conjugative transfer and microcin production. *Int J Med Microbiol* 294: 141-148.
- [180] Hofreuter D, Haas R (2002) Characterization of two cryptic *Helicobacter pylori* plasmids: a putative source for horizontal gene transfer and gene shuffling. *J Bacteriol* 184: 2755-2766.
- [181] De Ungria MC, Kolesnikow T, Cox PT, Lee A (1999) Molecular characterization and interstrain variability of pHP51, a plasmid isolated from the Sydney strain (SS1) of *Helicobacter pylori*. *Plasmid* 41: 97-109.
- [182] Heuermann D, Haas R (1995) Genetic organization of a small cryptic plasmid of *Helicobacter pylori*. *Gene* 165: 17-24.
- [183] Joo JS, Song JY, Baik SC, Lee WK, Cho MJ, et al. (2012) Genetic organization and conjugal plasmid DNA transfer of pHP69, a plasmid from a Korean isolate of *Helicobacter pylori*. *J Microbiol* 50: 955-961.



- [184] Kleanthous H, Clayton CL, Tabaqchali S (1991) Characterization of a plasmid from *Helicobacter pylori* encoding a replication protein common to plasmids in gram-positive bacteria. *Mol Microbiol* 5: 2377-2389.
- [185] Minnis JA, Taylor TE, Knesek JE, Peterson WL, McIntire SA (1995) Characterization of a 3.5-kbp plasmid from *Helicobacter pylori*. *Plasmid* 34: 22-36.
- [186] Backert S, Kwok T, König W (2005) Conjugative plasmid DNA transfer in *Helicobacter pylori* mediated by chromosomally encoded relaxase and TraG-like proteins. *Microbiology* 151: 3493-3503.
- [187] Rohrer S, Holsten L, Weiss E, Benghezal M, Fischer W, et al. (2012) Multiple pathways of plasmid DNA transfer in *Helicobacter pylori*. *PLoS One* 7: e45623.
- [188] Quiñones M, Knesek JE, McIntire SA (2001) Sequence and gene expression analyses of plasmid pHPM8 from *Helicobacter pylori* reveal the presence of two operons with putative roles in plasmid replication and antibiotic activity. *Plasmid* 46: 223-228.
- [189] Kuipers EJ, Israel DA, Kusters JG, Blaser MJ (1998) Evidence for a conjugation-like mechanism of DNA transfer in *Helicobacter pylori*. *J Bacteriol* 180: 2901-2905.
- [190] Oyarzabal OA, Rad R, Backert S (2007) Conjugative transfer of chromosomally encoded antibiotic resistance from *Helicobacter pylori* to *Campylobacter jejuni*. *J Clin Microbiol* 45: 402-408.
- [191] Fischer W, Windhager L, Rohrer S, Zeiller M, Karnholz A, et al. (2010) Strain-specific genes of *Helicobacter pylori*: genome evolution driven by a novel type IV secretion system and genomic island transfer. *Nucleic Acids Res* 38: 6089-6101.
- [192] Kersulyte D, Velapatino B, Mukhopadhyay AK, Cahuayme L, Bussalleu A, et al. (2003) Cluster of type IV secretion genes in *Helicobacter pylori*'s plasticity zone. *J Bacteriol* 185: 3764-3772.
- [193] Kersulyte D, Lee W, Subramaniam D, Anant S, Herrera P, et al. (2009) *Helicobacter Pylori*'s plasticity zones are novel transposable elements. *PLoS One* 4: e6859.
- [194] Occhialini A, Marais A, Alm R, Garcia F, Sierra R, et al. (2000) Distribution of open reading frames of plasticity region of strain J99 in *Helicobacter pylori* strains isolated from gastric carcinoma and gastritis patients in Costa Rica. *Infect Immun* 68: 6240-6249.
- [195] Romo-Gonzalez C, Salama NR, Burgeno-Ferreira J, Ponce-Castaneda V, Lazcano-Ponce E, et al. (2009) Differences in genome content among *Helicobacter pylori* isolates from patients with gastritis, duodenal ulcer, or gastric cancer reveal novel disease-associated genes. *Infect Immun* 77: 2201-2211.
- [196] de Jonge R, Kuipers EJ, Langeveld SC, Loffeld RJ, Stoof J, et al. (2004) The *Helicobacter pylori* plasticity region locus jhp0947-jhp0949 is associated with duodenal ulcer

disease and interleukin-12 production in monocyte cells. *FEMS Immunol Med Microbiol* 41: 161-167.

- [197] Lehours P, Dupouy S, Bergey B, Ruskone-Foumestroux A, Delchier JC, et al. (2004) Identification of a genetic marker of *Helicobacter pylori* strains involved in gastric extranodal marginal zone B cell lymphoma of the MALT-type. *Gut* 53: 931-937.
- [198] Proenca Modena JL, Lopes Sales AI, Olszanski Acrani G, Russo R, Vilela Ribeiro MA, et al. (2007) Association between *Helicobacter pylori* genotypes and gastric disorders in relation to the *cag* pathogenicity island. *Diagn Microbiol Infect Dis* 59: 7-16.
- [199] Rhead JL, Letley DP, Mohammadi M, Hussein N, Mohagheghi MA, et al. (2007) A new *Helicobacter pylori* vacuolating cytotoxin determinant, the intermediate region, is associated with gastric cancer. *Gastroenterology* 133: 926-936.
- [200] Rizwan M, Alvi A, Ahmed N (2008) Novel protein antigen (JHP940) from the genomic plasticity region of *Helicobacter pylori* induces tumor necrosis factor alpha and interleukin-8 secretion by human macrophages. *J Bacteriol* 190: 1146-1151.
- [201] Santos A, Queiroz DM, Menard A, Marais A, Rocha GA, et al. (2003) New pathogenicity marker found in the plasticity region of the *Helicobacter pylori* genome. *J Clin Microbiol* 41: 1651-1655.
- [202] Sugimoto M, Watada M, Jung SW, Graham DY, Yamaoka Y (2012) Role of *Helicobacter pylori* plasticity region genes in development of gastroduodenal diseases. *J Clin Microbiol* 50: 441-448.
- [203] Wozniak RA, Waldor MK (2010) Integrative and conjugative elements: mosaic mobile genetic elements enabling dynamic lateral gene flow. *Nat Rev Microbiol* 8: 552-563.
- [204] Brüssow H, Canchaya C, Hardt WD (2004) Phages and the evolution of bacterial pathogens: from genomic rearrangements to lysogenic conversion. *Microbiol Mol Biol Rev* 68: 560-602, table of contents.
- [205] Marshall BJ, Armstrong JA, Francis GJ, Nokes NT, Wee SH (1987) Antibacterial action of bismuth in relation to *Campylobacter pyloridis* colonization and gastritis. *Digestion* 37 Suppl 2: 16-30.
- [206] Schmid EN, von Recklinghausen G, Ansorg R (1990) Bacteriophages in *Helicobacter* (*Campylobacter*) *pylori*. *J Med Microbiol* 32: 101-104.
- [207] Heintschel von Heinegg E, Nalik HP, Schmid EN (1993) Characterisation of a *Helicobacter pylori* phage (HP1). *J Med Microbiol* 38: 245-249.
- [208] Vale FF, Matos APA, Carvalho P, Vítor JMB (2008) *Helicobacter pylori* phage screening. *Microsc Microanal* 14.
- [209] Lehours P, Vale FF, Bjursell MK, Melefors O, Advani R, et al. (2011) Genome sequencing reveals a phage in *Helicobacter pylori*. *MBio* 2.

- [210] Eppinger M, Baar C, Linz B, Raddatz G, Lanz C, et al. (2006) Who ate whom? Adaptive *Helicobacter* genomic changes that accompanied a host jump from early humans to large felines. *PLoS Genet* 2: e120.
- [211] Luo CH, Chiou PY, Yang CY, Lin NT (2012) Genome, integration, and transduction of a novel temperate phage of *Helicobacter pylori*. *J Virol* 86: 8781-8792.
- [212] Uchiyama J, Takeuchi H, Kato S, Takemura-Uchiyama I, Ujihara T, et al. (2012) Complete genome sequences of two *Helicobacter pylori* bacteriophages isolated from Japanese patients. *J Virol* 86: 11400-11401.
- [213] Uchiyama J, Takeuchi H, Kato S, Gamoh K, Takemura-Uchiyama I, et al. (2013) Characterization of *Helicobacter pylori* bacteriophage KHP30. *Appl Environ Microbiol* 79: 3176-3184.
- [214] Chen J, Novick RP (2009) Phage-mediated intergeneric transfer of toxin genes. *Science* 323: 139-141.
- [215] Sharma CM, Hoffmann S, Darfeuille F, Reignier J, Findeiss S, et al. (2010) The primary transcriptome of the major human pathogen *Helicobacter pylori*. *Nature* 464: 250-255.
- [216] Gottesman S, Storz G (2011) Bacterial small RNA regulators: versatile roles and rapidly evolving variations. *Cold Spring Harb Perspect Biol* 3.
- [217] Storz G, Vogel J, Wassarman KM (2011) Regulation by small RNAs in bacteria: expanding frontiers. *Mol Cell* 43: 880-891.
- [218] Geissmann T, Possedko M, Huntzinger E, Fechter P, Ehresmann C, et al. (2006) Regulatory RNAs as mediators of virulence gene expression in bacteria. *Handb Exp Pharmacol*: 9-43.
- [219] Romby P, Vandenesch F, Wagner EG (2006) The role of RNAs in the regulation of virulence-gene expression. *Curr Opin Microbiol* 9: 229-236.
- [220] Waters LS, Storz G (2009) Regulatory RNAs in bacteria. *Cell* 136: 615-628.
- [221] Babitzke P, Romeo T (2007) CsrB sRNA family: sequestration of RNA-binding regulatory proteins. *Curr Opin Microbiol* 10: 156-163.
- [222] Wassarman KM (2007) 6S RNA: a regulator of transcription. *Mol Microbiol* 65: 1425-1431.
- [223] Henkin TM (2008) Riboswitch RNAs: using RNA to sense cellular metabolism. *Genes Dev* 22: 3383-3390.
- [224] Zhang J, Lau MW, Ferre-D'Amare AR (2010) Ribozymes and riboswitches: modulation of RNA function by small molecules. *Biochemistry* 49: 9123-9131.
- [225] Brennan RG, Link TM (2007) Hfq structure, function and ligand binding. *Curr Opin Microbiol* 10: 125-133.

- [226] Gorke B, Vogel J (2008) Noncoding RNA control of the making and breaking of sugars. *Genes Dev* 22: 2914-2925.
- [227] Storz G, Altuvia S, Wassarman KM (2005) An abundance of RNA regulators. *Annu Rev Biochem* 74: 199-217.
- [228] Vanderpool CK, Balasubramanian D, Lloyd CR (2011) Dual-function RNA regulators in bacteria. *Biochimie* 93: 1943-1949.
- [229] Pernitzsch SR, Sharma CM (2012) Transcriptome complexity and riboregulation in the human pathogen *Helicobacter pylori*. *Front Cell Infect Microbiol* 2: 14.
- [230] Parkhill J, Wren BW, Mungall K, Ketley JM, Churcher C, et al. (2000) The genome sequence of the food-borne pathogen *Campylobacter jejuni* reveals hypervariable sequences. *Nature* 403: 665-668.
- [231] Lasa I, Toledo-Arana A, Dobin A, Villanueva M, de los Mozos IR, et al. (2011) Genome-wide antisense transcription drives mRNA processing in bacteria. *Proc Natl Acad Sci U S A* 108: 20172-20177.
- [232] Thibonnier M, Thiberge JM, De Reuse H (2008) Trans-translation in *Helicobacter pylori*: essentiality of ribosome rescue and requirement of protein tagging for stress resistance and competence. *PLoS One* 3: e3810.
- [233] Wassarman KM, Storz G (2000) 6S RNA regulates *E. coli* RNA polymerase activity. *Cell* 101: 613-623.
- [234] Barrick JE, Sudarsan N, Weinberg Z, Ruzzo WL, Breaker RR (2005) 6S RNA is a widespread regulator of eubacterial RNA polymerase that resembles an open promoter. *RNA* 11: 774-784.
- [235] Faucher SP, Friedlander G, Livny J, Margalit H, Shuman HA (2010) *Legionella pneumophila* 6S RNA optimizes intracellular multiplication. *Proc Natl Acad Sci U S A* 107: 7533-7538.
- [236] Wen Y, Feng J, Scott DR, Marcus EA, Sachs G (2011) A cis-encoded antisense small RNA regulated by the HP0165-HP0166 two-component system controls expression of ureB in *Helicobacter pylori*. *J Bacteriol* 193: 40-51.
- [237] Williams SM, Chen YT, Andermann TM, Carter JE, McGee DJ, et al. (2007) *Helicobacter pylori* chemotaxis modulates inflammation and bacterium-gastric epithelium interactions in infected mice. *Infect Immun* 75: 3747-3757.
- [238] McGee DJ, Langford ML, Watson EL, Carter JE, Chen YT, et al. (2005) Colonization and inflammation deficiencies in Mongolian gerbils infected by *Helicobacter pylori* chemotaxis mutants. *Infect Immun* 73: 1820-1827.
- [239] Croxen MA, Sisson G, Melano R, Hoffman PS (2006) The *Helicobacter pylori* chemotaxis receptor TlpB (HP0103) is required for pH taxis and for colonization of the gastric mucosa. *J Bacteriol* 188: 2656-2665.

- [240] Schreiber S, Konradt M, Groll C, Scheid P, Hanauer G, et al. (2004) The spatial orientation of *Helicobacter pylori* in the gastric mucus. *Proc Natl Acad Sci U S A* 101: 5024-5029.
- [241] Shimomura H, Hayashi S, Yokota K, Oguma K, Hirai Y (2004) Alteration in the composition of cholesteryl glucosides and other lipids in *Helicobacter pylori* undergoing morphological change from spiral to coccoid form. *FEMS Microbiol Lett* 237: 407-413.
- [242] Chaput C, Ecobichon C, Cayet N, Girardin SE, Werts C, et al. (2006) Role of AmiA in the morphological transition of *Helicobacter pylori* and in immune escape. *PLoS Pathog* 2: e97.
- [243] Nagai S, Mimuro H, Yamada T, Baba Y, Moro K, et al. (2007) Role of Peyer's patches in the induction of *Helicobacter pylori*-induced gastritis. *Proc Natl Acad Sci U S A* 104: 8971-8976.
- [244] Wang G, Maier SE, Lo LF, Maier G, Dosi S, et al. (2010) Peptidoglycan deacetylation in *Helicobacter pylori* contributes to bacterial survival by mitigating host immune responses. *Infect Immun* 78: 4660-4666.
- [245] Harris AG, Wilson JE, Danon SJ, Dixon MF, Donegan K, et al. (2003) Catalase (KatA) and KatA-associated protein (KapA) are essential to persistent colonization in the *Helicobacter pylori* SS1 mouse model. *Microbiology* 149: 665-672.
- [246] Dorer MS, Cohen IE, Sessler TH, Fero J, Salama NR (2013) Natural competence promotes *Helicobacter pylori* chronic infection. *Infect Immun* 81: 209-215.
- [247] Schoep TD, Fulurija A, Good F, Lu W, Himbeck RP, et al. (2010) Surface properties of *Helicobacter pylori* urease complex are essential for persistence. *PLoS One* 5: e15042.
- [248] Ha NC, Oh ST, Sung JY, Cha KA, Lee MH, et al. (2001) Supramolecular assembly and acid resistance of *Helicobacter pylori* urease. *Nat Struct Biol* 8: 505-509.
- [249] Pinkse MW, Maier CS, Kim JI, Oh BH, Heck AJ (2003) Macromolecular assembly of *Helicobacter pylori* urease investigated by mass spectrometry. *J Mass Spectrom* 38: 315-320.
- [250] Marcus EA, Scott DR (2001) Cell lysis is responsible for the appearance of extracellular urease in *Helicobacter pylori*. *Helicobacter* 6: 93-99.
- [251] Hu LT, Mobley HL (1990) Purification and N-terminal analysis of urease from *Helicobacter pylori*. *Infect Immun* 58: 992-998.
- [252] Marshall BJ, Barrett LJ, Prakash C, McCallum RW, Guerrant RL (1990) Urea protects *Helicobacter* (*Campylobacter*) *pylori* from the bactericidal effect of acid. *Gastroenterology* 99: 697-702.
- [253] Cao P, McClain MS, Forsyth MH, Cover TL (1998) Extracellular release of antigenic proteins by *Helicobacter pylori*. *Infect Immun* 66: 2984-2986.



- [254] Voland P, Weeks DL, Marcus EA, Prinz C, Sachs G, et al. (2003) Interactions among the seven *Helicobacter pylori* proteins encoded by the urease gene cluster. *Am J Physiol Gastrointest Liver Physiol* 284: G96-G106.
- [255] Scott DR, Marcus EA, Weeks DL, Lee A, Melchers K, et al. (2000) Expression of the *Helicobacter pylori* ureI gene is required for acidic pH activation of cytoplasmic urease. *Infect Immun* 68: 470-477.
- [256] Weeks DL, Eskandari S, Scott DR, Sachs G (2000) A H<sup>+</sup>-gated urea channel: the link between *Helicobacter pylori* urease and gastric colonization. *Science* 287: 482-485.
- [257] Weeks DL, Gushansky G, Scott DR, Sachs G (2004) Mechanism of proton gating of a urea channel. *J Biol Chem* 279: 9944-9950.
- [258] Scott DR, Marcus EA, Wen Y, Singh S, Feng J, et al. (2010) Cytoplasmic histidine kinase (HP0244)-regulated assembly of urease with UreI, a channel for urea and its metabolites, CO<sub>2</sub>, NH<sub>3</sub>, and NH<sub>4</sub>(+), is necessary for acid survival of *Helicobacter pylori*. *J Bacteriol* 192: 94-103.
- [259] Eaton KA, Krakowka S (1994) Effect of gastric pH on urease-dependent colonization of gnotobiotic piglets by *Helicobacter pylori*. *Infect Immun* 62: 3604-3607.
- [260] Williams CL, Preston T, Hossack M, Slater C, McColl KEL (1996) *Helicobacter pylori* utilises urease for amino acid synthesis. *FEMS Immunol Med Microbiol* 13: 87-94.
- [261] Kuwahara H, Miyamoto Y, Akaike T, Kubota T, Sawa T, et al. (2000) *Helicobacter pylori* urease suppresses bactericidal activity of peroxynitrite via carbon dioxide production. *Infect Immun* 68: 4378-4383.
- [262] Schwartz JT, Allen LA (2006) Role of urease in megasome formation and *Helicobacter pylori* survival in macrophages. *J Leukoc Biol* 79: 1214-1225.
- [263] Makristathis A, Rokita E, Labigne A, Willinger B, Rotter ML, et al. (1998) Highly significant role of *Helicobacter pylori* urease in phagocytosis and production of oxygen metabolites by human granulocytes. *J Infect Dis* 177: 803-806.
- [264] Rokita E, Makristathis A, Presterl E, Rotter ML, Hirschl AM (1998) *Helicobacter pylori* urease significantly reduces opsonization by human complement. *J Infect Dis* 178: 1521-1525.
- [265] Celli JP, Turner BS, Afdhal NH, Keates S, Ghiran I, et al. (2009) *Helicobacter pylori* moves through mucus by reducing mucin viscoelasticity. *Proc Natl Acad Sci U S A* 106: 14321-14326.
- [266] Harris PR, Ernst PB, Kawabata S, Kiyono H, Graham MF, et al. (1998) Recombinant *Helicobacter pylori* urease activates primary mucosal macrophages. *J Infect Dis* 178: 1516-1520.

- [267] Harris PR, Mobley HL, Perez-Perez GI, Blaser MJ, Smith PD (1996) *Helicobacter pylori* urease is a potent stimulus of mononuclear phagocyte activation and inflammatory cytokine production. *Gastroenterology* 111: 419-425.
- [268] Wroblewski LE, Shen L, Ogden S, Romero-Gallo J, Lapierre LA, et al. (2009) *Helicobacter pylori* dysregulation of gastric epithelial tight junctions by urease-mediated myosin II activation. *Gastroenterology* 136: 236-246.
- [269] Tanahashi T, Kita M, Kodama T, Yamaoka Y, Sawai N, et al. (2000) Cytokine expression and production by purified *Helicobacter pylori* urease in human gastric epithelial cells. *Infect Immun* 68: 664-671.
- [270] Fan X, Gunasena H, Cheng Z, Espejo R, Crowe SE, et al. (2000) *Helicobacter pylori* urease binds to class II MHC on gastric epithelial cells and induces their apoptosis. *J Immunol* 165: 1918-1924.
- [271] Boneca IG, Ecobichon C, Chaput C, Mathieu A, Guadagnini S, et al. (2008) Development of inducible systems to engineer conditional mutants of essential genes of *Helicobacter pylori*. *Appl Environ Microbiol* 74: 2095-2102.
- [272] Zawilak-Pawlik A, Kois A, Stingl K, Boneca IG, Skrobuk P, et al. (2007) HobA--a novel protein involved in initiation of chromosomal replication in *Helicobacter pylori*. *Mol Microbiol* 65: 979-994.
- [273] Bertram R, Hillen W (2008) The application of Tet repressor in prokaryotic gene regulation and expression. *Microb Biotechnol* 1: 2-16.
- [274] Sun Y, Chen X, Xiao D (2007) Tetracycline-inducible expression systems: new strategies and practices in the transgenic mouse modeling. *Acta Biochim Biophys Sin (Shanghai)* 39: 235-246.
- [275] Debowski AW, Verbrugghe P, Sehnal M, Marshall BJ, Benghezal M (2013) Development of a Tetracycline-Inducible Gene Expression System for the Study of *Helicobacter pylori* Pathogenesis. *Appl Environ Microbiol* 79: 7351-7359.
- [276] McClain MS, Duncan SS, Gaddy JA, Cover TL (2013) Control of gene expression in *Helicobacter pylori* using the Tet repressor. *J Microbiol Methods* 95: 336-341.
- [277] Debowski AW, Himbeck R, Middleton M, Camilleri T, Sehnal M, et al. (2011) A tetracycline-based conditional knockout demonstrates the role of urease in colonization and maintenance of *Helicobacter pylori* infection. *CHRO A114*.
- [278] Garcia-Vallve S, Janssen PJ, Ouzounis CA (2002) Genetic variation between *Helicobacter pylori* strains: gene acquisition or loss? *Trends Microbiol* 10: 445-447.
- [279] Saunders NJ, Boonmee P, Peden JF, Jarvis SA (2005) Inter-species horizontal transfer resulting in core-genome and niche-adaptive variation within *Helicobacter pylori*. *BMC Genomics* 6: 9.

- [280] Pupo E, Padron A, Santana E, Sotolongo J, Quintana D, et al. (2005) Preparation of plasmid DNA-containing liposomes using a high-pressure homogenization--extrusion technique. *J Control Release* 104: 379-396.
- [281] Wong BC, Yin Y, Berg DE, Xia HH, Zhang JZ, et al. (2001) Distribution of distinct *vacA*, *cagA* and *iceA* alleles in *Helicobacter pylori* in Hong Kong. *Helicobacter* 6: 317-324.
- [282] Owen RJ, Ferrus M, Gibson J (2001) Amplified fragment length polymorphism genotyping of metronidazole-resistant *Helicobacter pylori* infecting dyspeptics in England. *Clin Microbiol Infect* 7: 244-253.
- [283] Carroll IM, Ahmed N, Beesley SM, Khan AA, Ghousunnissa S, et al. (2004) Microevolution between paired antral and paired antrum and corpus *Helicobacter pylori* isolates recovered from individual patients. *J Med Microbiol* 53: 669-677.
- [284] Kuipers EJ, Israel DA, Kusters JG, Gerrits MM, Weel J, et al. (2000) Quasispecies development of *Helicobacter pylori* observed in paired isolates obtained years apart from the same host. *J Infect Dis* 181: 273-282.
- [285] Smeets LC, Arents NL, van Zwet AA, Vandenbroucke-Grauls CM, Verboom T, et al. (2003) Molecular patchwork: Chromosomal recombination between two *Helicobacter pylori* strains during natural colonization. *Infect Immun* 71: 2907-2910.
- [286] Salama N, Guillemin K, McDaniel TK, Sherlock G, Tompkins L, et al. (2000) A whole-genome microarray reveals genetic diversity among *Helicobacter pylori* strains. *Proc Natl Acad Sci U S A* 97: 14668-14673.
- [287] Gressmann H, Linz B, Ghai R, Pleissner KP, Schlapbach R, et al. (2005) Gain and loss of multiple genes during the evolution of *Helicobacter pylori*. *PLoS Genet* 1: e43.
- [288] Israel DA, Salama N, Arnold CN, Moss SF, Ando T, et al. (2001) *Helicobacter pylori* strain-specific differences in genetic content, identified by microarray, influence host inflammatory responses. *J Clin Invest* 107: 611-620.
- [289] Israel DA, Salama N, Krishna U, Rieger UM, Atherton JC, et al. (2001) *Helicobacter pylori* genetic diversity within the gastric niche of a single human host. *Proc Natl Acad Sci U S A* 98: 14625-14630.
- [290] Kraft C, Stack A, Josenhans C, Niehus E, Dietrich G, et al. (2006) Genomic changes during chronic *Helicobacter pylori* infection. *J Bacteriol* 188: 249-254.
- [291] Morelli G, Didelot X, Kusecek B, Schwarz S, Bahlawane C, et al. (2010) Microevolution of *Helicobacter pylori* during prolonged infection of single hosts and within families. *PLoS Genet* 6: e1001036.
- [292] Graham DY, Opekun AR, Osato MS, El-Zimaity HM, Lee CK, et al. (2004) Challenge model for *Helicobacter pylori* infection in human volunteers. *Gut* 53: 1235-1243.
- [293] Aebischer T, Bumann D, Epple HJ, Metzger W, Schneider T, et al. (2008) Correlation of T cell response and bacterial clearance in human volunteers challenged with *Helicobacter pylori*.

cobacter pylori revealed by randomised controlled vaccination with Ty21a-based Salmonella vaccines. Gut 57: 1065-1072.

- [294] Lundin A, Bjorkholm B, Kupersmidt I, Unemo M, Nilsson P, et al. (2005) Slow genetic divergence of *Helicobacter pylori* strains during long-term colonization. Infect Immun 73: 4818-4822.
- [295] Salama NR, Gonzalez-Valencia G, Deatherage B, Aviles-Jimenez F, Atherton JC, et al. (2007) Genetic analysis of *Helicobacter pylori* strain populations colonizing the stomach at different times postinfection. J Bacteriol 189: 3834-3845.
- [296] Beji A, Vincent P, Darchis I, Husson MO, Cortot A, et al. (1989) Evidence of gastritis with several *Helicobacter pylori* strains. Lancet 2: 1402-1403.
- [297] Ghose C, Perez-Perez GI, van Doorn LJ, Dominguez-Bello MG, Blaser MJ (2005) High frequency of gastric colonization with multiple *Helicobacter pylori* strains in Venezuelan subjects. J Clin Microbiol 43: 2635-2641.
- [298] Patra R, Chattopadhyay S, De R, Ghosh P, Ganguly M, et al. (2012) Multiple infection and microdiversity among *Helicobacter pylori* isolates in a single host in India. PLoS One 7: e43370.
- [299] Marshall BJ, Armstrong JA, McGeachie DB, Glancy RJ (1985) Attempt to fulfil Koch's postulates for pyloric Campylobacter. Med J Aust 142: 436-439.
- [300] Kavermann H, Burns BP, Angermuller K, Odenbreit S, Fischer W, et al. (2003) Identification and characterization of *Helicobacter pylori* genes essential for gastric colonization. J Exp Med 197: 813-822.
- [301] Kersulyte D, Chalkauskas H, Berg DE (1999) Emergence of recombinant strains of *Helicobacter pylori* during human infection. Mol Microbiol 31: 31-43.
- [302] Krebs J, Bunk B, Spröer C, Luong K, Morgan RD, et al. (2013) Comprehensive methylome analysis of the human gastric pathogen, *Helicobacter pylori*. 17th International Workshop on Campylobacter, Helicobacter and Related Organisms Abstract O57a.
- [303] Takeuchi H, Israel DA, Miller GG, Donahue JP, Krishna U, et al. (2002) Characterization of expression of a functionally conserved *Helicobacter pylori* methyltransferase-encoding gene within inflamed mucosa and during in vitro growth. J Infect Dis 186: 1186-1189.
- [304] Koonin EV, Wolf YI (2010) Constraints and plasticity in genome and molecular-phenome evolution. Nat Rev Genet 11: 487-498.
- [305] Levy SF, Siegal ML (2012) The robustness continuum. Adv Exp Med Biol 751: 431-452.
- [306] Masel J, Siegal ML (2009) Robustness: mechanisms and consequences. Trends Genet 25: 395-403.

- [307] Masei J, Trotter MV (2010) Robustness and evolvability. *Trends Genet* 26: 406-414.
- [308] Wagner A (2012) The role of robustness in phenotypic adaptation and innovation. *Proc Biol Sci* 279: 1249-1258.
- [309] Wagner GP, Altenberg L (1996) Perspective: Complex Adaptations and the Evolution of Evolvability. *Evolution* 50: 967-976.
- [310] Kussell E, Leibler S (2005) Phenotypic diversity, population growth, and information in fluctuating environments. *Science* 309: 2075-2078.
- [311] Vakil N, Vaira D (2013) Treatment for *H. pylori* infection: new challenges with antimicrobial resistance. *J Clin Gastroenterol* 47: 383-388.
- [312] Debowski AW, Carnoy C, Verbrugghe P, Nilsson HO, Gauntlett JC, et al. (2012) Xer recombinase and genome integrity in *Helicobacter pylori*, a pathogen without topoisomerase IV. *PLoS One* 7: e33310.
- [313] Strugatsky D, McNulty R, Munson K, Chen CK, Soltis SM, et al. (2013) Structure of the proton-gated urea channel from the gastric pathogen *Helicobacter pylori*. *Nature* 493: 255-258.
- [314] Nilsson C, Skoglund A, Moran AP, Annuk H, Engstrand L, et al. (2006) An enzymatic ruler modulates Lewis antigen glycosylation of *Helicobacter pylori* LPS during persistent infection. *Proc Natl Acad Sci U S A* 103: 2863-2868.
- [315] Gunasekara S, Vrielink A, Stubbs KA (2010) Preliminary studies into the inhibition of the cholesterol alpha-glucosyltransferase from *Helicobacter pylori* using azasugars. *Carbohydr Res* 345: 960-964.
- [316] Thomson N, Sebahia M, Cerdano-Tarraga A, Bentley S, Crossman L, et al. (2003) The value of comparison. *Nat Rev Microbiol* 1: 11-12.