

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

185,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Natural Compound-Generated Oxidative Stress: From Bench to Bedside

Aloran Mazumder and Marc Diederich

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/64183>

Abstract

Oxidants are constantly generated in a biological system as a result of physiological processes. However, an imbalance between oxidants and antioxidants can lead to a pathophysiological condition known as oxidative stress. Natural compounds as inducers of oxidative stress are able to modulate physiological functions of cancer cells leading to cell death or survival. This chapter aims at providing an overview of pro- and antioxidant activities of natural compounds related to cancer and related therapies.

Keywords: natural compounds, cancer, oxidative stress, clinical use

1. Natural compound anticancer agents

In the search of improved cytotoxic agents against cancer, natural compounds possess advantages with regard to availability, low toxicity, and suitability for oral application and metabolite likeliness [1]. Moreover, new technologies of combinatorial chemistry and high-throughput screening are used to design different synthetic drugs with natural compounds that serve as templates for development of novel molecules with enhanced biological properties.

In 1960, the National Cancer Institute (NCI) began a large-scale screening program for anti-tumor agents, and 35,000 plant species samples were tested primarily on mouse leukemia cells [2, 3]. The most promising drug to emerge from this program was paclitaxel, a microtubule disruptive agent obtained from the bark of the Pacific yew *Taxus brevifolia*. This finding served as the springboard for further investigations with natural compounds, and in the late 1960s, vinblastine and vincristine were reported from *Catharanthus roseus*. Both drugs major-

ly contributed to long-term remission and cures for childhood leukemia, Hodgkin's lymphoma, testicular teratoma, etc. Other anticancer agents to enter clinics, which are derived from natural sources, include etoposide, which has been proven as an effective treatment against testicular teratoma and small cell lung cancer, whereas teniposide was shown to be effective against acute lymphocytic leukemia (ALL) and neuroblastoma in children and non-Hodgkin's lymphoma [1]. A comprehensive study published on new medicines approved by US Food and Drug Administration between 1981 and 2010 revealed that 34% of those medicines based on small molecules were either natural products or a direct derivative which mainly included statins, immunosuppressant, and tubulin-binding anticancer drugs [4, 5].

Natural compound constituents demonstrated anticancer activity according to a combination of epidemiological and experimental studies [6]. Mechanistic insights underlined that the chemotherapeutic potential of these agents may be a combination of antioxidant, anti-inflammatory, immune-promoting, cytostatic, differentiating, and cytotoxic effects. Altogether, natural compounds efficiently prevent initiation, promotion, and progression of cancer development thus interfering with all 10 hallmarks and enabling characteristics of cancer [7–10].

Increasing technological advancements led to the development of better purification techniques with defined molecular assays, which can efficiently exclude “distracting molecules” such as tannins and saponins, thereby increasing the chances of identifying the critical agent with specific anticancer activity. The diverse bioactivity potential of natural compounds can be related to the huge structural diversity existing in nature. This compound repertoire is available for further modifications to improve the therapeutic potential of lead compounds. In addition, combinatorial biosynthesis further modulates the functional groups of lead compounds and can be complemented with high-throughput screening, computational chemistry,




Figure 1. Molecular scaffolds of plant anticancer agents. Molecular structures were drawn with ChemDraw13 (Perkin Elmer).

and bioinformatics to generate structural analogues with improved pharmacological activity and reduced toxicity [1] (**Figure 1**).

2. Natural compounds as scavengers of free radicals

Oxidants are constantly generated in a biological system as a result of physiological processes. However, an imbalance between oxidants and antioxidants can lead to a pathophysiological condition known as oxidative stress [11]. In light of this knowledge, oxidative stress has been defined as perturbations in redox homeostasis. Broadly, the cellular redox level is regulated by three different systems, two of which are dependent on glutathione that includes glutathione (GSH), glutathione reductase (GR), glutathione peroxidases (GPX), and glutathione S-transferases (GST) [12–14]. Glutathione undergoes oxidation to form glutathione disulfide (GSSG), thereby reducing the disulfide bonds of cytoplasmic proteins to cysteine and protects the cell against oxidative stress [15]. Under normal conditions, GSH exists in reduced form due to constitutive activity of GR. GSTs act as detoxifying enzymes that conjugate GSH to various electrophilic compounds [16].

Reactive oxygen species (ROS) have been reported in both solid and hematopoietic cancers where they are associated with tumor development and progression [17, 18]. However, cancer cells also express antioxidant proteins to detoxify ROS, suggesting that the fine-tuning of intracellular ROS signaling is critical for cancer. Therefore, understanding the susceptibility of cancer cells to oxidative signals could open new therapeutic window for rational design of new anticancer agents [19]. In addition to their well-characterized effects on cell division and viability, cytotoxic agents can induce oxidative stress by modulating levels of ROS such as the superoxide anion radical, hydrogen peroxide, and hydroxyl radicals. Eukaryotic cells have highly organized pathways to orchestrate the many extracellular stimuli received and convert them into specific physiological processes. This classical cascade also termed as signal transduction pathways includes a series of events occurring constitutively and initiated by interaction of a ligand with its receptor on the cell membrane. ROS in this cascade have been proposed as second messengers in the activation of signaling events that lead to survival or death [20]. Moreover, redox-sensitive cysteine residues are known to sense and transduce changes in cellular redox status caused by ROS production and the presence of oxidized thiols. Various dietary phytochemicals have been shown to exhibit beneficial effects including the prevention of cancer by modulating the cellular redox status by acting as either an antioxidant or pro-oxidant. They function as detoxifying enzyme inducers, which mainly include phenolic and sulfur-containing compounds. Phenolic compounds are classified as polyphenols or flavonoids, whereas sulfur-containing compounds may be classified into isothiocyanates and organosulfur compounds. Epigallocatechin-3-gallate (EGCG) from green tea, curcumin [21–24] from turmeric, and resveratrol [25, 26] from grapes are the classical examples of polyphenols, whereas flavonoids include quercetin from citrus fruits [26–28] and genistein from soya. Isothiocyanates represent a group of compounds such as sulforaphane from broccoli and phenethyl isothiocyanate from turnips. Organosulfur compounds mainly include diallyltrasulfide derived from garlic [29–34]. Cells respond to these phytochemicals by a non-classical

receptor-sensing mechanism of electrophilic chemical stress characterized as “thiol-modulated cellular signaling” events leading to gene expression commending the pharmacological activity (**Figure 2**).




Figure 2. Natural compounds as scavengers of free radicals. Molecular structures were drawn with ChemDraw13 (Perkin Elmer).

3. Survival pathways activated by free radicals

ROSs are tumorigenic as elevated levels of ROS-sensitive signaling pathways have been implicated in various cancers where they are involved in sustenance of cell growth, prolifer-

ation, survival, migration, and by inducing DNA damage leading to formation of genetic lesions initiating tumorigenesis [35, 36]. Low levels of hydrogen peroxide (H_2O_2) stimulation have been shown to propagate cell proliferation in an array of cancer cell types. Role of hormones in endocrine cancers is well documented. In hormone-dependent breast cancer cells, one of the functions of estrogen is to translocate to mitochondria, thereby initiating mitochondrial ROS production that can be impaired by inhibition of mitochondrial uniporter, which prevents estrogen-induced cell proliferation [37, 38]. Sodium arsenic in MCF-7 was shown to mimic the effect of estrogen and potentiated S phase progression and proliferation by inducing ROS production and ROS-related depolarization of the mitochondrial membrane [39]. Moreover, estrogen-induced cell proliferation of MCF-7 was strongly inhibited by antioxidants such as *N*-acetyl-L-cysteine (NAC) or mitochondrial blockers of protein synthesis such as chloramphenicol [40]. ROS generation was shown to augment G1/S transition by increasing the expression levels of cyclins D1, D3, E1, E2, and B2 [41]. In contingent to these finding, cytochrome P450B1-mediated conversion of estrogen to a putative carcinogenic metabolite 4-hydroxyestradiol in human mammary epithelial cells MCF-10 leads to intracellular ROS production and neoplastic transformation. ROS overproduction was shown to activate I κ B kinase (IKK) signaling with increased nuclear translocation and NF- κ B activity [42].

Since deregulation of NF- κ B is related to increased cell survival, proliferation, and development of drug resistance in different cancers, series of work conducted in this direction showed that NF- κ B is a redox-regulated sensor for oxidative stress and is activated by low doses of H_2O_2 [43, 44]. In MCF-7 cells, interleukin (IL)-1 β stimulation of NF- κ B is partially regulated by H_2O_2 -mediated activation of NF- κ B inducing kinase (NIK)-mediated phosphorylation of IKK α [45]. Moreover, overexpression of manganese superoxide dismutase (MnSOD) in MCF-7 cells completely abolished tumor necrosis factor (TNF) α -mediated NF- κ B activation, I κ B α degradation, p65 nuclear translocation, and NF- κ B-dependent reporter gene expression [40]. In other forms of cancer such as oral squamous carcinoma, a mild difference in endogenous ROS functions as a physiological signaling modulator of the NF- κ B signaling cascades through its ability to activate NIK [46]. Besides solid tumors, redox regulation of NF- κ B has also been implicated in hematopoietic cancers. Our group for the first time reported that in U937 cells, melatonin a pineal hormone might induce ROS generation, which ultimately is involved in transactivation of NF- κ B-promoting survival of these cells [47–50]. Moreover, myeloid leukemia, which often maintains a high intracellular ROS level and uses redox signal for survival, is sensitive to NF- κ B inhibition since NF- κ B is involved in moderating the ROS level, which prevent activation of c-Jun N-terminal kinase (JNK) and cell death [51–54] (**Figure 3**).

Apart from NF- κ B, ROS-mediated regulation of tyrosine phosphatases, protein tyrosine kinases, and receptor tyrosine kinases, which is critical for cell survival and cancer such as mitogen-activated protein (MAP) kinase/extracellular-regulated kinase (Erk) cascade and phosphoinositide-3-kinase (PI3K)/Akt-regulated signaling cascade, is well documented in the literature [55, 56]. Activation of MAPK/Erk1/2, which is mediated through growth factors, and K-ras is functionally linked to increased cell proliferation. Several studies have shown how ROS activate Erk1/2 pathway by modulating and activating its upstream target such as Ras. For instance, oxidative modification at its cysteine 118 residue leads to the inhibition of GDP/

GTP exchange [57]. Moreover, ROS activates p90^{RSK} that acts as an upstream kinase of Erk1/2 [58, 59]. In ovarian cancer, sustained Erk1/2 activity was linked to increased concentration of endogenous ROS resulting from ubiquitination and loss of endogenous mitogen-activated protein kinase phosphatase 3 (MKP3), which negatively regulates Erk1/2 [58, 59].




Figure 3. Molecular scaffolds of physiological antioxidants Molecular structures were drawn with ChemDraw13 (Perkin Elmer).

Oxidative stress regulation of PI3K/Akt pathway has been implicated in different cancers. In ovarian cancers, H₂O₂ produced in response to epithelial growth factor signaling (EGF) activates Akt and p70 S6k1, a substrate of Akt involved in regulating protein synthesis [60]. In pancreatic cancer PANC-1 cells, NADPH oxidase (NOX)-4-mediated generation of intracellular ROS was related to survival of these cells, which undergo apoptosis in response to diphenylene iodonium (DPI), an inhibitor of NOX that inhibited superoxide production and impaired levels of phosphorylated Akt [61]. Moreover, benzo(a)pyrene (BaP), a known mammary carcinogen in rodents, increased cell proliferation in human mammary epithelial cells MCF-10A through H₂O₂ generation and activation of epidermal growth factor receptor (EGFR), Akt, and ERK phosphorylation, which was strongly inhibited by NAC treatment [62].

4. Reactive oxygen species contribute in tumor progression

Intracellular redox status aids tumor progression by modulating the processes of metastasis, angiogenesis, survival of cells under hypoxic conditions, and maintenance of cancer stem cell (CSC) subpopulation [63]. Decreased cell adhesion to extracellular matrix, anchorage-independent survival, and invasion of tumor cells are well documented to be influenced by ROS [64]. Perturbation of mitochondrial respiratory chain in breast cancer cells leads to generation of a cellular subpopulation with increased levels of ROS, which are highly metastatic and maintain increased invasive property in vivo [65]. ROS induction was shown to influence overexpression of chemokine CXCL14 through the activator protein (AP)-1-signaling pathway and promote cell motility through elevation of cytosolic Ca²⁺ by binding to the inositol 1,4,5 triphosphate receptor on the endoplasmic reticulum [65]. DNA methylation and histone modification leading to epigenetic silencing of superoxide dismutase (SOD)-2 alter the expression of antioxidant enzyme MnSOD, which promotes invasion of breast cancers [66].

Moreover, a decreased MnSOD level was also associated with increased pancreatic tumor invasion [67]. Degradation of the extracellular matrix (ECM) and activated matrix metalloproteinases (MMPs) are a prerequisite of cancer cell migration and invasion. Binding of several integrins to the ECM results in increased expression of several MMP proteins. Since integrins signal by a vast array of kinases, phosphatases, GTPases, and transcription factors, it is likely that an elevated level of ROS has an effect on integrin-mediated signaling. Several studies reported the inactivation of critical phosphatases such as protein tyrosine phosphatase (PTP)-PEST (PTPN12), SHP-2 (Src homology 2 [SH2] domain-containing non-transmembrane PTP), and low molecular weight protein tyrosine phosphatases (LMW-PTPs) by oxidation [68]. Catalase, a H₂O₂ scavenger, binds SHP-2 and growth factor receptor-bound protein-2 (Grb2) adapter protein upon integrin ligand binding and therefore protects them against H₂O₂-mediated oxidation [69]. In non-transformed intestinal epithelial cells, elevated ROS increased the expression of α 2 β 1-integrin, which subsequently increased the levels of cyclooxygenase-2 (COX-2) and promoted cell migration [64]. These results also suggest a mechanism where ROS-induced modulation of ECM promotes cancer formation in intestinal epithelial cells. ROSs have also been implicated in promoting tumor progression by modulating the processes involved in epithelial mesenchymal transition (EMT). Several transcription factors, which promote metastasis such as AP-1, Ets, Smad, and Snail, are regulated by ROS, inducing an effect on upstream target molecules involved in activation of these transcription factors such as protein kinase (PK) C and PTPs [70].




Figure 4. Molecular mechanisms of hypoxia affected by natural compounds. Scheme was drawn with ScienceSlides Suite 2105 (Visiscience).

In a given tumor mass, cancer cells often are exposed to an environment with reduced levels of tissue oxygen, a condition known as hypoxia. Prolonged limitation in oxygen supply can result in cell death. Therefore, cancer cells often undergo genetic and adaptive changes that contribute to a malignant phenotype and adopt characteristics of an aggressive tumor. Cancer cells mimic a phenomenon known as the “Warburg effect” that is to switch to anaerobic glycolysis when adequate oxygen supply is absent [71]. ROSs have been implicated to facilitate the tumor survival under hypoxic conditions by modulating different transcription factors involved. Hypoxia inducible transcription factor (HIF)-1 is most widely studied for its role in tumor promotion under hypoxic conditions. HIF-1 is a heterodimer that consists of hypoxic response factor HIF-1 α and constitutively expressed aryl hydrocarbon receptor nuclear translocator (ARNT) also known as HIF-1 β [72]. Under reduced oxygen levels, HIF-1 binds to hypoxia response elements, thereby activating hypoxia response genes such as the pro-angiogenic vascular endothelial growth factor (VEGF) [73]. Moreover, HIF-1 has been shown to regulate expression of all enzymes of the glycolysis pathway as well as glucose transporters GLUT1 and GLUT3 [74]. In human breast carcinoma, increased MnSOD activity is reported to inhibit HIF-1 α along with suppression VEGF protein that impaired tumor metastasis [75]. Suppression of endogenous ROS by NADPH oxidase inhibitor DPI and mitochondrial electron chain inhibitor rotenone decreased HIF-1 induction and VEGF expression in ovarian and prostate cancer cells [75]. Moreover, growth factor such as epidermal growth factor (EGF)-induced ROS production may lead to activation of AKT/p70S6K1 pathway resulting in increased expression of VEGF stimulating tumor angiogenesis [60] (**Figure 4**).

In any given tumor, subpopulations of cells have the ability to self-renew and drive tumorigenesis. This population of cells is termed as cancer stem cells (CSCs), which are isolated from most cancers such as hematopoietic, breast, lung, colon, etc. CSCs are characterized by the expression of specific stem cell markers and are of clinical relevance as they are highly drug resistant and mostly initiate recurrence after chemo- or radiotherapy [76]. Studies have shown that normal hematopoietic and epithelial stem cells maintained a lower level of ROS than mature progeny to prevent cellular differentiation and maintain long-term cellular self-renewable. Similarly, CSCs unlike cancer cells have reduced level of ROS. Moreover, compared to tumor cell counterparts, CSCs showed increased expression of enzymes, which are associated with ROS scavenging [76]. Particularly, glutathione synthetase that is involved in glutathione synthesis is upregulated along with Forkhead transcription factor (FOXO)-1 to confer resistance to oxidative stress in hematopoietic stem cells [77]. Also, activation of antioxidant response that is frequently reported in CSCs prevents DNA damage in these cells exposed to ionizing radiations, thereby protecting CSCs against irradiation-induced cell death [78]. Based on these findings, it is widely accepted that cancer recurrence in response to withdrawal of conventional therapies is majorly dependent on existence of a resistant CSC subpopulation within the patients. Therefore, further identification of key molecular drivers that regulate the redox balance in CSCs might provide a possibility to eliminate these cells, which may contribute in overcoming the limitations of cancer relapse in future.

5. Cell death pathways activated by reactive oxygen species

As mentioned above, cancer cells in particular generate increased ROS levels; now severe accumulation of cellular ROS in response to chemotherapy may induce cell cycle arrest, senescence, or lethal toxicity inducing apoptosis [79]. Electrons leaking from the respiratory complexes in mitochondria are a major source for ROS production [80]. For instance, As_2O_3 which impair the function of respiratory chain increases the production of superoxide ions [65]. Alternatively, drugs, which act as redox cyclers such as anthracyclines daunorubicin and doxorubicin, react with cytochrome p450 reductase and NAD(P)H dehydrogenase [quinone] 1(NQO1) in the presence of reduced NADPH to generate superoxide in the presence of oxygen [81].

Apoptosis is linked to an increase in mitochondrial oxidative stress that causes a series of hallmark events such as release of cytochrome c followed by caspases activation ultimately leading to cell death. Sodium salicylate and non-steroidal anti-inflammatory drugs were reported to induce apoptosis in cancers such as colon, breast prostate, and leukemia through ROS production and activation of intrinsic cell death pathway measured by cleavage of caspase-9 and caspase-3 [82]. However, apoptosis was subsequently that a Rac1-NADPH oxidase-dependent pathway is activated in response to treatments that produce ROS and triggers apoptosis [82]. Mitochondrial release of H_2O_2 has been associated with activation of different stress kinases such as c-Jun N-terminal kinase (JNK) and p38. In response to ROS production, JNK mediates phosphorylation and downregulation of anti-apoptotic proteins B-cell lymphoma (Bcl)-2 and Bcl-extra large (xL) [79]. Moreover, several studies reported that both Bcl-2 and Bcl-xL antagonize ROS generation and protect cells against apoptosis [44, 83]. p38 MAPKs are also implicated in apoptosis induction in response to increased ROS production [84]. p38 is activated through apoptosis signal regulating kinase (Ask)-1. Activity of Ask-1 is dependent on a redox-regulated protein thioredoxin that in its reduced form binds to and conserves Ask-1 in an inactivated form. Increased ROS production uncouples thioredoxin from Ask-1 leading to its activation and phosphorylation of p38 required for $\text{TNF}\alpha$ -mediated apoptosis [84]. Studies conducted on L929 fibrosarcoma cells revealed that mitochondrial ROS play a key role in inducing $\text{TNF}\alpha$ cytotoxicity presumably by ROS-mediated caspase activation and cell death [85]. Moreover, TNF receptor associated factor 4 (TNFR4), a component of the TNF signaling chain, binds to NADPH and activates JNK suggesting different mechanisms by which death receptors induce ROS activation in cells [86]. Additionally, different studies have reported the significance of ROS-mediated signaling pathway regulated by protein kinase D1. PDK1 is activated by direct binding to Src and by phosphorylation, which promotes proliferation [35]. Inhibition of this pathway sensitizes cancer cells to ROS. Furthermore, beyond the conventional therapy to induce cytotoxicity to cancer cells and overcome the limitations associated with therapy resistance and risk of developing metastatic phenotype, recent advancement is made to explore the phenomenon of senescence, which inhibits the proliferation of cancer cells and restricts them in a dormant phase [87]. Senescence in cancer cells is mainly characterized by increased activity of β -galactosidase along with modulation of several cell cycle regulators such as cyclin-dependent kinases (CDKs), p16, and p27 [87]. Different

polyphenolic compounds extracted from artichokes (*Cynara cardunculus*) or ginseng (*Panax ginseng*) were described to trigger ROS-dependent senescence.

6. Pathological alterations triggered by free radicals

Intracellular ROS generation may lead to damage of cellular macromolecules such as DNA, proteins, and lipid bilayer. Studies have indicated that H_2O_2 is not very reactive towards DNA; however, the damage to DNA is mainly caused by hydroxyl ions that are generated by the Fenton reaction where transition metals such as iron or copper donate or accept free electrons during intracellular reactions [88]. H_2O_2 acts as a catalyst in the reaction in the formation of free radicals. The generated hydroxyl ions are highly diffusible and lead to DNA damage like oxidation, single-, and double-strand breakage. Under normal physiological conditions, such DNA defects are repaired by base excision repair (BER) or nucleotide excision repair (NER). Cells unable to repair the DNA lesions undergo apoptosis to ensure that the mutations are not passed on during cell division. However, failure in either process of DNA repair or apoptosis may harbor the possibility of formation of cancerous growth.




Figure 5. Molecular mechanisms of ROS-induced macromolecule damage. Scheme was drawn with ScienceSlides Suite 2105 (Visiscience).

ROS-mediated damage of proteins is mainly associated with modifications in specific amino acid residues leading to altered function [89]. Beside, some ROS-mediated modifications of protein also includes increased protein carbonylation, nitration of tyrosine and phenylalanine

residues or formation of cross-linked and glycated proteins [89]. The oxidized amino acid residues in proteins may influence their activity in a signal transduction pathway. For instance, oxidation of phosphatases within the catalytic sites impairs their enzymatic activity [90].

Moreover, ROSs react with polyunsaturated or polyunsaturated fatty acids to trigger lipid peroxidation that has also been used as a tumor biomarker in clinical studies [91]. For instance, in colorectal cancer patients, the presence of thiobarbituric acid reactivates has been linked to high levels of lipid peroxidation [63] (Figure 5).

7. Natural compounds as pharmacological antioxidants

It has been reported in several studies that dietary phytochemicals can interfere with every stage of cancer development. Therefore, antioxidant functions of phytonutrients have been investigated thoroughly for their role in pathophysiology associated with cancer. Dietary antioxidant compounds with significant anticancer activity mainly include anthocyanidins (and their glycosides termed anthocyanins) from berries [92], catechins from green tea, curcumin from turmeric, genistein from soy, resveratrol from grapes and red wine, all-trans lycopene from tomatoes [93], indole-3-carbinol from broccoli, sulforaphane from asparagus, quercetin from red onions and apples. Beside this, carotenoids, flavonoids, and isothiocyanates have also exhibited strong antioxidant properties.




Figure 6. Pharmacological antioxidants of plant origins. Molecular structures were drawn with ChemDraw13 (Perkin Elmer).

Epigallocatechin gallate (EGCG) is the most abundant catechin found in green tea and curcumin-induced anticancer activity promoting cell cycle arrest, polyamine synthesis, and

affecting transglutaminase (TG) activity along with regulation of signaling pathways mediated by NF- κ B, AP-1, and MAPKs [94]. In a recent study, EGCG was shown to inhibit cell proliferation of cervical carcinoma Hela cells by promoting depolymerization of cellular microtubule and disrupting tubulin-microtubule equilibrium. Spectroscopic analysis revealed that EGCG bound to the α -subunit of tubulin at the interphase of α - and β -heterodimers preventing colchicine binding to the colchicine-binding site [95]. Also, in osteosarcoma cells, EGCG treatment induced cell cycle arrest, promoted apoptosis, and inhibited growth of transplanted tumors *in vivo* by regulating miR1/c-MET interaction [96] (Figures 6 and 7).




Figure 7. Molecular mechanisms involved in ROS-triggered survival. Scheme was drawn with ScienceSlides Suite 2105 (Visiscience).

Eugenol (4-allyl-2 methoxyphenol) is a naturally occurring phenolic compound that exhibits antioxidant properties. The antioxidant activity of eugenol was evaluated by the extent of protection offered against free radical-mediated lipid peroxidation using both *in vitro* and *in vivo* studies [97]. The chemopreventive and anticancer role of eugenol was evaluated on *N*-methyl-*N'*-nitro-*N*-nitrosoguanidine (MNNG)-induced gastric cancer in Wistar rats by analyzing the markers of apoptosis, invasion, and angiogenesis. Rats exposed to MNNG developed gastric cancer with upregulation of pro-invasive and angiogenic factors. Eugenol inhibited cell proliferation by suppression of NF- κ B signaling. Apoptosis in these cells following eugenol treatment was mitochondrial pathway mediated that decreased the expression of Bcl-2, following release of cytochrome c and caspases activation. Anti-angiogenic and inhibition of invasion was evidenced by decreased expression of VEGF, its receptor VEGFR1 changes in the activities of MMPs and the expression levels of MMP-2 and MMP-9, VEGF, VEGFR1, tissue inhibitor of metalloproteinases (TIMP)-2 and reversion-inducing cysteine-rich protein with kazal motifs (RECK), a metastasis inhibitor [97].

Several studies aim toward proving the anticancer properties of flavonoids on an array of cancer cell types. Hirano and co-workers tested the anticancer activity of 28 flavonoids on human acute myeloid leukemia (AML) cell line HL-60. Eight of these flavonoids showed strong inhibition of cell proliferation with IC₅₀ values in a nanomolar range [98]. In contingent to this finding, Kuntz et al. showed strong inhibition of proliferation induced by flavonoids on two colon cancer cell models with Caco-2 displaying features of small intestinal epithelial cells and

HT-29, resembling colonic cryptic cells [99]. Moreover, *in vivo* studies on mice strongly inhibited the growth and metastatic potential of melanoma cells B16-BL6 in response to flavonoid treatment [100].

Epigenetic modifications resulting in heritable changes into gene expression without changing the DNA sequence have been marked as key player in promoting cancer [101]. The most common types of epigenetic modifications that may contribute to tumor promotion are DNA methylation and histone acetylation or methylation. Antioxidant compounds mainly isoflavones, flavonols, and catechins have shown to modulate epigenetic features, thereby showing antitumor activity [102–104]. EGCG was shown to affect DNA methyltransferase by inhibiting DNMT and reactivating tumor suppressor genes RAR α , p16, and O⁶-methylguanine methyltransferase in esophageal cancer KYSE 510 cells [105]. Treatment with caffeic acid (3,4-dihydroxycinnamic acid) or chlorogenic acid [106] of hormone-dependent MCF-7 and hormone-independent MDA-MB-231 breast cancer cell lines partially inhibited the methylation of promoter region of the RAR β gene, thereby restoring its function [107]. Furthermore, studies also indicated that dietary antioxidants such as genistein, quercetin, parthenolide, and lycopene may affect DNA methylation status of different genes associated with cancer [108–111].

In addition to this, synergistic or additive effects of phytochemicals could be achieved when administered along with conventional chemotherapy or radiation therapy. This could be explained due to the fact that phytochemicals, which target different biochemical pathways, may enhance the efficacy of conventional therapies. Moreover, different studies have reported the synergistic cytotoxicity on different cancers when phytochemicals are administered together. Apple extracts and quercetin 3- β -D-glucoside combination showed synergistic antiproliferative effect on MCF-7 breast cancer cells [112]. Genistein a major phytoestrogen which has higher affinity for ER β compared to ER α showed synergistic cytotoxicity in combination with indole-3-carbinol in HT-29 cells by simultaneously inhibiting Akt phosphorylation and progression of autophagic process [113]. Combination of δ -tocopherol and resveratrol showed strong inhibition of HMC-1 mastocytoma cell proliferation. The two compounds together strongly inhibited Ser473-phosphorylation of Akt, thereby reducing its activity compared to individual treatment [114]. Gagliano et al. suggested that the use of quercetin in combination with other antioxidants such as resveratrol or sulforaphane might be a novel approach for the treatment of human glioma, which has poor clinical prognosis in both adults and children [115].

Additionally, pharmacological implications of polyphenols have also been explored with respect to inhibition of cancer stem cells and self-renewal. It has been demonstrated that polyphenols can efficiently target pathways such as Wnt/ β -catenin, Hedgehog, and Notch, which are critical for cancer stem, cells self-renewal [116]. Sulforaphane has been demonstrated to target cancer stem cells by modulating the pathways such as NF- κ B, Hedgehog, and Wnt/ β -catenin in different cancers such as breast, pancreas, and prostate and has been proposed as an adjuvant of chemotherapy in different pre-clinical studies [117, 118]. As discussed earlier, cancer stem cells are characterized by a glycolytic metabolism with lower mitochondrial respiration compared to the tumor cells. Therefore, a proposed strategy to counteract CSCs

population is to impair their metabolism by inhibiting glycolysis or by forcing CSCs into mitochondrial metabolism and oxidative phosphorylation. To this purpose, polyphenols have been implicated to regulate the cancer metabolism. For instance, EGCG in human breast cancer have been shown to target the 5' adenosine monophosphate-activated protein kinase (AMPK) pathway, which is involved in maintaining cellular energy status, cell cycle, and protein synthesis [119] (**Figures 8 and 9**).




Figure 8. Pharmacological antioxidants of plant origins. Molecular structures were drawn with ChemDraw13 (Perkin Elmer).




Figure 9. Molecular mechanisms involved in Wnt signaling. Scheme was drawn with ScienceSlides Suite 2105 (Visi-science).

8. Natural compounds as pharmacological pro-oxidants

As discussed earlier, cancer cells produce high levels of ROS that allow these cells to maintain a state of increased basal oxidative stress. The increased state of oxidative stress promotes survival but on the other hand makes the cancer cells vulnerable to further increase in ROS levels over a cancer-specific threshold. Accordingly, pro-oxidant agents and increased oxidative stress levels could then selectively target cancer cells. Different compounds of natural origins modulate the intracellular ROS levels and induce both chemopreventive and anticancer effect in different cancer types.

Polyphenolic extracts from artichokes (*Cynara cardunculus*) at high doses induce apoptosis and decrease the invasive potential of human metastatic breast cancer. Apoptosis was regulated in a caspase-independent manner. Additionally, sublethal concentrations of artichoke increased ROS and induced significant increase in senescence-associated β -galactosidase along with upregulation of tumor suppressor genes p16^{INK4} and p21^{Cip1/Waf1}. Altogether, NAC attenuated the antiproliferative effect induced by artichoke extracts, which suggests that induction of premature senescence and apoptosis is regulated in a ROS-dependent manner [120].

20(S)-ginsenoside Rg3 [20(S)-Rg3], a chemical compound extracted from *Panax ginseng*, induced senescence in glioma cells at sublethal concentrations, which was abrogated by NAC treatment suggesting involvement of ROS. Moreover, depletion of Akt and inactivation of the p53/p21 pathway attenuated the compound-induced senescence. These results suggest that ROS is playing a role in activation of Akt and p53/p21, which leads to growth arrest in human glioma cancer [121].




Figure 10. Molecular scaffolds involved ROS generation. Molecular structures were drawn with ChemDraw13 (Perkin Elmer).

Bisdemethoxycurcumin, a curcuminoid from turmeric, demonstrated potential chemotherapeutic activities by inhibiting proliferation and decreasing the cell viability of hormone-dependent breast cancer. Bisdemethoxycurcumin treatment leads to increased ROS production, which disrupted mitochondrial membrane potential assessed using mitochondrial potential sensor JC-1. Moreover, the compound induced increased expression of pro-apoptotic protein p53 and its downstream effector p21 along with cell cycle regulator p16 and its downstream regulator retinoblastoma protein (pRb). The results overall suggested bisdemethoxycurcumin-induced ROS accumulation, which leads to inhibition of hormone-dependent breast cancer [122].

We have previously reported that garlic-derived organosulfur compounds including diallyl-tetrasulfide induce growth arrest and apoptosis in colon cancer cells by disrupting the redox status in the cells. Drug-induced cell cycle arrest in G2/M phase followed by apoptosis was further associated with decreased Cdc25c expression, one of the key enzymes responsible for G2/M transition [32]. Moreover, we have also shown that plumbagin, a plant naphthoquinone, reduces cell viability and induces apoptosis in a series of hematopoietic cancer cell lines including HL-60, Jurkat, K562, Raji, and U937 with a most pronounced effect on AML U937 cells by 10-fold increase in ROS production. This was followed by decreased expression of anti-apoptotic proteins Mcl-1 and Bcl-2 along with activation of caspases-8, caspases-9, caspases-7, and caspases-3 [123]. Recently, we have also demonstrated ROS induction in neuroblastic and stromal neuroblastoma cells by hemisynthetic cardenolide UNBS1450. ROS induction was followed by autophagic response eventually leading to apoptosis or necroptosis. Time-dependent increase in ROS affected lysosomal integrity of the cells inducing lysosome-associated membrane protein (LAMP)-2 degradation leading to cathepsin B and L activation [124] (**Figure 10**).

9. Conclusion

Natural compounds or their derivatives comprise of more than 50% of cancer chemotherapeutic agents available in the clinics. Information encoded by the human genome project would definitely lead to identification of several gene products, which could potentially be targeted by novel anticancer drugs. Due to various advantages associated with the use of natural compounds such as high availability and reduced toxicity, it is likely that the natural products templates combined with chemistry will allow the generation of novel analogues with enhanced pharmacological benefits to enter clinics.

Malignant cells, which often exhibit increased ROS generation that is associated with tumor proliferation and drug resistance, highlight the crucial role of ROS stress in cancer. Therefore, targeting the redox-modulated biochemical properties of cancer cell may allow to develop a feasible therapeutic approach to overcome challenges associated with cancer treatment. Furthermore, not critically explored unique redox biology of cancer stem cells suggests the use of redox modulating strategies to eradicate these cells.

Acknowledgements

Aloran Mazumder and Marc Diederich are supported by RSL (Luxemburg) and by the Research Institute of Pharmaceutical Sciences, College of Pharmacy, Seoul National University, by the National Research Foundation grant funded by the MEST of Korea for Tumor Micro-environment GCRC 2012-0001184 and by Brain Korea (BK21) PLUS program.

Author details

Aloran Mazumder and Marc Diederich*

*Address all correspondence to: marcdiederich@snu.ac.kr

College of Pharmacy, Seoul National University, Seoul, Korea

References

- [1] Mann J. Natural products in cancer chemotherapy: past, present and future. *Nat Rev Cancer*. 2002;2(2):143–8.
- [2] Zubrod CG. Origins and development of chemotherapy research at the National-Cancer-Institute. *Cancer Treat Rep*. 1984;68(1):9–19.
- [3] Kinghorn AD, Balandrin MF. Human medicinal agents from plants. American Chemical Society (ACS); Oxford University Press UK, 1993.
- [4] Newman DJ, Cragg GM. Natural products as sources of new drugs over the 30 years from 1981 to 2010. *J Nat Prod*. 2012;75(3):311–35.
- [5] Harvey AL, Edrada-Ebel R, Quinn RJ. The re-emergence of natural products for drug discovery in the genomics era. *Nat Rev Drug Discov*. 2015;14(2):111–29.
- [6] Tsuda H, Ohshima Y, Nomoto H, Fujita K-I, Matsuda E, Iigo M, et al. Cancer prevention by natural compounds. *Drug Metab Pharmacokinet*. 2004;19(4):245–63.
- [7] Cerella C, Teiten MH, Radogna F, Dicato M, Diederich M. From nature to bedside: pro-survival and cell death mechanisms as therapeutic targets in cancer treatment. *Biotechnol Adv*. 2014;32(6):1111–22.
- [8] Morceau F, Chateauvieux S, Orsini M, Trecul A, Dicato M, Diederich M. Natural compounds and pharmaceuticals reprogram leukemia cell differentiation pathways. *Biotechnol Adv*. 2015;33(6 Pt 1):785–97.

- [9] Schnekenburger M, Dicato M, Diederich M. Plant-derived epigenetic modulators for cancer treatment and prevention. *Biotechnol Adv.* 2014;32(6):1123–32.
- [10] Schumacher M, Kelkel M, Dicato M, Diederich M. Gold from the sea: marine compounds as inhibitors of the hallmarks of cancer. *Biotechnol Adv.* 2011;29(5):531–47.
- [11] Moriarty-Craige SE, Jones DP. Extracellular thiols and thiol/disulfide redox in metabolism. *Annu Rev Nutr.* 2004;24:481–509.
- [12] Duvoix A, Schmitz M, Schnekenburger M, Dicato M, Morceau F, Galteau MM, et al. Transcriptional regulation of glutathione S-transferase P1-1 in human leukemia. *BioFactors.* 2003;17(1–4):131–8.
- [13] Duvoix A, Schnekenburger M, Delhalle S, Blasius R, Borde-Chiche P, Morceau F, et al. Expression of glutathione S-transferase P1-1 in leukemic cells is regulated by inducible AP-1 binding. *Cancer Lett.* 2004;216(2):207–19.
- [14] Schnekenburger M, Karius T, Diederich M. Regulation of epigenetic traits of the glutathione S-transferase P1 gene: from detoxification toward cancer prevention and diagnosis. *Front Pharmacol.* 2014;5:170.
- [15] Brigelius-Flohe R. Tissue-specific functions of individual glutathione peroxidases. *Free Radic Biol Med.* 1999;27(9–10):951–65.
- [16] Ursini F, Maiorino M, Brigelius-Flohe R, Aumann K, Roveri A, Schomburg D, et al. Diversity of glutathione peroxidases. *Methods Enzymol.* 1994;252:38–53.
- [17] Townsend DM, Tew KD. The role of glutathione-S-transferase in anti-cancer drug resistance. *Oncogene.* 2003;22(47):7369–75.
- [18] Sharma R, Yang Y, Sharma A, Awasthi S, Awasthi YC. Antioxidant role of glutathione S-transferases: protection against oxidant toxicity and regulation of stress-mediated apoptosis. *Antioxid Redox Signal.* 2004;6(2):289–300.
- [19] Montero AJ, Jassem J. Cellular redox pathways as a therapeutic target in the treatment of cancer. *Drugs.* 2011;71(11):1385–96.
- [20] Forman HJ, Torres M, Fukuto J. Redox signaling. Oxygen/nitrogen radicals: cell injury and disease. Springer Science + Business Media New York., 2002. p. 49–62.
- [21] Duvoix A, Blasius R, Delhalle S, Schnekenburger M, Morceau F, Henry E, et al. Chemopreventive and therapeutic effects of curcumin. *Cancer Lett.* 2005;223(2):181–90.
- [22] Reuter S, Eifes S, Dicato M, Aggarwal BB, Diederich M. Modulation of anti-apoptotic and survival pathways by curcumin as a strategy to induce apoptosis in cancer cells. *Biochem Pharmacol.* 2008;76(11):1340–51.
- [23] Teiten MH, Gaascht F, Eifes S, Dicato M, Diederich M. Chemopreventive potential of curcumin in prostate cancer. *Genes Nutr.* 2010;5(1):61–74.

- [24] Teiten MH, Dicato M, Diederich M. Curcumin as a regulator of epigenetic events. *Mol Nutr Food Res*. 2013;57(9):1619–29.
- [25] Kelkel M, Jacob C, Dicato M, Diederich M. Potential of the dietary antioxidants resveratrol and curcumin in prevention and treatment of hematologic malignancies. *Molecules*. 2010;15(10):7035–74.
- [26] Teiten MH, Gaascht F, Dicato M, Diederich M. Targeting the wingless signaling pathway with natural compounds as chemopreventive or chemotherapeutic agents. *Curr Pharm Biotechnol*. 2012;13(1):245–54.
- [27] Spagnuolo C, Cerella C, Russo M, Chateauvieux S, Diederich M, Russo GL. Quercetin downregulates Mcl-1 by acting on mRNA stability and protein degradation. *Br J Cancer*. 2011;105(2):221–30.
- [28] Cerella C, Gagneaux A, Dicato M, Diederich M. Antagonistic role of natural compounds in mTOR-mediated metabolic reprogramming. *Cancer Lett*. 2015;356(2 Pt A):251–62.
- [29] Cerella C, Dicato M, Jacob C, Diederich M. Chemical properties and mechanisms determining the anti-cancer action of garlic-derived organic sulfur compounds. *Anticancer Agents Med Chem*. 2011;11(3):267–71.
- [30] Kelkel M, Cerella C, Mack F, Schneider T, Jacob C, Schumacher M, et al. ROS-independent JNK activation and multisite phosphorylation of Bcl-2 link diallyl tetrasulfide-induced mitotic arrest to apoptosis. *Carcinogenesis*. 2012;33(11):2162–71.
- [31] Czepukojs B, Baltes AK, Cerella C, Kelkel M, Viswanathan UM, Salm F, et al. Synthetic polysulfane derivatives induce cell cycle arrest and apoptotic cell death in human hematopoietic cancer cells. *Food Chem Toxicol*. 2014;64:249–57.
- [32] Busch C, Jacob C, Anwar A, Burkholz T, Aicha Ba L, Cerella C, et al. Diallylpolysulfides induce growth arrest and apoptosis. *Int J Oncol*. 2010;36(3):743–9.
- [33] Viry E, Anwar A, Kirsch G, Jacob C, Diederich M, Bagrel D. Antiproliferative effect of natural tetrasulfides in human breast cancer cells is mediated through the inhibition of the cell division cycle 25 phosphatases. *Int J Oncol*. 2011;38(4):1103–11.
- [34] Anwar A, Burkholz T, Scherer C, Abbas M, Lehr C-M, Diederich M, et al. Naturally occurring reactive sulfur species, their activity against Caco-2 cells, and possible modes of biochemical action. *J Sulfur Chem*. 2011;29(3–4):251–68.
- [35] Storz P. Reactive oxygen species in tumor progression. *Front Biosci*. 2005;10(1–3):1881–96.
- [36] Burdon RH, Gill V, Rice-Evans C. Oxidative stress and tumour cell proliferation. *Free Radic Res Commun*. 1990;11(1–3):65–76.
- [37] Reddy KB, Glaros S. Inhibition of the MAP kinase activity suppresses estrogen-induced breast tumor growth both in vitro and in vivo. *Int J Oncol*. 2007;30(4):971–5.

- [38] Parkash J, Felty Q, Roy D. Estrogen exerts a spatial and temporal influence on reactive oxygen species generation that precedes calcium uptake in high-capacity mitochondria: implications for rapid nongenomic signaling of cell growth. *Biochemistry*. 2006;45(9):2872–81.
- [39] Li Q, Engelhardt JF. Interleukin-1 β induction of NF κ B is partially regulated by H₂O₂-mediated activation of NF κ B-inducing kinase. *J Biol Chem*. 2006;281(3):1495–505.
- [40] Manna SK, Zhang HJ, Yan T, Oberley LW, Aggarwal BB. Overexpression of manganese superoxide dismutase suppresses tumor necrosis factor-induced apoptosis and activation of nuclear transcription factor- κ B and activated protein-1. *J Biol Chem*. 1998;273(21):13245–54.
- [41] Felty Q, Singh KP, Roy D. Estrogen-induced G1/S transition of G0-arrested estrogen-dependent breast cancer cells is regulated by mitochondrial oxidant signaling. *Oncogene*. 2005;24(31):4883–93.
- [42] Park S-A, Na H-K, Kim E-H, Cha Y-N, Surh Y-J. 4-Hydroxyestradiol induces anchorage-independent growth of human mammary epithelial cells via activation of I κ B kinase: potential role of reactive oxygen species. *Cancer Res*. 2009;69(6):2416–24.
- [43] Basseres DS, Baldwin AS. Nuclear factor- κ B and inhibitor of κ B kinase pathways in oncogenic initiation and progression. *Oncogene*. 2006;25(51):6817–30.
- [44] Li N, Karin M. Is NF- κ B the sensor of oxidative stress? *FASEB J*. 1999;13(10):1137–43.
- [45] Chen CL, Lin CF, Chiang CW, Jan MS, Lin YS. Lithium inhibits ceramide- and etoposide-induced protein phosphatase 2A methylation, Bcl-2 dephosphorylation, caspase-2 activation, and apoptosis. *Mol Pharmacol*. 2006;70(2):510–7.
- [46] Wang Y, Huang X, Cang H, Gao F, Yamamoto T, Osaki T, et al. The endogenous reactive oxygen species promote NF- κ B activation by targeting on activation of NF- κ B-inducing kinase in oral squamous carcinoma cells. *Free Radical Res*. 2007;41(9):963–71.
- [47] Cristofanon S, Uguccioni F, Cerella C, Radogna F, Dicato M, Ghibelli L, et al. Intracellular prooxidant activity of melatonin induces a survival pathway involving NF- κ B activation. *Ann N Y Acad Sci*. 2009;1171:472–8.
- [48] Radogna F, Diederich M, Ghibelli L. Melatonin: a pleiotropic molecule regulating inflammation. *Biochem Pharmacol*. 2010;80(12):1844–52.
- [49] Radogna F, Cristofanon S, Paternoster L, D'Alessio M, De Nicola M, Cerella C, et al. Melatonin antagonizes the intrinsic pathway of apoptosis via mitochondrial targeting of Bcl-2. *J Pineal Res*. 2008;44(3):316–25.
- [50] Radogna F, Albertini MC, De Nicola M, Diederich M, Bejarano I, Ghibelli L. Melatonin promotes Bax sequestration to mitochondria reducing cell susceptibility to apoptosis via the lipoxygenase metabolite 5-hydroxyeicosatetraenoic acid. *Mitochondrion*. 2015;21:113–21.

- [51] Cerella C, D'Alessio M, Cristofanon S, De Nicola M, Radogna F, Dicato M, et al. Subapoptogenic oxidative stress strongly increases the activity of the glycolytic key enzyme glyceraldehyde 3-phosphate dehydrogenase. *Ann N Y Acad Sci.* 2009;1171:583–90.
- [52] Cristofanon S, Nuccitelli S, D'Alessio M, Radogna F, De Nicola M, Bergamaschi A, et al. Oxidative upregulation of Bcl-2 in healthy lymphocytes. *Ann N Y Acad Sci.* 2006;1091:1–9.
- [53] Cristofanon S, Morceau F, Scovassi AI, Dicato M, Ghibelli L, Diederich M. Oxidative, multistep activation of the noncanonical NF-kappaB pathway via disulfide Bcl-3/p50 complex. *FASEB J.* 2009;23(1):45–57.
- [54] Cristofanon S, Dicato M, Ghibelli L, Diederich M. Oxidative stress and apoptosis. Redox signaling and regulation in biology and medicine. Wiley-VCH Verlag GmbH & Co. KGaA Weinheim, 2009. p. 373–83.
- [55] Brunet A, Bonni A, Zigmond MJ, Lin MZ, Juo P, Hu LS, et al. Akt promotes cell survival by phosphorylating and inhibiting a Forkhead transcription factor. *Cell.* 1999;96(6):857–68.
- [56] Roberts P, Der C. Targeting the Raf-MEK-ERK mitogen-activated protein kinase cascade for the treatment of cancer. *Oncogene.* 2007;26(22):3291–310.
- [57] Lander HM, Hajjar DP, Hempstead BL, Mirza UA, Chait BT, Campbell S, et al. A molecular redox switch on p21ras structural basis for the nitric oxide-p21ras interaction. *J Biol Chem.* 1997;272(7):4323–26.
- [58] Chan DW, Liu VW, Tsao GS, Yao K-M, Furukawa T, Chan KK, et al. Loss of MKP3 mediated by oxidative stress enhances tumorigenicity and chemoresistance of ovarian cancer cells. *Carcinogenesis.* 2008;29(9):1742–50.
- [59] McCubrey JA, Steelman LS, Chappell WH, Abrams SL, Wong EW, Chang F, et al. Roles of the Raf/MEK/ERK pathway in cell growth, malignant transformation and drug resistance. *Biochimica et Biophysica Acta (BBA) MolCell Res.* 2007;1773(8):1263–84.
- [60] Liu L-Z, Hu X-W, Xia C, He J, Zhou Q, Shi X, et al. Reactive oxygen species regulate epidermal growth factor-induced vascular endothelial growth factor and hypoxia-inducible factor-1 α expression through activation of AKT and P70S6K1 in human ovarian cancer cells. *Free Radical Biol Med.* 2006;41(10):1521–33.
- [61] Mochizuki T, Furuta S, Mitsushita J, Shang W, Ito M, Yokoo Y, et al. Inhibition of NADPH oxidase 4 activates apoptosis via the AKT/apoptosis signal-regulating kinase 1 pathway in pancreatic cancer PANC-1 cells. *Oncogene.* 2006;25(26):3699–707.
- [62] Burdick AD, Davis JW, 2nd, Liu KJ, Hudson LG, Shi H, Monske ML, et al. Benzo(a)pyrene quinones increase cell proliferation, generate reactive oxygen species, and trans-activate the epidermal growth factor receptor in breast epithelial cells. *Cancer Res.* 2003;63(22):7825–33.

- [63] Liou G-Y, Storz P. Reactive oxygen species in cancer. *Free Radical Res.* 2010;44(5):479–96.
- [64] Broom OJ, Massoumi R, Sjolander A. Alpha2beta1 integrin signalling enhances cyclooxygenase-2 expression in intestinal epithelial cells. *J Cell Physiol.* 2006;209(3):950–8.
- [65] Pelicano H, Lu W, Zhou Y, Zhang W, Chen Z, Hu Y, et al. Mitochondrial dysfunction and reactive oxygen species imbalance promote breast cancer cell motility through a CXCL14-mediated mechanism. *Cancer Res.* 2009;69(6):2375–83.
- [66] Hitchler MJ, Oberley LW, Domann FE. Epigenetic silencing of SOD2 by histone modifications in human breast cancer cells. *Free Radical Biol Med.* 2008;45(11):1573–80.
- [67] Lewis A, Du J, Liu J, Ritchie JM, Oberley LW, Cullen JJ. Metastatic progression of pancreatic cancer: changes in antioxidant enzymes and cell growth. *Clin Exp Metastasis.* 2005;22(7):523–32.
- [68] Heneberg P. Reactive nitrogen species and hydrogen sulfide as regulators of protein tyrosine phosphatase activity. *Antioxidants Redox Signal.* 2014;20(14):2191–209.
- [69] Yano S, Arroyo N, Yano N. SHP2 binds catalase and acquires a hydrogen peroxide-resistant phosphatase activity via integrin-signaling. *FEBS Lett.* 2004;577(3):327–32.
- [70] Werner E, Werb Z. Integrins engage mitochondrial function for signal transduction by a mechanism dependent on Rho GTPases. *J Cell Biol.* 2002;158(2):357–68.
- [71] Lopez-Lazaro M. The warburg effect: why and how do cancer cells activate glycolysis in the presence of oxygen? *Anti-Cancer Agents Med Chem (Formerly Current Medicinal Chemistry-Anti-Cancer Agents).* 2008;8(3):305–12.
- [72] Harris AL. Hypoxia—a key regulatory factor in tumour growth. *Nat Rev Cancer.* 2002;2(1):38–47.
- [73] Spitz DR, Sim JE, Ridnour LA, Galoforo SS, Lee YJ. Glucose deprivation-induced oxidative stress in human tumor cells. A fundamental defect in metabolism? *Ann N Y Acad Sci.* 2000;899(1):349–62.
- [74] Semenza GL, Roth PH, Fang H-M, Wang GL. Transcriptional regulation of genes encoding glycolytic enzymes by hypoxia-inducible factor 1. *J Biol Chem.* 1994;269(38):23757–63.
- [75] Xia C, Meng Q, Liu L-Z, Rojanasakul Y, Wang X-R, Jiang B-H. Reactive oxygen species regulate angiogenesis and tumor growth through vascular endothelial growth factor. *Cancer Res.* 2007;67(22):10823–30.
- [76] Visvader JE, Lindeman GJ. Cancer stem cells in solid tumours: accumulating evidence and unresolved questions. *Nat Rev Cancer.* 2008;8(10):755–68.

- [77] Tothova Z, Kollipara R, Huntly BJ, Lee BH, Castrillon DH, Cullen DE, et al. FoxOs are critical mediators of hematopoietic stem cell resistance to physiologic oxidative stress. *Cell*. 2007;128(2):325–39.
- [78] Diehn M, Cho RW, Lobo NA, Kalisky T, Dorie MJ, Kulp AN, et al. Association of reactive oxygen species levels and radioresistance in cancer stem cells. *Nature*. 2009;458(7239):780–3.
- [79] Cadenas E. Mitochondrial free radical production and cell signaling. *Mol Aspects Med*. 2004;25(1):17–26.
- [80] Szatrowski TP, Nathan CF. Production of large amounts of hydrogen peroxide by human tumor cells. *Cancer Res*. 1991;51(3):794–8.
- [81] Vásquez-Vivar J, Martasek P, Hogg N, Masters BSS, Pritchard KA, Kalyanaraman B. Endothelial nitric oxide synthase-dependent superoxide generation from adriamycin. *Biochemistry*. 1997;36(38):11293–7.
- [82] Chung YM, Bae YS, Lee SY. Molecular ordering of ROS production, mitochondrial changes, and caspase activation during sodium salicylate-induced apoptosis. *Free Radical Biol Med*. 2003;34(4):434–42.
- [83] Gottlieb E, Vander Heiden MG, Thompson CB. Bcl-xL prevents the initial decrease in mitochondrial membrane potential and subsequent reactive oxygen species production during tumor necrosis factor alpha-induced apoptosis. *Mol Cell Biol*. 2000;20(15):5680–9.
- [84] Saitoh M, Nishitoh H, Fujii M, Takeda K, Tobiume K, Sawada Y, et al. Mammalian thioredoxin is a direct inhibitor of apoptosis signal-regulating kinase (ASK) 1. *EMBO J*. 1998;17(9):2596–606.
- [85] Schulze-Osthoff K, Beyaert R, Vandevoorde V, Haegeman G, Fiers W. Depletion of the mitochondrial electron transport abrogates the cytotoxic and gene-inductive effects of TNF. *EMBO J*. 1993;12(8):3095.
- [86] Xu YC, Wu RF, Gu Y, Yang Y-S, Yang M-C, Nwariaku FE, et al. Involvement of TRAF4 in oxidative activation of c-Jun N-terminal kinase. *J Biol Chem*. 2002;277(31):28051–57.
- [87] Perez-Mancera PA, Young AR, Narita M. Inside and out: the activities of senescence in cancer. *Nat Rev Cancer*. 2014;14(8):547–58.
- [88] Imlay JA, Chin SM, Linn S. Toxic DNA damage by hydrogen peroxide through the Fenton reaction in vivo and in vitro. *Science*. 1988;240(4852):640–2.
- [89] Levine RL. Carbonyl modified proteins in cellular regulation, aging, and disease 2, 3. *Free Radical Biol Med*. 2002;32(9):790–6.
- [90] Squier TC, Bigelow DJ. Protein oxidation and age-dependent alterations in calcium homeostasis. *Front Biosci*. 2000;5:D504–26.

- [91] Lauschke H, Tolba R, Burger B, Minor T, Hirner A. Lipid peroxidation as additional marker in patients with colorectal cancer. Results of a preliminary study. *Eur Surg Res.* 2002;34(5):346–50.
- [92] Folmer F, Basavaraju U, Jaspars M, Hold G, El-Omar E, Dicato M, et al. Anticancer effects of bioactive berry compounds. *Phytochem Rev.* 2014;13(1):295–322.
- [93] Kelkel M, Schumacher M, Dicato M, Diederich M. Antioxidant and anti-proliferative properties of lycopene. *Free Radic Res.* 2011;45(8):925–40.
- [94] Lamoral-Theys D, Pottier L, Dufrasne F, Neve J, Dubois J, Kornienko A, et al. Natural polyphenols that display anticancer properties through inhibition of kinase activity. *Curr Med Chem.* 2010;17(9):812–25.
- [95] Chakrabarty S, Ganguli A, Das A, Nag D, Chakrabarti G. Epigallocatechin-3-gallate shows anti-proliferative activity in HeLa cells targeting tubulin-microtubule equilibrium. *Chem-Biol Interact.* 2015;242:380–9.
- [96] Zhu K, Wang W. Green tea polyphenol EGCG suppresses osteosarcoma cell growth through upregulating miR-1. *Tumor Biol.* 2015;37:1–10.
- [97] Manikandan P, Murugan RS, Priyadarsini RV, Vinothini G, Nagini S. Eugenol induces apoptosis and inhibits invasion and angiogenesis in a rat model of gastric carcinogenesis induced by MNNG. *Life Sci.* 2010;86(25):936–41.
- [98] Hirano T, Gotoh M, Oka K. Natural flavonoids and lignans are potent cytostatic agents against human leukemic HL-60 cells. *Life Sci.* 1994;55(13):1061–9.
- [99] Kuntz S, Wenzel U, Daniel H. Comparative analysis of the effects of flavonoids on proliferation, cytotoxicity, and apoptosis in human colon cancer cell lines. *Eur J Nutr.* 1999;38(3):133–42.
- [100] Caltagirone S, Rossi C, Poggi A, Ranelletti FO, Natali PG, Brunetti M, et al. Flavonoids apigenin and quercetin inhibit melanoma growth and metastatic potential. *Int J Cancer.* 2000;87(4):595–600.
- [101] Rodriguez-Paredes M, Esteller M. Cancer epigenetics reaches mainstream oncology. *Nat Med.* 2011;17(3):330–9.
- [102] Pudenz M, Roth K, Gerhauser C. Impact of Soy isoflavones on the epigenome in cancer prevention. *Nutrients.* 2014;6(10):4218–72.
- [103] Gilbert ER, Liu D. Flavonoids influence epigenetic-modifying enzyme activity: structure – function relationships and the therapeutic potential for cancer. *Curr Med Chem.* 2010;17(17):1756–68.
- [104] Henning SM, Wang P, Carpenter CL, Heber D. Epigenetic effects of green tea polyphenols in cancer. *Epigenomics.* 2013;5(6):729–41.

- [105] Yang CS, Fang M, Lambert JD, Yan P, Huang TH. Reversal of hypermethylation and reactivation of genes by dietary polyphenolic compounds. *Nutr Rev.* 2008;66(Suppl. 1):S18–20.
- [106] Gaascht F, Dicato M, Diederich M. Coffee provides a natural multitarget pharmacopeia against the hallmarks of cancer. *Genes Nutr.* 2015;10(6):51.
- [107] Lee WJ, Zhu BT. Inhibition of DNA methylation by caffeic acid and chlorogenic acid, two common catechol-containing coffee polyphenols. *Carcinogenesis.* 2006;27(2):269–77.
- [108] Xie Q, Bai Q, Zou LY, Zhang QY, Zhou Y, Chang H, et al. Genistein inhibits DNA methylation and increases expression of tumor suppressor genes in human breast cancer cells. *Genes Chromosomes Cancer.* 2014;53(5):422–31.
- [109] Liu C-M, Ma J-Q, Xie W-R, Liu S-S, Feng Z-J, Zheng G-H, et al. Quercetin protects mouse liver against nickel-induced DNA methylation and inflammation associated with the Nrf2/HO-1 and p38/STAT1/NF-κB pathway. *Food Chem Toxicol.* 2015;82:19–26.
- [110] Liu Z, Liu S, Xie Z, Pavlovicz RE, Wu J, Chen P, et al. Modulation of DNA methylation by a sesquiterpene lactone parthenolide. *J Pharmacol Exp Ther.* 2009;329(2):505–14.
- [111] Fu LJ, Ding YB, Wu LX, Wen CJ, Qu Q, Zhang X, et al. The effects of lycopene on the methylation of the GSTP1 promoter and global methylation in prostatic cancer cell lines PC3 and LNCaP. *Int J Endocrinol.* 2014;2014:620165.
- [112] Yang J, Liu RH. Synergistic effect of apple extracts and quercetin 3-β-D-glucoside combination on antiproliferative activity in MCF-7 human breast cancer cells in vitro. *J Agric Food Chem.* 2009;57(18):8581–6.
- [113] Nakamura Y, Yogosawa S, Izutani Y, Watanabe H, Otsuji E, Sakai T. A combination of indol-3-carbinol and genistein synergistically induces apoptosis in human colon cancer HT-29 cells by inhibiting Akt phosphorylation and progression of autophagy. *Mol Cancer.* 2009;8(100):100.
- [114] Reiter E, Azzi A, Zingg JM. Enhanced anti-proliferative effects of combinatorial treatment of α-tocopherol and resveratrol in human HMC?1 cells. *BioFactors.* 2007;30(2):67–77.
- [115] Gagliano N, Aldini G, Colombo G, Rossi R, Colombo R, Gioia M, et al. The potential of resveratrol against human gliomas. *Anti-Cancer Drugs.* 2010;21(2):140–50.
- [116] Takahashi-Yanaga F, Kahn M. Targeting Wnt signaling: can we safely eradicate cancer stem cells? *Clin Cancer Res.* 2010;16(12):3153–62.
- [117] Li Y, Yuan YY, Meeran SM, Tollefsbol TO. Synergistic epigenetic reactivation of estrogen receptor-α (ERα) by combined green tea polyphenol and histone deacetylase inhibitor in ERα-negative breast cancer cells. *Mol Cancer.* 2010;9:274.

- [118] Rausch V, Liu L, Kallifatidis G, Baumann B, Mattern J, Gladkich J, et al. Synergistic activity of sorafenib and sulforaphane abolishes pancreatic cancer stem cell characteristics. *Cancer Res.* 2010;70(12):5004–13.
- [119] Hadad SM, Fleming S, Thompson AM. Targeting AMPK: a new therapeutic opportunity in breast cancer. *Crit Rev Oncol/Hematol.* 2008;67(1):1–7.
- [120] Mileo AM, Di Venere D, Abbruzzese C, Miccadei S. Long term exposure to polyphenols of artichoke (*Cynara scolymus* L.) exerts induction of senescence driven growth arrest in the MDA-MB231 human breast cancer cell line. *Oxid Med Cell Longev.* 2014;2014:1–17.
- [121] Sin S, Kim SY, Kim SS. Chronic treatment with ginsenoside Rg3 induces Akt-dependent senescence in human glioma cells. *Int J Oncol.* 2012;41(5):1669–74.
- [122] Li Y-B, Gao J-L, Zhong Z-F, Hoi P-M, Lee SM-Y, Wang Y-T. Bisdemethoxycurcumin suppresses MCF-7 cells proliferation by inducing ROS accumulation and modulating senescence-related pathways. *Pharmacol Rep.* 2013;65(3):700–9.
- [123] Gaascht F, Teiten MH, Cerella C, Dicato M, Bagrel D, Diederich M. Plumbagin modulates leukemia cell redox status. *Molecules.* 2014;19(7):10011–32.
- [124] Radogna F, Cerella C, Gaigneaux A, Christov C, Dicato M, Diederich M. Cell type-dependent ROS and mitophagy response leads to apoptosis or necroptosis in neuroblastoma. *Oncogene.* 2015.